Ciencia Latina Revista Científica Multidisciplinar, Ciudad de México, México.
ISSN 2707-2207 / ISSN 2707-2215 (en línea), enero-febrero 2026,
Volumen 10, Número 1.

https://doi.org/10.37811/cl_rcm.v10i1

TEORÍA CUÁNTICA DE CAMPOS RELATIVISTAS: UNA ALTERNATIVA DE SOLUCIÓN AL PROBLEMA DEL MILENIO DE YANG – MILLS. UN INTENTO POR UNIFICAR LA RELATIVIDAD GENERAL Y LA MECÁNICA CUÁNTICA. VOLUMEN X.

RELATIVISTIC QUANTUM FIELD THEORY: AN ALTERNATIVE SOLUTION TO THE YANG–MILLS MILLENNIUM PROBLEM. AN ATTEMPT TO UNIFY GENERAL RELATIVITY AND QUANTUM MECHANICS. VOLUME X.

Manuel Ignacio Albuja Bustamante
Universidad Laica Eloy Alfaro de Manabí

Arturo Damián Rodríguez Zambrano
Universidad Laica Eloy Alfaro de Manabí

DOI: https://doi.org/10.37811/cl_rcm.v10i1.22952

TEORÍA CUÁNTICA DE CAMPOS RELATIVISTAS: UNA ALTERNATIVA DE SOLUCIÓN AL PROBLEMA DEL MILENIO DE YANG – MILLS. UN INTENTO POR UNIFICAR LA RELATIVIDAD GENERAL Y LA MECÁNICA CUÁNTICA. VOLUMEN X.

Manuel Ignacio Albuja Bustamante[1]

ignaciomanuelalbujabustamante@gmail.com

https://orcid.org/0009-0005-0115-767X

Investigador Independiente

RESUMEN

En este trabajo, compuesto por diez volúmenes, abordaremos aspectos esenciales de la Teoría Cuántica de Campos Relativistas (TCCR), con propósitos de optimización de los cálculos expuestos en trabajos anteriores pero sobre todo, posicionar la referida teoría, como una alternativa de solución al problema del milenio de Yang – Mills y la brecha de masa. La idea esencial es la misma, todo espacio – tiempo cuántico, es decir, todo campo cuántico, es curvo y esa deformación ocurre por la gravedad y supergravedad cuánticas, según sea el caso, que provocan las partículas oscuras o estrella, al momento de interactuar con un campo gravitónico o supergravitónico, según corresponda, o en relación a la criticidad de su centro de masa y/o energía, lo que afecta su espín, velocidad y momento angular y por ende, sus trayectorias orbitales. Por tanto, la TCCR, no es un intento por cuantizar la gravedad, sino por introducir la gravedad, como principio de mínima acción de un sistema cuántico y de sus estados fundamentales. Las métricas siguen siendo las mismas, es decir, que para un campo cuántico curvo o geométricamente deformado, la densidad lagrangiana/hamiltoniana equivale a: $\mathcal{LH}_{curvature} = \langle \int \hat{e}^{i\hbar t} \sqrt{\bar{g}^{\mu\nu}} \otimes \overleftrightarrow{m\psi}\bar{\psi} - \partial^2 \triangle' \rangle' \langle \otimes_{\Re}^{\mathfrak{G}} |d^4x/\partial\mathcal{R}\rangle' \int \left\| \widetilde{\frac{\partial \phi_{\sigma\rho}^{I'}}{\partial \phi_{\sigma\rho}^{\dagger}}} \right\| -$

$\left\langle \frac{\partial \phi_{\sigma\rho}^{\star}}{\partial_{\mu\nu}^{\S}} \right| \partial \uparrow / \partial t \setminus \partial \downarrow / \partial t \partial^2 \square \left| \begin{matrix} \blacksquare^{\circlearrowleft} \\ \blacksquare_{\circlearrowright} \end{matrix} \partial^2 \varphi / \partial \psi \blacksquare \right\rangle \wedge_{\nu}^{\mu} \sum_{\substack{0 \leq \iota \leq m \\ 0<J<n}} P(\iota, J) \prod_{k=1}^{n} A_k \cup_{n=1}^{m} (X_n \cap Y_n) \cup_{n=1}^{m} (X_n \cap Y_n) \circledcirc \Lambda_{\nu}^{\mu} \odot \Gamma_{\nu}^{\mu},$

respecto de una partícula pesada ρ, sea oscura o blanca (partícula estrella), según corresponda, a propósito de la criticidad de su masa y/o energía $\langle 0 | \sum_{\delta} \partial \mathfrak{m} / \partial \mathfrak{e} \rangle$ o de su interacción con un gravitón o un gravitino, según corresponda, en coordenadas $\langle \rho^{\mu} \rho^{\nu} \rho^{\sigma} \rho^{\varrho} \rangle$, esto último, lo que ocurre por permeabilización del campo gravitónico o supergravitónico en $\blacksquare = \int \langle \partial \mathfrak{G} / \partial \mathfrak{S} \mathfrak{G} \rangle$, lo que corresponde al espacio – cuántico deformado en $\mathfrak{C}_{\mathfrak{H}\Re} = \langle \sum_{\blacksquare}^{\sigma\rho} \mathcal{R}_{\nu}^{\mu\dagger} | \otimes \mathcal{H}_{\mu}^{\nu\star} \rangle$ lo que en dimensiones $\mathbb{R}^{\eta}$, representa, gravedad o supergravedad cuánticas por curvatura o supercurvatura del espacio - tiempo cuántico multidimensional.

Palabras Clave: Supergravedad cuántica, gravedad cuántica, partícula oscura, partícula estrella, hiperpartículas, suprapartículas, teoría cuántica de campos relativistas, problema del milenio de Yang – Mills y la brecha de masa, partículas ligeras, curvatura, supercurvatura, multidimensiones, agujeros cuánticos.

[1] Autor principal
Correspondencia: ignaciomanuelalbujabustamante@gmail.com

RELATIVISTIC QUANTUM FIELD THEORY: AN ALTERNATIVE SOLUTION TO THE YANG–MILLS MILLENNIUM PROBLEM. AN ATTEMPT TO UNIFY GENERAL RELATIVITY AND QUANTUM MECHANICS. VOLUME X.

ABSTRACT.

In this work, composed of ten volumes, we will address essential aspects of the Quantum Theory of Relativistic Fields (TCCR), with the purpose of optimizing the calculations exposed in previous works but above all, positioning the aforementioned theory as an alternative solution to the Yang-Mills millennium problem and the mass gap. The essential idea is the same, all quantum space-time, that is, every quantum field, is curved and that deformation occurs due to quantum gravity and supergravity, as the case may be, caused by dark particles or stars, when interacting with a gravitonic or supergravitonic field, as appropriate, or in relation to the criticality of its center of mass and/or energy which affects their spin, velocity and angular momentum and therefore, their orbital trajectories. Therefore, the TCCR is not an attempt to quantize gravity, but to introduce gravity, as the principle of least action of a quantum system and its fundamental states.

The metrics remain the same, i.e., for a curved or geometrically warped quantum field, the Lagrangian/Hamiltonian density is equal to: $\mathcal{LH}_{curvature} = \langle \int \hat{e}^{i\hbar t} \sqrt{\bar{g}^{\mu\nu}} \otimes \overleftrightarrow{m\psi}\bar{\psi} - \partial^2 \triangle' \rangle' \langle \otimes_{\Re}^{\mathfrak{G}} \left| d^4x/\partial\mathcal{R} \right\rangle' \int \left\| \widetilde{\frac{\partial \phi'_{\sigma\rho}}{\partial \phi^{\dagger}_{\sigma\rho}}} \right\| -$

$\left\langle \frac{\partial \phi^{\star}_{\sigma\rho}}{\partial^{\mathfrak{H}}_{\mu\nu}} \right| \partial \uparrow / \partial t \setminus \partial \downarrow / \partial t \partial^2 \square \left| \begin{matrix} \blacksquare^{\circlearrowleft} \\ \blacksquare_{\circlearrowright} \end{matrix} \partial^2 \varphi / \partial \psi \blacksquare \right\rangle \Lambda^{\mu}_{\nu} \sum_{\substack{0 \leq \iota \leq m \\ 0<J<n}} P(\iota, j) \prod_{k=1}^{n} A_k \cup_{n=1}^{m} (X_n \cap Y_n) \cup_{n=1}^{m} (X_n \cap Y_n) \circledcirc \Lambda^{\mu}_{\nu} \odot \Gamma^{\mu}_{\nu}$ with respect to a heavy particle ρ, whether dark or white (star particle), as appropriate, regarding the criticality of its mass and/or energy $\langle 0 | \sum_{\delta} \partial \mathfrak{m} / \partial \mathfrak{e} \rangle$ or its interaction with a graviton or a gravitin, as appropriate, in coordinates $\langle \rho^{\mu} \rho^{\nu} \rho^{\sigma} \rho^{\varrho} \rangle$, the latter, which occurs by permeabilization of the gravitonic or supergravitonic field in $\blacksquare = \int \langle \partial \mathfrak{G} / \partial \mathfrak{S} \mathfrak{G} \rangle$, what corresponds to the space – quantum deformed in $\mathfrak{C}_{\mathfrak{H}\Re} = \langle \sum_{\blacksquare}^{\sigma\rho} \mathcal{R}_{\nu}^{\mu\dagger} \big| \otimes \mathcal{H}_{\mu}^{\nu\star} \rangle$ the which in dimensions $\mathbb{R}^{\eta}$, represents, quantum gravity or supergravity by curvature or supercurvature of multidimensional quantum space-time.

Keywords: Quantum supergravity, quantum gravity, dark particle, star particle, quantum theory of relativistic fields.

Artículo recibido 02 enero 2026
Aceptado para publicación: 30 enero 2026

INTRODUCCIÓN.

En este punto, es indispensable establecer las bases teóricas que conforman la Teoría Cuántica de Campos Relativistas (TCCR) y que se encuentran desarrolladas en trabajos previos. Por tanto, estos son los puntos más relevantes.

1. Todo campo cuántico, es curvo por acción inmediata de una partícula cuya masa y/o energía alcanzan el mayor grado de criticidad. En este caso, la gravedad es endógena o implícita, es decir, una cualidad propia de la partícula interactuante.

2. Siguiendo lo dicho, en el numeral que antecede, las partículas se dividen en:

2.1. Partículas Supermasivas (Tipo IA): Son aquellas, cuyo centro de masa/energía en unidades de Planck dados en $\mathcal{M}_p = \sqrt{\frac{\hbar c}{\mathfrak{G}}} \approx 2{,}18 \times 10^{-8}\, kg$ (masa) y $E_p = \frac{\hbar}{t_P}$, $E_p = m_P c^2$, $E_p = \sqrt{\frac{\hbar c^5}{G}} \approx 1.956 \times 10^9$ J $\approx 1.22 \times 10^{19}$GeV ≈ 0.5433MWh$\sqrt{\frac{\hbar c^5}{8\pi G}} \approx 0.390 \times 10^9$ J $\approx 2.43 \times 10^{18}$GeV (energía $\approx 10^{-120}$), alcanza el mayor grado de criticidad, deformando el espacio – tiempo cuántico, lo que afecta el estado fundamental de los orbitales (espín, momentum, velocidad, trayectorias, etc), desplegados por las partículas repercutidas. Esta partícula también se la denomina "partícula oscura", en la medida en que, su centro de energía/masa, es oscuro. Principal candidata para explicar la materia oscura, en la medida en que, la gravedad converge en su centro, absorbiendo energía y materia.

2.2. Partículas Blancas (Tipo IB): Son aquellas, cuyo centro de masa/ energía en unidades de Planck, alcanza el mayor grado de criticidad, deformando el espacio – tiempo cuántico, lo que afecta el estado fundamental de los orbitales (espín, momentum, velocidad, trayectorias, etc), desplegados por las partículas repercutidas. Esta partícula también se la denomina "partícula estrella", en la medida en que, su centro de masa/energía es extremadamente denso, superando la masa, temperatura y energía de Planck, en $\mathcal{M}_p = \sqrt{\frac{\hbar c}{\mathfrak{G}}} \approx 2{,}18 \times 10^{-8}\, kg$ (masa), $\mathcal{M}_p = \sqrt{\frac{\hbar c^5}{\mathfrak{G}\hbar^2}}$ $T_P \approx 1.416784(16) \times 10^{32}$K (temperatura) y $E_p = \frac{\hbar}{t_P}$, $E_p = m_P c^2$, $E_p = \sqrt{\frac{\hbar c^5}{G}} \approx 1.956 \times 10^9$ J $\approx 1.22 \times 10^{19}$GeV ≈ 0.5433MWh$\sqrt{\frac{\hbar c^5}{8\pi G}} \approx 0.390 \times 10^9$ J $\approx 2.43 \times 10^{18}$GeV. También se la denomina "partícula estrella".

2.3. Hiperpartículas (Tipo IIA): Son aquellas, cuyo centro de masa/energía es extremadamente bajo, en unidades de Planck, más sin embargo, son capaces de igualar o superar la velocidad de la luz.

2.4. Suprapartículas (Tipo IIB): Son aquellas, cuyo centro de masa/energía es el equivalente al de una partícula oscura o blanca, más sin embargo, éstas, a diferencia de las referidas en los numerales 2.1 y 2.2, ésta igual o supera la velocidad de la luz.

3. Agujero negro cuántico: Fenómeno que ocurre en un espacio cuántico de Sitter, esto es, cuando una partícula oscura colisiona con otra o en su defecto, cuando una partícula blanca colisiona con otra o cuando una partícula blanca y una partícula oscura colisionan entre sí. Los agujeros negros cuánticos, también se forman por el colapso (por compresión gravitacional) o por la aniquilación (por interacción) de una partícula oscura o de una partícula blanca. Lo primero, ocurre cuando se atraen mutuamente por gravedad en tanto que lo segundo, ocurre cuando su centro de masa/energía alcanza el mayor grado de criticidad posible. En el centro del agujero negro cuántico, se encuentra la masa de la partícula aniquilada o comprimida, la que comporta condiciones gravitatorias extremas. Ahí es donde radica la singularidad de un agujero negro cuántico. La información que ingresa al agujero negro cuántico, no se destruye, muy al contrario, se transforma en materia y energía, las mismas que son repulsadas por el agujero negro cuántico blanco que se encuentra en el otro extremo del agujero cuántico de gusano. Por tanto, la materia y energía atrapada por el agujero negro cuántico, se convierte en materia y energía oscuras interferidas por gravedad extrema.

4. Agujero cuántico de gusano: Túnel cuántico por el cual, se conectan un agujero negro cuántico y un agujero blanco cuántico. A través de este túnel, por teletransportación cuántica, la información es procesada y convertida en materia y energía, todo esto, en un espacio de Sitter.

5. Agujero blanco cuántico: Fenómeno que ocurre en un espacio cuántico de Sitter, volviéndose la región de salida o repulsión de materia y energía, a propósito de lo que devora el agujero negro cuántico y de lo que procesa el canal cuántico de gusano. Lo que repulsa el agujero blanco cuántico, es materia y energía procesadas.

6. Espacio – tiempo cuántico: Entiéndase por espacio – tiempo cuántico, al campo en sí mismo, cuya Longitud de Planck, es superior a $\ell_{\mathcal{P}} = \sqrt{\frac{\hbar G}{c^3}} \approx 1{,}616199(97) \times 10^{-35}\ metros$**.** La métrica es la

curvatura escalar de Ricci, así: $\mathcal{R} = \sum_{\alpha,\beta=0}^{3} g^{\alpha\beta}\mathcal{R}_{\alpha\beta} \approx o\left(\mathcal{L}_p^{-2}\right) \approx 3{,}828 \cdot 10^{69} m^{-2}$. Ahora bien, el espacio – tiempo cuántico puede ser, bien de Sitter (dS) o bien, anti de Sitter (AdS). En el primero, se forma la curvatura cuántica y sus subniveles, subespacios o subcapas, en tanto que en el segundo, se forman los agujeros cuánticos y las multidimensiones.

7. Todo campo cuántico, es curvo por acción inmediata de la gravedad, esto a propósito de la existencia (Modelo – Higgs):

7.1. De un campo gravitónico, es decir, cuando una partícula cualquiera, interactúa con un gravitón, lo que supone la permeabilidad del campo cuántico, por un campo gravitónico que transfiere gravedad al campo primario, curvándolo.

7.2. De un campo supergravitónico, es decir, cuando una partícula cualquiera, interactúa con un gravitino o supergravitón, lo que supone la permeabilidad del campo cuántico, por un campo gravitónico que transfiere gravedad al campo primario, deformándolo.

7.3. Lo referido en este numeral se denomina gravedad exógena.

8. La gravedad cuántica, sea endógena o exógena comporta la curvatura del espacio – tiempo cuántico, en tanto que, la supergravedad cuántica, sea endógena o exógena, comporta la deformación (supercurvatura) del espacio – tiempo cuántico, formándose pliegues multidimensionales (en alta configuración – membranas dimensionales) en rango superior a $\mathbb{R}^4 - AdS$. Cabe indicar que las membranas dimensionales, se dividen en TIPO I y TIPO II respectivamente, la primera a propósito de la curvatura del campo en gravedad cuántica y la segunda, la deformación del campo en supergravedad cuántica, todo esto, lo cual también depende de la naturaleza de la gravedad que interfiere, es decir, si es exógena o endógena, lo que llamaríamos membranas dimensionales tipo IA, IB, IIA y IIB respectivamente, las cuales, pueden contener dimensiones y subdimensiones infinitas, en relación a las interacciones de la partícula que provoca de la deformación del espacio – tiempo cuántico. Esto es lo que llamamos supersimetrías de gauge en dimensiones altas a $\mathbb{R}^4$, es decir, cuando estamos ante membranas dimensionales tipo IA, IB y IIB, según sea el caso en tanto que, las membranas dimensionales del tipo IIA, contienen dimensiones infinitas en $\mathbb{R}^4 - dS$.

9. Cuando una partícula colisiona con otra y se aniquilan o cuando la partícula pesada colapsa por compresión, la extinción provoca ondas cuánticas que se desplazan en longitud sobre el campo cuántico deformado el mismo que, es superfluido.

10. El puente ER, en esta teoría, explica la superposición y el entrelazamiento cuánticos en sentido estricto, en un espacio AdS.

11. Los enunciados antes referidos, aplican a la antimateria, es decir, a la región de antipartículas.

12. La brecha de masa, provoca la curvatura del espacio – tiempo cuántico pero no lo deforma por completo, pues este fenómeno, no ocurre con una partícula deformante, sino en partículas ligeras como las hiperpartículas, esto en la medida en que, no registran estado de vacío.

13. Adicionalmente, es importante, establecer las siguientes reglas:

13.1. La gravedad cuántica relativista, ocurre concretamente en un espacio cuántico de Sitter, en el que se pueden formar subdimensiones o subespacios dentro del límite de $\mathbb{R}^4$.

13.2. La supergravedad cuántica relativista, ocurre concretamente en un espacio cuántico anti de Sitter, en el que se pueden formar hiperespacios o dimensiones más altas, superiores a $\mathbb{R}^4$.

13.3. Las partículas propuestas, viajan en gravedad cuántica más, interactúan en supergravedad cuántica por permeabilización.

13.4. Cualquier partícula, de las aquí propuestas, se puede convertir en otra, por aniquilación, siguiendo los diagramas de Feynman.

13.5. Las dimensiones en alta configuración así como las de ensamble, son infinitas.

13.6. La materia y energía oscuras, están formadas esencialmente por partículas aniquiladas o colapsadas por gravedad. En consecuencia, es la criticidad de la masa la que las vuelve compatibles.

13.7. Los agujeros cuánticos, absorben partículas ligeras y pesadas, sin distinción, lo que explica la expansión del universo por acción gravitacional en la materia.

13.8. Las partículas aquí propuestas, son susceptibles de enganche, como ocurre con un diquark.

13.9. En esta teoría, se incorpora el concepto de cuerda, pero en un espacio cuántico anti de Sitter.

13.10. Las partículas pesadas, cuando se desplazan de un punto a otro en forma infinita hasta su aniquilación o colapso, lo hacen por medio de gravedad, deformando, en el caso de las partículas blancas

y las hiperpartículas, un espacio de Sitter, creando capas dimensionales en límite de $\mathbb{R}^4$ en tanto que, la partícula oscura, crea capas dimensiones en alta configuración a $\mathbb{R}^4$ en un espacio anti de Sitter.

13.11. La hiperpartícula es la única en este modelo, que no tiene masa, es por ello que puede viajar a la velocidad de la luz.

13.12. La suprapartícula es por excepción, un caso de mutación por aniquilación, en la medida en que, pese a tratarse de una partícula pesada, con un centro de masa/energía extremadamente crítico y denso, es capaz de viajar a la velocidad de la luz. La suprapartícula solamente existe por aniquilación en entre dos o más partículas pesadas, quedando excluidas las partículas ligeras. Adicionalmente, la suprapartícula, tiene la capacidad de desplazarse entre dimensiones dS y AdS, lo que esta teoría denomina dimensiones en $\mathbb{R}^{\eta}$. En consecuencia, las dimensiones por gravedad y supergravedad, pueden intersecarse por gravedad. En este punto, es pertinente para efectos de ejemplificar, citar el diagrama de Penrose expandido al infinito.

13.13. Los campos de las partículas ligeras, son deformados por acción a distancia, debido a las interacciones de una partícula pesada, esto es, por gravedad.

13.14. Solamente las partículas pesadas pueden deformar el campo propio y de las partículas ligeras, por acción de la gravedad que se desprende de su centro de masa/energía extremo. En consecuencia, la gravedad endógena, se materializa por impermeabilización del campo de Braut – Englert – Higgs respecto de la partícula pesada. El bosón de Higgs es el que transfiere la masa, a las partículas pesadas, aniquilándose con éstas.

13.15. La gravedad exógena, se vuelve posible, por permeabilización de un campo cuántico arbitrario, lo que, como ha quedado explicado en esta teoría, funciona como un mecanismo de Higgs.

13.16. El colapso de una partícula pesada, ocurre por la expansión de su centro de masa/energía, debido a la gravedad interferente, ditalación que es comprimida en contrario, por los límites del campo de la partícula de que se trate, lo que provoca, la deformación del plano cuántico e incluso la formación de agujeros cuánticos, según la criticidad de los valores de masa/energía involucrados.

13.17. La fusión de campos cuánticos, es posible, por acción de la gravedad entre ambos, lo que vuelve posible, su aniquilación.

13.18. Las ondas en un plano cuántico, no solamente se forman por la aniquilación o colpaso de una partícula pesada, sino también, cuando viaja de un punto a otro.

13.19. Las partículas ligeras, crean gravedad mínima a propósito de su centro de masa/energía, la cual sin embargo, es imperceptible aunque superior a cero, pues, contribuye a la aniquilación con otro campo más pesado.

13.20. La gravedad endógena, se debe a que, el campo de Higgs, y por ende, el bosón de Higgs, no solamente transfiere masa a las partículas pesadas y ligeras, con excepción de la hiperpartícula, sino que también, le dota de gravedad, a propósito de la masa transferida.

13.21. Esta teoría es estrictamente de gauge.

RESULTADOS Y DISCUSIÓN:

Suponemos que, en un mapa cuántico de Einstein – Hilbert, una partícula deformante $\alpha\beta\gamma\delta$ se desplaza en el espacio cuántico, en el que interactúa, deformando el plano por gravedad, y por ende, creando, bien dimensiones altas en $\mathbb{R}^4 - AdS$ por supercurvatura (supergravedad cuántica) o bien, dimensiones en $\mathbb{R}^4 - ds$ por curvatura, esto es, en condiciones de gravedad. Para estos efectos, una partícula deformante debe colapsar por compresión gravitacional, aniquilarse cuando interactúa con otras más inestables o con otra partícula pesada, o por permeabilidad del campo gravitónico o supergravitónico en el espacio cuántico curvo, esto último, lo cual ocurre, cuando una partícula pesada interactúa con el gravitón o el gravitino (supergravitino), según sea el caso. Por tanto, la gravedad actúa a nivel cuántico, sea por aniquilación, compresión, ésta última gravitacional o por permeabilización. Suponemos en simultáneo, que una vez, causada la aniquilación o compresión por gravedad, de una partícula pesada o cuando ocurre la permeabilización, se produce, bien la curvatura cuántica, cuya métrica es el tensor es de Riemann – Ricci – Einstein, incluyendo el flujo de la simetría, o en su defecto, la supercurvatura de Weyl, cuya métrica es la de Chern-Simons-Nambu-Goto para supergravedad. La primera, produce subcampos que son subdimensiones de un mismo plano de Sitter (dS), en tanto que la segunda, produce campos en dualidad holográfica, que son dimensiones altas al plano cuatridimensional en un espacio anti de Sitter (AdS). En este sentido, el campo pasa a ser no homeomorfo, difeomorfo e isométrico, afectando los orbitales de las partículas cuyo centro de masa/energía es inferior en unidades de Planck (partículas ligeras) en relación a la partícula que deforma el plano. La interacción y/o aniquilación de

estas partículas deformantes, provoca un agujero negro cuántico (con excepción de las interacciones dadas por las hiperpartículas tipo IIA), formado por materia y energía oscuras, cuya naturaleza es fermiónica/bosónica, esto a propósito de que, la partícula aniquilada o comprimida, engendra materia y energía oscuras, lo que no ocurre en escenarios de permeabilización gravitónica más sí, en escenarios de permeabilización supergravitónica. El agujero cuántico de salida, es blanco, por ende, repulsivo de materia y energía transformada por la gravedad, a través del tracto Einstein – Rosen. Cuando la materia y la energía son transformadas en oscuras, por la gravedad, éstas se comprimen hasta un punto de no retorno/densidad supermasiva, causando dos especies de singularidad inherentes al agujero negro cuántico, siendo éstas, primaria y secundaria, la primera en la que la gravedad es extrema y deforma la materia y la energía, fundiéndose con el núcleo del agujero negro cuántico (que contiene la partícula muerta) y la segunda, en la que la gravedad transforma la materia y la energía, desplazándola a través del tracto Einstein – Rosen y expulsándola a través de un agujero blanco cuántico. Esto es lo que ocurre en escenarios de entrelazamiento y túneles cuánticos supermasivos en los que, la partícula deformante genera gravedad extrema. Llámese también, gravedad absoluta. Queda claro entonces, que el sistema cuántico de agujeros, no se produce en condiciones de gravedad relativa, esto es, cuando ocurre únicamente la curvatura cuántica por gravedad moderada, lo que sucede por ejemplo, con las interacciones dadas por las hiperpartículas tipo IIA o en el caso de la brecha de masa de las partículas ligeras respecto del estado de vacío.

Dicho lo anterior, es que, propongo una posible alternativa de solución al problema del milenio de Yang – Mills y la brecha de masa, a partir de la Teoría Cuántica de Campos Relativistas, la cual se constituye además, como un intento por reconciliar la relatividad general y la mecánica cuántica.

A partir de aquí, sugerimos los cálculos de instantones (para regular la brecha de masa y la densidad de energía por carga), osciladores, propagadores, operadores, mapas, coordenadas vectoriales, orbitales, correladores, propulsores, tensores de stress por curvatura, torsión, escalares, spinors, potenciadores, simetrías y supersimetrías de calibre abelianas y no abelianas en relación a las partículas pesadas y sus interacciones con el espacio cuántico deformado, en tanto que respecto de éste último, los cálculos están vinculados a su geometría e hipergeometría (análisis cohomológico), incluyendo los agujeros cuánticos,

no sin antes aclarar, que las demostraciones matemáticas contenidas en trabajos anteriores, son interdependientes a éste manuscrito y sus diez volúmenes.

Aclarado lo anterior, pasamos a precisar que el Modelo aquí referido, se divide en:

1. Supergravedad cuántica en SYM (Super Yang – Mills).

2. Gravedad cuántica en YM (Yang – Mills).

3. Agujeros cuánticos en YM (Yang – Mills).

4. Modelo de Unificación.

Las métricas usadas son, entre otras:

- Espacios de Einstein – Hilbert.
- Métrica de Chern – Simons.
- Métrica de Kaluza – Klein.
- Métrica de Nambu – Goto.
- Métrica de Feynman – Wheeler.
- Métrica de Born – Oppenheimer.
- Métrica de Hartree – Fock.
- Métrica de Yang – Mills.
- Métrica de Kerr – Newman.
- Espacios de Sitter y anti de Sitter.
- Espacios de Riemann – Perelman – Poincaré.
- Tensores y flujo de Ricci.
- Métrica de Green.
- Métrica de Goldstone.
- Métrica de Brout – Englert – Higgs.
- Métrica de Schwinger – Dyson.
- Métrica de Yukawa.
- Métrica de Von Neumann
- Métrica de Friedman.

MODELO CUÁNTICO DE UNIFICACIÓN RELATIVISTA SYM (YANG – MILLS – PARTE IV).

$F^{\dagger}_{\kappa jk} = \langle j|c^{\dagger}_{\kappa}|k\rangle$, where $|j\rangle$ is the j th local many-body basis state

$$\Sigma = \sum_{m=1}^{\infty} \sum_{j=1}^{C(m)} \sum_{k=1}^{2^m} \Sigma_{j,k}^{(m)}$$

$$\Sigma_{1,1}^{(2)}(\tau) = \quad \tau \quad \tau_2 \quad \tau_1 \quad 0$$

$$= c_{211} \int_0^{\tau} d\tau_2 \int_0^{\tau_2} d\tau_1 \, \Delta_{\nu\lambda}(\tau - \tau_1)\Delta_{\mu\kappa}(\tau_2)$$
$$\times F^{\dagger}_{\nu} \, \mathcal{G}(\tau - \tau_2) \, F^{\dagger}_{\mu} \, \mathcal{G}(\tau_2 - \tau_1) \, F_{\lambda} \, \mathcal{G}(\tau_1) \, F_{\kappa}.$$

$$\Sigma_{1,2}^{(2)}(\tau) = \quad \tau \quad \tau_2 \quad \tau_1 \quad 0$$

$$= c_{212} \int_0^{\tau} d\tau_2 \int_0^{\tau_2} d\tau_1 \, \Delta_{\nu\lambda}(\tau - \tau_1)\Delta_{\kappa\mu}(-\tau_2)$$
$$\times F^{\dagger}_{\nu} \, \mathcal{G}(\tau - \tau_2) \, F_{\kappa} \, \mathcal{G}(\tau_2 - \tau_1) \, F_{\lambda} \, \mathcal{G}(\tau_1) \, F^{\dagger}_{\mu}.$$

$$K(\tau, \omega) = -\frac{e^{-\omega\tau}}{1 + e^{-\beta\omega}}$$

$$K(\tau, \omega) = -K(\beta + \tau, \omega) = -K(-\tau, -\omega) \text{ for } \tau \in (-\beta, 0)$$

$$\Delta_{\nu\lambda}(\tau) \approx \sum_{l=1}^{p} \Delta_{\nu\lambda l} K(\tau, \omega_l).$$

$$\mathcal{O}\big((np)^{m-1}(mr^2N^3 + nrN^3)\big).$$

$r = \mathcal{O}\big(\log(\Lambda)\log(\epsilon^{-1})\big)$, where $\Lambda = \beta\omega_{\max}$ is the dimensionless product

$$p = \mathcal{O}\big(\log(\Lambda)\log(\epsilon^{-1})\big)$$

$\rho(\omega) = \delta(\omega - \omega_0)$, then $\Delta(\tau) = K(\tau, \omega_0)$

$$\Delta_{\nu\lambda}(i\nu_n) \approx \sum_{l=1}^{p} \Delta_{\nu\lambda l} K(i\nu_n, \omega_l) = \sum_{l=1}^{p} \frac{\Delta_{\nu\lambda l}}{i\nu_n - \omega_l},$$

$$K(i\nu_n,\omega)=\int_0^\beta d\tau e^{i\nu_n\tau}K(\tau,\omega)=\frac{1}{i\nu_n-\omega}$$

$$i\nu_n=(2n+1)\pi i/\beta$$

$$D=\{i\nu_n\}_{n=-\infty}^{\infty}$$

$$r^{(k)}(z)=\frac{n(z)}{d(z)}=\sum_{j=1}^{k}\frac{w_jf_j}{z-z_j}\Big/\sum_{j=1}^{k}\frac{w_j}{z-z_j}.$$

$$z_k=\underset{z\in Z\setminus Z^{(k-1)}}{\arg\max}\left|r^{(k-1)}(z)-f(z)\right|$$

$$w^{(k)}=\underset{\|w^{(k)}\|_2=1}{\arg\min}\sum_{z\in Z\setminus Z^{(k)}}|d(z)f(z)-n(z)|^2$$

$$d(z)f(z)-n(z)=\sum_{j=1}^{k}\frac{f(z)-f_j}{z-z_j}w_j$$

$$w^{(k)}=\underset{\|w^{(k)}\|_2=1}{\arg\min}\left\|Aw^{(k)}\right\|_2$$

$$A_{ij}=\frac{f(\zeta_i)-f_j}{\zeta_i-z_j},\zeta_i\in Z\setminus Z^{(k)}$$

$$\begin{pmatrix}0 & w_1 & w_2 & \cdots & w_k\\ 1 & z_1 & & & \\ 1 & & z_2 & & \\ \vdots & & & \ddots & \\ 1 & & & & z_k\end{pmatrix}v=\lambda\begin{pmatrix}0 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & \ddots & \\ & & & & 1\end{pmatrix}v$$

$$\Delta(-i\nu_n)=\Delta(i\nu_n)^\dagger$$

$$\Delta_{\nu\lambda}(i\nu_n)=\int_{-\infty}^{\infty}d\omega\frac{\rho_{\nu\lambda}(\omega)}{i\nu_n-\omega}$$

$$\mathrm{Err}\left(\{\omega_l,\Delta_l\}_{l=1}^{p}\right)=\sum_{n=-\infty}^{\infty}\left\|\Delta(i\nu_n)-\sum_{l=1}^{p}\frac{\Delta_l}{i\nu_n-\omega_l}\right\|_F^2,$$

$$\mathcal{E}\left(\{\omega_l\}_{l=1}^{p}\right)=\min_{\{\Delta_l\}_{l=1}^{p}}\mathrm{Err}\left(\{\omega_l,\Delta_l\}_{l=1}^{p}\right)$$

$$\left(\sum_{\nu,\lambda=1}^{n}\frac{1}{\beta}\int_0^\beta d\tau|f_{\nu\lambda}(\tau)|^2\right)^{1/2}.$$

$$\rho(\omega)=\sum_{k=1}^{3}c_k\delta(\omega-\omega_k)$$

$$\rho(\omega) = \frac{2}{\pi}\sqrt{1-\omega^2}$$

$$\rho(\omega) = \sum_{k=1}^{3} c_k \mathrm{e}^{-(a_k(\omega-\omega_k))^2}$$


$$\rho(\omega) = \sum_{j=1}^{N} \delta(\omega-\omega_j) v_j v_j^{\dagger},$$

$$\sum_{j=1}^{N} \|v_j\|_2^2 = 1$$

$$K(\tau-\tau',\omega) = \frac{K(\tau,\omega)K(\tau',-\omega)}{K(0,-\omega)}$$
$$K(\tau-\tau',\omega) = \frac{K(\tau-\tau'',\omega)K(\tau''-\tau',\omega)}{K(0,\omega)}$$


$$\Delta_{\nu\lambda}(\tau-\tau') \approx \sum_{\omega_l \le 0}^{p} \frac{\Delta_{\nu\lambda l}}{K_l^-(0)} K_l^+(\tau) K_l^-(\tau')$$
$$+ \sum_{\omega_l > 0}^{p} \frac{\Delta_{\nu\lambda l}}{K_l^+(0)} K_l^+(\tau-\tau'') K_l^+(\tau''-\tau').$$

$$K_l^{\pm}(\tau) = K(\tau, \pm\omega_l) \oslash \frac{1}{K^-(0)} = 1 + e^{\beta\omega}$$




$$= \sum_{\omega_l \le 0} \frac{K_l^+(\tau)}{K_l^-(0)} \overline{F}_{\lambda l}^{\dagger} \int_0^{\tau} d\tau_2\, \mathcal{G}(\tau-\tau_2) F_\mu^{\dagger} \Delta_{\mu\kappa}(\tau_2) \int_0^{\tau_2} d\tau_1\, \mathcal{G}(\tau_2-\tau_1) F_\lambda (\mathcal{G}K_l^-)(\tau_1) F_\kappa$$
$$+ \sum_{\omega_l > 0} \frac{1}{K_l^+(0)} \overline{F}_{\lambda l}^{\dagger} \int_0^{\tau} d\tau_2\, (\mathcal{G}K_l^+)(\tau-\tau_2) F_\mu^{\dagger} \Delta_{\mu\kappa}(\tau_2) \int_0^{\tau_2} d\tau_1\, (\mathcal{G}K_l^+)(\tau_2-\tau_1) F_\lambda \mathcal{G}(\tau_1) F_\kappa$$

$$= \sum_{\omega_l \le 0} \frac{1}{K_l^-(0)}$$



$$+ \sum_{\omega_l > 0} \frac{1}{K_l^+(0)}$$



.

$$\bar{F}_{\lambda l}^{\dagger} = \sum_{\nu=1}^{n} \Delta_{\nu\lambda l} F_\nu^{\dagger}$$

$$\left(\mathcal{G}K_l^{\pm}\right)(\tau) = \mathcal{G}(\tau) K_l^{\pm}(\tau)$$

Compute $F_\lambda \mathcal{G}(\tau_1) K_l^-(\tau_1) \left(\mathcal{O}(rN^3)\right)$.

Convolve by $\mathcal{G}\left(\mathcal{O}(r^2N^3)\right)$.

For each κ, multiply by F_κ from the right $\left(\mathcal{O}(nrN^3)\right)$.

For each μ, κ, multiply by $\Delta_{\mu\kappa}$, and sum over $\kappa\big(\mathcal{O}(n^2 r N^2)\big)$.

Multiply by $F_\mu^\dagger$ and sum over $\mu\big(\mathcal{O}(nrN^3)\big)$.

Convolve by $\mathcal{G}\big(\mathcal{O}(r^2 N^3)\big)$.

Multiply by $K_l^+ \bar{F}_{\lambda l}^\dagger\big(\mathcal{O}(rN^3)\big)$.


$$\mathcal{O}\big(np(r^2 N^3 + nrN^3)\big).$$

$$K(\tau' - \tau, \omega) = -K(\tau - \tau', -\omega) \text{ for } \tau > \tau'$$

$$\Delta_{\lambda\nu}(\tau' - \tau) \approx -\sum_{\omega_l > 0}^{p} \frac{\Delta_{\lambda\nu l}}{K_l^+(0)} K_l^-(\tau) K_l^+(\tau') - \sum_{\omega_l \le 0}^{p} \frac{\Delta_{\lambda\nu l}}{K_l^-(0)} K_l^-(\tau - \tau'') K_l^-(\tau'' - \tau')$$

$$\bar{F}_{\nu l} = \sum_{\lambda=1}^{n} \Delta_{\lambda\nu l} F_\lambda$$


$\Sigma_{1,1}^{(3)}(\tau) =$



$$K(\tau - \tau', \omega) = \frac{K\big(\tau - \tau^{(1)}, \omega\big) K\big(\tau^{(1)} - \tau^{(2)}, \omega\big) \cdots K\big(\tau^{(j)} - \tau', \omega\big)}{K^j(0, \omega)}$$

$$\Delta_{\xi\lambda}(\tau_4 - \tau_1) \approx \sum_{\omega_l \le 0}^{p} \frac{\Delta_{\xi\lambda l}}{K_l^-(0)} K_l^+(\tau_4) K_l^-(\tau_1) + \sum_{\omega_l > 0}^{p} \frac{\Delta_{\xi\lambda l}}{(K_l^+(0))^2} K_l^+(\tau_4 - \tau_3) \times K_l^+(\tau_3 - \tau_2) K_l^+(\tau_2 - \tau_1)$$

$$K(\tau' - \tau, \omega) = \frac{(-1)^j}{K^j(0, -\omega)} K\big(\tau' - \tau^{(j)}, -\omega\big) \times K\big(\tau^{(j)} - \tau^{(j-1)}, -\omega\big) \cdots K\big(\tau^{(1)} - \tau, -\omega\big)$$



the hybridization line $\Delta_{\mu\kappa}(\tau)\big(\Delta_{\kappa\mu}(-\tau)\big)$ connecting by $F_\mu^\dagger\Delta_{\mu\kappa}\big(F_\kappa\Delta_{\kappa\mu}\big)$ on its vertex, and $F_\kappa\big(F_\mu^\dagger\big)$ at the zero vertex.

For term l of the $\omega_l \le 0$ sum, place $K_l^+\bar{F}_{\nu l}^\dagger(K_l^-\bar{F}_{\pi l})$ at the primary vertex of the hybridization, $K_l^-F_\nu\big(K_l^+F_\pi^\dagger\big)$ at the secondary vertex, and divide by $K_l^-(0)(-K_l^+(0))$.

For term l of the $\omega_l > 0$ sum, place $\bar{F}_{\nu l}^\dagger(\bar{F}_{\pi l})$ at the primary vertex of the hybridization $F_\nu\big(F_\pi^\dagger\big)$ at the secondary vertex, $K_l^+(K_l^-)$ and backbone between the two vertices, and divide by

$$(K_l^+(0))^j\big((-K_l^-(0))^j\big)$$

$$\mathcal{O}\big((np)^{m-1}(mr^2N^3+nrN^3)\big)$$

$$\begin{aligned}
\hat{H} &= \hat{H}_{\text{dimer}} + \hat{H}_{\text{bath}} + \hat{H}_{\text{coupling}} \\
\hat{H}_{\text{dimer}} &= -v\big(\hat{c}_0^\dagger\hat{c}_1 + \hat{c}_1^\dagger\hat{c}_0\big) + U\hat{n}_0\hat{n}_1 \\
\hat{H}_{\text{bath}} &= -t_{\text{b}}\sum_{k=0}^{1}\big(\hat{b}_{0k}^\dagger\hat{b}_{1k} + \hat{b}_{1k}^\dagger\hat{b}_{0k}\big) \\
\hat{H}_{\text{coupling}} &= -t\sum_{\lambda=0}^{1}\sum_{k=0}^{1}\big(\hat{c}_\lambda^\dagger\hat{b}_{\lambda k} + \hat{b}_{\lambda k}^\dagger\hat{c}_\lambda\big)
\end{aligned}$$

$\Delta(i\nu_n) = 2t^2(i\nu_n I - H_0)^{-1}$, where $H_0 = \begin{pmatrix} 0 & t_{\text{b}} \\ t_{\text{b}} & 0 \end{pmatrix}$ and I is the 2×2 identity matrix

$$\hat{H}_{\text{loc}} = U\sum_{\kappa=0}^{1}\hat{n}_{\kappa\uparrow}\hat{n}_{\kappa\downarrow} + \sum_{\sigma,\sigma'\in\{\uparrow,\downarrow\}}(U' - J_H\delta_{\sigma\sigma'})\hat{n}_{0\sigma}\hat{n}_{1\sigma'}$$
$$+J_H\sum_{\kappa\neq\lambda\in\{0,1\}}\left(\hat{c}^{\dagger}_{\kappa\uparrow}\hat{c}^{\dagger}_{\kappa\downarrow}\hat{c}_{\lambda\downarrow}\hat{c}_{\lambda\uparrow} + \hat{c}^{\dagger}_{\kappa\uparrow}\hat{c}^{\dagger}_{\lambda\downarrow}\hat{c}_{\kappa\downarrow}\hat{c}_{\lambda\uparrow}\right)$$

$$\hat{n}_{\kappa\sigma} = \hat{c}^{\dagger}_{\kappa\sigma}\hat{c}_{\kappa\sigma}$$

$$\rho_{\kappa\lambda}(\omega) = \left(\delta_{\kappa\lambda} + s(1-\delta_{\kappa\lambda})\right)t^2 J(\omega)$$

$$\mu = (3U - 5J_H)/2 + \Delta\mu, \text{ where } \Delta\mu = -1.5$$

$$J(\omega) = \sum_{k=0}^{1}\delta(\omega - \epsilon_k), \text{ with } \epsilon_0 = -2.3t, \epsilon_1 = 2.3t$$

$$J(\omega) = \frac{2}{\pi D^2}\sqrt{D^2 - \omega^2}, \text{ with } D = 2t, s = 1, \text{ and } \beta = 8$$

$$H_{\text{int}} = U\sum_{a}\hat{n}_{a\uparrow}\hat{n}_{a\downarrow} + \frac{1}{2}\sum_{a\neq b}\sum_{\sigma,\sigma'}(U' - J\delta_{\sigma\sigma'})\hat{n}_{a\sigma}\hat{n}_{b\sigma'}$$
$$-\sum_{a\neq b}\left(J\hat{c}^{\dagger}_{a\uparrow}\hat{c}_{a\downarrow}\hat{c}^{\dagger}_{b\downarrow}\hat{c}_{b\uparrow} + J'\hat{c}^{\dagger}_{b\uparrow}\hat{c}^{\dagger}_{b\downarrow}\hat{c}_{a\uparrow}\hat{c}_{a\downarrow}\right),$$

Density operator $\hat{n}_{a\sigma} = \hat{c}^{\dagger}_{a\sigma}\hat{c}_{a\sigma}$, with spin($\sigma \in \{\uparrow,\downarrow\}$) and orbital ($a, b \in \{xz, yz, xy\}$) indices.

$$H_{\text{soc}} = \lambda_{\text{soc}}\sum_{ij}\hat{\Psi}^{\dagger}_{i}[h_{\text{soc}}]_{ij}\hat{\Psi}_{j}$$

$\hat{\Psi} = \left[\hat{c}_{xz\uparrow}, \hat{c}_{yz,\uparrow}, \hat{c}_{xy,\downarrow}, \hat{c}_{xz\downarrow}, \hat{c}_{yz,\downarrow}, \hat{c}_{xy,\uparrow}\right]$ and h_{soc} is the matrix

$$h_{soc} = \begin{pmatrix} 0 & -i & i & & & \\ i & 0 & -1 & & 0 & \\ -i & -1 & 0 & & & \\ & & & 0 & i & i \\ 0 & & -i & 0 & & 1 \\ & & & -i & 1 & 0 \end{pmatrix}.$$

$$\Delta_A(\tau) = \mathbf{t}\cdot G_B(\tau)\cdot\mathbf{t}$$
$$\Delta_B(\tau) = \mathbf{t}\cdot G_A(\tau)\cdot\mathbf{t}$$

$$G_{\{A,B\}}(\tau) = \text{diag}\left(t_{xz}, t_{yz}, t_{xy}, t_{xz}, t_{yz}, t_{xy}\right)$$

$$xz, yz\left(\langle\hat{n}_{xz,\sigma}\rangle = \langle\hat{n}_{yz,\sigma}\rangle \approx 0.5\right)\left(\langle\hat{n}_{xy,\sigma}\rangle \approx 1\right)$$

$$\left(\langle\hat{n}_{xy,\sigma}\rangle \approx 0.9850\right)\left(\langle\hat{n}_{xz,\sigma}\rangle = \langle\hat{n}_{yz,\sigma}\rangle \approx 0.5075\right)$$

$$r(-i\nu_n) = r^{\dagger}(i\nu_n)$$

$\left\{i\nu_{n_j}, -i\nu_{n_j}\right\}$, and then $\mathcal{Z}^{(k)} = \mathcal{Z}^{(k-1)} \cup \left\{\text{i}\nu_{n_j}, -\text{i}\nu_{n_j}\right\}$

$$w^{(k)} = (w_1, \overline{w_1}, \ldots, w_k, \overline{w_k})$$

$$A^{\pm,\nu\lambda} \in \mathbb{C}^{k\times k}$$

$$A_{ij}^{\pm,\nu\lambda} = \frac{\Delta_{\nu\lambda}(\zeta_i) - \Delta_{\nu\lambda}\left(\pm \mathrm{i}\nu_{n_j}\right)}{\zeta_i \mp \mathrm{i}\nu_{n_j}},$$

$$\nu, \lambda = 1, \ldots, n, \zeta_i \in Z \setminus Z^{(k)}, \text{ and } j = 1, \ldots, k$$

$$\min_{w_1,\ldots,w_k\in\mathbb{C}} \sum_{\nu,\lambda=1}^{n} \sum_{i=1}^{k} \left| \sum_{j=1}^{k} A_{ij}^{+,\nu\lambda} w_j + \sum_{j=1}^{k} A_{ij}^{-,\nu\lambda} \overline{w_j} \right|^2$$

$$\text{subject to } \sum_{j=1}^{k} \left|w_j\right|^2 = 1$$

$$\min_{u,v\in\mathbb{R}^k} \sum_{\nu,\lambda=1}^{n} \left\| \mathcal{A}^{\nu\lambda} \begin{pmatrix} u \\ v \end{pmatrix} \right\|_2^2 = \left\| \mathcal{A} \begin{pmatrix} u \\ v \end{pmatrix} \right\|_2^2$$

$$w_j = u_j + i v_j$$

$$\mathcal{A}^{\nu\lambda} = \begin{pmatrix} \mathrm{Re}\left(A^{+,\nu\lambda} + A^{-,\nu\lambda}\right) & -\mathrm{Im}\left(A^{+,\nu\lambda} - A^{-,\nu\lambda}\right) \\ \mathrm{Im}\left(A^{+,\nu\lambda} + A^{-,\nu\lambda}\right) & \mathrm{Re}\left(A^{+,\nu\lambda} - A^{-,\nu\lambda}\right) \end{pmatrix},$$

$$\mathcal{A} \in \mathbb{R}^{2kn^2} \times 2k$$


$$G_{j,k}^{(m)}(\tau) = \sum_{m=1}^{\infty} \sum_{j=1}^{C(m)} \sum_{k=1}^{2^{m-1}} G_{j,k}^{(m)}$$

$$G^{(3)}_{1,2,\nu\kappa}(\tau) = \tau, \nu \quad \tau_3 \quad \tau_4 \quad \Delta_{\xi\mu} \quad \Delta_{\lambda\pi} \quad \tau_2 \quad \tau_1 \quad 0, \kappa = d_{312} \int_{\tau}^{\beta} d\tau_4 \int_{\tau}^{\tau_4} d\tau_3 \int_{0}^{\tau} d\tau_2 \int_{0}^{\tau_2} d\tau_1 \, \Delta_{\lambda\pi}(\tau_1 - \tau_4) \Delta_{\xi\mu}(\tau_3 - \tau_2)$$


$$\times \mathrm{Tr}\left[\mathcal{G}(\beta - \tau_4)\, F_\lambda\, \mathcal{G}(\tau_4 - \tau_3)\, F_\xi^\dagger\, \mathcal{G}(\tau_3 - \tau)\, F_\nu\, \mathcal{G}(\tau - \tau_2)\, F_\mu\, \mathcal{G}(\tau_2 - \tau_1)\, F_\pi^\dagger\, \mathcal{G}(\tau_1 - 0)\, F_\kappa^\dagger\right].$$

$$G^{(3)}_{1,2,\nu\kappa}(\tau) = \quad \Delta_{\lambda\pi} \quad \Delta_{\xi\mu} \quad \beta \quad \tau_4 \quad \tau_3 \quad \tau, \nu \quad \tau_2 \quad \tau_1 \quad 0, \kappa \quad .$$


$d_{312}G^{(3)}_{1,2,\nu\kappa}(\tau) =$




$= -\sum_{\omega_l \le 0} \frac{1}{K_l^+(0)}$




$-\sum_{\omega_l > 0} \frac{1}{(K_l^-(0))^3}$




$= -\sum_{\omega_l \le 0, \omega_{l'} \le 0} \frac{1}{K_l^+(0) K_{l'}^-(0)}$




$-\sum_{\omega_l \le 0, \omega_{l'} > 0} \frac{1}{K_l^+(0) K_{l'}^+(0)}$



$-\sum_{\omega_l > 0, \omega_{l'} \le 0} \frac{1}{(K_l^-(0))^3 K_{l'}^-(0)}$



$-\sum_{\omega_l > 0, \omega_{l'} > 0} \frac{1}{(K_l^-(0))^3 K_{l'}^+(0)}$



.

$$-\sum_{\omega_l \le 0, \omega_{l'} > 0} \frac{1}{K_l^+(0) K_{l'}^+(0)} \mathrm{Tr} \int_\tau^\beta d\tau_4 \mathcal{G}(\beta - \tau_4) K_l^-(\tau_4) \bar{F}_{\pi l} \int_\tau^{\tau_4} d\tau_3 \mathcal{G}(\tau_4 - \tau_3) \bar{F}^\dagger_{\mu l'} \left(\mathcal{G}K_{l'}^+\right)(\tau_3 - \tau)$$

$$\times F_\nu \int_0^\tau d\tau_2 \left(\mathcal{G}K_{l'}^+\right)(\tau - \tau_2) F_\mu \int_0^{\tau_2} d\tau_1 \mathcal{G}(\tau_2 - \tau_1) F_\pi^\dagger (\mathcal{G}K_l^+)(\tau_1) F_\kappa^\dagger$$

$\int_\tau^\beta d\tau' F(\tau - \tau') G(\tau')$ into the usual form $\int_0^\tau d\tau' F(\tau - \tau') G(\tau')$

$$\mathcal{O}\left((np)^{m-1}(mr^2N^3 + nrN^3)\right)$$

$$\left(-\partial_\tau - \hat{H}_{\mathrm{loc}} - \eta_0 I\right)\mathcal{G}_0(\tau) = 0, \mathcal{G}_0(0) = -I$$

$$\mathcal{G}_0(\tau) \equiv -e^{-\tau(\hat{H}_{\mathrm{loc}} + \eta_0 I)}$$

$$Z_0 = -\mathrm{Tr}[\mathcal{G}_0(\beta)].$$

$$\eta_0 = \frac{1}{\beta} \log \mathrm{Tr}\left[e^{-\beta \hat{H}_{\mathrm{loc}}}\right]$$

$$\left(-\partial_\tau - \hat{H}_{\mathrm{loc}} - (\eta_0 + \eta) I - \Sigma *\right)\mathcal{G} = 0, \mathcal{G}(0) = -I,$$

$$(\Sigma * \mathcal{G})(\tau) = \int_0^\tau d\bar{\tau}\Sigma(\tau - \bar{\tau})\mathcal{G}(\tau)$$

$$(I - \eta\mathcal{G}_0 * -\mathcal{G}_0 * \Sigma *)\mathcal{G} = \mathcal{G}_0$$

$$Z = -\mathrm{Tr}[\mathcal{G}(\beta)] = 1$$

$$\Omega(\eta) \equiv -\frac{1}{\beta}\log\, Z(\eta) = \frac{1}{\beta}\log\, \mathrm{Tr}[\mathcal{G}(\beta)] = 0,$$

$$\frac{d\Omega}{d\eta} = -\frac{1}{\beta Z}\frac{dZ}{d\eta} = \frac{1}{\beta Z}\mathrm{Tr}\left[\frac{\partial \mathcal{G}}{\partial \eta}(\beta)\right].$$

$\partial\mathcal{G}/\partial\eta$ in short form $\mathcal{L}\mathcal{G} = \mathcal{G}_0$ with $\mathcal{L} \equiv I - \eta\mathcal{G}_0 * -\mathcal{G}_0 * \Sigma *$

Taking the η-derivative $\frac{d\mathcal{L}}{d\eta}\mathcal{G} + \mathcal{L}\frac{\partial \mathcal{G}}{\partial \eta} = 0$

$$\frac{d\mathcal{L}}{d\eta} = -\mathcal{G}_0 * \left(I + \frac{\partial \Sigma}{\partial \eta} *\right) = -\mathcal{L}\mathcal{G} * \left(I + \frac{\partial \Sigma}{\partial \eta} *\right)$$

$$\mathcal{L}\frac{\partial \mathcal{G}}{\partial \eta} = -\frac{d\mathcal{L}}{d\eta}\mathcal{G} = \mathcal{L}\mathcal{G} * \left(I + \frac{\partial \Sigma}{\partial \eta} *\right)\mathcal{G}$$

$$\frac{\partial \mathcal{G}}{\partial \eta} = \mathcal{G} * \left(I + \frac{\partial \Sigma^*}{\partial \eta}\right)\mathcal{G} = \mathcal{G} * \mathcal{G},$$

$$\frac{d\Omega}{d\eta} = \frac{1}{\beta Z}\mathrm{Tr}[(\mathcal{G} * \mathcal{G})(\beta)] = -\frac{1}{\beta \mathrm{Tr}[\mathcal{G}(\beta)]}\mathrm{Tr}[(\mathcal{G} * \mathcal{G})(\beta)]$$

$$\hat{H}_{\mathrm{loc}} \to \hat{H}_{\mathrm{loc}} + \mu\hat{N}$$

$$\left(I - \eta\mathcal{G}_0 * -\mu\mathcal{G}_0\hat{N} * -\mathcal{G}_0 * \Sigma *\right)\mathcal{G} = \mathcal{G}_0$$

$$\langle\hat{N}\rangle = -\frac{1}{Z}\mathrm{Tr}[\hat{N}\mathcal{G}(\beta)]$$

$$\mathbf{F}(\mathbf{x}) \equiv \begin{bmatrix}\Omega \\ \Delta n\end{bmatrix} = \mathbf{0}, \mathbf{x} = \begin{bmatrix}\eta \\ \mu\end{bmatrix},$$

$$\Delta n(\eta, \mu) \equiv N - \langle\hat{N}\rangle \equiv N + \frac{1}{Z}\mathrm{Tr}[\hat{N}\mathcal{G}(\beta)]$$

$$\Omega(\eta, \mu) \equiv \frac{1}{\beta}\log\, \mathrm{Tr}[\mathcal{G}(\beta)].$$

$$J_{\mathbf{F}} = \frac{d\mathbf{F}}{d\mathbf{x}} = \begin{bmatrix}\frac{\partial\Omega}{\partial\eta} & \frac{\partial\Omega}{\partial\mu} \\ \frac{\partial\Delta n}{\partial\eta} & \frac{\partial\Delta n}{\partial\mu}\end{bmatrix}$$

$$\frac{\partial \Omega}{\partial \eta} = \frac{1}{\beta Z}\mathrm{Tr}[(\mathcal{G} * \mathcal{G})(\beta)]$$
$$\frac{\partial \Omega}{\partial \mu} = \frac{1}{\beta Z}\mathrm{Tr}[(\mathcal{G} * \hat{N}\mathcal{G})(\beta)]$$

$$\frac{\partial \Delta n}{\partial \eta} = \frac{1}{Z^2}\mathrm{Tr}[(\mathcal{G} * \mathcal{G})(\beta)]\mathrm{Tr}[\hat{N}\mathcal{G}(\beta)] + \frac{1}{Z}\mathrm{Tr}[\hat{N}(\mathcal{G} * \mathcal{G})(\beta)]$$

$$\frac{\partial \Delta n}{\partial \mu} = \frac{1}{Z^2}\mathrm{Tr}[(\mathcal{G} * \hat{N}\mathcal{G})(\beta)]\mathrm{Tr}[\hat{N}\mathcal{G}(\beta)] + \frac{1}{Z}\mathrm{Tr}[\hat{N}(\mathcal{G} * \hat{N}\mathcal{G})(\beta)]$$

$$\frac{\partial \mathcal{G}}{\partial \mu} = \mathcal{G} * \left(\hat{N} + \frac{\partial \Sigma}{\partial \mu}\right) * \mathcal{G} = (\mathcal{G} * \hat{N}\mathcal{G})(\tau)$$

$$\mathbf{x} \leftarrow \mathbf{x} - [J_{\mathbf{F}}(\mathbf{x})]^{-1}\mathbf{F}(\mathbf{x})$$

$$\begin{aligned}-\mathcal{L}_{\text{eff}} = \sum_{u_j=u,c} \bar{u}_j \Bigg[& \frac{g_s}{2m_t} T^A \sigma^{\mu\nu} \left(\xi_L^{u_j t} P_L + \xi_R^{u_j t} P_R\right) G^A_{\mu\nu} + \frac{e}{2m_t} \sigma^{\mu\nu} \left(\lambda_L^{u_j t} P_L + \lambda_R^{u_j t} P_R\right) F_{\mu\nu} \\ & + \frac{g_W}{2c_W m_t} \sigma^{\mu\nu} \left(\kappa_L^{u_j t} P_L + \kappa_R^{u_j t} P_R\right) Z_{\mu\nu} - \frac{g_W}{2c_W} \gamma^\mu \left(X_L^{u_j t} P_L + X_R^{u_j t} P_R\right) Z_\mu \\ & - \frac{1}{\sqrt{2}} \left(\eta_L^{u_j t} P_L + \eta_R^{u_j t} P_R\right) H \Bigg] t + \text{ H.C.}\end{aligned}$$

$$\mathcal{L}_{\text{Standar Model Effective Field Theory}} = \mathcal{L}^{(4)}_{\text{SM}} + \sum_{d>4} \frac{\mathcal{C}_i^{(d)}}{\Lambda^{d-4}} \mathcal{O}_i^{(d)}$$

$$\mathcal{L}^{i(6)}_{\text{SMEFT}} = \sum_{p,r} \left(\frac{\mathcal{C}^i_{pr}}{\Lambda^2} \mathcal{O}^i_{pr} + \text{ h.c.}\right)$$

$\mathcal{O}_{pr}^{uG}$	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r)\tilde{\phi} G^A_{\mu\nu}$	$\mathcal{O}_{pr}^{\phi q\,(1)}$	$(\phi^\dagger i \overleftrightarrow{D}_\mu \phi)(\bar{q}_p \gamma^\mu q_r)$
$\mathcal{O}_{pr}^{uB}$	$(\bar{q}_p \sigma^{\mu\nu} u_r)\tilde{\phi} B_{\mu\nu}$	$\mathcal{O}_{pr}^{\phi q\,(3)}$	$(\phi^\dagger i \overleftrightarrow{D}^I_\mu \phi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$\mathcal{O}_{pr}^{uW}$	$(\bar{q}_p \sigma^{\mu\nu} u_r)\tau^I \tilde{\phi} W^I_{\mu\nu}$	$\mathcal{O}_{pr}^{\phi u}$	$(\phi^\dagger i \overleftrightarrow{D}_\mu \phi)(\bar{u}_p \gamma^\mu u_r)$
		$\mathcal{O}_{pr}^{u\phi}$	$(\phi^\dagger \phi)(\bar{q}_p u_r \tilde{\phi})$

$$\Gamma_u = U_u \Gamma_u^{\text{diag}}, \Gamma_d = U_d \Gamma_d^{\text{diag}}, q = \begin{pmatrix} U_{u_L}^\dagger u_L \\ U_{d_L}^\dagger d_L \end{pmatrix}$$

$$\Gamma_u = \Gamma_u^{\text{diag}}, \Gamma_d = V_{\text{CKM}} \Gamma_d^{\text{diag}}, q = \begin{pmatrix} u_L \\ V_{\text{CKM}} d_L \end{pmatrix}$$

$V_{\text{CKM}} = \left(U_{d_L}^\dagger U_{u_L}\right)$, and V_{CKM} is the Cabibbo-Kobayashi-Maskawa (CKM) matrix

$$(\xi_L)_{pr}= \sqrt{2}v\frac{m_t}{g_s}\left(\mathcal{C}_{rp}^{uG}\right)^*, (\xi_R)_{pr} = \sqrt{2}v\frac{m_t}{g_s}\mathcal{C}_{pr}^{uG}$$
$$(\lambda_L)_{pr}= \sqrt{2}v\frac{m_t}{e}\left(s_W\left(\mathcal{C}_{rp}^{uW}\right)^* + c_W\left(\mathcal{C}_{rp}^{uB}\right)^*\right), (\lambda_R)_{pr} = \sqrt{2}v\frac{m_t}{e}\left(s_W\mathcal{C}_{pr}^{uW} + c_W\mathcal{C}_{pr}^{uB}\right)$$
$$(\kappa_L)_{pr}= \sqrt{2}v\frac{c_W m_t}{g_W}\left(c_W\left(\mathcal{C}_{rp}^{uW}\right)^* - s_W\left(\mathcal{C}_{rp}^{uB}\right)^*\right), (\kappa_R)_{pr} = \sqrt{2}v\frac{c_W m_t}{g_W}\left(c_W\mathcal{C}_{pr}^{uW} - s_W\mathcal{C}_{pr}^{uB}\right)$$
$$(X_L)_{pr} = v^2\left(\mathcal{C}_{pr}^{\phi q(1)} - \mathcal{C}_{pr}^{\phi q(3)}\right) \equiv v^2\mathcal{C}_{pr}^{\phi(-)}, (X_R)_{pr} = v^2\mathcal{C}_{pr}^{\phi u}$$
$$(\eta_L)_{pr} = \frac{3}{2}v^2\left(\mathcal{C}_{rp}^{u\phi}\right)^*, (\eta_R)_{pr} = \frac{3}{2}v^2\mathcal{C}_{pr}^{u\phi}$$

$$\mathcal{C}_i(\mu) = \left(1 + \frac{\gamma_{ii}}{16\pi^2}\log\left(\frac{\mu}{\Lambda}\right)\right)\mathcal{C}_i(\Lambda) + \sum_{i\neq j}\frac{\gamma_{ij}}{16\pi^2}\log\left(\frac{\mu}{\Lambda}\right)\mathcal{C}_j(\Lambda)$$

$$\begin{pmatrix} \mathcal{C}_{23}^{uB} \\ \mathcal{C}_{23}^{uW} \\ \mathcal{C}_{23}^{uG} \\ \mathcal{C}_{23}^{\phi u} \\ \mathcal{C}_{23}^{\phi q(1)} \\ \mathcal{C}_{23}^{\phi q(3)} \\ \mathcal{C}_{23}^{u\phi} \end{pmatrix}_{\mu_{\mathrm{EW}}} = \begin{pmatrix} 0.888659 & 0.001755 & -0.025980 & 0 & 0 & 0 & 0 \\ 0.000585 & 0.921365 & -0.027466 & 0 & 0 & 0 & 0 \\ -0.019485 & -0.061798 & 1.04996 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.886147 & 0.000081 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0.908814 & 0.050908 & 0 \\ 0 & 0 & 0 & 0 & 0.016969 & 0.955807 & 0 \\ 0.017356 & 0.007568 & -0.368257 & -0.010302 & 0 & -0.000030 & 0.977383 \end{pmatrix} \begin{pmatrix} \mathcal{C}_{23}^{uB} \\ \mathcal{C}_{23}^{uW} \\ \mathcal{C}_{23}^{uG} \\ \mathcal{C}_{23}^{\phi u} \\ \mathcal{C}_{23}^{\phi(1)} \\ \mathcal{C}_{23}^{\phi q(3)} \\ \mathcal{C}_{23}^{u\phi} \end{pmatrix}_{\mu_{1\mathrm{TeV}}}$$


$$\frac{d\vec{L}_i}{d\log\,\mu} = \frac{1}{16\pi^2}\sum_j \beta_{ij}\vec{L}_j = \frac{\alpha_s}{4\pi}\sum_j \beta_{ij}^s\vec{L}_j + \frac{\alpha_e}{4\pi}\sum_j \beta_{ij}^e\vec{L}_j$$

$$\vec{L}_i(\mu) = U_{ij}(\mu,\mu_0)\vec{L}_j = \left[U_{ij}^s(\mu,\mu_0) + U_{ij}^e(\mu,\mu_0)\right]\vec{L}_j(\mu_0)$$

$$-2\log\,\mathcal{L}\left(\mathcal{C}_i(\mu_{\text{strong gravity}})\right) = \chi^2\left(\mathcal{C}_i(\mu_{\text{strong gravity}})\right) = \chi^2_{\text{High}}\left(\mathcal{C}_i(\mu_{\text{strong gravity}})\right) + \chi^2_{\text{High}}\left(\mathcal{C}_i(\mu_{\text{strong gravity}})\right) + \chi^2_{\text{others}}\left(\mathcal{C}_i(\mu_{\text{strong gravity}})\right)$$
$$= \sum_{ij}\left(\mathcal{O}_i^{\text{theo}}(\mathcal{C}_i) - \mathcal{O}_i^{\text{exp}}\right)(\sigma^2)_{ij}^{-1}\left(\mathcal{O}_j^{\text{theo}}(\mathcal{C}_j) - \mathcal{O}_j^{\text{exp}}\right)$$

$$\mathcal{H}_{\text{eff}}^{b\to d_i\gamma} = -\frac{4G_F}{\sqrt{2}}V_{tb}V_{td_i}^*\sum_{i=7,8}\left(C_i(\mu)O_i + C_i'(\mu)O_i'\right)$$

$$O_7 = \frac{e}{16\pi^2}m_b\left(\bar{d}_j\sigma_{\mu\nu}P_R b\right)F^{\mu\nu}, O_7' = \frac{e}{16\pi^2}m_b\left(\bar{d}_j\sigma_{\mu\nu}P_L b\right)F^{\mu\nu}$$
$$O_8 = \frac{g_s}{16\pi^2}m_b\left(\bar{d}_j\sigma_{\mu\nu}T^a P_R b\right)G^{\mu\nu a}, O_8' = \frac{g_s}{16\pi^2}m_b\left(\bar{d}_j\sigma_{\mu\nu}T^a P_L b\right)G^{\mu\nu a}$$



$$C_i(\mu) = C_i^{\mathrm{SM}}(\mu) + C_i^{\mathrm{NP}}(\mu).$$

$$\begin{pmatrix} C_7 \\ C_8 \end{pmatrix}_{\mu_b} = \begin{pmatrix} 0.66301 & 0.09259 \\ 0.00326 & 0.69877 \end{pmatrix} \begin{pmatrix} C_7 \\ C_8 \end{pmatrix}_{\mu_{\mathrm{EW}}}.$$

$$C_7^{\mathrm{NP}}(\mu) = N_{d_i b}^{\mathrm{rad}} \frac{e}{m_b} L_{d_i b}^{d\gamma}(\mu), C_8^{\mathrm{NP}}(\mu) = N_{d_i b}^{\mathrm{rad}} \frac{e^2}{g_s m_b} L_{d_i b}^{dG}(\mu)$$

$$\left(N_{d_i b}^{\mathrm{rad}}\right)^{-1} = \frac{4G_F}{\sqrt{2}} \frac{\alpha_{\mathrm{em}}}{4\pi} V_{ts}^* V_{tb}$$

$$\mathcal{B}(B \to X_s\gamma) \times 10^4 = (3.40 \pm 0.17) - 8.25 C_7^{\mathrm{NP}}(\mu_{\mathrm{EW}}) - 2.10 C_8^{\mathrm{NP}}(\mu_{\mathrm{EW}})$$

$$\mathcal{B}\left(B_q \to V\gamma\right) = \tau_{B_q} \frac{G_F^2 \alpha_{\mathrm{em}^2} m_{B_q}^3 m_b^2}{32\pi^3} \left(1 - \frac{m_V^2}{m_{B_q}^2}\right)^3 |\lambda_t|^2 (|C_7(\mu_b)|^2 + |C_7'(\mu_b)|^2) T_1(0)$$

$$\mathcal{H}_{eff}^{b \to d_i \ell\ell} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{td_j}^* \left[\sum_{i=1}^{6} C_i(\mu) O_i(\mu) + \sum_{i=7,8,9,10,P,S} (C_i(\mu) O_i + C_i'(\mu) O_i')\right],$$

$$O_9 = \frac{e^2}{16\pi^2} (\bar{d}_j \gamma_\mu P_L b)(\bar{\ell}\gamma^\mu \ell), O_9' = \frac{e^2}{16\pi^2} (\bar{d}_j \gamma_\mu P_R b)(\bar{\ell}\gamma^\mu \ell),$$
$$O_{10} = \frac{e^2}{16\pi^2} (\bar{d}_j \gamma_\mu P_L b)(\bar{\ell}\gamma^\mu \gamma_5 \ell), O_{10}' = \frac{e^2}{16\pi^2} (\bar{d}_j \gamma_\mu P_R b)(\bar{\ell}\gamma^\mu \gamma_5 \ell),$$
$$O_S = \frac{e^2}{16\pi^2} m_b (\bar{d}_j P_R b)(\bar{\ell}\ell), O_S' = \frac{e^2}{16\pi^2} m_b (\bar{d}_j P_L b)(\bar{\ell}\ell).$$

$$C_{10}^{\mathrm{NP}}(\mu)=\lambda_1\left(-L^{V,LL}_{\substack{ed\\ \mu\mu d_i b}}+L^{V,LR}_{\substack{de\\ d_i b\mu\mu}}\right), C_{10}'^{\mathrm{NP}}(\mu)=\lambda_1\left(-L^{V,LR}_{\substack{ed\\ \mu\mu d_i b}}+L^{V,RR}_{\substack{ed\\ \mu\mu d_i b}}\right),$$

$$C_S^{\mathrm{NP}}(\mu)=\frac{\lambda_1}{m_b}\left(L^{S,RR}_{\substack{ed\\ \mu\mu d_i b}}+L^{S,RL*}_{\substack{ed\\ \mu\mu b d_i}}\right), C_S'^{\mathrm{NP}}(\mu)=\frac{\lambda_1}{m_b}\left(L^{S,RL}_{\substack{ed\\ \mu\mu d_i b}}+L^{S,RR*}_{\substack{ed\\ \mu\mu b d_i}}\right),$$

$$(\lambda_1)^{-1}=-\frac{4G_F}{\sqrt{2}}V_{tb}V_{ts}^*\frac{e^2}{16\pi^2}.$$

$$\begin{pmatrix}C_9\\ C_{10}\\ C_S\end{pmatrix}_{\mu_b}=\begin{pmatrix}0.99522 & 0.00716 & 0\\ 0.00716 & 1.0 & 0\\ 0 & 0 & 1.37433\end{pmatrix}\begin{pmatrix}C_9\\ C_{10}\\ C_S\end{pmatrix}_{\mu_{\mathrm{EW}}}$$

$$B\to K^{(*)}\mu^+\mu^-, B\to\phi\mu^+\mu^-$$

$$B_s^0\to\ell^+\ell^-, B^0\to\ell^+\ell^-, K_{L,S}\to\ell^+\ell^-$$

$$\mathcal{B}\left(B_q\to\mu^+\mu^-\right)=\tau_{B_q}f_{B_q}^2m_{B_q}^3\frac{G_F^2\alpha^2}{64\pi^3}\left|V_{tq}^*V_{tb}\right|^2\beta_\mu\left(m_{B_q}^2\right)\left[\frac{m_{B_q}^2}{m_b^2}|C_s-C_s'|^2\left(1-\frac{4m_\mu^2}{m_{B_q}^2}\right)\right.$$
$$\left.+\left|\frac{m_{B_q}}{m_b}\left(C_p-C_p'\right)+2\frac{m_\mu}{m_{B_q}}\left(C_{10}-C_{10}'\right)\right|^2\right]$$

$$\beta_\mu(q^2)=\sqrt{1-\frac{4m_\mu^2}{q^2}}$$

$$\langle 0|\bar{s}\gamma_\mu P_L b|B_q(p)\rangle=\frac{i}{2}f_{B_q}p_\mu$$

$$\mathcal{B}(B^0\to\mu^+\mu^-)^{\mathrm{Exp}}=(1.2^{+0.8}_{-0.7}\pm0.1)\times10^{-10}, \mathcal{B}(B_s^0\to\mu^+\mu^-)^{\mathrm{Exp}}=(3.83^{+0.38\ +0.24}_{-0.36\ -0.21})\times10^{-9}$$

$$\mathcal{B}(B^0\to\mu^+\mu^-)^{\mathrm{SM}}=(1.03\pm0.05)\times10^{-10}, \mathcal{B}(B_s^0\to\mu^+\mu^-)^{\mathrm{SM}}=(3.66\pm0.14)\times10^{-9}.$$

$$\mathcal{B}(K_L\to\mu^+\mu^-)_{\mathrm{SD}}=\tau_{K_L}f_K^2m_K\frac{G_F^2\alpha^2}{8\pi^3}m_\mu^2\beta_\mu(m_K^2)$$
$$\times\left\{\left[\Re\left(V_{ts}^*V_{td}\hat{P}\right)\right]^2+\left[\Im\left(V_{ts}^*V_{td}\hat{S}\right)\right]^2\right\}$$

$$\hat{P}(K)\equiv(C_{10}-C_{10}')+\frac{m_K^2}{2m_\mu}\frac{m_s}{m_d+m_s}(C_P-C_P')$$
$$\hat{S}(K)\equiv\beta_\mu(m_K^2)\frac{m_K^2}{2m_\mu}\frac{m_s}{m_d+m_s}(C_s-C_s')$$

$$\mathcal{B}(K_L\to\mu^+\mu^-)^{\mathrm{Exp}}=(6.84\pm0.11)\times10^{-9}$$

$$\mathcal{B}(K_L\to\mu^+\mu^-)_{\mathrm{SD}}^{\mathrm{SM}}=(0.79\pm0.12)\times10^{-9}$$

$$(K_L\to\mu^+\mu^-)_{\mathrm{SD}}<2.5\times10^{-9}$$

$$\mathcal{H}^{b\to d_i\nu\bar{\nu}} = \frac{4G_F}{\sqrt{2}} V_{tb}V_{td_i}^*(\mathcal{C}_L^\nu \mathcal{O}_L^\nu + \mathcal{C}_R^\nu \mathcal{O}_R^\nu)$$

Branching Ratio	SM values	Experimental values
$\mathcal{B}(B^+ \to K^+\nu\bar{\nu})$	$(5.22 \pm 0.15 \pm 0.28) \times 10^{-6}$	$\left(2.3 \pm 0.5^{+0.5}_{-0.4}\right) \times 10^{-5}$
$\mathcal{B}(B^0 \to K^{*0}\nu\bar{\nu})$	$(9.47 \pm 1.28 \pm 0.57) \times 10^{-6}$	$(3.8^{+2.9}_{-2.6}) \times 10^{-5}$ $< 1.8 \times 10^{-5}$
$\mathcal{B}(K^+ \to \pi^+\nu\bar{\nu})$	$(8.60 \pm 0.42) \times 10^{-11}$	$(1.06^{+0.40}_{-0.34} \pm 0.09) \times 10^{-10}$

$$\mathcal{O}_L^\nu = \frac{e^2}{16\pi^2}\left(\bar{d}_j\gamma_\mu P_L b\right)(\bar{\nu}\gamma^\mu P_L \nu), \qquad \mathcal{O}_R^\nu = \frac{e^2}{16\pi^2}\left(\bar{d}_j\gamma_\mu P_R b\right)(\bar{\nu}\gamma^\mu P_L \nu)$$

$$\frac{d\mathcal{B}(B^+ \to K^+\nu\bar{\nu})}{dq^2} = \tau_B \frac{G_F^2\alpha^2}{256\pi^5}\frac{\lambda(q^2, m_B^2, m_K^2)^{3/2}}{m_B^3}|V_{tb}V_{ts}^*|^2[f_+(q^2)]^2|C_L^\nu + C_R^\nu|^2$$


$$\lambda(a^2, b^2, c^2) = (a^2 - (b-c)^2)(a^2 - (b+c)^2), q^2$$

$$\begin{aligned}\frac{d\mathcal{B}(B \to K^*\nu\bar{\nu})}{dq^2} = \tau_B \frac{G_F^2\alpha^2}{128\pi^5}\frac{\lambda\left(q^2, m_B^2, m_{K^*}^2\right)^{1/2} q^2}{m_B^3}(m_B + m_{K^*})^2|V_{tb}V_{ts}^*|^2 \\ \times\left\{\left([A_1(q^2)]^2 + \frac{32m_B^2 m_{K^*}^2}{q^2(m_B + m_{K^*})^2}[A_{12}(q^2)]^2\right)|C_L^\nu - C_R^\nu|^2\right. \\ \left. + \frac{\lambda\left(q^2, m_B^2, m_{K^*}^2\right)}{(m_B + m_{K^*})^4}[V(q^2)]^2|C_L^\nu + C_R^\nu|^2\right\}\end{aligned}$$

$$\mathcal{B}(K^+ \to \pi\nu\bar{\nu}) = 3\lambda_2^{-2} J_V^{K^+}|C_L^\nu + C_R^\nu|^2$$

$$J_V^{K^+} = \frac{1}{\Gamma_{K^+}^{\text{Exp}}}\frac{1}{3\cdot 2^9\pi^3 m_{K^+}^3}\int ds\lambda^{3/2}\left(s, m_{K^+}^2, m_{\pi^+}^2\right)\left|f_+^{K^+}(s)\right|^2 = 0.23G_F^{-2}$$

$$(\lambda_2)^{-1} = \frac{4G_F}{\sqrt{2}}V_{td}V_{ts}^*\frac{e^2}{16\pi^2}, C_L^{\nu\text{SM}} = \lambda_2 \times (1.30 \times 10^{-10}\text{GeV}^{-2}), C_R^{\nu\text{SM}} = 0$$



$$\mathcal{H}^{d_i \to u_j \ell \nu_\ell} = \frac{4G_F}{\sqrt{2}} V_{u_j d_i} \left[(1 + C_{V_L}) O_{V_L} + C_{V_R} O_{V_R} \right]$$

$$O_{V_L} = (\bar{u}_j \gamma^\mu P_L d_i)(\bar{\ell} \gamma_\mu P_L \nu_\ell), O_{V_R} = (\bar{u}_j \gamma^\mu P_R d_i)(\bar{\ell} \gamma_\mu P_L \nu_\ell)$$


$$\begin{pmatrix} C_{V_L} \\ C_{V_R} \end{pmatrix}_{\mu_b} = \begin{pmatrix} 1.00716 & 0 \\ 0 & 1.00358 \end{pmatrix} \begin{pmatrix} C_{V_L} \\ C_{V_R} \end{pmatrix}_{\mu_{\text{EW}}}$$

$$d_i(u_i) \to u_j(d_j) \ell \nu_\ell$$

$$\frac{d\Gamma(P \to M \ell \nu_\ell)}{dq^2} = \frac{G_F^2 \left| V_{u_j d_i} \right|^2}{192\pi^3 m_P^3} q^2 \sqrt{\lambda(q^2, m_P^2, m_M^2)} \left(1 - \frac{m_\ell^2}{q^2}\right)^2 \left| 1 + C_{V_L} + C_{V_R} \right|^2 \times \left\{ \left(1 + \frac{m_\ell^2}{2q^2}\right) H_{V,0}^{s2} + \frac{3}{2} \frac{m_\ell^2}{q^2} H_{V,t}^{s2} \right\}$$

$$\mathcal{B}(P \to \ell \nu_\ell) = \frac{\tau_P}{8\pi} m_P m_\ell^2 f_P^2 G_F^2 \left(1 - \frac{m_\ell^2}{m_P^2}\right)^2 \left| V_{u_j d_i} (1 + C_{V_1}^\ell - C_{V_2}^\ell) \right|^2$$

$$\left| V_{u_j d_i} \right| \longrightarrow \left| V_{u_j d_i} (1 + C_{V_L} \pm C_{V_R}) \right|$$



$$\mathcal{L}_{tWb} = -\frac{g_2}{\sqrt{2}}\left(\bar{b}\gamma_\mu(V_L P_L + V_R P_R)tW^{\mu-} + \bar{b}\frac{i\sigma_{\mu\nu}q^\nu}{m_W}(g_L P_L + g_R P_R)tW^{\mu-}\right) + \text{ h.c.}$$


$$\Pi_{V_iV_j}(q^2) = \Pi^{\text{SM}}_{V_iV_j}(q^2) + \delta\Pi_{V_iV_j}(q^2)$$

$$\left(\Pi_{V_iV_j}\right)_{\mu\nu}(q^2) = -ig_{\mu\nu}(q^2 - m^2_{V_i}) - i\left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}\right)\Sigma^T_{V_iV_j}(q^2) - i\frac{q_\mu q_\nu}{q^2}\Sigma^L_{V_iV_j}(q^2)$$

$$S = \left(\frac{4s^2_W c^2_W}{\alpha_e}\right)\left(\left[\frac{\delta\Sigma^T_{ZZ}(m^2_Z) - \delta\Sigma^T_{ZZ}(0)}{m^2_Z}\right] - \frac{c^2_W - s^2_W}{c_W s_W}\frac{\delta\Sigma^T_{\gamma Z}(m^2_Z)}{m^2_Z} - \frac{\delta\Sigma^T_{\gamma\gamma}(m^2_Z)}{m^2_Z}\right)$$

$$T = \frac{1}{\alpha_e}\left(\frac{\delta\Sigma^T_{WW}(0)}{m^2_W} - \frac{\delta\Sigma^T_{ZZ}(0)}{m^2_Z}\right)$$


$$U = \frac{4s^2_W}{\alpha_e}\left(\left[\frac{\delta\Sigma^T_{WW}(m^2_W) - \delta\Sigma^T_{WW}(0)}{m^2_W}\right] - \frac{c_W}{s_W}\frac{\delta\Sigma^T_{Z\gamma}(m^2_Z)}{m^2_Z} - \frac{\delta\Sigma^T_{\gamma\gamma}(m^2_Z)}{m^2_Z}\right) - S$$



$$\mathcal{L}_{Zf\bar{f}} = \frac{g_2}{2\cos\,\theta_W}\sum_f \bar{f}\gamma^\mu\left(g^f_{Z,v} - g^f_{Z,a}\gamma^5\right)fZ_\mu,$$

$$g^f_{Z,a} \rightarrow g^{f,\text{SM}}_{Z,a} + \delta g^{f,\text{NP}}_{Z,a}$$
$$g^f_{Z,v} \rightarrow g^{f,\text{SM}}_{Z,v} + \delta g^{f,\text{NP}}_{Z,v}$$

$g^{f,\text{SM}}_{Z,v} = \left(I^3 - 2Q_f \sin^2\,\theta\right)$ and $g^{f,\text{SM}}_{Z,a} = I^3$, where I^3 and Q_f represent the component of the isospin and the charge of the fermion, respectively



$$\Gamma(Z \to f\bar{f}) = \frac{N_c^b}{48}\frac{\alpha}{s_W^2 c_W^2} m_Z \sqrt{1-\mu_f^2}\left(\left|g_{Z,a}^f\right|^2(1-\mu_f^2) + \left|g_{Z,v}^f\right|^2\left(1+\frac{\mu_f^2}{2}\right)\right)(1+\delta_f^0)(1+\delta_b)$$
$$(1+\delta_{\text{QCD}})(1+\delta_{\text{QED}})(1+\delta_\mu^f)$$


$$\mu_f^2 = \frac{4m_f^2}{m_Z^2}$$

Γ_Z(GeV)	(2.4961 ± 0.0010)	(2.4955 ± 0.0023)	$\sum_f \Gamma(Z \to ff)$
σ_{had} (nb)	(41.484)	(41.541 ± 0.037)	$\frac{12\pi}{m_Z^2}\frac{\Gamma(Z \to e^+e^-)\Gamma(Z \to q\bar{q})}{\Gamma_Z^2}$
A_e	(0.1475 ± 0.0010)	(0.1516 ± 0.0021)	$\frac{\Gamma(Z \to e_L^+e_L^-) - \Gamma(Z \to e_R^+e_R^-)}{\Gamma(Z \to e^+e^-)}$
A_μ	0.1472	0.142 ± 0.015	$\frac{\Gamma(Z \to \mu_L^+\mu_L^-) - \Gamma(Z \to \mu_R^+\mu_R^-)}{\Gamma(Z \to \mu^+\mu^-)}$
A_τ	0.1472	0.143 ± 0.004	$\frac{\Gamma(Z \to \tau_L^+\tau_L^-) - \Gamma(Z \to \tau_R^+\tau_R^-)}{\Gamma(Z \to \tau^+\tau^-)}$
A_b	0.935	0.923 ± 0.020	$\frac{\Gamma(Z \to b_L\bar{b}_L) - \Gamma(Z \to b_R\bar{b}_R)}{\Gamma(Z \to b\bar{b})}$

A_s	0.936	0.90 ± 0.09	$\frac{\Gamma(Z \to s_L \bar{s}_L) - \Gamma(Z \to s_R \bar{s}_R)}{\Gamma(Z \to s\bar{s})}$
A_c	0.667	0.670 ± 0.027	$\frac{\Gamma(Z \to c_L \bar{c}_L) - \Gamma(Z \to c_R \bar{c}_R)}{\Gamma(Z \to c\bar{c})}$
A_e^{FB}	$(1.606 \pm 0.006)\%$	$(1.45 \pm 0.25)\%$	$\frac{3}{4}A_e^2$
A_μ^{FB}	1.63%	$(1.69 \pm 0.13)\%$	$\frac{3}{4}A_e A_\mu$
A_τ^{FB}	1.63%	$(1.88 \pm 0.17)\%$	$\frac{3}{4}A_e A_\tau$
A_b^{FB}	$(10.34 \pm 0.07)\%$	$(9.92 \pm 0.16)\%$	$\frac{3}{4}A_e A_b$
A_s^{FB}	$(10.35 \pm 0.07)\%$	$(9.8 \pm 1.1)\%$	$\frac{3}{4}A_e A_s$
A_c^{FB}	$(7.35 \pm 0.02)\%$	$(7.07 \pm 0.35)\%$	$\frac{3}{4}A_e A_c$
R_e	20.736 ± 0.010	20.804 ± 0.050	$\frac{\Gamma(Z \to q\bar{q})}{\Gamma(Z \to e^+e^-)}$
R_μ	20.736 ± 0.010	20.784 ± 0.034	$\frac{\Gamma(Z \to q\bar{q})}{\Gamma(Z \to \mu^+\mu^-)}$
R_τ	20.781 ± 0.010	20.764 ± 0.045	$\frac{\Gamma(Z \to q\bar{q})}{\Gamma(Z \to \tau^+\tau^-)}$
R_b	0.21581	(0.21629 ± 0.00066)	$\frac{\Gamma(Z \to b\bar{b})}{\sum_q (Z \to q\bar{q})}$
R_c	0.1722	(0.1721 ± 0.0030)	$\frac{\Gamma(Z \to c\bar{c})}{\sum_q (Z \to q\bar{q})}$

Γ_W(GeV)	2.088	2.085 ± 0.042	$\sum_f \Gamma(W \to ff')$


R_{Wc}	0.50	0.49 ± 0.04	$\frac{\Gamma(W \to cs)}{\Gamma(W \to cs) + \Gamma(W \to ud)}$



$$\mathcal{L}_{Hbb} = C_L^H(\bar{b}P_L b) + C_R^H(\bar{b}P_R b)$$

$$C_{L(R)}^H = C_{L(R)}^{H,\mathrm{SM}} + C_{L(R)}^{H,\mathrm{NP}}$$

$$C_{L,R}^{H,\mathrm{SM}} = \frac{m_b}{v}$$



$$\Gamma(H \to b\bar{b}) = \frac{N_c}{16\pi m_H^2}\sqrt{m_H^2 - 2m_b^2}\left[m_H^2((C_L^H)^2 + (C_R^H)^2) - 2m_b^2(C_L^H + C_R^H)^2\right].$$

$$\Gamma(H \to b\bar{b}) = (1.961 \pm 0.923) \times 10^{-3}\mathrm{GeV}.$$

$$\delta\mathcal{L}_{h\gamma\gamma} = -\frac{\delta_{\gamma\gamma}}{2\Lambda^2}e^2 vhF_{\mu\nu}F^{\mu\nu}$$
$$\delta\mathcal{L}_{h\gamma Z} = -\frac{\delta_{\gamma Z}}{2\Lambda^2}e^2 vhF_{\mu\nu}Z^{\mu\nu}$$


$$\frac{\Gamma(h \to \gamma\gamma)}{\Gamma^{\mathrm{SM}}(h \to \gamma\gamma)} \simeq \left|1 - \frac{4\pi^2 v^2 \delta_{\gamma\gamma}}{\Lambda^2 I^{\gamma\gamma}}\right|^2$$
$$\frac{\Gamma(h \to \gamma Z)}{\Gamma^{\mathrm{SM}}(h \to \gamma Z)} \simeq \left|1 - \frac{4\pi^2 v^2 \delta_{\gamma Z}}{\Lambda^2 I^{\gamma Z}}\right|^2$$

$$\delta\mathcal{L}_{ggh} = -\frac{\delta_{gg}}{2\Lambda^2}g_s^2 vhG_{\mu\nu}^a G^{a\mu\nu}$$


$$\frac{\sigma(gg \to h)}{\sigma^{\mathrm{SM}}(gg \to h)} \simeq \frac{\Gamma(h \to gg)}{\Gamma^{\mathrm{SM}}(h \to gg)} \simeq \left|1 - \frac{8\pi^2 v^2 \delta_{gg}}{\Lambda^2 I^{gg}}\right|^2$$

$$\kappa_i^2 = \frac{\Gamma_i^{\mathrm{Total}}}{\Gamma_i^{SM}}$$

$$\kappa_\gamma = (1.02^{+0.08}_{-0.07}), \kappa_{Z\gamma} = 1.38^{+0.31}_{-0.37}, \kappa_g = 1.01^{+0.11}_{-0.09}$$



$$\begin{aligned}\mathcal{L}_{WWV} = i g_{WWV}\Big(& g_1^V\left(W_{\mu\nu}^+ W^{-\mu} W^{+\nu} - W_\mu^+ W_\nu^- W^{\mu\nu}\right) + \kappa_V W_\mu^+ W_\nu^- V^{\mu\nu} \\ & + \frac{\lambda_V}{M_W^2} W_{\lambda\mu}^+ W_\nu^{-\mu} V^{\nu\lambda} + i g_4^V W_\mu^+ W_\nu^- (\partial^\mu V^\nu + \partial^\nu V^\mu) - i g_5^V \epsilon^{\mu\nu\rho\sigma}\left(W_\mu^+ \overleftrightarrow{\partial}_\rho W_\nu^-\right) V_\rho \\ & + \tilde{\kappa}_V W_\mu^+ W_\nu^- \tilde{V}^{\mu\nu} + \frac{\tilde{\lambda}_V}{M_W^2} W_{\lambda\mu}^+ W^{-\mu} \tilde{V}^{\nu\lambda}\Big)\end{aligned}$$



$$g_i \in \{g_Z^1, \kappa_\gamma, \lambda_\gamma, \kappa_Z, \lambda_Z, g_Z^4, g_Z^5, \tilde{\kappa}_Z, \tilde{\lambda}_Z.\}$$

$$g_i = g_i^{\mathrm{SM}}(1 + \delta g_i)$$

$\hat{\mu}_t^{\mathrm{SM}}(m_Z^2) = -0.0224$, fourpoint vertex corrections yielding $\left\|\hat{\mu}_t^{\mathrm{SM}}(m_Z^2)\right\| = -0.0253$

$$\hat{\mu}_t^{\mathrm{Exp}}(m_Z^2) = -0.024^{+0.013+0.016}_{-0.009-0.011}$$

$$\mathcal{L}_{t\bar{t}g} = -g_s\left(\bar{t}\gamma^\mu G_\mu t + i\frac{\hat{d}_t}{2m_t}\bar{t}\sigma^{\mu\nu}\gamma_5 G_{\mu\nu}^a T^a t + \frac{\hat{\mu}_t}{2m_t}\bar{t}\sigma^{\mu\nu} G_{\mu\nu}^a T^a t\right)$$

$G_\mu \equiv G_\mu^a T^a$, where G_μ^a are the gluon fields

strength tensor is $G_{\mu\nu} = G_{\mu\nu}^a T^a$, where $G_{\mu\nu}^a \equiv \partial_\mu G_\nu^a - \partial_\nu G_\mu^a - g_s f^{abc} G_\mu^b G_\nu^c$, and f^{abc} are the structure constants

$\mu_t = g_s\hat{\mu}_t/m_t$ and $d_t = g_s\hat{d}_t/m_t$

$$\mu_t = \frac{g_s}{m_t}\left(\hat{\mu}_t^{\mathrm{SM}} + \hat{\mu}_t^{\mathrm{NP}}\right)$$
$$d_t = \frac{g_s}{m_t}\left(\hat{d}_t^{\mathrm{SM}} + \hat{d}_t^{\mathrm{NP}}\right)$$

$\mathcal{B}(t \to \gamma c)$	5×10^{-14}	4.16×10^{-5}(LH) 4.16×10^{-5}(RH) 1.51×10^{-5}
$\mathcal{B}(t \to \gamma u)$	4×10^{-16}	0.85×10^{-5}(LH) 1.2×10^{-5}(RH) 0.95×10^{-5}
$\mathcal{B}(t \to Zc)$	1×10^{-14}	1.3×10^{-4}(LH) 1.2×10^{-4}(RH)
$\mathcal{B}(t \to Zu)$	7×10^{-17}	6.2×10^{-5}(LH) 6.6×10^{-5}(RH)
$\mathcal{B}(t \to Hc)$	3×10^{-15}	3.4×10^{-4}
$\mathcal{B}(t \to Hu)$	2×10^{-17}	2.8×10^{-4}
$\mathcal{B}(t \to gc)$	5×10^{-12}	3.7×10^{-4}
$\mathcal{B}(t \to gu)$	4×10^{-14}	6.1×10^{-5}

$$\mathcal{B}(t \to u_j g) = C_F \frac{m_t}{16\pi\Gamma_t}\left[\left|\xi_L^{u_j t}\right|^2 + \left|\xi_R^{u_j t}\right|^2\right],$$
$$\mathcal{B}(t \to u_j \gamma) = e^2 \frac{m_t}{16\pi\Gamma_t}\left[\left|\lambda_L^{u_j t}\right|^2 + \left|\lambda_R^{u_j t}\right|^2\right],$$

$$\mathcal{B}(t \to u_j Z) = \frac{g_W^2 m_t}{128\pi c_W^2 \Gamma_t}\left(1 - \frac{M_Z^2}{m_t^2}\right)^2 \left[\left(2 + \frac{m_t^2}{M_Z^2}\right)\left(\left|X_L^{u_j t}\right|^2 + \left|X_R^{u_j t}\right|^2\right)\right.$$
$$+4\left(2 + \frac{M_Z^2}{m_t^2}\right)\left(\left|\kappa_L^{u_j t}\right|^2 + \left|\kappa_R^{u_j t}\right|^2\right)$$
$$\left.+6\left(X_L^{u_j t}\kappa_R^{*u_j t} + X_L^{*u_j t}\kappa_R^{u_j t} + X_R^{u_j t}\kappa_L^{*u_j t} + X_R^{*u_j t}\kappa_L^{u_j t}\right)\right]$$
$$\mathcal{B}(t \to u_j H) = \frac{m_t}{64\pi\Gamma_t}\left(1 - \frac{M_H^2}{m_t^2}\right)^2 \left(\left|\eta_L^{u_j t}\right|^2 + \left|\eta_R^{u_j t}\right|^2\right)$$

$$[|\xi_L^{ct}|^2 + |\xi_R^{ct}|^2] < 0.77 \times 10^{-4}, \quad [|\xi_L^{ut}|^2 + |\xi_R^{ut}|^2] < 1.27 \times 10^{-5}$$
$$[|\lambda_L^{ct}|^2 + |\lambda_R^{ct}|^2] < 1.75 \times 10^{-4}, \quad [|\lambda_L^{ut}|^2 + |\lambda_R^{ut}|^2] < 4.0 \times 10^{-5}$$
$$[|X_L^{ct}|^2 + |X_R^{ct}|^2] < 2.71 \times 10^{-4}, \quad [|X_L^{ut}|^2 + |X_R^{ut}|^2] < 1.38 \times 10^{-4}$$
$$[|\kappa_L^{ct}|^2 + |\kappa_R^{ct}|^2] < 1.65 \times 10^{-4}, \quad [|\kappa_L^{ut}|^2 + |\kappa_R^{ut}|^2] < 0.84 \times 10^{-4}$$
$$[|\eta_L^{ct}|^2 + |\eta_R^{ct}|^2] < 3.08 \times 10^{-3}, \quad [|\eta_L^{ut}|^2 + |\eta_R^{ut}|^2] < 2.54 \times 10^{-3}$$

Scenario	Values $\times 10^2$	Scenario	Values $\times 10^2$
$\mathrm{Re}(\xi_L^{ct}), \mathrm{Im}(\xi_L^{ct})$	$(12.3 \pm 19.7), (8.5 \pm 23.9)$	$\mathrm{Re}(\xi_L^{ut}), \mathrm{Im}(\xi_L^{ut})$	$(6.6 \pm 25.1), (-10.9 \pm 22.0)$
$\mathrm{Re}(\xi_R^{ct}), \mathrm{Im}(\xi_R^{ct})$	$(0.07 \pm 1.71), (0.09 \pm 1.60)$	$\mathrm{Re}(\xi_R^{ut}), \mathrm{Im}(\xi_R^{ut})$	$(-0.33 \pm 1.40), (1.57 \pm 4.55)$
$\mathrm{Re}(\xi_L^{ct}), \mathrm{Re}(\xi_R^{ct})$	$(13.2 \pm 16.9), (0.37 \pm 0.97)$	$\mathrm{Re}(\xi_L^{ut}), \mathrm{Re}(\xi_R^{ut})$	$(-8.6 \pm 23.5), (-0.38 \pm 1.29)$
$\mathrm{Re}(\xi_L^{ct}), \mathrm{Im}(\xi_L^{ct}),$ $\mathrm{Re}(\xi_R^{ct}), \mathrm{Im}(\xi_R^{ct}).$	$(-12.3 \pm 19.5), (8.2 \pm 24.4),$ $(0.16 \pm 0.98), (0.01 \pm 1.83).$	$\mathrm{Re}(\xi_L^{ut}), \mathrm{Im}(\xi_L^{ut}),$ $\mathrm{Re}(\xi_R^{ut}), \mathrm{Im}(\xi_R^{ut}).$	$(-6.4 \pm 52.7), (-3.3 \pm 85.4),$ $(-0.19 \pm 2.05), (-0.40 \pm 5.03).$

Scenario	Values $\times 10^2$	Scenario	Values $\times 10^2$
$\mathrm{Re}(\lambda_L^{ct}), \mathrm{Im}(\lambda_L^{ct})$	$(4.8 \pm 75.8), (-3.9 \pm 78.2)$	$\mathrm{Re}(\lambda_L^{ut}), \mathrm{Im}(\lambda_L^{ut})$	$(1.0 \pm 89.0), (-0.2 \pm 90.0)$
$\mathrm{Re}(\lambda_R^{ct}), \mathrm{Im}(\lambda_R^{ct})$	$(0.0 \pm 0.22), (0.0 \pm 0.21)$	$\mathrm{Re}(\lambda_R^{ut}), \mathrm{Im}(\lambda_R^{ut})$	$(-0.04 \pm 1.72), (0.0 \pm 1.17)$
$\mathrm{Re}(\lambda_L^{ct}), \mathrm{Re}(\lambda_R^{ct})$	$(1.3 \pm 90.0), (0.17 \pm 0.21)$	$\mathrm{Re}(\lambda_L^{ut}), \mathrm{Re}(\lambda_R^{ut})$	$(1.1 \pm 89.4), (0.30 \pm 0.87)$
$\mathrm{Re}(\lambda_L^{ct}), \mathrm{Im}(\lambda_L^{ct}),$ $\mathrm{Re}(\lambda_R^{ct}), \mathrm{Im}(\lambda_R^{ct}).$	$(-2.7 \pm 76.4), (-7.7 \pm 64.6),$ $(0.0 \pm 0.22), (0.0 \pm 0.22).$	$\mathrm{Re}(\lambda_L^{ut}), \mathrm{Im}(\lambda_L^{ut}),$ $\mathrm{Re}(\lambda_R^{ut}), \mathrm{Im}(\lambda_R^{ut}).$	$(1.0 \pm 89.0), (0.0 \pm 90.0),$ $(-0.05 \pm 1.71), (0.0 \pm 1.17).$

Different scenarios:

1. Single-Chirality.

2. Dual-Chirality.

3. Real Coupling.

Scenario	Values × **10**	Scenario	Values × **10**
$\mathrm{Re}(\kappa_L^{ct}), \mathrm{Im}(\kappa_L^{ct})$	$(-0.20 \pm 2.38), (-1.92 \pm 2.84)$	$\mathrm{Re}(\kappa_L^{ut}), \mathrm{Im}(\kappa_L^{ut})$	$(0.29 \pm 5.05), (-2.89 \pm 2.12)$
$\mathrm{Re}(\kappa_R^{ct}), \mathrm{Im}(\kappa_R^{ct})$	$(0.16 \pm 0.41), (2.75 \pm 2.04)$	$\mathrm{Re}(\kappa_R^{ut}), \mathrm{Im}(\kappa_R^{ut})$	$(0.89 \pm 2.03), (2.04 \pm 2.13)$
$\mathrm{Re}(\kappa_L^{ct}), \mathrm{Re}(\kappa_R^{ct})$	$(-0.29 \pm 2.92), (0.03 \pm 0.36)$	$\mathrm{Re}(\kappa_L^{ut}), \mathrm{Re}(\kappa_R^{ut})$	$(0.29 \pm 0.99), (-0.04 \pm 0.15)$
$\mathrm{Re}(X_L^{ct}), \mathrm{Re}(X_R^{ct})$	$(-0.13 \pm 0.39), (3.1 \pm 3.52)$	$\mathrm{Re}(X_L^{ut}), \mathrm{Re}(X_R^{ut})$	$(0.10 \pm 0.27), (-0.92 \pm 2.63)$
$\mathrm{Re}(\kappa_L^{ct}), \mathrm{Im}(\kappa_L^{ct})$, $\mathrm{Re}(\kappa_R^{ct}), \mathrm{Im}(\kappa_R^{ct})$.	$(-1.44 \pm 4.12), (1.40 \pm 4.16)$, $(2.64 \pm 6.90), (-4.59 \pm 4.49)$	$\mathrm{Re}(\kappa_L^{ut}), \mathrm{Im}(\kappa_L^{ut})$, $\mathrm{Re}(\kappa_R^{ut}), \mathrm{Im}(\kappa_R^{ut})$.	$(-2.31 \pm 2.15), (0.77 \pm 2.93)$, $(1.16 \pm 0.51), (-2.75 \pm 0.49)$

Scenario	Values	Scenario	Values
$\mathrm{Re}(\eta_L^{ct}), \mathrm{Im}(\eta_L^{ct})$	$(-1.05 \pm 12.96), (1.1 \pm 288.9)$	$\mathrm{Re}(\eta_L^{ut}), \mathrm{Im}(\eta_L^{ut})$	$(0.13 \pm 5.01), (0.46 \pm 63.17)$
$\mathrm{Re}(\eta_R^{ct}), \mathrm{Im}(\eta_R^{ct})$	$(0.37 \pm 1.71), (3.09 \pm 7.57)$	$\mathrm{Re}(\eta_R^{ut}), \mathrm{Im}(\eta_R^{ut})$	$(-0.02 \pm 0.07), (0.02 \pm 1.63)$
$\mathrm{Re}(\eta_L^{ct}), \mathrm{Re}(\eta_R^{ct})$	$(-1.05 \pm 12.95), (0.01 \pm 0.17)$	$\mathrm{Re}(\eta_L^{ut}), \mathrm{Re}(\eta_R^{ut})$	$(1.49 \pm 5.27), (-0.02 \pm 0.07)$
$\mathrm{Re}(\eta_L^{ct}), \mathrm{Im}(\eta_L^{ct})$, $\mathrm{Re}(\xi_L^{ct}), \mathrm{Im}(\xi_L^{ct})$	$(-0.95 \pm 16.92), (0.96 \pm 46.78)$, $(-0.04 \pm 0.76), (0.06 \pm 0.53)$	$\mathrm{Re}(\eta_R^{ct}), \mathrm{Im}(\eta_R^{ct})$, $\mathrm{Re}(\xi_R^{ct}), \mathrm{Im}(\xi_R^{ct})$	(0.38 ± 1.59), (3.10 ± 6.98), $(-0.001 \pm 0.011), (0.0 \pm 0.0267)$
$\mathrm{Re}(\eta_L^{ut}), \mathrm{Im}(\eta_L^{ut})$, $\mathrm{Re}(\xi_L^{ut}), \mathrm{Im}(\xi_L^{ut})$	$(0.02 \pm 5.56), (-0.89 \pm 9.36)$, $(0.10 \pm 0.27), (-0.10 \pm 0.23)$	$\mathrm{Re}(\eta_R^{ut}), \mathrm{Im}(\eta_R^{ut})$, $\mathrm{Re}(\xi_R^{ut}), \mathrm{Im}(\xi_R^{ut})$	(0.06 ± 0.97), (0.01 ± 1.62), $(0.003 \pm 0.16), (0.09 \pm 0.07)$

Scenarios	Value $(10^7 \times \mathrm{GeV}^{-2})$	Scenarios	Value $(10^7 \times \mathrm{GeV}^{-2})$
$\mathrm{Re}(\mathcal{C}_{23}^{uG}), \mathrm{Im}(\mathcal{C}_{23}^{uG})$	$(0.23 \pm 6.51), (0.73 \pm 3.0)$	$\mathrm{Re}(\mathcal{C}_{13}^{uG}), \mathrm{Im}(\mathcal{C}_{13}^{uG})$	$(-0.70 \pm 2.90), (3.27 \pm 9.45)$
$\mathrm{Re}(\mathcal{C}_{23}^{uB}), \mathrm{Im}(\mathcal{C}_{23}^{uB})$	$(0.0 \pm 0.17), (0.0 \pm 0.29)$	$\mathrm{Re}(\mathcal{C}_{13}^{uB}), \mathrm{Im}(\mathcal{C}_{13}^{uB})$	$(0.03 \pm 0.54), (0.0 \pm 0.86)$
$\mathrm{Re}(\mathcal{C}_{23}^{uW}), \mathrm{Im}(\mathcal{C}_{23}^{uW})$	$(0.0 \pm 0.60), (0.01 \pm 0.30)$	$\mathrm{Re}(\mathcal{C}_{13}^{uW}), \mathrm{Im}(\mathcal{C}_{13}^{uW})$	$(-0.14 \pm 0.87), (0.0 \pm 1.14)$
$\mathrm{Re}\left(\mathcal{C}_{23}^{u\phi}\right), \mathrm{Im}\left(\mathcal{C}_{23}^{u\phi}\right)$	$(1.54 \pm 18.94), (0.01 \pm 568)$	$\mathrm{Re}\left(\mathcal{C}_{13}^{u\phi}\right), \mathrm{Im}\left(\mathcal{C}_{13}^{u\phi}\right)$	$(-5.84 \pm 8.48), (-11.2 \pm 38.4)$
$\mathrm{Re}(\mathcal{C}_{23}^{uB}), \mathrm{Re}(\mathcal{C}_{23}^{uW})$	$(-0.22 \pm 2.20), (0.32 \pm 4.00)$	$\mathrm{Re}(\mathcal{C}_{13}^{uB}), \mathrm{Re}(\mathcal{C}_{13}^{uW})$	$(-0.12 \pm 1.00), (-0.55 \pm 1.62)$

$$C_{32}^{\phi q(-)}(\mu_{\mathrm{EW}}) = (-0.58 \pm 2.02) \times 10^{-10}, C_{31}^{\phi q(-)}(\mu_{\mathrm{EW}}) = (0.27 \pm 4.95) \times 10^{-10}.$$

$$C_{32}^{\phi q(-)}(\mu_{\mathrm{EW}}) = (-2.20 \pm 6.46) \times 10^{-7}, C_{31}^{\phi q(-)}(\mu_{\mathrm{EW}}) = (1.70 \pm 4.45) \times 10^{-7}.$$

$$C^{\phi q(-)} \equiv C^{\phi q(1)} - C^{\phi q(3)}$$

Coupling (GeV^{-2})	μ_b	μ_t	$\mu = 1\mathrm{TeV}$	$\mu = 5\mathrm{TeV}$	$\mu = 10\mathrm{TeV}$
$C_{23}^{uG} \times 10^7$	(-0.76 ± 2.29)	(-0.66 ± 1.97)	(-0.63 ± 1.88)	(-0.61 ± 1.82)	(-0.60 ± 1.80)
$C_{23}^{uB} \times 10^8$	(0.23 ± 0.20)	(-0.10 ± 1.45)	(-0.24 ± 1.97)	(-0.37 ± 2.42)	(-0.42 ± 2.62)
$C_{23}^{uW} \times 10^8$	(-0.35 ± 0.91)	(-0.88 ± 2.48)	(-1.09 ± 3.10)	(-1.27 ± 3.63)	(-1.35 ± 3.85)
$C_{23}^{\phi q(1)} \times 10^{10}$	(-0.43 ± 1.86)	(-0.61 ± 2.05)	(-0.68 ± 2.11)	(-0.74 ± 2.17)	(-0.77 ± 2.19)
$C_{23}^{\phi q(3)} \times 10^{10}$	(-0.27 ± 1.87)	(0.59 ± 2.05)	(0.62 ± 2.09)	(0.65 ± 2.12)	(0.66 ± 2.13)
$C_{23}^{\phi u} \times 10^6$	(0.59 ± 7.16)	(0.79 ± 9.32)	(0.85 ± 10.2)	(0.92 ± 10.97)	(0.94 ± 11.29)
$C_{23}^{u\phi} \times 10^6$	(0.12 ± 1.78)	(0.18 ± 1.91)	(0.19 ± 1.91)	(0.20 ± 1.90)	(0.20 ± 1.89)
$C_{13}^{uG} \times 10^7$	(-1.16 ± 2.97)	(-1.00 ± 2.63)	(-0.96 ± 2.55)	(-0.93 ± 2.50)	(-0.91 ± 2.48)
$C_{13}^{uB} \times 10^8$	(0.35 ± 3.31)	(-0.15 ± 5.79)	(-0.36 ± 6.82)	(-0.55 ± 7.72)	(-0.63 ± 8.09)
$C_{13}^{uW} \times 10^7$	(-0.06 ± 0.59)	(-0.15 ± 0.92)	(-0.18 ± 1.05)	(-0.21 ± 1.17)	(-0.22 ± 1.21)
$C_{13}^{\phi q(1)} \times 10^{10}$	(0.20 ± 4.57)	(0.28 ± 5.02)	(0.32 ± 5.18)	(0.34 ± 5.32)	(0.36 ± 5.37)
$C_{13}^{\phi q(3)} \times 10^{10}$	(-0.23 ± 4.59)	(-0.28 ± 5.00)	(-0.29 ± 5.12)	(-0.50 ± 5.20)	(-0.31 ± 5.23)
$C_{13}^{\phi u} \times 10^6$	(-1.19 ± 3.45)	(-1.56 ± 4.49)	(-1.70 ± 4.92)	(-1.83 ± 5.29)	(-1.88 ± 5.44)
$C_{13}^{u\phi} \times 10^7$	(-3.17 ± 8.29)	(-2.97 ± 7.70)	(-2.86 ± 7.40)	(-2.76 ± 7.10)	(-2.71 ± 6.97)

$$\left|\mathcal{C}_{13}^{uG}\right| < 0.57 \times 10^{-7}, \left|\mathcal{C}_{23}^{uG}\right| < 1.4 \times 10^{-7},$$
$$\left|\mathcal{C}_{13}^{uW} + \mathcal{C}_{13}^{uB}\right| < 0.123 \times 10^{-6}, \left|\mathcal{C}_{23}^{uW} + \mathcal{C}_{23}^{uB}\right| < 0.235 \times 10^{-6},$$
$$\left|(\mathcal{C}_{31}^{uW})^* + (\mathcal{C}_{31}^{uB})^*\right| < 0.103 \times 10^{-6}, \left|(\mathcal{C}_{32}^{uW})^* + (\mathcal{C}_{32}^{uB})^*\right| < 0.227 \times 10^{-6},$$
$$\left|\mathcal{C}_{13}^{u\phi}\right| < 0.68 \times 10^{-6}, \left|\mathcal{C}_{23}^{u\phi}\right| < 0.78 \times 10^{-6},$$

$$d\Gamma_{A\to b,q^*(c,d)} = \frac{|\mathcal{M}|^2}{512\pi^3 m_A^3 q^2}\lambda^{1/2}(m_A^2, m_b^2, q^2)\lambda^{1/2}(q^2, m_c^2, m_d^2)dq^2 d\cos\,\theta$$


$$\left.\frac{d\Gamma(q^2)}{dq^2}\right|_{t\to u_j\gamma^*(\ell\ell)} = \frac{e^2v^2m_t^3}{96\pi^3q^2}\left(1-\frac{q^2}{m_t^2}\right)^2\left(2+\frac{q^2}{m_t^2}\right)\left|c_W\mathcal{C}_{pq}^{uB} + s_W\mathcal{C}_{pq}^{uW}\right|^2$$

$$\left.\frac{d\Gamma(q^2)}{dq^2}\right|_{t\to u_jZ^*(\ell\ell)} = \frac{g_Z^2m_t^3v^2\left(g_{Z_L}^2+g_{Z_R}^2\right)}{1536\pi^3(q^2-M_Z^2)^2}\left(1-\frac{q^2}{m_t^2}\right)^2\left\{8q^2\left(2+\frac{q^2}{m_t^2}\right)\left|c_W\mathcal{C}_{pq}^{uW} - s_W\mathcal{C}_{pq}^{uB}\right|^2\right.$$
$$\left. + g_Z^2v^2\left(1+\frac{2q^2}{m_t^2}\right)\left|\mathcal{C}_{pq}^{\phi q(-)}\right|^2\right\}$$

$$\left.\frac{d\Gamma(q^2)}{dq^2}\right|_{t\to u_jH^*(\ell\ell)} = \frac{9m_tm_\ell^2v^2q^2}{1024\pi^3(q^2-M_H^2)^2}\left(1-\frac{q^2}{m_t^2}\right)^2\left|\mathcal{C}_{pq}^{u\phi}\right|^2$$

$$g_Z = \frac{e}{s_Wc_W}, g_{Z_L} = I_W^{(3)} - Q_fs_W^2, g_{Z_R} = -Q_fs_W^2$$

$$\mathcal{B}(t\to ce^+e^-) = 8.48\times10^{-15}, \qquad \mathcal{B}(t\to c\mu^+\mu^-) = 9.55\times10^{-15}$$
$$\mathcal{B}(t\to ue^+e^-) = 6.81\times10^{-17}, \qquad \mathcal{B}(t\to u\mu^+\mu^-) = 7.68\times10^{-17}$$
$$\mathcal{B}(t\to cv\bar{v}) = 2.99\times10^{-14}, \qquad \mathcal{B}(t\to uv\bar{v}) = 2.40\times10^{-16}$$



$$i\mathcal{M}(t\to u_j+\gamma_\pm) = i\bar{u}(p_f)\Gamma_{fi,\gamma}^{\mu}(q^2)u(p_i)\varepsilon_{\pm,\mu}^*(q)$$

$$\Gamma_{fi,\gamma}^{\mu}(q^2) = i\sigma^{\mu\nu}q_\nu\left(f_{fi,\gamma}^LP_L + f_{fi,\gamma}^RP_R\right)$$

$$f_{fi,\gamma}^L = \sum_\alpha V_{t\alpha}^*V_{u_j\alpha}m_{u_j}\mathcal{F}_\alpha^L + \sum_{l_{\rm NP}}\mathcal{C}_{l_{\rm NP}}\mathcal{K}_{l_{\rm NP}}^L$$
$$f_{fi,\gamma}^R = \sum_\alpha V_{t\alpha}^*V_{u_j\alpha}m_t\mathcal{F}_\alpha^R + \sum_{l_{\rm NP}}\mathcal{C}_{l_{\rm NP}}\mathcal{K}_{l_{\rm NP}}^R$$

$$i\mathcal{M}(\bar{t}\to\bar{u}_j+\gamma_\mp) = i\mathcal{M}^{CP}(t\to u_j+\gamma_\pm) = i\bar{v}(p_i)\bar{\Gamma}_{if,\gamma}^{\mu}(q^2)v(p_f)\varepsilon_{\mp,\mu}^*(q)$$

$$\bar{\Gamma}^{\mu}_{if,\gamma}(q^2) = i\sigma^{\mu\nu}q_\nu\left(\bar{f}^L_{if,\gamma}P_L + \bar{f}^R_{if,\gamma}P_R\right)$$

$$\bar{f}^L_{if,\gamma} = \sum_{\alpha} V_{t\alpha}V^*_{u_j\alpha}m_t\mathcal{F}^R_\alpha + \sum_{l_{\rm NP}} \mathcal{C}^*_{l_{\rm NP}}\mathcal{K}^R_{l_{\rm NP}}$$

$$\bar{f}^R_{if,\gamma} = \sum_{\alpha} V_{t\alpha}V^*_{u_j\alpha}m_{u_j}\mathcal{F}^L_\alpha + \sum_{l_{\rm NP}} \mathcal{C}^*_{l_{\rm NP}}\mathcal{K}^L_{l_{\rm NP}}$$

$$\Delta_{\rm CP,+} = \frac{\Gamma(t\to u_j\gamma_+) - \Gamma(\bar{t}\to\bar{u}_j\gamma_-)}{\Gamma(t\to u_j\gamma) + \Gamma(\bar{t}\to\bar{u}_j\gamma)} = \frac{|f^L_{fi}|^2 - |\bar{f}^R_{if}|^2}{|f^L_{fi}|^2 + |f^R_{fi}|^2 + |\bar{f}^L_{if}|^2 + |\bar{f}^R_{if}|^2}$$

$$\Delta_{\rm CP,-} = \frac{\Gamma(t\to u_j\gamma_-) - \Gamma(\bar{t}\to\bar{u}_j\gamma_+)}{\Gamma(t\to u_j\gamma) + \Gamma(\bar{t}\to\bar{u}_j\gamma)} = \frac{|f^R_{fi}|^2 - |\bar{f}^L_{if}|^2}{|f^L_{fi}|^2 + |f^R_{fi}|^2 + |\bar{f}^L_{if}|^2 + |\bar{f}^R_{if}|^2}$$

$$\Delta_{\rm CP} = \frac{\Gamma(t\to u_j\gamma_+) - \Gamma(\bar{t}\to\bar{u}_j\gamma_-) + \Gamma(t\to u_j\gamma_-) - \Gamma(\bar{t}\to\bar{u}_j\gamma_+)}{\Gamma(t\to u_j\gamma) + \Gamma(\bar{t}\to\bar{u}_j\gamma)}$$
$$= \frac{\Gamma(t\to u_j\gamma) - \Gamma(\bar{t}\to\bar{u}_j\gamma)}{\Gamma(t\to u_j\gamma) + \Gamma(\bar{t}\to\bar{u}_j\gamma)} = \frac{|f^L_{fi}|^2 + |f^R_{fi}|^2 - |\bar{f}^R_{if}|^2 - |\bar{f}^L_{if}|^2}{|f^L_{fi}|^2 + |f^R_{fi}|^2 + |\bar{f}^L_{if}|^2 + |\bar{f}^R_{if}|^2}$$


$$\Delta_{\rm CP,+} = -\frac{m^2_{u_j}\sum_{\alpha,\beta}\mathcal{J}_{\alpha\beta}\,{\rm Im}\left(\mathcal{F}^L_\alpha\mathcal{F}^{L*}_\beta\right) + 2m_{u_j}\sum_{\alpha,l_{\rm NP}}{\rm Im}\left(V_{t\alpha}V^*_{u_j\alpha}\mathcal{C}_{l_{\rm NP}}\right){\rm Im}\left(\mathcal{K}^L_{l_{\rm NP}}\mathcal{F}^{L*}_\alpha\right) + \sum_{l_{\rm NP}\neq l'_{\rm NP}}{\rm Im}\left(\mathcal{C}_{l_{\rm NP}}\mathcal{C}^*_{l'_{\rm NP}}\right){\rm Im}\left(\mathcal{K}^L_{l_{\rm NP}}\mathcal{K}^{L*}_{l'_{\rm NP}}\right)}{\mathcal{D}}$$

$$\Delta_{\rm CP,-} = -\frac{m^2_{t}\sum_{\alpha,\beta}\mathcal{J}_{\alpha\beta}\,{\rm Im}\left(\mathcal{F}^R_\alpha\mathcal{F}^{R*}_\beta\right) + 2m_{t}\sum_{\alpha,l_{\rm NP}}{\rm Im}\left(V_{t\alpha}V^*_{u_j\alpha}\mathcal{C}_{l_{\rm NP}}\right){\rm Im}\left(\mathcal{K}^R_{l_{\rm NP}}\mathcal{F}^{R*}_\alpha\right) + \sum_{l_{\rm NP}\neq l'_{\rm NP}}{\rm Im}\left(\mathcal{C}_{l_{\rm NP}}\mathcal{C}^*_{l'_{\rm NP}}\right){\rm Im}\left(\mathcal{K}^R_{l_{\rm NP}}\mathcal{K}^{R*}_{l'_{\rm NP}}\right)}{\mathcal{D}}$$

$$\mathcal{D} = \sum_{a,\beta}\mathcal{R}_{\alpha\beta}\left\{m^2_t{\rm Re}\left(\mathcal{F}^R_\alpha\mathcal{F}^{R*}_\beta\right) + m^2_{u_j}{\rm Re}\left(\mathcal{F}^L_\alpha\mathcal{F}^{L*}_\beta\right)\right\}$$
$$+2\sum_{\alpha,l_{\rm NP}}{\rm Re}\left(V_{t\alpha}V^*_{u_j\alpha}\mathcal{C}_{l_{\rm NP}}\right)\left\{m_t{\rm Re}\left(\mathcal{F}^{R*}_\alpha\mathcal{K}^R_{l_{\rm NP}}\right) + m_{u_j}{\rm Re}\left(\mathcal{F}^{L*}_\alpha\mathcal{K}^L_{l_{\rm NP}}\right)\right\}$$
$$+\sum_{l_{\rm NP},l'_{\rm NP}}{\rm Re}\left(\mathcal{C}_{l_{\rm NP}}\mathcal{C}^*_{l'_{\rm NP}}\right)\left\{{\rm Re}\left(\mathcal{K}^L_{l_{\rm NP}}\mathcal{K}^{L*}_{l'_{\rm NP}}\right) + {\rm Re}\left(\mathcal{K}^R_{l_{\rm NP}}\mathcal{K}^{R*}_{l'_{\rm NP}}\right)\right\}$$

$$\mathcal{J}_{\alpha\beta} = {\rm Im}\left(V^*_{t\alpha}V_{u_j\alpha}V_{t\beta}V^*_{u_j\beta}\right), \qquad \mathcal{R}_{\alpha\beta} = {\rm Re}\left(V^*_{t\alpha}V_{u_j\alpha}V_{t\beta}V^*_{u_j\beta}\right).$$

Decay Channel	$\Delta_{CP,+}$	$\Delta_{CP,-}$
$t \to c\gamma$	$(1.05 \pm 4.93) \times 10^{-4}$	$(-0.11 \pm 6.06) \times 10^{-3}$
$t \to u\gamma$	$(-2.31 \pm 6.56) \times 10^{-9}$	$(-0.26 \pm 2.24) \times 10^{-4}$
$t \to cg$	$(-0.05 \pm 1.81) \times 10^{-4}$	$(-0.95 \pm 6.30) \times 10^{-3}$
$t \to ug$	$(-1.18 \pm 3.52) \times 10^{-8}$	$(-0.32 \pm 1.19) \times 10^{-4}$



$$\mathcal{L}_{\text{eff}} \supset +d_q(\mu)\frac{i}{2}\bar{q}\sigma^{\mu\nu}\gamma_5 q F_{\mu\nu} + \tilde{d}_q(\mu)\frac{i}{2}g_s(\mu)\bar{q}\sigma^{\mu\nu}T^a\gamma_5 q G^a_{\mu\nu} + w(\mu)\frac{1}{3}G^a_{\mu\sigma}G^{b,\sigma}_{\nu}\tilde{G}^{c,\mu\nu}$$

$$\tilde{G}^{a\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\alpha\beta}G^a_{\alpha\beta}$$

$$\frac{d_n}{e} = \left(g_T^d\frac{d_u}{e} + g_T^u\frac{d_d}{e} + g_T^s\frac{d_s}{e}\right) + (1 \pm 0.5) \times 1.1\left(\tilde{d}_d + 0.5\tilde{d}_u\right) + (22 \pm 10) \times 10^{-3}\text{GeV}.w$$

$$g_T^u = 0.784(28)(10), g_T^d = -0.204(11)(10), g_T^s = -0.0027(16)$$

$$\left|\frac{d_n}{e}\right| < 1.8 \times 10^{-26}\ \text{cm}$$

$$d_q(\mu) = 2\text{Im}\left[L_{ii}^{q\gamma}\right](\mu), \tilde{d}_q(\mu) = \frac{2}{g_s(\mu)}\text{Im}\left[L_{ii}^{qG}\right](\mu)$$

$$\frac{d_n}{e}(\mu) = -\frac{3.94 \times 10^{-14}}{e}\left(g_T^d\text{Im}\left[L_{11}^{u\gamma}\right](\mu_{\text{had}}) + g_T^u\text{Im}\left[L_{11}^{d\gamma}\right](\mu_{\text{had}}) + g_T^s\text{Im}\left[L_{22}^{d\gamma}\right](\mu_{\text{had}})\right) - \frac{4.3 \times 10^{-14}}{g_s(\mu)}(1 \pm 0.5)\left\{\text{Im}\left[L_{11}^{dG}\right](\mu_{\text{had}}) + 0.5\text{Im}\left[L_{11}^{uG}\right](\mu_{\text{had}})\right\}$$

$$\begin{pmatrix} L^{u\gamma} \\ L^{uG} \\ L^{d\gamma} \\ L^{dG} \end{pmatrix}_{\mu_{\text{had}}} = \begin{pmatrix} 0.87 & -0.064 & 0 & 0 \\ -0.05 & 1.23 & 0 & 0 \\ 0 & 0 & 0.88 & 0.03 \\ 0 & 0 & 0.024 & 1.23 \end{pmatrix} \begin{pmatrix} L^{u\gamma} \\ L^{uG} \\ L^{d\gamma} \\ L^{dG} \end{pmatrix}_{\mu_{\text{strong gravity}}}$$

$$\begin{aligned}
\frac{d_d(\mu_{\text{had}})}{e} &= 2.3 \times 10^{-8}\tilde{d}_c(\mu_t) + 1.0 \times 10^{-4}\text{GeV} \cdot w(\mu_t), \\
\frac{d_u(\mu_{\text{had}})}{e} &= -2.1 \times 10^{-8}\tilde{d}_c(\mu_t) - 9.1 \times 10^{-5}\text{GeV} \cdot w(\mu_t), \\
\tilde{d}_d(\mu_{\text{had}}) &= 1.8 \times 10^{-6}\tilde{d}_c(\mu_t) + 7.0 \times 10^{-4}\text{GeV} \cdot w(\mu_t), \\
\tilde{d}_u(\mu_{\text{had}}) &= 8.2 \times 10^{-7}\tilde{d}_c(\mu_t) + 3.1 \times 10^{-4}\text{GeV} \cdot w(\mu_t), \\
w(\mu_{\text{had}}) &= 1.7 \times 10^{-2}\text{GeV}^{-1}\tilde{d}_c(\mu_t) + 0.41w(\mu_t).
\end{aligned}$$

$$\left|\frac{d_n}{e}\right| = \left|(3.7 \pm 1.7) \times 10^{-4}.\tilde{d}_c(\mu_t) + (9.3 \pm 4.1) \times 10^{-3}\text{GeV}.w(\mu_t)\right|$$

$$\begin{aligned}
&|\text{Im}(\xi_L^{ut}\xi_R^{tu*})| < 2.20 \times 10^{-8}, \quad |\text{Im}(\xi_L^{ct}\xi_R^{tc*})| < 0.89 \times 10^{-4}, \\
&|\text{Im}(\lambda_L^{ut}\lambda_R^{tu*})| < 2.19 \times 10^{-7}, \quad |\text{Im}(\lambda_L^{ct}\lambda_R^{tc*})| < 2.39 \times 10^{-4}, \\
&|\text{Im}(\eta_L^{ut}\eta_R^{tu*})| < 1.61 \times 10^{-7}, \quad |\text{Im}(\eta_L^{ct}\eta_R^{tc*})| < 1.76 \times 10^{-4}, \\
&|\text{Im}(\kappa_L^{ut}\kappa_R^{tu*})| < 3.63 \times 10^{-8}, \quad |\text{Im}(\kappa_L^{ct}\kappa_R^{tc*})| < 3.97 \times 10^{-5}, \\
&|\text{Im}(X_L^{ut}X_R^{tu*})| < 2.54 \times 10^{-7}, \quad |\text{Im}(X_L^{ct}X_R^{tc*})| < 2.79 \times 10^{-4}.
\end{aligned}$$

$$\begin{aligned}
&|\text{Im}(\xi_L^{ut})\text{Re}(\xi_R^{tu*})|, |\text{Re}(\xi_L^{ut})\text{Im}(\xi_R^{tu*})| < 2.20 \times 10^{-8}, \\
&|\text{Im}(\xi_L^{ct})\text{Re}(\xi_R^{tc*})|, |\text{Re}(\xi_L^{ct})\text{Im}(\xi_R^{tc*})| < 0.89 \times 10^{-4}, \\
&|\text{Im}(\lambda_L^{ut})\text{Re}(\lambda_R^{tu*})|, |\text{Re}(\lambda_L^{ut})\text{Im}(\lambda_R^{tu*})| < 2.19 \times 10^{-7}, \\
&|\text{Im}(\lambda_L^{ct})\text{Re}(\lambda_R^{tc*})|, |\text{Re}(\lambda_L^{ct})\text{Im}(\lambda_R^{tc*})| < 2.39 \times 10^{-4}, \\
&|\text{Im}(\eta_L^{ut})\text{Re}(\eta_R^{tu*})|, |\text{Re}(\eta_L^{ut})\text{Im}(\eta_R^{tu*})| < 1.61 \times 10^{-7}, \\
&|\text{Im}(\eta_L^{ct})\text{Re}(\eta_R^{tc*})|, |\text{Re}(\eta_L^{ct})\text{Im}(\eta_R^{tc*})| < 1.76 \times 10^{-4}, \\
&|\text{Im}(\kappa_L^{ut})\text{Re}(\kappa_R^{tu*})|, |\text{Re}(\kappa_L^{ut})\text{Im}(\kappa_R^{tu*})| < 3.63 \times 10^{-8}, \\
&|\text{Im}(\kappa_L^{ct})\text{Re}(\kappa_R^{tc*})|, |\text{Re}(\kappa_L^{ct})\text{Im}(\kappa_R^{tc*})| < 3.97 \times 10^{-5}, \\
&|\text{Im}(X_L^{ut})\text{Re}(X_R^{tu*})|, |\text{Re}(X_L^{ut})\text{Im}(X_R^{tu*})| < 2.54 \times 10^{-7}, \\
&|\text{Im}(X_L^{ct})\text{Re}(X_R^{tc*})|, |\text{Re}(X_L^{ct})\text{Im}(X_R^{tc*})| < 2.79 \times 10^{-4}.
\end{aligned}$$

$$\left|\mathrm{Re}\left(\mathcal{C}_{31}^{uG}\right)\mathrm{Im}\left(\mathcal{C}_{13}^{uG}\right)\right|,\left|\mathrm{Im}\left(\mathcal{C}_{31}^{uG}\right)\mathrm{Re}\left(\mathcal{C}_{13}^{uG}\right)\right| < 0.91\times10^{-17}$$
$$\left|\mathrm{Re}\left(\mathcal{C}_{32}^{uG}\right)\mathrm{Im}\left(\mathcal{C}_{23}^{uG}\right)\right|,\left|\mathrm{Im}\left(\mathcal{C}_{32}^{uG}\right)\mathrm{Re}\left(\mathcal{C}_{23}^{uG}\right)\right| < 3.68\times10^{-14}$$
$$\begin{bmatrix}\left|\mathrm{Re}\left(\mathcal{C}_{31}^{uW}\right)\mathrm{Im}\left(\mathcal{C}_{13}^{uW}\right)\right|,\left|\mathrm{Im}\left(\mathcal{C}_{31}^{uW}\right)\mathrm{Re}\left(\mathcal{C}_{13}^{uW}\right)\right|,\\ \left|\mathrm{Re}\left(\mathcal{C}_{31}^{uB}\right)\mathrm{Im}\left(\mathcal{C}_{13}^{uB}\right)\right|,\left|\mathrm{Im}\left(\mathcal{C}_{31}^{uB}\right)\mathrm{Re}\left(\mathcal{C}_{13}^{uB}\right)\right|\end{bmatrix} < 1.15\times10^{-17}$$
$$\begin{bmatrix}\left|\mathrm{Re}\left(\mathcal{C}_{32}^{uW}\right)\mathrm{Im}\left(\mathcal{C}_{23}^{uW}\right)\right|,\left|\mathrm{Im}\left(\mathcal{C}_{32}^{uW}\right)\mathrm{Re}\left(\mathcal{C}_{23}^{uW}\right)\right|,\\ \left|\mathrm{Re}\left(\mathcal{C}_{32}^{uB}\right)\mathrm{Im}\left(\mathcal{C}_{23}^{uB}\right)\right|,\left|\mathrm{Im}\left(\mathcal{C}_{32}^{uB}\right)\mathrm{Re}\left(\mathcal{C}_{23}^{uB}\right)\right|\end{bmatrix} < 1.26\times10^{-14}$$
$$\left|\mathrm{Re}\left(\mathcal{C}_{31}^{u\phi}\right)\mathrm{Im}\left(\mathcal{C}_{13}^{u\phi}\right)\right|,\left|\mathrm{Im}\left(\mathcal{C}_{31}^{u\phi}\right)\mathrm{Re}\left(\mathcal{C}_{13}^{u\phi}\right)\right| < 1.95\times10^{-17}$$
$$\left|\mathrm{Re}\left(\mathcal{C}_{32}^{u\phi}\right)\mathrm{Im}\left(\mathcal{C}_{23}^{u\phi}\right)\right|,\left|\mathrm{Im}\left(\mathcal{C}_{32}^{u\phi}\right)\mathrm{Re}\left(\mathcal{C}_{23}^{u\phi}\right)\right| < 2.14\times10^{-14}$$

$$[\beta^{uG}]_{rs} = -\frac{1}{36}(81g_2^2+19g_1^2+204g_s^2)[\mathcal{C}^{uG}]_{rs}+6g_2g_s[\mathcal{C}^{uW}]_{rs}+\frac{10}{3}g_1g_s[\mathcal{C}^{uB}]_{rs}$$
$$+2\left[\Gamma_u\Gamma_u^\dagger\mathcal{C}^{uG}\right]_{rs}-2\left[\Gamma_d\Gamma_d^\dagger\mathcal{C}^{uG}\right]_{rs}+\left[\mathcal{C}_{uG}\Gamma_u^\dagger\Gamma_u\right]+\gamma_H^{(Y)}[\mathcal{C}^{uG}]_{rs}+\left[\gamma_q^{(Y)}\mathcal{C}^{uG}\right]_{rs}+\left[\mathcal{C}^{uG}\gamma_u^{(Y)}\right]_{rs},$$
$$[\beta^{uW}]_{rs} = -\frac{1}{36}(33g_2^2+19g_1^2-96g_s^2)[\mathcal{C}^{uW}]_{rs}+\gamma_H^{(Y)}[\mathcal{C}^{uW}]_{rs}+\left[\gamma_q^{(Y)}\mathcal{C}^{uW}\right]_{rs}+\left[\mathcal{C}^{uW}\gamma_u^{(Y)}\right]_{rs}$$
$$+\left[\mathcal{C}^{uW}\Gamma_u^\dagger\Gamma_u\right]_{rs}+2\left[\Gamma_d\Gamma_d^\dagger\mathcal{C}^{uW}\right]_{rs}-\frac{1}{6}g_1g_2[\mathcal{C}^{uB}]_{rs}+\frac{8}{3}g_2g_s[\mathcal{C}^{uG}]_{rs}$$
$$[\beta^{uB}]_{rs} = -\frac{1}{36}(81g_2^2-313g_1^2-96g_s^2)[\mathcal{C}^{uB}]_{rs}+\gamma_H^{(Y)}[\mathcal{C}^{uB}]_{rs}+\left[\gamma_q^{(Y)}\mathcal{C}^{uB}\right]_{rs}+\left[\mathcal{C}^{uB}\gamma_u^{(Y)}\right]_{rs}$$
$$-\frac{1}{2}g_1g_2[\mathcal{C}^{uW}]_{rs}-2\left[\Gamma_d\Gamma_d^\dagger\mathcal{C}^{uB}\right]_{rs}+2\left[\Gamma_u\Gamma_u^\dagger\mathcal{C}^{uB}\right]_{rs}+\left[\mathcal{C}^{uB}\Gamma_u^\dagger\Gamma_u\right]_{rs}+\frac{40}{9}g_1g_s[\mathcal{C}^{uG}]_{rs}$$
$$\left[\beta^{\phi q(1)}\right]_{rs} = \frac{1}{3}g_1^2\left[\mathcal{C}^{\phi q(1)}\right]_{rs}+\frac{3}{2}\Big(\left[\Gamma_d\Gamma_d^\dagger\mathcal{C}^{\phi q(1)}\right]_{rs}+\left[\Gamma_u\Gamma_u^\dagger\mathcal{C}^{\phi q(1)}\right]_{rs}+\left[\mathcal{C}^{\phi q(1)}\Gamma_d\Gamma_d^\dagger\right]_{rs}+\left[\mathcal{C}^{\phi q(1)}\Gamma_u\Gamma_u^\dagger\right]_{rs}$$
$$+3\left[\Gamma_d\Gamma_d^\dagger\mathcal{C}^{\phi q(3)}\right]_{rs}-3\left[\Gamma_u\Gamma_u^\dagger\mathcal{C}^{\phi q(3)}\right]_{rs}+3\left[\mathcal{C}^{\phi q(3)}\Gamma_d\Gamma_d^\dagger\right]_{rs}-3\left[\mathcal{C}^{\phi q(3)}\Gamma_u\Gamma_u^\dagger\right]_{rs}\Big)$$
$$+2\gamma_H^{(Y)}\left[\mathcal{C}^{\phi q(1)}\right]_{rs}+\left[\gamma_q^{(Y)}\mathcal{C}^{\phi q(1)}\right]_{rs}+\left[\mathcal{C}^{\phi q(1)}\gamma_q^{(Y)}\right]_{rs}$$
$$\left[\beta^{\phi q(3)}\right]_{rs} = -\frac{17}{3}g_2^2\left[\mathcal{C}^{\phi q(3)}\right]_{rs}+\frac{1}{2}\Big(3\left[\Gamma_d\Gamma_d^\dagger\mathcal{C}^{\phi q(1)}\right]_{rs}-3\left[\Gamma_u\Gamma_u^\dagger\mathcal{C}^{\phi q(1)}\right]_{rs}+3\left[\mathcal{C}^{\phi q(1)}\Gamma_d\Gamma_d^\dagger\right]_{rs}$$
$$-3\left[\mathcal{C}^{\phi q(1)}\Gamma_u\Gamma_u^\dagger\right]_{rs}+\left[\Gamma_d\Gamma_d^\dagger\mathcal{C}^{\phi q(3)}\right]+\left[\Gamma_u\Gamma_u^\dagger\mathcal{C}^{\phi q(3)}\right]_{rs}+\left[\mathcal{C}^{\phi q(3)}\Gamma_d\Gamma_d^\dagger\right]_{rs}+\left[\mathcal{C}^{\phi q(3)}\Gamma_u\Gamma_u^\dagger\right]_{rs}\Big)$$
$$+2\gamma_H^{(Y)}\left[\mathcal{C}^{\phi q(3)}\right]_{rs}+\left[\gamma_q^{(Y)}\mathcal{C}^{\phi q(3)}\right]_{rs}+\left[\mathcal{C}^{\phi q(3)}\gamma_q^{(Y)}\right]_{rs}$$
$$\left[\beta^{u\phi}\right]_{rs} = -\left(\frac{35}{12}g_1^2+\frac{27}{4}g_2^2+8g_s^2\right)\left[\mathcal{C}^{u\phi}\right]_{rs}-g_1(5g_1^2-3g_2^2)[\mathcal{C}^{uB}]_{rs}+g_2(5g_1^2-9g_2^2)[\mathcal{C}^{uW}]_{rs}$$
$$-(3g_2^2-g_1^2)\left[\Gamma_u\mathcal{C}^{\phi u}\right]_{rs}+4g_1^2\left[\mathcal{C}^{\phi q(1)}\Gamma_u\right]_{rs}-4g_1^2\left[\mathcal{C}^{\phi q(3)}\Gamma_u\right]_{rs}-5g_1\left[\mathcal{C}^{uB}\Gamma_u^\dagger\Gamma_u+\Gamma_u\Gamma_u^\dagger\mathcal{C}^{uB}\right]_{rs}$$
$$-3g_2\left[\mathcal{C}^{uW}\Gamma_u^\dagger\Gamma_u-\Gamma_u\Gamma_u^\dagger\mathcal{C}^{uW}\right]_{rs}-12g_2\left[\Gamma_d\Gamma_d^\dagger\mathcal{C}^{uW}\right]_{rs}-16g_s\left[\mathcal{C}^{uG}\Gamma_u^\dagger\Gamma_u+\Gamma_u\Gamma_u^\dagger\mathcal{C}^{uG}\right]_{rs}$$
$$+\lambda\left(12\left[\mathcal{C}^{u\phi}\right]_{rs}-2\left[\mathcal{C}^{\phi q(1)}\Gamma_u\right]_{rs}+6\left[\mathcal{C}^{\phi q(3)}\Gamma_u\right]+2\left[\Gamma_u\mathcal{C}^{\phi u}\right]_{rs}\right)$$
$$-2\left[\mathcal{C}^{\phi q(1)}\Gamma_u\Gamma_u^\dagger\Gamma_u\right]_{rs}+6\left[\mathcal{C}^{\phi q(3)}\Gamma_d\Gamma_d^\dagger\Gamma_u\right]_{rs}+2\left[\Gamma_u\Gamma_u^\dagger\Gamma_u\mathcal{C}^{\phi u}\right]_{rs}+4\left[\mathcal{C}^{u\phi}\Gamma_u^\dagger\Gamma_u\right]_{rs}$$
$$+5\left[\Gamma_u\Gamma_u^\dagger\mathcal{C}^{u\phi}\right]_{rs}-2\left[\Gamma_d\Gamma_d^\dagger\mathcal{C}^{u\phi}\right]_{rs}+3\gamma_H^{(Y)}\left[\mathcal{C}^{u\phi}\right]_{rs}+\left[\gamma_q^{(Y)}\mathcal{C}^{u\phi}\right]_{rs}+\left[\mathcal{C}^{u\phi}\gamma_u^{(Y)}\right]_{rs}$$

$$\gamma_H^{(Y)} = \frac{3}{4}(g_1^2 + 3g_2^2) + \mathrm{Tr}\left(3\Gamma_u\Gamma_u^\dagger + 3\Gamma_d\Gamma_d^\dagger + \Gamma_e\Gamma_e^\dagger\right)$$

$$\left[\gamma_q^{(Y)}\right]_{rs} = \frac{1}{2}\left[\Gamma_u\Gamma_u^\dagger + \Gamma_d\Gamma_d^\dagger\right]$$

$$\left[\gamma_u^{(Y)}\right]_{rs} = \left[\Gamma_u^\dagger\Gamma_u\right]_{rs}$$

$$\left[\gamma_d^{(Y)}\right]_{rs} = \left[\Gamma_d^\dagger\Gamma_d\right]_{rs}$$

$$\left[\gamma_e^{(Y)}\right]_{rs} = \left[\Gamma_e^\dagger\Gamma_e\right]_{rs}$$

$$g_1 = 0.3576, g_2 = 0.6515, g_s = 1.220, \lambda = 0.2813,$$

$$m_u = 1.27 \times 10^{-3}, m_d = 2.7 \times 10^{-3}, m_s = 5.51 \times 10^{-2}, m_c = 0.635, m_b = 2.85,$$

$$\mu_{\mathrm{had}} = 2\mathrm{GeV}, \mu_b = 4.18\mathrm{GeV}, \mu_{\mathrm{EW}} = 91.1876\mathrm{GeV}, \mu_t = 172.69\mathrm{GeV}, \mu_\Lambda = \mathcal{O}$$

$$L_{d_ib}^{d\gamma}(\mu) = -i\frac{em_bg_2^2}{16\pi^2M_W^2}\left[\frac{V_{tb}V_{u_is}^*(1+x_t)\lambda_R^{u_it}(\mu) - 2V_{ts}^*V_{u_ib}(1-x_t)\lambda_R^{tu_i*}(\mu)}{8(x_t-1)} + \frac{V_{tb}V_{u_is}^*\lambda_R^{u_it}(\mu)}{4}\log\left(\frac{\mu^2}{M_W^2}\right) - \frac{V_{tb}V_{u_is}^*x_t^2\lambda_R^{u_it}(\mu)}{4(x_t-1)^2}\log x_t\right]$$

$$L_{d_ib}^{dG}(\mu) = -i\frac{g_sm_bg_2^2}{16\pi^2M_W^2}\left[\frac{V_{tb}V_{u_is}^*(1+x_t)\xi_R^{u_it}(\mu) - 2V_{ts}^*V_{u_ib}(1-x_t)\xi_R^{tu_i*}(\mu)}{8(x_t-1)} + \frac{V_{tb}V_{u_is}^*\xi_R^{u_it}(\mu)}{4}\log\left(\frac{\mu^2}{M_W^2}\right) - \frac{V_{tb}V_{u_is}^*x_t^2\xi_R^{u_it}(\mu)}{4(x_t-1)^2}\log x_t\right]$$

$$L_{\mu\mu d_ib}^{V,LL}(\mu) = -\frac{1}{2}\frac{e^2g_2^2}{16\pi^2M_W^2}\left[\frac{V_{tb}V_{u_is}^*\lambda_R^{u_it}(\mu) - V_{ts}^*V_{ub}\lambda_R^{tu_i*}(\mu)}{2(x_t-1)}\log x_t\right] - \frac{g_2^4g_{Z_L}}{16\pi^2M_Z^2\cos^2\theta_W}\left[\left(V_{tb}V_{u_is}^*X_L^{u_it}(\mu) + V_{ts}^*V_{u_ib}X_L^{tu_i*}(\mu)\right)\left\{\frac{2x_t+3}{16} + \frac{x_t}{8}\log\left(\frac{\mu^2}{m_t^2}\right)\right\}\right]$$

$$g_{ed}^{V,LR}(\mu) = -\frac{1}{2}\frac{e^2g_2^2}{16\pi^2M_W^2}\left[\frac{V_{tb}V_{u_is}^*\lambda_R^{u_it}(\mu) - V_{ts}^*V_{ub}\lambda_R^{tu_i*}(\mu)}{2(x_t-1)}\log x_t\right] - \frac{}{16\pi^2M_Z^2\cos^2\theta_W}\left[\left(V_{tb}V_{u_is}^*X_L^{u_it}(\mu) + V_{ts}^*V_{u_ib}X_L^{tu_i*}(\mu)\right)\left\{\frac{2x_t+3}{16} + \frac{x_t}{8}\log\left(\frac{\mu^2}{m_t^2}\right)\right\}\right]$$

$$L^{S,RR}(\mu) = L_{\mu\mu d_ib}^{ed\ S,RL*}(\mu) = \frac{1}{2d}\frac{m_bm_\mu m_tg_2^2}{16\pi^2vM_H^2M_W^2}\left[\frac{V_{tb}V_{u_is}^*\eta_R^{u_it}(\mu) + 2V_{ts}^*V_{u_ib}\eta_R^{tu_i*}(\mu)}{4\sqrt{2}}\log\left(\frac{\mu^2}{m_t^2}\right) + \frac{3V_{tb}V_{u_is}^*\eta_R^{u_it}}{4\sqrt{2}}(x_t-1)^2\log x_t + \frac{V_{tb}V_{u_is}^*(x_t-7)\eta_R^{u_it} + 4(x_t-1)V_{ts}^*V_{u_ib}\eta_R^{tu_i*}}{8\sqrt{2}(x_t-1)}\right]$$

(B.1e)

$$L^{V,LL}(\mu) = -\frac{g_2^4g_{Z_L}}{16\pi^2M_Z^2\cos^2\theta_W}\left[\left(V_{tb}V_{u_is}^*X_L^{u_it}(\mu) + V_{ts}^*V_{u_ib}X_L^{tu_i*}(\mu)\right)\left\{\frac{2x_t+3}{16} + \frac{x_t}{8}\log\left(\frac{\mu^2}{m_t^2}\right)\right\}\right]$$

$$L_{vebu_i}^{V,LL} = \frac{g_2^2}{16\sqrt{2}\pi^2 M_W^2 G_F}\left|\frac{V_{tb}}{V_{u_ib}}\right|\left[g_s^2\left(\frac{5}{6}+\log\frac{\mu^2}{m_t^2}\right)\xi_R^{u_it}\right.$$
$$+e^2\left(\frac{3}{8(x_t-1)}\log x_t+\frac{15x_t-34}{96}+\frac{3x_t-14}{16}\log\frac{\mu^2}{m_t^2}\right)\lambda_R^{u_it}$$
$$+\frac{3g_2^2}{82\sqrt{2}}\frac{vm_t}{M_W^2}\left(\frac{1}{2}-\frac{\log x_H}{(x_H-1)(x_t-1)}-\frac{1}{3}\frac{x_t(x_t-4)}{(x_H-x_t)(x_t-1)}\log\frac{x_H}{x_t}+\frac{1}{3}\log\frac{\mu^2}{M_H^2}\right)\eta_R^{u_it}$$
$$+\frac{m_t}{4\sqrt{2}v}\left(1-\frac{x_H}{(x_H-x_t)}\log\frac{x_H}{x_t}+\log\frac{\mu^2}{m_t^2}\right)\eta_R^{u_it}$$
$$-\frac{g_2^2}{16}(2X_L^{u_it}-3\kappa_R^{u_it})\left(\frac{5}{2}+\frac{3\log x_t}{(x_t-1)(x_Z-1)}+\frac{3x_Z^2}{(x_Z-1)(x_Z-x_t)}\log\frac{x_t}{x_Z}+3\log\frac{\mu^2}{m_t^2}\right)$$
$$-\frac{(g_1^2-3g_2^2)}{16}\left(\frac{3x_Z\left(x_tX_L^{u_it}-2x_Z\kappa_R^{u_it}\right)}{x_t(x_t-x_Z)}\log\frac{x_t}{x_Z}+\left(\frac{x_t-x_Z}{x_Z}+\frac{x_t-3x_Z}{x_Z}\log\frac{\mu^2}{m_t^2}\right)X_L^{u_it}\right.$$
$$\left.2\frac{x_t+x_Z}{x_t}\left(1+3\log\frac{\mu^2}{m_t^2}\right)\kappa_R^{u_it}\right)+\frac{(g_1^2+3g_2^2)}{2}\left(-6\kappa_R^{u_it}+\left(2\frac{x_t}{x_Z}\left(1+\log\frac{\mu^2}{m_t^2}\right)+3\right)X_L^{u_it}\right)\Bigg]$$

$$L_{vebu_i}^{V,LR} = \frac{g_2^2}{16\sqrt{2}\pi^2 M_W^2 G_F}\left|\frac{V_{tb}}{V_{u_ib}}\right|\left[\frac{m_b}{m_t}\left(\frac{g_s^2}{12}\left(-1+6\log\frac{\mu^2}{m_t^2}\right)\xi_L^{u_it}+\frac{e^2}{48}\left(1-6\log\frac{\mu^2}{m_t^2}\right)\lambda_L^{u_it}\right.\right.$$
$$\left.+\frac{g_1^2}{96}\left(\frac{6\left(x_tX_R^{u_it}-2x_Z\kappa_L^{u_it}\right)}{(x_t-x_Z)}\log\frac{x_t}{x_Z}+\frac{x_tX_R^{u_it}-2x_Z\kappa_L^{u_it}}{x_Z}+\frac{2\left(x_tX_R^{u_it}+6x_Z\kappa_L^{u_it}\right)}{x_Z}\log\frac{\mu^2}{m_t^2}\right)\right)$$
$$+\frac{m_b}{16\sqrt{2}v}\left(\frac{2x_t}{(x_H-x_t)}\log\frac{x_H}{x_t}-\left(1+2\log\frac{\mu^2}{M_H^2}\right)\right)\eta_L^{u_it}+\frac{g_1^2}{12}\left(\frac{x_t-x_Z}{x_Z}X_R^{u_it}+2\frac{x_t+x_Z}{x_t}\kappa_L^{u_it}\right.$$
$$\left.+3\frac{x_Z\left(x_tX_R^{u_it}-2x_Z\kappa_L^{u_it}\right)}{x_t(x_t-x_Z)}\log\frac{x_t}{x_Z}+\left(\frac{x_t-3x_Z}{x_Z}X_R^{u_it}+6\frac{x_Z+x_t}{x_t}\kappa_L^{u_it}\right)\log\frac{\mu^2}{m_t^2}\right)\Bigg]$$

$$C_{V_L} = \frac{V_{u_ib}}{8\pi^2}\left[\left(C_Fg_s^2\xi_R^{u_it}+\frac{e^2}{3}\lambda_R^{u_it}\right)\left(1+\frac{3x_t+1}{4(x_t-1)}\log\left(-\frac{\mu^2}{m_t^2}\right)-\frac{1}{(x_t-1)}\log\left(-\frac{\mu^2}{M_W^2}\right)\right)\right.$$
$$+\frac{(g_1^2+3g_2^2)}{192\pi^2}\left(\left(X_L^{u_it}+3\kappa_R^{u_it}\right)\log\frac{\mu^2}{M_Z^2}-\frac{(x_t+2)(x_t-x_Z-1)}{(x_t-1)^2}X_L^{u_it}-\frac{4x_t^2-x_t(3x_Z+4)-x_Z}{x_t(x_t-1)}\kappa_R^{u_it}\right.$$
$$\left.\left(+\frac{(2x_t^2-4x_tx_Z-2x_Z+x_t-3)}{(x_t-1)^3}X_L^{u_it}+4\frac{x_t-2x_Z-1}{(x_t-1)^2}\kappa_R^{u_it}\right)\log(-x_Z)\right)$$
$$+e^2\left(\frac{3x_t+5}{8(x_t-1)}-\frac{x_t^2+6x_t+1}{4(x_t-1)^2}\log(-x_t)\right)\lambda_R^{u_it}+\frac{g_2^2}{4\sqrt{2}}\frac{m_tv}{M_W^2}\left(\frac{1}{x_t-1}+\frac{x_t-x_H+1}{(x_t-1)^2}\log x_H\right.$$
$$\left.+\frac{M_W^2}{m_t}\frac{(x_t-x_H)(x_t-1)}{(x_t-1)^2}\log\frac{x_H}{x_H-x_t}\right)\eta_R^{u_it}+\frac{g_2^2}{8}\left(\frac{\left(2x_t^2X_L^{u_it}+(3x_t^2+x_t(5-6x_Z)+2x_Z)\kappa_R^{u_it}\right.}{x_t(x_t-1)}\right.$$
$$-2\frac{(x_t^2-x_t-x_Z+2)X_L^{u_it}+(1+x_t^2+x_t(6-4x_Z)-4x_Z+2x_Z^2)\kappa_R^{u_it}}{(x_t-1)^2}\log x_Z$$
$$\left.+\frac{2(x_t-x_Z)\left(X_L^{u_it}x_t(2-x_t+x_t^2)+(x_t+x_t^3-3x_t^2(x_Z-2)-x_Z)\kappa_R^{u_it}\right.}{x_t^2(x_t-1)^2}\log\frac{x_Z}{x_Z-x_t}\right)\Bigg]$$

$$\Sigma_{\gamma\gamma}^{T}(q^2) = -\frac{e^2}{8\pi^2}\left(\left|\lambda_L^{u_i t}\right|^2 + \left|\lambda_R^{u_i t}\right|^2\right) q^2$$

$$\Sigma_{\gamma Z}^{T}(q^2) = -\frac{e g_2}{64\pi^2 c_W}\Big[\left(X_L^{tu_i *}\lambda_L^{u_i t} + X_R^{tu_i *}\lambda_R^{u_i t} + X_R^{u_i t}\lambda_L^{tu_i *} + X_L^{u_i t}\lambda_R^{tu_i *}\right)\left(1 + 2\log\frac{\mu^2}{m_t^2}\right)$$
$$+8\mathrm{Re}\left(\kappa_L^{tu_i *}\lambda_L^{u_i t} + \kappa_R^{tu_i *}\lambda_R^{u_i t}\right)\Big] q^2$$

$$\Sigma_{ZZ}^{T}(q^2) = -\frac{g_2^2 m_t^2}{64\pi^2 c_W^2}\left(\left|X_L^{u_i t}\right|^2 + \left|X_R^{u_i t}\right|^2\right)\left(1 + 2\log\frac{\mu^2}{m_t^2}\right)$$
$$+\frac{g_2^2}{576\pi^2 c_W^2}\Big[4\left(\left|X_L^{u_i t}\right|^2 + \left|X_R^{u_i t}\right|^2\right)\left(1 + 3\log\frac{\mu^2}{m_t^2}\right) - 72\left(\left|\kappa_L^{u_i t}\right|^2 + \left|\kappa_R^{u_i t}\right|^2\right)$$
$$-18\mathrm{Re}\left(X_L^{tu_i *}\kappa_L^{u_i t} + X_R^{tu_i *}\kappa_L^{u_i t} + X_L^{tu_i *}\kappa_R^{u_i t} + X_R^{tu_i *}\kappa_R^{u_i t}\right)\left(1 + 2\log\frac{\mu^2}{m_t^2}\right)\Big] q^2$$

$$\delta g_{Z,v}^{b,\mathrm{NP}} = -\delta g_{Z,a}^{b,\mathrm{NP}} = \frac{g_2^2}{32\pi^2}\sum_{u_i=u,c} V_{tb}V_{u_i b}^{*}\Big\{\mathrm{Re}\left(X_L^{u_i t}\right)\Big[\frac{2\left(5x_Z + 2 - x_t(1-x_Z)\right)}{x_Z}$$
$$+2x_t\log\frac{\mu^2}{m_t^2} + \frac{2\left(2 + 3x_Z - 2x_t(1+x_Z)\right)}{(x_t-1)x_Z}\log x_t + \frac{2(x_t - x_Z)(x_t - 3x_Z - 2)}{x_Z^2}\log\left(\frac{x_t}{x_t - x_z}\right)\Big]$$
$$-8i\mathrm{Im}\left(\kappa_R^{u_i t}\right)\left[1 - \log x_t - \frac{x_t - x_Z}{x_Z}\log\left(\frac{x_t}{x_t - x_Z}\right)\right]\Big\}$$

$$V_{tb}V_{u_i b}^{*} \Leftrightarrow V_{ts}V_{u_i s}^{*}$$

$$C_L^{H,\mathrm{NP}} = \sum_{u_i=u,c}\frac{g_2^2 m_b m_t}{128\sqrt{2}\pi^2 M_H^2}V_{tb}V_{u_i b}^{*}\Bigg(-8(x_H+1)\eta_R^{u_i t} + \frac{8 - x_H(x_t-3) - 8x_t}{x_t - 1}\eta_R^{tu_i *}$$
$$-2x_H\left(2\eta_R^{u_i t} + \eta_R^{tu_i *}\right)\log\frac{\mu^2}{m_t^2} + 2\left(4\eta_R^{u_i t} - \frac{4x_t + x_H - 4}{(x_t-1)^2}\eta_R^{tu_i *}\right)\log x_t$$
$$-\frac{4(x_H + x_t)\left((2+x_H)\eta_R^{u_i t} + 2\eta_R^{tu_i *}\right)}{x_H}\log\frac{x_t}{x_t + x_H}\Bigg)$$

$$C_R^{H,\mathrm{NP}} = \sum_{u_i=u,c}\frac{g_2^2 m_b m_t}{128\sqrt{2}\pi^2 M_H^2}V_{tb}V_{u_i b}^{*}\Bigg(\left(8 + x_H\frac{x_t - 3}{x_t - 1}\right)\eta_R^{u_i t} + 8(1+x_H)\eta_R^{tu_i *}$$
$$+2x_H\left(\eta_R^{u_i t} + 2\eta_R^{tu_i *}\right)\log\frac{\mu^2}{m_t^2} - 2\left(4\eta_R^{tu_i *} - \frac{4x_t + x_H - 4}{(x_t-1)^2}\eta_R^{u_i t}\right)\log x_t$$
$$+\frac{4(x_H + x_t)\left((2+x_H)\eta_R^{tu_i *} + 2\eta_R^{u_i t}\right)}{x_H}\log\frac{x_t}{x_t + x_H}\Bigg)$$

$$\delta_{\gamma\gamma} = \frac{1}{12\sqrt{2}\pi^2 m_t v}\Bigg(2i\mathrm{Im}\left(\eta_L^{tu_i *}\lambda_L^{u_i t} + \eta_R^{tu_i *}\lambda_R^{u_i t}\right)\left(\log\frac{\mu^2}{m_t^2} + \log\frac{m_t^2}{m_t^2 - M_H^2}\right)$$
$$-2i\mathrm{Im}\left(\eta_L^{u_i t}\lambda_L^{tu_i *} + \eta_R^{u_i t}\lambda_R^{tu_i *}\right) + \frac{3\sqrt{2}m_t}{v}\left(\left|\lambda_L^{u_i t}\right|^2 + \left|\lambda_R^{u_i t}\right|^2\right)\Bigg)$$

$$\delta_{\gamma Z}=\frac{1}{192\pi^2v^2m_t}\Bigg[\Bigg(-2\sqrt{2}iv\frac{5g_1^2-3g_2^2}{g_1g_2}\mathrm{Im}\left(\eta_L^{tu_i*}\lambda_L^{u_it}+\eta_R^{tu_i*}\lambda_R^{u_it}\right)+6i\frac{m_t}{g_1g_2}\mathrm{Im}\left(X_R^{tu_i*}\lambda_L^{u_it}+X_L^{tu_i*}\lambda_R^{u_it}\right)$$
$$+4\sqrt{2}e^2g_1g_2\left(2i\mathrm{Im}\left(\eta_L^{tu_i*}\kappa_L^{u_it}+\eta_R^{tu_i*}\kappa_R^{u_it}\right)-\left(\eta_L^{u_it}\kappa_L^{tu_i*}+\eta_R^{u_it}\kappa_R^{tu_i*}\right)\right)\Bigg)\left(1+\log\frac{\mu^2}{m_t^2}\right)$$
$$+\frac{24m_t}{g_1g_2}\mathrm{Re}\left(\kappa_L^{tu_i*}\lambda_L^{u_it}+\kappa_R^{tu_i*}\lambda_R^{u_it}\right)+\Bigg(2\sqrt{2}iv\frac{(5g_1^2-3g_2^2)}{g_1g_2}\mathrm{Im}\left(\eta_L^{tu_i*}\lambda_L^{u_it}+\eta_R^{tu_i*}\lambda_R^{u_it}\right)$$
$$-4\sqrt{2}e^2g_1g_2\left(2i\mathrm{Im}\left(\eta_L^{tu_i*}\kappa_L^{u_it}+\eta_R^{tu_i*}\kappa_R^{u_it}\right)-\left(\eta_L^{u_it}\kappa_L^{tu_i*}+\eta_R^{u_it}\kappa_R^{tu_i*}\right)\right)\Bigg)\log\frac{m_t^2}{m_t^2-M_H^2}\Bigg]$$

$$\delta_{gg}=\frac{1}{8\sqrt{2}\pi^2m_tv}\Bigg(i\mathrm{Im}\left(\eta_L^{tu_i*}\xi_L^{u_it}+\eta_R^{tu_i*}\xi_R^{u_it}\right)\left(\log\frac{\mu^2}{m_t^2}+\log\frac{m_t^2}{m_t^2-M_H^2}\right)$$
$$-i\mathrm{Im}\left(\eta_L^{u_it}\xi_L^{tu_i*}+\eta_R^{u_it}\xi_R^{tu_i*}\right)+\frac{\sqrt{2}m_t}{v}\left(\left|\xi_L^{u_it}\right|^2+\left|\xi_R^{u_it}\right|^2\right)\Bigg)$$

$$\frac{d_{u_i}}{e}(\mu)=\sum_j\frac{g_2^2V_{td_j}V_{u_id_j}^*}{32\pi^2}\frac{m_{u_i}}{m_t^2}\left[\frac{5x_{d_j}^2-5x_{d_j}-6}{2\left(x_{d_j}-1\right)}+3x_{d_j}\log\left(\frac{\mu^2}{m_{d_j}^2}\right)+3\frac{x_{d_j}\log\left(x_{d_j}\right)}{\left(x_{d_j}-1\right)^2}\right]\mathrm{Re}\left(\lambda_R^{u_it}\right)$$
$$+\frac{q_{u_i}}{16\pi^2m_t}\left\{C_Fg_s^2(\mu)\mathrm{Im}\left(\xi_L^{u_it}\xi_R^{tu_i*}\right)\log\left(\frac{\mu^2}{m_t^2}\right)+e^2\mathrm{Im}\left(\lambda_L^{u_it}\lambda_R^{tu_i*}\right)\log\left(\frac{\mu^2}{m_t^2}\right)\right\},$$
$$+\frac{q_{u_i}}{32\pi^2}\frac{m_t}{M_H^2}\left\{\frac{\left(x_{t/H}-3\right)}{2\left(x_{t/H}-1\right)^2}+\frac{\log\,x_{t/H}}{\left(x_{t/H}-1\right)^3}\right\}\mathrm{Im}\left(\eta_L^{u_it}\eta_R^{tu_i*}\right)$$
$$+\frac{g_2^2}{16\pi^2c_W^2}\frac{q_{u_i}}{m_t}\Bigg[\mathrm{Im}\left(\kappa_L^{u_it}X_L^{tu_i*}+X_R^{u_it}\kappa_R^{tu_i*}+\kappa_L^{u_it}\kappa_R^{tu_i*}\right)\log\left(\frac{\mu^2}{m_t^2}\right)$$
$$+\frac{\log\,x_t}{4\left(x_{t/Z}-1\right)^2}\left\{x_{t/Z}^2\mathrm{Im}\left(2X_L^{tu_i*}X_R^{u_it}+X_L^{u_it}\kappa_L^{tu_i*}+X_R^{tu_i*}\kappa_R^{u_it}\right)\right.$$
$$+2\mathrm{Im}\left(X_L^{u_it}\kappa_L^{tu_i*}+X_R^{tu_i*}\kappa_R^{u_it}+\kappa_L^{u_it}\kappa_R^{tu_i*}\right)$$
$$\left.+x_{t/Z}\mathrm{Im}\left(2X_L^{tu_i*}X_R^{u_it}+9\left(X_L^{tu_i*}\kappa_L^{u_it}+X_R^{u_it}\kappa_R^{tu_i*}\right)+10\kappa_L^{u_it}\kappa_R^{tu_i*}\right)\right\}$$
$$+\frac{\log\,x_{t/Z}}{2\left(x_{t/Z}-1\right)^3}\left\{2x_{t/Z}^2\mathrm{Im}\left(X_L^{tu_i*}X_R^{u_it}\right)+x_{t/Z}\mathrm{Im}\left(5\left(X_L^{tu_i*}\kappa_L^{tu_i*}+X_R^{tu_i*}\kappa_R^{u_it}\right)+8\kappa_L^{u_it}\kappa_R^{tu_i*}\right)\right.$$
$$\left.+2\mathrm{Im}\left(X_L^{u_it}\kappa_L^{tu_i*}+X_R^{tu_i*}\kappa_R^{u_it}+\kappa_L^{tu_i*}\kappa_R^{u_it}\right)\right\}\Bigg]$$
$$w(\mu)=\frac{g_s^3(\mu)}{(32\pi^2)^2}\frac{1}{M_H^2}\left\{-\frac{x_{t/H}^2-5x_{t/H}-2}{3\left(x_{t/H}-1\right)^2}-\frac{2x_{t/H}}{\left(x_{t/H}-1\right)^4}\log\,x_{t/H}\right\}\mathrm{Im}\left(\eta_L^{ct}\eta_R^{tc*}\right)$$

$x_{d_j}=m_{d_j}^2/M_W^2, x_{t/Z}=m_t^2/M_Z^2$, and $x_{t/H}=m_t^2/M_H^2$

$$\frac{d_{d_i}}{e}(\mu)=\frac{g_2^2m_{d_i}}{32\pi^2M_W^2}V_{td_i}\left(V_{cd_i}^*\mathrm{Re}(\lambda_R^{ct})+V_{ud_i}^*\mathrm{Re}(\lambda_R^{ut})\right)\left\{\frac{1}{(x_t-1)}+\frac{x_t}{(x_t-1)^2}\log\,x_t\right\}$$
$$\frac{\tilde{d}_{d_i}}{g_s}(\mu)=\frac{g_2^2m_{d_i}}{32\pi^2M_W^2}V_{td_i}\left(V_{cd_i}^*\mathrm{Re}(\xi_R^{ct})+V_{ud_i}^*\mathrm{Re}(\xi_R^{ut})\right)\left\{\frac{1}{(x_t-1)}+\frac{x_t}{(x_t-1)^2}\log\,x_t\right\}$$

$$\hat{\mu}_t^{\mathrm{NP}}=\sum_{i,j}\frac{g_2^2 m_t}{64\pi^2 M_W^2}V_{td_j}V_{u_i d_j}\left(-m_{u_i}\mathrm{Im}\left(\xi_L^{u_i t}\right)+m_t\mathrm{Im}\left(\xi_R^{u_i t}\right)\right)\left(\frac{2x_t^2-x_t-2}{x_t^2}+\log\frac{\mu^2}{M_W^2}\right.$$
$$\left.+\frac{(x_t-1)^2(x_t+2)}{x_t^3}\log\frac{M_W^2}{M_W^2-m_t^2}\right)$$
$$\hat{d}_t^{\mathrm{NP}}=\sum_{i,j}\frac{g_2^2 m_t}{64\pi^2 M_W^2}V_{td_j}V_{u_i d_j}\left(m_{u_i}\mathrm{Re}\left(\xi_L^{u_i t}\right)+m_t\mathrm{Re}\left(\xi_R^{u_i t}\right)\right)\left(\frac{2x_t^2-x_t-2}{x_t^2}+\log\frac{\mu^2}{M_W^2}\right.$$
$$\left.+\frac{(x_t-1)^2(x_t+2)}{x_t^3}\log\frac{M_W^2}{M_W^2-m_t^2}\right)$$

$$R\left(D^{(*)}\right)=\frac{\Gamma\left(B\to D^{(*)}\tau\nu\right)}{\Gamma\left(B\to D^{(*)}\mu\nu\right)}$$

$$\partial_m^2\log\langle W\rangle|_{m=0}=\int d^4x_1 d^4x_2\hat{\mu}(x_1,x_2)\langle\mathcal{O}_2(x_1)\mathcal{O}_2(x_2)\rangle_W$$

$$\mathcal{Z}=\int da\mathrm{e}^{-\frac{8\pi^2 N}{\lambda}\mathrm{tr}a^2}\left|\mathcal{Z}_{1-\mathrm{loop}}(a)\mathcal{Z}_{\mathrm{inst}}(a)\right|^2$$

$$a\mapsto\sqrt{\frac{\lambda}{8\pi^2 N}}a,$$

$$\mathcal{Z}_{\mathcal{N}=2^*}(m)=\left(\frac{\lambda}{8\pi^2 N}\right)^{\frac{N(2N+1)}{2}}\int da\exp\left[-\mathrm{tr}a^2+m^2\widetilde{M}+O(m^4)\right]$$

$$\widetilde{M}=\widetilde{M}^{(1)}+\widetilde{M}^{(2)}$$

$$\widetilde{M}^{(1)}=-\frac{1}{2}\sum_{n=1}^{\infty}(-1)^n(2n+1)\zeta_{2n+1}\left(\frac{\lambda}{2\pi^2 N}\right)^n\mathrm{tr}a^{2n}$$
$$\widetilde{M}^{(2)}=-\frac{1}{2}\sum_{n=1}^{\infty}\sum_{k=0}^{n}(-1)^n(2n+1)\zeta_{2n+1}\left(\frac{\lambda}{8\pi^2 N}\right)^n\binom{2n}{2k}\mathrm{tr}a^{2n-2k}\mathrm{tr}a^{2k}$$

$$\mathcal{Z}\left(m_f\right)=\left(\frac{\lambda}{8\pi^2 N}\right)^{\frac{N(2N+1)}{2}}\int da\exp\left[-\mathrm{tr}a^2-S_0+\sum_{f=1}^{4}m_f^2 M+O\left(m_f^4\right)\right]$$

$$S_0 = 4\sum_{n=1}^{\infty}(-1)^{n+1}\left(\frac{\lambda}{8\pi^2 N}\right)^{n+1}(2^{2n}-1)\frac{\zeta_{2n+1}}{n+1}\mathrm{tr}a^{2n+2},$$

$$M = -\sum_{n=1}^{\infty}(-1)^{n}(2n+1)\zeta_{2n+1}\left(\frac{\lambda}{8\pi^2 N}\right)^{n}\mathrm{tr}a^{2n}.$$

$$W(a,\lambda) = \sum_{k=0}^{\infty}\frac{1}{k!}\left(\frac{\lambda}{2N}\right)^{\frac{k}{2}}\mathrm{tr}a^{k}$$

$$\tilde{\mathcal{J}}(\lambda) \equiv \partial_m^2 \log\,\widetilde{\mathcal{W}}(m,\lambda)\big|_{m=0} = 2\frac{\langle W\widetilde{M}\rangle_0 - \langle W\rangle_0\langle\widetilde{M}\rangle_0}{\langle W\rangle_0}$$

$$\mathcal{J}(\lambda) \equiv \partial_{m_f}^2 \log\,\mathcal{W}(m_f,\lambda)\Big|_{m_f=0} = 2\frac{\langle WM\rangle - \langle W\rangle\langle M\rangle}{\langle W\rangle}$$

$$\langle f(a)\rangle = \frac{\langle f(a)\mathrm{e}^{-S_0}\rangle_0}{\langle \mathrm{e}^{-S_0}\rangle_0}$$

$$\partial_y^2 \log\,\mathcal{Z}_N(y) = \frac{\mathcal{Z}_{N+1}(y)\mathcal{Z}_{N-1}(y)}{\mathcal{Z}_N(y)^2}$$

$$\mathcal{Z}_{N=-1}(y) = 0, \mathcal{Z}_{N=0}(y) = 1$$

$$y = \frac{(4\pi)^2 N}{\lambda}$$

$$\partial_y^2 F_N(y) = -\exp\left[-F_{N+1}(y) + 2F_N(y) - F_{N-1}(y)\right]$$

$$F_N(\lambda) \equiv F_N^{\mathcal{N}=4}(\lambda) + \Delta F_N(\lambda)$$
$$= F_N^{\mathcal{N}=4}(\lambda) + NF^{(1)}(\lambda) + F^{(2)}(\lambda) + \frac{F^{(3)}(\lambda)}{N} + O\left(\frac{1}{N^2}\right)$$

$$F_N^{\mathcal{N}=4}(y) = \frac{N(2N+1)}{2}\log\,y - \log\left(\frac{G(N+1)G(N+3/2)}{G(3/2)}\right)$$

$$F^{(1)}(\lambda) = \frac{\log\,2}{2\pi^2}\lambda + 4\int_0^{\infty}\frac{dt}{t}\frac{\mathrm{e}^t}{(\mathrm{e}^t+1)^2}\left[\left(\frac{2\pi}{\sqrt{\lambda}t}\right)J_1\left(\frac{t\sqrt{\lambda}}{\pi}\right) - 1\right]$$

$$K_N = M_N W_N - \mathcal{M}_N \mathcal{W}_N$$

$$\mathcal{M}_N = N\mathcal{M}^{(0)}(\lambda) + \mathcal{M}^{(1)}(\lambda) + \frac{\mathcal{M}^{(2)}(\lambda)}{N} + O\left(\frac{1}{N^2}\right)$$
$$\mathcal{W}_N = N\mathcal{W}^{(0)}(\lambda) + \mathcal{W}^{(1)}(\lambda) + \frac{\mathcal{W}^{(2)}(\lambda)}{N} + O\left(\frac{1}{N^2}\right)$$

$$\mathcal{W}^{(0)}(\lambda) = \frac{4I_1(\sqrt{\lambda})}{\sqrt{\lambda}}$$

$$\mathcal{M}^{(0)}(\lambda) = -2\int_0^\infty dt\,\frac{\mathrm{e}^t t}{(\mathrm{e}^t-1)^2}\left[\left(\frac{4\pi}{\sqrt{\lambda}t}\right)J_1\left(\frac{\sqrt{\lambda}t}{2\pi}\right)-1\right]$$

$$\partial_\lambda\mathcal{M}^{(1)}(\lambda) = -\frac{1}{4}\left(2\lambda^2\partial_\lambda^2 F^{(1)}(\lambda)+4\lambda\partial_\lambda F^{(1)}(\lambda)-1\right)\left(\lambda\partial_\lambda^2\mathcal{M}^{(0)}(\lambda)+2\partial_\lambda\mathcal{M}^{(0)}(\lambda)\right)$$

$$\mathcal{K}_N(\lambda) = \mathcal{K}^{(0)}(\lambda)+\frac{\mathcal{K}^{(1)}(\lambda)}{N}+\frac{\mathcal{K}^{(2)}(\lambda)}{N^2}+O\left(\frac{1}{N^3}\right)$$

$$\partial_\lambda\mathcal{K}^{(0)}(\lambda) = \frac{1}{2\lambda}\left[\left(\lambda^2\partial_\lambda^2+2\lambda\partial_\lambda\right)\mathcal{W}^{(0)}(\lambda)\right]\left[\left(\lambda^2\partial_\lambda^2+2\lambda\partial_\lambda\right)\mathcal{M}^{(0)}(\lambda)\right]$$

$$\mathcal{K}^{(0)}(\lambda) = 2\pi\int_0^\infty dt\,\frac{\mathrm{e}^t t^2}{(1-\mathrm{e}^t)^2}\frac{1}{4\pi^2+t^2}\mathcal{B}(t)$$

$$\mathcal{B}(t) \equiv \sqrt{\lambda}I_0(\sqrt{\lambda})J_1\left(\frac{t\sqrt{\lambda}}{2\pi}\right)-\frac{t\sqrt{\lambda}}{2\pi}I_1(\sqrt{\lambda})J_0\left(\frac{t\sqrt{\lambda}}{2\pi}\right)$$

$$\mathcal{J}(\lambda) = \mathcal{J}^{(0)}(\lambda)+\frac{1}{N}\mathcal{J}^{(1)}(\lambda)+\frac{1}{N^2}\mathcal{J}^{(2)}(\lambda)+O\left(\frac{1}{N^3}\right)$$

$$\mathcal{J}^{(n)}(\lambda) = \frac{2\mathcal{K}^{(n)}(\lambda)}{\mathcal{W}^{(0)}(\lambda)}-\sum_{i=1}^{n}\frac{\mathcal{W}^{(i)}(\lambda)}{\mathcal{W}^{(0)}(\lambda)}\mathcal{J}^{(n-i)}(\lambda)$$

$$\partial_\lambda F^{(1)}(\lambda) \underset{\lambda\to\infty}{\sim} \frac{\log(2)}{2\pi^2}-\frac{1}{2\lambda}+\frac{\pi^2}{2\lambda^2},$$
$$\partial_\lambda\mathcal{M}^{(0)}(\lambda) \underset{\lambda\to\infty}{\sim} \frac{1}{\lambda}-\frac{4\pi^2}{3\lambda^2},$$
$$\mathcal{K}^{(0)}(\lambda) \underset{\lambda\to\infty}{\sim} I_0(\sqrt{\lambda}).$$

$$\partial_\lambda F_1(\lambda) = \frac{\log(2)}{2\pi^2}-\frac{1}{2\lambda}+\frac{\pi^2}{2\lambda^2}+\frac{1}{\pi^2}\sum_{n=0}^{\infty}\left(K_0((2n+1)\sqrt{\lambda})-K_4((2n+1)\sqrt{\lambda})\right)$$

$$\partial_\lambda\mathcal{M}^{(0)}(\lambda) = \frac{2}{\lambda}\int_0^\infty dt\,\frac{\mathrm{e}^t t}{(\mathrm{e}^t-1)^2}J_2\left(\frac{t\sqrt{\lambda}}{\pi}\right)$$

$$\frac{\mathrm{e}^t}{(\mathrm{e}^t-1)^2} = \sum_{n=1}^{\infty} n\mathrm{e}^{-nt}$$

$$\partial_\lambda\mathcal{M}^{(0)}(\lambda) = \frac{1}{2\pi^2 a^2}\sum_{n=1}^{\infty} n\int_0^\infty dtt\mathrm{e}^{-nt}J_2(at)$$

$$\mathcal{A}_1 = \frac{1}{\pi^2 a^4}\sum_{n=1}^{\infty} n$$
$$\mathcal{A}_2 = -\frac{1}{2\pi^2 a^2}\sum_{n=1}^{\infty}\left(\frac{n^2}{(a^2+n^2)^{3/2}}+\frac{2}{a^2}\frac{n^2}{(a^2+n^2)^{1/2}}\right)$$

$$\mathcal{A}_1 = -\frac{1}{12\pi^2 a^4}$$

$$\frac{1}{(n^2+a^2)^\alpha} = \frac{1}{\Gamma(\alpha)}\int_0^\infty dt t^{\alpha-1}\mathrm{e}^{-(n^2+a^2)t}$$

$$\mathcal{A}_2 = -\frac{1}{a^4\pi^{\frac{5}{2}}}\int_0^\infty dt\frac{1+a^2t}{\sqrt{t}}\mathrm{e}^{-a^2t}\sum_{n=1}^\infty n^2\mathrm{e}^{-n^2t}$$

$$\sum_{n=1}^\infty n^2\mathrm{e}^{-n^2t} = \frac{\sqrt{\pi}}{4t^{\frac{3}{2}}} + \frac{\sqrt{\pi}}{4t^{\frac{5}{2}}}\sum_{k=1}^\infty e^{-\frac{\pi^2k^2}{t}}(2t-4\pi^2k^2)$$

$$\mathcal{A}_2 = D - \frac{1}{4a^4\pi^2}\int_0^\infty dt\frac{1+a^2t}{t^3}\mathrm{e}^{-a^2t}\sum_{k=1}^\infty (2t-4\pi^2k^2)\mathrm{e}^{-\frac{\pi^2k^2}{t}}$$

$$D = -\frac{1}{4a^2\pi^2}\int_0^\infty dt\frac{1+t}{t^2}e^{-t}$$

$$\mathcal{D}(z) \equiv -\frac{1}{4a^2\pi^2}\int_0^\infty dt\frac{1+t}{t^2}t^z e^{-t} = -\frac{z}{4a^2\pi^2}\Gamma(z-1)$$

$$D = \mathcal{D}(0) = \frac{1}{4a^2\pi^2}$$

$$K_\nu(z) = \frac{1}{2}\left(\frac{z}{2}\right)^\nu\int_0^\infty dt t^{-\nu-1}e^{-t-\frac{z^2}{4t}}$$

$$\mathcal{A}_2 = \frac{1}{4\pi^2a^2} - \frac{2}{a^2\pi^2}\sum_{k=1}^\infty\left(\frac{K_1(2ak\pi)}{2ak\pi} - K_2(2ak\pi)\right) - \frac{1}{a^2\pi^2}\sum_{k=1}^\infty (K_0(2ak\pi) - 2ak\pi K_1(2ak\pi)).$$

$$\partial_\lambda\mathcal{M}^{(0)}(\lambda) = \frac{1}{\lambda} - \frac{4\pi^2}{3\lambda^2} + \frac{4}{\lambda}\sum_{k=1}^\infty\left(K_0(\sqrt{\lambda}k) + \frac{(2+\lambda k^2)K_1(\sqrt{\lambda}k)}{\sqrt{\lambda}k}\right).$$

$$\partial_\lambda\mathcal{K}^{(0)}(\lambda) = \frac{I_1(\sqrt{\lambda})}{2\sqrt{\lambda}}\left(1 + 2\sqrt{\lambda}\sum_{k=1}^\infty\left(kK_1(\sqrt{\lambda}k) - k^2\sqrt{\lambda}K_0(\sqrt{\lambda}k)\right)\right).$$

$$\mathcal{K}^{(0)}(\lambda) = \int_0^\lambda dq\partial_q\mathcal{K}^{(0)}(q)$$

$$\mathcal{K}^{(0)}(\lambda) = I_0(\sqrt{\lambda}) + \sum_{k=1}^\infty (\mathcal{J}_0(k,\lambda) + \mathcal{J}_1(k,\lambda)) - \frac{7}{4} - \frac{\pi^2}{3},$$

$$\mathcal{J}_0(k,\lambda)=\begin{cases}-\dfrac{1}{2}\lambda^{3/2}\big(I_1(\sqrt{\lambda})K_0(\sqrt{\lambda})+I_2(\sqrt{\lambda})K_1(\sqrt{\lambda})\big), & k=1\\ -\dfrac{2\lambda k^2 I_0(\sqrt{\lambda})K_0(k\sqrt{\lambda})}{(k^2-1)^2}+\dfrac{4\sqrt{\lambda}k^2 I_1(\sqrt{\lambda})K_0(k\sqrt{\lambda})}{(k^2-1)^2}+\dfrac{2\lambda k^5 I_1(\sqrt{\lambda})K_1(k\sqrt{\lambda})}{(k^2-1)^2} & \\ +\dfrac{2\lambda k^4 I_0(\sqrt{\lambda})K_0(k\sqrt{\lambda})}{(k^2-1)^2}+\dfrac{4\sqrt{\lambda}k^3 I_0(\sqrt{\lambda})K_1(k\sqrt{\lambda})}{(k^2-1)^2}-\dfrac{2\lambda k^3 I_1(\sqrt{\lambda})K_1(k\sqrt{\lambda})}{(k^2-1)^2}, & k\geq 2\end{cases}$$

$$\mathcal{J}_1(k,\lambda)=\begin{cases}\lambda I_1(\sqrt{\lambda})K_1(\sqrt{\lambda})+\lambda I_0(\sqrt{\lambda})\left(K_0(\sqrt{\lambda})+\dfrac{2}{z}K_1(\sqrt{\lambda})\right), & k=1\\ -\dfrac{2k\sqrt{\lambda}\big(kI_1(\sqrt{\lambda})K_0(k\sqrt{\lambda})+I_0(\sqrt{\lambda})K_1(k\sqrt{\lambda})\big)}{k^2-1}, & k\geq 2\end{cases}$$

$$\begin{aligned}\mathcal{W}^{(1)}(\lambda)=&I_0(\sqrt{\lambda})-\lambda\frac{\log(2)}{\pi^2}I_2(\sqrt{\lambda})+\frac{1}{2\pi^2}-\frac{2}{3}\\ &-\frac{2}{\pi^2}\sum_{k=0}^{\infty}\frac{1}{(2k+1)^2}(\mathcal{J}_1(2k+1,\lambda)-\mathcal{J}_0(2k+1,\lambda))\end{aligned}$$

$$\begin{aligned}\mathcal{I}^{(0)}(\lambda)=&\frac{\sqrt{\lambda}}{2}\frac{I_0(\sqrt{\lambda})}{I_1(\sqrt{\lambda})}-\frac{\sqrt{\lambda}}{2I_1(\sqrt{\lambda})}\left(\frac{7}{4}+\frac{\pi^2}{3}\right)+\lambda^{3/2}K_1(\sqrt{\lambda})-\frac{\lambda^2}{4}K_0(\sqrt{\lambda})\\ &-\frac{I_0(\sqrt{\lambda})}{4I_1(\sqrt{\lambda})}\left[(\lambda-4)\lambda K_1(\sqrt{\lambda})-2\lambda^{3/2}K_0(\sqrt{\lambda})\right]\\ &+\lambda\sum_{k=2}^{\infty}\frac{k}{(k^2-1)^2}\left[\frac{I_0(\sqrt{\lambda})}{I_1(\sqrt{\lambda})}A_k(\lambda)+kB_k(\lambda)\right]\end{aligned}$$

$$\begin{aligned}A_k(\lambda)&=k(k^2-1)\sqrt{\lambda}K_0(k\sqrt{\lambda})+(k^2+1)K_1(k\sqrt{\lambda}),\\ B_k(\lambda)&=(3-k^2)K_0(k\sqrt{\lambda})+k(k^2-1)\sqrt{\lambda}K_1(k\sqrt{\lambda}).\end{aligned}$$

$$\begin{aligned}\mathcal{I}^{(0)}(\lambda)\underset{\lambda\to\infty}{\sim}&\frac{\sqrt{\lambda}}{2}+\frac{1}{4}+\frac{3}{16\sqrt{\lambda}}+\frac{3}{16\lambda}+\frac{63}{256\lambda^{3/2}}+O\left(\frac{1}{\lambda^2}\right)\\ &+\mathrm{e}^{-\sqrt{\lambda}}\sqrt{\frac{\pi}{2}}\left[-\frac{\lambda^{7/4}}{2}+\frac{21}{16}\lambda^{5/4}-\frac{243+256\pi^2}{768}\lambda^{3/4}+\frac{351-256\pi^2}{2048}\lambda^{1/4}+O\left(\frac{1}{\lambda^{1/4}}\right)\right]\\ &+\mathrm{e}^{-2\sqrt{\lambda}}\left[2\sqrt{\pi}\lambda^{5/4}+\frac{7}{8}\sqrt{\pi}\lambda^{3/4}+i\sqrt{\lambda}+\frac{153}{256}\sqrt{\pi}\lambda^{1/4}+\frac{3i}{4}+O\left(\frac{1}{\lambda^{1/4}}\right)\right]\\ &+O\left(\mathrm{e}^{-3\sqrt{\lambda}}\right)\end{aligned}$$

$$\mathcal{J}^{(1)}(\lambda) \underset{\lambda\to\infty}{\sim} -\frac{\lambda^{3/2}\log 2}{8\pi^2} - \frac{\sqrt{\lambda}}{8} + \frac{3\sqrt{\lambda}\log 2}{64\pi^2} - \frac{1}{8} + \frac{3\log 2}{32\pi^2} + O\left(\lambda^{-1/2}\right)$$
$$+\mathrm{e}^{-\sqrt{\lambda}}\left[-\frac{\lambda^{13/4}\log 2}{8\sqrt{2}\pi^{3/2}} + \frac{49\lambda^{11/4}\log 2}{64\sqrt{2}\pi^{3/2}}\right.$$
$$\left.+\left(\frac{1}{4\sqrt{2}\pi^{3/2}} + \frac{1}{8}\sqrt{\frac{\pi}{2}} - \frac{921\log 2}{1024\sqrt{2}\pi^{3/2}} - \frac{1}{12}\sqrt{\frac{\pi}{2}}\log 2\right)\lambda^{9/4} + O\left(\lambda^{7/4}\right)\right]$$
$$+\mathrm{e}^{-2\sqrt{\lambda}}\left[-\frac{\lambda^{7/2}}{8\pi} + \frac{5\lambda^3}{32\pi} + \frac{\lambda^{11/4}\log 2}{\pi^{3/2}}\right.$$
$$\left.-\frac{13\lambda^{9/4}\log 2}{16\pi^{3/2}} + \left(\frac{155}{256\pi} - \frac{\pi}{6}\right)\lambda^{5/2} + O(\lambda^2)\right] + O\left(\mathrm{e}^{-3\sqrt{\lambda}}\right),$$
$$\mathcal{J}^{(2)}(\lambda) \underset{\lambda\to\infty}{\sim} \frac{3\log^2 2}{64\pi^4}\lambda^{5/2} + \left(-\frac{1}{768} - \frac{3\log^2 2}{512\pi^4} + \frac{\log 256}{256\pi^2}\right)\lambda^{3/2} + \frac{3}{256}\lambda + O(\sqrt{\lambda})$$
$$+\mathrm{e}^{-\sqrt{\lambda}}\left[-\frac{\log 2}{64\sqrt{2}\pi^{7/2}}\lambda^{19/4} + \frac{101\log^2 2}{512\sqrt{2}\pi^{7/2}}\lambda^{17/4} + \left(-\frac{5233\log^2 2}{8192\sqrt{2}\pi^{7/2}} - \frac{\log^2 2}{96\sqrt{2}\pi^{3/2}}\right.\right.$$
$$\left.\left.+\frac{\log 2}{16\sqrt{2}\pi^{7/2}} + \frac{\log 2}{32\sqrt{2}\pi^{3/2}}\right)\lambda^{15/4} + O\left(\lambda^{13/4}\right)\right]$$
$$+\mathrm{e}^{-2\sqrt{\lambda}}\left[-\frac{\log 2}{16\pi^3}\lambda^5 + \frac{19\log 2}{64\pi^3}\lambda^{9/2} + \frac{\log^2 2}{4\pi^{7/2}}\lambda^{17/4} + O(\lambda^4)\right] + O\left(\mathrm{e}^{-3\sqrt{\lambda}}\right)$$

$$\tilde{\mathcal{J}}(\lambda) = \tilde{\mathcal{J}}^{(0)}(\lambda) + \frac{1}{N}\tilde{\mathcal{J}}^{(1)}(\lambda) + O\left(\frac{1}{N^2}\right)$$

$$\tilde{\mathcal{J}}^{(0)}(\lambda) = \frac{8\pi^2}{I_1(\sqrt{\lambda})}\int_0^\infty dt\,\frac{\mathrm{e}^t t}{(\mathrm{e}^t-1)^2}\frac{1}{4\pi^2+t^2}J_1\left(\frac{\sqrt{\lambda}t}{2\pi}\right)\mathcal{B}(t)$$

$$\tilde{\mathcal{J}}^{(1)}(\lambda) = \frac{1}{\mathcal{W}^{(0)}(\lambda)}\left[2\widetilde{\mathcal{K}}^{(1)}(\lambda) - \frac{1}{2}\left(I_0(\sqrt{\lambda})-1\right)\tilde{\mathcal{J}}^{(0)}(\lambda)\right]$$

$$\begin{aligned}\widetilde{\mathcal{K}}^{(1)}(\lambda) =& \frac{\pi}{2}\int_0^\infty dt\,\frac{\mathrm{e}^t t^2}{(1-\mathrm{e}^t)^2}\frac{1}{\pi^2+t^2}\mathcal{B}(2t)\\ &+\frac{\sqrt{\lambda}}{2}I_1(\sqrt{\lambda})\int_0^\infty dt\,\frac{\mathrm{e}^t t}{(1-\mathrm{e}^t)^2}J_1\left(\frac{t\sqrt{\lambda}}{2\pi}\right)^2\\ &-\pi\int_0^\infty dt\,\frac{\mathrm{e}^t t^2}{(1-\mathrm{e}^t)^2}\frac{1}{4\pi^2+t^2}\left[1-J_0\left(\frac{t\sqrt{\lambda}}{2\pi}\right)\right]\mathcal{B}(t)\end{aligned}$$

$$\tilde{\mathcal{J}}^{(0)}(\lambda) \underset{\lambda\to\infty}{\sim} \mathcal{Q}_{(1,0)}(\lambda) + \frac{I_0(\sqrt{\lambda})}{I_1(\sqrt{\lambda})}\mathcal{Q}_{(1,1)}(\lambda) \equiv \sum_{n=0}^{\infty} a_n \lambda^{\frac{1-n}{2}}$$

$$\mathcal{Q}_{(1,0)}(\lambda) = -1 + \frac{1}{3}\left(\frac{\lambda}{\pi}\right)^{3/2}\sum_{s=1}^{\infty}\frac{1}{\lambda^s}\frac{\Gamma\left(s-\frac{3}{2}\right)^2\Gamma\left(s-\frac{1}{2}\right)}{\Gamma(s-1)}\left(12\sum_{i=1}^{s-2} i\zeta_{2i+1} + \pi^2 - 3\right)$$

$$\mathcal{Q}_{(1,1)}(\lambda) = \sqrt{\lambda} + \frac{\lambda^2}{3\pi^{3/2}}\sum_{s=1}^{\infty}\frac{1}{\lambda^s}\frac{\Gamma\left(s-\frac{5}{2}\right)\Gamma\left(s-\frac{3}{2}\right)\Gamma\left(s-\frac{1}{2}\right)}{\Gamma(s-1)}\left(12\sum_{i=1}^{s-2} i\zeta_{2i+1} + \pi^2 - 3\right)$$

$$\sum_{n\geq 0} a_n z^n \mapsto \mathbf{B}\left[\frac{1}{\sqrt{\lambda}}\tilde{\jmath}^{(0)}\right] \equiv \sum_{n\geq 0} a_n \frac{t^n}{n!}$$

$$a_n \underset{n\gg 1}{\sim} \frac{\mathcal{S}_1}{2\pi i}\left(\frac{f(b_1)}{A^{n+b_1}}\sum_{m\geq 0} A^m c_m^{(1)}\Gamma(n+b_1-m) + \frac{\Gamma(n+1)}{A^{n+b_1+1}}\right)$$
$$+\frac{\mathcal{S}_2}{2\pi i}\left(\frac{f(b_2)}{(-A)^{n+b_2}}\sum_{m\geq 0} (-A)^m c_m^{(2)}\Gamma(n+b_2-m) + \frac{\Gamma(n+1)}{(-A)^{n+b_2+1}}\right)$$

$$A_n^{\text{even}} \equiv n\sqrt{\frac{a_n}{a_{n+2}}} \quad \text{with} \quad n = 0,2,4,6\ldots$$

$$A_n^{\text{odd}} \equiv n\sqrt{\frac{a_n}{a_{n+2}}} \quad \text{with} \quad n = 1,3,5,7\ldots$$

$$s_n = \sum_{k=0}^{\infty} \frac{g_k}{n^k},$$

$$s_n^{(M)} = \sum_{\ell=0}^{M} \frac{s_{n+\ell}(n+\ell)^M(-1)^{\ell+M}}{\ell!\,(M-\ell)!}.$$

$$r_n^{\text{even}} = \frac{a_{n+1}}{a_n}\frac{2}{n}, n = 0,2,4,6,\ldots$$
$$r_n^{\text{odd}} = \frac{a_{n+1}}{a_n}\frac{2}{n}, n = 1,3,5,7,\ldots.$$

$$r^{\text{even}} = 1+\frac{3}{n}-\frac{1}{2n^2}-\frac{21}{n^3}-\frac{393}{8n^4}-\left(72\zeta_3+\frac{1113}{8}\right)\frac{1}{n^5}-\left(444\zeta_3+\frac{11163}{16}\right)\frac{1}{n^6}$$
$$-\left(2304\zeta_3+720\zeta_5+\frac{40509}{8}\right)\frac{1}{n^7}-\left(14997\zeta_3+10890\zeta_5+\frac{5305437}{128}\right)\frac{1}{n^8}+O\left(\frac{1}{n^9}\right)$$
$$r^{\text{odd}} = 1-\frac{1}{n}-\frac{1}{2n^2}+\frac{1}{n^3}-\frac{129}{8n^4}-\left(24\zeta_3+\frac{181}{8}\right)\frac{1}{n^5}-\left(108\zeta_3+\frac{4243}{16}\right)\frac{1}{n^6}$$
$$-\left(804\zeta_3+360\zeta_5+\frac{15503}{8}\right)\frac{1}{n^7}-\left(6129\zeta_3+4770\zeta_5+\frac{2211405}{128}\right)\frac{1}{n^8}+O\left(\frac{1}{n^9}\right).$$

$$c_i^{(2)} = \left\{1,-1,\frac{3}{32},-\frac{3}{16},-\frac{969}{2048},-\frac{2439}{2048},-\frac{228933}{65536},-\frac{393309}{32768},\ldots\right\}$$
$$c_i^{(1)} = \left\{\frac{5}{4},\frac{61}{32},\frac{213}{128},\frac{3\zeta_3}{4}+\frac{2715}{2048},\frac{9\zeta_3}{16}+\frac{13671}{8192},\frac{153\zeta_3}{128}+\frac{45\zeta_5}{32}+\frac{173961}{65536},\ldots\right\},$$

$$\frac{\mathrm{e}^{-2\sqrt{\lambda}}}{\sqrt{\lambda}}\left[8+(8i)\sum_{n\geq 0} c_n^{(1)}\left(\frac{1}{\sqrt{\lambda}}\right)^n\right]$$

$$\mathcal{Q}_{(0,0)}(\lambda) = \frac{\sqrt{\lambda}}{2\pi}\int_0^{\infty} dt\,\frac{\mathrm{e}^t t^3}{(1-\mathrm{e}^t)^2}\frac{1}{4\pi^2+t^2}J_0\left(\frac{t\sqrt{\lambda}}{2\pi}\right)J_0\left(\frac{t\sqrt{\lambda}}{2\pi}\right)$$

$$\mathcal{Q}_{(0,0)}(\lambda) \underset{\lambda\to\infty}{\sim} \frac{\lambda}{4\pi^{3/2}} \sum_{s=1}^{\infty} \frac{(2\pi)^{2s}}{\lambda^s} \frac{1}{\Gamma\left(\frac{3}{2}-s\right)^3 \Gamma(s)} \mathcal{J}(s)$$

$$\mathcal{J}(s) = (-1)^{s-1} 4^{1-s} \pi^{-2(s-1)} \left(\sum_{i=1}^{s-2} i\zeta_{2i+1} - \frac{1}{2}\delta_{s,1} + \frac{\pi^2}{12} - \frac{1}{4} \right)$$

$$\tilde{\jmath}^{(1)}(\lambda) \underset{\lambda\to\infty}{\sim} \sqrt{\lambda} \sum_{n=0}^{\infty} b_n \lambda^{-n/2}$$

$$b_0 = \frac{1}{8}, b_1 = 0, b_2 = -\frac{3}{64}, b_3 = -\frac{3}{32}(1 + 4\zeta_3)$$

$$\frac{2}{n}\frac{b_{n+1}}{b_n} \underset{n\gg 1}{\sim} 1 + \frac{4}{n} + \frac{7}{2}\frac{1}{n^2} - \frac{35}{2}\frac{1}{n^3} - \frac{533}{8}\frac{1}{n^4} - \left(\frac{823}{4} + 72\zeta_3\right)\frac{1}{n^5} - \left(\frac{14455}{16} + 516\zeta_3\right)\frac{1}{n^6}$$
$$- \left(2820\zeta_3 + 720\zeta_5 + \frac{95473}{16}\right)\frac{1}{n^7} + O\left(\frac{1}{n^8}\right)$$

$$\frac{2}{n}\frac{b_{n+1}}{b_n} \underset{n\gg 1}{\sim} 1 - \frac{1}{2}\frac{1}{n^2} + \frac{1}{2}\frac{1}{n^3} - \frac{125}{8}\frac{1}{n^4} - \left(\frac{153}{4} + 24\zeta_3\right)\frac{1}{n^5} - \left(\frac{4855}{16} + 132\zeta_3\right)\frac{1}{n^6}$$

$$- \left(936\zeta_3 + 360\zeta_5 + \frac{35861}{16}\right)\frac{1}{n^7} + O\left(\frac{1}{n^8}\right)$$

$$b_n \underset{n\gg 1}{\sim} \frac{\mathcal{S}_1}{2\pi i 2^n}\left[\sum_{j\geq 0} d_j^{(1)} 2^j \Gamma(n-j) + \frac{d}{2}\Gamma(n+1) - \frac{1}{4}\Gamma(n+2)\right]$$
$$+ \frac{\mathcal{S}_2}{2\pi i(-2)^n}\left[\sum_{j\geq 0} d_j^{(2)} (-2)^j \Gamma(n-j) - \frac{1}{2}\Gamma(n+1)\right]$$

$$\mathcal{S}_2 = \frac{\mathcal{S}_1}{2} = i, d = -\frac{1}{4}$$

$$d_j^{(2)} = \left\{-\frac{1}{2}, \frac{3}{32}, -\frac{15}{64}, -\frac{585}{2048}, \frac{423\zeta_3}{32} + \frac{27\zeta_5}{8} + \frac{562983}{20480}\right\}$$
$$d_j^{(1)} = \left\{-\frac{41}{32}, -\frac{213}{128}, -\frac{3\zeta_3}{4} - \frac{4419}{2048}, -\frac{81\zeta_3}{8} - \frac{9\zeta_5}{4} - \frac{886439}{40960}\right\}$$

$$\mathrm{e}^{-\sqrt{\lambda}}\left[\sqrt{\frac{\pi}{2}}\left(\frac{1}{4}\frac{1}{\lambda^{7/4}} - \frac{21}{32}\frac{1}{\lambda^{5/4}} + \left(\frac{243 + 64\pi^2}{1536}\right)\frac{1}{\lambda^{3/4}} + O\left(\frac{1}{\lambda^{1/4}}\right)\right)\right]$$
$$+ \mathrm{e}^{-2\sqrt{\lambda}}\left[-\frac{2}{\lambda} + \frac{1}{2\sqrt{\lambda}} + (2i)\sum_{n\geq 0} d_n^{(1)}\left(\frac{1}{\sqrt{\lambda}}\right)^{n+1}\right] + O\left(\mathrm{e}^{-3\sqrt{\lambda}}\right)$$

$$S_{\mathrm{F}_1} = -T_{\mathrm{F}_1}\int d^{\mathcal{D}}\sigma\sqrt{-\det G_{\mu\nu}(X)\partial_a X^\mu \partial_b X^\nu} + \cdots$$

$$\mathrm{Z}_k^{(p)} = \int_0^\infty dt \frac{\mathrm{e}^t t^p}{(\mathrm{e}^t - 1)^2} J_k\left(\frac{t\sqrt{\lambda}}{2\pi}\right)$$

$$\frac{d^{\mathcal{D}}}{d\lambda^2}(\lambda F_1) = \frac{\log\ 2}{\pi^2} - \frac{2}{\pi\sqrt{\lambda}}\int_0^{\infty} dt\,\frac{\mathrm{e}^t}{(\mathrm{e}^t+1)^2} J_1\left(\frac{t\sqrt{\lambda}}{\pi}\right)$$

$$\frac{d^{\mathcal{D}}}{d\lambda^2}(\lambda F_1) = \frac{\log\ 2}{\pi^2} - \frac{2}{\pi\sqrt{\lambda}}\int_{-i\infty}^{i\infty} \frac{ds}{2\pi i}\frac{\Gamma(-s)\Gamma(2s+2)}{\Gamma(s+2)}\eta(2s+1)\left(\frac{\sqrt{\lambda}}{2\pi}\right)^{2s+1}$$

$$\frac{d^{\mathcal{D}}}{d\lambda^2}(\lambda F_1) \underset{\lambda\to\infty}{\sim} \frac{\log\ 2}{\pi^2} - \frac{1}{2\lambda}$$

$$\mathcal{T}(b) = \frac{\log\ 2}{\pi^2} - \frac{2}{\pi\sqrt{\lambda}}\int_{-i\infty}^{i\infty} \frac{ds}{2\pi i}\frac{\Gamma(-s)\Gamma(2s+2)}{\Gamma(s+b)}\eta(2s+1)\left(\frac{\sqrt{\lambda}}{2\pi}\right)^{2s+1}$$

$$\mathcal{T}(b) \underset{\lambda\to\infty}{\sim} \mathcal{T}^{(\mathrm{p})}(b) = \frac{\log\ 2}{\pi^2} - \frac{1}{2\lambda\Gamma(b-1)} + \frac{\sin\,(\pi b)}{\pi^3}\sum_{n=2}^{\infty}(4^n-1)(2n-1)\zeta(2n)\Gamma(n)\Gamma(-b+n+1)\lambda^{-n}$$

$$\widetilde{\mathbf{B}}\colon \sum_{n=1}^{\infty} c_n\lambda^{-n} \longrightarrow \phi(w,b) = \sum_{n=1}^{\infty}\frac{2c_n}{\zeta(2n)\Gamma(2n+1)}(2w)^{2n}$$

$$\phi(w,b) = \phi_1(w,b) + \phi_2(w,b) + \phi_3(w,b)$$

$$\phi_1(w,b) = -\frac{4\sin\,(\pi b)}{\pi^3}\Gamma(2-b)\left(2w^2\,{}_2F_1\left(1,2-b;\frac{3}{2};w^2\right) - w^2\,{}_3F_2\left(1,1,2-b;\frac{3}{2},2;w^2\right)\right)$$

$$\phi_2(w,b) = -\frac{4\sin\,(\pi b)}{\pi^3}\Gamma(2-b)\left(-8w^2\,{}_2F_1\left(1,2-b;\frac{3}{2};4w^2\right) + 4w^2\,{}_3F_2\left(1,1,2-b;\frac{3}{2},2;4w^2\right)\right)$$

$$\phi_3(w,b) = -\frac{12w^2\Gamma(2-b)}{\pi^3}.$$

$$\mathcal{S}_\theta\left[\mathcal{T}^{(\mathrm{p})}(b)\right] = \frac{\log\ 2}{\pi^2} - \frac{1}{2\lambda\Gamma(b-1)} + \sqrt{\lambda}\int_0^{e^{i\theta}\infty}\frac{dw}{4\sinh^2\,(\sqrt{\lambda}w)}\phi(w,b)$$

$$\mathcal{S}_{\pm}\left[\mathcal{T}^{(p)}(b)\right] = \lim_{\theta\to 0^{\pm}}\mathcal{S}_\theta[\mathcal{T}(b)]$$

$$\mathcal{S}_{\mathrm{med}}\left[\mathcal{T}^{(p)}(b)\right] = \mathcal{S}_{\pm}\left[\mathcal{T}^{(p)}(b)\right] \mp \frac{1}{2}\Delta\left[\mathcal{T}^{(p)}(b)\right]$$

$$\Delta\left[\mathcal{T}^{(p)}(b)\right] \equiv (\mathcal{S}_+ - \mathcal{S}_-)[\mathcal{J}(b)] = \sqrt{\lambda}\int_0^{\infty}\frac{dw}{4\sinh^2\,(\sqrt{\lambda}w)}\mathrm{Disc}\phi(w,b)$$

$$\mathrm{Disc}_2F_1(a,b;c;z) = \frac{2\pi i\Gamma(c)}{\Gamma(a)\Gamma(b)\Gamma(c-a-b+1)}z^{1-c}(z-1)^{c-a-b}\,{}_2F_1(1-a,1-b;-a-b+c+1;1-z)$$

$$\mathrm{Disc}_3F_2\left(1,1,2-b;\frac{3}{2},2;z\right) = -i\sin\,(\pi b)\sqrt{\pi}z^{-\frac{1}{2}}(z-1)^{b-\frac{1}{2}}\frac{\Gamma(b-1)}{\Gamma\left(b+\frac{1}{2}\right)}\,{}_2F_1\left(1,b;b+\frac{1}{2};1-z\right)$$

$$\mathrm{Disc}\phi(w,b) = -2i\sin\,(\pi b)(\Delta_1(w,b) + \Delta_2(w,b))$$

$$\Delta_1(w,b)=\frac{2w}{\pi^{3/2}}\left[-\frac{(w^2-1)^{b-\frac{1}{2}}}{\Gamma\left(b+\frac{1}{2}\right)}\,{}_2F_1\left(1,b;b+\frac{1}{2};1-w^2\right)+\frac{2(w^2-1)^{b-\frac{3}{2}}}{\Gamma\left(b-\frac{1}{2}\right)}\right]$$

$$\Delta_2(w,b)=\frac{2w}{\pi^{3/2}}\left[2\frac{(4w^2-1)^{b-\frac{1}{2}}}{\Gamma\left(b+\frac{1}{2}\right)}\,{}_2F_1\left(1,b;b+\frac{1}{2};1-4w^2\right)-\frac{4(4w^2-1)^{b-\frac{3}{2}}}{\Gamma\left(b-\frac{1}{2}\right)}\right].$$

$$\mathcal{S}_{\text{med}}\left[\mathcal{T}^{(p)}(b)\right]=\mathcal{S}_{\pm}\left[\mathcal{T}^{(p)}(b)\right]\pm i\sin(\pi b)\left(\mathcal{T}_1^{(\text{np})}(b,\lambda)+\mathcal{T}_2^{(\text{np})}(b,\lambda)\right)$$

$$\mathcal{T}_1^{(\text{np})}(b,\lambda)=\sqrt{\lambda}\int_1^{\infty}\frac{dw}{4\sinh^2(\sqrt{\lambda}w)}\Delta_1(w,b)$$

$$\mathcal{T}_2^{(\text{np})}(b,\lambda)=\sqrt{\lambda}\int_{\frac{1}{2}}^{\infty}\frac{dw}{4\sinh^2(\sqrt{\lambda}w)}\Delta_2(w,b)$$

$$\mathcal{T}(b)=\mathcal{S}_{\pm}\left[\mathcal{T}^{(p)}(b)\right]+\sigma(b)\left(\mathcal{T}_1^{(\text{np})}(b,\lambda)+\mathcal{T}_2^{(\text{np})}(b,\lambda)\right)$$

$$\mathcal{T}^{(np)}\equiv\mathcal{T}_1^{(np)}+\mathcal{T}_2^{(np)}$$

$$\mathcal{T}_i^{(np)}\equiv\mathcal{T}_i^{(np)}(2,\lambda)\;i=1,2$$

$$\mathcal{T}_1^{(np)}=\frac{4\sqrt{\lambda}}{\pi^2}\sum_{n=1}^{\infty}n\mathrm{e}^{-2n\sqrt{\lambda}}\int_0^{\infty}dw\mathrm{e}^{-2nw\sqrt{\lambda}}\left((w+1)\sqrt{w(w+2)}+\sinh^{-1}\left(\sqrt{w(w+2)}\right)\right)$$

$$\int_0^{\infty}\mathrm{e}^{-2\sqrt{\lambda}nw}w^z dw=\frac{\Gamma(z+1)}{(2n\sqrt{\lambda})^{z+1}}$$

$$\mathcal{T}_1^{(np)}=\frac{2}{\pi^2}\sqrt{\lambda}\sum_{n=1}^{\infty}n\mathrm{e}^{-2n\sqrt{\lambda}}\sum_{k=0}^{\infty}\frac{(-1)^k(4k^2+3)\Gamma\left(k-\frac{3}{2}\right)}{2^{k+\frac{1}{2}}\pi^{5/2}(2k+1)\Gamma(k+1)}\frac{\Gamma\left(k+\frac{3}{2}\right)}{(n\sqrt{\lambda})^{k+\frac{3}{2}}}.$$

$$\mathcal{T}_2^{(np)}=-\frac{2}{\pi^2}\sqrt{\lambda}\sum_{n=1}^{\infty}n\mathrm{e}^{-n\sqrt{\lambda}}\sum_{k=0}^{\infty}\frac{(-1)^k(4k^2+3)\Gamma\left(k-\frac{3}{2}\right)}{2^{k+\frac{1}{2}}\pi^{5/2}(2k+1)\Gamma(k+1)}\frac{\Gamma\left(k+\frac{3}{2}\right)}{(n\sqrt{\lambda})^{k+\frac{3}{2}}}$$

$$\mathcal{T}^{(np)}=-\frac{2\sqrt{2}}{\pi^2}\sum_{n=0}^{\infty}\mathrm{e}^{-(2n+1)\sqrt{\lambda}}\sum_{k=0}^{\infty}\frac{(-1)^k\left(k^2+\frac{3}{4}\right)\Gamma\left(k-\frac{3}{2}\right)}{2^k\sqrt{\pi}\Gamma(k+1)}\frac{\Gamma\left(k+\frac{1}{2}\right)}{((2n+1)\sqrt{\lambda})^{k+1/2}}.$$

$$\widehat{\mathcal{K}}^{(0)}=4\pi\int_0^{\infty}dt\frac{\mathrm{e}^t t^2}{(1-\mathrm{e}^t)^2}\frac{1}{\pi^2+t^2}\mathcal{B}(2t)$$

$$\widehat{\mathcal{K}}^{(0)}=2\pi\int_0^{\infty}dt\frac{\mathrm{e}^{t/2}t^2}{(1-\mathrm{e}^{t/2})^2}\frac{1}{4\pi^2+t^2}\mathcal{B}(t)$$

$$\partial_\lambda\widehat{\mathcal{M}}^{(0)}=\frac{2}{\lambda}\int_0^{\infty}dt\frac{\mathrm{e}^{t/2}t}{(1-\mathrm{e}^{t/2})^2}J_2\left(\frac{t\sqrt{\lambda}}{2\pi}\right)$$

$$\partial_\lambda \widehat{\mathcal{M}}^{(0)} = \frac{2}{\lambda}\int_0^\infty dt \sum_{n=1}^\infty n\mathrm{e}^{-nt/2} t J_2(at)$$
$$= \frac{2}{\lambda}\sum_{n=1}^\infty \left[\frac{2n}{a^2} - \frac{4n^2}{(n^2+4a^2)^{3/2}} - \frac{2}{a^2}\frac{n^2}{\sqrt{n^2+4a^2}}\right]$$

$$\frac{2}{\lambda}\sum_{n=0}^\infty \frac{2n}{a^2} = -\frac{4\pi^2}{3\lambda^2}$$

$$-\frac{8}{\lambda}\sum_{n=1}^\infty \frac{n^2}{(n^2+4a^2)^{3/2}} = 4\sum_{n=1}^\infty \left[-\frac{2}{\lambda}\frac{n^2}{(n^2+\tilde{a}^2)^{3/2}}\right]$$

$$-\frac{8}{\lambda}\sum_{n=1}^\infty \frac{n^2}{(n^2+4a^2)^{3/2}} = -\frac{4}{\lambda}\int_0^\infty \frac{dt}{t}\mathrm{e}^{-4a^2 t} - \frac{16}{\lambda}\sum_{k=1}^\infty [K_0(4ak\pi) - (4ak\pi)K_1(4ak\pi)]$$

$$-\frac{4}{\lambda a^2}\sum_{n=1}^\infty \frac{n^2}{\sqrt{n^2+4a^2}} = -\frac{16}{\lambda\tilde{a}^2}\sum_{n=1}^\infty \frac{n^2}{\sqrt{n^2+\tilde{a}^2}} = 4\left[-\frac{4}{\lambda\tilde{a}^2}\sum_{n=1}^\infty \frac{n^2}{\sqrt{n^2+\tilde{a}^2}}\right]$$

$$\partial_\lambda \widehat{\mathcal{M}}^{(0)} = \frac{4}{\lambda} - \frac{4\pi^2}{3\lambda^2} + \frac{16}{\lambda}\sum_{k=1}^\infty \left[\frac{(2\lambda k^2+1)K_1(2k\sqrt{\lambda})}{k\sqrt{\lambda}} + K_0(2k\sqrt{\lambda})\right]$$

$$\partial_\lambda \widehat{\mathcal{K}}^{(0)} = \frac{2I_1(\sqrt{\lambda})}{\sqrt{\lambda}}\left[1 + 4\sqrt{\lambda}\sum_{k=1}^\infty \left(kK_1(2k\sqrt{\lambda}) - 2k^2\sqrt{\lambda}K_0(2k\sqrt{\lambda})\right)\right].$$

$$\widehat{\mathcal{K}}^{(0)}(\lambda) = \int_0^\lambda dq\,\partial_q \widehat{\mathcal{K}}^{(0)}(q)$$
$$= 4\big(I_0(\sqrt{\lambda}) - 1\big) + 4\sum_{k=1}^\infty (\mathcal{J}_1(2k,\lambda) + \mathcal{J}_0(2k,\lambda)) + (4-\pi^2)$$
$$= 4I_0(\sqrt{\lambda}) + 4\sum_{k=1}^\infty (\mathcal{J}_1(2k,\lambda) + \mathcal{J}_0(2k,\lambda)) - \pi^2$$

$$\mathcal{K}^{(0)}(\lambda) = I_0(\sqrt{\lambda}) + \int_0^\lambda dq \lim_{N\to\infty}\left(\sum_{k=1}^N f_k(q)\right)$$

$$f_k(q) = I_1(\sqrt{q})\big(kK_1(k\sqrt{q}) - k^2\sqrt{q}K_0(k\sqrt{q})\big)$$

$$\lim_{q\to 0^+} f_k(q) = \frac{1}{2},\ \lim_{q\to\infty} f_k(q) \sim \mathrm{e}^{(1-k)\sqrt{q}}$$

$$S_N(q) \equiv \sum_{k=2}^N f_k(q)$$

$$|S_N(q)| \leq g(q)\ \forall N,\ \text{for almost all } q \in [0,\infty)$$

$$S_N(0) = \frac{N-1}{2}$$

$$\lim_{N\to\infty} S_N(\epsilon) = -\frac{3}{4}$$

$$g(q) = \begin{cases} \mathcal{C}, & 0 < q \leq 1 \\ \dfrac{2}{\mathrm{e}^q - 1}, & 1 < q < \infty \end{cases}$$

$$\varphi(z) = \sum_{n\geq 0} a_n z^n$$

$$\hat{\varphi}(t) \underset{t\to A}{\simeq} \frac{\mathcal{S}a}{2\pi}\frac{1}{t-A}.$$

$$\left(\frac{\mathcal{S}a}{2\pi}\right)\lim_{\epsilon\to 0}\left[\int_0^{\infty+i\epsilon}\frac{dt\mathrm{e}^{-zt}}{t-A} - \int_0^{\infty-i\epsilon}\frac{dt\mathrm{e}^{-zt}}{t-A}\right]$$

$$\lim_{\epsilon\to 0}\frac{1}{(x\pm i\epsilon)^n} = \mathcal{P}\left(\frac{1}{x^n}\right) \mp \frac{i\pi(-1)^{n-1}}{(n-1)!}\delta^{(n-1)}(x)$$

$$\lim_{\epsilon\to 0}\left[\mathrm{e}^{-i\epsilon z}\int_0^{\infty}\frac{dt\mathrm{e}^{-zt}}{t-A+i\epsilon} - \mathrm{e}^{i\epsilon z}\int_0^{\infty}\frac{dt\mathrm{e}^{-zt}}{t-A-i\epsilon}\right]$$

$$= \int_0^{\infty} dt\mathrm{e}^{-zt}\left[\mathcal{P}\left(\frac{1}{t-A}\right) - i\pi\delta(t-A) - \mathcal{P}\left(\frac{1}{t-A}\right) - i\pi\delta(t-A)\right]$$

$$= -2\pi i\int_0^{\infty} dt\mathrm{e}^{-zt}\delta(t-A) = -2\pi i\mathrm{e}^{-Az}$$

$$\hat{\varphi}(t) \underset{t\to A}{\simeq} \frac{\mathcal{S}a}{2\pi}\frac{iaz\mathcal{S}\mathrm{e}^{-Az}}{(t-A)^2} - i\mathcal{S}a\mathrm{e}^{-Az},$$

$$a_{\frac{\mathrm{e}^t}{(\mathrm{e}^t+1)^2}} = \sum_{b=1}^{N(2N+1)} a_b T^b, da = \prod_{b=1}^{N(2N+1)}\frac{da_b}{\sqrt{2\pi}}$$

$$\hat{H} = J_s\sum_{\langle \boldsymbol{i},\boldsymbol{j}\rangle_{\text{square}}} \hat{\mathbf{S}}_{\boldsymbol{i}}\cdot\hat{\mathbf{S}}_{\boldsymbol{j}} + J_d\sum_{\langle \boldsymbol{i},\boldsymbol{j}\rangle_{\text{dimer}}} \hat{\mathbf{S}}_{\boldsymbol{i}}\cdot\hat{\mathbf{S}}_{\boldsymbol{j}}.$$

$$\mathbf{s}_i = \frac{1}{2}f_{i\alpha}^{\dagger}\boldsymbol{\sigma}_{\alpha\beta}f_{i\beta}.$$

$$\psi_i = \begin{pmatrix} f_{i\uparrow} \\ f_{i\downarrow}^{\dagger} \end{pmatrix},$$

$$H = -\sum_{ij}\psi_i^{\dagger}u_{ij}\psi_j.$$

$$u_{ij} = iu_{ij}^0\sigma^0 + u_{ij}^a\sigma^a,$$

$$u_{ji}^0 = -u_{ij}^0, u_{ji}^a = u_{ij}^a.$$

$$\mathrm{SU}(2)_g: \psi_i \to U_{g,g(i)}\psi_{g(i)}$$
$$u_{ij} \to U_{g,g(i)}u_{g(i),g(j)}U_{g,g(j)}^\dagger$$

$$u_{\mathbf{r},A,A,\mathbf{r},B,A} = \begin{bmatrix} -te^{-i\theta} & 0 \\ 0 & te^{i\theta} \end{bmatrix} = u_s$$
$$u_{\mathbf{r},A,A,\mathbf{r},A,B} = \begin{bmatrix} te^{i\theta} & 0 \\ 0 & -te^{-i\theta} \end{bmatrix} = -u_s^\dagger$$
$$u_{\mathbf{r},A,B,\mathbf{r},B,B} = u_{\mathbf{r},B,B,\mathbf{r},B,A} = u_{\mathbf{r},B,A,\mathbf{r}+\hat{\mathbf{x}},A,A} = -u_s^\dagger$$
$$u_{\mathbf{r},A,B,\mathbf{r}+\hat{\mathbf{y}},A,A} = u_{\mathbf{r}+\hat{\mathbf{y}},B,A,\mathbf{r},B,B} = u_{\mathbf{r},B,B,\mathbf{r}+\hat{\mathbf{x}},A,B} = u_s$$
$$u_{\mathbf{r},A,A,\mathbf{r},B,B} = \Delta_d\sigma^x, u_{\mathbf{r}+\hat{\mathbf{x}},A,B,\mathbf{r}+\hat{\mathbf{y}},B,A} = \Delta_d\sigma^y$$
$$u_{\mathbf{r},B,B,\mathbf{r}+\hat{\mathbf{x}}+\hat{\mathbf{y}},A,A} = \Delta_{d'}\sigma^x, u_{\mathbf{r},B,A,\mathbf{r},A,B} = \Delta_{d'}\sigma^y$$
$$u_{\mathbf{r},B,A,\mathbf{r}+\hat{\mathbf{x}},A,B} = -\Delta_{g,2}\sigma^x + \Delta_{g,1}\sigma^y$$
$$u_{\mathbf{r},A,B,\mathbf{r}+\hat{\mathbf{y}},B,A} = \Delta_{g,2}\sigma^x + \Delta_{g,1}\sigma^y$$
$$u_{\mathbf{r}+\hat{\mathbf{x}},A,A,\mathbf{r},B,B} = \Delta_{g,1}\sigma^x + \Delta_{g,2}\sigma^y$$
$$u_{\mathbf{r},B,B,\mathbf{r}+\hat{\mathbf{y}},A,A} = \Delta_{g,1}\sigma^x - \Delta_{g,2}\sigma^y$$

$$\mathcal{F}_i = \begin{pmatrix} f_{i\uparrow} & -f_{i\downarrow} \\ f_{i\downarrow}^\dagger & f_{i\uparrow}^\dagger \end{pmatrix}$$

$$\mathcal{F}_i^\dagger = \sigma^y\mathcal{F}_i^T\sigma^y$$

$$\mathcal{F}_i \to U_{g,g(i)}\mathcal{F}_{g(i)}$$

$$\mathcal{F}_i \to \mathcal{F}_i\sigma^z\Omega_i^T\sigma^z$$

$$H = \sum_{\langle ij\rangle} i\alpha_{ij}\mathrm{Tr}\left[\mathcal{F}_i^\dagger\mathcal{F}_j\right] + \beta_{ij}^a\mathrm{Tr}\left[\sigma^a\mathcal{F}_i^\dagger\mathcal{F}_j\right] + i\gamma_{ij}\mathrm{Tr}\left[\sigma^a\mathcal{F}_i^\dagger\sigma^a\mathcal{F}_j\right]$$

$$u_{ij} = i\alpha_{ij}\sigma^0 + \beta_{ij}^a\sigma^a$$

$$\alpha_{ij} = -\alpha_{ji}, \alpha_{i,i+\hat{\mathrm{x}}} = t, \alpha_{i,i+\hat{\mathrm{y}}} = (-1)^{i_x}t$$

$$H = -2t\sum_k \mathrm{Tr}\left[\mathcal{F}_k^\dagger\left(\sin\left(k_x/2\right)\rho^x + \sin\left(k_y/2\right)\rho^z\kappa^x\right)\mathcal{F}_k\right]$$

$$\mathcal{F}_{\mathbf{r},m_x,m_y} = \sum_{m_x'} \rho_{m_x,m_x'}^x\mathcal{X}_{\mathbf{r},m_x',0} + \kappa_{m_y,m_y}^z\mathcal{X}_{\mathbf{r},m_x,1}$$

$$H \approx it\sum_v \mathrm{Tr}\left[\mathcal{X}_v^\dagger\left(\rho^x\partial_x - \rho^z\partial_y\right)\mathcal{X}_v\right]$$

$$\mathcal{L}_{\mathrm{MF}} = i\mathrm{Tr}\left[\overline{\mathcal{X}}\gamma^\mu\partial_\mu\mathcal{X}\right]$$

$\overline{\mathcal{X}} = \mathcal{X}^\dagger\gamma^0$ and the gamma matrices $\gamma^0 = \rho^y, \gamma^x = i\rho^z, \gamma^y = i\rho^x$

$$L^T = \sigma^y L^\dagger \sigma^y$$

Lie generators $M = M^\dagger, L = e^{iM}$, this is equivalent to

$$M^T = -\sigma^y M \sigma^y$$

$$T^j = \{\mu^y, \sigma^a, \mu^x\sigma^a, \mu^z\sigma^a\}$$

$$\Gamma^j = \{\mu^x, \mu^z, \mu^y\sigma^a\}$$

$$\beta^z_{i,i+\hat{\mathbf{x}}} = -\delta\theta(-1)^{i_x+i_y}, \beta^z_{i,i+\hat{\mathbf{y}}} = \delta\theta(-1)^{i_y}$$

$$\begin{aligned}
\delta H =& -\delta\theta\sum_{\boldsymbol{r}}\left[\mathcal{X}^\dagger_{\boldsymbol{r},0}\rho^x + \mathcal{X}^\dagger_{\boldsymbol{r},1}\kappa^z\right]\rho^x\kappa^z\left[\rho^x\mathcal{X}_{\boldsymbol{r},0} + \kappa^z\mathcal{X}_{\boldsymbol{r},1}\right] \\
&+\frac{1}{2}\delta\theta\sum_{\boldsymbol{r}}\left[\mathcal{X}^\dagger_{\boldsymbol{r},0}\rho^x + \mathcal{X}^\dagger_{\boldsymbol{r},1}\kappa^z\right](\rho^x - i\rho^y)\kappa^z\left[\rho^x\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{x}},0} + \kappa^z\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{x}},1}\right] \\
&+\frac{1}{2}\delta\theta\sum_{\boldsymbol{r}}\left[\mathcal{X}^\dagger_{\boldsymbol{r},0}\rho^x + \mathcal{X}^\dagger_{\boldsymbol{r},1}\kappa^z\right](\rho^x + i\rho^y)\kappa^z\left[\rho^x\mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{x}},0} + \kappa^z\mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{x}},1}\right] \\
\approx& 2\delta\theta\int \mathrm{d}^2\boldsymbol{r}\left[\mathcal{X}^\dagger_{\boldsymbol{r},0}\rho^z\partial_x\mathcal{X}_{\boldsymbol{r},1} - \mathcal{X}^\dagger_{\boldsymbol{r},1}\rho^z\partial_x\mathcal{X}_{\boldsymbol{r},0}\right] \\
\Rightarrow \delta\mathcal{L} =& -2i\delta\theta\mathrm{Tr}\left[\sigma^z\overline{\mathcal{X}}\mu^y\gamma^y\partial_x\mathcal{X}\right]
\end{aligned}$$

$$\begin{aligned}
\delta H =& +\delta\theta\sum_{\boldsymbol{r}}\left[\mathcal{X}^\dagger_{\boldsymbol{r},0}\rho^x + \mathcal{X}^\dagger_{\boldsymbol{r},1}\kappa^z\right]\kappa^x\left[\rho^x\mathcal{X}_{\boldsymbol{r},0} + \kappa^z\mathcal{X}_{\boldsymbol{r},1}\right] \\
&-\frac{1}{2}\delta\theta\sum_{\boldsymbol{r}}\left[\mathcal{X}^\dagger_{\boldsymbol{r},0}\rho^x + \mathcal{X}^\dagger_{\boldsymbol{r},1}\kappa^z\right](\kappa^x - i\kappa^y)\left[\rho^x\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{y}},0} + \kappa^z\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{y}},1}\right] \\
&-\frac{1}{2}\delta\theta\sum_{\boldsymbol{r}}\left[\mathcal{X}^\dagger_{\boldsymbol{r},0}\rho^x + \mathcal{X}^\dagger_{\boldsymbol{r},1}\kappa^z\right](\kappa^x + i\kappa^y)\left[\rho^x\mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{y}},0} + \kappa^z\mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{y}},1}\right] \\
\approx& 2\delta\theta\int \mathrm{d}^2\boldsymbol{r}\left[\mathcal{X}^\dagger_{\boldsymbol{r},0}\rho^z\partial_x\mathcal{X}_{\boldsymbol{r},1} - \mathcal{X}^\dagger_{\boldsymbol{r},1}\rho^z\partial_x\mathcal{X}_{\boldsymbol{r},0}\right] \\
\Rightarrow \delta\mathcal{L} =& -2i\delta\theta\mathrm{Tr}\left[\sigma^z\overline{\mathcal{X}}\mu^y\gamma^x\partial_y\mathcal{X}\right]
\end{aligned}$$

$$\delta\mathcal{L} = \Phi^a_3\mathrm{Tr}\left[\sigma^a\overline{\mathcal{X}}\mu^y\left(\gamma^y i\partial_x + \gamma^x i\partial_y\right)\mathcal{X}\right]$$

$$\begin{aligned}
\delta H =& \Delta_d\mathrm{Tr}\Bigg\{\sum_{\boldsymbol{r}}\sigma^x\left[\mathcal{X}^\dagger_{\boldsymbol{r},B,0} + \mathcal{X}^\dagger_{\boldsymbol{r},A,1}\right]\left[\mathcal{X}_{\boldsymbol{r},A,0} - \mathcal{X}_{\boldsymbol{r},B,1}\right] + \text{h.c.} \\
&+\sigma^y\left[\mathcal{X}^\dagger_{\boldsymbol{r},B,0} - \mathcal{X}^\dagger_{\boldsymbol{r},A,1}\right]\left[\mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{x}}+\hat{\mathbf{y}},A,0} + \mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{x}}+\hat{\mathbf{y}},B,1}\right] + \text{h.c.}\Bigg\} \\
\approx& \Delta_d\int \mathrm{d}^2\boldsymbol{r}\mathrm{Tr}\left\{\sigma^x\left(\mathcal{X}^\dagger\rho^x\mu^z\mathcal{X} + \mathcal{X}^\dagger\rho^z\mu^x\mathcal{X}\right)\right\} + \mathrm{Tr}\left\{\sigma^y\left(\mathcal{X}^\dagger_{\boldsymbol{r}}\rho^x\mu^z\mathcal{X}_{\boldsymbol{r}} - \mathcal{X}^\dagger_{\boldsymbol{r}}\rho^z\mu^x\mathcal{X}_{\boldsymbol{r}}\right)\right\} \\
\Rightarrow \delta\mathcal{L} =& \Delta_d\mathrm{Tr}\left[\sigma^x\overline{\mathcal{X}}(\gamma^x\mu^z - \gamma^y\mu^x)\mathcal{X}\right] + \Delta_d\mathrm{Tr}\left[\sigma^y\overline{\mathcal{X}}(\gamma^x\mu^z + \gamma^y\mu^x)\mathcal{X}\right]
\end{aligned}$$

$$\begin{aligned}
\delta\mathcal{L} =& \left(\Delta_d + \Delta_{d'} + 2\Delta_{g,1}\right)\left(\mathrm{Tr}\left[\sigma^x\overline{\mathcal{X}}(\gamma^x\mu^z - \gamma^y\mu^x)\mathcal{X}\right] + \mathrm{Tr}\left[\sigma^y\overline{\mathcal{X}}(\gamma^x\mu^z + \gamma^y\mu^x)\mathcal{X}\right]\right) \\
&+\Delta_{g,2}\left(\mathrm{Tr}\left[\sigma^x\overline{\mathcal{X}}(-\gamma^0\mu^y + 1)\left(i\partial_x + i\partial_y\right)\mathcal{X}\right] + \mathrm{Tr}\left[\sigma^y\overline{\mathcal{X}}(\gamma^0\mu^y + 1)\left(i\partial_y - i\partial_x\right)\mathcal{X}\right]\right)
\end{aligned}$$

$$\Phi^a_1\mathrm{Tr}\left[\sigma^a\overline{\mathcal{X}}\gamma^x\mu^z\mathcal{X}\right] + \Phi^a_2\mathrm{Tr}\left[\sigma^a\overline{\mathcal{X}}\gamma^y\mu^x\mathcal{X}\right]$$

$$\Phi_1^a \mathrm{Tr}\left[\sigma^a \overline{\mathcal{X}}\left(\gamma^x \mu^z - \delta\left(\gamma^0 \mu^y i\partial_x - i\partial_y\right)\right)\mathcal{X}\right]$$

$$\Phi_2^a \mathrm{Tr}\left[\sigma^a \overline{\mathcal{X}}\left(\gamma^y \mu^x + \delta\left(\gamma^0 \mu^y i\partial_y - i\partial_x\right)\right)\mathcal{X}\right]$$

$$\delta \equiv \Delta_{g,2}/\left(\Delta_d + \Delta_{d'} + 2\Delta_{g,1}\right)$$

$$\langle \Phi_1 \rangle \propto \left(\Delta_d + \Delta_{d'} + 2\Delta_{g,1}\right)(1,1,0)$$
$$\langle \Phi_2 \rangle \propto \left(\Delta_d + \Delta_{d'} + 2\Delta_{g,1}\right)(-1,1,0)$$

$$\begin{aligned}\mathcal{L} =& i\mathrm{Tr}\left[\overline{\mathcal{X}}\gamma^\mu \partial_\mu \mathcal{X}\right] + \Phi_1^a \mathrm{Tr}\left[\sigma^a \overline{\mathcal{X}}\left(\gamma^x \mu^z - \delta\left(\gamma^0 \mu^y i\partial_x - i\partial_y\right)\right)\mathcal{X}\right] \\ &+ \Phi_2^a \mathrm{Tr}\left[\sigma^a \overline{\mathcal{X}}\left(\gamma^y \mu^x + \delta\left(\gamma^0 \mu^y i\partial_y - i\partial_x\right)\right)\mathcal{X}\right] \\ &+ \Phi_3^a \mathrm{Tr}\left[\sigma^a \overline{\mathcal{X}} \mu^y \left(\gamma^y i\partial_x + \gamma^x i\partial_y\right)\mathcal{X}\right] + V(\Phi)\end{aligned}$$

$$\begin{aligned}V(\Phi) =& s(\Phi_1^a \Phi_1^a + \Phi_2^a \Phi_2^a) + \tilde{s}\Phi_3^a \Phi_3^a + w\epsilon_{abc}\Phi_1^a \Phi_2^b \Phi_3^c \\ &+ u(\Phi_1^a \Phi_1^a + \Phi_2^a \Phi_2^a)^2 + \tilde{u}(\Phi_3^a \Phi_3^a)^2 + v_1(\Phi_1^a \Phi_2^a)^2 \\ &+ v_2(\Phi_1^a \Phi_1^a)\left(\Phi_2^b \Phi_2^b\right) + v_3[(\Phi_1^a \Phi_3^a)^2 + (\Phi_2^a \Phi_3^a)^2] \\ &+ v_4(\Phi_1^a \Phi_1^a + \Phi_2^a \Phi_2^a)\left(\Phi_3^b \Phi_3^b\right)\end{aligned}$$

$$\begin{aligned}T_x&: (i_x, i_y) \mapsto (i_x + 1, i_y), \\ T_y&: (i_x, i_y) \mapsto (i_x, i_y + 1), \\ P_x&: (i_x, i_y) \mapsto (-i_x, i_y), \\ P_y&: (i_x, i_y) \mapsto (i_x, -i_y), \\ R_{\pi/2}&: (i_x, i_y) \mapsto (-i_y, i_x).\end{aligned}$$

$$\begin{aligned}T_{2x}&: (i_x, i_y) \mapsto (i_x + 2, i_y), \\ T_{2y}&: (i_x, i_y) \mapsto (i_x, i_y + 2), \\ G_x&: (i_x, i_y) \mapsto (i_x + 1, -i_y), \\ G_y&: (i_x, i_y) \mapsto (-i_x, i_y + 1), \\ \sigma_{xy}&: (i_x, i_y) \mapsto (i_y, i_x), \\ \sigma_{x\bar{y}}&: (i_x, i_y) \mapsto (-i_y + 1, -i_x + 1), \\ C_4&: (i_x, i_y) \mapsto (-i_y + 2, i_x - 1).\end{aligned}$$

$$\begin{aligned}T_x&: \mathcal{X} \to \mu^x \mathcal{X} \\ T_y&: \mathcal{X} \to \mu^z \mathcal{X} \\ P_x&: \mathcal{X} \to -\rho^z \mu^z \mathcal{X}(-x, y) \\ P_y&: \mathcal{X} \to \rho^x \mu^x \mathcal{X}(x, -y) \\ \mathcal{T}&: \mathcal{X} \to \rho^y \mu^y \mathcal{X}, i \to -i \\ R_{\pi/2}&: \mathcal{X} \to e^{-i\pi\rho^y/4} e^{i\pi\mu^y/4} \mathcal{X}(-y, x)\end{aligned}$$

$$G_x = P_y \circ T_x, \sigma_{xy} = P_x \circ R_{\pi/2}$$

$$\begin{aligned}G_x&: \mathcal{X} \to \rho^x \mathcal{X}(x, -y) \\ \mathcal{T}&: \mathcal{X} \to \rho^y \mu^y \mathcal{X}, i \to -i \\ \sigma_{xy}&: \mathcal{X} \to -\rho^z \mu^z e^{-i\pi\rho^y/4} e^{i\pi\mu^y/4} \mathcal{X}(y, x)\end{aligned}$$

$\mu^y\gamma^\mu i\partial_v$	G_x	$\mathcal{T}$	σ_{xy}	$\mu^{0,x,y}i\partial_\mu$	G_x	$\mathcal{T}$	σ_{xy}
$\mu^y\gamma^0 i\partial_0$	+	+	-	$i\partial_0$	-	+	-
$\mu^y\gamma^0 i\partial_x$	+	-	$-\mu^y\gamma^0 i\partial_y$	$i\partial_x$	-	-	$-i\partial_y$
$\mu^y\gamma^0 i\partial_y$	-	-	$-\mu^y\gamma^0 i\partial_x$	$i\partial_y$	+	-	$-i\partial_x$
$\mu^y\gamma^x i\partial_0$	+	-	$-\mu^y\gamma^y i\partial_0$	$\mu^x i\partial_0$	-	-	$-\mu^z i\partial_0$
$\mu^y\gamma^x i\partial_x$	+	+	$-\mu^y\gamma^y i\partial_y$	$\mu^x i\partial_x$	-	+	$-\mu^z i\partial_y$
$\mu^y\gamma^x i\partial_y$	-	+	$-\mu^y\gamma^y i\partial_x$	$\mu^x i\partial_y$	+	+	$-\mu^z i\partial_x$
$\mu^y\gamma^y i\partial_0$	-	-	$-\mu^y\gamma^x i\partial_0$	$\mu^z i\partial_0$	-	-	$-\mu^x i\partial_0$
$\mu^y\gamma^y i\partial_x$	-	+	$-\mu^y\gamma^x i\partial_y$	$\mu^z i\partial_x$	-	+	$-\mu^x i\partial_y$
$\mu^y\gamma^y i\partial_y$	+	+	$-\mu^y\gamma^x i\partial_x$	$\mu^z i\partial_y$	+	+	$-\mu^x i\partial_x$

Transformation of $\mathrm{Tr}\left[\sigma^a\overline{\mathcal{X}}i\partial_\mu\mathcal{X}\right]$, $\mathrm{Tr}\left[\sigma^a\overline{\mathcal{X}}\Gamma^j i\partial_\mu\mathcal{X}\right]$, and $\mathrm{Tr}\left[\sigma^a\overline{\mathcal{X}}T^j\gamma^\mu i\partial_\nu\mathcal{X}\right]$

$\langle\Phi_3^a\rangle \propto w\epsilon_{abc}\langle\Phi_1^b\rangle\langle\Phi_2^c\rangle$ condenses

$$\psi_{a,m_x,v} = i\sigma_{ab}^y\left[\mathcal{X}_{m_x,v}\right]_{1,b}$$

$$\mathcal{L} = \mathcal{L}_\psi + \mathcal{L}_\Phi + \mathcal{L}_{\Phi\psi}$$
$$\mathcal{L}_\psi = i\bar{\psi}_v\gamma^\mu\left(\partial_\mu - iA_\mu^a\sigma^a\right)\psi_v$$
$$\mathcal{L}_\Phi = \frac{1}{2g}\left[\left(\partial_x\Phi_1^a - 2\epsilon_{abc}A_x^b\Phi_1^c\right)^2 + \left(\partial_y\Phi_2^a - 2\epsilon_{abc}A_y^b\Phi_2^c\right)^2\right] + \cdots$$
$$\mathcal{L}_{\Phi\psi} = y\sum_\alpha\left(\Phi_{1\alpha}^a\bar{\psi}\mu^z\gamma^x\sigma^a\psi + \Phi_{2\alpha}^a\bar{\psi}\mu^x\gamma^y\sigma^a\psi\right)$$

$$\mathcal{L}_\psi = i\bar{\psi}_v\gamma^\mu\left(\partial_\mu - iA_\mu^a\sigma^a\right)\psi_v$$
$$\mathcal{L}_\Phi = \frac{1}{2g}\left[\left(\delta_{ac}\partial_\mu - 2\epsilon_{abc}A_\mu^b\right)\Phi_{s\alpha}^c\right]^2$$
$$\mathcal{L}_{\Phi\psi} = y\sum_{s,\alpha}\Phi_{s\alpha}^a\bar{\psi}X^s\sigma^a\psi$$

$$\sum_{s,\alpha}(\Phi_{s\alpha}^a)^2 = 1$$

$$\Phi^a_{s\alpha} \to \sqrt{\frac{g}{3N_b}}\,\Phi^a_{s\alpha}$$

$$\mathcal{L}_\Phi = \frac{1}{2g}\left[\left(\left(\delta_{ac}\partial_\mu - 2\epsilon_{abc}A^b_\mu\right)\Phi^c_{s\alpha}\right)^2 + i\lambda\left((\Phi^a_{s\alpha})^2 - \frac{3N_b}{g}\right)\right]$$

$$S_b = \frac{3}{g}\int \mathrm{d}^{\mathcal{D}}r\left(-\frac{N_b i\lambda}{2g} - \frac{N_b}{2}\log\det\left|-\partial_\mu^2 + i\lambda\right|\right)$$

$$\frac{N_b i}{2g} = -\frac{N_b}{2}\frac{\delta}{\delta\lambda}\log\det\left|-\partial_\mu^2 + i\lambda\right| = -\frac{N_b i}{2}G_\Phi(x,x),$$

$$G_\Phi = \left(-\partial_\mu^2 + i\lambda\right)^{-1}.$$

$$G_\Phi^{-1} = k^2 + i\lambda$$

$$\frac{1}{g} = \int \frac{\mathrm{d}^{\mathcal{D}}k}{8\pi^3}\frac{1}{k^2 + i\lambda}$$

$$G_\Phi(k) = (k^2 + r)^{-1}$$

$$\mathcal{F} = \frac{1}{2}\int \frac{\mathrm{d}^{\mathcal{D}}p}{(2\pi)^3}\left(\Pi_\lambda\lambda^2 + \Pi_A\left(\delta_{\mu v} - \frac{p_\mu p_v}{p^2}\right)A^a_\mu A^a_v\right)$$

$$\Pi_\lambda(p,r) = \frac{3N_b}{4\pi p}\arctan\frac{p}{2\sqrt{r}}$$

$$\Pi_A(p,r) = N_f\frac{p}{16} + 8N_b\left[\frac{p^2 + 4r}{8\pi p}\arctan\frac{p}{2\sqrt{r}} - \frac{\sqrt{r}}{4\pi}\right]$$

$$k_\mu A_\mu = 1 - \zeta$$


$$D_\lambda = \langle\lambda\lambda\rangle = \frac{1}{\Pi_\lambda}$$

$$D^{ab}_{\mu v} = \left\langle A^a_\mu A^b_v\right\rangle = \frac{\delta_{ab}}{\Pi_A}\left(\delta_{\mu v} - \zeta\frac{p_\mu p_v}{p^2}\right)$$

$$G^{ab}_\Phi = \frac{\delta_{ab}}{k^2 + r}$$

$$G_\psi = \frac{k}{k^2}$$

$$\Pi_\lambda(p) = \frac{3N_b}{8p}, \Pi_A(p) = \left(N_f + 8N_b\right)\frac{p}{16}.$$



$$\dim[\psi] = \frac{D-1+\eta_\psi}{2}$$

$$\Sigma_\psi(k) = \sigma^a\sigma^a \int \frac{d^{\mathcal{D}}q}{(2\pi)^3}\gamma_\mu G_\psi(k+q)\gamma_\nu D_{\mu\nu}(-q) \to \frac{8}{\left(N_f+8N_b\right)\pi^2}(1-3\zeta)\not{k}\log\, k.$$

$$\eta_\psi = \frac{8(1-3\zeta)}{\left(N_f+8N_b\right)\pi^2}$$

$$\dim[\psi] = 1+\frac{\eta_\psi}{2} = 1+\frac{4}{\left(N_f+8N_b\right)\pi^2}(1-3\zeta).$$

$$\dim[\Phi] = \frac{D-2+\eta_\Phi}{2}\langle i\mathrm{Tr}[\overline{\mathcal{X}}\Gamma^j\mathcal{X}]\rangle$$

$$I_{A;1} = 8\int \frac{d^{\mathcal{D}}q}{(2\pi)^3}G_\Phi(k+q)D_{\mu\nu}(-q)(2k+q)_\mu(2k+q)_\nu$$
$$\to -\frac{8\cdot 4}{\left(N_f+8N_b\right)\pi^2}\left(\frac{10}{3}+2\zeta\right)k^2\log\, k$$

$$2\epsilon_{abc}\cdot 2\epsilon_{fbc} = 8\delta_{af},$$

$$I_{\lambda;1} = i^2\int \frac{d^{\mathcal{D}}q}{(2\pi)^3}G_\Phi(k+q)D_\lambda(-q) \to \frac{\partial N^z\partial\bar{\psi}\partial\mu^y\partial^2\psi.}{9N_b\pi^2}k^2\log\, k.$$

$$\eta_{\Phi} = -\frac{32}{\left(N_f + 8N_b\right)\pi^2}\left(\frac{10}{3} + 2\zeta\right) + \frac{N^z = \bar{\psi}\mu^y\psi.}{9N_b\pi^2}.$$

$$\Gamma^j = \{\mu^x, \mu^z, \mu^y\sigma^a\}.$$

$$V^i = (\bar{\psi}\mu^x\psi, \bar{\psi}\mu^z\psi).$$

$$I_{A;2} = 3\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\gamma_\mu G_\psi(k_1+q)G_\psi(k_2+q)\gamma_\nu D_{\mu\nu}(-q)$$
$$\to \frac{3\cdot 8\mu^i}{\left(N_f + 8N_b\right)\pi^2}(\zeta - 3)\log\, k$$

$$\dim\left[\bar{\psi}\mu^i\psi\right] = 2\dim[\psi] + \eta_{\mathrm{vrtx}} = 2 - \frac{64}{\left(N_f + 8N_b\right)\pi^2}$$

$$\chi(k) = \left\langle\bar{\psi}\Gamma^i\psi(k)\bar{\psi}\Gamma^j\psi(-k)\right\rangle \sim |k|^{1-\frac{128}{(N_f+8N_b)\pi^2}} \sim |k|^{0.279}$$


$$\chi(r) \sim |r|^{-4+\frac{128}{(N_f+8N_b)\pi^2}} \approx |r|^{-3.279}$$

$$\xi \propto (g - g_c)^{-v}$$

$$v = \frac{\gamma_\Phi}{2 - \eta_\Phi}$$

$$G_\Phi^{-1}(k=0) \sim (g - g_c)^{\gamma_\Phi}.$$

$$\frac{1}{g_c} - \frac{1}{g} = \frac{\sqrt{r_g}}{4\pi}$$



$$r_g \log\left(r_g\right) \text{ of } \Sigma(0,r) - \frac{\Pi_\lambda(0,0)}{\Pi_\lambda(0,r)}\Sigma(0,0)$$

$$\gamma_\Phi = 2(1-\alpha).$$

$$\Sigma^{(a)} = I_{A;1}$$

$$\Sigma^{(b)} = 8\sum_{\mu,v}\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{1}{\Pi_A(q)}\left(\delta_{\mu v} - \zeta\frac{q_\mu q_v}{q^2}\right)$$

$$\Sigma^{(c)} = I_{\lambda;1}$$

$$\Sigma^{(d)} = \frac{i^2}{\Pi_\lambda(0,r)}\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}3N_b G_\Phi(q)^2 I_{A;1}(q)$$

$$\Sigma^{(e)} = \frac{i^2}{\Pi_\lambda(0,r)}\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}3N_b G_\Phi(q)^2 I_{\lambda;1}(q)$$

$$\Sigma^{(f)} = \frac{i^2\Sigma^{(b)}}{\Pi_\lambda(0,r)}\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}3N_b G_\Phi(q)^2 = -\Sigma^{(b)}$$

$$\alpha_\lambda = \frac{2}{N_b\pi^2}.$$

gauge contributions $\Sigma = \Sigma^{(a)} + \Sigma^{(d)}$

$$\Sigma(0,r) - \frac{\Pi(0,0)}{\Pi(0,r)}\Sigma(0,0) = 8\int \frac{q^{\mathcal{D}}\,\mathrm{d}q}{2\pi^2}\left[\frac{1-\zeta}{\Pi_A(q,r_g)}\frac{q^{\mathcal{D}}}{q^{\mathcal{D}}+r_g} - \frac{3N_b}{\Pi_\lambda(0,r_g)}\frac{I_A(q,r_g)}{\Pi_A(q,r_g)} + \frac{3N_b}{\Pi_\lambda(0,r_g)}\frac{1}{4q\Pi_A(q,0)}\right]$$

$$I_A(q,r_g) \approx (1-\zeta)\frac{\Pi_\lambda(0,r_g)}{3N_b} + \frac{1}{4q} - \frac{\sqrt{r_g}}{\pi q^2}.$$

$$\alpha_A = -\frac{8\cdot 4}{\pi^2(N_f+8N_b)}\left(\zeta + \frac{7N_f-8N_b}{N_f+8N_b}\right).$$

$$\gamma_\Phi = 2(1-\alpha) = 2 - \frac{4}{N_b\pi^2} + \frac{64}{\pi^2}\left(\frac{7N_f-8N_b}{(N_f+8N_b)^2} + \frac{\zeta}{N_f+8N_b}\right).$$

$$v = \frac{\gamma_\Phi}{2-\eta_\Phi} \approx \frac{\gamma_\Phi}{2}\left(1+\frac{\eta_\Phi}{2}\right)$$

$$= \left[1 - \frac{2}{N_b\pi^2} + \frac{32}{\pi^2}\left(\frac{7N_f-8N_b}{(N_f+8N_b)^2} + \frac{\zeta}{N_f+8N_b}\right)\right]\left[1 + \frac{2}{9N_b\pi^2} - \frac{16}{(N_f+8N_b)\pi^2}\left(\frac{10}{3}+2\zeta\right)\right]$$

$$\approx 1 - \frac{16}{9N_b\pi^2} + \frac{32}{\pi^2}\frac{7N_f-8N_b}{(N_f+8N_b)^2} - \frac{160}{3(N_f+8N_b)\pi^2}$$

$$I_{\Phi\psi} = (-i\sigma^b)\sigma^c\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\gamma_\mu G_\psi(q)X^s G_\Phi(q)q_v(2\epsilon_{abc})D_{\mu v}(-q)$$

$$= \frac{4\cdot 16\sigma^a}{N_f+8N_b}\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\left[\frac{qX^s q}{q^5} - \zeta\frac{X^s}{q^3}\right] \to \frac{4\cdot 16\sigma^a X^s}{6(N_f+8N_b)\pi^2}(1+3\zeta)\log\, k$$

$$
\begin{aligned}
I_{\psi\psi} &= \sigma^b \sigma^a \sigma^b \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \gamma_\mu G_\psi(q) X^s G_\psi(q) \gamma_\nu D_{\mu\nu}(-q) \\
&= -\sigma^a \frac{16}{N_f + 8N_b} \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \gamma_\mu \frac{q X^s q}{q^5} \gamma_\nu \left(\delta_{\mu\nu} - \zeta \frac{q_\mu q_\nu}{q^2} \right) \\
&= -\sigma^a \frac{16}{N_f + 8N_b} \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \left[\frac{\gamma_\mu q X^s q \gamma_\mu}{q^5} - \zeta \frac{X^s}{q^3} \right] \\
&= \sigma^a \frac{16}{N_f + 8N_b} \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \left[\frac{q X^s q}{q^5} + \zeta \frac{X^s}{q^3} \right] \\
&\to \sigma^a X^s \frac{16}{6(N_f + 8N_b)\pi^2} (1 - 3\zeta) \log\, k
\end{aligned}
$$

$$
\begin{aligned}
\dim[\Phi_s^a \sigma^a X^s \bar{\psi}\psi] &= 2\dim[\psi] + \dim[\Phi] + \eta_{\mathrm{vrtx}} \\
&= \frac{5}{2} + \frac{2}{9N_b\pi^2} - \frac{64}{3(N_f + 8N_b)\pi^2}
\end{aligned}
$$

$$
\dim[y] = \frac{1}{2} - \frac{2}{9N_b\pi^2} + \frac{64}{3(N_f + 8N_b)\pi^2}
$$

$$
\begin{aligned}
&U_{T_{2x}}(\boldsymbol{r}, m_x, m_y) = g_{T_{2x}}, U_{T_{2y}}(\boldsymbol{r}, m_x, m_y) = g_{T_{2y}} \\
&U_{G_x}(\boldsymbol{r}, m_x, m_y) = g_{G_x} \\
&U_{\sigma_{xy}}(\boldsymbol{r}, m_x, m_y) = (-1)^{\delta_{(m_x,m_y),(B,B)}} g_{\sigma_{xy}} \\
&U_{\mathcal{T}}(\boldsymbol{r}, m_x, m_y) = -(-1)^{\delta_{m_x,m_y}} g_{\mathcal{T}}
\end{aligned}
$$

$$
u_{i,i+\hat{\mathbf{x}}} = it, u_{i,i+\hat{\mathbf{y}}} = (-1)^{i_x} it.
$$

$$
\begin{aligned}
&U_{T_x}(\boldsymbol{i}) = (-1)^{i_y} g_{T_x}, U_{T_y}(\boldsymbol{i}) = g_{T_y}, \\
&U_{P_x}(\boldsymbol{i}) = (-1)^{i_x} g_{P_x}, U_{P_y}(x, y) = (-1)^{i_y} g_{P_y}, \\
&U_{R_{\pi/2}}(\boldsymbol{i}) = (-1)^{i_x + i_x i_y} g_{R_{\pi/2}}, \\
&U_{\mathcal{T}}(\boldsymbol{i}) = (-1)^{i_x + i_y} g_{\mathcal{T}}.
\end{aligned}
$$

$g_{T_{2x}} = e^{i\phi_{T_{2x}}\sigma^z}$, $g_{T_{2y}} = e^{i\phi_{T_{2y}}\sigma^z}$, $g_{G_x} = e^{i\phi_{G_x}\sigma^z} i\sigma^x$, $g_{\sigma_{xy}} = e^{i\phi_{\sigma_{xy}}\sigma^z} i\sigma^x$ and $g_{\mathcal{T}} = e^{i\phi_{\mathcal{T}}\sigma^z}$

$$
\begin{aligned}
&U_{T_{2x}}(\boldsymbol{r}, m_x, m_y) = e^{i\phi_{T_{2x}}\sigma^z}, U_{T_{2y}}(\boldsymbol{r}, m_x, m_y) = e^{i\phi_{T_{2y}}\sigma^z} \\
&U_{G_x}(\boldsymbol{r}, m_x, m_y) = e^{i\phi_{G_x}\sigma^z} i\sigma^x \\
&U_{\sigma_{xy}}(\boldsymbol{r}, m_x, m_y) = (-1)^{\delta_{(m_x,m_y),(B,B)}} e^{i\phi_{\sigma_{xy}}\sigma^z} i\sigma^x \\
&U_{\mathcal{T}}(\boldsymbol{r}, m_x, m_y) = -(-1)^{\delta_{m_x,m_y}} e^{i\phi_{\mathcal{T}}\sigma^z}
\end{aligned}
$$

$$
\begin{aligned}
u_{i,i+\hat{\mathbf{x}}} &= it_{s,0}\sigma^0 - (-1)^{i_x + i_y} t_{s,z}\sigma^z \\
u_{i,i+\hat{\mathbf{y}}} &= (-1)^{i_x} it_{s,0}\sigma^0 + (-1)^{i_y} t_{s,z}\sigma^z
\end{aligned}
$$

$$U_{T_x}(\boldsymbol{i}) = (-1)^{i_y} e^{i\phi_{T_x}\sigma^z} i\sigma^x, U_{T_y}(\boldsymbol{i}) = e^{i\phi_{T_y}\sigma^z} i\sigma^x$$
$$U_{P_x}(\boldsymbol{i}) = (-1)^{i_x} e^{i\phi_{P_x}\sigma^z}, U_{P_y}(\boldsymbol{i}) = (-1)^{i_y} e^{i\phi_{P_y}\sigma^z}$$
$$U_{R_{\pi/2}}(\boldsymbol{i}) = (-1)^{i_x + i_x i_y} e^{i\phi_{R_{\pi/2}}\sigma^z} i\sigma^x$$
$$U_{\mathcal{T}}(\boldsymbol{i}) = (-1)^{i_x + i_y} e^{i\phi_{\mathcal{T}}\sigma^z}$$

the hoppings are $t_{s,0} = t\cos\left(\frac{\varphi-\pi}{4}\right)$ and $t_{s,z} = t\sin\left(\frac{\varphi-\pi}{4}\right)$

$\phi_{T_{2x}} = \phi_{T_{2y}} = 0$, $\phi_{G_x} = -\pi/4, \phi_{\sigma_{xy}} = -\pi/2$, and $\phi_{\mathcal{T}} = \pi/2$ and

$$U_{T_{2x}}(\boldsymbol{r}, m_x, m_y) = U_{T_{2y}}(\boldsymbol{r}, m_x, m_y) = \sigma^0,$$
$$U_{G_x}(\boldsymbol{r}, m_x, m_y) = i\frac{1}{\sqrt{2}}(\sigma^x + \sigma^y),$$
$$U_{\sigma_{xy}}(\boldsymbol{r}, m_x, m_y) = (-1)^{\delta_{(m_x,m_y),(BB)}} i\sigma^y,$$
$$U_{\mathcal{T}}(\boldsymbol{r}, m_x, m_y) = -(-1)^{\delta_{m_x,m_y}} i\sigma^z.$$

$$\begin{aligned}
\delta H = \sum_{\boldsymbol{r}} \mathrm{Tr}\{&(\Delta_{g,1}\sigma^y - \Delta_{g,2}\sigma^x)[\mathcal{X}^\dagger_{r,A,0} + \mathcal{X}^\dagger_{\boldsymbol{r},B,1}][\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{x}},B,0} - \mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{x}},A,1}] + \text{h.c.} \\
&+(\Delta_{g,1}\sigma^x + \Delta_{g,2}\sigma^y)[\mathcal{X}^\dagger_{r,B,0} + \mathcal{X}^\dagger_{\boldsymbol{r},A,1}][\mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{x}},A,0} - \mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{x}},B,1}] + \text{h.c.} \\
&+(\Delta_{g,1}\sigma^y + \Delta_{g,2}\sigma^x)[\mathcal{X}^\dagger_{\boldsymbol{r},B,0} - \mathcal{X}^\dagger_{\boldsymbol{r},A,1}][\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{y}},A,0} + \mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{y}},B,1}] + \text{h.c.} \\
&+(\Delta_{g,1}\sigma^x - \Delta_{g,2}\sigma^y)[\mathcal{X}^\dagger_{\boldsymbol{r},A,0} - \mathcal{X}^\dagger_{\boldsymbol{r},B,1}][\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{y}},B,0} + \mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{y}},A,1}] + \text{ h.c.}\} \\
\approx \int \mathrm{d}^{\mathcal{D}}\boldsymbol{r} \; &+(\Delta_{g,1}\sigma^x + \Delta_{g,2}\sigma^y)(\Delta_{g,1}\sigma^y - \Delta_{g,2}\sigma^x)(\mathcal{X}^\dagger\rho^x\mu^z\mathcal{X} - \mathcal{X}^\dagger\rho^z\mu^x\mathcal{X}) \\
&+(\Delta_{g,1}\sigma^y + \mathcal{X}^\dagger_{g,2}\sigma^z\mu^x\mathcal{X})(\mathcal{X}^\dagger\rho^x\mu^z\mathcal{X} - \mathcal{X}^\dagger\rho^z\mu^x\mathcal{X}) \\
&+(\Delta_{g,1}\sigma^x - \Delta_{g,2}\sigma^y)(\mathcal{X}^\dagger\rho^x\mu^z\mathcal{X} + \mathcal{X}^\dagger\rho^z\mu^x\mathcal{X})\} \\
\Rightarrow \delta\mathcal{L} = 2\Delta_{g,1}\mathrm{Tr}&[\sigma^x\overline{\mathcal{X}}(\gamma^x\mu^z - \gamma^y\mu^x)\mathcal{X}] + 2\Delta_{g,1}\mathrm{Tr}[\sigma^y\overline{\mathcal{X}}(\gamma^x\mu^z + \gamma^y\mu^x)\mathcal{X}]
\end{aligned}$$

$$\begin{aligned}
\delta_H = \sum_{\boldsymbol{r}} &\Delta_{g,2}\mathrm{Tr}\{\sigma^x[\mathcal{X}^\dagger_{\boldsymbol{r},A,0} + \mathcal{X}^\dagger_{\boldsymbol{r},B,1}][\mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{y}},B,0} - \mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{y}},A,1}] + [\mathcal{X}^\dagger_{\boldsymbol{r},B,0} - \mathcal{X}^\dagger_{\boldsymbol{r},A,1}][\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{y}},A,0} + \mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{y}},B,1}] \\
&-[\mathcal{X}^\dagger_{\boldsymbol{r},A,0} + \mathcal{X}^\dagger_{\boldsymbol{r},B,1}][\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{x}},B,0} - \mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{x}},A,1}] - [\mathcal{X}^\dagger_{\boldsymbol{r},B,0} - \mathcal{X}^\dagger_{\boldsymbol{r},A,1}][\mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{x}},A,0} + \mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{x}},B,1}]\} \\
&+\Delta_{g,2}\mathrm{Tr}\{\sigma^y[\mathcal{X}^\dagger_{\boldsymbol{r},B,0} + \mathcal{X}^\dagger_{\boldsymbol{r},A,1}][\mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{x}},A,0} - \mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{x}},B,1}] + [\mathcal{X}^\dagger_{\boldsymbol{r},A,0} - \mathcal{X}^\dagger_{\boldsymbol{r},B,1}][\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{x}},B,0} + \mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{x}},A,1}] \\
&-[\mathcal{X}^\dagger_{\boldsymbol{r},B,0} + \mathcal{X}^\dagger_{\boldsymbol{r},A,1}][\mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{y}},A,0} - \mathcal{X}_{\boldsymbol{r}-\hat{\mathbf{y}},B,1}] - [\mathcal{X}^\dagger_{\boldsymbol{r},A,0} - \mathcal{X}^\dagger_{\boldsymbol{r},B,1}][\mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{y}},B,0} + \mathcal{X}_{\boldsymbol{r}+\hat{\mathbf{y}},A,1}]\} \\
\approx &\Delta_{g,2}\int \mathrm{d}^{\mathcal{D}}\boldsymbol{r}\mathrm{Tr}\{\sigma^x\mathcal{X}^\dagger(\mu^y - \rho^y)(i\partial_x + i\partial_y)\mathcal{X}\} + \mathrm{Tr}\{\sigma^x\mathcal{X}^\dagger(-\mu^y - \rho^y)(i\partial_y - i\partial_x)\mathcal{X}\} \\
\Rightarrow \delta\mathcal{L} = &\Delta_{g,2}\mathrm{Tr}[\sigma^x\overline{\mathcal{X}}(-\gamma^0\mu^y + 1)(i\partial_x + i\partial_y)\mathcal{X} + \sigma^y\bar{X}(i\gamma^0\mu^y + 1)(i\partial_y - i\partial_x)\mathcal{X}]
\end{aligned}$$

$$N_b \int \frac{\mathrm{d}^{\mathcal{D}}p}{(2\pi^3)}\frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi^3)}\Big(8G_\Phi(q)\delta_{p,0}\int \frac{\mathrm{d}^{\mathcal{D}}p'}{(2\pi)^3}A^a_\mu(p')A^a_\mu(-p')$$
$$-\frac{1}{2}G_\Phi(q)G_\Phi(q-p)[-3\lambda(p)\lambda(-p) + 8(2q-p)_\mu A^a_\mu(p)(2q-p)_\nu A^a_\nu(-p)]\Big)$$


$$-\frac{N_f}{2}\int \frac{\mathrm{d}^{\mathcal{D}}p}{(2\pi)^3}\frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\mathrm{Tr}[G_\psi(q)\gamma^\mu A^a_\mu(p)\sigma_a G_\psi(p+q)\gamma^\nu A^b_\nu(-p)\sigma_b]$$

$$\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{1}{q^2+r}=-\frac{\sqrt{r}}{4\pi}$$

$$\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{1}{(q^2+r)((q-p)^2+r)}=\frac{1}{4\pi p}\arctan\frac{p}{2\sqrt{r}}$$

$$\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{(2q-p)_\mu(2q-p)_\nu}{(q^2+r)((q-p)^2+r)}=-\left(\delta_{\mu\nu}+\frac{p_\mu p_\nu}{p^2}\right)\frac{\sqrt{r}}{4\pi}$$
$$-\left(\delta_{\mu\nu}-\frac{p_\mu p_\nu}{p^2}\right)\left(\frac{4r+p^2}{8\pi p}\arctan\frac{p}{2\sqrt{r}}\right)$$

$$\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{\mathrm{Tr}[\gamma^\mu q\gamma^\nu(\not{p}+q)]}{q^2(p+q)^2}=-\frac{1}{16p}$$



$$\Sigma_\psi(k)=\sigma^a\sigma^a\int\frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\gamma_\mu G_\psi(k+q)\gamma_\nu D_{\mu\nu}(-q)$$
$$=3\int\frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{16}{N_f+8N_b}\gamma_\mu\frac{q+\not{k}}{(k+q)^2}\gamma_\nu\frac{1}{q}\left(\delta_{\mu\nu}-\zeta\frac{q_\mu q_\nu}{q^2}\right)$$
$$=3\frac{16}{N_f+8N_b}\int\frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\left[\gamma_\mu\frac{q+\not{k}}{(k+q)^2q}\gamma_\mu-\zeta\frac{q(q+\not{k})q}{(k+q)^2q^3}\right]$$
$$=3\frac{16}{N_f+8N_b}\int\frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\left[-\frac{q+\not{k}}{(k+q)^2q}+\zeta\frac{q(q+\not{k})q}{(k+q)^2q^3}\right]$$

$$\int\frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{q+\not{k}}{(k+q)^2q}=\frac{1}{2}\int_0^1\mathrm{d}x\int\frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{x^{-1/2}(q+\not{k})}{[(1-x)(k+q)^2+xq^2]^{3/2}}$$
$$=\frac{1}{2}\int_0^1\mathrm{d}x\int\frac{\mathrm{d}^{\mathcal{D}}l}{(2\pi)^3}\frac{x^{-1/2}(l+x\not{k})}{[l^2+x(1-x)k^2]^{3/2}}$$
$$=\int_0^1\mathrm{d}x\frac{1}{2(4\pi)^{3/2}}\left[\frac{\sqrt{x}\not{k}}{\Gamma(3/2)}\frac{\Gamma(0)}{[x(1-x)k^2]^0}\right]$$
$$\to-\frac{1}{6\pi^2}\not{k}\log k$$

$$\Delta=x(1-x)k^2\text{ and }\epsilon=\frac{D}{2}-\frac{3}{2}$$

$$\frac{\Gamma(\epsilon)}{\Delta^\epsilon}\approx\frac{1/\epsilon}{1+\epsilon\log\Delta}\approx\frac{1}{\epsilon}-\log\Delta\to-\log k^2=-2\log k.$$

$$\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{q(q+\not{k})q}{(k+q)^2q^3}=\frac{3}{2}\int_0^1 \mathrm{d}x\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{\sqrt{x}q(q+\not{k})q}{[(1-x)(k+q)+xq^2]^{5/2}}$$
$$=\frac{3}{2}\int_0^1 \mathrm{d}x\sqrt{x}\int \frac{\mathrm{d}^{\mathcal{D}}l}{(2\pi)^3}\frac{(l-(1-x)\not{k})(l+x\not{k})(l-(1-x)\not{k})}{[l^2+x(1-x)k^2]^{5/2}}$$
$$\to\frac{3}{2}\int_0^1 \mathrm{d}x\frac{\sqrt{x}(-2\log\,k)}{8\pi\sqrt{\pi}\cdot 2\Gamma(5/2)}\delta_{\mu\nu}[-(1-x)(\gamma^\mu\gamma^\nu\not{k}+\not{k}\gamma^\mu\gamma^\nu)+x\gamma^\mu\not{k}\gamma^\nu]$$
$$=\int_0^1 \mathrm{d}x\frac{-2\sqrt{x}\log\,k}{8\pi^2}[-3(1-x)\not{k}-x\not{k}-3(1-x)\not{k}]$$
$$=\int_0^1 \mathrm{d}x\frac{-2\sqrt{x}(5x-6)}{8\pi^2}\not{k}\log\,k$$
$$=-\frac{1}{2\pi^2}\not{k}\log\,k$$

$$I_{A;1}=8\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}G_\Phi(k+q)D_{\mu\nu}(-q)(2k+q)_\mu(2k+q)_\nu$$
$$=\frac{8\cdot 16}{N_f+8N_b}\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{(2k+q)_\mu(2k+q)_\nu}{q(k+q)^2}\left(\delta_{\mu\nu}-\zeta\frac{q_\mu q_\nu}{q^2}\right)$$
$$=\frac{8\cdot 16}{N_f+8N_b}\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\left[\frac{(2k+q)^2}{q(k+q)^2}-\zeta\frac{(2kq+q^2)^2}{q^3(k+q)^2}\right]$$

$$\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{(2k+q)^2}{q(k+q)^2}=\frac{1}{2}\int_0^1 \mathrm{d}x\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{x^{-1/2}(2k+q)^2}{[(1-x)(k+q)^2+xq^2]^{3/2}}$$
$$=\frac{1}{2}\int_0^1 \mathrm{d}x\int \frac{\mathrm{d}^{\mathcal{D}}l}{(2\pi)^3}\frac{x^{-1/2}(l+(1+x)k)^2}{[l^2+x(1-x)k^2]^{3/2}}$$
$$\to\frac{k^2\log\,k}{16\pi\sqrt{\pi}\Gamma(3/2)}\int_0^1 \mathrm{d}x\frac{1}{\sqrt{x}}[-2(1+x)^2+3x(1-x)]$$
$$=-\frac{10}{12\pi^2}k^2\log\,k$$

$$\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{(2kq+q^2)^2}{q^3(k+q)^2}=\frac{3}{2}\int_0^1 \mathrm{d}x\int \frac{\mathrm{d}^{\mathcal{D}}q}{(2\pi)^3}\frac{\sqrt{x}(2kq+q^2)^2}{[(1-x)(k+q)^2+xq^2]^{5/2}}$$
$$=\frac{3}{2}\int_0^1 \mathrm{d}x\int \frac{\mathrm{d}^{\mathcal{D}}l}{(2\pi)^3}\frac{\sqrt{x}\left[(l+(1+x)k)_\mu(l-(1-x)k)_\mu\right]^2}{[l^2+x(1-x)k^2]^{5/2}}$$
$$=\frac{3}{2}\int_0^1 \mathrm{d}x\sqrt{x}\int \frac{\mathrm{d}^{\mathcal{D}}l}{(2\pi)^3}\frac{(l^2)^2+4x^2(k\cdot l)^2+(1-x^2)^2k^4-2(1-x^2)l^2k^2}{[l^2+x(1-x)k^2]^{5/2}}$$
$$\to\frac{3}{2}\frac{k^2\log\,k}{8\pi\sqrt{\pi}\Gamma(5/2)}\int_0^1 \mathrm{d}x\sqrt{x}\left[\frac{15x(1-x)}{2}-4x^2+6(1-x^2)\right]$$
$$=\frac{1}{2\pi^2}k^2\log\,k$$

$$
\begin{aligned}
I_{\lambda;1} &= i^2 \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} G_{\Phi}(k+q) D_{\lambda}(-q) \\
&= -\frac{8}{3N_b} \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \frac{q^2}{q(k+q)^2} \\
&= -\frac{4}{3N_b} \int_0^1 \mathrm{d}x \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \frac{x^{-1/2} q^2}{[(1-x)(k+q)^2 + xq^2]^{3/2}} \\
&= -\frac{4}{3N_b} \int_0^1 \mathrm{d}x \int \frac{\mathrm{d}^{\mathcal{D}} l}{(2\pi)^3} x^{-1/2} \frac{l^2 + (1-x)^2 k^2 - 2(1-x) l \cdot k}{[l^2 + x(1-x)k^2]^{3/2}} \\
&\to -\frac{4}{3N_b \cdot 8\pi\sqrt{\pi}\Gamma(3/2)} \int_0^1 \mathrm{d}x \frac{1}{\sqrt{x}} \left[\frac{6x(1-x)}{2} - 2(1-x)^2 \right] k^2 \log\, k \\
&= \frac{4}{9N_b \pi^2} k^2 \log\, k
\end{aligned}
$$

$$
\begin{aligned}
I_{A;2} &= 3 \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \gamma_{\mu} G_{\psi}(k_1+q) G_{\psi}(k_2+q) \gamma_{\nu} D_{\mu\nu}(-q) \\
&= \frac{3 \cdot 16}{N_f + 8N_b} \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \gamma_{\mu} \frac{\not{k}_1 + q}{(k_1+q)^2} \frac{\not{k}_2 + q}{(k_2+q)^2} \frac{1}{q} \gamma_{\nu} \left(\delta_{\mu\nu} - \zeta \frac{q_{\mu} q_{\nu}}{q^2} \right) \\
&= \frac{3 \cdot 16}{N_f + 8N_b} \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \left[\frac{-(k_1+q)(k_2+q) + 4(k_1+q) \cdot (k_2+q)}{(k_1+q)^2 (k_2+q)^2 q} - \zeta \frac{q(k_1+q)(k_2+q)q}{(k_1+q)^2 (k_2+q)^2 q^3} \right].
\end{aligned}
$$

$$
\begin{aligned}
I_{A;2,1} &= \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \frac{3}{(k+q)^2 q} \\
&= \frac{1}{2} \int_0^1 \mathrm{d}x \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \frac{3x^{-1/2}}{[(1-x)(k+q)^2 + xq^2]^{3/2}} \\
&= \frac{3}{2} \int_0^1 \mathrm{d}x \frac{1}{\sqrt{x}} \int \frac{\mathrm{d}^{\mathcal{D}} l}{(2\pi)^3} \frac{1}{[l^2 + x(1-x)k^2]^{3/2}} \\
&\to \frac{3}{2} \int_0^1 \mathrm{d}x \frac{1}{\sqrt{x}} \frac{1}{8\pi\sqrt{\pi}\Gamma(3/2)} (-2\log\, k) = \frac{-3i}{2\pi^2} \log\, k
\end{aligned}
$$

$$
I_{A;2,2} = \int \frac{\mathrm{d}^{\mathcal{D}} q}{(2\pi)^3} \frac{q^2}{(k+q)^2 q^3} = \frac{1}{3} I_{A;2,1}
$$

$$
\Gamma_k = \int_x \sqrt{g'} \{ -\frac{F}{2} R' + U + \frac{K}{2} D^{\mu} \chi D_{\mu} \chi + Z_H D^{\mu} H^{\dagger} D_{\mu} H + \cdots \}
$$

$$
F = \xi \chi^2 + \xi_H H^{\dagger} H + f_{\infty} k^2
$$

$$
U_E = \frac{M^4}{F^2} U
$$

$$
g_{\mu\nu} = \frac{F}{M^2} g'_{\mu\nu}
$$

$$
\frac{\partial U_E}{\partial H}(H_0) = 0, \varphi_0 = |H_0|.
$$

$$U_E = \frac{M^4 U}{\xi^2 \chi^4}\left(1 - \frac{2\xi_H H^\dagger H}{\xi\chi^2} - \frac{2f_\infty k^2}{\xi\chi^2}\right)$$

$\partial U_E/\partial\left(H^\dagger H\right) = 0$ at $H^\dagger H = \varphi_0^2$ and $k = 0$ (gauge symmetry)

$$\frac{\partial U}{\partial H^\dagger H}\left(\chi, H^\dagger H = \varphi_0^2\right) = \frac{2\xi_H}{\xi\chi^2} U\left(\chi, H^\dagger H = \varphi_0^2\right)$$

$$\frac{\varphi_{0E}^2}{M^2} = \frac{\varphi_0^2}{F} = \frac{\varphi_0^2}{\xi\chi^2}$$

$$h_0 = \frac{2\varphi_0^2}{\chi^2} = 2\xi\frac{\varphi_{0E}^2}{M^2}$$

$$\frac{m_H^2}{M^2} = \frac{2\lambda_h\varphi_0^2}{\xi\chi^2} = \frac{\lambda_{h,0}}{\xi}h_0$$

$$\lambda_{h,0} = \frac{\partial^2 U}{\partial(H^\dagger H)^2}\left(\chi, H^\dagger H = \varphi_0^2\right)$$

$$k\partial_k U = \frac{\mathcal{N}k^4}{32\pi^2} = 4c_U k^4$$

$$u = \frac{U}{k^4}, \tilde{\rho} = \frac{\chi^2}{2k^2}, \tilde{h} = \frac{H^\dagger H}{k^2}, h = \frac{\tilde{h}}{\tilde{\rho}} = \frac{2H^\dagger H}{\chi^2}$$

$$\frac{\partial u}{\partial h}(\tilde{\rho}, h_0) = 0$$

$$\hat{\lambda}_h = \frac{\partial^2 u}{\partial h^2} = \lambda_h\tilde{\rho}^2$$

$$\lambda_{h,0} = \frac{1}{\tilde{\rho}^2}\partial_{\tilde{h}}^2 u(\tilde{\rho}, h_0)$$

$$k\partial_k u = -4u + 2\tilde{\rho}\partial_{\tilde{\rho}}u + 4c_U$$

$$c_U = \frac{1}{128\pi^2}\left(\bar{N}_S + 2\bar{N}_V - 2\bar{N}_F + 2\right)$$

$$m_f^2 = y_f^2 H^\dagger H$$

$$\bar{N}_F = \sum_f \left(1 + \frac{m_f^2}{k^2}\right)^{-1} + 3$$
$$= \sum_f \left(1 + y_f^2 h\tilde{\rho}\right)^{-1} + 3$$

$$2\bar{N}_V = \frac{6}{1 + \tilde{m}_W^2} + \frac{3}{1 + \tilde{m}_Z^2} - 3 + 18$$

$$\tilde{m}_W^2 = \frac{g_2^2 h\tilde{\rho}}{2}, \tilde{m}_Z^2 = \left(\frac{g_2^2}{2} + \frac{3g_1^2}{10}\right) h\tilde{\rho},$$

$$\bar{N}_s = \sum_s \left(1 + \frac{m_s^2}{k^2}\right)^{-1}$$

$$M_{ij}^2 = \frac{\partial^2 U}{\partial\varphi_i \partial\varphi_j},$$

where $\varphi_i = (\chi, \tilde{H}_\gamma)$ with $\tilde{H}_\gamma$ the four fields in H, i.e. $H_1 = \frac{1}{\sqrt{2}}(\tilde{H}_1 + i\tilde{H}_2), H_2 = \frac{1}{\sqrt{2}}(\tilde{H}_3 + i\tilde{H}_4)$,

$$H^\dagger H = \tfrac{1}{2}\textstyle\sum_\gamma \tilde{H}_\gamma^2$$

$$\tilde{\rho}\partial_{\tilde{\rho}} u = 2(u - c_U) \int_{\tilde{h}\partial_{\tilde{h}} u = 2(u - c_U),}^{\blacksquare} \tilde{h}\partial_{\tilde{h}} = \tilde{\rho}\partial_{\tilde{\rho}}$$

$$k\partial_k Z_H|_{\chi,H} = -\eta_H Z_H$$

$$\eta_H = -k\partial_k \ln Z_H = \frac{3}{8\pi^2}\left(y_t^2 - \frac{3}{4}g_2^2 - \frac{3}{20}g_1^2\right)$$

$$\tilde{h} = \frac{H_R^\dagger H_R}{k^2} = \frac{Z_H H^\dagger H}{k^2}, h = \frac{\tilde{h}}{\tilde{\rho}}$$

$$k\partial_{k|\chi,H} = k\partial_{k|\chi,H_R} - \frac{\eta_H}{2} H_R \frac{\partial}{\partial H_{R|\chi,k}},$$

$$k\partial_{k|\tilde{\rho},h} = k\partial_{k|\chi,H} + \eta_H h\partial_{h|\tilde{\rho}} + 2\tilde{\rho}\partial_{\tilde{\rho}|h}.$$

$$2\tilde{\rho}\partial_{\tilde{\rho}|h} \to 2\tilde{\rho}\partial_{\tilde{\rho}|h} + \eta_H h\partial_{h|\tilde{\rho}},$$
$$2\tilde{h}\partial_{\tilde{h}|h} \to (2 + \eta_H)\tilde{h}\partial_{\tilde{h}|h} + \eta_H h\partial_{h|\tilde{h}}.$$

$$\tilde{\rho}\partial_{\tilde{\rho}}(\partial_h u) = 2\partial_h u - \frac{3y_t^2\tilde{\rho}}{16\pi^2(1 + y_t^2 h\tilde{\rho})^2}\left(1 + \frac{\partial \ln y_t^2}{\partial \ln h}\right).$$

$$|\partial \ln y_t^2 / \partial \ln h| \ll 1$$

$$\partial_{\tilde{\rho}}\partial_h u = \frac{2}{\tilde{\rho}}\partial_h u - \frac{3y_t^2}{16\pi^2(1 + y_t^2 h\tilde{\rho})^2}.$$

$$\hat{\lambda}_h = \partial_h^2 u, \partial_{\tilde{\rho}}\hat{\lambda}_h = \frac{2}{\tilde{\rho}}\hat{\lambda}_h + \frac{3y_t^4\tilde{\rho}}{8\pi^2(1 + y_t^2 h\tilde{\rho})^3}.$$

$$\partial_h u = \partial_h L_-(h)\tilde{\rho}^2 + \frac{3y_t^2\tilde{\rho}}{16\pi^2} + \frac{3y_t^4 h\tilde{\rho}^2}{8\pi^2}\ln(h\tilde{\rho}) + \cdots$$
$$\partial_h^2 u = \partial_h^2 L_-(h)\tilde{\rho}^2 + \frac{3y_t^4\tilde{\rho}^2}{8\pi^2}(\ln(h\tilde{\rho}) + 1) + \cdots$$

$$u = L_{-}(h)\tilde{\rho}^2 - \frac{59}{128\pi^2} + \frac{3y_t^2\tilde{h}}{16\pi^2} + \frac{3y_t^4\tilde{h}^2}{16\pi^2}\left(\ln(\tilde{h}) - \frac{1}{2}\right) + \cdots$$

$$U = \frac{1}{4}L_{-}(h)\chi^4 - \frac{59k^4}{128\pi^2} + \frac{3y_t^2k^2H^{\dagger}H}{16\pi^2} + \frac{3y_t^4\left(H^{\dagger}H\right)^2}{16\pi^2}\left(\ln\left(\frac{H^{\dagger}H}{k^2}\right) - \frac{1}{2}\right)$$

$$k\partial_k U_{\chi,H} = -\frac{59}{32\pi^2}k^4 + \frac{3y_t^2k^2H^{\dagger}H}{8\pi^2} - \frac{3y_t^4\left(H^{\dagger}H\right)^2}{8\pi^2}$$

$$k\partial_k\lambda_h = -\frac{3y_t^2}{4\pi^2}$$

$$\tilde{h}\partial_{\tilde{h}}u = 2u + \frac{1}{64\pi^2}\left(47 + \frac{12}{1+y_t^2\tilde{h}}\right)$$

$$u = \tilde{L}(h)\tilde{h}^2 - \frac{47}{128\pi^2} - \frac{3}{32\pi^2}t_u(\tilde{m}_t^2), \tilde{m}_t^2 = y_t^2\tilde{h}$$

$$t_u(\tilde{m}^2) = 1 - 2\tilde{m}^2 - 2\tilde{m}^4\ln\frac{\tilde{m}^2}{1+\tilde{m}^2}$$

$$\tilde{m}^2\partial_{\tilde{m}^2}t_u = 2t_u - \frac{2}{1+\tilde{m}^2}$$

$$\frac{\partial t_u}{\partial\tilde{m}^2} = -\left(2 + \frac{2\tilde{m}^2}{1+\tilde{m}^2} + 4\tilde{m}^2\ln\left(\frac{\tilde{m}^2}{1+\tilde{m}^2}\right)\right)$$

$$\frac{\partial^2 t_u}{(\partial\tilde{m}^2)^2} = -\left(4\ln\left(\frac{\tilde{m}^2}{1+\tilde{m}^2}\right) + \frac{4}{1+\tilde{m}^2} + \frac{2}{(1+\tilde{m}^2)^2}\right)$$

$$\lambda_h'(\tilde{h}) = \frac{\partial^2 u}{\partial\tilde{h}^2}\,{}_{|h}\bar{\lambda}_h(h) - \frac{3y_t^4}{32\pi^2}\frac{\partial^2 t_u}{(\partial\tilde{m}_t^2)^2}$$

$$\lambda_h' = \bar{\lambda}_h' + \frac{3y_t^4}{8\pi^2}\left[\ln\left(y_t^2\frac{H^{\dagger}H}{k^2}\right) - \ln\left(1 + y_t^2\frac{H^{\dagger}H}{k^2}\right) + \frac{k^2}{y_t^2H^{\dagger}H + k^2} + \frac{k^4}{2(y_t^2H^{\dagger}H + k^2)^2}\right]$$

$$k^2 \gg y_t^2H^{\dagger}H$$

$$\lambda_h' = \bar{\lambda}_h' + \frac{3y_t^4}{8\pi^2}\left[\ln\left(y_t^2\frac{H^{\dagger}H}{k^2}\right) + \frac{3}{2}\right]$$

$$k\partial_k\lambda_h' = -\frac{3y_t^4}{4\pi^2}$$

$$\tilde{h}\partial_{\tilde{h}}\lambda'_h = \tilde{h}\partial_{\tilde{h}}\left(\partial_{\tilde{h}}^2 u\right) = \tilde{\partial}_h^2\left(\tilde{h}\partial_{\tilde{h}} u\right) - 2\lambda'_h$$
$$= \frac{3}{16\pi^2}\partial_{\tilde{h}}^2\left(1 + y_t^2\tilde{h}\right)^{-1} = \frac{3y_t^4}{8\pi^2\left(1 + y_t^2\tilde{h}\right)^3}$$

$$\lambda'_h = \bar{\lambda}'_h + \frac{3y_t^4}{8\pi^2}\left(\ln\left(\frac{y_t^2\tilde{h}}{1 + y_t^2\tilde{h}}\right) + \frac{1}{1 + y_t^2\tilde{h}} + \frac{1}{2\left(1 + y_t^2\tilde{h}\right)^2}\right)$$

$$\mu'_h(\tilde{h}) = \partial_{\tilde{h}} u(\tilde{h}, h)_{|h}$$

$$\tilde{h}\partial_{\tilde{h}}\mu'_{h|h} = \partial_{\tilde{h}}\left(\tilde{h}\partial_{\tilde{h}} u\right) - \mu'_h$$
$$= \mu'_h - \frac{3y_t^2}{16\pi^2\left(1 + y_t^2\tilde{h}\right)^2},$$

$$\mu'_h = c_\mu\tilde{h} + \frac{3y_t^2}{16\pi^2}\left(1 + \frac{y_t^2\tilde{h}}{1 + y_t^2\tilde{h}} + 2y_t^2\tilde{h}\ln\left(\frac{y_t^2\tilde{h}}{1 + y_t^2\tilde{h}}\right)\right)$$

$$\mu'_h(\tilde{h} = 0) = \frac{3y_t^2}{16\pi^2}$$

$$m_H^2 = \frac{\partial U}{\partial(H^\dagger H)_{|H=0}} = \frac{3y_t^2 k^2}{16\pi^2}$$

$$\tilde{L}(h) = \frac{L_-(h)}{h^2} = \frac{L_+(h)}{h^2} = \frac{L(h)}{h^2}.$$

$$\mu_h = \partial_{\tilde{h}} u_{|\tilde{\rho}} = k^{-2}\frac{\partial U}{\partial H^\dagger H}\ {}_{|\chi},$$

$$\lambda_h = \partial_{\tilde{h}}^2 u_{|\tilde{\rho}} = \left\langle \frac{\partial^2 U}{(\partial H^\dagger H)^2}\right|_{\partial_{\tilde{h}} = \tilde{\rho}^{-1}\partial_h}$$

$$\partial_{\tilde{h}|\tilde{\rho}} = \partial_{\tilde{h}|h} + \frac{h}{\tilde{h}}\partial_{h|\tilde{h}}$$

$$L(h) = \frac{1}{2}\bar{\lambda}_\chi + \bar{\lambda}_m h + \frac{1}{2}\bar{\lambda}_h h^2(1 + \Delta(h)).$$

$$U = \frac{1}{8}\bar{\lambda}_\chi\chi^4 + \frac{1}{2}\bar{\lambda}_m\chi^2 H^\dagger H$$
$$+ \frac{1}{2}\bar{\lambda}_h\left(H^\dagger H\right)^2\left(1 + \Delta\left(\frac{2H^\dagger H}{\chi^2}\right)\right)$$
$$- \frac{k^4}{128\pi^2}\left[47 + 12t_u\left(\frac{y_t^2 H^\dagger H}{k^2}\right)\right].$$

$$\lim_{\tilde{m}^2\to\infty} t_u(\tilde{m}^2) = \frac{2}{3\tilde{m}^2}$$

$$U_{k\to 0} = \frac{1}{2}\bar{\lambda}_h\left(H^\dagger H - \varphi_0^2\right)^2 + \frac{\delta}{8}\chi^4 + \Delta U,$$

$$\partial_h L(h)_{|h_0} - \partial_h L(h)_{|0} = \bar{\lambda}_h h_0,$$

$$\bar{\lambda}_m = \partial_h L(h)_{|0}$$

$$\Delta(h_0) + \frac{1}{2}h\partial_h\Delta(h_0) = \partial^2 U/\partial\left(H^\dagger H\right)^2.$$

$$\Delta U = \frac{1}{2}\bar{\lambda}_h\left(H^\dagger H\right)^2\Delta$$

$$\partial U/\partial\left(H^\dagger H\right) \text{ at } \varphi_0^2$$

$$\varphi_0^2 = -\frac{\bar{\lambda}_m}{2\bar{\lambda}_h}\chi^2, \delta = \bar{\lambda}_\chi - \frac{\bar{\lambda}_m^2}{\bar{\lambda}_h}$$

$$\lambda_h = \frac{\partial^2 U}{\partial(H^\dagger H)^2}_{|\varphi_0} = \bar{\lambda}_h\left(1 + \frac{3}{2}h\partial_h\Delta + \frac{1}{2}h^2\partial_{h^2}\Delta\right)_{|h_0}.$$

$$h_0 = \frac{2\varphi_0^2}{\chi^2} = -\frac{\bar{\lambda}_m}{\bar{\lambda}_h}$$

$$\tilde{H}_1 = \sqrt{2\tilde{h}k^2}, \tilde{H}_2 = \tilde{H}_3 = \tilde{H}_4 = \tilde{m}_g^2 \setminus \partial_{\tilde{h}}u$$

$$\tilde{m}_s^2 = \begin{pmatrix} \partial_{\tilde{h}}u + 2\tilde{h}\tilde{\partial}_h^2 u & 2\sqrt{\tilde{h}\tilde{\rho}}\partial_{\tilde{\rho}}\partial_{\tilde{h}}u \\ 2\sqrt{\tilde{h}\tilde{\rho}}\partial_{\tilde{\rho}}\partial_{\tilde{h}}u & \partial_{\tilde{\rho}}u + 2\tilde{\rho}\partial_{\tilde{\rho}}^2 u \end{pmatrix} = \begin{pmatrix} \tilde{m}_r^2 & \tilde{m}_{rc}^2 \\ \tilde{m}_{rc}^2 & \tilde{m}_c^2 \end{pmatrix}.$$

$$\tilde{m}_\pm^2 = \frac{1}{2}\left(\tilde{m}_r^2 + \tilde{m}_c^2 \pm \sqrt{(\tilde{m}_r^2 - \tilde{m}_c^2)^2 + 4\tilde{m}_{rc}^4}\right)$$

$$\tilde{m}_r^2 = \frac{1}{\tilde{\rho}}\left(\partial_h u + 2\hat{\lambda}_h h\right)$$

$$\tilde{m}_{rc}^2 = 2\sqrt{h}\partial_{h|\tilde{\rho}}\partial_{\tilde{\rho}|\tilde{h}}u$$

$$= 2\sqrt{h}\left(\partial_{\tilde{\rho}}\partial_h u(\tilde{\rho}, h) - \frac{1}{\tilde{\rho}}\partial_h u - \lambda_h h\tilde{\rho}\right)$$

$$\partial_{\tilde{\rho}|\tilde{h}} = \partial_{\tilde{\rho}|h} - \frac{h}{\tilde{\rho}}\partial_{h|\tilde{\rho}}$$

$$\bar{N}_s = \frac{3}{1 + \tilde{\rho}^{-1}\partial_h u} + \frac{1}{1 + \tilde{m}_+^2} + \frac{1}{1 + \tilde{m}_-^2}.$$

$$\partial_h u(\tilde{\rho}, h_0(\tilde{\rho})) = \tilde{m}_r^2 \sim h_0\tilde{\rho}$$

$$\tilde{m}_g^2 = 0, \tilde{m}_r^2 = 2\lambda_h h_0 \tilde{\rho}$$

$$\begin{aligned}\tilde{m}_{rc}^2 &= -\sqrt{h_0}\left(\tilde{m}_r^2 - 2\partial_{\tilde{\rho}}\partial_h u(\tilde{\rho}, h_0)\right) \\ &= -\sqrt{h_0}\left(\tilde{m}_r^2 + \frac{3y_t^2}{8\pi^2(1+y_t^2 h\tilde{\rho})^2}\right).\end{aligned}$$

$$\tilde{m}_r^2 = 2\lambda_h \tilde{h}_0 = 2\lambda_h h_0 \tilde{\rho}$$

$$u(\tilde{\rho}, h) = \bar{u}(\tilde{\rho}, h) + L(h)\tilde{\rho}^2$$

$$u = L(h)\tilde{\rho}^2 + \frac{1}{128\pi^2}\left\{17 - 2\sum_f n_f t_u(\tilde{m}_f^2) + 3\sum_v t_u(\tilde{m}_v^2) + \sum_s t_u(\tilde{m}_s^2)\right\}$$

$$L(h)\tilde{\rho}^2 = \bar{\lambda}_m \tilde{\rho}\tilde{h} + \frac{1}{2}\bar{\lambda}_m \tilde{h}^2 + \frac{1}{2}\bar{\lambda}_\chi \tilde{\rho}^2$$

$$\begin{aligned}u = \bar{\lambda}_m \tilde{\rho}\tilde{h} + \frac{1}{2}\bar{\lambda}_h \tilde{h}^2 + \frac{1}{2}\bar{\lambda}_\chi \tilde{\rho}^2 + \frac{1}{128\pi^2}\{&3t_u(\tilde{m}_Z^2) \\ &+6t_u(\tilde{m}_W^2) + 3t_u(\tilde{m}_g^2) + t_u(\tilde{m}_r^2) - 12t_u(\tilde{m}_t^2) \\ &-12t_u(\tilde{m}_b^2) - 12t_u(\tilde{m}_{ch}^2) - 4t_u(\tilde{m}_\tau^2) - 32\}\end{aligned}$$

$$\tilde{m}_W^2 = \frac{1}{2}g_2^2\tilde{h}, \tilde{m}_Z^2 = \left(\frac{1}{2}g_2^2 + \frac{3}{10}g_1^2\right)\tilde{h}$$

$$\tilde{m}_t^2 = y_t^2\tilde{h}, \tilde{m}_b^2 = y_b^2\tilde{h}, \tilde{m}_{ch}^2 = y_{ch}^2\tilde{h},$$

$$\tilde{m}_g^2 = \partial_{\tilde{h}}u, \tilde{m}_r^2 = \tilde{\partial}_h u + 2\tilde{h}\partial_{\tilde{h}}^2 u$$

$$\begin{aligned}\partial_{\tilde{h}}u = \bar{\lambda}_m + \bar{\lambda}_h\tilde{h} - \frac{1}{64\pi^2}\{&3g_2^2 s_u(\tilde{m}_W^2) \\ &+\left(\frac{3}{2}g_2^2 + \frac{9}{10}g_1^2\right)s_u(\tilde{m}_Z^2) - 12y_t^2 s_u(\tilde{m}_t^2) \\ &-12y_b^2 s_u(\tilde{m}_b^2) - 12s_u(\tilde{m}_{ch}^2) - 4s_u(\tilde{m}_\tau^2) \\ &+3\partial_{\tilde{h}}^2 u s_u(\tilde{m}_g^2) + \left(3\partial_{\tilde{h}}^2 + 2\tilde{h}\partial_{\tilde{h}}^3\right)u s_u(\tilde{m}_r^2)\}\end{aligned}$$

$$s_u(\tilde{m}^2) = 1 + 2\tilde{m}^2 \ln\left(\frac{\tilde{m}^2}{1+\tilde{m}^2}\right) + \frac{\tilde{m}^2}{1+\tilde{m}^2} = -\frac{1}{\tilde{m}^2}\left(t_u(\tilde{m}^2) - \frac{1}{1+\tilde{m}^2}\right)$$

$$s_u(0) = 1 \text{ and } s_u(\tilde{m}^2 \to \infty) \Longrightarrow \frac{1}{3\tilde{m}^4}$$

$$k\partial_k \lambda_{h|H} = -2\tilde{h}\partial_{\tilde{h}}\lambda_h(\tilde{h}) - 2\eta_H \lambda_h(\tilde{h}).$$

$$2\tilde{m}_W^2 + \tilde{m}_Z^2 + \tilde{m}_h^2 > 4\left(\tilde{m}_t^2 + \tilde{m}_b^2\right)$$

$\tilde{m}_h^2 = 2\lambda_h\tilde{h} = 2\partial_{\tilde{h}}^2 u$, and $\tilde{h}\partial_{\tilde{h}}^3 u$ is neglected

$$U = m_h^2 H^\dagger H + \cdots = \tilde{m}_h^2 k^2 H^\dagger H + \cdots,$$

$$\tilde{m}_h^2 = \partial_{\tilde{h}} u|_{\tilde{h}=0} = \frac{3}{64\pi^2}\{4y_t^2 + 4y_b^2 + 4y_{ch}^2 + y_\tau^2 - \left(\frac{3}{2}g_2^2 + \frac{3}{10}g_1^2 + 2\partial_{\tilde{h}}^2 u(0)\right)\}$$

$$L(h) = \left(\bar{\lambda}_\chi + \bar{\lambda}_h h^2\right)/2, \text{ i.e. } \bar{\lambda}_m = 0$$

$$\tilde{\mu}_h^2 = \partial_{\tilde{h}} u(\tilde{h} = 0) > 0 \text{ (symmetric regime)}$$

$$\varphi_0^2 = \tilde{h}_0 k^2 \to 0 \text{ or } m_H^2 = \tilde{\mu}_h^2 k^2 \to 0$$

$$L(h)\tilde{\rho}^2 = \tilde{L}(h)\tilde{h}^2 = \frac{1}{2}\bar{\lambda}_\chi \tilde{\rho}^2 + \frac{1}{2}\bar{\lambda}_h \tilde{h}^2 + \bar{\lambda}_m \tilde{h}\tilde{\rho},$$

$$m_H^2 = \frac{1}{2}\bar{\lambda}_m \chi^2$$

$$\partial_{\tilde{\rho}}\partial_h u(h_0) + \hat{\lambda}_h \partial_{\tilde{\rho}} h_0 = 0.$$

$$\partial_{\tilde{\rho}}\partial_h u(\tilde{\rho}, h_0) = -\frac{3y_t^2}{16\pi^2(1 + y_t^2 h_0 \tilde{\rho})^2}.$$

$y_t^2 h_0 \tilde{\rho} \lesssim 1$, and settles to a constant $y_t^2 h_0 \tilde{\rho} \gg 1$

$$\partial_{\tilde{\rho}} h_0(\tilde{\rho}) = \frac{3y_t^2}{16\pi^2 \hat{\lambda}_h (1 + y_t^2 h_0 \tilde{\rho})^2}.$$

$$\lambda_h = \partial_{\tilde{h}^2} u = \frac{\hat{\lambda}_h}{\tilde{\rho}^2},$$

$$\tilde{\rho}\partial_{\tilde{\rho}} h_0 = \frac{3y_t^2}{16\pi^2 \lambda_h \tilde{\rho}(1 + y_t^2 h_0 \tilde{\rho})^2}$$

$$\frac{\partial \ln\, h_0}{\partial \ln\, \tilde{\rho}} = \frac{3}{16\pi^2 \lambda_h \tilde{m}_t^2 \tilde{\rho}^2}.$$

$$\bar{u}(\tilde{\rho}) = u(\tilde{\rho}, h_0(\tilde{\rho})).$$

$$\partial_h u(\tilde{\rho}, h_0(\tilde{\rho}) = 0,$$

$$\partial_{\tilde{\rho}} \bar{u}(\tilde{\rho}) = \partial_{\tilde{\rho}} u(\tilde{\rho}, h_0) + \partial_h u(\tilde{\rho}, h_0)\partial_{\tilde{\rho}} h_0 = \partial_{\tilde{\rho}} u(\tilde{\rho}, h_0),$$

$$\partial_{\tilde{\rho}}^2 \bar{u}(\tilde{\rho}) = \partial_{\tilde{\rho}}^2 u(\tilde{\rho}, h_0) + 2\partial_{\tilde{\rho}}\partial_h u(\tilde{\rho}, h_0)\partial_{\tilde{\rho}} h_0 + \partial_h^2 u(\tilde{\rho}, h_0)\left(\partial_{\tilde{\rho}} h_0\right)^2$$

$$\partial_{\tilde{\rho}}^2 \bar{u}(\tilde{\rho}) = \partial_{\tilde{\rho}}^2 u(\tilde{\rho}, h_0) - \frac{1}{\hat{\lambda}_h}\left(\partial_{\tilde{\rho}}\partial_h u(\tilde{\rho}, h_0)\right)^2.$$

$$\bar{m}_c^2 = \partial_\rho \bar{u} + 2\tilde{\rho}\partial_{\tilde{\rho}}^2 \bar{u} = \partial_{\tilde{\rho}} u(\tilde{\rho}, h_0) + 2\tilde{\rho}\partial_{\tilde{\rho}}^2 u(\tilde{\rho}, h_0) - \frac{2\tilde{\rho}}{\hat{\lambda}_h}\left(\partial_{\tilde{\rho}}\partial_h u\right)^2,$$

$$\tilde{m}_{rc}^4 \ll (\hat{m}_r^2 - \tilde{m}_c^2)^2$$

$$\tilde{m}_-^2 = \tilde{m}_c^2 - \frac{\tilde{m}_{rc}^4}{\tilde{m}_r^2 - \tilde{m}_c^2},$$

$$\tilde{m}_c^2 = \left(\partial_{\tilde{\rho}} + 2\tilde{\rho}\partial_{\tilde{\rho}}^2\right)u(\tilde{\rho}, h_0) + h_0\tilde{m}_r^2 - 4h_0\partial_{\tilde{\rho}}\partial_h u(\tilde{\rho}, h_0).$$

$\tilde{m}_c^2 \ll \tilde{m}_r^2, \tilde{m}_r^2 = 2\hat{\lambda}_h h_0/\tilde{\rho}$ the mass eigenvalue $\tilde{m}_-^2$

$$\tilde{m}_f^2 = y_f^2 h_0\tilde{\rho} = 2h_f^2\tilde{\rho}$$

$$\tilde{m}_r^2 = 2\lambda_h h_0\tilde{\rho}$$

$$\bar{f}(\tilde{\rho}) = F(\tilde{\rho}, h_0(\tilde{\rho}))/k^2$$

$$\lim_{\tilde{\rho}\to\infty}\bar{u}(\tilde{\rho}) = u_\infty = \frac{5}{128\pi^2}$$

$$U_E = \frac{u_\infty h^4 M^4}{\xi^2\chi^4}$$

$$\lambda_{\chi,0} = 4\frac{\partial^2 U}{(\partial\chi^2)^2}, \lambda_{m,0} = 2\frac{\partial^2 U}{\partial\chi^2\partial(H^\dagger H)},$$
$$\lambda_{h,0} = \frac{\partial^2 U}{\partial(H^\dagger H)^2},$$

$$k\partial_k g_3^2 = \bar{\beta}_3 g_3^4$$

$$\bar{\beta}_3 = -\frac{33}{24\pi^2} + \frac{1}{12\pi^2}\sum_q \frac{k^2}{k^2 + m_q^2}.$$

$$\Gamma_k^{(g)} = \frac{1}{4}\int_x \sqrt{g'}Z_F F^{\mu\nu}F_{\mu\nu}$$

$$Z_F(\chi, H) = g^{-2}(\chi, H)$$

$$\tilde{h}\partial_{\tilde{h}} g_3^2 = -\frac{1}{2}\bar{\beta}_3 g_3^4$$

$$\bar{\beta}_3 = -\frac{33}{24\pi^2} + \frac{1}{12\pi^2}\sum_q \frac{1}{1+\tilde{m}_q^2} = -\frac{1}{24\pi^2}\left(33 - 2\sum_q \frac{1}{1+y_q^2\tilde{h}}\right)$$

$$\tilde{h}\partial_{\tilde{h}} g_3^{-2} = -\frac{7}{16\pi^2}$$

$$\frac{1}{g_3^2} = \frac{1}{\bar{g}_3^2} - \frac{7}{16\pi^2}\ln\frac{\tilde{h}}{\bar{h}} = \frac{7}{16\pi^2}\ln\frac{\tilde{h}_c}{\tilde{h}},$$

$$\bar{g}_3^2 = g_3^2(\tilde{h} = \bar{h}, h)$$

$$\tilde{h}_c = \exp\left(\frac{16\pi^2}{7\bar{g}_3^2}\right)\bar{h}$$

$$k_c^2 = \frac{H^\dagger H}{\tilde{h}_c} = \frac{H^\dagger H}{\bar{h}}\exp\left(-\frac{16\pi^2}{7\bar{g}_3^2}\right).$$

$$k_c^2 = \frac{\chi^2}{2\tilde{\rho}_c}$$

$$\bar{y}_q = y_q\left(\tilde{h} = \bar{y}_q^{-2}\right).$$

$$\tilde{h}\partial_{\tilde{h}} g_3^{-2} = -\frac{33}{48\pi^2} + \frac{1}{24\pi^2}\sum_q \frac{1}{1+\bar{y}_q^2\tilde{h}}$$

$$g_3^{-2}(\tilde{h}) = \frac{1}{\bar{g}_3^2} - \frac{7}{16\pi^2}\ln\left(\frac{\tilde{h}}{\bar{h}}\right) - \frac{1}{24\pi^2}\sum_q \ln\left(\frac{1+\bar{y}_q^2\tilde{h}}{1+\bar{y}_q^2\bar{h}}\right)$$

$$\Lambda_{\mathrm{QCD}}^2 = \frac{H^\dagger H}{\tilde{h}_c}$$

$$u = \bar{u}_1(\tilde{h}) + \bar{u}_2(\tilde{\rho}) + \bar{\lambda}_m\tilde{h}\tilde{\rho}$$

$$U = \frac{1}{8}\bar{\lambda}_\chi\chi^4 + \frac{1}{2}\bar{\lambda}_m\chi^2 H^\dagger H + \frac{1}{2}\bar{\lambda}_h(1+\Delta)\left(H^\dagger H\right)^2$$

$$\bar{\beta}_3 = -\frac{c}{24\pi^2}$$

$$y_t^2 h\tilde{\rho}_t = 1, \tilde{\rho}_t = \frac{1}{y_t^2 h}$$

$$g_3^{-2}(\tilde{\rho}) = g_{3,0}^{-2} - \frac{7}{16\pi^2}\ln\frac{\tilde{\rho}_t}{\tilde{\rho}_0} - \frac{23}{48\pi^2}\ln\frac{\tilde{\rho}_b}{\tilde{\rho}_t} - \frac{25}{48\pi^2}\ln\frac{\tilde{\rho}_{ch}}{\tilde{\rho}_b} - \frac{9}{16\pi^2}\ln\frac{\tilde{\rho}}{\tilde{\rho}_{ch}}$$

$$\hat{g}_3^{-2} = g_{3,0}^{-2} + \frac{7}{16\pi^2}\ln\frac{y_t^2}{y_{ch}^2} - \frac{23}{48\pi^2}\ln\frac{y_t^2}{y_b^2} - \frac{25}{48\pi^2}\ln\frac{y_b^2}{y_{ch}^2} - \frac{1}{8\pi^2}\ln y_{ch}^2$$

$$g_3^{-2}(\tilde{\rho}, h) = \hat{g}_3^{-2} - \frac{1}{8\pi^2}\ln(h\tilde{\rho}_0) - \frac{9}{16\pi^2}\ln\frac{\tilde{\rho}}{\tilde{\rho}_0}$$

$$\ln \frac{\tilde{\rho}_c}{\tilde{\rho}_0} = \frac{16\pi^2}{9\hat{g}_3^2} - \frac{2}{9}\ln(h\tilde{\rho}_0)$$

$$\tilde{\rho}_c = \exp\left(\frac{16\pi^2}{9\hat{g}_3^2}\right) h^{-\frac{2}{9}} \tilde{\rho}_0^{\frac{7}{9}}$$

$$\Lambda_{\mathrm{QCD}}^2 = \frac{\chi^2}{2\tilde{\rho}_c} = \exp\left(-\frac{16\pi^2}{9\hat{g}_3^2}\right)\left(H^\dagger H\right)^{\frac{2}{9}} k_0^{\frac{14}{9}},$$

$$\Lambda_{\mathrm{QCD}}^2 = \exp\left(-\frac{16\pi^2}{9g_{3,0}^2}\right)(2\tilde{\rho}_0)^{-\frac{7}{9}} A_q \left(H^\dagger H\right)^{\frac{2}{9}} \chi^{\frac{14}{9}},$$

$$A_q = \exp\left\{-\frac{1}{27}\left(21\ln\frac{y_t^2}{y_{ch}^2} - 23\ln\frac{y_t^2}{y_b^2} - 25\ln\frac{y_b^2}{y_{ch}^2} - 6\ln y_{ch}^2\right)\right\}$$

$$m_{\mathrm{QCD}}^2 = c_s \Lambda_{\mathrm{QCD}}^2, \tilde{m}_{\mathrm{QCD}}^2 = \frac{c_s\tilde{\rho}}{\tilde{\rho}_c}$$

$$c_U = -\frac{1}{128\pi^2}\left\{\frac{12}{1+y_t^2\tilde{h}} + \frac{12}{1+y_b^2\tilde{h}} + \frac{12}{1+y_{ch}^2\tilde{h}} + \frac{4}{1+y_\tau^2\tilde{h}} + \frac{12}{1+\tilde{m}_s^2} - \frac{6}{1+\tilde{m}_W^2} - \frac{3}{1+\tilde{m}_Z^2} - \frac{1}{1+\tilde{m}_r^2} - \frac{3}{1+\partial_h u/\tilde{\rho}} + 20\right\}$$

$$\tilde{m}_W^2 = \frac{g_2^2\tilde{h}}{2}, \quad \tilde{m}_Z^2 = \left(\frac{g_2^2}{2} + \frac{3g_1^2}{10}\right)\tilde{h},$$

$$\tilde{m}_r^2 = \frac{\partial_h u + 2h\partial_{\tilde{h}}^2 u}{\tilde{\rho}}, \quad \tilde{m}_s^2 = \tilde{m}_{\mathrm{QCD}}^2 + y_s^2\tilde{h}.$$

$$k\partial_k y_t^2 = \beta_t = \frac{k^2}{16\pi^2(k^2+m_t^2)}\left[9y_t^4 - \left(16g_3^2 + \frac{9}{2}g_2^2 + \frac{17}{10}g_1^2\right)y_t^2\right]$$

$$R_t = \left(g_3^2 + \frac{9}{32}g_2^2 + \frac{17}{160}g_1^2\right)/y_t^2$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\left(\frac{y_t^2}{g_3^2}\right) = -\frac{9y_t^2}{32\pi^2}\left(\frac{y_t^2}{g_3^2} - \frac{2}{9}\right),$$

$$\tilde{\rho}\partial_{\tilde{\rho}} y_{t|h}^2 = -\frac{b_t y_t^4}{2(1+y_t^2 h\tilde{\rho})}, b_t = \frac{9-16R_t}{16\pi^2}$$

$$y_t^{-2}(\tilde{\rho}, h) = y_t^{-2}(\tilde{\rho}_0, h) + \frac{b_t}{2}\left[\ln\left(\frac{\tilde{\rho}}{1+\bar{y}_t^2 h\tilde{\rho}}\right) - \ln\left(\frac{\tilde{\rho}_0}{1+\bar{y}_t^2 h\tilde{\rho}_0}\right)\right]$$

$$\Gamma_{2\chi,2t,H} = \frac{\partial y_t}{\partial\rho} = \frac{2}{\chi^2}\tilde{\rho}\partial_{\tilde{\rho}} y_t = -\frac{1}{2y_t\chi^2}\beta_t.$$

$$\Gamma_{2\chi,2A} \sim k^2 \frac{\partial Z_F}{\partial \rho} = \frac{\partial Z_F}{\partial \tilde{\rho}} = \frac{1}{\tilde{\rho}} \left(\tilde{\rho} \partial_{\tilde{\rho}} g_3^{-2} \right) = \frac{1}{2\tilde{\rho}} \left(\bar{\beta}_3 + \cdots \right)$$

$$k\partial_k g^2 = -B_g g^2 - B_F g^4$$

$$M_p^2(k) = 2w_0 k^2 + \xi \chi^2 = 2w(\tilde{\rho}) k^2.$$

$$B_g \approx \frac{5k^2}{144\pi^2 M_p^2(k)} \left(\frac{4}{1 - v(k)} - \frac{3}{(1 - v(k))^2} \right),$$

$$v(k) = \frac{u(k)}{w(k)}$$

$$B_g(\tilde{\rho} \gg 1) \approx \frac{5}{288\pi^2 \xi \tilde{\rho}}$$

$$B_g \approx \frac{5}{288\pi^2 w_0} \left(\frac{4}{1 - v_0} - \frac{3}{(1 - v_0)^2} \right), v_0 = \frac{u_0}{w_0}.$$

$$\tilde{\rho} \partial_{\tilde{\rho}} g_{|h}^{-2} = -\frac{B_g}{2} g^{-2} - \frac{B_F}{2}$$

$$g^{-2}(\tilde{\rho}) = \left(\frac{\tilde{\rho}}{\hat{\rho}} \right)^{-\frac{1}{2} B_g} - \frac{B_F}{B_g}.$$

$$g^{-2}(\tilde{\rho}) = -\left(\frac{\tilde{\rho}}{\hat{\rho}} \right)^{-\frac{1}{2} B_g} - \frac{B_F}{B_g},$$

$$g^{-2} = \hat{g}^{-2} - \frac{B_F}{2} \ln \frac{\tilde{\rho}}{\hat{\rho}}$$

$$\tilde{\rho} \partial_{\tilde{\rho}} \partial_h g_3^{-2} = -\frac{1}{2} \partial_h B_F = \frac{1}{2} \partial_h \bar{\beta}_3 = \frac{1}{24\pi^2} \partial_h (1 + y_t^2 h \tilde{\rho})^{-1}$$

$$\tilde{\rho} \partial_{\tilde{\rho}} \partial_h g_3^{-2} = -\frac{y_t^2 \tilde{\rho}}{24\pi^2 (1 + y_t^2 h \tilde{\rho})^2}$$

$$\partial_h g_3^{-2} = -\frac{\bar{y}_t^2 \tilde{\rho}}{24\pi^2}$$

$$g_3^{-2}(\tilde{\rho}, h) = g_3^{-2}(\tilde{\rho}, 0) - \frac{\bar{y}_t^2 h \tilde{\rho}}{24\pi^2}$$

$$\partial_h g_3^{-2} = -\frac{\bar{y}_t^2 \tilde{\rho}}{24\pi^2 (1 + \bar{y}_t^2 h \tilde{\rho})},$$

$$g_3^{-2}(\tilde{\rho}, h) = g_3^{-2}(\tilde{\rho}, 0) - \frac{1}{24\pi^2} \ln (1 + \bar{y}_t^2 h \tilde{\rho})$$

$$g_3^2(\tilde{h} = h\tilde{\rho}, h) = g_3^2(\tilde{\rho}, h)$$

$$g_{3,0}^2 = g_3^2(\tilde{\rho}_0, h) = g_3^2\big(\tilde{h} = h\tilde{\rho}_0, h\big)$$

$$\bar{g}_3^2 = g_3^2(\tilde{h} = \bar{h}, h)$$

$$\bar{g}_3^{-2} = g_{3,0}^{-2} + \frac{1}{2}\bar{\beta}_3 \ln\left(\frac{\bar{h}}{h\tilde{\rho}_0}\right).$$

$$h\partial_h \bar{g}_3^{-2} = -\frac{1}{2}\bar{\beta}_3$$

$$g_3^2(\tilde{h} = \bar{h}, h) = \bar{g}_3^2(h)$$

$$u = L(h)\tilde{\rho}^2 + \frac{1}{128\pi^2}\sum_i \hat{n}_i t_u^{(i)}\big(\tilde{m}_i^2\big)$$

$$\tilde{\rho}\partial_{\tilde{\rho}} t_u^{(i)} = 2\left(t_u^{(i)} - \frac{1}{1+\tilde{m}_i^2}\right)$$

$$\tilde{v}(\tilde{\rho}, h) = \partial_h u(\tilde{\rho}, h) = \tilde{\rho}\mu_h(\tilde{\rho}, h),$$

$$\tilde{v}(\tilde{\rho}) = \tilde{v}(\tilde{\rho}, 0).$$

$$\mu_h = \partial_{\tilde{h}}\mu(\tilde{\rho}, \tilde{h})$$

$$\lambda_m(\tilde{\rho}, h) = \frac{1}{\tilde{\rho}^2}\big(\tilde{\rho}\partial_{\tilde{\rho}} - h\partial_h - 1\big)\tilde{v}(\tilde{\rho}, h)$$

$$\lambda_m(\tilde{\rho}) = \lambda_m(\tilde{\rho}, 0) = \left(\frac{1}{\tilde{\rho}}\partial_{\tilde{\rho}} - \frac{1}{\tilde{\rho}^2}\right)\tilde{v}(\tilde{\rho}) = \partial_{\tilde{\rho}}\left(\frac{\tilde{v}(\tilde{\rho})}{\tilde{\rho}}\right).$$

$$\partial_h u_{|\tilde{\rho}} = \tilde{v} + \lambda_h h_0 \tilde{\rho}^2 = 0,$$

$$\varphi_0^2 = \frac{1}{2}h_0(\tilde{\rho} \to \infty)\chi^2,$$
$$h_0 = \frac{2\varphi_0^2}{\chi^2} = -\frac{\tilde{v}(\tilde{\rho})}{\lambda_h(\tilde{\rho})\tilde{\rho}^2}(\tilde{\rho} \to \infty).$$

$$v(\tilde{\rho}, h) = \tilde{v}(\tilde{\rho}, h) - \partial_h L(h)\tilde{\rho}^2 = \frac{1}{128\pi^2}\sum_i \hat{n}_i \partial_h \tilde{m}_i^2 \frac{\partial t_u^{(i)}}{\partial \tilde{m}_i^2}.$$

$$v = -\frac{1}{64\pi^2}\sum_i \hat{n}_i \partial_h \tilde{m}_i^2 s_u^{(i)}\big(\tilde{m}_i^2\big),$$

$$s_u^{(i)}\big(\tilde{m}_i^2\big) = -\frac{1}{2}\frac{\partial t_u^{(i)}}{\partial \tilde{m}_i^2} = 1 + \frac{\tilde{m}_i^2}{1+\tilde{m}_i^2} - 2\tilde{m}_i^2 \ln\frac{1+\tilde{m}_i^2}{\tilde{m}_i^2}$$

$$\lim_{\tilde{m}_i^2 \to \infty} s_u^{(i)}\big(\tilde{m}_i^2\big) = \frac{1}{3\tilde{m}_i^4}$$

$$\lim_{\tilde{\rho}\to\infty} v = 0, \lim_{\tilde{\rho}\to\infty} \tilde{v}(\tilde{\rho},h) = \partial_h L\tilde{\rho}^2.$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\tilde{v}(\tilde{\rho},h) = \partial_h\left(\tilde{\rho}\partial_{\tilde{\rho}}u\right) = 2\partial_h(u-c_U) = 2\tilde{v} - 2\partial_h c_U.$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\tilde{v}(\tilde{\rho},h) = 2\partial_h L + \frac{1}{64\pi^2}\sum_i \hat{n}_i\partial_h\left(t_u^{(i)} - \frac{1}{1+\tilde{m}_i^2}\right),$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\tilde{v}(\tilde{\rho},h) = 2\tilde{v}(\tilde{\rho},h) - B(\tilde{\rho},h),$$

$$B = 2\partial_h c_U = \frac{1}{64\pi^2}\sum_i \hat{n}_i\partial_h\left(1+\tilde{m}_i^2\right)^{-1} = -\frac{1}{64\pi^2}\sum_i \hat{n}_i\partial_h\tilde{m}_i^2\left(1+\tilde{m}_i^2\right)^{-2}$$

$$\tilde{m}_i^2 = \alpha_i(\tilde{h},h)\tilde{h} = \alpha_i h\tilde{\rho}$$

$$\partial_h\tilde{m}_{i|\tilde{\rho}}^2 = \partial_h\tilde{m}_{i|\tilde{h}}^2 + \tilde{\rho}\partial_{\tilde{h}}\tilde{m}_{i|h}^2 = \alpha_i\left(1 + \frac{\partial\ln\,\alpha_i}{\partial\ln\,\tilde{h}} + \frac{\partial\ln\,\alpha_i}{\partial\ln\,h}\right)\tilde{\rho}$$

$$\tilde{m}_i^2 = \alpha_i(\tilde{\rho},h)h\tilde{\rho}$$

$$\partial_h\tilde{m}_i^2(\tilde{\rho},h) = \alpha_i(\tilde{\rho},h)\left(1 + \frac{\partial\ln\,\alpha_i(\tilde{\rho},h)}{\partial\ln\,h}\right)\tilde{\rho}$$

$$\partial_{\tilde{\rho}}\left(\frac{\tilde{v}}{\tilde{\rho}^2}\right) = \frac{\beta_v}{\tilde{\rho}^2},$$

$$\tilde{v}(\tilde{\rho},h) = \partial_h L\tilde{\rho}^2 - \tilde{\rho}^2\int_{\tilde{\rho}}^{\infty} \mathrm{d}\rho'\,\frac{\beta_v(\rho',h)}{\rho'^2}$$

$$\partial_h L = \lim_{\tilde{\rho}\to\infty}\frac{\tilde{v}(\tilde{\rho},h)}{\tilde{\rho}^2}$$

$$\beta_v = \frac{1}{64\pi^2}\sum_i \hat{n}_i\alpha_i(1+\alpha_i h\tilde{\rho})^{-2}$$

$$\tilde{v} = -\beta_v\tilde{\rho} + \partial_h L\tilde{\rho}^2$$

$$\beta_v = \frac{1}{64\pi^2}\sum_i \frac{\hat{n}_i}{\alpha_i h^2\tilde{\rho}^2}$$

$$\tilde{v}(\tilde{\rho},h) = \partial_h L\tilde{\rho}^2 - \frac{1}{192\pi^2}\sum_i \frac{\hat{n}_i}{\alpha_i h^2\tilde{\rho}}$$

$$\tilde{v}(\tilde{\rho},h) = \partial_h L\tilde{\rho}^2 - \tilde{\beta}_v(\tilde{\rho},h)\tilde{\rho}$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\tilde{\beta}_v = \tilde{\beta}_v - \beta_v$$

$$\tilde{\beta}_\nu(\tilde{h},h)=\tilde{h}\int_{\tilde{h}}^{\infty}\frac{\mathrm{d}h'}{h'^2}\beta_\nu(h',h)\approx\frac{\tilde{h}}{64\pi^2}\sum_i\hat{n}_i\int_{\tilde{h}}^{\infty}\frac{\mathrm{d}h'\alpha_i}{h'^2(1+\alpha_i h')^2}.$$

$$\tilde{\beta}_\nu=\frac{1}{64\pi^2}\sum_i\hat{n}_i\alpha_i\left[1+\alpha_i\tilde{h}\left(2\ln\left(\frac{\alpha_i\tilde{h}}{1+\alpha_i\tilde{h}}\right)+\frac{1}{1+\alpha_i\tilde{h}}\right)\right]=\frac{1}{64\pi^2}\sum_i\hat{n}_i\alpha_i s_u(\tilde{m}_i^2)$$

$$\partial_h\tilde{\beta}_{\nu|\tilde{h}}=\frac{1}{64\pi^2}\sum_i\hat{n}_i\partial_h\alpha_{i|\tilde{h}}\left[1+4\alpha_i\tilde{h}\ln\left(\frac{\alpha_i\tilde{h}}{1+\alpha_i\tilde{h}}\right)+\frac{4\alpha_i\tilde{h}}{1+\alpha_i\tilde{h}}-\frac{\alpha_i^2\tilde{h}^2}{\left(1+\alpha_i\tilde{h}\right)^2}\right].$$

$$\tilde{\beta}_\nu=\frac{1}{192\pi^2\tilde{h}^2}\sum_i\frac{\hat{n}_i}{\alpha_i}$$

$$\partial_h\tilde{\beta}_{\nu|\tilde{h}}=-\frac{1}{192\pi^2\tilde{h}^2}\sum_i\frac{\hat{n}_i\partial_h\alpha_{i|\tilde{h}}}{\alpha_i^2}.$$

$$\frac{\partial U}{\partial(H^\dagger H)}=\frac{1}{2}\chi^2\partial_h L-\tilde{\beta}_\nu k^2,$$

$$\lambda_h=\frac{\partial^2 U}{\partial(H^\dagger H)^2}=\partial_h^2 L-\frac{1}{\tilde{\rho}}\partial_h\tilde{\beta}_{\nu|\tilde{\rho}}$$

$$\frac{\partial U}{\partial(H^\dagger H)}\approx\frac{\bar{\lambda}_m}{2}\chi^2-\frac{k^2}{64\pi^2}\sum_i\hat{n}_i\alpha_i.$$

$$\frac{1}{\tilde{\rho}}\partial_h\tilde{\beta}_\nu(\tilde{\rho},h)=\lambda_h(\tilde{\rho},h)-\bar{\lambda}_h$$

$$=\frac{1}{64\pi^2}\sum_i\hat{n}_i\left\{\alpha_i^2\left[2\ln\left(\frac{\tilde{m}_i^2}{1+\tilde{m}_i^2}\right)\right.\right.$$

$$\left.\left.+\frac{3+2\tilde{m}_i^2}{\left(1+\tilde{m}_i^2\right)^2}\right]+\frac{1}{\tilde{\rho}}\partial_h\alpha_{i|\tilde{\rho}}\left[1+\tilde{m}_i^2\left(4\ln\left(\frac{\tilde{m}_i^2}{1+\tilde{m}_i^2}\right)+\frac{4+3\tilde{m}_i^2}{\left(1+\tilde{m}_i^2\right)^2}\right)\right]\right\}$$

$$\frac{1}{\tilde{\rho}}\partial_h y_{t|\tilde{\rho}}^2\approx\frac{b_t\bar{y}_t^2y_t^4}{2(1+\bar{y}_t^2h\tilde{\rho})}$$

$$\tilde{v}(\tilde{\rho},h)=\partial_h L(h)\tilde{\rho}^2+\frac{3y_t^2}{16\pi^2}\tilde{\rho}s_u(y_t^2h\tilde{\rho}),$$

$$\partial_h L(h)=\frac{\tilde{v}(\tilde{\rho}_0,h)}{\tilde{\rho}_0^2}-\frac{3y_t^2}{16\pi^2\tilde{\rho}_0}s_u(y_t^2h\tilde{\rho}_0).$$

$$\bar{\lambda}_h(h) = \partial_h^2 L(h) = \frac{\partial_h \tilde{v}(\tilde{\rho}_0, h)}{\tilde{\rho}_0^2} - \frac{3y_t^4}{16\pi^2}\frac{\partial s_u(\tilde{m}_t^2)}{\partial \tilde{m}_t^2}$$

$$= \frac{\partial_h \tilde{v}(\tilde{\rho}_0, h)}{\tilde{\rho}_0^2} - \frac{3y_t^4}{8\pi^2}\left[\ln\left(\frac{\tilde{m}_{t,0}^2}{1+\tilde{m}_{t,0}^2}\right) + \frac{1}{1+\tilde{m}_{t,0}^2} + \frac{1}{2\left(1+\tilde{m}_{t,0}^2\right)^2}\right]$$

$$\bar{\lambda}_h(h) = \frac{3y_t^4}{8\pi^2}\left[\ln\left(\frac{1+y_t^2 h\tilde{\rho}_0}{y_t^2 h\tilde{\rho}_0}\right) - \frac{1}{1+y_t^2 h\tilde{\rho}_0} - \frac{1}{2(1+y_t^2 h\tilde{\rho}_0)^2}\right]$$

$$\bar{\lambda}_h(h) \approx \frac{3y_t^4}{8\pi^2}\ln\left(\frac{1}{y_t^2 h\tilde{\rho}_0}\right).$$

$$\bar{\lambda}_h(h) = \frac{1}{8\pi^2 y_t^2 h^3 \tilde{\rho}_0^3}$$

$$\lambda_m(\tilde{\rho}, \tilde{h}) = \partial_{\tilde{\rho}}\partial_{\tilde{h}} u(\tilde{\rho}, \tilde{h})$$

$$\lambda_m(\tilde{\rho}, \tilde{h}) = \partial_{\tilde{\rho}}\mu_h(\tilde{\rho}, \tilde{h}), \mu_h(\tilde{\rho}, \tilde{h}) = \partial_{\tilde{h}} u(\tilde{\rho}, \tilde{h})$$

$$\lambda_m(\tilde{\rho}, h) = \frac{1}{\tilde{\rho}^2}\partial_h\left(\tilde{\rho}\partial_{\tilde{\rho}} - h\partial_h\right)u(\tilde{\rho}, h)$$

$$\partial_{\tilde{\rho}|\tilde{h}} = \partial_{\tilde{\rho}|h} + \frac{\partial h}{\partial\tilde{\rho} \mid \tilde{h}}\partial_{h|\tilde{\rho}} = \partial_{\tilde{\rho}|h} - \frac{h}{\tilde{\rho}}\partial_{h|\tilde{\rho}}$$

$$\tilde{m}_{rc}^2 = 2\sqrt{\tilde{h}\tilde{\rho}}\lambda_m(\tilde{\rho}, \tilde{h})$$

$$\lambda_m(\tilde{\rho} \to 0, h = 0) = \lambda_{m,0}$$
$$\lambda_m(\tilde{\rho} \to \infty, h = 0) = \bar{\lambda}_m$$

$$\lambda_m(\tilde{\rho}) = \lambda_m(\tilde{\rho}, h = 0).$$

$$\lambda_m(\tilde{\rho}, h) = \partial_{\tilde{\rho}|\tilde{h}}\left(\frac{\tilde{v}}{\tilde{\rho}}\right) = \left(\partial_{\tilde{\rho}|h} - \frac{h}{\tilde{\rho}}\partial_{h|\tilde{\rho}}\right)\left(\frac{\tilde{v}}{\tilde{\rho}}\right).$$

$$\lambda_m(\tilde{\rho}, h) = \partial_h L - h\partial_h^2 L + \lambda_m^{(\mathrm{cr})}(\tilde{\rho}, h).$$

$$\lambda_m^{(\mathrm{cr})}(\tilde{\rho}, h) = -\partial_{\tilde{\rho}}\tilde{\beta}_{v|\tilde{h}} = \frac{h}{\tilde{\rho}}\partial_h\tilde{\beta}_{v|\tilde{h}} = \frac{h^2}{\tilde{h}}\partial_h\tilde{\beta}_{v|\tilde{h}}$$

$$\lambda_m^{(\mathrm{cr})}(\tilde{\rho}) = \lambda_m^{(\mathrm{cr})}(\tilde{\rho}, h = 0)$$

$$\lambda_m^{(\mathrm{cr})} = \frac{1}{64\pi^2\tilde{\rho}}\sum_i \hat{n}_i h\partial_h\alpha_{i|\tilde{h}} = -\frac{1}{64\pi^2\tilde{\rho}}\sum_i \hat{n}_i\tilde{\rho}\partial_{\tilde{\rho}}\alpha_{i|\tilde{h}=0} = \frac{1}{128\pi^2\tilde{\rho}}\sum_i \hat{n}_i\beta_i$$

$$\beta_i = -2\tilde{\rho}\partial_{\tilde{\rho}}\alpha_i = k\partial_k\alpha_{i|\rho}$$

$$\lambda_m^{(\mathrm{cr})}(\tilde{\rho},h)=-\frac{1}{64\pi^2\tilde{\rho}}\sum_i \hat{n}_i\left(\tilde{\rho}\partial_{\tilde{\rho}}-h\partial_h\right)\alpha_i T_i\left(\alpha_i\tilde{h}\right),$$

$$T_i\left(\alpha_i\tilde{h}\right)=1+4\alpha_i\tilde{h}\left(\ln\frac{\alpha_i\tilde{h}}{1+\alpha_i\tilde{h}}+\frac{1}{1+\alpha_i\tilde{h}}\right)-\frac{\alpha_i^2\tilde{h}^2}{\left(1+\alpha_i\tilde{h}\right)^2}$$

$$\left(\tilde{\rho}\partial_{\tilde{\rho}}-h\partial_h\right)y_t^2\approx-\frac{b_t}{2}y_t^4$$

$$\lambda_m^{(\mathrm{cr})t}=-\frac{3\beta_t}{32\pi^2\tilde{\rho}},$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m=-2\lambda_m+\frac{1}{\tilde{\rho}^2}\partial_h\left(\tilde{\rho}\partial_{\tilde{\rho}}-h\partial_h\right)\tilde{\rho}\partial_{\tilde{\rho}}u=-2\lambda_m+\frac{2}{\tilde{\rho}^2}\partial_h\left(\tilde{\rho}\partial_{\tilde{\rho}}-h\partial_h\right)(u-c_U)$$

$$=-\frac{2}{\tilde{\rho}^2}\partial_h\left(\tilde{\rho}\partial_{\tilde{\rho}}-h\partial_h\right)c_U=-2\left[\left(\frac{1}{\tilde{\rho}}\partial_{\tilde{\rho}}-\frac{1}{\tilde{\rho}^2}\right)\partial_h-\frac{h}{\tilde{\rho}^2}\partial_h^2\right]c_U.$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m(\tilde{\rho},h)=-2\partial_{\tilde{\rho}}\partial_{\tilde{h}}c_U(\tilde{\rho},\tilde{h})$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m(\tilde{\rho},h)=\frac{1}{64\pi^2}\partial_{\tilde{\rho}}\{6\partial_{\tilde{h}}\tilde{m}_W^2(1+\tilde{m}_W^2)^{-2}+3\partial_{\tilde{h}}\tilde{m}_Z^2(1+\tilde{m}_Z^2)^{-2}$$

$$+\sum_s \partial_{\tilde{h}}\tilde{m}_s^2(1+\tilde{m}_s^2)^{-2}\ -2\sum_f n_f\partial_{\tilde{h}}\tilde{m}_f^2\left(1+\tilde{m}_f^2\right)^{-2}\Bigg\}$$

$$=\frac{1}{64\pi^2}\partial_{\tilde{\rho}}\sum_i \hat{n}_i\partial_{\tilde{h}}\tilde{m}_i^2\left(1+\tilde{m}_i^2\right)^{-2}$$

$$=\frac{1}{64\pi^2}\sum_i \hat{n}_i\left\{\partial_{\tilde{\rho}}\partial_{\tilde{h}}\tilde{m}_i^2\left(1+\tilde{m}_i^2\right)^{-2}\ -2\partial_{\tilde{h}}\tilde{m}_i^2\partial_{\tilde{\rho}}\tilde{m}_i^2\left(1+\tilde{m}_i^2\right)^{-3}\right\}$$

$$\partial_{\tilde{\rho}}\tilde{m}_{i|\tilde{h}}^2=h\left(\tilde{\rho}\partial_{\tilde{\rho}}-h\partial_h\right)\alpha_i(\tilde{\rho},h)$$
$$\partial_{\tilde{h}}\tilde{m}_{i|\tilde{\rho}}^2=\alpha_i(\tilde{\rho},h)+h\partial_h\alpha_i(\tilde{\rho},h)$$
$$\partial_{\tilde{\rho}}\partial_{\tilde{h}}\tilde{m}_i^2=\frac{1}{\tilde{\rho}}\left(\tilde{\rho}\partial_{\tilde{\rho}}-h\partial_h\right)(1+h\partial_h)\alpha_i(\tilde{\rho},h)$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m=-\frac{1}{128\pi^2\tilde{\rho}}\sum_i{}' \hat{n}_i\beta_i+\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m^{(g)}+\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m^{(r)}$$

$$\tilde{m}_g^2=\partial_{\tilde{h}}u_{|\tilde{\rho}}$$

$$\lambda_m(\tilde{\rho},h)=\partial_{\tilde{\rho}|\tilde{h}}\partial_{\tilde{h}|\tilde{\rho}}u,\lambda_h(\tilde{\rho},h)=\partial_{\tilde{h}|\tilde{\rho}}^2u$$

$$\partial_{\tilde{\rho}}\tilde{m}^2_{g|\tilde{h}}= \lambda_m, \partial_{\tilde{h}}\tilde{m}^2_{g|\tilde{\rho}} = \lambda_h$$

$$\partial_{\tilde{\rho}}\partial_{\tilde{h}}\tilde{m}^2_g(\tilde{\rho},\tilde{h}) = \frac{1}{\tilde{\rho}}\left(\tilde{\rho}\partial_{\tilde{\rho}} - h\partial_h\right)\lambda_h(\tilde{\rho},h)$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m^{(g)} = \frac{3}{64\pi^2\tilde{\rho}}\left(\tilde{\rho}\partial_{\tilde{\rho}} - h\partial_h\right)\lambda_h\left(1+\tilde{m}_g^2\right)^{-2} - \frac{3}{32\pi^2}\lambda_m\lambda_h\left(1+\tilde{m}_g^2\right)^{-3}$$

$$\tilde{m}_r^2 = \tilde{m}_g^2 + 2\lambda_h\tilde{h}$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m^{(r)} = \frac{1}{64\pi^2\tilde{\rho}}\left(\tilde{\rho}\partial_{\tilde{\rho}} - h\partial_h\right)(3+2h\partial_h)\lambda_h(1+\tilde{m}_r^2)^{-2}$$

$$-\frac{1}{32\pi^2}\left(\lambda_m + 2h\left(\tilde{\rho}\partial_{\tilde{\rho}} - h\partial_h\right)\lambda_h\right)\times(3+2h\partial_h)\lambda_h(1+\tilde{m}_r^2)^{-3}$$

$$\lambda_m(\tilde{\rho},h) = \lambda_m^{(\mathrm{cr})}(\tilde{\rho},h) + \delta\lambda_m$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m = -\frac{1}{2}\beta_m = -\frac{1}{2}\left(\tilde{\beta}_m + A_m\lambda_m\right),$$

$$\left(\tilde{\rho}\partial_{\tilde{\rho}}\lambda_h = -\beta_h/2\right)$$

$$\tilde{\beta}_m = \frac{1}{64\pi^2\tilde{\rho}}\left\{\sum_i{}' \hat{n}_i\frac{\beta_i}{\left(1+\tilde{m}_i^2\right)^2} + 3\beta_h\left(\frac{1}{\left(1+\tilde{m}_g^2\right)^2} + \frac{1}{(1+\tilde{m}_r^2)^2}\right)\right\}$$

$$A_m = \frac{3\lambda_h}{16\pi^2}\left(\frac{1}{\left(1+\tilde{m}_g^2\right)^3} + \frac{1}{(1+\tilde{m}_r^2)^3}\right) + \eta_H$$

$$\lambda_m = \frac{1}{2}(1+g)\tilde{\beta}_m.$$

$$\tilde{\beta}_m= \frac{b_m}{64\pi^2\tilde{\rho}}$$

$$b_m = \sum_i{}' \hat{n}_i\beta_i + \frac{6\beta_h}{(1+\mu_h)^2}$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\tilde{\beta}_m= -\tilde{\beta}_m - \frac{d_m}{128\pi^2\tilde{\rho}}$$

$$d_m = -2\tilde{\rho}\partial_{\tilde{\rho}}b_m$$

$$\frac{1}{2}(1+g)\tilde{\rho}\partial_{\tilde{\rho}}\tilde{\beta}_m + \frac{1}{2}\tilde{\beta}_m\tilde{\rho}\partial_{\tilde{\rho}}g = -\frac{1}{2}\left(\tilde{\beta}_m + A_m\frac{1+g}{2}\tilde{\beta}_m\right)$$

$$2\tilde{\beta}_m\tilde{\rho}\partial_{\tilde{\rho}}g = (2g - A_m(1+g))\tilde{\beta}_m + \frac{(1+g)d_m}{64\pi^2\tilde{\rho}}.$$

$$\tilde{\rho}\partial_{\tilde{\rho}}g= g + (1+g)f_m$$

$$f_m = \frac{1}{2}\left(\frac{d_m}{b_m} - A_m\right)$$

$$g_1(\tilde{\rho}) = \left(g(\tilde{\rho}_0) + \frac{f_m}{1+f_m}\right)\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{1+f_m} - \frac{f_m}{1+f_m}.$$

$$g_1^{(\mathrm{cr})}(\tilde{\rho}) = -\frac{f_m(\tilde{\rho})}{1+f_m(\tilde{\rho})}$$

$$\lambda_m^{(\mathrm{cr})}(\tilde{\rho}) = \frac{b_m(\tilde{\rho})}{128\pi^2\tilde{\rho}(1+f_m(\tilde{\rho}))}$$

$$\lambda_m(\tilde{\rho}) = \frac{1}{2}\tilde{\beta}_m\left(1 - \frac{f_m(\tilde{\rho})}{1+f_m(\tilde{\rho})} + h(\tilde{\rho})\right)$$

$$\lambda_m^{(\mathrm{cr})}(\tilde{\rho}) = \frac{\tilde{\beta}_m(\tilde{\rho})}{1-f_m(\tilde{\rho})}\left(1 - \frac{\tilde{\rho}\partial_{\tilde{\rho}} f_m(\tilde{\rho})}{(1-f_m(\tilde{\rho}))^2}\right)$$

$$\begin{aligned}
b_m =& -12\beta_t + 6\beta_h + \frac{9}{2}\beta_2 + \frac{9}{10}\beta_1 \\
=& \frac{1}{16\pi^2}\left\{-12\left[9y_t^4 - \left(16g_3^2 + \frac{9}{2}g_2^2 + \frac{17}{10}g_1^2\right)y_t^2\right]\right. \\
&+6\left[12\lambda_h^2 - 12y_t^4 + \frac{9}{4}g_2^4 + \frac{9}{10}g_2^2g_1^2 + \frac{27}{100}g_1^4 + 12y_t^2\lambda_h\right. \\
&\left.\left.-9g_2^2\lambda_h - \frac{9}{5}g_1^2\lambda_h\right] + \frac{9}{2}\left[-\frac{19}{3}g_2^4\right] + \frac{9}{10}\left[\frac{41}{5}g_1^4\right]\right\} \\
=& \frac{1}{16\pi^2}\left\{-180y_t^4 + 192y_t^2g_3^2 + 54y_t^2g_2^2 + \frac{102}{5}y_t^2g_1^2\right. \\
&+72\lambda_h^2 + 72y_t^2\lambda_h - 54g_2^2\lambda_h - \frac{54}{5}g_1^2\lambda_h \\
&\left.-15g_2^4 + 9g_1^4 + \frac{27}{5}g_2^2g_1^2\right\}.
\end{aligned}$$

$$\lambda_m^{(1)} = \frac{b_m}{128\pi^2\tilde{\rho}}$$

$$\begin{aligned}
d_m =& \frac{1}{16\pi^2}\{192y_t^2\beta_3 \\
&+\left(192g_3^2 - 360y_t^2 + 54g_2^2 + \frac{102}{5}g_1^2 + 72\lambda_h\right)\beta_t \\
&+\left(72y_t^2 - 54g_2^2 - \frac{54}{5}g_1^2 + 144\lambda_h\right)\beta_h \\
&+\left(54y_t^2 - 30g_2^2 + \frac{27}{5}g_1^2 - 54\lambda_h\right)\beta_2 \\
&+\left(\frac{102}{5}y_t^2 + \frac{27}{5}g_2^2 + 18g_1^2 - \frac{54}{5}\lambda_h\right)\beta_1\Big\}.
\end{aligned}$$

$$\hat{\delta}_m = \lambda(\tilde{\rho}_0) - \lambda_m^{(\mathrm{cr})}(\tilde{\rho}_0).$$

$$\delta_m(\tilde{\rho}) = \lambda_m(\tilde{\rho}) - \lambda_m^{(\mathrm{cr})}(\tilde{\rho}),$$

$$\delta_m(\tilde{\rho}_0) = \hat{\delta}_m, \delta_m(\tilde{\rho} \to \infty) = \bar{\lambda}_m.$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m = -\frac{1}{2}\beta_m$$
$$\tilde{\rho}\partial_{\tilde{\rho}}\delta_m = -\frac{1}{2}\frac{\partial\beta_m}{\partial\lambda_m}\delta_m$$

$$\beta_m = \tilde{\beta}_m(\tilde{\rho}) - A_m(\tilde{\rho})\lambda_m,$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\delta_m = -\frac{1}{2}A_m(\tilde{\rho})\delta_m,$$

$$\delta_m(\tilde{\rho}) = C_m(\tilde{\rho})\hat{\delta},$$

$$C_m(\tilde{\rho}) = \exp\left\{-\frac{1}{2}\int_{\tilde{\rho}_0}^{\tilde{\rho}} \mathrm{d}\rho' \frac{A_m(\rho')}{\rho'}\right\}.$$

$$\bar{A}_m = \int_{\tilde{\rho}_0}^{\tilde{\rho}} \frac{\mathrm{d}\rho'}{\rho'} \frac{A_m(\rho')}{\ln\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)},$$

$$C_m(\tilde{\rho}) = \left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{-\frac{\bar{A}_m}{2}}.$$

$$A_m(\tilde{\rho}) = \frac{3\lambda_h(\tilde{\rho})}{8\pi^2}(1+\partial_{\tilde{h}}u)^{-3}$$

$$h_0 = D|\hat{\delta}|^{2\beta}$$

$$h_0 = -\frac{\lambda_m}{\lambda_h} = \frac{C_m(\tilde{\rho})}{\lambda_h}|\hat{\delta}| = \frac{|\hat{\delta}|}{\lambda_h}\tilde{\rho}^{\bar{A}_m/2}h_0^{\bar{A}_m/2},$$

$$\beta = \frac{1}{2}\left(1-\frac{\bar{A}_m}{2}\right)^{-1}$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\mu = \mu - \frac{1}{2}\tilde{\beta}_\mu(\mu,\alpha),$$
$$\tilde{\rho}\partial_{\tilde{\rho}}\alpha = -\frac{1}{2}\beta_\alpha(\mu,\alpha).$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\begin{pmatrix}\delta\mu\\ \delta\alpha\end{pmatrix} = -\frac{1}{2}\begin{pmatrix}S_{\mu\mu} & S_{\mu\alpha}\\ S_{\alpha\mu} & S_{\alpha\alpha}\end{pmatrix}\begin{pmatrix}\delta\mu\\ \delta\alpha\end{pmatrix}$$

$$S_{\mu\mu} = -2 + \frac{\partial\tilde{\beta}_\mu}{\partial\mu}, \quad S_{\mu\alpha} = \frac{\partial\tilde{\beta}_\mu}{\partial\alpha}$$
$$S_{\alpha\mu} = \frac{\partial\beta_\alpha}{\partial\mu}, \qquad S_{\alpha\alpha} = \frac{\partial\beta_\alpha}{\partial\alpha}$$

$$\tilde{\beta}_\mu = -2\beta_\nu = -n\alpha/(32\pi^2)$$

$$\frac{\partial\beta_\alpha}{\partial\mu} = -\frac{9\lambda_h^2}{2\pi^2(1+\mu)^4}, \frac{\partial\beta_\alpha}{\partial\alpha} = \frac{3\lambda_h}{2\pi^2(1+\mu)^3}$$

$$\frac{\partial\tilde{\beta}_\mu}{\partial\alpha}=-\frac{3}{32\pi^2(1+\mu)^2},\frac{\partial\tilde{\beta}_\mu}{\partial\mu}=-2+\frac{3\lambda_h}{8\pi^2(1+\mu)^3}$$

$$Z=3\lambda_h/(16\pi^2(1+\mu)^3)$$

$$\lambda_+=\frac{3\lambda_h}{2\pi^2(1+\mu)^3}-18\left(\frac{3\lambda_h}{16\pi^2(1+\mu)^3}\right)^2$$
$$\lambda_-=-2+\frac{3\lambda_h}{8\pi^2(1+\mu)^3}+18\left(\frac{3\lambda_h}{16\pi^2(1+\mu)^3}\right)^2$$

$$\delta\mu=\delta\mu(\tilde{\rho}_0)\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{-\frac{\lambda_-}{2}}$$

$$\delta\mu(\tilde{\rho})=\delta\mu(\tilde{\rho}_0)\left[(1-\tilde{c}Z)\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{-\frac{\lambda_-}{2}}+\tilde{c}Z\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{-\frac{\lambda_+}{2}}\right].$$

$$\delta_m(\tilde{\rho})=\partial_{\tilde{\rho}}\delta\mu(\tilde{\rho})=-\frac{\delta\mu(\tilde{\rho}_0)}{2\tilde{\rho}_0}\left[\lambda_-(1-\tilde{c}Z)\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{-\frac{1}{2}\hat{A}_m}+\lambda_+\tilde{c}Z\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{-\left(1+\frac{\lambda_+}{2}\right)}\right]$$

$$\hat{A}_m=A_m\left(1+\frac{27\lambda_h}{8\pi^2(1+\mu)^3}\right)$$

$$A_m^{(\mathrm{gr})}=\frac{5}{12\pi^2 f}\left(1-\frac{2u}{f}\right)^{-2},f=\frac{F}{k^2}=\frac{M_p^2(k)}{k^2}$$

$$A_m^{(\mathrm{gr})}\lambda_m=4\partial_{\tilde{\rho}}\partial_{\tilde{h}}c_U^{(\mathrm{gr})}(\tilde{\rho},\tilde{h}).$$

$$c_U^{(\mathrm{gr})}=\frac{1}{128\pi^2}\left\{\left(1+\frac{1}{3}\left(1-\frac{\partial\ln f}{\partial\ln\tilde{\rho}}\right)\right)\left(\frac{5}{1-v}+\frac{1}{1-v/4}\right)-4\right\}$$

$$Fq^2-2U+Fk^2r_k(q^2/k^2)$$

$$\left(1+\tilde{m}_{\mathrm{gr}}^2\right)^{-1}=(1-v)^{-1}\sim\frac{1}{3}\left(1-\frac{\partial\ln f}{\partial\ln\tilde{\rho}}\middle|\ h\right)$$
$$v=\frac{u}{w}=\frac{2u}{f}$$

$$\delta_g=\frac{1}{4}\frac{\partial\ln f}{\partial\ln\tilde{\rho}}\Big|_{\hbar}$$

$$\partial_{\tilde{h}}c_U^{(\mathrm{gr})}(\tilde{\rho},\tilde{h})=\frac{(1-\delta_g)\partial_{\tilde{h}}v}{96\pi^2}\left(\frac{5}{(1-v)^2}+\frac{1}{4(1-v/4)^2}\right).$$

$$\partial_{\tilde{h}}c_U^{(\mathrm{gr})}(\tilde{\rho},\tilde{h})=\frac{1-\delta_g}{48\pi^2(1-v)^2f}(\partial_{\tilde{h}}u-u\partial_{\tilde{h}}\ln f).$$

$$\partial_{\tilde{\rho}}\partial_{\tilde{h}}c_U^{(\mathrm{gr})}(\tilde{\rho},\tilde{h}) = \frac{\lambda_m(1-\delta_g)}{48\pi^2(1-v)^2 f} - \frac{\partial_{\tilde{\rho}}u\partial_{\tilde{h}}u(1-\delta_g)}{12\pi^2(1-v)^3 f^2} + \Delta_{\mathrm{gr}},$$

$$u(\tilde{\rho},\tilde{h}) = \lambda_m\tilde{\rho}\tilde{h} + \bar{u}(\tilde{\rho},\tilde{h})$$

$$\begin{aligned}\partial_{\tilde{\rho}}u\partial_{\tilde{h}}u &= \lambda_m^2\tilde{\rho}\tilde{h} + \lambda_m(\tilde{\rho}\partial_{\tilde{\rho}} + \tilde{h}\partial_{\tilde{h}})\bar{u} + \partial_{\tilde{\rho}}\bar{u}\partial_{\tilde{h}}\bar{u} \\ &= -\lambda_m^2\tilde{\rho}\tilde{h} + 2\lambda_m(u - c_U) + \partial_{\tilde{\rho}}\bar{u}\partial_{\tilde{h}}\bar{u}\end{aligned}$$

$$f = 2w_0 + 2\xi\tilde{\rho} + \xi_H\tilde{h},$$

$$\delta_g = \frac{2\xi\tilde{\rho} + \xi_H\tilde{h}}{4(2w_0 + 2\xi\tilde{\rho} + \xi_H\tilde{h})}.$$

$$\partial_{\tilde{h}}\ln f = \frac{\xi_H}{f}, \partial_{\tilde{\rho}}\ln f = \frac{2\xi}{f}, \partial_{\tilde{h}}\partial_{\tilde{\rho}}\ln f = -2\frac{\xi\xi_H}{f^2},$$

$$\partial_{\tilde{h}}\delta_g = \frac{\xi_H}{4f}(1 - 4\delta_g)$$

$\partial_{\tilde{h}}\ln f = \xi_H/(2w_0), \partial_{\tilde{\rho}}\ln f = \xi/w_0$ contribute to $\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m = -\beta_m/2$

$$\lambda_m(\tilde{\rho}) = \lambda_m^{(r)}(\tilde{\rho}) + d_m\tilde{\rho}^{-\frac{A_m}{2}},$$

$$\lambda_m(\tilde{\rho}) - \lambda_m^{(r)}(\tilde{\rho}) \star (\tilde{\rho}/\tilde{\rho}_0)^{-A_m/2}$$

$$\mu_h = \partial_{\tilde{h}}u(\tilde{\rho},\tilde{h} = 0)$$

$$\lambda_m(\tilde{\rho}) = \partial_{\tilde{\rho}}\mu_h(\tilde{\rho})$$

$$k\partial_k\mu_h = 2\tilde{\rho}\partial_{\tilde{\rho}}\mu_h - 2\mu_h + 4\partial_{\tilde{h}}c_U$$

$$k\partial_k\mu_h = 2\tilde{\rho}\partial_{\tilde{\rho}}\mu_h + (A_\mu - 2)\mu_h$$

$$\partial_{\tilde{h}}c_U^{(\mathrm{gr})} = \frac{1}{4}A_\mu^{(\mathrm{gr})}\mu_h + \cdots, A_\mu^{(\mathrm{gr})} = A_m^{(\mathrm{gr})}.$$

$$\mu_h(k) = \mu_h^{(\mathrm{cr})}(k) + \delta_\mu(k), \delta_\mu(k) \sim k^{A_m - 2}.$$

$$\mu_h(\tilde{\rho}) = \mu_h^{(\mathrm{cr})}(\tilde{\rho}) + \delta_\mu(\tilde{\rho}), \delta_\mu(\tilde{\rho}) = d_\mu\tilde{\rho}^{1-\frac{A_m}{2}}$$

$$\lambda_m^{(\mathrm{cr})}(\tilde{\rho}) = \partial_{\tilde{\rho}}\mu_h^{(\mathrm{cr})}(\tilde{\rho}), \delta\lambda_m(\tilde{\rho}) = \partial_{\tilde{\rho}}\delta_\mu(\tilde{\rho}),$$

$$d_m = d_\mu\left(1 - \frac{A_m}{2}\right)$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\mu_h = -\frac{1}{2}\bar{\beta}_\mu(\tilde{\rho}) + \left(1 - \frac{A_\mu}{2}\right)\mu_h$$

$$\mu_h(0) = \frac{\bar{\beta}_\mu(0)}{2 - A_\mu}$$

$$\frac{\varphi_0}{M_p} = \sqrt{\frac{h_0}{2\xi}} = \left(-\frac{\bar{\lambda}_m}{2\xi\bar{\lambda}_h}\right)^{\frac{1}{2}}$$

$$\mu_h(\tilde{\rho} = 0) = \partial_{\tilde{h}} u|_{\tilde{\rho}=\tilde{h}=0}, \lambda_m(\tilde{\rho} = 0) = \partial_{\tilde{\rho}}\partial_{\tilde{h}} u\big|_{\tilde{\rho}=\tilde{h}=0} \text{ or } \lambda_h(\tilde{\rho} = 0) = \partial_{\tilde{h}}^2 u\big|_{\tilde{\rho}=\tilde{h}=0}$$

$$\partial_{\tilde{\rho}}\partial_{\tilde{h}}^2 u\big|_{\tilde{\rho}=\tilde{h}=0} \text{ or } \partial_{\tilde{\rho}}^2\partial_{\tilde{h}} u\big|_{\tilde{\rho}=\tilde{h}=0}$$

$$\mu_h(\tilde{\rho}) = \partial_{\tilde{h}} u(\tilde{\rho}, h = 0)$$
$$\tilde{\rho}\partial_{\tilde{\rho}}\mu_h = -\frac{1}{2}\beta_\mu$$

$$\partial_{\tilde{\rho}}\mu_h = \lambda_m$$

$$\tilde{\rho}\partial_{\tilde{\rho}}^2\mu_h + \lambda_m = -\frac{1}{2}\partial_{\tilde{\rho}}\beta_\mu$$

$$\beta_\mu(\tilde{\rho}) = \beta_\mu^{(1)}\tilde{\rho} + \cdots = -2\lambda_m\tilde{\rho} + \cdots$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\sigma = -\beta_\sigma/2$$

$$\tilde{h}\partial_{\tilde{h}}\mu_h = -\frac{1}{2}\beta_\mu = \tilde{h}\lambda_h,$$

$$\tilde{h}\partial_{\tilde{h}}^2\mu_h + \lambda_h = -\frac{1}{2}\partial_{\tilde{h}}\beta_\mu,$$

$$\partial_{\tilde{h}}\beta_\mu\big|_{\tilde{h}=0} = -2\lambda_h(\tilde{h} = 0)$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m = -\frac{1}{2}\beta_m = -\frac{1}{128\pi^2\tilde{\rho}}\sum_i \hat{n}_i\beta_i - \frac{1}{2}A_m\lambda_m$$

$$\lambda_m^* = -\frac{\sum_i \hat{n}_i c_i}{64\pi^2 A_m}$$

$$\beta_i = c_i\tilde{\rho} + \frac{d_i}{2}\tilde{\rho}^2 + \cdots$$

$$\partial_{\tilde{\rho}}\beta_m = \frac{1}{128\pi^2}\sum_i \hat{n}_i d_i + A_m\partial_{\tilde{\rho}}\lambda_m + \partial_{\tilde{\rho}}A_m\lambda_m$$

$$\sigma(\tilde{\rho}) = \sigma^{(r)}(\tilde{\rho}) + \delta_\sigma(\tilde{\rho}),$$
$$\tilde{\rho}\partial_{\tilde{\rho}}\delta_\sigma = -\frac{1}{2}A_\sigma\delta_\sigma,$$
$$A_\sigma = \partial_\sigma\beta_\sigma.$$

$$\sigma(\tilde{\rho}) = \sigma^{(r)}(\tilde{\rho}) + c_\sigma\tilde{\rho}^{-\frac{A_\sigma}{2}}$$

$$\frac{\partial \lambda_m}{\partial x} = -\frac{1}{2}\left(\tilde{\beta}_m(x) + A(x)\lambda_m\right).$$

$$\lambda_m(x) = \lambda_c(x) + \tilde{C}(x)\lambda_m(x_0),$$

$$\tilde{C}(x) = \exp\left\{-\frac{1}{2}\int_{x_0}^{x} dx' A(x')\right\}.$$

$$\lambda_c(x) = -\frac{\tilde{C}(x)}{2}\int_{x_0}^{x} dx' \frac{\tilde{\beta}_m(x')}{\tilde{C}(x')}.$$

$$\frac{1}{2}\tilde{\beta}_m(x) = \frac{b_m(x)}{128\pi^2}e^{-x},$$

$$\mu_h(\tilde{\rho}) = \mu_0 + \lambda_0\tilde{\rho} + \cdots.$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\mu_h = \left(1 - \frac{A}{2}\right)\mu_h + \beta_v,$$

$$\mu_0 = -\beta_0\left(1 - \frac{A_0}{2}\right)^{-1}$$

$$\lambda_0 = \frac{2\beta_1 - A_1\mu_0}{A_0} = \frac{2}{A_0}\left(\beta_1 + \frac{A_1\beta_0}{2 - A_0}\right).$$

$$\lambda_m(\tilde{\rho}) = \partial_{\tilde{\rho}}\mu_h(\tilde{\rho})$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m(\tilde{\rho}) = \partial_{\tilde{\rho}}\beta_v(\tilde{\rho}) - \frac{1}{2}\partial_{\tilde{\rho}}A(\tilde{\rho})\mu_h(\tilde{\rho}) - \frac{1}{2}A(\tilde{\rho})\lambda_m(\tilde{\rho}),$$

$$\tilde{\beta}_m(\tilde{\rho}) = -2\partial_{\tilde{\rho}}\beta_v(\tilde{\rho}) + \partial_{\tilde{\rho}}A_m(\tilde{\rho})\mu_h(\tilde{\rho}).$$

$$\tilde{\beta}_0 = \tilde{\beta}_m(\tilde{\rho} = 0) = -2\beta_1 + A_1\mu_0.$$

$$\ln\,\tilde{C} = -\frac{1}{2}\int_{\tilde{\rho}_0}^{\tilde{\rho}} \frac{d\rho'}{\rho'}(A_0 + A_1\rho') = -\frac{1}{2}\left(A_0 \ln\frac{\tilde{\rho}}{\tilde{\rho}_0} + A_1(\tilde{\rho} - \tilde{\rho}_0)\right)$$

$$\tilde{C} = \left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{-\frac{A_0}{2}} \exp\left(-\frac{A_1}{2}(\tilde{\rho} - \tilde{\rho}_0)\right)$$

$$\begin{aligned}\lambda_c(\tilde{\rho}) =& -\frac{1}{2}\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{-\frac{A_0}{2}} \exp\left\{-\frac{A_1}{2}(\tilde{\rho} - \tilde{\rho}_0)\right\} \\ &\times \int_{\tilde{\rho}_0}^{\tilde{\rho}} \frac{d\rho'}{\rho'}\left(\tilde{\beta}_0 + \tilde{\beta}_1\rho'\right)\left(\frac{\rho'}{\tilde{\rho}_0}\right)^{\frac{A_0}{2}} \exp\left\{\frac{A_1}{2}(\rho' - \tilde{\rho}_0)\right\} \\ =& \frac{\tilde{\beta}_0}{A_0}\left[\left(\frac{\tilde{\rho}_0}{\tilde{\rho}}\right)^{\frac{A_0}{2}} - 1\right]\left(1 - \frac{A_1}{2}\tilde{\rho}\right) \\ &+ \frac{2\tilde{\beta}_1 + A_1\tilde{\beta}_0}{4 + 2A_0}\left(\tilde{\rho}_0^{\,1+\frac{A_0}{2}}\tilde{\rho}^{-\frac{A_0}{2}} - \tilde{\rho}\right).\end{aligned}$$

$$\lambda_m(\tilde{\rho}) = \left(\frac{\tilde{\beta}_0}{A_0} + \frac{2\tilde{\beta}_1 + A_1\tilde{\beta}_0}{4 + 2A_0}\tilde{\rho}_0 + \lambda_m(\tilde{\rho}_0)\left(1 + \frac{A_1}{2}\tilde{\rho}_0\right)\right)\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{-\frac{A_0}{2}} - \frac{A_1}{2}\left(\frac{\tilde{\beta}_0}{A_0} + \lambda_m(\tilde{\rho}_0)\right)\tilde{\rho}\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{-\frac{A_0}{2}}$$

$$-\frac{\tilde{\beta}_0}{A_0} + \left(\frac{A_1\tilde{\beta}_0}{2A_0} - \frac{2\tilde{\beta}_1 + A_1\tilde{\beta}_0}{4 + 2A_0}\right)\tilde{\rho}$$

$$\lambda_m(\tilde{\rho}_0) = -\left(\frac{\tilde{\beta}_0}{A_0} + \frac{2\tilde{\beta}_1 + A_1\tilde{\beta}_0}{4 + 2A_0}\tilde{\rho}_0\right)\left(1 + \frac{A_1}{2}\tilde{\rho}_0\right)^{-1}$$

$$\lambda_m(\tilde{\rho}) = -\frac{\tilde{\beta}_0}{A_0} + \left(\frac{A_1\tilde{\beta}_0}{2A_0} - \frac{2\tilde{\beta}_1 + A_1\tilde{\beta}_0}{4 + 2A_0}\right)\tilde{\rho}$$

$$\lambda_m(\tilde{\rho}) = -\lim_{\tilde{\rho}_0 \to 0}\left[\exp\left\{-\frac{1}{2}\int_{\tilde{\rho}_0}^{\tilde{\rho}}\frac{d\rho'}{\rho'}A(\rho')\right\}\frac{\tilde{\beta}_0}{A_0} + \frac{1}{2}\int_{\tilde{\rho}_0}^{\tilde{\rho}}\frac{d\rho'}{\rho'}\tilde{\beta}_m(\rho')\exp\left\{\frac{1}{2}\int_{\tilde{\rho}}^{\rho'}\frac{d\rho''}{\rho''}A(\rho'')\right\}\right]$$

$$\lambda_m(\tilde{\rho}) = -\frac{1}{2}\lim_{\tilde{\rho}_0 \to 0}\int_{\tilde{\rho}_0}^{\tilde{\rho}}\frac{d\rho'}{\rho'}\left(\tilde{\beta}_0 + \tilde{\beta}_1\rho'\right)\left(\frac{\rho'}{\tilde{\rho}}\right)^{\frac{A_0}{2}} = -\left(\frac{\tilde{\beta}_0}{A_0} + \frac{\tilde{\beta}_1}{2 + A_0}\tilde{\rho}\right)$$

$$\lambda_m(\tilde{\rho}) = \exp\left\{\frac{1}{2}\int_{\tilde{\rho}}^{\infty}\frac{d\rho'}{\rho'}A(\rho')\right\}\left(\bar{\lambda}_m + \Delta_m(\tilde{\rho})\right)$$

$$\Delta_m(\tilde{\rho}) = \frac{1}{2}\int_{\tilde{\rho}}^{\infty}\frac{d\rho'}{\rho'}\tilde{\beta}_m(\rho')\exp\left\{-\frac{1}{2}\int_{\rho'}^{\infty}\frac{d\rho''}{\rho''}A(\rho'')\right\}$$

$$D(\tilde{\rho}) = \exp\left\{\frac{1}{2}\int_{\tilde{\rho}}^{\infty}\frac{d\rho'}{\rho'}A(\rho')\right\}$$

$$\bar{\lambda}_m = -\Delta_m(0) = -\frac{1}{2}\int_{0}^{\infty}\frac{d\rho'}{\rho'}\tilde{\beta}_m(\rho')\exp\left\{-\frac{1}{2}\int_{\rho'}^{\infty}\frac{d\rho''}{\rho''}A(\rho'')\right\}$$

$$\lambda_m^{(R)}(\tilde{\rho}) = D^{-1}(\tilde{\rho})\lambda_m(\tilde{\rho})$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m^{(R)} = -\frac{1}{2}\tilde{\beta}_m^{(R)} = -\frac{1}{2}D^{-1}(\tilde{\rho})\tilde{\beta}_m(\tilde{\rho}).$$

$$\bar{\lambda}_m = -\frac{1}{2}\int_{-\infty}^{\infty}dx\tilde{\beta}_m^{(R)}(x) = -\frac{1}{2}\int_{0}^{\infty}\frac{d\rho'}{\rho'}\tilde{\beta}_m^{(R)}(\rho')$$

$$\tilde{\beta}_m^{(R)}(\rho') = \left(\tilde{\beta}_0 + \tilde{\beta}_1\rho'\right)\left(\frac{\rho'}{\bar{\rho}}\right)^{\frac{A_0}{2}}D^{-1}(\bar{\rho}),$$

$$\sigma_m(\tilde{\rho}) = \tilde{\rho}\lambda_m(\tilde{\rho}, h = 0)$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\sigma_m = \sigma_m + \tilde{\rho}^2\partial_{\tilde{\rho}}\lambda_m = \sigma_m - \frac{1}{2}\tilde{\rho}\beta_m = \sigma_m - \frac{1}{128\pi^2}\left\{\sum_{\tilde{\rho}\partial_{\tilde{\rho}}\mu_h}{}' \hat{n}_i\beta_i - 12\tilde{\rho}\partial_{\tilde{\rho}}\lambda_h(1+\partial_{\tilde{h}}u)^{-2}\right.$$

$$\left. = +24\lambda_h\sigma_m(1+\partial_{\tilde{h}}u)^{-3}\right\}$$

$$\tilde{m}_i^2 = \alpha_i h\tilde{\rho}, \tilde{\rho}\partial_{\tilde{\rho}}\alpha_i = -\frac{1}{2}\beta_i\lambda_m\sigma_m.$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\sigma_m = \left(1-\frac{A_m}{2}\right)\sigma_m - \frac{1}{128\pi^2}\left\{\sum_{\partial_{\tilde{\rho}}\mu_h}{}' \hat{n}_i\beta_i + 6\beta_h(1+\mu_h)^{-2}\right\}$$

$$\mu_h(\tilde{\rho}) = \partial_{\tilde{h}}u(\tilde{\rho}, \tilde{h}=0), \lambda_h(\tilde{\rho}) = \partial_{\tilde{h}}^2 u(\tilde{\rho}, \tilde{h}=0)$$
$$\beta_h = -2\tilde{\rho}\partial_{\tilde{\rho}}\lambda_h(\tilde{\rho})$$

$$A_m = \frac{3\lambda_h}{8\pi^2(1+\mu_h)^3} + \eta_H$$

$$\eta_H = \frac{3}{8\pi^2}\left(y_t^2 - \frac{3(5g_2^2+g_1^2)}{20(1+\mu_h)}\right).$$

$$\beta_t = -2\tilde{\rho}\partial_{\tilde{\rho}}y_t^2$$
$$= \frac{1}{16\pi^2}\left\{\frac{3y_t^4}{1+\mu_h} - \left(16g_3^2 + \frac{4}{5}g_1^2\right)y_t^2\right\} + \eta_H y_t^2$$
$$\beta_3 = -2\tilde{\rho}\partial_{\tilde{\rho}}g_3^2 = -\frac{7g_3^4}{8\pi^2}$$
$$\beta_2 = -2\tilde{\rho}\partial_{\tilde{\rho}}g_2^2 = -\frac{1}{12\pi^2}\left(5 - \frac{1}{4(1+\mu_h)^3}\right)g_2^4$$
$$\beta_1 = -2\tilde{\rho}\partial_{\tilde{\rho}}g_1^2 = \frac{1}{2\pi^2}\left(1 + \frac{1}{40(1+\mu_h)^3}\right)g_1^4$$

$$\beta_h = \frac{3}{4\pi^2}\left\{\frac{\lambda_h^2}{(1+\mu_h)^3} - y_t^4 + \frac{3}{16}g_2^4 + \frac{3}{40}g_2^2g_1^2 + \frac{9}{400}g_1^4\right\} + 2\eta_H\lambda_h$$

$$x_F = \ln\left(\frac{M_p^2}{2m_Z^2}\right) = 74.9.$$

$$\mu(x_F) = \sigma_m(x_F) + \delta\mu(x_F).$$

$$\mu_{\mathrm{cr}} = -\frac{3}{32\pi^2}\left(\lambda_h + \frac{3}{4}g_2^2 + \frac{3}{20}g_1^2 - 2y_t^2\right).$$

$$\lambda_m^{(\mathrm{cr})} = \frac{1}{2}\tilde{\beta}_m\left(1-\frac{A_m}{2}\right)^{-1},$$

$$\tilde{\beta}_m = \frac{1}{64\pi^2\tilde{\rho}}\left(\sum_{e^{-i\omega t}}{}'' \hat{n}_i\beta_i + \frac{6\beta_h}{(1+\mu_h)^2}\right).$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\lambda_m = -\frac{1}{2}\tilde{\beta}_m - \frac{1}{2}A_m\lambda_m.$$

$$\beta_i^{(\mathrm{gr})} = -B_g g_i^2$$
$$B_g = \frac{5}{288\pi^2 w}\left(\frac{4}{1-v} - \frac{3}{(1-v)^2}\right)$$
$$w = w_0 + \tilde{\rho}, v = \frac{u_0}{w}$$

$$\Delta\beta_t = \frac{y_t^2}{16\pi^2}\left\{\frac{a_y y_t^2 - a_g g_3^2}{1 + c_F\tilde{\rho}} + \frac{2b_t}{3w(1-v)^2}\right\}.$$

$$\tilde{H}_t = \bar{t}_R\begin{pmatrix} t_L \\ b_L \end{pmatrix}$$

$$\mathcal{L}_{Y,t} = y_t H^\dagger \tilde{H}_t + \text{ h.c.}$$

$$\mathcal{L}_{4,t} = -\frac{1}{2}\bar{\lambda}_t \tilde{H}_t^\dagger \tilde{H}_t$$

$$k\partial_k\bar{\lambda}_t = -\frac{D}{16\pi^2 k^2}$$
$$D = \frac{80}{3}g_3^4 - 5y_t^4$$

$$k\partial_k\Gamma_k|_{H_k} = k\partial_k\Gamma_k|_H + \Delta_k$$
$$\Delta_k = -\int_x\left(\frac{\partial\Gamma_k}{\partial H_k}\partial_t H_k + \partial_t H_k^\dagger\frac{\partial\Gamma_k}{\partial H_k^\dagger}\right)$$

$$\partial_t H_k = k\partial_k H_k|_{H,\tilde{H}_t} = k\partial_k\alpha_k\tilde{H}_t$$

$$\frac{\partial\Gamma_k}{\partial H_k} = \frac{\partial U}{\partial\left(H_k^\dagger H_k\right)}H_k^\dagger + y_t\tilde{H}_t^\dagger$$
$$\frac{\partial\Gamma_k}{\partial H_k^\dagger} = \frac{\partial U}{\partial\left(H_k^\dagger H_k\right)}H_k + y_t\tilde{H}_t$$

$$\Delta_k = -\int_x k\partial_k\alpha_k\left\{\frac{\partial U}{\partial\left(H_k^\dagger H_k\right)}H_k^\dagger\tilde{H}_t + y_t\tilde{H}_t^\dagger\tilde{H}_t + h.c.\right\}$$

$$-y_t k\partial_k\alpha_k = \frac{1}{4}k\partial_k\bar{\lambda}_t = -\frac{D}{64\pi^2 k^2},$$

$$\tilde{\Delta}_k = -\int_x\frac{D}{64\pi^2 k^2 y_t}\frac{\partial U}{\partial(H^\dagger H)}\left(H^\dagger\tilde{H}_t + \text{ h.c. }\right)$$

$$\Delta(k\partial_k y_t) = -\frac{D}{64\pi^2 k^2 y_t}\frac{\partial U}{\partial(H^\dagger H)}.$$

$$k\partial_k y_t^2 = \beta_{t,H} - \frac{D\partial_{\tilde{h}}u}{32\pi^2} = \beta_{t,H} - \frac{D\mu_h}{32\pi^2},$$

$$\tilde{\rho}\partial_{\tilde{\rho}} y_t^2 = -\frac{1}{32\pi^2}\left\{9y_t^4 - \left(16g_3^2 + \frac{9}{2}g_2^2 + \frac{17}{10}g_1^2\right)y_t^2 - \frac{1}{2}D\mu_h\right\} - \frac{1}{2}\eta_{tc}y_t^2$$

$$\tilde{\epsilon}_t = \frac{\mu_h}{y_t^2}$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\tilde{\epsilon}_t = \frac{1}{y_t^2}\left(\tilde{\rho}\partial_{\tilde{\rho}}\mu_h - \tilde{\epsilon}_t\tilde{\rho}\partial_{\tilde{\rho}}y_t^2\right)$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\mu_h = \mu_h + \beta_v - \frac{1}{2}(\eta_H + \eta_{ct})\mu_h$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\tilde{\epsilon}_t = -\frac{D}{64\pi^2}\tilde{\epsilon}_t^2 + E\tilde{\epsilon}_t - \frac{F}{64\pi^2}$$

$$E = 1 + \frac{1}{32\pi^2}\left[3y_t^2 - \left(16g_3^2 + \frac{4}{5}g_1^2\right)\right],$$
$$F = 12 - \left(\frac{9}{2}\frac{g_2^2}{y_t^2} + \frac{9}{10}\frac{g_1^2}{y_t^2} + \frac{2\lambda_h}{y_t^2}\right).$$

$$\tilde{\epsilon}_{t,\mathrm{UV}} = \frac{F}{64\pi^2 E}$$

$$\mu_h = \frac{3}{16\pi^2}\left[y_t^2 - \left(\frac{3}{8}g_2^2 + \frac{3}{40}g_1^2 + \frac{1}{6}\lambda_h\right)\right],$$

$$\epsilon_{t*} = \frac{32\pi^2}{D}\left(E \pm \sqrt{E^2 - DF/(32\pi^2)^2}\right)$$

$$\tilde{\epsilon}_{t,\mathrm{IR}} = \frac{64\pi^2 E}{D}$$

$$\mu_H \approx \frac{64\pi^2 y_t^2}{D}$$

$$\tilde{h}_0 \approx \frac{64\pi^2 y_t^2}{\lambda_h |D|}$$

$$\tilde{\epsilon}_t = -\lambda_h \tilde{h}_0 / y_t^2$$

$$\tilde{m}_t^2 = -y_t^4 \tilde{\epsilon}_t / \lambda_h$$

$$(1 + \tilde{m}_t^2)^{-1} = (1 - y_t^4 \tilde{\epsilon}_t / \lambda_h)^{-1}$$

$$\tilde{\epsilon}_t = \sqrt{\frac{F}{D}} = \frac{F}{32\pi^2 E}, \mu_h = \sqrt{\frac{F}{D}}y_t^2 = \frac{F y_t^2}{32\pi^2 E}$$

$$E = 1 - \frac{g_3^2}{2\pi^2}\left(1 - \frac{9}{16R_t}\right)R_g,$$

$$R_g = 1 + \frac{9g_2^2}{32g_3^2} + \frac{17g_1^2}{160g_3^2} \approx 1.1$$

$$F = 12 - \frac{6m_W^2}{m_t^2} - \frac{3m_Z^2}{m_t^2} - \frac{m_H^2}{m_t^2}$$

$$D_3 = \frac{80}{3} g_3^4 t_{D3}(\tilde{m}_t^2)$$

$$t_{D3}(\tilde{m}_t^2) = (1+\tilde{m}_t^2)^{-2}\left(\frac{1}{3} + \frac{2}{3(1+\tilde{m}_t^2)}\right)$$

$$\alpha_s^2 = \frac{12\pi^2 E^2}{5F} = \frac{12\pi^2}{5F}\left[1 - \frac{2\alpha_s}{\pi}\left(1 - \frac{9}{16R_t}\right)R_g\right]^2.$$

$$\left(1 - \frac{48B_t^2}{5F}\right)\alpha_s^2 + \frac{48\pi B_t}{5F}\alpha_s - \frac{12\pi^2}{5F} = 0$$

$$S_t = \frac{\lambda_h}{y_t^2}, k\partial_k S_t = \beta_{S,p} + \delta\beta_S$$

$$\delta\beta_S = \frac{D\mu_h\lambda_h}{32\pi^2 y_t^4} = \frac{(80g_3^4 - 15y_t^4)\mu_h\lambda_h}{96\pi^2 y_t^4}$$

$$S_t = S_{t,p} + \delta_S, k\partial_k S_{t,p} = \beta_{S,p}$$

$$k\partial_k\delta_S = \delta\beta_S$$

$$t_u^{(i)}(\tilde{\rho},h) = 2\tilde{\rho}^2\int_{\tilde{\rho}}^{\infty}\frac{\mathrm{d}\rho'}{\rho'^3\left(1+\tilde{m}_i^2(\rho',h)\right)} = 1 - 2\tilde{\rho}^2\int_{\tilde{\rho}}^{\infty}\frac{\mathrm{d}\rho'\tilde{m}_i^2(\rho',h)}{\rho'^3\left(1+\tilde{m}_i^2(\rho',h)\right)}$$

$$\partial_h t_u^{(i)}(\tilde{\rho},h) = -2\tilde{\rho}^2\int_{\tilde{\rho}}^{\infty}\mathrm{d}\rho'\frac{\partial_h\tilde{m}_i^2}{\rho'^3\left(1+\tilde{m}_i^2\right)^2}$$

$$\tilde{m}_i^2(\tilde{\rho},h) = \alpha_i(\tilde{\rho},h)h\tilde{\rho}$$

$$\partial_h t_u^{(i)} = -2\tilde{\rho}^2\int_{\tilde{\rho}}^{\infty}\mathrm{d}\rho'\frac{\alpha_i + h\partial_h\alpha_i}{\rho'^2(1+\alpha_i h\rho')^2}$$

$$\tilde{m}_i^2(\tilde{\rho},h) \sim \tilde{\rho}, \text{ with } \alpha_i(\tilde{\rho}\to\infty,h) \to \bar{\alpha}_i(h)$$

$$t_u^{(i)}(\tilde{\rho},h) = 2\tilde{\rho}^2\int_{\tilde{\rho}}^{\infty}\frac{\mathrm{d}\rho'}{\rho'^3(1+\bar{\alpha}_i(h)h\rho')} = \frac{2}{3\bar{\alpha}_i(h)h\tilde{\rho}} = \frac{2}{3\tilde{m}_i^2(\tilde{\rho},h)}$$

$$\partial_h t_u^{(i)}(\tilde{\rho},h) = -\frac{2(\bar{\alpha}_i + h\partial_h\bar{\alpha}_i)}{3\bar{\alpha}_i^2 h^2\tilde{\rho}} = -\frac{2}{3\tilde{m}_i^2}\left(\frac{1}{h} + \partial_h\ln\,\bar{\alpha}_i\right)$$

$$L(h)\tilde{\rho}^2 = u_{\mathrm{UV}}(\tilde{\rho}) - \frac{1}{128\pi^2}\sum_i \hat{n}_i t_u^{(i)}\left(\tilde{m}_i^2\right)$$

$$\int_{\tilde{\rho}}^{\infty} \mathrm{d}\rho' \frac{\alpha_i + h\partial_h \alpha_i}{\rho'^3 (1+\alpha_i h\rho')^2} = \Delta_1 + \Delta_2$$

$$\Delta_1 = \int_{\tilde{\rho}}^{\tilde{\rho}_{\mathrm{tr}}} \frac{\mathrm{d}\rho' \alpha_i(\rho')}{\rho'^2}$$

$$\Delta_2 = \int_{\tilde{\rho}_{\mathrm{tr}}}^{\infty} \mathrm{d}\rho' \frac{\alpha_i + h\partial_h \alpha_i}{\rho'^2 (1+\alpha_i h\rho')^2}$$

$$\frac{\alpha(\rho')}{\rho^2}\, \mathrm{d}\rho' = -\mathrm{d}\left(\frac{f(\rho')}{\rho'}\right),$$

$$f\left(1 - \frac{\partial \ln f}{\partial \ln \rho'}\right) = \alpha(\rho').$$

$$f_0(\rho') = \alpha(\rho')$$

$$f_1(\rho') = \frac{\alpha(\rho')}{1 - \frac{\partial \ln \alpha(\rho')}{\partial \ln \rho'}}$$

$$f_n(\rho') = \frac{\alpha(\rho')}{1 - \frac{\partial \ln f_{n-1}(\rho')}{\partial \ln \rho'}}$$

$$\frac{\partial \alpha_i}{\partial \ln \tilde{\rho}} = -\frac{1}{2}\beta_i$$

$$\Delta_1 = \frac{\alpha_i(\tilde{\rho})}{\tilde{\rho}(1 - (\beta_i/\alpha_i)(\tilde{\rho}))} - \frac{\alpha_i(\tilde{\rho}_{\mathrm{tr}})h}{c_{\mathrm{tr}}\left(1 - (\beta_i/\alpha_i)(\tilde{\rho}_{\mathrm{tr}})\right)}.$$

$$\Delta_2 = \Delta_3 + \Delta_4 + \Delta_5$$

$$\Delta_3 = \int_{\tilde{\rho}_{\mathrm{tr}}}^{\infty} \frac{\mathrm{d}\rho' \alpha_i(\rho')}{\rho'^2}, \Delta_4 = \int_{\tilde{\rho}_{\mathrm{tr}}}^{\infty} \mathrm{d}\rho' \frac{h\partial_h \alpha_i}{\rho'^2 (1+\alpha_i h\rho')^2}$$

$$\Delta_5 = -h \int_{\tilde{\rho}_{\mathrm{tr}}}^{\infty} \mathrm{d}\rho' \frac{2\alpha_i^2 + \alpha_i^3 h\rho'}{\rho'(1+\alpha_i h\rho')^2}$$

$$\Delta_5 = -h\alpha_i^2 \int_{\tilde{\rho}_{\mathrm{tr}}}^{\infty} \mathrm{d}\rho' \left(\frac{2}{\rho'(1+\alpha_i h\rho')} - \frac{\alpha_i h}{(1+\alpha_i h\rho')^2}\right) = h\alpha_i^2 \left(2\ln\left(\frac{\alpha_i c_{\mathrm{tr}}}{1+\alpha_i c_{\mathrm{tr}}}\right) + \frac{1}{1+\alpha_i c_{\mathrm{tr}}}\right)$$

$$\partial_h t_u^{(i)}(\tilde{\rho}, h) \approx -\frac{2\alpha_i(\tilde{\rho})\tilde{\rho}}{1 - (\beta_i/\alpha_i)(\tilde{\rho})} - 2\tilde{\rho}^2 \Delta_4,$$

$$\hat{h}\partial_{\hat{h}} t_{u|h}^{(i)} = 2\left(t_u^{(i)} - \frac{1}{1+\tilde{m}_i^2}\right),$$

$$\tilde{h}\partial_{\tilde{h}}\left(\frac{t_u^{(i)}}{\tilde{h}^2}\right) = -\frac{2}{\left(1+\tilde{m}_i^2\right)\tilde{h}^2},$$

$$t_u^{(i)}(\tilde{h}, h) = 2\tilde{h}^2 \int_{\tilde{h}}^{\infty} \frac{\mathrm{d}h'}{h'^3\left(1+\tilde{m}_i^2(h', h)\right)} = 1 - 2\tilde{h}^2 \int_{\tilde{h}}^{\infty} \frac{\mathrm{d}h' \tilde{m}_i^2(h', h)}{h'^3\left(1+\tilde{m}_i^2(h', h)\right)}$$

$$\tilde{m}_i^2(\tilde{h},h)=\alpha_i(\tilde{h},h)\tilde{h},$$

$$t_u^{(i)}(\tilde{h},h)=1-2\tilde{h}^2\int_{\tilde{h}}^{\infty}\frac{\mathrm{d}h'\alpha_i(h',h)}{h'^2(1+\alpha_i(h',h)h')}$$

$$\partial_h t_u^{(i)}(\tilde{h},h)=-\frac{2\tilde{h}^2}{h}\int_{\tilde{h}}^{\infty}\frac{\mathrm{d}h'h\partial_h\alpha_i(h',h)}{h'^2(1+\alpha_i h')^2}.$$

$$h\partial_{h|\tilde{h}}=h\partial_{h|\tilde{\rho}}-\tilde{\rho}\partial_{\tilde{\rho}|h},$$

$$\partial_h t_{u|\tilde{h}}^{(i)}=2\tilde{\rho}^2\int_{\tilde{\rho}}^{\infty}\frac{\mathrm{d}\rho'\left(\rho'\partial_{\rho'}-h\partial_h\right)\alpha_i(\rho',h)}{\rho'^2(1+\alpha_i h\rho')^2}$$

$$k\partial_k y_t^2=\frac{b_t y_t^4}{1+y_t^2\tilde{h}},b_t=\frac{9}{16\pi^2}$$

$$\tilde{h}\partial_{\tilde{h}}y_{t|h}^2=-\frac{1}{2}b_t y_t^4\left(1+y_t^2\tilde{h}\right)^{-1}$$

$$\left(1+y_t^2\tilde{h}\right)^{-1}\text{ by }\left(1+\bar{y}_t^2\tilde{h}\right)^{-1}$$

$$y_t^{-2}(\tilde{h})=y_t^{-2}\left(\tilde{h}_{\mathrm{in}}\right)+\frac{b_t}{2}\ln\left(\frac{\tilde{h}\left(1+\bar{y}_t^2\tilde{h}_{\mathrm{in}}\right)}{\tilde{h}_{\mathrm{in}}\left(1+\bar{y}_t^2\tilde{h}\right)}\right)$$

$$y_t^{-2}(\tilde{h})=y_t^{-2}\left(\tilde{h}_{\mathrm{in}}\right)+\frac{b_t}{2}\ln\frac{1+\bar{y}_t^2\tilde{h}_{\mathrm{in}}}{\bar{y}_t^2\tilde{h}_{\mathrm{in}}}-\frac{b_t}{2\bar{y}_t^2\tilde{h}}.$$

$$h\partial_h\alpha_t=h\partial_h y_t^2=-y_t^4h\partial_h y_t^{-2}(\tilde{h},h)=-y_t^4h\partial_h y_t^{-2}\left(\tilde{h}_{\mathrm{in}}\right)$$

$$y_t^2(\tilde{\rho}_0,h)=\bar{y}_{t,0}^2$$

$$y_t^{-2}(\tilde{h})=\bar{y}_{t,0}^{-2}-\frac{b_t}{2}\ln\left(\frac{h\tilde{\rho}_0}{1+\bar{y}_t^2h\tilde{\rho}_0}\right)+\frac{b_t}{2}\ln\left(\frac{\tilde{h}}{1+\bar{y}_t^2\tilde{h}}\right),$$

$$h\partial_h y_t^2(\tilde{h},h)=\frac{b_t}{2}y_t^4(\tilde{h},h)$$

$$h\partial_h t_u^{(t)}(\tilde{h},h)=-b_t\tilde{h}^2\int_{\tilde{h}}^{\infty}\frac{\mathrm{d}h'y_t^4(h',h)}{h'^2(1+y_t^2(h',h)h')^2}$$

$$h\partial_h t_u^{(t)}(\tilde{h},h)=-b_t\tilde{h}y_t^4(\tilde{h})J(\tilde{h})$$

$$J(\tilde{h})=\tilde{h}\int_{\tilde{h}}^{\infty}\frac{\mathrm{d}h'}{h'^2(1+\bar{y}_t^2h')^2}=1+\bar{y}_t^2\tilde{h}\left(2\ln\left(\frac{\bar{y}_t^2\tilde{h}}{1+\bar{y}_t^2\tilde{h}}\right)+\frac{1}{1+\bar{y}_t^2\tilde{h}}\right)$$

$$t_u^{(t)}(\tilde{h},h)=t_u^{(t)}\left(\tilde{h},h_0\right)-b_t y_t^4(\tilde{h})\tilde{h}J(\tilde{h})\ln\left(\frac{h}{h_0}\right)$$

$$J(\tilde{h} \gg \bar{y}_t^2) = \frac{1}{3\bar{y}_t^4 \tilde{h}^2},$$

$$\Delta U = \frac{1}{4}L(h)\chi^4 + \frac{3b_t}{32\pi^2}y_t^4\left(\frac{H^\dagger H}{k^2}\right)J\left(\frac{H^\dagger H}{k^2}\right)\ln\left(\frac{2H^\dagger H}{\chi^2}\right)H^\dagger H k^2$$

$$\Delta U = \frac{b_t k^6}{32\pi^2 H^\dagger H}\ln\left(\frac{2H^\dagger H}{\chi^2}\right)$$

$$\nu = \frac{A_\nu \tilde{\rho}}{64\pi^2}$$

$$A_\nu = 12y_t^2 s_u^{(t)}(y_t^2 h\tilde{\rho}) - 3g_2^2 s_u^{(W)}\left(\frac{1}{2}g_2^2 h\tilde{\rho}\right) - \left(\frac{3g_2^2}{2} + \frac{9g_1^2}{10}\right)s_u^{(Z)}\left(\left(\frac{g_2^2}{2} + \frac{3g_1^2}{10}\right)h\tilde{\rho}\right)$$

$$- 3\lambda_h s_u^{(H)}\left(2\lambda_h h\tilde{\rho} + \frac{\tilde{v}}{\tilde{\rho}}\right)$$

$$\Delta_+ = \frac{1}{\bar{\varphi}_0}(12\bar{m}_t^2 - 6\bar{m}_W^2 - 3\bar{m}_Z^2 - 3\bar{m}_H^2) > 0$$

$$\Delta_- = \bar{\varphi}_0^2\left(\frac{4}{\bar{m}_t^2} - \frac{2}{\bar{m}_W^2} - \frac{1}{\bar{m}_Z^2} - \frac{1}{\bar{m}_H^2}\right) < 0$$

$$h_{\max} = \frac{c_{\max}}{y_t^2 \tilde{\rho}}$$

$$\lim_{h\to\infty} \nu(\tilde{\rho}, h) = 0$$

$$12\tilde{m}_b^2 + 12\tilde{m}_c^2 + 4\tilde{m}_\tau^2 = \frac{2}{\tilde{m}_W^2} + \frac{1}{\tilde{m}_Z^2} + \frac{2}{\tilde{m}_H^2} - \frac{4}{\tilde{m}_t^2}$$

$$\hat{h}_0(\tilde{\rho}) = \sqrt{-\Delta_-/\Delta_b}\,\tilde{\rho}^{-1}$$

$$\Delta_b = \frac{12\bar{m}_b^2 + 12\bar{m}_c^2 + 4\bar{m}_\tau^2}{\bar{\varphi}_0^2}$$

$$\nu(\tilde{\rho}, h_0) + \left(\bar{\lambda}_m + \bar{\lambda}_h h_0\right)\tilde{\rho}^2 = 0$$

$$h_0 = -\frac{\nu(\tilde{\rho}, h_0)}{\bar{\lambda}_h \tilde{\rho}^2}$$

$$\tilde{h}_0 = h_0\tilde{\rho} = -\frac{A_\nu(\tilde{h}_0)}{64\pi^2\bar{\lambda}_h}$$

$$\tilde{m}_{t|0}^2 = y_t^2\tilde{h}_0 = -\frac{A_\nu}{64\pi^2}\frac{\bar{m}_t^2}{\bar{m}_H^2}$$

$$u= u_0 + \tilde{v}h + \frac{1}{2}\partial_h \tilde{v} h^2 + \cdots$$
$$= u_0 + \left(h\partial_h + \frac{1}{2}h^2\partial_h^2\right) L_{|h=0}\tilde{\rho}^2 - \tilde{\beta}_v \tilde{h} - \frac{1}{2}h\partial_h \tilde{\beta}_v \tilde{h}$$

$$h\partial_h \tilde{\beta}_{v|\tilde{\rho}} = h\partial_h \tilde{\beta}_{v|\tilde{h}} + \tilde{h}\partial_{\tilde{h}} \tilde{\beta}_{v|h}$$

$$U= u_0(\tilde{\rho})k^4 + \frac{1}{2}\bar{\lambda}_m \chi^2 H^\dagger H + \frac{1}{2}\bar{\lambda}_h \left(H^\dagger H\right)^2$$
$$-\tilde{\beta}_v(\tilde{h})\left(1 + \frac{1}{2}\frac{\partial \ln\, \tilde{\beta}_v}{\partial \ln\, \tilde{h}}\right) k^2 H^\dagger H - \partial_h \tilde{\beta}_v(\tilde{h}) \frac{k^2}{\chi^2}\left(H^\dagger H\right)^2$$

$$\tilde{\beta}_v(\tilde{h}) = \tilde{\beta}_v(\tilde{h}, h=0), \partial_h \tilde{\beta}_v(\tilde{h}) = \partial_h \tilde{\beta}_v(\tilde{h}, h=0), \text{ with } \tilde{h} = H^\dagger H/k^2$$

$$u = u_0(\tilde{\rho}, \bar{h}_0) + \tilde{v}(\tilde{\rho}, \bar{h}_0)(h - \bar{h}_0) + \frac{1}{2}\partial_h \tilde{v}(\tilde{\rho}, \bar{h}_0)(h - \bar{h}_0)^2 + \cdots$$

$$= u_0 + \left[(h - \bar{h}_0)\partial_h + \frac{1}{2}(h - \bar{h}_0)^2 \partial_{h^2}\right] L_{|h=\bar{h}_0}\tilde{\rho}^2$$

$$-\left[\tilde{\beta}_v(\tilde{h}, \bar{h}_0) + \frac{1}{2}(h\partial_h + \tilde{h}\partial_{\tilde{h}})\tilde{\beta}_v(\tilde{h}, h_0)\left(1 - \frac{\bar{h}_0}{h}\right)\right] \times (\tilde{h} - \bar{h}_0\tilde{\rho})$$

$$\partial_h L(\bar{h}_0) = 0, \partial_h^2 L(\bar{h}_0) = \bar{\lambda}_{h,0}$$

$$U = u_0 k^4 + \frac{1}{2}\left[\bar{\lambda}_{h,0} - \frac{1}{\tilde{\rho}}\partial_h \tilde{\beta}_v(\tilde{\rho}, \bar{h}_0)\right]\left(H^\dagger H - \frac{1}{2}\bar{h}_0\chi^2\right)^2 - \tilde{\beta}_v(\tilde{\rho}, \bar{h}_0)k^2\left(H^\dagger H - \frac{1}{2}\bar{h}_0\chi^2\right)$$

$$\frac{\partial U}{\partial (H^\dagger H)} = \left[\bar{\lambda}_h - \frac{1}{\tilde{\rho}}\partial_h \tilde{\beta}_v(\tilde{\rho}, \bar{h}_0)\right]\left(H^\dagger H - \frac{1}{2}\bar{h}_0\chi^2\right) - \tilde{\beta}_v(\tilde{\rho}, \bar{h}_0)k^2$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\partial_{\tilde{h}}\left(\tilde{\rho}, \tilde{h}_0(\tilde{\rho})\right) + \partial_{\tilde{h}}^2 u \tilde{\rho}\partial_{\tilde{\rho}}\tilde{h}_0(\tilde{\rho}) = 0$$

$$\lambda_m(\tilde{\rho}) = \partial_{\tilde{\rho}}\partial_{\tilde{h}} u\big|_{\tilde{h}_0}, \lambda_h = \partial_{\tilde{h}}^2 u\big|_{\tilde{h}_0}.$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\tilde{h}_0 = -\frac{\tilde{\rho}\lambda_m}{\lambda_h} = -\frac{\sigma_m}{\lambda_h} = \zeta$$
$$\sigma_m = \tilde{\rho}\lambda_m, \zeta = -\frac{\sigma_m}{\lambda_h}$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\zeta = -\frac{1}{\lambda_h}\tilde{\rho}\partial_{\tilde{\rho}}\sigma_m + \frac{\sigma_m}{\lambda_h^2}\tilde{\rho}\partial_{\tilde{\rho}}\lambda_h$$
$$= \zeta + \frac{\tilde{\rho}\beta_m}{2\lambda_h} + \frac{\zeta\beta_h}{2\lambda_h}$$

$$\tilde{\rho}\beta_m = \tilde{\rho}\tilde{\beta}_m + A_m\sigma_m$$

$$\tilde{\rho}\tilde{\beta}_m = -\frac{1}{64\pi^2}\left\{\frac{12\beta_t}{\left(1+y_t^2\tilde{h}_0\right)^2} - 3\beta_h\left(1+\frac{1}{\left(1+2\lambda_h\tilde{h}_0\right)^2}\right) - \frac{3\beta_2}{\left(1+g_2^2\tilde{h}_0/2\right)^2}\right.$$

$$\left. - \frac{3\beta_2/2 + 9\beta_1/10}{\left((1+g_2^2/2+3g_1^2/10)\tilde{h}_0\right)^2}\right\}$$

$$\tilde{\rho}\partial_{\tilde{\rho}}\zeta = \gamma_\zeta(\tilde{h}_0)\zeta + \eta_\zeta(\tilde{h}_0)$$

$$\gamma_\zeta = 1 - \frac{A_m}{2} + \frac{\beta_h}{2\lambda_h}, \eta_\zeta = \frac{\tilde{\rho}\tilde{\beta}_m}{2\lambda_h}$$

$$\partial_x^2\tilde{h}_0 = \partial_x\zeta = \gamma_\zeta(\tilde{h}_0)\partial_x\tilde{h}_0 + \eta_\zeta(\tilde{h}_0).$$

$$\delta\tilde{h} = \tilde{h}_0 - \hat{h}_0$$

$$\eta_\zeta(\hat{h}_0) = \eta'_\zeta\delta\tilde{h}$$

$$\partial_x^2\delta\tilde{h} = \gamma_\zeta\partial_x\delta\tilde{h} - \eta'_\zeta\delta\tilde{h}$$

$$\delta\tilde{h} = \delta\tilde{h}(x_0)\exp\{\omega(x-x_0)\}$$

$$\omega^2 - \gamma_\zeta\omega - \eta'_\zeta = 0$$

$$\omega_\pm = \frac{\gamma_\zeta}{2}\left(1 \pm \sqrt{1+\frac{4\eta'}{\gamma_\zeta^2}}\right),$$

$$\omega_+ = \gamma_\zeta + \frac{\eta'}{\gamma_\zeta}, \omega_- = -\frac{\eta'}{\gamma_\zeta}.$$

$$\delta\tilde{h}(\tilde{\rho}) = \delta\tilde{h}(\tilde{\rho}_0)\left(\frac{\tilde{\rho}}{\tilde{\rho}_0}\right)^{\omega_+},$$

$$\Phi = \begin{pmatrix} \phi^+ \\ \frac{1}{\sqrt{2}}(v_\phi + \phi + i\chi) \end{pmatrix}, \Delta = \begin{pmatrix} \frac{1}{\sqrt{2}}\Delta^+ & \Delta^{++} \\ \Delta^0 & -\frac{1}{\sqrt{2}}\Delta^+ \end{pmatrix} \text{ with } \Delta^0 = \frac{1}{\sqrt{2}}(v_\Delta + \delta + i\eta),$$

$$\begin{aligned} V(\Phi,\Delta) =& m^2\Phi^\dagger\Phi + M^2\mathrm{Tr}\left(\Delta^\dagger\Delta\right) + \left[\mu\Phi^T i\sigma_2\Delta^\dagger\Phi + \text{ h.c. }\right] \\ &+\lambda_1\left(\Phi^\dagger\Phi\right)^2 + \lambda_2\left[\mathrm{Tr}\left(\Delta^\dagger\Delta\right)\right]^2 + \lambda_3\mathrm{Tr}\left[\left(\Delta^\dagger\Delta\right)^2\right] \\ &+\lambda_4\left(\Phi^\dagger\Phi\right)\mathrm{Tr}\left(\Delta^\dagger\Delta\right) + \lambda_5\Phi^\dagger\Delta\Delta^\dagger\Phi \end{aligned}$$

$$m^2 = -\lambda_1 v_\phi^2 - \frac{1}{2}(\lambda_4+\lambda_5)v_\Delta^2 + \sqrt{2}\mu v_\Delta$$

$$M^2 = -(\lambda_2+\lambda_3)v_\Delta^2 - \frac{1}{2}(\lambda_4+\lambda_5)v_\phi^2 + \frac{\mu v_\phi^2}{\sqrt{2}v_\Delta}$$

$$\mathcal{M}_{\pm}^{2} = \left(\frac{\mu v_{\phi}^{2}}{\sqrt{2}v_{\Delta}} - \frac{\lambda_5}{4}v_{\phi}^{2}\right)\begin{pmatrix} 2v_{\Delta}^{2}/v_{\phi}^{2} & -\sqrt{2}v_{\Delta}/v_{\phi} \\ -\sqrt{2}v_{\Delta}/v_{\phi} & 1 \end{pmatrix}$$

$$\mathcal{M}_{\text{odd}}^{2} = \frac{\mu v_{\phi}^{2}}{\sqrt{2}v_{\Delta}}\begin{pmatrix} 4v_{\Delta}^{2}/v_{\phi}^{2} & -2v_{\Delta}/v_{\phi} \\ -2v_{\Delta}/v_{\phi} & 1 \end{pmatrix}, \mathcal{M}_{\text{even}}^{2} = \begin{pmatrix} \mathcal{M}_{11}^{2} & \mathcal{M}_{12}^{2} \\ \mathcal{M}_{21}^{2} & \mathcal{M}_{22}^{2} \end{pmatrix},$$

$$\mathcal{M}_{11}^{2} = 2\lambda_1 v_{\phi}^{2}, \mathcal{M}_{22}^{2} = \frac{\mu v_{\phi}^{2}}{\sqrt{2}v_{\Delta}} + 2(\lambda_2 + \lambda_3)v_{\Delta}^{2}$$
$$\mathcal{M}_{12}^{2} = \mathcal{M}_{21}^{2} = (\lambda_4 + \lambda_5)v_{\phi}v_{\Delta} - \sqrt{2}\mu v_{\phi}$$

$$\begin{pmatrix} \phi^{\pm} \\ \Delta^{\pm} \end{pmatrix} = R(\beta)\begin{pmatrix} G^{\pm} \\ H^{\pm} \end{pmatrix}, \begin{pmatrix} \chi \\ \eta \end{pmatrix} = R(\beta')\begin{pmatrix} G^{0} \\ A \end{pmatrix}, \begin{pmatrix} \phi \\ \delta \end{pmatrix} = R(\alpha)\begin{pmatrix} h \\ H \end{pmatrix},$$

$$R(\theta) = \begin{pmatrix} c_\theta & -s_\theta \\ s_\theta & c_\theta \end{pmatrix}$$

$$\tan\,\beta = \frac{\sqrt{2}v_{\Delta}}{v_{\phi}}, \tan\,\beta' = \frac{2v_{\Delta}}{v_{\phi}}, \tan\,2\alpha = \frac{2\mathcal{M}_{12}^{2}}{\mathcal{M}_{11}^{2} - \mathcal{M}_{22}^{2}}.$$

$$m_{H^{\pm\pm}}^{2} = \frac{\mu v_{\phi}^{2}}{\sqrt{2}v_{\Delta}} - \lambda_3 v_{\Delta}^{2} - \frac{\lambda_5}{2}v_{\phi}^{2}$$
$$m_{H^{\pm}}^{2} = \left(\frac{\mu v_{\phi}^{2}}{\sqrt{2}v_{\Delta}} - \frac{\lambda_5}{4}v_{\phi}^{2}\right)\left(1 + \frac{2v_{\Delta}^{2}}{v_{\phi}^{2}}\right), m_{A}^{2} = \frac{\mu v_{\phi}^{2}}{\sqrt{2}v_{\Delta}}\left(1 + \frac{4v_{\Delta}^{2}}{v_{\phi}^{2}}\right)$$
$$m_{h}^{2} = \mathcal{M}_{11}^{2}c_{\alpha}^{2} + \mathcal{M}_{22}^{2}s_{\alpha}^{2} + 2\mathcal{M}_{12}^{2}s_{\alpha}c_{\alpha}, m_{H}^{2} = \mathcal{M}_{11}^{2}s_{\alpha}^{2} + \mathcal{M}_{22}^{2}c_{\alpha}^{2} - 2\mathcal{M}_{12}^{2}s_{\alpha}c_{\alpha}$$

$$\sin(2\alpha) = \frac{2v_{\phi}v_{\Delta}}{m_{H}^{2} - m_{h}^{2}}\left[\frac{4m_{H^{\pm}}^{2}}{v_{\phi}^{2} + 2v_{\Delta}^{2}} - \frac{2m_{A}^{2}}{v_{\phi}^{2} + 4v_{\Delta}^{2}} - \lambda_4\right]$$

$$v, m_{h}^{2}, m_{H}^{2}, m_{A}^{2}, m_{H^{\pm}}^{2}, m_{H^{\pm\pm}}^{2}, \lambda_4, \beta'$$

$$\left(m_{H}^{2} - m_{h}^{2}\right)^{2} \geq 4(\mathcal{M}_{12}^{2})^{2}$$

$$\mathcal{L}_{\text{Kin}} = \left(D_\mu \Phi\right)^{\dagger}(D^\mu \Phi) + \text{Tr}\left[\left(D_\mu \Delta\right)^{\dagger} D^\mu \Delta\right]$$

$$D_\mu \Phi = \left(\partial_\mu - ig\tau^a W_\mu^a - ig' Y B_\mu\right)\Phi, D_\mu \Delta = \partial_\mu \Delta - ig\left[\tau^a W_\mu^a, \Delta\right] - ig' Y B_\mu \Delta,$$

$$m_{W}^{2} = \frac{g^2}{4}\left(v_{\phi}^{2} + 2v_{\Delta}^{2}\right), m_{Z}^{2} = \frac{g_Z^2}{4}\left(v_{\phi}^{2} + 4v_{\Delta}^{2}\right)$$

$$\rho_0 = \frac{m_{W}^{2}}{m_{Z}^{2}c_{W}^{2}} = \frac{v_{\phi}^{2} + 2v_{\Delta}^{2}}{v_{\phi}^{2} + 4v_{\Delta}^{2}}$$

$$\mathcal{L}_{\Delta}^{Y} = h_{ij}\left(\overline{L_L^c}\right)_i i\tau_2 \Delta (L_L)_j + \text{ h.c.}$$

$$\mathcal{L}_{\Phi}^{Y} \supset -\sum_{f=u,d,e} \frac{m_f}{v}\left[\bar{f}f(\zeta_h h + \zeta_H H) - 2iI_f \bar{f}\gamma_5 f(\zeta_{G^0} G^0 + \zeta_A A)\right]$$
$$+\left[\left(\frac{\sqrt{2}m_u}{v} V_{ud}\bar{u}P_L d - \frac{\sqrt{2}m_d}{v} V_{ud}\bar{u}P_R d\right)\left(\zeta_{G^\pm} G^+ + \zeta_{H^\pm} H^+\right) + \text{h.c.}\right]$$
$$-\left[\frac{\sqrt{2}m_e}{v}\bar{\nu}P_R e\left(\zeta_{G^\pm} G^+ + \zeta_{H^\pm} H^+\right) + \text{h.c.}\right]$$

$$\zeta_h = c_\alpha/c_\beta, \qquad \zeta_{G^0} = c_{\beta'}/c_\beta, \qquad \zeta_{G^\pm} = 1$$
$$\zeta_H = -s_\alpha/c_\beta, \qquad \zeta_A = -s_{\beta'}/c_\beta, \qquad \zeta_{H^\pm} = -s_\beta/c_\beta$$

$$\lambda_1 > 0, \lambda_2 + \text{MIN}\left[\lambda_3, \frac{\lambda_3}{2}\right] > 0$$

$$\lambda_4 + \text{MIN}[0, \lambda_5] + 2\text{MIN}\left[\sqrt{\lambda_1(\lambda_2 + \lambda_3)}, \sqrt{\lambda_1\left(\lambda_2 + \frac{\lambda_3}{2}\right)}\right] > 0$$

$$y_1 = 2\lambda_1, y_2 = 2(\lambda_2 + \lambda_3), y_3 = 2\lambda_2$$
$$y_{4,\pm} = \lambda_1 + \lambda_2 + 2\lambda_3 \pm \sqrt{(\lambda_1 - \lambda_2 - 2\lambda_3)^2 + \lambda_5^2}$$
$$y_{5,\pm} = 3\lambda_1 + 4\lambda_2 + 3\lambda_3 \pm \sqrt{(3\lambda_1 - 4\lambda_2 - 3\lambda_3)^2 + \frac{3}{2}(2\lambda_4 + \lambda_5)^2}$$
$$y_6 = \lambda_4, y_7 = \lambda_4 + \lambda_5, y_8 = \frac{1}{2}(2\lambda_4 + 3\lambda_5)$$
$$y_9 = \frac{1}{2}(2\lambda_4 - \lambda_5), y_{10} = 2\lambda_2 - \lambda_3$$

$$v_\Delta = \frac{\mu v_\phi^2}{\sqrt{2}\left[M^2 + (\lambda_2 + \lambda_3)v_\Delta^2 + (\lambda_4 + \lambda_5)v_\phi^2/2\right]}$$

$$G_F = \frac{1}{\sqrt{2}v^2(1 - \Delta r)} \text{ with } v^2 = \frac{m_W^2}{\pi\alpha_{\text{em}}}\left(1 - \frac{m_W^2}{\rho_0 m_Z^2}\right)$$

$$\Delta r = \frac{\text{Re}\widehat{\Pi}_{WW}(0)}{m_W^2} + \delta_{VB}$$

$$\delta_{VB} = \frac{\alpha_{\text{em}}}{4\pi s_W^2}\left[6 + \frac{10(1 - s_W^2) - 3(R/c_W^2)(1 - 2s_W^2)}{2(1 - R)}\ln R\right]_{\Gamma_{Zf\bar{f}}^{V/A,\,\text{loop}}} \text{ with } R = \frac{m_W^2}{m_Z^2}$$

$$\bar{m}_W^2 = \rho_0 m_Z^2 \bar{c}_W^2 \text{ with } \bar{c}_W^2 = 1 - \bar{s}_W^2,$$

$$\bar{s}_W^2 = \frac{1}{2}\left(1 - \sqrt{1 - \frac{4A_0}{\rho_0 m_Z^2}}\right) \text{ with } A_0 = \frac{\pi\alpha_{\text{em}}}{\sqrt{2}G_F}.$$

$$s^2_{W,G_F} = \frac{1}{2}\left[1 - \sqrt{1 - \frac{4A_0}{\rho_0 m_Z^2(1-\Delta r)}}\right], m^2_{W,G_F} = \rho_0 m_Z^2 c_W^2$$

$$\bar{v}_\phi^2 + 2\bar{v}_\Delta^2 = \bar{v}^2 \text{ with } \bar{v}^2 = \frac{1}{\sqrt{2}G_F},$$

$$\tan\beta' = 2\bar{v}_\Delta/\bar{v}_\phi$$

$$v^2_{\Delta,G_F} = \frac{s^2_{\beta'}}{2\left(1+c^2_{\beta'}\right)}\frac{1}{\sqrt{2}G_F(1-\Delta r)}$$

$$g_V^f = 2\left(\frac{\rho_0(1-\Delta r)}{1+\hat{\Pi}'_Z(m_Z^2)}\right)^{1/2} \times\left[\frac{I_f}{2} - Q_f s_W^2\left(1 + \frac{c_W}{s_W}\frac{\hat{\Pi}_{Z\gamma}(m_Z^2)}{m_Z^2 + \hat{\Pi}_{\gamma\gamma}(m_Z^2)}\right) + \Gamma^{V,\,\text{loop}}_{Zf\bar{f}}(0,0,m_Z^2)\right]$$

$$g_A^f = 2\left(\frac{\rho_0(1-\Delta r)}{1+\hat{\Pi}'_Z(m_Z^2)}\right)^{1/2}\left[\frac{I_f}{2} + \Gamma^{A,\,\text{loop}}_{Zf\bar{f}}(0,0,m_Z^2)\right]$$

$$\hat{\Pi}'_Z(p^2) = \text{Re}\frac{\mathrm{d}\hat{\Pi}_Z(p^2)}{\mathrm{d}p^2}\bigg|_{p^2=m_Z^2} \text{ with } \hat{\Pi}_Z(p^2) = \hat{\Pi}_{ZZ}(p^2) - \frac{\left[\hat{\Pi}_{Z\gamma}(p^2)\right]^2}{p^2 + \hat{\Pi}_{\gamma\gamma}(p^2)}.$$

$$s_f^2 = \frac{1}{4|Q_f|}\left(1 - \frac{\text{Re}\left(g_V^f\right)}{\text{Re}\left(g_A^f\right)}\right)$$

$$\Gamma(Z\to\ell\bar{\ell}) = \frac{\sqrt{2}m_Z^3 G_F}{12\pi}\left[\left(g_V^\ell\right)^2 + \left(g_A^\ell\right)^2\right]\left(1 + Q_\ell^2\frac{3\alpha_{\text{em}}}{4\pi}\right).$$

$$\rho = \rho_0 + \Delta\rho$$

$$\Delta\rho = \text{Re}\left[\frac{\Pi_{ZZ}(0)}{m_Z^2} - \frac{\Pi_{WW}(0)}{m_W^2} + \frac{2s_W}{c_W}\frac{\Pi_{Z\gamma}(0)}{m_Z^2} - \frac{s_{2\beta'}}{1+c^2_{\beta'}}\delta\beta'\right],$$

$$\tilde{\Pi}_{ij}(p^2) = \Pi_{ij}^{1\text{PI}}(p^2) + \Pi_{ij}^{\text{Tad}} + \Pi_{ij}^{\text{PT}}(p^2).$$

$$\delta\alpha = -\frac{1}{2(m_H^2 - m_h^2)}\left(\tilde{\Pi}_{Hh}(m_H^2) + \tilde{\Pi}_{Hh}\left(m_h^2\right)\right)$$


$$\delta\beta' = -\frac{1}{2m_A^2}\left(\tilde{\Pi}_{AG^0}(m_A^2) + \tilde{\Pi}_{AG^0}(0)\right)$$


$$\delta\beta = \left(1 + s_\beta^2\right)\delta\beta'/\sqrt{2}$$


$$\mathcal{O}_{\xi_V} = \mathcal{O}|_{\xi_V=1} + \mathcal{O}|_{\text{Gauge Decay (G.D.)}}$$

$$C_0(p^2;X,Y) \equiv \frac{1}{m_X^2 - m_Y^2}[B_0(p^2;X,X) - B_0(p^2;Y,Y)],$$

$$C_0(p^2;X,Y,Z) \equiv \frac{1}{m_X^2 - m_Y^2}[B_0(p^2;X,Z) - B_0(p^2;Y,Z)],$$







$$\mathcal{M} = \overline{\mathcal{M}}\left(\frac{m_f}{v}\right)^2 (\bar{f}f) \times (\bar{f}f)$$

$$\overline{\mathcal{M}}^{hh}_{Z-15}\Big|_{\text{G.D.}} + \overline{\mathcal{M}}^{hh}_{W-11}\Big|_{\text{G.D.}} = \frac{g_Z^2}{128\pi^2}\frac{\zeta_h^2}{p^2 - m_h^2}(1-\xi_Z)[-2c_{hhZZ}B_0(0;Z,G^0)$$
$$+c^2_{G^0hZ}\left(p^2+m_h^2\right)C_0(p^2;Z,G^0) + 2c^2_{AhZ}\left(p^2-2m_A^2+m_h^2\right)C_0(p^2;Z,G^0,A)]$$
$$+\frac{g^2}{64\pi^2}\frac{\zeta_h^2}{p^2-m_h^2}(1-\xi_W)[-2c_{hhWW^2}B_0(0;W,G^\pm)$$
$$+2c^2_{H^\pm hW^\mp}\left(p^2+m_h^2-2m^2_{H^\pm}\right)C_0(p^2;W,G^\pm,H^\pm) + c^2_{G^\pm hW^\mp}\left(p^2+m_h^2\right)C_0(p^2;W,G^\pm)]$$
$$\overline{\mathcal{M}}^{HH}_{Z-15}\Big|_{\text{G.D.}} + \overline{\mathcal{M}}^{HH}_{W-11}\Big|_{\text{G.D.}} = \frac{g_Z^2}{128\pi^2}\frac{\zeta_h^2}{p^2 - m_H^2}(1-\xi_Z)[-2c_{HHZZ}B_0(0;Z,G^0)$$
$$+c^2_{G^0HZ}(p^2+m_H^2)C_0(p^2;Z,G^0) - 2c^2_{AHZ}(2m_A^2-m_H^2-p^2)C_0(p^2;Z,G^0,A)]$$
$$+\frac{g^2}{64\pi^2}\frac{\zeta_H^2}{p^2-m_H^2}(1-\xi_W)[-2c_{HHWW}B_0(0;W,G^\pm)$$
$$+2c^2_{H^\pm HW^\mp}\left(p^2+m_H^2-2m^2_{H^\pm}\right)C_0(p^2;W,G^\pm,H^\pm) + c^2_{G^\pm HW^\mp}(p^2+m_H^2)C_0(p^2;W,G^\pm)]$$
$$\overline{\mathcal{M}}^{hH}_{Z-15}\Big|_{\text{G.D.}} + \overline{\mathcal{M}}^{hH}_{W-11}\Big|_{\text{G.D.}} = \frac{g_Z^2}{256\pi^2}\frac{\zeta_H\zeta_h}{(p^2-m_h^2)(p^2-m_H^2)}(1-\xi_Z)$$
$$\times\left[-c_{hHZZ}\left(2p^2-m_h^2-m_H^2\right)B_0(0;Z,G^0) + 2c_{G^0hZ^c}c_{G^0HZ}\left(p^4-m_h^2m_H^2\right)C_0(p^2;Z,G^0)\right.$$
$$\left.+4c_{AhZ^c}c_{AHZ}\left\{p^4+m_A^2\left(m_h^2+m_H^2-2p^2\right)-m_h^2m_H^2\right\}C_0(p^2;Z,G^0,A)\right]$$
$$+\frac{g^2}{128\pi^2}\frac{\zeta_h\zeta_H}{(p^2-m_h^2)(p^2-m_H^2)}(1-\xi_W)\left[-c_{hHWW}\left(2p^2-m_h^2-m_H^2\right)B_0(0;W,G^\pm)\right.$$
$$+4c_{H^\pm hW^\mp}c_{H^\mp HW^\pm}\left\{p^2\left(p^2-2m^2_{H^\pm}\right)-m_h^2m_H^2+m_h^2m^2_{H^\pm}+m_H^2m_H^2\right\}C_0(p^2;W,G^\pm,H^\pm)$$
$$\left.+2c_{G^\pm hW^\mp}c_{G^\mp HW}\left(p^4-m_h^2m_H^2\right)C_0(p^2;W,G^\pm)\right]$$

$$\sum_{i=1,2}\overline{\mathcal{M}}_{Z-i}\Big|_{\text{G.D.}}=\frac{c_{\beta'}}{c_\beta}\left(\zeta_h c_{hZZ}\overline{\mathcal{M}}_{\text{ver1}}\left[\frac{g_Z}{\sqrt{2}},Z,h\right]\Big|_{\text{G.D.}}+\zeta_H c_{HZZ}\overline{\mathcal{M}}_{\text{ver1}}\left[\frac{g_Z}{\sqrt{2}},Z,H\right]\Big|_{\text{G.D.}}\right)$$

$$\begin{aligned}\sum_{i=3-6}\overline{\mathcal{M}}_{Z-i}\Big|_{\text{G.D.}}&=\zeta_{G^0}\zeta_h c_{G^0hZ}\overline{\mathcal{M}}_{\text{ver2}}\left[\frac{g_Z}{\sqrt{2}},Z,h\right]\Big|_{\text{G.D.}}+\zeta_{G^0}\zeta_H c_{G^0HZ}\overline{\mathcal{M}}_{\text{ver2}}\left[\frac{g_Z}{\sqrt{2}},Z,H\right]\Big|_{\text{G.D.}}\\&+\zeta_A\zeta_h c_{AhZ}\overline{\mathcal{M}}_{\text{ver3}}\left[\frac{g_Z}{\sqrt{2}},Z,h\right]\Big|_{\text{G.D.}}+\zeta_A\zeta_H c_{AHZ}\overline{\mathcal{M}}_{\text{ver3}}\left[\frac{g_Z}{\sqrt{2}},Z,H\right]\Big|_{\text{G.D.}}\end{aligned}$$

$$\overline{\mathcal{M}}_{Z-7}\big|_{\text{G.D.}}=\frac{c_{\beta'}^2}{c_\beta^2}\overline{\mathcal{M}}_{\text{box1}}\left[\frac{g_Z}{\sqrt{2}},Z\right]\Big|_{\text{G.D.}}$$

$$\sum_{i=8-11}\overline{\mathcal{M}}_{Z-i}\Big|_{\text{G.D.}}=\zeta_{G^0}^2\overline{\mathcal{M}}_{\text{box2}}\left[\frac{g_Z}{\sqrt{2}},Z,G^0\right]\Big|_{\text{G.D.}}+\zeta_A^2\overline{\mathcal{M}}_{\text{box2}}\left[\frac{g_Z}{\sqrt{2}},Z,A\right]\Big|_{\text{G.D.}}$$

$$\sum_{i=12-14}\overline{\mathcal{M}}_{Z-i}\Big|_{\text{G.D.}}=\zeta_h^2\overline{\mathcal{M}}_{\text{ver}+\delta}\left[\frac{g_Z}{\sqrt{2}},Z,h\right]\Big|_{\text{G.D.}}+\zeta_H^2\overline{\mathcal{M}}_{\text{ver}+\delta}\left[\frac{g_Z}{\sqrt{2}},Z,H\right]\Big|_{\text{G.D.}}$$

$$\sum_{i=1,2}\overline{\mathcal{M}}_{W-i}\Big|_{\text{G.D.}}=\zeta_h c_{hWW}\overline{\mathcal{M}}_{\text{ver1}}[g,W,h]\big|_{\text{G.D.}}+\zeta_H c_{HWW}\overline{\mathcal{M}}_{\text{ver1}}[g,W,H]\big|_{\text{G.D.}}$$

$$\begin{aligned}\sum_{i=3-6}\overline{\mathcal{M}}_{W-i}\Big|_{\text{G.D.}}&=\zeta_{G^\pm}\zeta_h c_{G^\pm hW^\mp}\overline{\mathcal{M}}_{\text{ver2}}[g,W,h]\big|_{\text{G.D.}}+\zeta_{G^\pm}\zeta_H c_{G^\pm HW^\mp}\overline{\mathcal{M}}_{\text{ver2}}[g,W,H]\big|_{\text{G.D}}\\&+\zeta_{H^\pm}\zeta_h c_{H^\pm hW^\mp}\overline{\mathcal{M}}_{\text{ver3}}[g,W,h]\big|_{\text{G.D.}}+\zeta_{H^\pm}\zeta_H c_{H^\pm HW^\mp}\overline{\mathcal{M}}_{\text{ver3}}[g,W,H]\big|_{\text{G.D.}}\end{aligned}$$

$$\overline{\mathcal{M}}_{W-7}\big|_{\text{G.D.}}=\overline{\mathcal{M}}_{\text{box1}}[g,W]\big|_{\text{G.D.}}$$

$$\sum_{i=8,9}\overline{\mathcal{M}}_{W-i}\Big|_{\text{G.D.}}=\zeta_{G^\pm}^2\overline{\mathcal{M}}_{\text{box2}}[g,W,G^\pm]\big|_{\text{G.D.}}+\zeta_{H^\pm}^2\overline{\mathcal{M}}_{\text{box2}}[g,W,H^\pm]\big|_{\text{G.D.}}$$

$$\overline{\mathcal{M}}_{W-10}\big|_{\text{G.D.}}=\zeta_h^2\overline{\mathcal{M}}_{\text{ver}+\delta}[g,W,h]\big|_{\text{G.D.}}+\zeta_{H^2}^2\overline{\mathcal{M}}_{\text{ver}+\delta}[g,W,H]\big|_{\text{G.D.}}$$

$$\begin{aligned}\overline{\mathcal{M}}_{\text{ver1}}[g_i,V,\phi]\big|_{\text{G.D.}}&=\frac{g_i^2}{16\pi^2}\frac{1}{p^2-m_\phi^2}\left[-\left(1+\frac{p^2}{2m_V^2}\right)B_0(p^2;V,V)\right.\\&\left.+\left(1-\xi_V+\frac{p^2}{m_V^2}\right)B_0(p^2;V,G_V)+\left(\xi_V-\frac{p^2}{2m_V^2}\right)B_0(p^2;G_V,G_V)\right],\\\overline{\mathcal{M}}_{\text{ver 2}}[g_i,V,\phi]\big|_{\text{G.D.}}&=\frac{g_i^2}{16\pi^2}\frac{1}{p^2-m_\phi^2}\left[B_0(p^2;V,V)-\left(1-\xi_V+\frac{p^2}{m_V^2}\right)B_0(p^2;V,G_V)\right.\\&\left.-\left(\xi_V-\frac{p^2}{m_V^2}\right)B_0(p^2;G_V,G_V)+(1-\xi_V)B_0(0;V,G_V)\right],\\\overline{\mathcal{M}}_{\text{ver3}}[g_i,V,\phi]\big|_{\text{G.D.}}&=\frac{g_i^2}{16\pi^2}\frac{1}{p^2-m_\phi^2}(1-\xi_V)\left[B_0(0;V,G_V)-\left(p^2-m_\phi^2\right)C_0(p^2;V,G_V,\phi)\right]\\\overline{\mathcal{M}}_{\text{box1}}[g_i,V]\big|_{\text{G.D.}}&=\frac{g_i^2}{64\pi^2}\frac{1}{m_V^2}[B_0(p^2;V,V)-2B_0(p^2;V,G_V)+B_0(p^2;G_V,G_V)]\\\overline{\mathcal{M}}_{\text{box2}}[g_i,V,\phi]\big|_{\text{G.D.}}&=\frac{g_i^2}{32\pi^2}\frac{1}{m_V^2}[B_0(p^2;V,\phi)-B_0(p^2;G_V,\phi)]\end{aligned}$$


$$\overline{\mathcal{M}}_{\text{ver}+\delta}[g_i,V,\phi]\big|_{\text{G.D.}}=-\frac{g_i^2}{32\pi^2}\frac{1}{p^2-m_\phi^2}(1-\xi_V)B_0(0;V,G_V)$$


$$\overline{\mathcal{M}}_{Z-1}\big|_{\text{G.D.}}\left(\overline{\mathcal{M}}_{W-1}\big|_{\text{G.D.}}\right)\text{ to }\overline{\mathcal{M}}_{Z-15}\big|_{\text{G.D}}\cdot\left(\overline{\mathcal{M}}_{W-11}\big|_{\text{G.D.}}\right)$$

$$\left(p^2-m_\phi^2\right)\left(p^2-m_{\phi'}^2\right)/\left(\zeta_\phi\zeta_{\phi'}\right)$$

$$\begin{aligned}
\Pi_{hh}^{\text{PT}}(p^2) =& -\frac{g^2}{16\pi^2}\left(p^2-m_h^2\right)\left[c_{G^\pm hW^\mp}^2 B_0(p^2;W,W)+c_{H^\pm HW^\mp}^2 B_0(p^2;W,H^\pm)\right]\\
&-\frac{g_Z^2}{32\pi^2}\left(p^2-m_h^2\right)\left[c_{G^0hZ}^2 B_0(p^2;Z,Z)+c_{AhZ}^2 B_0(p^2;Z,A)\right]\\
\Pi_{HH}^{\text{PT}}(p^2) =& -\frac{g^2}{16\pi^2}\left(p^2-m_H^2\right)\left[c_{G^\pm HW^\mp}^2 B_0(p^2;W,W)+c_{H^\pm HW^\mp}^2 B_0(p^2;W,H^\pm)\right]\\
&-\frac{g_Z^2}{32\pi^2}\left(p^2-m_H^2\right)\left[c_{G^0HZ}^2 B_0(p^2;Z,Z)+c_{AHZ}^2 B_0(p^2;Z,A)\right]\\
\Pi_{hH}^{\text{PT}}(p^2) =& -\frac{g^2}{32\pi^2}\left(2p^2-m_h^2-m_H^2\right)\left[c_{G^\pm hW^\mp}c_{G^\mp HW^\pm}B_0(p^2;W,W)\right.\\
&\left.+c_{H^\pm hW^\mp}c_{H^\mp HW^\pm}B_0(p^2;W,H^\pm)\right]\\
&-\frac{g_Z^2}{64\pi^2}\left(2p^2-m_h^2-m_H^2\right)\left[c_{G^0hZ^c}c_{G^0HZ}B_0(p^2;Z,Z)+c_{AhZ}c_{AHZ}B_0(p^2;Z,A)\right]
\end{aligned}$$

$$\begin{aligned}
\zeta_h &= \zeta_A c_{AhZ}+\zeta_{G^0}c_{G^0hZ}=\zeta_{G^\pm}c_{G^\pm hW^\mp}+\zeta_{H^\pm}c_{H^\pm hW^\mp}\\
\zeta_H &= \zeta_A c_{AHZ}+\zeta_{G^0}c_{G^0HZ}=\zeta_{G^\pm}c_{G^\pm HW^\mp}+\zeta_{H^\pm}c_{H^\pm HW^\mp}\\
\zeta_{G^0} &= \zeta_h c_{G^0hZ}+\zeta_H c_{G^0HZ}=\zeta_{G^\pm}c_{G^\pm G^0W}+\zeta_{H^\pm}c_{H^\pm G^0W^\mp}\\
\zeta_A &= \zeta_h c_{AhZ}+\zeta_H c_{AHZ}=\zeta_{G^\pm}c_{G^\pm AW}+\zeta_{H^\pm}c_{H^\pm AW^\mp}\\
\zeta_{G^\pm} &= \zeta_h c_{G^\pm hW^\mp}+\zeta_H c_{G^\pm HW^\mp}\\
\zeta_{H^\pm} &= \zeta_h c_{H^\pm hW^\mp}+\zeta_H c_{H^\pm HW^\mp}
\end{aligned}$$





$$\mathcal{M} = -\overline{\mathcal{M}}\left(\frac{m_f}{v}\right)^2 (\bar{f}\gamma_5 f) \times (\bar{f}\gamma_5 f).$$

$$\overline{\mathcal{M}}^{AA}_{W-8}\Big|_{\text{G.D.}} + \overline{\mathcal{M}}^{AA}_{Z-12}\Big|_{\text{G.D.}} = \frac{g^2}{16\pi^2}\frac{\zeta_A^2\zeta_{G^0}^2}{p^2 - m_A^2}(1-\xi_W)$$
$$\times\left[-B_0(0;W,G^\pm) + \left(p^2 + m_A^2 - 2m_{H^\pm}^2\right)C_0(p^2;W,G^\pm,H^\pm)\right]$$
$$+\frac{g_Z^2}{64\pi^2}\frac{\zeta_A^2}{p^2 - m_A^2}(1-\xi_Z)\left[-\left(c_{AHZ}^2 + c_{AhZ}^2\right)B_0(0;Z,G^0)\right.$$
$$+c_{AHZ}^2(p^2 + m_A^2 - 2m_H^2)C_0(p^2;Z,G^0,H)$$
$$\left.+c_{AhZ}^2\left(p^2 + m_A^2 - 2m_h^2\right)C_0(p^2;Z,G^0,h)\right]$$
$$\overline{\mathcal{M}}^{AG^0}_{W-8}\Big|_{\text{G.D.}} + \overline{\mathcal{M}}^{AG^0}_{Z-12}\Big|_{\text{G.D.}} = \frac{g^2}{32\pi^2}\frac{\zeta_A\zeta_{G^0}c_{H^\pm AW^\mp}c_{H^\pm G^0W^\mp}}{(p^2 - m_A^2)\left(p^2 - m_{G^0}^2\right)}(1-\xi_W)$$
$$\times\left[-(2p^2 - m_A^2)B_0(0;W,G^\pm) + 2\left\{p^2\left(p^2 - 2m_{H^\pm}^2\right) + m_A^2 m_{H^\pm}^2\right\}C_0(p^2;W,G^\pm,H^\pm)\right]$$

$$+\frac{g_Z^2}{128\pi^2}\frac{2\zeta_A\zeta_{G^0}}{(p^2-m_A^2)(p^2-m_{G^0}^2)}(1-\xi_Z)$$
$$\times[-(c_{AhZ}c_{G^0hZ}+c_{AHZ}c_{G^0HZ})(2p^2-m_A^2)B_0(0;Z,G^0)$$
$$+2c_{AhZ}c_{G^0hZ}\{m_A^2m_h^2+p^2(p^2-2m_h^2)\}C_0(p^2;Z,G^0,h)$$
$$+2c_{AHZ}c_{G^0HZ}\{m_A^2m_H^2+p^2(p^2-2m_H^2)\}C_0(p^2;Z,G^0,H)]$$

$$\sum_{i=1-11}\overline{\mathcal{M}}_{Z-i}\Big|_{\text{G.D.}}=\frac{g_Z^2}{64\pi^2}(1-\xi_Z)\left[\frac{\zeta_A}{p^2-m_A^2}\{\zeta_AB_0(0;Z,G^0)\right.$$
$$-2c_{AHZ}\zeta_H(p^2-m_H^2)C_0(p^2;Z,G^0,H)-2(p^2-m_h^2)c_{AhZ}\zeta_hC_0(p^2;Z,G^0,h)\}$$
$$+\frac{\zeta_{G^0}}{p^2-m_{G^0}^2}\{\zeta_{G^0}B_0(0;Z,G^0)-2c_{G^0HZ}\zeta_H(p^2-m_H^2)C_0(p^2;Z,G^0,H)$$
$$-2(p^2-m_h^2)c_{G^0hZ}\zeta_hC_0(p^2;Z,G^0,h)\}$$
$$\left.+\zeta_h^2C_0(p^2;Z,G^0,h)+\zeta_H^2C_0(p^2;Z,G^0,H)\right]$$


$$\sum_{i=1-7}\overline{\mathcal{M}}_{W-i}\Big|_{\text{G.D.}}=\frac{g^2(1-\xi_W)}{32\pi^2}\left[\frac{\zeta_A\zeta_{H^\pm}c_{H^\pm AW^\mp}}{p^2-m_A^2}\{B_0(0;W,G^\pm)-2(p^2-m_{H^+}^2)C_0(p^2;W,G^\pm,H^\pm)\}\right.$$
$$+\frac{\zeta_{G^0}}{p^2-m_{G^0}^2}\{-(\zeta_{G^0}-2\zeta_{H^\pm}c_{H^\pm G^0W^\mp})B_0(0;W,G^\pm)$$
$$-2(p^2-m_{H^+}^2)c_{H^\pm G^0W^\mp}\zeta_{H^\pm}C_0(p^2;W,G^\pm,H^\pm)\}$$
$$\left.+\zeta_{H^\pm}^2C_0(p^2;W,G^\pm,H^\pm)\right]$$

$$\Lambda_{G^0}=\frac{p^2}{p^2-m_{G^0}^2}\Lambda_{G^0}-im_Z\Lambda_Z^\mu(\Delta_Z)_{\mu\nu}p^\nu$$

$$\Lambda_{G^0}=-\frac{m_f}{v}\bar{f}\gamma_5f,\Lambda_Z^\mu=ig_Z\bar{f}\gamma^\mu(v_f-a_f\gamma_5)f$$
$$(\Delta_Z)_{\mu\nu}=\frac{1}{p^2-m_Z^2}\left(-g_{\mu\nu}+\frac{(1-\xi_Z)p_\mu p_\nu}{p^2-\xi_Zm_Z^2}\right)$$

$$\left(\frac{1}{m^2}+\frac{1}{p^2-m_{G^0}^2}+\frac{1}{p^2-m_A^2}\right)\Lambda_{G^0}=\left(\frac{p^2(p^2-m_A^2)}{m^2(p^2-m_{G^0}^2)(p^2-m_A^2)}+\frac{p^2-m_A^2}{(p^2-m_{G^0}^2)(p^2-m_A^2)}\right.$$
$$\left.+\frac{p^2}{(p^2-m_{G^0}^2)(p^2-m_A^2)}\right)\Lambda_{G^0}+\cdots$$

$$\Pi_{AA}^{\text{PT}}(p^2)=-\frac{g^2}{16\pi^2}c_{H^\pm AW^\mp}^2(p^2-m_A^2)B_0(p^2;W,H^\pm)$$
$$-\frac{g_Z^2}{32\pi^2}(p^2-m_A^2)\left[c_{AhZ}^2B_0(p^2;Z,h)+c_{AHZ}^2B_0(p^2;Z,H)\right]$$
$$\Pi_{AG^0}^{\text{PT}}(p^2)=-\frac{g^2}{32\pi^2}c_{H^\pm G^0W^\mp}c_{H^\pm AW^\mp}(2p^2-m_A^2)B_0(p^2;W,H^\pm)$$
$$-\frac{g_Z^2}{64\pi^2}(2p^2-m_A^2)[c_{hG^0Z^c}c_{AhZ}B_0(p^2;Z,h)+c_{HG^0Z^c}c_{AHZ}B_0(p^2;Z,H)]$$



$$\hat{\Gamma}_{hf\bar{f}}(p_1,p_2,q) = \hat{\Gamma}^{S}_{hf\bar{f}} + \gamma_5\hat{\Gamma}^{P}_{hf\bar{f}} + \not{p}_1\hat{\Gamma}^{V_1}_{hf\bar{f}} + \not{p}_2\hat{\Gamma}^{V_2}_{hf\bar{f}} + \not{p}_1\gamma_5\hat{\Gamma}^{A_1}_{hf\bar{f}} + \not{p}_2\gamma_5\hat{\Gamma}^{A_2}_{hf\bar{f}} + \not{p}_1\not{p}_2\hat{\Gamma}^{T}_{hf\bar{f}} + \not{p}_1\not{p}_2\gamma_5\hat{\Gamma}^{PT}_{hf\bar{f}}$$

$$\hat{\Gamma}^{P}_{hf\bar{f}} = \hat{\Gamma}^{PT}_{hf\bar{f}} = 0, \hat{\Gamma}^{V_1}_{hf\bar{f}} = -\hat{\Gamma}^{V_2}_{hf\bar{f}}, \hat{\Gamma}^{A_1}_{hf\bar{f}} = -\hat{\Gamma}^{A_2}_{hf\bar{f}} \text{ with } p_1^2 = p_2^2 = m_f^2.$$

$$\hat{\Gamma}^{X}_{hf\bar{f}} = \Gamma^{X,\text{tree}}_{hf\bar{f}} + \Gamma^{X,\text{loop}}_{hf\bar{f}}, (X = S, P, V_1, V_2, A_1, A_2, T, PT),$$

$$\Gamma^{S,\text{tree}}_{hf\bar{f}} = -\frac{m_f}{v}\zeta_h, \Gamma^{X,\text{tree}}_{hf\bar{f}} = 0 \text{ for } X \neq S.$$

$$\Gamma^{X,\text{loop}}_{hf\bar{f}} = \Gamma^{X,1PI}_{hf\bar{f}} + \delta\Gamma^{X}_{hf\bar{f}}.$$

$$\delta\Gamma^{S}_{hf\bar{f}} = \Gamma^{S,\text{tree}}_{hf\bar{f}}\left(\frac{\delta m_f}{m_f} - \frac{\delta v}{v} + \tan\beta\delta\beta + \delta Z_V^f + \frac{\delta Z_h}{2}\right) + \Gamma^{S,\text{tree}}_{Hf\bar{f}}\left(\delta\alpha + \frac{\delta Z_{Hh}}{2}\right)$$

$$\text{where } \Gamma^{S,\text{tree}}_{Hf\bar{f}} = -m_f\zeta_H/v, \text{ and } \delta\Gamma^{X}_{hf\bar{f}} = 0 \text{ for } X \neq S$$

$$\hat{\Gamma}^{\mu\nu}_{hVV}(p_1,p_2,q) = g^{\mu\nu}\hat{\Gamma}^{1}_{hVV} + \frac{p_1^\nu p_2^\mu}{m_V^2}\hat{\Gamma}^{2}_{hVV} + i\epsilon^{\mu\nu\rho\sigma}\frac{p_{1\rho}p_{2\sigma}}{m_V^2}\hat{\Gamma}^{3}_{hVV}$$

$$\hat{\Gamma}^{i}_{hVV} = \Gamma^{i,\text{tree}}_{hVV} + \Gamma^{i,\text{loop}}_{hVV}, (i = 1,2,3)$$

$$\Gamma^{1,\text{tree}}_{hVV} = c_V g_{hVV}, \Gamma^{2,,\text{tree}}_{hVV} = \Gamma^{3,\text{tree}}_{hVV} = 0$$

$$g_{hZZ} = \frac{1}{2}c_{hZZ}g_Z m_Z, g_{hWW} = c_{hWW}g m_W$$

$$\Gamma^{i,\text{loop}}_{hVV} = \Gamma^{i,1\text{PI}}_{hVV} + \Gamma^{i,\text{Tad}}_{hVV} + \delta\Gamma^{i}_{hVV}$$

$$\Gamma^{1,\text{Tad}}_{hVV} = c_V\left(2g_{hhVV}\frac{\Gamma_h^{1\text{PI}}}{m_h^2} + g_{hHVV}\frac{\Gamma_H^{1\text{PI}}}{m_H^2}\right), \Gamma^{2,\text{Tad}}_{hVV} = \Gamma^{3,\text{Tad}}_{hVV} = 0,$$

$$g_{hhZZ} = \frac{1}{8}c_{hhZZ}g_Z^2, g_{hhWW} = \frac{1}{4}c_{hhWW}g^2$$
$$g_{hHZZ} = \frac{1}{8}c_{hHZZ}g_Z^2, g_{hHWW} = \frac{1}{4}c_{hHWW}g^2$$

$$\hat{\Gamma}^{\mu\nu}_{hVV'}(p_1,p_2,q) = g^{\mu\nu}\hat{\Gamma}^{1}_{hVV'} + p_1^\nu p_2^\mu\hat{\Gamma}^{2}_{hVV'} + i\epsilon^{\mu\nu\rho\sigma}p_{1\rho}p_{2\sigma}\hat{\Gamma}^{3}_{hVV'}$$

$$\hat{\Gamma}^{1}_{hVV'} = -p_1\cdot p_2\hat{\Gamma}^{2}_{hVV'}$$

$$\hat{\Gamma}^{i}_{hVV'} = \Gamma^{i,\,\mathrm{loop}}_{hVV'}, (i = 1,2,3)$$

$$\Gamma^{i,\,\mathrm{loop}}_{hVV'} = \Gamma^{i,1\mathrm{PI}}_{hVV'} + \delta\Gamma^{i}_{hVV'}, (i = 1,2,3)$$

$$\delta\Gamma^{i}_{hgg} = \delta\Gamma^{i}_{h\gamma\gamma} = 0 (i = 1,2,3)$$

$$\delta\Gamma^{1}_{hZ\gamma} = g_{hZZ}\left(\delta Z_{Z\gamma} - \frac{\delta s^2_W}{c_W s_W}\right), \delta\Gamma^{2}_{hZ\gamma} = \delta\Gamma^{3}_{hZ\gamma} = 0$$

$$\Gamma(h \to f\bar{f}) = \Gamma_{\mathrm{LO}}(h \to f\bar{f})(1 - \Delta r) + \Gamma^{\mathrm{EW}}(h \to f\bar{f}) + \Gamma^{\mathrm{QCD}}(h \to f\bar{f})$$

$$\Gamma_{\mathrm{LO}}(h \to f\bar{f}) = N^f_c \frac{m_h}{8\pi}\left|\Gamma^{S,\,\mathrm{tree}}_{hf\bar{f}}\right|^2 \lambda^{3/2}\left(\frac{m^2_f}{m^2_h}, \frac{m^2_f}{m^2_h}\right)$$

$$\lambda(x, y) = (1 - x - y)^2 - 4xy$$

$$\Gamma^{\text{strong gravity}}(h \to f\bar{f}) = N^f_c \frac{m_h}{8\pi}\lambda^{3/2}\left(\frac{m^2_f}{m^2_h}, \frac{m^2_f}{m^2_h}\right) \times 2\mathrm{Re}\left\{\Gamma^{S,\,\mathrm{tree}}_{hf\bar{f}}\left[\Gamma^{S,\,\mathrm{loop}}_{hf\bar{f}} + 2m_f\Gamma^{V_1,\,\mathrm{loop}}_{hf\bar{f}} + m^2_h\left(1 - \frac{m^2_f}{m^2_h}\right)\Gamma^{T,\,\mathrm{loop}}_{hf\bar{f}}\right]^*\right\} + \Gamma(h \to f\bar{f}\gamma)$$

$$\Gamma(h \to f\bar{f}\gamma) = N^f_c \frac{2\alpha_{\mathrm{em}}Q^2_f}{16\pi^2 m_h}\left|\Gamma^{S,\,\mathrm{tree}}_{hf\bar{f}}\right|^2 [\Omega^{LL}_{11} + \Omega^{LR}_{11} + \Omega^{LL}_{22} + \Omega^{LR}_{22} + \Omega^{LL}_{12} + \Omega^{LR}_{12}],$$

$$h(p_h) \to Z(k_Z) + f(k_f) + \bar{f}(k_{\bar{f}})$$

$s = (k_{\bar{f}} + k_f)^2$, $t = (k_Z + k_{\bar{f}})^2$, and $u = (k_Z + k_f)^2$ and satisfy $s + t + u = m^2_h + m^2_Z$.

$$t = \frac{1}{2}\left[m^2_h + m^2_Z - s + m^2_h\lambda^{1/2}\left(\frac{m^2_Z}{m^2_h}, \frac{s}{m^2_h}\right)\cos\,\theta\right]$$

$$\Gamma(h \to ZZ^*) = \sum_{f \neq t}\Gamma(h \to Zf\bar{f})$$

$$\Gamma(h \to Zf\bar{f}) = \Gamma_{\mathrm{LO}}(h \to Zf\bar{f})(1 - 2\Delta r) + \Gamma^{\mathrm{EW}}(h \to Zf\bar{f}) + \Gamma^{\mathrm{QCD}}(h \to Zf\bar{f})$$

$$\Gamma_{\mathrm{LO}}(h \to Zf\bar{f}) = N^f_c\left(\frac{c_{hZZ}}{1 + s^2_\beta}\right)^2 \frac{G^2_F m_h m^4_Z}{24\pi^3}(v^2_f + a^2_f)F(\epsilon_Z)$$

$$F(x) = \frac{3(1 - 8x^2 + 20x^4)}{\sqrt{4x^2 - 1}}\arccos\left(\frac{3x^2 - 1}{2x^3}\right) - (1 - x^2)\left(\frac{47}{2}x^2 - \frac{13}{2} + \frac{1}{x^2} - 3(1 - 6x^2 + x^4)\ln\,x\right)$$

$$\Gamma^{\mathrm{EW}}(h \to Zf\bar{f}) = \int_0^{(m_h - m_Z)^2} \mathrm{d}s\,\frac{\mathrm{d}\Gamma_{hZZ}}{\mathrm{d}s} + \int_0^{(m_h - m_Z)^2} \mathrm{d}s\int_{-1}^{1} \mathrm{d}\cos\,\theta\,\frac{\mathrm{d}\Gamma_{hZZ}}{\mathrm{d}s\,\mathrm{d}\cos\,\theta} + \Gamma^{\mathrm{QED}}(h \to Zf\bar{f})$$

$$\frac{\mathrm{d}\Gamma_{hZZ}}{\mathrm{d}s} = N_c^f \frac{m_h^3}{96\pi^3}\frac{g_Z^2\Gamma_{hZZ}^{1,\text{ tree}}}{4m_Z^2}\left(v_f^2+a_f^2\right)\frac{c_a^{\text{kin}}}{(s-m_Z^2)^2}\,\mathrm{Re}\left\{-\Gamma_{hZZ}^{1,\text{ tree}}\frac{\hat{\Pi}'_{ZZ}(m_Z^2)}{2}\right.$$
$$+\left[\Gamma_{hZZ}^{1,\text{ loop}}+\frac{c_b^{\text{kin}}}{c_a^{\text{kin}}}\frac{\Gamma_{hZZ}^{2,\text{ loop}}}{m_Z^2}\right]\left(m_Z^2,s,m_h^2\right)-\Gamma_{hZZ}^{1,\text{ tree}}\frac{\hat{\Pi}_{ZZ}(s)}{s-m_Z^2}$$
$$+\frac{v_f Q_f c_W s_W}{v_f^2+a_f^2}\left(\frac{s-m_Z^2}{s}\left[\hat{\Gamma}_{hZ\gamma}^1+\frac{c_b^{\text{kin}}}{c_a^{\text{kin}}}\hat{\Gamma}_{hZ\gamma}^2\right]\left(m_Z^2,s,m_h^2\right)-\Gamma_{hZZ}^{1,\text{ tree}}\frac{\hat{\Pi}_{Z\gamma}(s)}{s}\right)$$
$$\left.+\Gamma_{hZZ}^{1,\text{ tree}}\frac{\left[v_f\Gamma_{Zff}^{V,\text{ loop}}+a_f\Gamma_{Zff}^{A,\text{ loop}}\right](0,0,s)}{v_f^2+a_f^2}\right\}$$

$$\hat{\Pi}'_{ZZ}(m_Z^2)=\mathrm{d}\hat{\Pi}_{ZZ}(p^2)/\mathrm{d}p^2|_{p^2=m_Z^2}$$

$$c_a^{\text{kin}}=\sqrt{x_1^2-4x_Z}\left[(x_1-6x_Z)^2+8x_Z(1-3x_Z)\right]$$
$$c_b^{\text{kin}}=\frac{m_h^2}{2}(x_1^2-4x_Z)^{3/2}(x_1-2x_Z)$$

with $x_Z=m_Z^2/m_h^2, x_s=s/m_h^2$ and $x_1=1+x_Z-x_s$

$$\frac{\mathrm{d}\Gamma_{hZZ}}{\mathrm{d}s\,\mathrm{d}\cos\,\theta}=N_c^f\frac{1}{512\pi^3 m_h}\sqrt{x_1^2-4x_Z}\,\frac{g_Z\Gamma_{hZZ}^{1,\text{ tree}}}{s-m_Z^2}\,\mathrm{Re}\left[c_1^{\text{kin}}T_{hf\bar{f}}^Z+B_Z\right]$$

$$c_1^{\text{kin}}=4s+2\left(tu-m_h^2m_Z^2\right)/m_Z^2.$$

$$T_{hf\bar{f}}^Z=\frac{2g_Z}{16\pi^2}\left\{4g_Z^3m_Z c_{hZZ}\left(v_f^4+6v_f^2a_f^2+a_f^4\right)\right.$$
$$\times\left[\left(\frac{C_0+C_{11}}{-2}\right)(t,0,m_h^2;Z,0,Z)+\left(\frac{C_{12}}{2}\right)(0,u,m_h^2;Z,0,Z)\right]$$
$$+g^3m_W c_{hWW}\left(v_f+a_f\right)^2$$
$$\left.\times\left[\left(\frac{C_0+C_{11}}{-2}\right)(t,0,m_h^2;W,0,W)+\left(\frac{C_{12}}{2}\right)(0,u,m_h^2;W,0,W)\right]\right\}$$

$$B_Z=\frac{1}{16\pi^2}\left[4\left(v_f+a_f\right)\sum_{i\neq 6}C^{BZi}B^{BZi}+8\left(v_f^4+6v_f^2a_f^2+a_f^4\right)C^{BZ6}B^{BZ6}\right]$$

$$\Gamma^{\text{QED}}(h\to Zf\bar{f})=\left(Q_f^2\frac{3\alpha_{\text{em}}}{4\pi}\right)\Gamma_{\text{LO}}(h\to Zf\bar{f}),$$
$$\Gamma^{\text{QCD}}(h\to Zf\bar{f})=\left(\frac{\alpha_s}{\pi}\right)\Gamma_{\text{LO}}(h\to Zf\bar{f})\ \text{ for }\ f=q$$

$$h(p_h)\to W^-(k_W)+f'\left(k_{f'}\right)+\bar{f}\left(k_{\bar{f}}\right)$$

$s=\left(k_{\bar{f}}+k_{f'}\right)^2$, $t=\left(k_W+k_{\bar{f}}\right)^2$, and $u=\left(k_W+k_{f'}\right)^2$ and satisfy $s+t+u=m_h^2+m_W^2$

$$\Gamma(h\to WW^*)=2\sum_{f,f'}\Gamma\left(h\to W^-f'\bar{f}\right)$$

$$\Gamma\left(h \to W^- f' \bar{f}\right) = \Gamma_{\mathrm{LO}}\left(h \to W^- f' \bar{f}\right)(1 - 2\Delta r) + \Gamma^{\text{strong gravity}}\left(h \to W^- f' \bar{f}\right) + \Gamma^{\mathrm{QCD}}\left(h \to W^- f' \bar{f}\right)$$

$$\Gamma_{\mathrm{LO}}\left(h \to W^- f' \bar{f}\right) = N_c^f c_{hWW}^2 \frac{G_F^2 m_h m_W^4}{96\pi^3} F(\epsilon_W),$$

$$\Gamma^{\text{strong gravity}}\left(h \to W^- f' \bar{f}\right) = R_{f'\bar{f}} \left[\int_0^{(m_h - m_W)^2} \mathrm{d}s \frac{\mathrm{d}\Gamma_{hWW}}{\mathrm{d}s} + \int_0^{(m_h - m_W)^2} \mathrm{d}s \int_{-1}^{1} \mathrm{d}\cos\,\theta \frac{\mathrm{d}\Gamma_{hWW}}{\mathrm{d}s\,\mathrm{d}\cos\,\theta} + \Gamma\left(h \to W^- f' \bar{f} \gamma\right) \right]$$

$$R_{f'\bar{f}} = \Gamma_{\mathrm{LO}}\left(h \to W^- f' \bar{f}\right)\Big|_{m_W = m_{W,G}} / \Gamma_{\mathrm{LO}}\left(h \to W^- f' \bar{f}\right)\Big|_{m_W = \bar{m}_W}$$

$$\frac{\mathrm{d}\Gamma_{hWW}}{\mathrm{d}s} = N_c^f \frac{m_h^3}{384\pi^3} \frac{g^2 \Gamma_{hWW}^{1,\,\mathrm{tree}}}{4m_W^2} \frac{c_a^{\mathrm{kin}}}{(s - m_W^2)^2} \mathrm{Re}\left\{ -\Gamma_{hWW}^{1,\,\mathrm{tree}} \frac{\widehat{\Pi}'_{WW}(m_W^2)}{2} + \left[\Gamma_{hWW}^{1,\,\mathrm{loop}} + \frac{c_b^{\mathrm{kin}}}{c_a^{\mathrm{kin}}} \frac{\Gamma_{hWW}^{2,\,\mathrm{loop}}}{m_W^2} \right] \left(m_W^2, s, m_h^2\right) - \Gamma_{hWW}^{1,\,\mathrm{tree}} \frac{\widehat{\Pi}_{WW}(s)}{s - m_W^2} + \Gamma_{hWW}^{1,\,\mathrm{tree}} \left[\Gamma_{Wff}^{V,\,\mathrm{loop}} + \Gamma_{Wff}^{A,\,\mathrm{loop}} \right](0,0,s) \right\}$$

$$\widehat{\Pi}'_{WW}(m_W^2) = \mathrm{d}\widehat{\Pi}_{WW}(p^2)/\mathrm{d}p^2\big|_{p^2 = m_W^2}$$

$$\frac{\mathrm{d}\Gamma_{hWW}}{\mathrm{d}s\,\mathrm{d}\cos\,\theta} = N_c^f \frac{1}{512\pi^3 m_h} \sqrt{x_1^2 - 4x_W} \frac{g_W \Gamma_{hWW}^{1,\,\mathrm{tree}}}{s - m_W^2} \mathrm{Re}\left[c_1^{\mathrm{kin}} T_{hf\bar{f}}^W + B_W \right]$$

$$T_{hf\bar{f}}^W = \frac{g_W}{16\pi^2} \left\{ 4g_Z^3 m_Z c_{hZZ} \left[\left(v_{f'} + a_{f'}\right)^2 \left(\frac{C_0 + C_{11}}{-2}\right)(t, 0, m_h^2; Z, 0, Z) + \left(v_f + a_f\right)^2 \left(\frac{C_{12}}{2}\right)(0, u, m_h^2; Z, 0, Z) \right] + 2g^3 m_W c_{hWW} \left[\left(\frac{C_0 + C_{11}}{-2}\right)(t, 0, m_h^2; W, 0, W) + \left(\frac{C_{12}}{2}\right)(0, u, m_h^2; W, 0, W) \right] \right\}$$

$$B_W = \frac{4}{16\pi^2} \sum C^{BWi} B^{BWi}$$

$$\Gamma^{\mathrm{QCD}}\left(h \to W^- f' \bar{f}\right) = \left(\frac{\alpha_s}{\pi}\right) \Gamma_{\mathrm{LO}}\left(h \to W^- f' \bar{f}\right) \text{ for } f = q$$

$$\Gamma_{\mathrm{LO}}(h \to \mathcal{V}\mathcal{V}') = \frac{1}{8\pi(1 + \delta_{\mathcal{V}\mathcal{V}'})m_h} \left|\hat{\Gamma}_{h\mathcal{V}\mathcal{V}'}^1\left(m_{\mathcal{V}}^2, m_{\mathcal{V}'}^2, m_h^2\right)\right|^2 \lambda^{1/2}\left(\frac{m_{\mathcal{V}}^2}{m_h^2}, \frac{m_{\mathcal{V}'}^2}{m_h^2}\right).$$

$$\Gamma_{\mathrm{LO}}(h \to \gamma\gamma) = \frac{\sqrt{2} G_F \alpha_{\mathrm{em}}^2 m_h^3}{256\pi^3} \left| \zeta_h \sum_f N_c^f Q_f^2 I_F^h(f) + c_{hWW} I_V^h(W) - \frac{\lambda_{H^+H^-h}}{v} I_S^h(H^\pm) - \frac{4\lambda_{H^{++}H^{--}h}}{v} I_S^h(H^{\pm\pm}) \right|^2$$

$$I_F^h(f) = -\frac{8m_f^2}{m_h^2}\left[1+\left(2m_f^2-\frac{m_h^2}{2}\right)C_0(0,0,h;f,f,f)\right]$$

$$I_V^h(W) = \frac{2m_W^2}{m_h^2}\left[6+\frac{m_h^2}{m_W^2}+\left(12m_W^2-6m_h^2\right)C_0(0,0,h;W,W,W)\right]$$

$$I_S^h(H^\pm) = \frac{2v^2}{m_h^2}\left[1+2m_{H^\pm}^2C_0(0,0,h;H^\pm,H^\pm,H^\pm)\right]$$

$$\Gamma_{\rm LO}(h\to Z\gamma)=\frac{\sqrt{2}G_F\alpha_{\rm em}^2m_h^3}{128\pi^3}\left(1-\frac{m_Z^2}{m_h^2}\right)^3\left|\zeta_h\sum_f N_c^f Q_f v_f J_F^h(f)+J_B^h\right|^2$$

$$J_F^h(f) = -\frac{4m_f^2}{s_W c_W(m_h^2-m_Z^2)}\left[2+\left(4m_f^2-m_h^2+m_Z^2\right)C_0(Z,0,h;f,f,f)+\frac{2m_Z^2}{m_h^2-m_Z^2}\left[B_0(h;f,f)-B_0(Z;f,f)\right]\right]$$

$$J_B^h=\frac{16\pi^2 v}{e^2(m_h^2-m_Z^2)}\hat{\Gamma}^1_{hZ\gamma}\left(m_Z^2,0,m_h^2\right)_B$$

$$\lambda_1=\frac{m_h^2}{2v^2},\lambda_2=-\frac{v^2}{2(m_\Phi^2-m_h^2)}\left(\lambda_4^2-\frac{4m_\Phi^2}{v^2}\lambda_4+\frac{4m_\Phi^2m_h^2}{v^4}\right),\lambda_3=\lambda_5=0$$

$$\lambda_4^-=\frac{m_h^2}{v^2},\lambda_4^+=\frac{4m_\Phi^2}{v^2}\left(1-\frac{m_h^2}{4m_\Phi^2}\right)$$

$$\bar{\Delta}_{\rm EW}(h\to XY)=\frac{\Gamma(h\to XY)}{\Gamma_{\rm LO}(h\to XY)}\Big|_{\rm CHTM}-\frac{\Gamma(h\to XY)}{\Gamma_{\rm LO}(h\to XY)}\Big|_{\rm SM},$$

$$\bar{\Delta}_{\rm EW}(h\to WW^*)$$

$$\mu\simeq\sqrt{2}m_\Phi^2\bar{v}_\Delta/\bar{v}^2$$

$$m_A^2=m_H^2,m_{H^\pm}^2-m_A^2=m_{H^{\pm\pm}}^2-m_{H^\pm}^2$$

$$m_A<m_{H^\pm}<m_{H^{\pm\pm}}$$

$$|\Delta m_W|\le 2.66\text{MeV},|\rho-1|\le 1.8\times10^{-3},|\Delta s_e^2|\le 6.6\times10^{-4}$$
$$|\Delta\Gamma(Z\to\bar{\ell})|\le 0.17\text{MeV}$$

$$h\to\gamma\gamma,\mu_{\gamma\gamma}=\sigma(gg\to h)\text{Br}(h\to\gamma\gamma)$$

$$\Delta\mu_{\gamma\gamma}=\frac{\mu_{\gamma\gamma}^{\rm CHTM}-\mu_{\gamma\gamma}^{\rm SM}}{\mu_{\gamma\gamma}^{\rm SM}}=\zeta_h^2\frac{\text{Br}(h\to\gamma\gamma)_{\rm CHTM}}{\text{Br}(h\to\gamma\gamma)_{\rm SM}}-1,$$

$$\sigma(gg\to h)_{\rm CHTM}=\zeta_h^2\sigma(gg\to h)_{\rm SM}$$

$$\bar{v}_\Delta, \lambda_3 (\simeq \lambda_5 \simeq 4(m_{H^\pm}^2 - m_{H^{\pm\pm}}^2)/v^2)$$

$$\lambda_4 + 2\sqrt{\lambda_1(\lambda_2 + \lambda_3/2)} > \Gamma(h \to XX)^{\mathrm{NP(SM)}}$$

$$\Delta R(h \to XX) = \frac{\Gamma(h \to XX)^{\mathrm{NP}}}{\Gamma(h \to XX)^{\mathrm{SM}}} - \Gamma(h \to XX)^{\mathrm{NP(SM)}},$$

$$\text{HS: } 300\text{GeV} \le m_A \le 1000\text{GeV}, 0 \le \lambda_4 \le 4\pi, 0 \le \Delta m \le 150\text{GeV},$$

$$\begin{aligned}
\text{2HDM Type-I: } & 400\text{GeV} < m_H < 1\text{TeV}, 0.99 < \sin(\beta - \alpha) < 1, \\
& 1.5 < \tan\beta < 10, 0 < M < m_H + 500\text{GeV}, \\
\text{2HDM Type-II: } & 800\text{GeV} < m_H < 1\text{TeV}, 0.995 < \sin(\beta - \alpha) < 1, \\
& 1 < \tan\beta < 10, 0 < M < m_H + 500\text{GeV}, \\
\text{HSM: } & 400\text{GeV} < m_H < 1000\text{GeV}, \lambda_S = 0.1, \mu_s = 0, \\
& 0.95 < \cos\alpha < 1, 0 < m_s < m_H + 500\text{GeV}, \\
\text{IDM: } & 100\text{GeV} < m_{H^\pm} < 1000\text{GeV}, m_H = 60\text{GeV}, m_A = m_{H^\pm} \\
& 0 < \lambda_2 < 4\pi, \mu_2^2 = 3581\text{GeV}^2.
\end{aligned}$$

$$\kappa_X^2 = \Gamma^{\mathrm{2HDM}}(h \to XX)/\Gamma^{\mathrm{SM}}(h \to XX)(X = \tau, b, g, \gamma) \text{ and } \kappa_\lambda = \lambda_{hhh}^{\mathrm{2HDM}}/\lambda_{hhh}^{\mathrm{SM}} 1314$$

$$\Delta\mu(h \to XX) = \frac{\mathrm{BR}(h \to XX)^{\mathrm{NP}}}{\mathrm{BR}(h \to XX)^{\mathrm{SM}}} - 1,$$

$$\Delta\mu(h \to XX) \simeq \Delta R(h \to XX) - \Delta R^{\mathrm{tot}},$$

$$\Delta R^{\mathrm{tot}} = \Gamma_h^{\mathrm{tot,\,NP}}/\Gamma_h^{\mathrm{tot,\,SM}} - 1$$

$$\Delta R^{\mathrm{tot}}, \Delta R(h \to XX) \ll 1$$

$$\Delta R^{\mathrm{tot}} = \Gamma_h^{\mathrm{tot,NP}}/\Gamma_h^{\mathrm{tot,SM}} - 1 \text{ with } \Gamma^{\mathrm{tot,NP(SM)}} \text{ being the total decay}$$

$$\text{LS: } 400\text{GeV} \le m_{H^{\pm\pm}} \le 1000\text{GeV}, -5 \le \lambda_4 \le 4\pi, 0 \le \Delta m \le 150\text{GeV},$$

$$\text{where } \Delta m = m_{H^\pm} - m_{H^{\pm\pm}}.$$

$$\begin{aligned}
\mathcal{L}_{\mathrm{kin}} \ni & \sum_{\phi\phi' = hh, hH, HH, AA, G^0A, G^0G^0} \left[\frac{g^2}{4} c_{\phi\phi' WW} \phi\phi' W^{\pm\mu} W_\mu^\mp + \frac{g_Z^2}{8} c_{\phi\phi' ZZ} \phi\phi' Z^\mu Z_\mu\right] \\
& + \sum_{\phi = h, H} \Big[g m_W c_{\phi WW} \phi W^{\pm\mu} W_\mu^\mp + \frac{g_Z}{2} m_Z c_{\phi ZZ} \phi Z^\mu Z_\mu \\
& \pm \frac{g}{2} c_{H^\pm \phi W^\mp} (p_1 - p_2)^\mu H^\pm \phi W_\mu^\mp \pm \frac{g}{2} c_{G^\pm \phi W^\mp} (p_1 - p_2)^\mu G^\pm \phi W_\mu^\mp \\
& - i\frac{g_Z}{2} c_{G^0\phi Z} (p_1 - p_2)^\mu G^0 \phi Z_\mu - i\frac{g_Z}{2} c_{A\phi Z} (p_1 - p_2)^\mu A \phi Z_\mu\Big] \\
& - \sum_{\phi = G^0, A} \left[i\frac{g}{2} c_{H^\pm \phi W^\mp} (p_1 - p_2)^\mu H^\pm \phi W_\mu^\mp + i\frac{g}{2} c_{G^\pm \phi W^\mp} (p_1 - p_2)^\mu G^\pm \phi W_\mu^\mp\right]
\end{aligned}$$

$$\left(\frac{m_f}{v}\right)^{-1} 16\pi^2\Gamma^{S,1\mathrm{PI}}_{hf\bar{f}}$$
$$= -4\zeta_h e^2 Q_f^2 C^{FVF}_{hff}(f,\gamma,f) - 4\zeta_h g_Z^2\left(v_f^2 - a_f^2\right)C^{FVF}_{hff}(f,Z,f)$$
$$+\frac{m_f^2}{v^2}\zeta_h\left[\zeta_h^2 C^{FSF}_{hff}(f,h,f) + \zeta_H^2 C^{FSF}_{hff}(f,H,f) - \zeta_{G^0}^2 C^{FSF}_{hff}(f,G^0,f) - \zeta_A^2 C^{FSF}_{hff}(f,A,f)\right]$$
$$-\frac{2m_{f'}^2}{v^2}\zeta_h\left[\zeta_{G^\pm}^2 C^{FSF}_{hff}(f',G^\pm,f') + \zeta_{H^\pm}^2 C^{FSF}_{hff}(f',H^\pm,f')\right]$$
$$-2g_Z^4 v^2\left(v_f^2 - a_f^2\right)\sqrt{1+s_\beta^2}\left(c_{\beta'}c_\alpha + 2s_{\beta'}s_\alpha\right)C_0(Z,f,Z)$$

Mixing factors	Expression	Mixing factors	Expression
c_{hhWW}	$\frac{1}{2}(3-c_{2\alpha})$	c_{hhZZ}	$\frac{1}{2}(5-3c_{2\alpha})$
c_{hHWW}	$s_{2\alpha}$	c_{hHZZ}	$3s_{2\alpha}$
c_{HHWW}	$\frac{1}{2}(3+c_{2\alpha})$	c_{HHZZ}	$\frac{1}{2}(5+3c_{2\alpha})$
c_{AAWW}	$\frac{1}{2}(3+c_{2\beta'})$	c_{AAZZ}	$\frac{1}{2}(5+3c_{2\beta'})$
c_{AG^0WW}	$s_{2\beta'}$	c_{AG^0ZZ}	$3s_{2\beta'}$
$c_{G^0G^0WW}$	$\frac{1}{2}(3-c_{2\beta'})$	$c_{G^0G^0ZZ}$	$\frac{1}{2}(5-3c_{2\beta'})$
c_{hWW}	$c_\beta c_\alpha + \sqrt{2}\,s_\beta s_\alpha$	c_{hZZ}	$c_{\beta'}c_\alpha + 2s_{\beta'}s_\alpha$
c_{HWW}	$-c_\beta s_\alpha + \sqrt{2}\,s_\beta c_\alpha$	c_{HZZ}	$-c_{\beta'}s_\alpha + 2s_{\beta'}c_\alpha$
$c_{G^\pm G^0W^\mp}$	$c_{\beta'}c_\beta + \sqrt{2}\,s_{\beta'}s_\beta$	$c_{G^\pm AW^\mp}$	0
$c_{H^\pm G^0W^\mp}$	$-c_{\beta'}s_\beta + \sqrt{2}\,s_{\beta'}c_\beta$	$c_{H^\pm AW^\mp}$	$s_{\beta'}s_\beta + \sqrt{2}\,c_{\beta'}c_\beta$
$c_{G^\pm hW^\mp}$	$c_\beta c_\alpha + \sqrt{2}\,s_\beta s_\alpha$	c_{G^0hZ}	$c_\alpha c_{\beta'} + 2s_\alpha s_{\beta'}$
$c_{H^\pm hW^\mp}$	$-s_\beta c_\alpha + \sqrt{2}\,c_\beta s_\alpha$	c_{G^0HZ}	$-c_{\beta'}s_\alpha + 2c_\alpha s_{\beta'}$
$c_{G^\pm HW^\mp}$	$-s_\alpha c_\beta + \sqrt{2}\,c_\alpha s_\beta$	c_{AhZ}	$-c_\alpha s_{\beta'} + 2s_\alpha c_{\beta'}$
$c_{H^\pm HW^\mp}$	$s_\alpha s_\beta + \sqrt{2}\,c_\alpha c_\beta$	c_{AHZ}	$2c_\alpha c_{\beta'} + s_\alpha s_{\beta'}$

$$-\frac{m_f^2}{v}\{6\lambda_{hhh}\zeta_h^2 C_0(h,f,h)+2\lambda_{HHh}\zeta_H^2 C_0(H,f,H)+2\lambda_{Hhh}\zeta_h\zeta_H[C_0(h,f,H)+C_0(H,f,h)]$$
$$-2\lambda_{G^0G^0h}\zeta_{G^0}^2 C_0(G^0,f,G^0)-2\lambda_{AAh}\zeta_A^2 C_0(A,f,A)-\lambda_{AG^0h}\zeta_{G^0}\zeta_A[C_0(A,f,G^0)+C_0(G^0,f,A)]\}$$
$$+\frac{2m_{f'}^2}{v}\{\lambda_{G^+G^-h}\zeta_{G^\pm}^2 C_0(G^\pm,f',G^\pm)+\lambda_{H^+H^-h}\zeta_{H^\pm}^2 C_0(H^\pm,f',H^\pm)$$
$$+\lambda_{H^+G^-h}\zeta_{G^\pm}\zeta_{H^\pm}[C_0(G^\pm,f',H^\pm)+C_0(H^\pm,f',G^\pm)]\}$$
$$-\frac{g^2}{4}\zeta_{G^\pm}\left(c_\beta c_\alpha+\sqrt{2}s_\beta s_\alpha\right)\left[C_{hff}^{VFS}(W,f',G^\pm)+C_{hff}^{SFV}(G^\pm,f',W)\right]$$
$$-\frac{g^2}{4}\zeta_{H^\pm}\left(-s_\beta c_\alpha+\sqrt{2}c_\beta s_\alpha\right)\left[C_{hff}^{VFS}(W,f',H^\pm)+C_{hff}^{SFV}(H^\pm,f',W)\right]$$
$$-\frac{g_Z^2}{8}\zeta_{G^0}\left(c_{\beta'}c_\alpha+2s_{\beta'}s_\alpha\right)\left[C_{hff}^{VFS}(Z,f,G^0)+C_{hff}^{SFV}(G^0,f,Z)\right]$$
$$-\frac{g_Z^2}{8}\zeta_A\left(-s_{\beta'}c_\alpha+2c_{\beta'}s_\alpha\right)\left[C_{hff}^{VFS}(Z,f,A)+C_{hff}^{SFV}(A,f,Z)\right]$$
$$\left(\frac{m_f}{v}\right)^{-1}16\pi^2\Gamma_{hf\bar{f}}^{P,1\mathrm{PI}}$$
$$=\frac{g^2}{4}\zeta_{G^\pm}\left(c_\beta c_\alpha+\sqrt{2}s_\beta s_\alpha\right)\left[C_{hff}^{VFS}(W,f',G^\pm)-C_{hff}^{SFV}(G^\pm,f',W)\right]$$
$$+\frac{g^2}{4}\zeta_{H^\pm}\left(-s_\beta c_\alpha+\sqrt{2}c_\beta s_\alpha\right)\left[C_{hff}^{VFS}(W,f',H^\pm)-C_{hff}^{SFV}(H^\pm,f',W)\right]$$

$$+2g_Z^2 v_f a_f \zeta_{G^0}\left(c_{\beta'}c_\alpha + 2s_{\beta'}s_\alpha\right)\left[C_{hff}^{VFS}(Z,f,G^0) - C_{hff}^{SFV}(G^0,f,Z)\right]$$
$$+2g_Z^2 v_f a_f \zeta_A\left(-s_{\beta'}c_\alpha + 2c_{\beta'}s_\alpha\right)\left[C_{hff}^{VFS}(Z,f,A) - C_{hff}^{SFV}(A,f,Z)\right],$$
$$16\pi^2\Gamma_{hf\bar{f}}^{V_1,1\mathrm{PI}}$$
$$= -\frac{2m_f^2}{v}\zeta_h\left[e^2Q_f^2(C_0+2C_{11})(f,\gamma,f) + g_Z^2\left(v_f^2+a_f^2\right)(C_0+2C_{11})(f,Z,f)\right]$$
$$-\frac{g^2}{2}\frac{m_{f'}^2}{v}\zeta_h(C_0+2C_{11})(f',W,f')$$
$$-\frac{m_f^4}{v^3}\zeta_h\left[\zeta_h^2(C_0+2C_{11})(f,h,f) + \zeta_H^2(C_0+2C_{11})(f,H,f)\right.$$
$$\left.+\zeta_{G^0}^2(C_0+2C_{11})(f,G^0,f) + \zeta_A^2(C_0+2C_{11})(f,A,f)\right]$$
$$-\frac{m_{f'}^2}{v^3}\left(m_f^2+m_{f'}^2\right)\zeta_h\left[\zeta_{G^\pm}^2(C_0+2C_{11})(f',G^\pm,f') + \zeta_{H^\pm}^2(C_0+2C_{11})(f',H^\pm,f')\right]$$
$$+\sqrt{1+s_\beta^2}\left(c_{\beta'}c_\alpha + 2s_{\beta'}s_\alpha\right)g_Z^4 v\left(v_f^2+a_f^2\right)(C_0+C_{11})(Z,f,Z)$$
$$+\left(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha\right)\frac{g^4}{4}v(C_0+C_{11})(W,f',W)$$
$$-\frac{m_f^2}{v^2}\left\{6\lambda_{hhh}\zeta_h^2(C_0+C_{11})(h,f,h) + 2\lambda_{HHh}\zeta_H^2(C_0+C_{11})(H,f,H)\right.$$
$$+2\lambda_{Hhh}\zeta_h\zeta_H[(C_0+C_{11})(H,f,h) + (C_0+C_{11})(h,f,H)]$$
$$+2\lambda_{G^0G^0h}\zeta_{G^0}^2(C_0+C_{11})(G^0,f,G^0) + 2\lambda_{AAh}\zeta_A^2(C_0+C_{11})(A,f,A)$$
$$\left.+\lambda_{AG^0h}\zeta_{G^0}\zeta_A[(C_0+C_{11})(A,f,G^0) + (C_0+C_{11})(G^0,f,A)]\right\}$$
$$-\frac{m_f^2+m_{f'}^2}{v^2}\left\{\lambda_{G^+G^-h}\zeta_{G^\pm}^2(C_0+C_{11})(G^\pm,f',G^\pm) + \lambda_{H^+H^-h}\zeta_{H^\pm}^2(C_0+C_{11})(H^\pm,f',H^\pm)\right.$$
$$\left.+\lambda_{H^+G^-h}\zeta_{G^\pm}\zeta_{H^\pm}[(C_0+C_{11})(G^\pm,f',H^\pm) + (C_0+C_{11})(H^\pm,f',G^\pm)]\right\}$$
$$+\frac{g^2}{4}\frac{m_{f'}^2}{v}\left[\zeta_{G^\pm}\left(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha\right)[(2C_0+C_{11})(W,f',G^\pm) + (-C_0+C_{11})(G^\pm,f',W)]\right.$$
$$\left.+\zeta_{H^\pm}\left(-s_\beta c_\alpha + \sqrt{2}c_\beta s_\alpha\right)[(2C_0+C_{11})(W,f',H^\pm) + (-C_0+C_{11})(H^\pm,f',W)]\right]$$
$$+\frac{g_Z^2}{8}\frac{m_f^2}{v}\left[\zeta_{G^0}\left(c_{\beta'}c_\alpha + 2s_{\beta'}s_\alpha\right)[(2C_0+C_{11})(Z,f,G^0) + (-C_0+C_{11})(G^0,f,Z)]\right.$$
$$\left.+\zeta_A\left(-s_{\beta'}c_\alpha + 2c_{\beta'}s_\alpha\right)[(2C_0+C_{11})(Z,f,A) + (-C_0+C_{11})(A,f,Z)]\right],$$
$$16\pi^2\Gamma_{hf\bar{f}}^{V_2,1\mathrm{PI}}$$
$$= -\frac{2m_f^2}{v}\zeta_h\left[e^2Q_f^2(C_0+2C_{12})(f,\gamma,f) + g_Z^2\left(v_f^2+a_f^2\right)(C_0+2C_{12})(f,Z,f)\right]$$
$$-\frac{g^2}{2}\frac{m_{f'}^2}{v}\zeta_h(C_0+2C_{12})(f',W,f')$$
$$-\frac{m_f^4}{v^3}\zeta_h\left[\zeta_h^2(C_0+2C_{12})(f,h,f) + \zeta_H^2(C_0+2C_{12})(f,H,f)\right.$$
$$\left.+\zeta_{G^0}^2(C_0+2C_{12})(f,G^0,f) + \zeta_A^2(C_0+2C_{12})(f,A,f)\right]$$

$$-\frac{m_{f'}^2}{v^3}\left(m_f^2+m_{f'}^2\right)\zeta_h\left[\zeta_{G^\pm}^2(C_0+2C_{12})(f',G^\pm,f')+\zeta_{H^\pm}^2(C_0+2C_{12})(f',H^\pm,f')\right]$$
$$+\sqrt{1+s_\beta^2}\left(c_{\beta'}c_\alpha+2s_{\beta'}s_\alpha\right)g_Z^4v\left(v_f^2+a_f^2\right)C_{12}(Z,f,Z)$$
$$+\left(c_\beta c_\alpha+\sqrt{2}s_\beta s_\alpha\right)\frac{g^4}{4}vC_{12}(W,f',W)$$
$$-\frac{m_f^2}{v^2}\{6\lambda_{hhh}\zeta_h^2C_{12}(h,f,h)+2\lambda_{HHh}\zeta_H^2C_{12}(H,f,H)$$
$$+2\lambda_{Hhh}\zeta_h\zeta_H[C_{12}(H,f,h)+C_{12}(h,f,H)]$$
$$+2\lambda_{G^0G^0h}\zeta_{G^0}^2C_{12}(G^0,f,G^0)+2\lambda_{AAh}\zeta_A^2C_{12}(A,f,A)$$
$$+\lambda_{AG^0h}\zeta_{G^0}\zeta_A[C_{12}(A,f,G^0)+C_{12}(G^0,f,A)]\}$$
$$-\frac{m_f^2+m_{f'}^2}{v^2}\{\lambda_{G^+G^-h}\zeta_{G^\pm}^2C_{12}(G^\pm,f',G^\pm)+\lambda_{H^+H^-h}\zeta_{H^\pm}^2C_{12}(H^\pm,f',H^\pm)$$
$$+\lambda_{H^+G^-h}\zeta_{G^\pm}\zeta_{H^\pm}[C_{12}(G^\pm,f',H^\pm)+C_{12}(H^\pm,f',G^\pm)]\}$$
$$+\frac{g^2}{4}\frac{m_{f'}^2}{v}\left[\zeta_{G^\pm}\left(c_\beta c_\alpha+\sqrt{2}s_\beta s_\alpha\right)[(2C_0+C_{12})(W,f',G^\pm)+(-C_0+C_{12})(G^\pm,f',W)]\right.$$
$$\left.+\zeta_{H^\pm}\left(-s_\beta c_\alpha+\sqrt{2}c_\beta s_\alpha\right)[(2C_0+C_{12})(W,f',H^\pm)+(-C_0+C_{12})(H^\pm,f',W)]\right]$$
$$+\frac{g_Z^2}{8}\frac{m_f^2}{v}\left[\zeta_{G^0}\left(c_{\beta'}c_\alpha+2s_{\beta'}s_\alpha\right)[(2C_0+C_{12})(Z,f,G^0)+(-C_0+C_{12})(G^0,f,Z)]\right.$$
$$\left.+\zeta_A\left(-s_{\beta'}c_\alpha+2c_{\beta'}s_\alpha\right)[(2C_0+C_{12})(Z,f,A)+(-C_0+C_{12})(A,f,Z)]\right],$$
$$16\pi^2\Gamma_{hf\bar{f}}^{A_1,1\mathrm{PI}}$$
$$=4g_Z^2v_fa_f\frac{m_f^2}{v}\zeta_h(C_0+2C_{11})(f,Z,f)+\frac{g^2}{2}\frac{m_{f'}^2}{v}\zeta_h(C_0+2C_{11})(f',W,f')$$
$$-\frac{m_{f'}^2}{v^3}\left(m_f^2-m_{f'}^2\right)\zeta_h\left[\zeta_{G^\pm}^2(C_0+2C_{11})(f',G^\pm,f')+\zeta_{H^\pm}^2(C_0+2C_{11})(f',H^\pm,f')\right]$$
$$-\sqrt{1+s_\beta^2}\left(c_{\beta'}c_\alpha+2s_{\beta'}s_\alpha\right)2g_Z^4v_fa_fv(C_0+C_{11})(Z,f,Z)$$
$$-\left(c_\beta c_\alpha+\sqrt{2}s_\beta s_\alpha\right)\frac{g^4}{4}v(C_0+C_{11})(W,f',W)$$
$$-\frac{m_f^2-m_{f'}^2}{v^2}\left[\zeta_{G^\pm}^2\lambda_{G^+G^-h}(C_0+C_{11})(G^\pm,f',G^\pm)+\zeta_{H^\pm}^2\lambda_{H^+H^-h}(C_0+C_{11})(H^\pm,f',H^\pm)\right.$$
$$\left.+\zeta_{G^\pm}\zeta_{H^\pm}\lambda_{H^+G^-h}[(C_0+C_{11})(G^\pm,f',H^\pm)+(C_0+C_{11})(H^\pm,f',G^\pm)]\right]$$
$$-\frac{g^2}{4}\frac{m_{f'}^2}{v}\left[\zeta_{G^\pm}\left(c_\beta c_\alpha+\sqrt{2}s_\beta s_\alpha\right)[(2C_0+C_{11})(W,f',G^\pm)+(-C_0+C_{11})(G^\pm,f',W)]\right.$$
$$\left.+\zeta_{H^\pm}\left(-s_\beta c_\alpha+\sqrt{2}c_\beta s_\alpha\right)[(2C_0+C_{11})(W,f',H^\pm)+(-C_0+C_{11})(H^\pm,f',W)]\right]$$
$$-2g_Z^2v_fa_f\frac{m_f^2}{v}\left[\zeta_{G^0}\left(c_{\beta'}c_\alpha+2s_{\beta'}s_\alpha\right)[(2C_0+C_{11})(Z,f,G^0)+(-C_0+C_{11})(G^0,f,Z)]\right.$$
$$\left.+\zeta_A\left(-s_{\beta'}c_\alpha+2c_{\beta'}s_\alpha\right)[(2C_0+C_{11})(Z,f,A)+(-C_0+C_{11})(A,f,Z)]\right],$$
$$16\pi^2\Gamma_{hf\bar{f}}^{A_2,1\mathrm{PI}}$$

$$= 4g_Z^2 v_f a_f \frac{m_f^2}{v}\zeta_h(C_0 + 2C_{12})(f,Z,f) + \frac{g^2}{2}\frac{m_{f'}^2}{v}\zeta_h(C_0 + 2C_{12})(f',W,f')$$
$$-\frac{m_{f'}^2}{v^3}\left(m_f^2 - m_{f'}^2\right)\zeta_h\left[\zeta_{G^\pm}^2(C_0 + 2C_{12})(f',G^\pm,f') + \zeta_{H^\pm}^2(C_0 + 2C_{12})(f',H^\pm,f')\right]$$
$$-\sqrt{1+s_\beta^2}\left(c_{\beta'}c_\alpha + 2s_{\beta'}s_\alpha\right)2g_Z^4 v_f a_f v C_{12}(Z,f,Z)$$
$$-\left(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha\right)\frac{g^4}{4}vC_{12}(W,f',W)$$
$$-\frac{m_f^2 - m_{f'}^2}{v^2}\left[\zeta_{G^\pm}^2\lambda_{G^+G^-h}C_{12}(G^\pm,f',G^\pm) + \zeta_{H^\pm}^2\lambda_{H^+H^-h}C_{12}(H^\pm,f',H^\pm)\right.$$
$$\left.+\zeta_{G^\pm}\zeta_{H^\pm}\lambda_{H^+G^-h}[C_{12}(G^\pm,f',H^\pm) + C_{12}(H^\pm,f',G^\pm)]\right]$$
$$-\frac{g^2}{4}\frac{m_{f'}^2}{v}\left[\zeta_{G^\pm}\left(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha\right)[(2C_0 + C_{12})(W,f',G^\pm) + (-C_0 + C_{12})(G^\pm,f',W)]\right.$$
$$\left.+\zeta_{H^\pm}\left(-s_\beta c_\alpha + \sqrt{2}c_\beta s_\alpha\right)[(2C_0 + C_{12})(W,f',H^\pm) + (-C_0 + C_{12})(H^\pm,f',W)]\right]$$
$$-2g_Z^2 v_f a_f \frac{m_f^2}{v}\left[\zeta_{G^0}\left(c_{\beta'}c_\alpha + 2s_{\beta'}s_\alpha\right)[(2C_0 + C_{12})(Z,f,G^0) + (-C_0 + C_{12})(G^0,f,Z)]\right.$$
$$\left.+\zeta_A\left(-s_{\beta'}c_\alpha + 2c_{\beta'}s_\alpha\right)[(2C_0 + C_{12})(Z,f,A) + (-C_0 + C_{12})(A,f,Z)]\right],$$

$$\left(\frac{m_f}{v}\right)^{-1} 16\pi^2\Gamma_{hf\bar{f}}^{T,1\mathrm{PI}}$$
$$= -\frac{m_f^2}{v^2}\zeta_h\left[\zeta_h^2(C_{11} - C_{12})(f,h,f) + \zeta_H^2(C_{11} - C_{12})(f,H,f)\right.$$
$$\left.-\zeta_{G^0}^2(C_{11} - C_{12})(f,G^0,f) - \zeta_A^2(C_{11} - C_{12})(f,A,f)\right]$$
$$+\frac{2m_{f'}^2}{v^2}\zeta_h\left[\zeta_{G^\pm}^2(C_{11} - C_{12})(f',G^\pm,f') + \zeta_{H^\pm}^2(C_{11} - C_{12})(f',H^\pm,f')\right]$$
$$-\frac{g^2}{4}\left[\zeta_{G^\pm}\left(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha\right)[(2C_0 + 2C_{11} - C_{12})(W,f',G^\pm) + (C_0 + C_{11} - 2C_{12})(G^\pm,f',W)]\right.$$
$$\left.+\zeta_{H^\pm}\left(-s_\beta c_\alpha + \sqrt{2}c_\beta s_\alpha\right)[(2C_0 + 2C_{11} - C_{12})(W,f',H^\pm) + (C_0 + C_{11} - 2C_{12})(H^\pm,f',W)]\right]$$
$$-\frac{g_Z^2}{8}\left[\zeta_{G^0}\left(c_{\beta'}c_\alpha + 2s_{\beta'}s_\alpha\right)[(2C_0 + 2C_{11} - C_{12})(Z,f,G^0) + (C_0 + C_{11} - 2C_{12})(G^0,f,Z)]\right.$$
$$\left.+\zeta_A\left(-s_{\beta'}c_\alpha + 2c_{\beta'}s_\alpha\right)[(2C_0 + 2C_{11} - C_{12})(Z,f,A) + (C_0 + C_{11} - 2C_{12})(A,f,Z)]\right]$$

$$\left(\frac{m_f}{v}\right)^{-1} 16\pi^2\Gamma_{hf\bar{f}}^{PT,1\mathrm{PI}}$$
$$= \frac{g^2}{4}\left[\zeta_{G^\pm}\left(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha\right)[(2C_0 + 2C_{11} - C_{12})(W,f',G^\pm) - (C_0 + C_{11} - 2C_{12})(G^\pm,f',W)]\right.$$
$$\left.+\zeta_{H^\pm}\left(-s_\beta c_\alpha + \sqrt{2}c_\beta s_\alpha\right)[(2C_0 + 2C_{11} - C_{12})(W,f',H^\pm) - (C_0 + C_{11} - 2C_{12})(H^\pm,f',W)]\right]$$
$$+2g_Z^2 v_f a_f\left[\zeta_{G^0}\left(c_{\beta'}c_\alpha + 2s_{\beta'}s_\alpha\right)[(2C_0 + 2C_{11} - C_{12})(Z,f,G^0) - (C_0 + C_{11} - 2C_{12})(G^0,f,Z)]\right.$$
$$\left.+\zeta_A\left(-s_{\beta'}c_\alpha + 2c_{\beta'}s_\alpha\right)[(2C_0 + 2C_{11} - C_{12})(Z,f,A) - (C_0 + C_{11} - 2C_{12})(A,f,Z)]\right]$$

$$C_{hff}^{FVF}(X,Y,Z)$$
$$= \left[m_f^2 C_0 + p_1^2(C_{11}+C_{21}) + p_2^2(C_{12}+C_{22}) + p_1\cdot p_2(2C_{23}-C_0) + 4C_{24}\right](X,Y,Z) - 1$$
$$C_{hff}^{FSF}(X,Y,Z)$$
$$= \left[m_f^2 C_0 + p_1^2(C_{11}+C_{21}) + p_2^2(C_{12}+C_{22}) + 2p_1\cdot p_2(C_{11}+C_{23}) + 4C_{24}\right](X,Y,Z) - \frac{1}{2}$$
$$C_{hff}^{VFS}(X,Y,Z)$$
$$= [p_1^2(2C_0+3C_{11}+C_{21}) + p_2^2(2C_{12}+C_{22}) + 2p_1\cdot p_2(2C_{12}+C_{23}) + 4C_{24}](X,Y,Z) - \frac{1}{2}$$

(B.10)

$$C_{hff}^{SFV}(X,Y,Z)$$
$$= [p_1^2(-C_0+C_{21}) + p_2^2(-C_{12}+C_{22}) + 2p_1\cdot p_2(C_{23}+C_{12}-C_0-C_{11}) + 4C_{24}](X,Y,Z)$$
$$-\frac{1}{2}.$$

$$16\pi^2\Gamma^{1,1\mathrm{PI}}_{hg^ag^b}(p_1^2,p_2^2,q^2)$$
$$= -\zeta_h\sum_q \frac{4g_s^2m_q^2}{v}[8C_{24}(q,q,q) - 2B_0(q^2;q,q) + (p_1^2+p_2^2-q^2)C_0(q,q,q)]\delta^{ab}$$
$$16\pi^2\Gamma^{2,1\mathrm{PI}}_{hg^ag^b}(p_1^2,p_2^2,q^2) = -\zeta_h\sum_q \frac{8g_s^2m_q^2}{v}q^2[C_0+4C_{1223}](q,q,q)\delta^{ab}$$

$$16\pi^2\Gamma^{1,1\mathrm{PI}}_{h\gamma\gamma}(p_1^2,p_2^2,q^2)_F$$
$$= -\zeta_h\sum_f N_c^f\frac{4e^2Q_f^2m_f^2}{v}[8C_{24}(f,f,f) - 2B_0(q^2;f,f) + (p_1^2+p_2^2-q^2)C_0(f,f,f)]$$
$$16\pi^2\Gamma^{2,1\mathrm{PI}}_{h\gamma\gamma}(p_1^2,p_2^2,q^2)_F = -\zeta_h\sum_f N_c^f\frac{8e^2Q_f^2m_f^2}{v}[C_0+4C_{1223}](f,f,f)$$
$$16\pi^2\Gamma^{1,1\mathrm{PI}}_{h\gamma\gamma}(p_1^2,p_2^2,q^2)_B$$
$$= \frac{2e^2m_W^2}{v}\left(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha\right)$$
$$\times\{(5p_1^2+5p_2^2-7q^2)C_0(W,W,W) - 6[B_0(q^2;W,W) - 4C_{24}(W,W,W)]$$
$$-\frac{m_h^2}{m_W^2}[B_0(q^2;W,W) - 4C_{24}(W,W,W)] - m_h^2C_0(W,W,W)\Big\}$$
$$-8e^2\lambda_{hH^{++}H^{--}}[4C_{24}(H^{\pm\pm},H^{\pm\pm},H^{\pm\pm}) - B_0(q^2;H^{\pm\pm},H^{\pm\pm})]$$
$$-2e^2\lambda_{hH^+H^-}[4C_{24}(H^{\pm},H^{\pm},H^{\pm}) - B_0(q^2;H^{\pm},H^{\pm})]$$
$$16\pi^2\Gamma^{2,1\mathrm{PI}}_{h\gamma\gamma}(p_1^2,p_2^2,q^2)_B$$
$$= \frac{8e^2m_W^2}{v}\left(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha\right)q^2\left[4C_0 + 6C_{1223} + \frac{m_h^2}{m_W^2}C_{1223}\right](W,W,W)$$
$$-32e^2\lambda_{hH^{++}H^{--}}q^2C_{1223}(H^{\pm\pm},H^{\pm\pm},H^{\pm\pm}) - 8e^2\lambda_{hH^+H^-}q^2C_{1223}(H^{\pm},H^{\pm},H^{\pm}).$$

$$16\pi^2\Gamma_{hZ\gamma}^{1,1\mathrm{PI}}(p_1^2,p_2^2,q^2)_F$$
$$=-\zeta_h\sum_f N_c^f v_f Q_f\frac{4eg_Z m_f^2}{v}[8C_{24}(f,f,f)-2B_0(q^2;f,f)+(p_1^2+p_2^2-q^2)C_0(f,f,f)]$$
$$16\pi^2\Gamma_{hZ\gamma}^{2,1\mathrm{PI}}(p_1^2,p_2^2,q^2)_F=-\zeta_h\sum_f N_c^f v_f Q_f\frac{8eg_Z m_f^2}{v}[C_0+4C_{1223}](f,f,f),$$
$$16\pi^2\Gamma_{hZ\gamma}^{1,1\mathrm{PI}}(p_1^2,p_2^2,q^2)_B$$
$$=eg^2m_W c_W\left(c_\alpha c_\beta+\sqrt{2}s_\alpha s_\beta\right)\left[2C_{hVV1}^{VVV}(W,W,W)-6B_0(q^2;W,W)+4-2C_{24}(c^\pm,c^\pm,c^\pm)\right]$$
$$+egg_Z m_W\left(s_W^2+s_\beta^2\right)\left(c_\alpha c_\beta+\sqrt{2}s_\alpha s_\beta\right)$$
$$\times\left[C_{hVV1}^{SVV}(G^\pm,W,W)+2m_W^2C_0(W,G^\pm,W)-2C_{24}(W,G^\pm,G^\pm)+B_0(p_1^2;G^\pm,W)\right]$$
$$+egg_Z m_W\left(c_{2W}-s_\beta^2\right)\left(c_\alpha c_\beta+\sqrt{2}s_\alpha s_\beta\right)C_{24}(G^\pm,G^\pm,W)$$
$$-eg^2m_W c_W\left(c_\alpha c_\beta+\sqrt{2}s_\alpha s_\beta\right)C_{hVV1}^{VVS}(W,W,G^\pm)$$
$$-2eg_Z m_W^2\left(s_W^2+s_\beta^2\right)\lambda_G+G^-h$$
$$-egC_0m_W\left[-c_\alpha c_\beta s_W^2+\sqrt{2}s_\alpha s_\beta(c_W^2-2)\right]B_0(p_2^2;G^\pm,W)$$
$$-4eg_Z\left(c_{2W}-s_\beta^2\right)\lambda_{G^+G}-h\left[C_{24}(G^\pm,G^\pm,G^\pm)-\frac{1}{4}B_0(q^2;G^\pm,G^\pm)\right]$$
$$+egg_Z m_W s_\beta c_\beta\left(-c_\alpha s_\beta+\sqrt{2}s_\alpha c_\beta\right)\left[C_{hVV1}^{SVV}(H^\pm,W,W)-2C_{24}(W,H^\pm,H^\pm)\right]$$
$$-2eg_Z m_W^2 s_\beta c_\beta\lambda_{H^+G^-h}C_0(H^\pm,W,G^\pm)$$
$$-egg_Z m_W s_\beta c_\beta\left(-c_\alpha s_\beta+\sqrt{2}s_\alpha c_\beta\right)[C_{24}(H^\pm,G^\pm,W)-B_0(p_1^2;H^\pm,W)]$$
$$-4eg_Z\left(c_{2W}-c_\beta^2\right)\lambda_{H^+H^-h}\left[C_{24}(H^\pm,H^\pm,H^\pm)-\frac{1}{4}B_0(q^2;H^\pm,H^\pm)\right]$$
$$-16eg_Z c_{2W}\lambda_{H^{++}H^{--}h}\left[C_{24}(H^{\pm\pm},H^{\pm\pm},H^{\pm\pm})-\frac{1}{4}B_0(q^2;H^{\pm\pm},H^{\pm\pm})\right]$$
$$+4eg_Z s_\beta c_\beta\lambda_{H^+G^-h}\left[C_{24}(H^\pm,G^\pm,G^\pm)+C_{24}(G^\pm,H^\pm,H^\pm)-\frac{1}{2}B_0(q^2;H^\pm,G^\pm)\right]$$

$$16\pi^2\Gamma_{hZ\gamma}^{2,1\mathrm{PI}}(p_1^2,p_2^2,q^2)_B$$
$$=eg^2m_W c_W\left(c_\alpha c_\beta+\sqrt{2}s_\alpha s_\beta\right)\left[2C_{hVV2}^{VVV}(W,W,W)-2C_{1223}(c^\pm,c^\pm,c^\pm)-C_{hVV2}^{VVS}(W,W,G^\pm)\right]$$
$$+egg_Z m_W\left(s_W^2+s_\beta^2\right)\left(c_\alpha c_\beta+\sqrt{2}s_\alpha s_\beta\right)\left[C_{hVV2}^{SVV}(G^\pm,W,W)-2C_{hVV2}^{VSS}(W,G^\pm,G^\pm)\right]$$
$$+egg_Z m_W\left(c_{2W}-s_\beta^2\right)\left(c_\alpha c_\beta+\sqrt{2}s_\alpha s_\beta\right)C_{hVV2}^{SSV}(G^\pm,G^\pm,W)$$
$$-4eg_Z\left(c_{2W}-s_\beta^2\right)\lambda_{G^+G^-h}C_{1223}(G^\pm,G^\pm,G^\pm)$$
$$+egg_Z m_W s_\beta c_\beta\left(-c_\alpha s_\beta+\sqrt{2}s_\alpha c_\beta\right)$$
$$\times\left[C_{hVV2}^{SVV}(H^\pm,W,W)-C_{hVV2}^{SSV}(H^\pm,G^\pm,W)-2C_{hVV2}^{VSS}(W,H^\pm,H^\pm)\right]$$
$$-4eg_Z\left(c_{2W}-c_\beta^2\right)\lambda_{H^+H^-h}C_{1223}(H^\pm,H^\pm,H^\pm)$$
$$-16eg_Z c_{2W}\lambda_{H^{++}H^{--}}C_{1223}(H^{\pm\pm},H^{\pm\pm},H^{\pm\pm})$$
$$+4eg_Z s_\beta c_\beta\lambda_{H^+G^-h}[C_{1223}(H^\pm,G^\pm,G^\pm)+C_{1223}(G^\pm,H^\pm,H^\pm)],$$

$$
\begin{aligned}
C_{hVV1}^{VVV}(X,Y,Z) =& [18C_{24} + p_1^2(2C_{21} + 3C_{11} + C_0) + p_2^2(2C_{22} + C_{12}) \\
&+ p_1 \cdot p_2(4C_{23} + 3C_{12} + C_{11} - 4C_0)](X,Y,Z) - 3 \\
C_{hVV2}^{VVV}(X,Y,Z) =& (10C_{23} + 9C_{12} + C_{11} + 5C_0)(X,Y,Z) \\
C_{hVV1}^{SVV}(X,Y,Z) =& [3C_{24} + p_1^2(C_{21} - C_0) + p_2^2(C_{22} - 2C_{12} + C_0) \\
&+ 2p_1 \cdot p_2(C_{23} - C_{11})](X,Y,Z) - \frac{1}{2}, \\
C_{hVV2}^{SVV}(X,Y,Z) =& (4C_{11} - 3C_{12} - C_{23})(X,Y,Z), \\
C_{hVV1}^{VVS}(X,Y,Z) =& [3C_{24} + p_1^2(C_{21} + 4C_{11} + 4C_0) + p_2^2(C_{22} + 2C_{12}) \\
&+ 2p_1 \cdot p_2(C_{23} + 2C_{12} + C_{11} + 2C_0)](X,Y,Z) - \frac{1}{2}, \\
C_{hVV2}^{VVS}(X,Y,Z) =& (2C_{11} - 5C_{12} - 2C_0 - C_{23})(X,Y,Z), \\
C_{hVV2}^{SSV}(X,Y,Z) =& (C_{23} - C_{12})(X,Y,Z), \\
C_{hVV2}^{VSS}(X,Y,Z) =& (C_{23} + C_{12} + 2C_{11} + 2C_0)(X,Y,Z).
\end{aligned}
$$


$k_Z = -p_Z, k_{\bar{f}} = -p_f$ and $k_f = -p_{\bar{f}}$

the Mandelstam variables are $s = (p_f + p_{\bar{f}})^2, t = (p_Z + p_f)^2$, and $u = (p_Z + p_{\bar{f}})^2$

The LO amplitude is:

$$\mathcal{M}_{\text{LO}} = C_{\text{tree}}^{Z}\, \bar{v}(p_{\bar{f}})\gamma^{\mu}(v_f - a_f\gamma_5)u(p_f)\epsilon_{\mu}^{*}(p_Z)$$

$$C_{\text{tree}}^{Z} = g_Z \Gamma_{hZZ}^{1,\,\text{tree}}/(s - m_Z^2)$$



$$\mathcal{M}_{BZi} = \frac{1}{16\pi^2}\, \bar{v}(p_{\bar{f}})\mathcal{M}_{BZi}^{\mu} u(p_f)\epsilon_{\mu}^{*}(p_Z),$$

$$\mathcal{M}^{\mu}_{BZi} = C^{BZi}\left[\not{p}_h\left(F^{BZi}_{1,3}p^{\mu}_f + F^{BZi}_{1,4}p^{\mu}_{\bar{f}}\right) + \gamma^{\mu}F^{BZi}_{\gamma}\right]P_L \ (i \neq 6)$$
$$\mathcal{M}^{\mu}_{BZ6} = C^{BZ6}\left[\not{p}_h\left(F^{BZ6}_{1,3}p^{\mu}_f + F^{BZ6}_{1,4}p^{\mu}_{\bar{f}}\right) + \gamma^{\mu}F^{BZ6}_{\gamma}\right](v_f - a_f\gamma_5)^3$$

$$(16\pi^2)2\mathrm{Re}\left[\sum \mathcal{M}^{*}_{\mathrm{tree}}\,\mathcal{M}_{BZi}\right] = 4(v_f + a_f)C^{Z}_{\mathrm{tree}}\,C^{BZi}B^{BZi} \ (i \neq 6),$$
$$(16\pi^2)2\mathrm{Re}\left[\sum \mathcal{M}^{*}_{\mathrm{tree}}\,\mathcal{M}_{BZ6}\right] = 8(v_f^4 + 6v_f^2a_f^2 + a_f^4)C^{Z}_{\mathrm{tree}}\,C^{BZ6}B^{BZ6},$$

$$B^{BZi} = sF^{BZi}_{\gamma} + \frac{tu - m_Z^2m_h^2}{4m_Z^2}\left[2F^{BZi}_{\gamma} - (t - m_Z^2)F^{BZi}_{1,3} - (u - m_Z^2)F^{BZi}_{1,4}\right].$$

$$D_{i,ij}(p_1^2, p_2^2, p_3^2, p_4^2, p_{12}^2, p_{23}^2; X, Y, Z, W)$$
$$= D_{i,ij}(p_1^2, p_2^2, p_3^2, p_{123}^2, p_{12}^2, p_{23}^2; m_X^2, m_Y^2, m_Z^2, m_W^2)$$

$$p_{123}^2 = (p_1 + p_2 + p_3)^2, p_{12}^2 = (p_1 + p_2)^2, p_{23}^2 = (p_2 + p_3)^2$$

$$C^{BZ1} = -2a_f g^4 m_W c_W(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha)$$
$$F^{BZ1}_{1,3} = F^{VFVV}_{p\cdot\gamma 1}(0,0, m_Z^2, m_h^2, s, t; W, 0, W, W)$$
$$F^{BZ1}_{1,4} = F^{VFVV}_{p\cdot\gamma 2}(0,0, m_Z^2, m_h^2, s, t; W, 0, W, W)$$
$$F^{BZ1}_{\gamma} = F^{VFVV}_{\gamma}(0,0, m_Z^2, m_h^2, s, t; W, 0, W, W)$$

$$F^{VFV}_{p\cdot\gamma 1} = 2(D_{13} - D_{12} + 2D_{26})$$
$$F^{VFVV}_{p\cdot\gamma 2} = 4(D_0 + D_{11} + D_{13} + D_{25})$$
$$F^{VFVV}_{\gamma} = -2C_0(p_{12}^2, p_3^2, p_{123}^2; m_{V_1}^2, m_{V_3}^2, m_{V_4}^2) - [4D_{27} + (2p_{12}^2 + p_{23}^2 - p_{123}^2)(D_0 + D_{11})$$
$$+2(p_{23}^2 - p_3^2)D_{12} + p_{23}^2D_{13}]$$

$$C^{BZ3} = a_f g^3 g_Z m_W(s_W^2 + s_\beta^2)(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha)$$
$$F^{BZ3}_{1,3} = F^{VFVS}_{p\cdot\gamma}(0,0, m_Z^2, m_h^2, s, t; W, 0, W, G^{\pm})$$
$$F^{BZ3}_{1,4} = 0$$
$$F^{BZ3}_{\gamma} = F^{VFVS}_{\gamma}(0,0, m_Z^2, m_h^2, s, t; W, 0, W, G^{\pm})$$

$$F^{VFVS}_{p\cdot\gamma} = -4(D_{12} - D_{13})$$
$$F^{VFVS}_{\gamma} = C_0(p_{12}^2, p_3^2, p_{123}^2; m_{V_1}^2, m_{V_3}^2, m_{S_4}^2)$$
$$+2[-(p_{23}^2 - p_{123}^2)(D_0 + D_{11}) + p_{23}^2D_{13}].$$

$$C^{BZ5} = -(v_{f'} + a_{f'})g^3 g_Z m_W(c_\beta c_\alpha + \sqrt{2}s_\beta s_\alpha)$$
$$F^{BZ5}_{1,3} = F^{VFFV}_{p\cdot\gamma 1}(0, m_Z^2, 0, m_h^2, t, u; W, 0, 0, W)$$
$$F^{BZ5}_{1,4} = F^{VFFV}_{p\cdot\gamma 2}(0, m_Z^2, 0, m_h^2, t, u; W, 0, 0, W)$$
$$F^{BZ5}_{\gamma} = F^{VFFV}_{\gamma}(0, m_Z^2, 0, m_h^2, t, u; W, 0, 0, W)$$

$$F^{VFFV}_{p\cdot\gamma 1} = -2(D_0 + D_{11} + D_{12} + D_{24})$$
$$F^{VFFV}_{p\cdot\gamma 2} = -2D_{26}$$
$$F^{VFFV}_{\gamma} = -C_0(p_{12}^2, p_3^2, p_{123}^2; m_{V_1}^2, m_{F_3}^2, m_{V_4}^2)$$
$$-[-2D_{27} + (p_{12}^2 - p_2^2)(D_0 + D_{11}) + p_2^2D_{12}].$$

$$C^{BZ6} = -2g_Z^4 m_Z\left(c_{\beta'}c_\alpha + 2s_{\beta'}s_\alpha\right)$$

$$C^{BZ7} = a_f g^3 g_Z m_W s_\beta c_\beta\left(-s_\beta c_\alpha + \sqrt{2}c_\beta s_\alpha\right)$$
$$F_{1,3}^{BZ7} = F_{p\cdot\gamma}^{VFVS}\left(0,0,m_Z^2,m_h^2,s,t;W,0,W,H^\pm\right)$$
$$F_{1,4}^{BZ7} = 0$$
$$F_\gamma^{BZ7} = F_\gamma^{VFVS}\left(0,0,m_Z^2,m_h^2,s,t;W,0,W,H^\pm\right)$$

$k_W = -p_W, k_{\bar{f}} = -p_f$ and $k_{f'} = -p_{\bar{f}'}$

the Mandelstam variables are obtained as $s = \left(p_f + p_{\bar{f}'}\right)^2$, $t = \left(p_W + p_f\right)^2$, and $u = \left(p_W + p_{\bar{f}'}\right)^2$.

The LO amplitude is:


$$\mathcal{M}_{\text{tree}} = C_{\text{tree}}^W\, \bar{v}\left(p_{\bar{f}'}\right)\gamma^\mu P_L u\left(p_f\right)\epsilon_\mu^*\left(p_W\right)$$

where $C_{\text{tree}}^W = g_W \Gamma_{hWW}^{1,\text{tree}}/(s - m_W^2)$

$$\mathcal{M}_{BWi} = \frac{1}{16\pi^2}\bar{v}\left(p_{\bar{f}'}\right)\mathcal{M}_{BWi}^\mu u\left(p_f\right)\epsilon_\mu^*\left(p_W\right)$$

$$\mathcal{M}_{BWi}^\mu = C^{BWi}\left[\not{p}_h\left(F_{1,3}^{BWi}p_f^\mu + F_{1,4}^{BWi}p_{\bar{f}'}^\mu\right) + \gamma^\mu F_\gamma^{BWi}\right]P_L$$

$$(16\pi^2)2\text{Re}\left[\sum \mathcal{M}_{\text{tree}}^*\,\mathcal{M}_{BWi}\right] = 4C_{\text{tree}}^W\, C^{BWi}\text{Re}\left[B^{BWi}\right]$$



$$B^{BWi} = sF_\gamma^{BWi} + \frac{tu - m_W^2 m_h^2}{4m_W^2}\left[2F_\gamma^{BWi} - (t - m_W^2)F_{1,3}^{BWi} - (u - m_W^2)F_{1,4}^{BWi}\right].$$

$$C^{BW1}= -4a_{f'}\left(v_{f'}+a_{f'}\right)\frac{g^2g_Z^2}{\sqrt{2}}c_W m_Z c_{hZZ},$$
$$F_{1,3}^{BW1}= F_{p\cdot\gamma 1}^{VFVV}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;Z,f',W,Z\right),$$
$$F_{1,4}^{BW1}= F_{p\cdot\gamma 2}^{VFVV}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;Z,f',W,Z\right),$$
$$F_{\gamma}^{BW1}= F_{\gamma}^{VFVV}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;Z,f',W,Z\right).$$

$$C^{BW3}= -4a_f\left(v_f+a_f\right)\frac{g^3g_Z}{\sqrt{2}}c_W m_W c_{hWW}$$
$$F_{1,3}^{BW3}= F_{p\cdot\gamma 1}^{VFVV}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;W,f,Z,W\right)$$
$$F_{1,4}^{BW3}= F_{p\cdot\gamma 2}^{VFVV}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;W,f,Z,W\right)$$
$$F_{\gamma}^{BW3}= F_{\gamma}^{VFVV}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;W,f,Z,W\right)$$

$$C^{BW5}= -4a_f eQ_f\frac{g^3}{\sqrt{2}}s_W m_W c_{hWW}$$
$$F_{1,3}^{BW5}= F_{p\cdot\gamma 1}^{VFVV}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;W,f,\gamma,W\right)$$
$$F_{1,4}^{BW5}= F_{p\cdot\gamma 2}^{VFVV}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;W,f,\gamma,W\right)$$
$$F_{\gamma}^{BW5}= F_{\gamma}^{VFVV}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;W,f,\gamma,W\right)$$

$$C^{BW7}= -2a_f\left(v_f+a_f\right)\frac{g^2g_Z^2}{\sqrt{2}}\left(s_W^2+s_\beta^2\right)m_W c_{hWW}$$
$$F_{1,3}^{BW7}= F_{p\cdot\gamma}^{VFVS}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;W,f,Z,G^{\pm}\right)$$
$$F_{1,4}^{BW7}= 0$$
$$F_{\gamma}^{BW7}= F_{\gamma}^{VFVS}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;W,f,Z,G^{\pm}\right)$$

$$C^{BW9}= 2a_f Q_f\frac{e^2g^2}{\sqrt{2}}m_W c_{hWW}$$
$$F_{1,3}^{BW9}= F_{p\cdot\gamma}^{VFVS}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;W,f,\gamma,G^{\pm}\right)$$
$$F_{1,4}^{BW9}= 0$$
$$F_{\gamma}^{BW9}= F_{\gamma}^{VFVS}\left(m_{f'}^2,m_f^2,m_W^2,m_h^2,s,t;W,f,\gamma,G^{\pm}\right)$$

$$C^{BW11}= -2\left(v_f+a_f\right)\left(v_{f'}+a_{f'}\right)\frac{g_Z^3g}{\sqrt{2}}m_Z c_{hZZ}$$
$$F_{1,3}^{BW11}= F_{p\cdot\gamma 1}^{VFFV}\left(m_f^2,m_W^2,m_{f'}^2,m_h^2,t,u;Z,f,f',Z\right)$$
$$F_{1,4}^{BW11}= F_{p\cdot\gamma 2}^{VFFV}\left(m_f^2,m_W^2,m_{f'}^2,m_h^2,t,u;Z,f,f',Z\right)$$
$$F_{\gamma}^{BW11}= F_{\gamma}^{VFFV}\left(m_f^2,m_W^2,m_{f'}^2,m_h^2,t,u;Z,f,f',Z\right)$$

$$C^{BW12} = -2a_f\left(v_f + a_f\right)\frac{g^2 g_Z^2}{\sqrt{2}} s_\beta c_\beta m_W c_{hH^\pm W^\mp}$$
$$F_{1,3}^{BW12} = F_{p\cdot\gamma}^{VFVS}\left(m_{f'}^2, m_f^2, m_W^2, m_h^2, s, t; W, f, Z, H^\pm\right)$$
$$F_{1,4}^{BW12} = 0$$
$$F_\gamma^{BW12} = F_\gamma^{VFVS}\left(m_{f'}^2, m_f^2, m_W^2, m_h^2, s, t; W, f, Z, H^\pm\right)$$

$\Phi = \frac{\varphi_1 + i\varphi_2}{\sqrt{2}}$, coupled to a gauge field A_μ

$$\mathcal{L} = \frac{1}{4}F_{\alpha\beta}F^{\alpha\beta} + \frac{1}{2\xi}(\nabla^\alpha A_\alpha)^2 + \nabla_\alpha \Phi \nabla^\alpha \Phi^* + m^2 \Phi\Phi^* - ieA_\alpha(\Phi\nabla^\alpha\Phi^* - \Phi^*\nabla^\alpha\Phi) + e^2 A_\alpha A^\alpha \Phi\Phi^*$$

$$m^2 = \Delta_1(\Delta_1 - d) = \Delta_2(\Delta_2 - d) \Longrightarrow \nu_1 = \pm\nu_2$$

$$S = \int_{AdS} \mathrm{d}^{d+1}x\sqrt{g}\left(\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2\xi}\left(\nabla_\mu A^\mu\right)^2\right)$$

$$\left[-\nabla_1^2 - d + \frac{1}{\frac{d-1}{2}}\left(1 - \frac{1}{\xi}\right)(W_1\cdot\nabla_1)(K_1\cdot\nabla_1)\right]\Pi(X_1, X_2; W_1, W_2) = (W_1\cdot W_2)\delta^{d+1}(X_1, X_2)$$

$$\Pi(X_1, X_2; W_1, W_2) = F_0(u)(W_1\cdot W_2) + F_1(u)(X_1\cdot W_2)(X_2\cdot W_1)$$

$$u := \frac{(X_1 - X_2)^2}{2} = -(1 + X_1\cdot X_2)$$

$$\Pi_{d-1}^{\mathcal{D}}(X_1, X_2; W_1, W_2) = \int_{-\infty}^{+\infty}\mathrm{d}\lambda \frac{1}{\lambda^2 + \left(\frac{d}{2} - 1\right)^2}\Omega_\lambda^{(1)}(X_1, X_2; W_1, W_2)$$
$$+ \int_{-\infty}^{+\infty}\mathrm{d}\lambda \frac{\xi}{\left(\lambda^2 + \frac{d^2}{4}\right)^2}(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega_\lambda^{(0)}(X_1, X_2).$$

$$M^2 = (\Delta - 1)(\Delta - d + 1)$$

$$\Pi_{d-1}^{\mathcal{D}}(X_1, X_2; W_1, W_2) = F_0^{\mathcal{D}}(u)(W_1\cdot W_2) + F_1^{\mathcal{D}}(u)(X_1\cdot W_2)(X_2\cdot W_1)$$

$$\Pi_{d-1,\mu\nu}^{\mathcal{D}}(x_1, x_2) = -F_0^{\mathcal{D}}(u)\frac{\partial^2 u}{\partial x_1^\mu \partial x_2^\nu} + F_1^{\mathcal{D}}(u)\frac{\partial u}{\partial x_1^\mu}\frac{\partial u}{\partial x_2^\nu}$$

$\Pi_{d-1,\mu\nu}^{\mathcal{D}}(x_1, x_2)$ to be:

$$F_0^{\mathcal{D}}(u) \underset{u\to\infty}{\sim} \frac{\Gamma\left(\frac{d+1}{2}\right)}{2\pi^{\frac{d+1}{2}}}\frac{1}{d-2}\frac{1}{u^{d-1}}$$
$$F_1^{\mathcal{D}}(u) \underset{u\to\infty}{\sim} \frac{1}{u}F_0^{\mathcal{D}}(u)$$

$$\int_{\lambda=\pm i\left(\frac{d}{2}-1\right)}^{\circlearrowleft} \mathrm{d}\lambda \frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\Omega_\lambda^{(1)}(X_1,X_2;W_1,W_2)$$
$$+\xi\int_{\lambda=\pm i\frac{d}{2}}^{\circlearrowleft} \mathrm{d}\lambda \frac{1}{\left(\lambda^2+\frac{d^2}{4}\right)^2}(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega_\lambda^{(0)}(X_1,X_2),$$

$$\square|_{\xi=\frac{d}{d-2}\frac{1}{u^{d-1}}} = F_0^{\mathrm{Hom}}(u)(W_1\cdot W_2)+F_1^{\mathrm{Hom}}(u)(X_1\cdot W_2)(X_2\cdot W_1),$$

$$F_0^{\mathrm{Hom}}(u) \underset{u\to\infty}{\sim} -\frac{\Gamma\left(\frac{d+1}{2}\right)}{2\pi^{\frac{d+1}{2}}(d-2)}\frac{1}{u^{d-1}}-\frac{1}{4\pi^{\frac{d}{2}}\Gamma\left(2-\frac{d}{2}\right)}\left[\left(1-\frac{d-2}{d}\xi\right)\log\left(\frac{u}{2}\right)+C\right]\frac{1}{u}$$
$$C:=-\frac{d-1}{d-2}+\left(1-\frac{d-2}{d}\xi\right)\left(\log(4)+\gamma_E+\psi^{(0)}\left(-\frac{d}{2}\right)-\frac{2}{d}\right)$$

$$F_1^{\mathrm{Hom}}(u) \underset{u\to\infty}{\sim} -\frac{\Gamma\left(\frac{d+1}{2}\right)}{2\pi^{\frac{d+1}{2}}(d-2)}\frac{1}{u^{d}}-\frac{1}{4\pi^{\frac{d}{2}}\Gamma\left(2-\frac{d}{2}\right)}\left[\left(1-\frac{d-2}{d}\xi\right)\log\left(\frac{u}{2}\right)+C+\frac{d-2}{d}\xi\right]\frac{1}{u^2}$$

$$\Pi_1^{\mathcal{N}}(X_1,X_2;W_1,W_2)$$
$$=\int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\Omega_\lambda^{(1)}(X_1,X_2;W_1,W_2)$$
$$+\int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda\frac{\xi}{\left(\lambda^2+\frac{d^2}{4}\right)^2}(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega_\lambda^{(0)}(X_1,X_2)$$

$$ds^2=\frac{1}{z^2}(\mathrm{d}\vec{x}^2+\mathrm{d}z^2)$$

$$\langle E_j(X_1)E_k(X_2)\rangle = \langle F_{zj}(X_1)F_{zk}(X_2)\rangle$$
$$\langle B_{i_1\dots i_{d-2}}(X_1)B_{j_1\dots j_{d-2}}(X_2)\rangle = \varepsilon^{ab}_{i_1\dots i_{d-2}}\varepsilon^{cd}_{j_1\dots j_{d-2}}\langle F_{ab}(X_1)F_{cd}(X_2)\rangle.$$

	Dirichlet	Neumann
$\langle E_j(X_1)E_k(X_2)\rangle$	$\mathcal{O}(z_1^{d-3})$	$\mathcal{O}(z_1)$
$\langle B_{i_1\dots i_{d-2}}(X_1)B_{j_1\dots j_{d-2}}(X_2)\rangle$	$\mathcal{O}(z_1^{d-2})$	$\mathcal{O}(z_1^0)$

$$\langle E_j(X_1)E_k(X_2)\rangle \underset{z_1\to 0}{\gg} \langle B_j(X_1)B_k(X_2)\rangle,$$

$$A_i \underset{z\to 0}{\sim} z^{d-2}j_i(\vec{x}), A_z \underset{z\to 0}{\sim} \mathcal{O}(z^{d-1})$$

$$\langle E_j(X_1)E_k(X_2)\rangle \underset{z_1\to 0}{\ll} \langle B_j(X_1)B_k(X_2)\rangle.$$

$$\langle \varphi_i(P_1)\varphi_j(P_2)\varphi_k(P_3)\varphi_l(P_4)\rangle_{\text{connected}} := \mathcal{A}_{\mathcal{D}}^{ijkl}$$

$\varphi_i(P_1) \leftrightarrow \varphi_j(P_2)$ or $\varphi_k(P_3) \leftrightarrow \varphi_l(P_4)$

$$T_{ij}^{A}(P_1,P_2,X_m) = \Pi^{(0)}_{\frac{d}{2}+i\nu_i}(P_1,X_m)\nabla_m^A\Pi^{(0)}_{\frac{d}{2}+i\nu_j}(P_2,X_m) - \begin{pmatrix} i\leftrightarrow j \\ P_1\leftrightarrow P_2\end{pmatrix},$$

$$\mathcal{A}_{\mathcal{D}}^{ijkl,\perp} = \frac{(-ie)^2}{\left(\frac{d-1}{2}\right)^2}\int_{AdS} \mathrm{d}^{d+1}X_1 \int_{AdS} \mathrm{d}^{d+1}X_2 K_{1,A}T_{ij}^{A}(P_1,P_2,X_1)$$
$$\Pi^{(1),\perp}_{d-1}(X_1,X_2;W_1,K_2)W_{2,B}T_{kl}^{B}(P_3,P_4,X_2)$$

$$W_1^A W_2^B \Pi^{(1),\perp}_{d-1,AB}(X_1,X_2) = \int_{\mathbb{R}^d}\mathrm{d}\lambda \frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\Omega^{(1)}_{\lambda}(X_1,X_2;W_1,W_2)$$

$$\mathcal{A}_{\mathcal{D}}^{ijkl,\perp} = \frac{1}{\pi\left(\frac{d}{2}-1\right)}\frac{(-ie)^2}{\left(\frac{d-1}{2}\right)^2}\int_{\mathbb{R}^d}\mathrm{d}\lambda \frac{\lambda^2\sqrt{\mathcal{C}_{\lambda}^{(1)}\mathcal{C}_{-\lambda}^{(1)}}}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\int_{\partial AdS}\mathrm{d}^dP_5$$
$$\int_{AdS}\mathrm{d}^{d+1}X_1 K_{1,A}T_{ij}^{A}(P_1,P_2,X_1)\Pi^{(1)}_{\frac{d}{2}+i\lambda}(X_1,P_5;W_1,D_Z)$$
$$\int_{AdS}\mathrm{d}^{d+1}X_2 \Pi^{(1)}_{\frac{d}{2}-i\lambda}(P_5,X_2;Z,K_2)W_{2,B}T_{kl}^{B}(P_3,P_4,X_2)$$

$$\frac{1}{J!\left(\frac{d-1}{2}\right)_J}\int_{AdS}\mathrm{d}^{d+1}X_1\Pi^{(0)}_{\frac{d}{2}+i\nu_i}(P_1,X_1)\left(K_{1,A}\nabla_1^A\right)^J\Pi^{(0)}_{\frac{d}{2}+i\nu_j}(P_2,X_1)\Pi^{(J)}_{\frac{d}{2}+i\lambda}(X_1,P_5;W_1,D_Z)$$
$$= \underbrace{\frac{\pi^{\frac{d}{2}}\Gamma\left(\frac{d}{4}+\frac{i\nu_i+i\nu_j+i\lambda+J}{2}\right)\Gamma\left(\frac{d}{4}+\frac{i\nu_i+i\nu_j-i\lambda+J}{2}\right)\Gamma\left(\frac{d}{4}+\frac{i\nu_i-i\nu_j+i\lambda+J}{2}\right)\Gamma\left(\frac{d}{4}+\frac{i\nu_j-i\nu_i+i\lambda+J}{2}\right)}{2^{1-J}\Gamma\left(\frac{d}{2}+i\nu_i\right)\Gamma\left(\frac{d}{2}+i\nu_j\right)\Gamma\left(\frac{d}{2}+i\lambda+J\right)}}_{b_{\text{Bulk}}\left(\nu_i,\nu_j,\lambda,J\right)}$$

$$\Pi^{(J)}_{\frac{d}{2}+i\nu}(X,P;W,Z) = \sqrt{\mathcal{C}_{\nu}^{(J)}}\frac{((-2P\cdot X)(W\cdot Z)+2(W\cdot P)(Z\cdot X))^J}{(-2P\cdot X)^{\frac{d}{2}+i\nu+J}}$$

$$\mathcal{C}_{\nu}^{(J)} = \frac{\left(J+\frac{d}{2}+i\nu-1\right)}{\left(\frac{d}{2}+i\nu-1\right)}\frac{\Gamma\left(\frac{d}{2}+i\nu\right)}{2\pi^{\frac{d}{2}}\Gamma(1+i\nu)}$$

$$\iiint \left\langle \mathcal{O}_{\nu_i}(P_1)\mathcal{O}_{\nu_j}(P_2)\mathcal{O}_{\lambda}^{(J)}(P_5,Z)\right\rangle = \frac{[(Z\cdot P_1)P_{25}-(Z\cdot P_2)P_{15}]^J}{P_{12}^{\frac{d}{4}+\frac{i\nu_i+i\nu_j-i\lambda+J}{2}}P_{25}^{\frac{d}{4}+\frac{i\nu_j-i\nu_i+i\lambda+J}{2}}P_{15}^{\frac{d}{4}+\frac{i\nu_i-i\nu_j+i\lambda+J}{2}}}$$

$$\mathcal{A}_{\mathcal{D}}^{ijkl,\perp}=\frac{(-ie)^2}{\pi}\frac{\sqrt{\mathcal{C}_{\nu_1}^{(0)}\mathcal{C}_{\nu_2}^{(0)}\mathcal{C}_{\nu_3}^{(0)}\mathcal{C}_{\nu_4}^{(0)}}}{\left(\frac{d}{2}-1\right)}\int_{\mathbb{R}^d}\mathrm{d}\lambda\frac{\lambda^2\mathcal{C}_{\lambda}^{(1)}\mathcal{C}_{-\lambda}^{(1)}}{\lambda^2+\left(\frac{d}{2}-1\right)^2}b_{\mathrm{Bulk}}(\nu_i,\nu_j,\lambda,1)b_{\mathrm{Bulk}}(\nu_k,\nu_l,-\lambda,1)$$
$$\int_{\partial AdS}\mathrm{d}^dP_5\left[\left(\left\langle\mathcal{O}_{\nu_i}(P_1)\mathcal{O}_{\nu_j}(P_2)\mathcal{O}_{\lambda}^{(1)}(P_5;D_Z)\right\rangle_{\blacksquare}-\left\langle\mathcal{O}_{\nu_j}(P_2)\mathcal{O}_{\nu_i}(P_1)\mathcal{O}_{\lambda}^{(1)}(P_5;D_Z)\right\rangle_{\blacksquare}\right)\right.$$
$$\left.\left(\left\langle\mathcal{O}_{\nu_k}(P_3)\mathcal{O}_{\nu_l}(P_4)\mathcal{O}_{-\lambda}^{(1)}(P_5;Z)\right\rangle_{\blacksquare}-\left\langle\mathcal{O}_{\nu_l}(P_4)\mathcal{O}_{\nu_k}(P_3)\mathcal{O}_{-\lambda}^{(1)}(P_5;Z)\right\rangle_{\blacksquare}\right)\right]$$

$$\left\langle\mathcal{O}_{\nu_i}(P_i)\mathcal{O}_{\nu_j}(P_j)\mathcal{O}_{\lambda}^{(1)}(P_5;D_Z)\right\rangle_{\blacksquare}=(-1)\left\langle\mathcal{O}_{\nu_j}(P_j)\mathcal{O}_{\nu_i}(P_i)\mathcal{O}_{\lambda}^{(1)}(P_5;D_Z)\right\rangle_{\blacksquare}$$

$$\mathcal{A}_{\mathcal{D}}^{ijkl,\perp}=\frac{4(-ie)^2}{\pi}\frac{\sqrt{\mathcal{C}_{\nu_i}^{(0)}\mathcal{C}_{\nu_j}^{(0)}\mathcal{C}_{\nu_k}^{(0)}\mathcal{C}_{\nu_l}^{(0)}}}{\left(\frac{d}{2}-1\right)}\int_{\mathbb{R}^d}\mathrm{d}\lambda\frac{\lambda^2\mathcal{C}_{\lambda}^{(1)}\mathcal{C}_{-\lambda}^{(1)}}{\lambda^2+\left(\frac{d}{2}-1\right)^2}b_{\mathrm{Bulk}}(\nu_i,\nu_j,\lambda,1)b_{\mathrm{Bulk}}(\nu_k,\nu_l,-\lambda,1)$$
$$\int_{\partial AdS}\mathrm{d}^dP_5\left\langle\mathcal{O}_{\nu_i}(P_1)\mathcal{O}_{\nu_j}(P_2)\mathcal{O}_{\lambda}^{(1)}(P_5;D_Z)\right\rangle_{\blacksquare}\left\langle\mathcal{O}_{\nu_k}(P_3)\mathcal{O}_{\nu_l}(P_4)\mathcal{O}_{-\lambda}^{(1)}(P_5;Z)\right\rangle_{\blacksquare}$$

$$\Psi_{\lambda,J}^{\{\Delta_i\}}(\{P_i\}):=\frac{1}{P_{12}^{\frac{\Delta_i+\Delta_j}{2}}P_{34}^{\frac{\Delta_k+\Delta_l}{2}}}\left(\frac{P_{24}}{P_{14}}\right)^{\frac{\Delta_{ij}}{2}}\left(\frac{P_{14}}{P_{13}}\right)^{\frac{\Delta_{kl}}{2}}\mathcal{F}_{\lambda,J}^{\{\Delta_i\}}(u,v)$$
$$=\frac{1}{\left(\frac{d}{2}-1\right)_J}\int_{\partial AdS}\mathrm{d}^dP_5\left\langle\mathcal{O}_{\nu_i}(P_1)\mathcal{O}_{\nu_j}(P_2)\mathcal{O}_{\frac{d}{2}+i\lambda}^{(J)}(P_5,D_Z)\right\rangle_{\blacksquare}\left\langle\mathcal{O}_{\frac{d}{2}-i\lambda}^{(J)}(P_5,Z)\mathcal{O}_{\nu_k}(P_3)\mathcal{O}_{\nu_l}(P_4)\right\rangle_{\blacksquare}$$

$$\mathcal{A}_{\mathcal{D}}^{ijkl,\perp}=\frac{(-ie)^2}{\pi}\sqrt{\prod_{a\in\{ijkl\}}\mathcal{C}_{\nu_a}^{(0)}}\left(\frac{1}{P_{12}}\right)^{\frac{d}{2}+i\frac{\nu_i+\nu_j}{2}}\left(\frac{1}{P_{34}}\right)^{\frac{d}{2}+i\frac{\nu_k+\nu_l}{2}}\left(\frac{P_{24}}{P_{14}}\right)^{i\frac{\nu_i-\nu_j}{2}}\left(\frac{P_{14}}{P_{13}}\right)^{i\frac{\nu_k-\nu_l}{2}}$$
$$\int_{\mathbb{R}^d}\mathrm{d}\lambda\frac{4\lambda^2}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\mathcal{C}_{\lambda}^{(1)}\mathcal{C}_{-\lambda}^{(1)}b_{\mathrm{Bulk}}\left(\nu_i,\nu_j,\lambda,1\right)b_{\mathrm{Bulk}}\left(\nu_k,\nu_l,-\lambda,1\right)\mathcal{F}_{\lambda}^{(1),\{\Delta_i\}}(u,v)$$

$$P_{ij}=-2P_i\cdot P_j,u=\frac{P_{12}P_{34}}{P_{13}P_{24}},v=\frac{P_{14}P_{23}}{P_{13}P_{24}}$$

$$\mathcal{F}_{\lambda,J}^{\{\Delta_i\}}(u,v)=\mathcal{K}_{d-\Delta_\lambda}^{(J),\{\Delta_k,\Delta_l\}}\hat{\mathcal{G}}_{\Delta_\lambda}^{(J)}(u,v)+\mathcal{K}_{\Delta_\lambda}^{(J),\{\Delta_i,\Delta_j\}}\hat{\mathcal{G}}_{d-\Delta_\lambda}^{(J)}(u,v)$$

$$\mathcal{K}_{\Delta_\lambda}^{(J),\{\Delta_1,\Delta_2\}}=\frac{\pi^{\frac{d}{2}}}{(-2)^J}\frac{\Gamma\left(\Delta_\lambda-\frac{d}{2}\right)\Gamma(\Delta_\lambda+J-1)\Gamma\left(\frac{d-\Delta_\lambda+\Delta_1-\Delta_2+J}{2}\right)\Gamma\left(\frac{d-\Delta_\lambda+\Delta_2-\Delta_1+J}{2}\right)}{\Gamma(\Delta_\lambda-1)\Gamma(d-\Delta_\lambda+J)\Gamma\left(\frac{\Delta_\lambda+\Delta_1-\Delta_2+J}{2}\right)\Gamma\left(\frac{\Delta_\lambda+\Delta_2-\Delta_1+J}{2}\right)}$$

$\mathcal{K}_{d-\Delta_\lambda,\{\Delta_i\}}^{(J)}\underset{\mathcal{F}_{\lambda,J}^{\{\Delta_i\}}(u,v)}{\triangle}\hat{\mathcal{G}}_{\Delta_\lambda}^{(J)}(u,v)$only depend on $\Delta_{ij}\equiv\Delta_i-\Delta_j$ and $\Delta_{kl}\equiv\Delta_k-\Delta_l$

$$\mathcal{A}_{\mathcal{D}}^{ijkl,\perp}=(-ie)^2\left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu}\int_{\mathbb{R}^d}\mathrm{d}\lambda\rho_{\mathcal{D}}^{J=1}\left(\lambda;\left\{\frac{d}{2}+i\nu\right\}\right)\mathcal{F}_{\lambda,1}^{\left\{\frac{d}{2}+i\nu\right\}}(u,v)$$

$\left\{\frac{d}{2}+i\nu\right\}$ denotes the four equal external scaling dimensions $\Delta_i=\frac{d}{2}+i\nu$

$$\rho_{\mathcal{D}}^{J=1}\left(\lambda;\left\{\frac{d}{2}+i\nu\right\}\right)=\Pi_{d-1}^{(1),\perp}(\lambda)\mathcal{Q}^{J=1}\left(\lambda,\left\{\frac{d}{2}+i\nu\right\}\right)$$

$$\Pi_{d-1}^{(1),\perp}=\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}$$

$$\mathcal{Q}^{J=1}\left(\lambda,\left\{\frac{d}{2}+i\nu\right\}\right)=\frac{\lambda\sinh(\pi\lambda)}{4\pi^{d+2}}\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\frac{\Gamma\left(\frac{d+2\pm 2i\lambda}{4}\right)^2\Gamma\left(\frac{d+2+4i\nu\pm 2i\lambda}{4}\right)^2}{\Gamma(1+i\nu)^2\Gamma\left(\frac{d}{2}+i\nu\right)^2}$$

$\rho_{\mathcal{D}}^{J=1}\left(\lambda;\left\{\frac{d}{2}+i\nu\right\}\right)$ is the product of the factor $\Pi_{d-1}^{(1),\perp}(\lambda)$

$$\rho_{\mathcal{D}}^{J=1}\left(\lambda;\left\{\frac{d}{2}+i\nu\right\}\right)\mathcal{K}_{d-\Delta_\lambda}^{(J=1)}$$

$$=-\frac{\lambda\sinh(\pi\lambda)}{2^{5+\frac{d}{2}+i\lambda}\pi^{\frac{d+3}{2}}}\frac{\Gamma\left(\frac{d-2+2i\lambda}{4}\right)^2}{\Gamma(1+i\nu)^2\Gamma\left(\frac{d}{2}+i\nu\right)^2}\frac{\Gamma(-i\lambda)\Gamma\left(\frac{d+2+2i\lambda}{4}\right)\Gamma\left(\frac{d+2+4i\nu\pm 2i\lambda}{4}\right)^2}{\left(\frac{d}{2}-1-i\lambda\right)\Gamma\left(\frac{d+4+2i\lambda}{4}\right)}$$

$$\Pi_{d-1}^{(1),\perp}\lambda=-i\left(\frac{d}{2}-1\right)$$

$$\rho_{\mathcal{D}}^{J=1}\left(\lambda,\left\{\frac{d}{2}+i\nu\right\}\right)\mathcal{K}_{d-\Delta_\lambda}^{(J=1)}\underset{\lambda\to -i\left(\frac{d}{2}-1\right)}{\sim}-\frac{\Gamma\left(\frac{d}{2}-1\right)^2}{2^{d+5}\pi^{\frac{d+1}{2}}\Gamma\left(\frac{d+1}{2}\right)}\frac{i(d-2)}{\lambda+i\left(\frac{d}{2}-1\right)}+\mathcal{O}(1)$$

$\Delta_i=\Delta_k=\frac{d}{2}+i\nu$ and $\Delta_j=\Delta_l=\frac{d}{2}-i\nu$, shortened as $\left\{\frac{d}{2}\pm i\nu\right\}$, the 4 -point function evaluates to

$$\mathcal{A}_{\mathcal{D}}^{ijkl,\perp}=(-ie)^2\left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}}\left(\frac{P_{24}}{P_{14}}\frac{P_{14}}{P_{13}}\right)^{i\nu}\int_{\mathbb{R}}\mathrm{d}\lambda\rho_{\mathcal{D}}^{J=1}\left(\lambda;\left\{\frac{d}{2}\pm i\nu\right\}\right)\mathcal{F}_{\lambda,1}^{\left\{\frac{d}{2}\pm i\nu\right\}}(u,v)$$

$$\rho_{\mathcal{D}}^{J=1}\left(\lambda;\left\{\frac{d}{2}\pm i\nu\right\}\right)=\Pi_{d-1}^{(1),\perp}(\lambda)\mathcal{Q}^{J=1}\left(\lambda,\left\{\frac{d}{2}\pm i\nu\right\}\right)$$

$$\Pi_{d-1}^{(1),\perp}=\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}$$

$$\mathcal{Q}^{J=1}\left(\lambda,\left\{\frac{d}{2}\pm i\nu\right\}\right)=\frac{\lambda\sinh(\pi\lambda)}{4\pi^{d+3}}\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\frac{\sinh(\pi\nu)}{\nu}\frac{\Gamma\left(\frac{d+2\pm 2i\lambda}{4}\right)^2\Gamma\left(\frac{d+2\pm 4i\nu\pm 2i\lambda}{4}\right)}{\Gamma\left(\frac{d}{2}\pm i\nu\right)}$$

$$\rho_{\mathcal{D}}^{J=1}\left(\lambda;\left\{\frac{d}{2}\pm i\nu\right\}\right)\mathcal{K}_{d-\Delta_\lambda}^{(J=1)}$$

$$=-\frac{\lambda\sinh(\pi\lambda)}{2^5\pi^{\frac{d}{2}+3}}\frac{\sinh(\pi\nu)}{\nu}\frac{\Gamma\left(\frac{d-2+2i\lambda}{4}\right)^2}{\Gamma\left(\frac{d}{2}\pm i\nu\right)}\frac{\Gamma(-i\lambda)\Gamma\left(\frac{d+2-2i\lambda}{4}\right)^2\Gamma\left(\frac{d+2\pm 4i\nu+2i\lambda}{4}\right)^2}{\left(\frac{d}{2}-1-i\lambda\right)\Gamma\left(\frac{d}{2}+i\lambda+1\right)}$$

$$\underset{\lambda\to -i\left(\frac{d}{2}-1\right)}{\sim}-\frac{\Gamma\left(\frac{d}{2}-1\right)\sinh(\pi\nu)\Gamma\left(\frac{d}{2}\pm i\nu\right)}{2^5\pi^{\frac{d+1}{2}}\nu\Gamma(d)}\frac{i}{\lambda+i\left(\frac{d}{2}-1\right)}+\mathcal{O}(1)$$

$$\mathcal{A}_{\mathcal{D}}^{ijkl,\|}=(-ie)^2\int_{AdS}\mathrm{d}^{d+1}X_1\,\mathrm{d}^{d+1}X_2T_{ij}^{A}(P_1,P_2,X_1)\Pi_{d-1,AB}^{(1),\|}(X_1,X_2)T_{kl}^{B}(P_3,P_4,X_2)$$

$$\Pi_{d-1,AB}^{(1),\|}(X_1,X_2)=\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{\xi}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\nabla_{1,A}\nabla_{2,B}\Omega_{\lambda}^{(0)}(X_1,X_2)$$

$$\mathcal{A}_{\mathcal{D}}^{ijkl,\|}=(-ie)^2\frac{\xi}{\pi}\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{\lambda^2\sqrt{\mathcal{C}_{\lambda}^{(0)}\mathcal{C}_{-\lambda}^{(0)}}}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\int_{\partial AdS}\mathrm{d}^dP_5\int_{AdS}\mathrm{d}^{d+1}X_1T_{ij}^{A}(P_1,P_2,X_1)\nabla_{1,A}\Pi_{\frac{d}{2}+i\lambda}^{(0)}(X_1,P_5)$$

$$\int_{AdS}\mathrm{d}^{d+1}X_2T_{kl}^{B}(P_3,P_4,X_2)\nabla_{2,B}\Pi_{\frac{d}{2}-i\lambda}^{(0)}(X_2,P_5)$$

$$\mathcal{C}_{\lambda}^{(J)}=\frac{\left(J+\frac{d}{2}+i\lambda-1\right)}{\left(\frac{d}{2}+i\lambda-1\right)}\frac{\Gamma\left(\frac{d}{2}+i\lambda\right)}{2\pi^{\frac{d}{2}}\Gamma(1+i\lambda)}$$

$$\mathcal{A}_{\mathcal{D}}^{ijkl,\|}=(-ie)^2\frac{\xi}{\pi}\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{\lambda^2\sqrt{\mathcal{C}_{\lambda}^{(0)}\mathcal{C}_{-\lambda}^{(0)}}}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\int_{\partial AdS}\mathrm{d}^dP_5\int\frac{\mathrm{d}^d\vec{x}_{\tilde{1}}}{z_1^{d+1}}\int\frac{\mathrm{d}^d\vec{x}_{\tilde{2}}}{z_2^{d+1}}$$

$$T_{ij}^{z}(P_1,P_2,X_1)\Pi_{\frac{d}{2}+i\lambda}^{(0)}(X_1,P_5)T_{kl}^{z}(P_3,P_4,X_2)\Pi_{\frac{d}{2}-i\lambda}^{(0)}(X_2,P_5),$$

$$\Pi_{\frac{d}{2}+i\nu}^{(0)}(X_1,P_2)=\sqrt{\mathcal{C}_{\nu}^{(0)}}\left(\frac{z_1}{z_1^2+\vec{x}_{12}^2}\right)^{\frac{d}{2}+i\nu},\mathcal{C}_{\nu}^{(0)}=\frac{\Gamma\left(\frac{d}{2}+i\nu\right)}{2\pi^{\frac{d}{2}}\Gamma(1+i\nu)}$$

$$\lim_{z\to 0}\frac{z^{d+2\alpha}}{(z^2+\vec{x}^2)^{d+\alpha}}=\pi^{\frac{d}{2}}\frac{\Gamma\left(\frac{d}{2}+\alpha\right)}{\Gamma(d+\alpha)}\delta^d(\vec{x})$$

$$\Pi_{\frac{d}{2}+i\nu}^{(0)}(X_1,P_1)\underset{z\to 0}{\sim}\sqrt{\mathcal{C}_{\nu}^{(0)}}\left[\tilde{\kappa}_{\nu}\delta^d(\vec{x}_{1\tilde{1}})z^{\frac{d}{2}-i\nu}+\left(\frac{z}{\vec{x}_{1\tilde{1}}^2}\right)^{\frac{d}{2}+i\nu}\right]+\mathcal{O}\left(z^{\frac{d}{2}+i\nu+2}\right)$$

$$\nabla^z\Pi_{\frac{d}{2}+i\nu}^{(0)}(X_1,P_1)\underset{z\to 0}{\sim}z^2\sqrt{\mathcal{C}_{\nu}^{(0)}}\left(\frac{d}{2}+i\nu\right)\left[\tilde{\kappa}_{\nu}^{\frac{d}{2}-i\nu}\delta^d(\vec{x}_{1\tilde{1}})z^{\frac{d}{2}-i\nu-1}+\frac{z^{\frac{d}{2}+i\nu-1}}{\left(\vec{x}_{1\tilde{1}}^2\right)^{\frac{d}{2}+i\nu}}+\mathcal{O}\left(z^{\frac{d}{2}+i\nu+1}\right)\right]$$

$$T_{12}^{z}(P_1,P_2,X_1) \underset{z\to 0}{\sim} z^{d+1}\left(\frac{\delta^d(\vec{x}_{1\tilde{1}})}{(\vec{x}_{2\tilde{1}}^2)^{\frac{d}{2}+i\nu}}-\frac{\delta^d(\vec{x}_{2\tilde{1}})}{(\vec{x}_{1\tilde{1}}^2)^{\frac{d}{2}+i\nu}}+\mathcal{O}(z^2)+\mathcal{O}\left(z^{2(1+i\nu)}\right)\right)$$

$$\tilde{\kappa}_\nu=\pi^{\frac{d}{2}}\frac{\Gamma(i\nu)}{\Gamma\left(\frac{d}{2}+i\nu\right)}$$


$$\left\langle\Phi(P_1)\Phi^\dagger(P_2)\Phi^\dagger(P_3)\Phi(P_4)\right\rangle=\frac{1}{4}\epsilon_{ij}\epsilon_{kl}\mathcal{A}_{\mathcal{N}}^{ijkl}$$

$$\begin{aligned}\Pi_1^{\mathcal{N}}(X_1,X_2;W_1,W_2)=\Pi_{d-1}^{\mathcal{D}}(X_1,X_2;W_1,W_2)\\+\int_{\lambda=\pm i\left(\frac{d}{2}-1\right)}^{\circlearrowleft}\mathrm{d}\lambda\,\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\Omega_\lambda^{(1)}(X_1,X_2;W_1,W_2)\\+\xi\int_{\lambda=\pm i\frac{d}{2}}^{\circlearrowleft}\mathrm{d}\lambda\,\frac{1}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\left(W_1\cdot\nabla_1\right)\left(W_2\cdot\nabla_2\right)\Omega_\lambda^{(0)}(X_1,X_2),\end{aligned}$$

$$\begin{aligned}\mathcal{A}_{\mathcal{N}}^{ijkl,\|}-\mathcal{A}_{\mathcal{D}}^{ijkl,\|}=(-ie)^2\frac{\xi}{\pi}\int_{\lambda=\pm i\frac{d}{2}}^{\circlearrowleft}\mathrm{d}\lambda\,\frac{\lambda^2}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\sqrt{\mathcal{C}_\lambda^{(0)}\mathcal{C}_{-\lambda}^{(0)}}\int_{\partial AdS}\mathrm{d}^dP_5\\\int_{AdS}\mathrm{d}^{d+1}X_1\,T_{ij}^A\left(P_1,P_2,X_1\right)\nabla_{1,A}\Pi_{\frac{d}{2}+i\lambda}^{(0)}(X_1,P_5)\\\int_{AdS}\mathrm{d}^{d+1}X_2\,T_{kl}^B\left(P_3,P_4,X_2\right)\nabla_{2,B}\Pi_{\frac{d}{2}-i\lambda}^{(0)}(X_2,P_5)\ .\end{aligned}$$

$$\begin{aligned}\mathcal{A}_{\mathcal{N}}^{ijkl,\|}=(-ie)^2\frac{\xi}{\pi}\frac{(2i\nu)^2\pi^d\left(\mathcal{C}_\nu^{(0)}\right)^2\Gamma(i\nu)^2}{\Gamma\left(\frac{d}{2}+i\nu\right)^2}(2\pi i)\mathrm{Res}_{\lambda=i\frac{d}{2}}\left[\frac{\lambda^2}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\frac{\mathcal{C}_{-\lambda}^{(0)}\mathcal{C}_\lambda^{(0)}}{|\vec{x}_{12}|^{2\left(\frac{d}{2}+i\nu\right)}|\vec{x}_{34}|^{2\left(\frac{d}{2}+i\nu\right)}}\right.\\\left(z^{d+2i\lambda}\frac{\pi^{\frac{d}{2}}\Gamma(-i\lambda)}{\Gamma\left(\frac{d}{2}-i\lambda\right)}\left(\frac{1}{|\vec{x}_{13}|^{2\left(\frac{d}{2}+i\lambda\right)}}-\frac{1}{|\vec{x}_{14}|^{2\left(\frac{d}{2}+i\lambda\right)}}-\frac{1}{|\vec{x}_{23}|^{2\left(\frac{d}{2}+i\lambda\right)}}+\frac{1}{|\vec{x}_{24}|^{2\left(\frac{d}{2}+i\lambda\right)}}\right)\right.\\\left.\left.+z^{d-2i\lambda}\frac{\pi^{\frac{d}{2}}\Gamma(i\lambda)}{\Gamma\left(\frac{d}{2}+i\lambda\right)}\left(\frac{1}{|\vec{x}_{13}|^{2\left(\frac{d}{2}-i\lambda\right)}}-\frac{1}{|\vec{x}_{14}|^{2\left(\frac{d}{2}-i\lambda\right)}}-\frac{1}{|\vec{x}_{23}|^{2\left(\frac{d}{2}-i\lambda\right)}}+\frac{1}{|\vec{x}_{24}|^{2\left(\frac{d}{2}-i\lambda\right)}}\right)\right)\right]\\\underset{d=3}{=}-(-ie)^2\frac{\xi}{6\pi^2}\frac{1}{|\vec{x}_{12}|^{2\left(\frac{3}{2}+i\nu\right)}|\vec{x}_{34}|^{2\left(\frac{3}{2}+i\nu\right)}}\left[\left(2\log(z)-\gamma_E-\psi^{(0)}\left(-\frac{1}{2}\right)\right)\log\left(\frac{|\vec{x}_{13}||\vec{x}_{24}|}{|\vec{x}_{14}||\vec{x}_{23}|}\right)\right.\\\left.-(\log^2|\vec{x}_{13}|-\log^2|\vec{x}_{14}|-\log^2|\vec{x}_{23}|+\log^2|\vec{x}_{24}|)\right]+\mathcal{O}(z^{2\alpha})\end{aligned}$$

$$\left\langle\Phi(P_1)e^{-ie\int_{P_1}^{P_2}dX_1^AA_A(X_1)}\Phi^\dagger(P_2)\Phi^\dagger(P_3)e^{ie\int_{P_3}^{P_4}dX_2^BA_B(X_2)}\Phi(P_4)\right\rangle$$



$$= \left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu}.$$

$$\mathcal{A}_{gg}^{\parallel} = (-ie)^2\xi\left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu}\int_{\mathbb{R}\oplus\circlearrowleft}\mathrm{d}\lambda\frac{1}{\left(\lambda^2+\frac{d^2}{4}\right)^2}$$

$$\int_{P_1}^{P_2}\mathrm{d}X_1^A\int_{P_3}^{P_4}\mathrm{d}X_2^B\nabla_{1,A}\nabla_{2,B}\Omega_\lambda^{(0)}(X_1,X_2)$$

$$= (-ie)^2\frac{\xi}{\pi}\left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu}\int_{\mathbb{R}\oplus\circlearrowleft}\mathrm{d}\lambda\frac{\lambda^2\sqrt{\mathcal{C}_{-\lambda}^{(0)}\mathcal{C}_\lambda^{(0)}}}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\int_{\mathbb{R}^d}\mathrm{d}^dP_5$$

$$\int_{P_1}^{P_2}\mathrm{d}X_1^A\nabla_{1,A}\Pi_{\frac{d}{2}+i\lambda}^{(0)}(X_1,P_5)\int_{P_3}^{P_4}\mathrm{d}X_2^B\nabla_{1,B}\Pi_{\frac{d}{2}-i\lambda}^{(0)}(X_2,P_5)$$

$$\mathcal{A}_{gg}^{\parallel} = (-ie)^2\frac{\xi}{\pi}\left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu}\int_{\mathbb{R}\oplus\circlearrowleft}\mathrm{d}\lambda\frac{\lambda^2}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\sqrt{\mathcal{C}_{-\lambda}^{(0)}\mathcal{C}_\lambda^{(0)}}\int_{\mathbb{R}^d}\mathrm{d}^dP_5$$

$$\left(\Pi_{\frac{d}{2}+i\lambda}^{(0)}(P_2,P_5)-\Pi_{\frac{d}{2}+i\lambda}^{(0)}(P_1,P_5)\right)\left(\Pi_{\frac{d}{2}-i\lambda}^{(0)}(P_4,P_5)-\Pi_{\frac{d}{2}-i\lambda}^{(0)}(P_3,P_5)\right)$$

$$\langle\varphi_1(X_1)\varphi_2(X_2)\dots\varphi_k(\vec{x}_k)\dots\varphi_n(X_n)\rangle = \lim_{z\to 0}\frac{z^{-\left(\frac{d}{2}+i\nu_k\right)}}{\sqrt{\mathcal{C}_{\nu_k}^{(0)}}}\langle\varphi_1(X_1)\varphi_2(X_2)\dots\varphi_k(z,\vec{x}_k)\dots\varphi_n(X_n)\rangle$$

$$\mathcal{A}_{gc}^{\parallel} = (-ie)^2\xi\left(\frac{1}{P_{12}}\right)^{\frac{d}{2}+i\nu}\int_{\mathbb{R}\oplus\circlearrowleft}\mathrm{d}\lambda\frac{1}{\left(\lambda^2+\frac{d^2}{4}\right)^2}$$

$$\int_{P_1}^{P_2}\mathrm{d}X_1^A\int_{AdS}\mathrm{d}^{d+1}X_2\nabla_{1,A}\nabla_{2,B}\Omega_\lambda^{(0)}(X_1,X_2)T_{34}^B(P_3,P_4,X_2)$$

$$\mathcal{A}_{gc}^{\parallel} = (-ie)^2\frac{\xi}{\pi}\left(\frac{1}{P_{12}}\right)^{\frac{d}{2}+i\nu}\int_{\mathbb{R}\oplus\circlearrowleft}\mathrm{d}\lambda\frac{\lambda^2}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\sqrt{\mathcal{C}_{-\lambda}^{(0)}\mathcal{C}_\lambda^{(0)}}\int_{\partial AdS}\mathrm{d}^dP_5$$

$$\int_{P_1}^{P_2}\mathrm{d}X_1^A\nabla_{1,A}\Pi_{\frac{d}{2}+i\lambda}^{(0)}(X_1,P_5)\int_{AdS}\mathrm{d}^{d+1}X_2\nabla_{2,B}\Pi_{\frac{d}{2}-i\lambda}^{(0)}(P_5,X_2)T_{34}^B(P_3,P_4,X_2)$$

$$\int_{\bar{P}_1}^{\bar{P}_2}\mathrm{d}X_1^A\nabla_{1,A}\Pi_{\frac{d}{2}+i\lambda}^{(0)}(X_1,P_5) = \Pi_{\frac{d}{2}+i\lambda}^{(0)}(\bar{P}_2,P_5)-\Pi_{\frac{d}{2}+i\lambda}^{(0)}(\bar{P}_1,P_5)$$

$$\int_{AdS} \mathrm{d}^{d+1}X_2 \nabla_{2,\beta} \Pi^{(0)}_{\frac{d}{2}-i\lambda}(P_5,X_2) T^{\beta}_{34}(P_3,P_4,X_2) = \int_{\partial AdS} \frac{\mathrm{d}^d \vec{x}_{\tilde{2}}}{z^{1+d}} \Pi^{(0)}_{\frac{d}{2}-i\lambda}(P_5,X_2) T^{z}_{34}(P_3,P_4,X_2)$$

$$\mathcal{A}^{\parallel}_{cc} + \mathcal{A}^{\parallel}_{gg} + \mathcal{A}^{\parallel}_{gc} + \mathcal{A}^{\parallel}_{cg} = \lim_{z\to 0}(-ie)^2 \frac{\xi}{\pi} \int_{\mathbb{R}\oplus\circlearrowleft\circlearrowright} \mathrm{d}\lambda \frac{\lambda^2}{\left(\lambda^2+\frac{d^2}{4}\right)^2} \sqrt{\mathcal{C}^{(0)}_{-\lambda}\mathcal{C}^{(0)}_{\lambda}} \int_{\partial AdS} \mathrm{d}^d P_5$$
$$\left(\int_{\partial AdS} \frac{\mathrm{d}^d \vec{x}_{\tilde{1}}}{z^{d+1}} \Pi^{(0)}_{\frac{d}{2}+i\lambda}(P_5,X_1) T^{z}_{12}(P_1,P_2,X_1)\Big|_{X_1=(z,\vec{x}_{\tilde{1}})} + \left(\frac{1}{P_{12}}\right)^{\frac{d}{2}+i\nu} \int_{\bar{P}_1}^{\bar{P}_2} \mathrm{d}X_1^A \nabla_{1,A} \Pi^{(0)}_{\frac{d}{2}+i\lambda}(X_1,P_5) \right)$$
$$\left(\int_{\partial AdS} \frac{\mathrm{d}^d \vec{x}_{\tilde{2}}}{z^{d+1}} \Pi^{(0)}_{\frac{d}{2}-i\lambda}(P_5,X_2) T^{z}_{34}(P_3,P_4,X_2)\Big|_{X_2=(z,\vec{x}_{\tilde{2}})} + \left(\frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu} \int_{\bar{P}_3}^{\bar{P}_4} \mathrm{d}X_2^A \nabla_{2,A} \Pi^{(0)}_{\frac{d}{2}-i\lambda}(X_2,P_5) \right)$$


$$\int_{\partial AdS} \frac{\mathrm{d}^d \vec{x}_{\tilde{1}}}{z^{d+1}} \Pi^{(0)}_{\frac{d}{2}+i\lambda}(P_5,X_1) T^{z}_{12}(P_1,P_2,X_1)\Big|_{X_1=(z,\vec{x}_{\tilde{1}})} + \left(\frac{1}{P_{12}}\right)^{\frac{d}{2}+i\nu} \int_{\bar{P}_1}^{\bar{P}_2} \mathrm{d}X_1^A \nabla_{1,A} \Pi^{(0)}_{\frac{d}{2}+i\lambda}(X_1,P_5)$$

$$\underset{z\to 0}{\sim} 0 \int_{\partial AdS} \mathrm{d}^d \vec{x}_{\tilde{1}} \Pi^{(0)}_{\frac{d}{2}+i\lambda}(P_5,X_1)\Big|_{X_1=(z,\vec{x}_{\tilde{1}})} \left(\frac{\delta^d(\vec{x}_{1\tilde{1}})}{\left(\vec{x}^2_{2\tilde{1}}\right)^{\frac{d}{2}+i\nu}} - \frac{\delta^d(\vec{x}_{2\tilde{1}})}{\left(\vec{x}^2_{1\tilde{1}}\right)^{\frac{d}{2}+i\nu}} + \mathcal{O}(z^2) + \mathcal{O}\left(z^{2(1+i\nu)}\right) \right)$$
$$+ \left(\frac{1}{P_{12}}\right)^{\frac{d}{2}+i\nu} \left(\Pi^{(0)}_{\frac{d}{2}+i\lambda}(\bar{P}_2,P_5) - \Pi^{(0)}_{\frac{d}{2}+i\lambda}(\bar{P}_1,P_5) \right)$$
$$= \int_{\partial AdS} \mathrm{d}^d \vec{x}_{\tilde{1}} \Pi^{(0)}_{\frac{d}{2}+i\lambda}(P_5,X_1)\Big|_{X_1=(z,\vec{x}_{\tilde{1}})} \left(\mathcal{O}(z^2) + \mathcal{O}\left(z^{2(1+i\nu)}\right)\right)$$

$$z^{4\alpha} \Pi^{(0)}_{\frac{d}{2}+i\lambda}(P_5,X_1) \Pi^{(0)}_{\frac{d}{2}-i\lambda}(P_5,X_2)\Big|_{X_a=(z,\vec{x}_{\tilde{a}})} \propto z^{4\alpha}\left(z^d + \cdots + z^{d\pm 2i\lambda} + \cdots\right)$$

$$\mathcal{A}_{\mathcal{N}}^{\perp} = \frac{4(-ie)^2}{\pi} \frac{\sqrt{\mathcal{C}_{\nu_1}^{(0)} \mathcal{C}_{\nu_2}^{(0)} \mathcal{C}_{\nu_3}^{(0)} \mathcal{C}_{\nu_4}^{(0)}}}{\left(\frac{d}{2}-1\right)}$$
$$\int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda \frac{\lambda^2 \mathcal{C}_{\lambda}^{(1)} \mathcal{C}_{-\lambda}^{(1)}}{\lambda^2 + \left(\frac{d}{2}-1\right)^2} b_{\mathrm{Bulk}}(\nu_1, \nu_2, \lambda, 1) b_{\mathrm{Bulk}}(\nu_3, \nu_4, -\lambda, 1)$$
$$\int_{\partial AdS} \mathrm{d}^d P_5 \left\langle \mathcal{O}_{\nu_1}(P_1) \mathcal{O}_{\nu_2}(P_2) \mathcal{O}_{\lambda}^{(1)}(P_5; D_Z) \right\rangle \left\langle \mathcal{O}_{\nu_3}(P_3) \mathcal{O}_{\nu_4}(P_4) \mathcal{O}_{-\lambda}^{(1)}(P_5; Z) \right\rangle$$

$$\varphi_{\frac{d}{2}+i\lambda}^{(1)}(P_1, P_2; P_5) =$$



$$\mathcal{A}_{\mathcal{N}}^{\perp} = (ie)^2 \left(\frac{1}{P_{12}} \frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu} \int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda \rho_{\mathcal{N},\,\mathrm{bare}}^{J=1}(\lambda) \mathcal{F}_{\lambda,1}^{\left\{\frac{d}{2}+i\nu \equiv \Delta_\nu\right\}}(u, v)$$
$$= (ie)^2 \left(\frac{1}{P_{12}} \frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu} \int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda \frac{\lambda \sinh(\pi\lambda)}{64\pi^{d+2}} \frac{\Gamma\left(\frac{d-2\pm 2i\lambda}{4}\right)^2 \Gamma\left(\frac{d+2+4i\nu \pm 2i\lambda}{4}\right)^2}{\Gamma(1+i\nu)^2 \Gamma\left(\frac{d}{2}+i\nu\right)^2} \mathcal{F}_{\lambda,1}^{\{\Delta_\nu\}}(u, v),$$

$$\lambda = \pm i\left(\frac{d}{2}-1\right) \text{of } \Gamma\left(\frac{d-2\pm 2i\lambda}{4}\right)^2$$

$$\varphi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}(P_1, P_2; P_5; Z) = \int_{P_1}^{P_2} \mathrm{d}X^A \Pi_{\frac{d}{2}+i\lambda, AB}^{\mathrm{e}^{-\pi\lambda}}(X, P_5) Z^B$$
$$= \int_{-\infty}^{+\infty} \mathrm{d}\sigma \Pi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}\left(X(\sigma), P_5; \frac{dX}{d\sigma}(\sigma), Z\right)$$

$$X(\sigma) = \frac{P_1 e^{-\sigma} + P_2 e^{+\sigma}}{\sqrt{-2P_1 \cdot P_2}} \text{ with } \sigma \in \mathbb{R}$$

$$\varphi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}(P_1, P_2; P_5; Z)$$
$$= \sqrt{\mathcal{C}_{\lambda}^{\mathrm{e}^{-\pi\lambda}}} \int_{-\infty}^{+\infty} \mathrm{d}\sigma \frac{\left[(-2X(\sigma) \cdot P_5)\left(\frac{dX(\sigma)}{d\sigma} \cdot Z\right) - (X(\sigma) \cdot Z)\left(-2P_5 \cdot \frac{dX(\sigma)}{d\sigma}\right)\right]}{(-2X(\sigma) \cdot P_5)^{\frac{d}{2}+i\lambda+1}}$$
$$= 2\sqrt{\mathcal{C}_{\lambda}^{\mathrm{e}^{-\pi\lambda}}} \frac{[(P_2 \cdot Z)P_{15} - (P_1 \cdot Z)P_{25}]}{(-2P_1 \cdot P_2)^{\frac{1-\Delta_\lambda}{2}}} \int_{-\infty}^{+\infty} \mathrm{d}\sigma \frac{1}{[-2P_1 \cdot P_5 e^{-\sigma} - 2P_2 \cdot P_5 e^{\sigma}]^{1+\Delta_\lambda}}$$

$$\int_{-\infty}^{+\infty} \mathrm{d}\sigma \frac{1}{[Ae^{-\sigma}+Be^{\sigma}]^{1+\Delta_\lambda}} = \frac{1}{(AB)^{\frac{1+\Delta_\lambda}{2}}} \int_{-\infty}^{+\infty} \mathrm{d}\sigma \frac{1}{\left[\sqrt{\frac{A}{B}}e^{-\sigma}+\sqrt{\frac{B}{A}}e^{\sigma}\right]^{1+\Delta_\lambda}}$$

$$= \frac{1}{(AB)^{\frac{1+\Delta_\lambda}{2}}} \int_0^{+\infty} \frac{\mathrm{d}x}{x} \frac{1}{\left[x+\frac{1}{x}\right]^{1+\Delta_\lambda}}$$

$$= \frac{1}{(AB)^{\frac{1+\Delta_\lambda}{2}}} \frac{1}{2} \frac{\Gamma\left(\frac{\Delta_\lambda+1}{2}\right)^2}{\Gamma(\Delta_\lambda+1)}$$

$$\varphi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}(P_1,P_2;P_5;Z) = \underbrace{-\frac{\Gamma\left(\frac{\Delta_\lambda+1}{2}\right)^2}{\Gamma(\Delta_\lambda+1)}}_{b_g(\lambda)} \sqrt{\mathcal{C}_\lambda^{\mathrm{e}^{-\pi\lambda}}} \frac{[(P_1\cdot Z)P_{25}-(P_2\cdot Z)P_{15}]}{(-2P_1\cdot P_2)^{\frac{1-\Delta_\lambda}{2}}(-2P_1\cdot P_5)^{\frac{1+\Delta_\lambda}{2}}(-2P_2\cdot P_5)^{\frac{1+\Delta_\lambda}{2}}}$$

$$= \sqrt{\mathcal{C}_\lambda^{\mathrm{e}^{-\pi\lambda}}} b_g(\lambda) \cdot \left\langle \mathcal{O}_{\Delta=0}^{J=0}(P_1)\mathcal{O}_{\Delta=0}^{J=0}(P_2)\mathcal{O}_{\Delta_\lambda}^{J=1}(P_5;Z)\right\rangle_{\blacksquare}$$

$$\begin{cases} \varphi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}(\alpha P_1,P_2;P_5;Z) = \varphi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}(P_1,\alpha P_2;P_5;Z) = \varphi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}(P_1,P_2;P_5;Z) & \Longrightarrow \Delta_1=\Delta_2=0 \\ \varphi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}(P_1,P_2;\alpha P_5;Z) = \alpha^{-\Delta_\lambda-1}\varphi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}(P_1,P_2;P_5;Z) & \Longrightarrow \Delta_3=\Delta_\lambda \end{cases}$$

$$\varphi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}(P_1,P_2;P_5;Z)$$

$$\mathcal{A}_{gg}^{\perp} = (-ie)^2 \left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu} \frac{1}{\pi\left(\frac{d}{2}-1\right)} \int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda \frac{\lambda^2}{\lambda^2+\left(\frac{d}{2}-1\right)^2} \sqrt{\mathcal{C}_\lambda^{\mathrm{e}^{-\pi\lambda}}\mathcal{C}_{-\lambda}^{\mathrm{e}^{-\pi\lambda}}}$$

$$\int_{\partial AdS} \mathrm{d}^d P_5 \varphi_{\frac{d}{2}+i\lambda}^{\mathrm{e}^{-\pi\lambda}}(P_1,P_2;P_5;D_Z)\varphi_{\frac{d}{2}-i\lambda}^{\mathrm{e}^{-\pi\lambda}}(P_3,P_4;P_5;Z)$$

$$= (-ie)^2 \left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu} \frac{1}{\pi\left(\frac{d}{2}-1\right)} \int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda \frac{\lambda^2}{\lambda^2+\left(\frac{d}{2}-1\right)^2} \mathcal{C}_\lambda^{\mathrm{e}^{-\pi\lambda}}\mathcal{C}_{-\lambda}^{\mathrm{e}^{-\pi\lambda}} b_g(\lambda)b_g(-\lambda)$$

$$\int_{\partial AdS} \mathrm{d}^d P_5 \left\langle \mathcal{O}_{\Delta=0}^{J=0}(P_1)\mathcal{O}_{\Delta=0}^{J=0}(P_2)\mathcal{O}_{\Delta\lambda}^{J=1}(P_5;D_Z)\right\rangle_1 \left\langle \mathcal{O}_{\Delta=0}^{J=0}(P_3)\mathcal{O}_{\Delta=0}^{J=0}(P_4)\mathcal{O}_{\Delta-\lambda}^{J=1}(P_5;Z)\right\rangle_1$$

$$\mathcal{A}_{gg}^{\perp} = \frac{(-ie)^2}{\pi}\left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+i\nu} \int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda \frac{\lambda^2}{\lambda^2+\left(\frac{d}{2}-1\right)^2} \mathcal{C}_\lambda^{\mathrm{e}^{-\pi\lambda}}\mathcal{C}_{-\lambda}^{\mathrm{e}^{-\pi\lambda}} b_g(\lambda)b_g(-\lambda)\mathcal{F}_{\lambda,1}^{\{\Delta_i=0\}}(u,v)$$


$$\mathcal{F}_{\lambda,1}^{\{\Delta_i=0\}}(u,v) = \mathcal{F}_{\lambda,1}^{\{\Delta_v\}}(u,v)$$

$$\mathcal{A}_{gc}^{\perp} = (-ie)^2 \left(\frac{1}{P_{12}}\right)^{\frac{d}{2}+iv} \frac{1}{\pi\left(\frac{d}{2}-1\right)} \int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda \frac{\lambda^2}{\lambda^2+\left(\frac{d}{2}-1\right)^2} \sqrt{c_{\lambda}^{\mathrm{e}^{-\pi\lambda}} c_{-\lambda}^{\mathrm{e}^{-\pi\lambda}}}$$
$$\int_{\partial AdS} \mathrm{d}^d P_5 \varphi_{\frac{d}{2}+i\lambda}^{(1)}(P_1, P_2; P_5; D_Z)$$
$$\int_{AdS} \mathrm{d}^{d+1} X_2 \frac{1}{\left(\frac{d-1}{2}\right)} \Pi_{\frac{d}{2}-i\lambda}^{(1)}(X_2, P_5; Z, K_2) W_{2,B} T_{34}^{B}(P_3, P_4; X_2)$$

$$\mathcal{A}_{gc}^{\perp} = 2(ie)^2 \sqrt{c_{\nu_3}^{\mathrm{e}^{-\pi\lambda}} c_{\nu_4}^{\mathrm{e}^{-\pi\lambda}}} \left(\frac{1}{P_{12}}\right)^{\frac{d}{2}+iv} \frac{1}{\pi\left(\frac{d}{2}-1\right)} \int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda \frac{\lambda^2 c_{\lambda}^{(1)} c_{-\lambda}^{(1)}}{\lambda^2+\left(\frac{d}{2}-1\right)^2} b_g(\lambda) b_{\mathrm{Bulk}}(\nu_3, \nu_4, -\lambda, 1)$$
$$\int_{\partial AdS} \mathrm{d}^d P_5 \left\langle \mathcal{O}_{\Delta=0}^{J=0}(P_1) \mathcal{O}_{\Delta=0}^{J=0}(P_2) \mathcal{O}_{\Delta_\lambda}^{J=1}(P_5; D_Z) \right\rangle \left\langle \mathcal{O}_{\Delta-\lambda}^{J=1}(P_5; Z) \mathcal{O}_{\Delta=\frac{d}{2}+i\nu_3}^{J=0}(P_3) \mathcal{O}_{\Delta=\frac{d}{2}+i\nu_4}^{J=0}(P_4) \right\rangle$$

$$\mathcal{A}_{gc}^{\perp} = 2\frac{(-ie)^2}{\pi} c_{\nu}^{\mathrm{e}^{-\pi\lambda}} \left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+iv}$$
$$\int_{\mathbb{R}\oplus\circlearrowleft} \mathrm{d}\lambda \frac{\lambda^2}{\lambda^2+\left(\frac{d}{2}-1\right)^2} c_{\lambda}^{\mathrm{e}^{-\pi\lambda}} c_{-\lambda}^{\mathrm{e}^{-\pi\lambda}} b_g(\lambda) b_{\mathrm{Bulk}}(\nu, \nu, -\lambda, 1) \mathcal{F}_{\lambda,1}^{\{\Delta_{1,2}=0, \Delta_{3,4}=\Delta_\nu\}}(u, v)$$

$\mathcal{F}_{\lambda,1}^{\{\Delta_{1,2}=0,\Delta_{3,4}=\Delta_\nu\}}(u,v)$ is $\Delta_1 - \Delta_2$ and $\Delta_3 - \Delta_4$, so $\mathcal{F}_{\lambda,1}^{\{\Delta_{1,2}=0,\Delta_{3,4}=\Delta_\nu\}}(u,v)$ is exactly the same function as $\mathcal{F}_{\lambda,1}^{\{\Delta_i=\Delta_\nu\}}(u,v)$.



$$\begin{aligned}\mathcal{A}_{\mathcal{N}} =&\mathcal{A}_{cc}^{\perp}+\mathcal{A}_{gg}^{\perp}+\mathcal{A}_{gc}^{\perp}+\mathcal{A}_{cg}^{\perp}\\ =&\frac{(-ie)^2}{\pi}\left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+iv}\int_{\mathbb{R}\oplus\mathcal{O}}^{\mathcal{O}}\mathrm{d}\lambda\frac{\lambda^2}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\mathcal{C}_{\lambda}^{(1)}\mathcal{C}_{-\lambda}^{(1)}\mathcal{F}_{\lambda,1}^{\{\Delta_i=\Delta_v\}}(u,v)\\ &\left[4\left(\mathcal{C}_{v}^{\mathrm{e}^{-\pi\lambda}}\right)^2 b_{\text{Bulk}}(v,v,\lambda,1)b_{\text{Bulk}}(v,v,-\lambda,1)+b_g(\lambda)b_g(-\lambda)\right.\\ &\left.+2\mathcal{C}_{v}^{\mathrm{e}^{-\pi\lambda}}b_g(\lambda)b_{\text{Bulk}}(v,v,-\lambda,1)+2\mathcal{C}_{v}^{\mathrm{e}^{-\pi\lambda}}b_{\text{Bulk}}(v,v,\lambda,1)b_g(-\lambda)\right]\\ =&(ie)^2\left(\frac{1}{P_{12}}\frac{1}{P_{34}}\right)^{\frac{d}{2}+iv}\int_{\mathbb{R}\oplus\mathcal{O}}^{\mathcal{O}}\mathrm{d}\lambda\rho_{\mathcal{N}}^{J=1}\left(\lambda;\left\{\frac{d}{2}+iv\right\}\right)\mathcal{F}_{\lambda,1}^{\{\Delta_v\}}(u,v),\end{aligned}$$

$$\rho_{\mathcal{N}}^{J=1}\left(\lambda;\left\{\frac{d}{2}+iv\right\}\right)$$

$$\rho_{\mathcal{N}}^{J=1}\left(\lambda;\left\{\frac{d}{2}+iv\right\}\right)=\Pi_1^{(1),\perp}(\lambda)\mathcal{Q}_{\text{Dressed}}^{J=1}\left(\lambda,\left\{\frac{d}{2}+iv\right\}\right)$$

$$\Pi_1^{(1),\perp}=\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2},$$

$$\begin{aligned}\mathcal{Q}_{\text{Dressed}}^{J=1}\left(\lambda,\left\{\frac{d}{2}+iv\right\}\right)=&\frac{\lambda\sinh(\pi\lambda)}{4\pi^{d+2}}\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\frac{\Gamma\left(\frac{d+2\pm2i\lambda}{4}\right)^2}{\Gamma(1+iv)^2\Gamma\left(\frac{d}{2}+iv\right)^2}\\ &\left[\Gamma\left(\frac{d+2+4iv\pm2i\lambda}{4}\right)-\Gamma(1+iv)\Gamma\left(\frac{d}{2}+iv\right)\right]^2\end{aligned}$$

$$\mathcal{K}_{d-\Delta}^{(1)}=-\frac{\pi^{\frac{d}{2}}\Gamma\left(\frac{d}{2}-\Delta\right)\Gamma(d-\Delta)\Gamma\left(\frac{\Delta+1}{2}\right)^2}{2\Gamma(d-\Delta-1)\Gamma(1+\Delta)\Gamma\left(\frac{1+d-\Delta}{2}\right)^2}$$

$$\begin{aligned}\rho_{\mathcal{N}}^{J=1}\left(\lambda;\left\{\frac{d}{2}+iv\right\}\right)\mathcal{K}_{d-\Delta_\lambda}^{(1)}=&-\frac{\lambda\sinh(\pi\lambda)}{2^{5+\frac{d}{2}+i\lambda}\pi^{\frac{d+3}{2}}}\frac{\Gamma\left(\frac{d-2+2i\lambda}{4}\right)^2}{\Gamma(1+iv)^2\Gamma\left(\frac{d}{2}+iv\right)^2}\frac{\Gamma(-i\lambda)\Gamma\left(\frac{d+2+2i\lambda}{4}\right)}{\left(\frac{d}{2}-1-i\lambda\right)\Gamma\left(\frac{d+4+2i\lambda}{4}\right)}\\ &\left[\Gamma\left(\frac{d+2+4iv\pm2i\lambda}{4}\right)-\Gamma(1+iv)\Gamma\left(\frac{d}{2}+iv\right)\right]^2\end{aligned}$$

$$\hat{G}_{\Delta_\lambda}^{(1)}(u,v)\left(\hat{G}_{d-\Delta_\lambda}^{(1)}(u,v)\right)\hat{G}_{\Delta_\lambda}^{(1)}(u,v)$$

the factor $\Gamma\left(\frac{d-2+2i\lambda}{4}\right)^2\Gamma\left(\frac{d+2+2i\lambda}{4}\right)/\left(\frac{d}{2}-1-i\lambda\right)$ has a double pole at $\lambda=i\left(\frac{d}{2}-1\right)$, a simple pole at $\lambda=-i\left(\frac{d}{2}-1\right)$, and third-order poles at $\lambda=i\left(\frac{d}{2}-1+2k\right),\forall k\in\mathbb{N}_{>0}$.

$$\lambda=-i\left(\frac{d}{2}-1\right)\otimes_{\mathcal{O}}^{\mathcal{O}}\Gamma\left(\frac{d+2+4iv\pm2i\lambda}{4}\right)-\Gamma(1+iv)\Gamma\left(\frac{d}{2}+iv\right)$$

$$\rho^{J=1}_{\mathcal{N},\,\text{bare}}\left(\lambda;\left\{\frac{d}{2}+i\nu\right\}\right)=\Pi_1^{(1),\perp}(\lambda)\mathcal{Q}^{J=1}\left(\lambda,\left\{\frac{d}{2}+i\nu\right\}\right)$$

$$\rho^{J=1}_{\mathcal{N},\,\text{bare}}\left(\lambda;\left\{\frac{d}{2}+i\nu\right\}\right)\mathcal{K}^{(J=1)}_{d-\Delta_\lambda}\underset{\lambda\to i\left(\frac{d}{2}-1\right)}{\sim}-\frac{1}{16\pi^{\frac{d}{4}}\Gamma\left(2-\frac{d}{2}\right)\left(\lambda-i\left(\frac{d}{2}-1\right)\right)^2}$$


$$+\frac{i\left(\psi^{(0)}\left(\frac{d}{2}+i\nu\right)-\psi^{(0)}(1+i\nu)+\psi^{(0)}\left(1-\frac{d}{2}\right)+\gamma_E+\frac{d-3}{d-2}\right)}{16\pi^{\frac{d}{4}}\Gamma\left(2-\frac{d}{2}\right)\left(\lambda-i\left(\frac{d}{2}-1\right)\right)}+\mathcal{O}\left[\Gamma\left(\frac{d+2+4i\nu\pm 2i\lambda}{4}\right)-\Gamma(1+i\nu)\Gamma\left(\frac{d}{2}+i\nu\right)\right]^2$$

$$f_{ij}=\partial_i a_j-\partial_j a_i$$

$$\Phi=\frac{\varphi_1+i\varphi_2}{\sqrt{2}}$$

$$\mathcal{L}^{int}=eA_\mu(\varphi_2\nabla^\mu\varphi_1-\varphi_1\nabla^\mu\varphi_2)+\frac{e^2}{2}A_\mu A^\mu(\varphi_1^2+\varphi_2^2)$$

$\langle W_1^\mu A_\mu(X_1)W_2^\nu A_\nu(X_2)\rangle_{\text{Full}}=$ 

$B^{(1)}_{\nu_1,\nu_2}(X_1,X_2;W_1,W_2)=$ 

$$B^{(1)}_{\nu_1,\nu_2}(X_1,X_2;W_1,W_2)=-e^2\int_{-\infty}^{+\infty}\mathrm{d}\lambda\,\underbrace{\left(\langle JJ\rangle_{\nu_1,\nu_2}(\lambda)+\mathcal{T}_{\nu_1}+\mathcal{T}_{\nu_2}\right)}_{B^{(1)}_{\nu_1,\nu_2}}\Omega^{(1)}_\lambda(X_1,X_2;W_1,W_2)$$

$$\langle JJ\rangle_{\nu_1,\nu_2}(\lambda)=\sum_{n=0}^{+\infty}\frac{8\pi a^{d+2}_{\nu_1,\nu_2}(n)}{2i\alpha_n^-}\left[\frac{1}{i(\lambda-\alpha_n)}+(\lambda\to-\lambda)\right],$$

$$\alpha_n=-i\left(\frac{d}{2}+i\nu_1+i\nu_2+2n+1\right),$$

$$a^{d+2}_{\nu_1,\nu_2}(n)=\frac{\left(\frac{d+2}{2}\right)_n(1+i\nu_1+i\nu_2+n)_n(2n+2+d+i\nu_1+i\nu_2)_{-\frac{d}{2}}}{2\pi^{\frac{d+2}{2}}n!\left(\frac{d}{2}+i\nu_1+n+1\right)_{-\frac{d}{2}}\left(\frac{d}{2}+i\nu_2+n+1\right)_{-\frac{d}{2}}\left(\frac{d}{2}+i\nu_1+i\nu_2+n+1\right)_n}.$$

$$\langle JJ\rangle_{\nu,\nu}(\lambda) = -\frac{2\,\Gamma\left(\frac{d}{2}+1+2i\nu\right)\Gamma\left(\frac{d}{2}+1+i\nu\right)^2}{\pi^{d/2}\Gamma(d+2+2i\nu)\Gamma(1+i\nu)^2}$$
$$\times\left(\frac{{}_5F_4\left(\begin{matrix}\frac{d}{2}+1, i\nu+\frac{1}{2}, \frac{d}{2}+i\nu+1, \frac{d}{4}+i\nu-\frac{i\lambda}{2}+\frac{1}{2}, \frac{d}{2}+2i\nu+1\\ i\nu+1, \frac{d}{2}+i\nu+\frac{3}{2}, \frac{d}{4}+i\nu-\frac{i\lambda}{2}+\frac{3}{2}, 2i\nu+1\end{matrix};1\right)}{\frac{d}{2}+1+2i\nu-i\lambda}+(\lambda\to-\lambda)\right),$$

$$\langle JJ\rangle_{\nu,-\nu}(\lambda) = -\frac{\Gamma\left(\frac{d}{2}+1+i\nu\right)\Gamma\left(\frac{d}{2}+1-i\nu\right)}{(4\pi)^{\frac{d-1}{2}}(d+1)\Gamma\left(\frac{d+1}{2}\right)\Gamma(1+i\nu)\Gamma(1-i\nu)}$$
$$\times\left(\frac{{}_5F_4\left(\begin{matrix}\frac{1}{2}, \frac{d}{2}+1, \frac{d}{4}-\frac{i\lambda}{2}+\frac{1}{2}, \frac{d}{2}-i\nu+1, \frac{d}{2}+i\nu+1\\ \frac{d}{2}+\frac{3}{2}, \frac{d}{4}-\frac{i\lambda}{2}+\frac{3}{2}, 1-i\nu, i\nu+1\end{matrix};1\right)}{\frac{d}{2}+1-i\lambda}+(\lambda\to-\lambda)\right).$$

$$\langle JJ\rangle_{\nu,\pm\nu}(\lambda) \underset{d\to3}{\sim} \frac{1}{3-d}\left(\frac{\lambda^2+\frac{1}{4}}{24\pi^2}-\frac{\nu^2+\frac{1}{4}}{4\pi^2}\right)+\overline{\langle JJ\rangle}_{\nu,\pm\nu}(\lambda)+\mathcal{O}(3-d).$$

$$\mathcal{T}_\nu = \frac{\Gamma\left(\frac{1-d}{2}\right)\Gamma\left(\frac{d}{2}+i\nu\right)}{(4\pi)^{\frac{d+1}{2}}\Gamma\left(-\frac{d}{2}+i\nu+1\right)}$$

$$\mathcal{T}_\nu \underset{d\to3}{\sim} \frac{\nu^2+\frac{1}{4}}{8\pi^2}\frac{1}{3-d}+\overline{\mathcal{T}}_\nu+\mathcal{O}(3-d)$$
$$\overline{\mathcal{T}}_\nu = \frac{\nu^2+\frac{1}{4}}{16\pi^2}\left(\psi(i\nu-1)-\psi\left(i\nu+\frac{3}{2}\right)-2\psi(2i\nu-2)-\gamma_{\mathrm{E}}+1+\log(16\pi)\right),$$

$$B^{(1)}_{\nu,\pm\nu}(\lambda) \underset{d\to3}{\sim} \frac{1}{3-d}\frac{\lambda^2+\frac{1}{4}}{24\pi^2}+\overline{\langle JJ\rangle}_{\nu,\pm\nu}(\lambda)+\overline{\mathcal{T}}_\nu+\overline{\mathcal{T}}_{\pm\nu}+\mathcal{O}(3-d)$$

$$\langle A(X_1,W_1)A(X_2,W_2)\rangle_{\text{one-loop}}$$
$$=\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2+e^2B^{(1)}_{\nu_1,\nu_2}(\lambda)}\Omega^{(1)}_\lambda(X_1,X_2;W_1,W_2)$$
$$+\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{\xi}{\left(\lambda^2+\frac{d^2}{4}\right)^2}(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega^{(0)}_\lambda(X_1,X_2)$$

$$A_\mu = \sqrt{Z_A}A^{\text{ren}}_\mu,\ e^2=\mu^{3-d}\frac{e^2_{\text{ren}}(\mu)}{Z_A}$$
$$\text{with } Z_A = 1+e^2_{\text{ren}}\underbrace{\left(\frac{1}{3-d}\delta z_A^{\text{div}}+\delta z_A^{\text{finite}}\right)}_{\delta z_A}+\mathcal{O}(e^4_{\text{ren}})$$

$$B^{(1),\mathrm{ren}}_{\nu,\pm\nu}(\lambda)= \mu^{3-d}B^{(1)}_{\nu,\pm\nu}(\lambda) + \delta z_A\left(\lambda^2+\left(\frac{d}{2}-1\right)^2\right)+\mathcal{O}(e^2_{\mathrm{ren}}) = \text{ finite}$$

$$\Rightarrow \delta z_A^{\mathrm{div}} = -\frac{1}{24\pi^2}$$

$$\beta_{e^2_{\mathrm{ren}}} = \frac{de^2_{\mathrm{ren}}}{d\log\,\mu} \underset{d=3}{=} \frac{e^4_{\mathrm{ren}}}{24\pi^2}+\mathcal{O}(e^6_{\mathrm{ren}})$$

$$B^{(1),\mathrm{ren}}_{\nu,\pm\nu}(\lambda) = \overline{\langle JJ\rangle}_{\nu,\pm\nu}(\lambda)+\overline{\mathcal{T}}_{\nu}+\overline{\mathcal{T}}_{\pm\nu}+\frac{1}{48\pi^2}+C\left(\lambda^2+\frac{1}{4}\right)$$

$$\begin{aligned}\langle A^{\mathrm{ren}}(X_1,W_1)A^{\mathrm{ren}}(X_2,W_2)\rangle_{\text{one-loop}} &= \int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{1}{\lambda^2+\frac{1}{4}+e^2_{\mathrm{ren}}B^{(1),\mathrm{ren}}_{\nu_1,\nu_2}(\lambda)}\Omega^{(1)}_{\lambda}(X_1,X_2;W_1,W_2)\\ &+\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{\xi}{\left(\lambda^2+\frac{9}{4}\right)^2}(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega^{(0)}_{\lambda}(X_1,X_2)\end{aligned}$$

$$B^{(1)}_{\nu,\nu}\left(\pm i\left(\frac{d}{2}-1\right)\right)=0.$$

$$B^{(1)}_{\nu,\nu}\left(\pm i\left(\frac{d}{2}-1\right)\right)$$

$$B^{(1),\,\mathrm{ren}}_{\nu,\nu}\left(\pm\frac{i}{2}\right)=0$$

$$\begin{aligned}\langle\Phi^2\rangle &= \frac{1}{2}\left(\langle\varphi_1^2\rangle-\langle\varphi_2^2\rangle\right)\\ &= \frac{1}{2}\left(\mathcal{T}_{\nu}-\mathcal{T}_{-\nu}\right)\\ &= -\frac{\sin(\pi i\nu)\Gamma\left(\frac{d}{2}+i\nu\right)\Gamma\left(\frac{d}{2}-i\nu\right)}{(4\pi)^{\frac{d+1}{2}}\Gamma\left(\frac{d+1}{2}\right)}\\ &\underset{d=3}{=} -\frac{(4\nu^2+1)\tan(\pi i\nu)}{64\pi}\end{aligned}$$

$$\begin{aligned}M^2_{\gamma} &= -e^2\frac{4i\nu}{d}\langle\Phi^2\rangle\\ &= e^2\frac{4i\nu\sin(\pi i\nu)\Gamma\left(\frac{d}{2}+i\nu\right)\Gamma\left(\frac{d}{2}-i\nu\right)}{(4\pi)^{\frac{d+1}{2}}d\Gamma\left(\frac{d+1}{2}\right)}+\mathcal{O}(e^4)\end{aligned}$$

$$\frac{1}{e^2_{\mathrm{ren}}}\left(\lambda^2+\frac{1}{4}\right)$$

$$B^{(1),\mathrm{ren}}_{\nu,-\nu}\left(\pm\frac{i}{2}\right)=\overline{\langle JJ\rangle}_{\nu,-\nu}\left(\pm\frac{i}{2}\right)+\overline{\mathcal{T}}_{\nu}+\overline{\mathcal{T}}_{-\nu}+\frac{1}{48\pi^2}=\frac{(4\nu^2+1)i\nu\tan(\pi i\nu)}{48\pi}$$

$$\overline{\langle JJ\rangle}_{\nu,-\nu}\left(\pm\frac{i}{2}\right)\bigotimes\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\longrightarrow\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2+e^2B^{(1)}_{\nu,\pm\nu}(\lambda)}$$

$$\Delta_j=d-1+e^2\gamma_j+\mathcal{O}(e^4)$$

$$M_\gamma^2=\big(\Delta_j-1\big)\big(\Delta_j-d+1\big)=(d-2)e^2\gamma_j+\mathcal{O}(e^4)$$

$$\delta O(x)=\delta\alpha O'(x)$$

$$\text{SSB: }\langle O'(x)\rangle=a_{O'}\neq 0$$

$$\delta S=-\int d^{d+1}x\sqrt{g(x)}\nabla_\mu\delta\alpha(x)J^\mu(x)\Rightarrow\nabla_\mu J^\mu(x)=\frac{\delta S}{\delta\alpha(x)}$$

$$\nabla_\mu^{x_1}\langle J^\mu(x_1)O(x_2)\rangle=-\frac{\int\mathcal{D}\Phi O(x_2)\frac{\delta}{\delta\alpha(x_1)}e^{-S}}{\int\mathcal{D}\Phi e^{-S}}$$
$$=\delta^{d+1}(x_1-x_2)\langle O'(x_2)\rangle$$

$$-\lim_{z_1\to 0}\int d^d\vec{x}_1 z_1^{-d-1}\langle J^z(z_1,\vec{x}_1)O(z_2,\vec{x}_2)\rangle=a_{O'}\neq 0$$

$$\langle t(\vec{x}_1)O(z_2,\vec{x}_2)\rangle=N\left(\frac{z_2}{z_2^2+(\vec{x}_1-\vec{x}_2)}\right)^d$$

$$\int d^d\vec{x}_1\left(\frac{z_2}{z_2^2+(\vec{x}_1-\vec{x}_2)}\right)^d=\frac{2^{1-d}\pi^{\frac{d+1}{2}}}{\Gamma\left(\frac{d+1}{2}\right)}$$

$$N=-\frac{2^{d-1}\Gamma\left(\frac{d+1}{2}\right)}{\pi^{\frac{d+1}{2}}}a_{O'}$$

$$\langle t(\vec{x}_1)t(\vec{x}_2)\rangle=\frac{C_t}{|\vec{x}_1-\vec{x}_2|^{2d}}$$

$$C_t=-\frac{2^{d-1}\Gamma\left(\frac{d+1}{2}\right)}{\pi^{\frac{d+1}{2}}}\frac{a_{O'}}{b_{Ot}}$$

$$\nabla_\mu J^\mu(x)=\frac{\delta S}{\delta\alpha(x)}$$

$$\delta\alpha(z,\vec{x})\underset{z\to 0}{\sim}\delta\alpha_\partial(\vec{x})\oplus\partial^\square\mathfrak{S}_{\text{subleading}}\oslash\partial\tilde{\psi}\hat{\psi}_\star^\dagger$$
$$\Rightarrow\delta S=-\int d^{d+1}x\sqrt{g(x)}\nabla_\mu\delta\alpha(x)J^\mu(x)-\int_{z=0}d^d\vec{x}\delta\alpha_\partial(\vec{x})t(\vec{x})$$
$$=\int d^{d+1}x\sqrt{g(x)}\delta\alpha(x)\nabla_\mu J^\mu(x)$$

$$S = S_m + \int d^{d+1}x\sqrt{g(x)} \bigotimes \langle \frac{1}{4}F^{\mu\nu}F_{\mu\nu} + eA_\mu J^\mu + \mathcal{O}(A_\mu^2)\rangle$$

$$\delta A_\mu(x) = \frac{1}{e}\nabla_\mu \delta\alpha(x)$$

$$A_i(z,\vec{x}) \underset{z\to 0}{\sim} z^{d-2}j_i(\vec{x}) + a_i(\vec{x}) + \cdots$$

$$\int_{z=0} d^d\vec{x}(d-2)a_i(\vec{x})j^i(\vec{x})$$

$$\delta\alpha(z,\vec{x}) \underset{z\to 0}{\sim} \delta\alpha_\partial(\vec{x}) \oplus \partial^\square \mathfrak{S}_{\text{subleading}} \oslash \partial\tilde{\psi}\hat{\psi}_\star^\dagger$$
$$\Rightarrow \delta a_i(\vec{x}) = \frac{1}{e}\partial_i \delta\alpha_\partial(\vec{x})$$

$$\delta S = \int_{z=0} d^d\vec{x}\partial_i\delta\alpha_\partial(\vec{x})\frac{1}{e}(d-2)j^i(\vec{x})$$

$$\langle j_i(\vec{x})j_k(0)\rangle = C_j^{(0)}\frac{I_{ik}(\vec{x})}{|\vec{x}|^{2(d-1)}}\left(1+\mathcal{O}(e^2)\right), I_{ik} \equiv \delta_{ik} - \frac{2x_i x_k}{\vec{x}^2}$$
$$C_j^{(0)} = \frac{\Gamma(d)}{2(d-2)\pi^{\frac{d}{2}}\Gamma\left(\frac{d}{2}\right)}$$

$$\delta S = \int_{z=0} d^d\vec{x}\delta\alpha_\partial(\vec{x})\left(-\frac{1}{e}(d-2)\partial_i j^i(\vec{x}) - t(\vec{x})\right)$$

$$\partial_i j^i(\vec{x}) = -\frac{e}{d-2}t(\vec{x})$$

$$\langle j_i(\vec{x})j_k(0)\rangle = C_j(e^2)\frac{I_{ik}(\vec{x})}{|\vec{x}|^{2\Delta_j(e^2)}}$$

$$\langle t(\vec{x})t(0)\rangle = 4\frac{(d-2)^2}{e^2}C_j(e^2)\left(\Delta_j(e^2)-d+1\right)\left(\Delta_j(e^2)-\frac{d}{2}+1\right)\frac{1}{|\vec{x}|^{2\Delta_j(e^2)+2}}$$

$C_j(e^2) = C_j^{(0)} + \mathcal{O}(e^2)$ and $\Delta_j(e^2) = d - 1 + e^2\gamma_j + \mathcal{O}(e^4)$

$$\gamma_j = \frac{C_t}{2d(d-2)^2 C_j^{(0)}}$$

$$\gamma_j = -\frac{1}{d(d-2)}\frac{a_{O'}}{b_{Ot}}$$

$$M_\gamma^2 = \left(\Delta_j(e^2)-1\right)\left(\Delta_j(e^2)-d+1\right) = (d-2)e^2\gamma_j + \mathcal{O}(e^4)$$

$$M_\gamma^2 = -\frac{e^2}{d}\frac{a_{O'}}{b_{Ot}} + \mathcal{O}(e^4)$$

$$\Phi = \frac{1}{\sqrt{2}}(\varphi_1 + i\varphi_2)$$

$$\Phi = (v + \delta\rho)e^{i\Omega}$$

$$J_\mu = iq\left(\Phi^*\nabla_\mu\Phi - \Phi\nabla_\mu\Phi^*\right) = -2qv^2\nabla_\mu\Omega + \cdots$$

$$J_z(z,\vec{x}) \underset{z\to 0}{\sim} -2qv^2 b_{\Omega t} dz^{d-1}(t(\vec{x}) \oplus \partial^\square \mathfrak{S}_{\text{subleading}} \oslash \partial\tilde{\psi}\hat{\psi}_\star^\dagger) \Rightarrow b_{\Omega t} = -\frac{1}{2qv^2 d}$$

$O = \varphi_2 = \sqrt{2}(v + \delta\rho)\sin\ \Omega = \sqrt{2}v\Omega + \cdots$, so that $b_{Ot} = \sqrt{2}vb_{\Omega t} = -\frac{1}{\sqrt{2}qvd}$, and $O' = q\varphi_1 = \sqrt{2}q(v + \delta\rho)\cos\ \Omega$ so that $a_{O'} = \sqrt{2}qv$

$$M_\gamma^2 = -\frac{e^2}{d}(-\sqrt{2}qvd)\sqrt{2}qv + \mathcal{O}(e^4) = 2e^2q^2v^2 + \mathcal{O}(e^4)$$

$$\Phi = \frac{\varphi_1 + \varphi_2}{\sqrt{2}}$$

$$\varphi_1(z,\vec{x}) \underset{z\to 0}{\sim} z^{\frac{d}{2}+iv}\left(O_1(\vec{x}) \oplus \partial^\square \mathfrak{S}_{\text{subleading}} \oslash \partial\tilde{\psi}\hat{\psi}_\star^\dagger\right),$$
$$\varphi_2(z,\vec{x}) \underset{z\to 0}{\sim} z^{\frac{d}{2}-iv}\left(O_2(\vec{x}) \oplus \partial^\square \mathfrak{S}_{\text{subleading}} \oslash \partial\tilde{\psi}\hat{\psi}_\star^\dagger\right),$$

$$J_z(z,\vec{x}) \underset{z\to 0}{\sim} z^{d-1}2iv(O_1O_2(\vec{x})) \Rightarrow t(\vec{x}) = 2ivO_1O_2(\vec{x})$$

$O = \mathrm{Im}\Phi^2 = \frac{1}{2}\varphi_1\varphi_2$, so that $b_{Ot} = \frac{1}{4iv}$, and $O' = \mathrm{Re}\Phi^2 = \frac{1}{2}(\varphi_1^2 - \varphi_2^2)$, which gives

$$\begin{aligned} a_{O'} &= \frac{1}{2}\left(\langle\varphi_1^2\rangle - \langle\varphi_2^2\rangle\right) \\ &= \frac{1}{2}\lim_{x_1\to x_2}\left(\Pi^{(0)}_{\frac{d}{2}+iv}(x_1,x_2) - \Pi^{(0)}_{\frac{d}{2}-iv}(x_1,x_2)\right) \\ &= -\frac{\sin(\pi iv)\Gamma\left(\frac{d}{2}+iv\right)\Gamma\left(\frac{d}{2}-iv\right)}{(4\pi)^{\frac{d+1}{2}}\Gamma\left(\frac{d+1}{2}\right)} \end{aligned}$$

$$\gamma_j = \frac{4iv\sin(\pi iv)\Gamma\left(\frac{d}{2}+iv\right)\Gamma\left(\frac{d}{2}-iv\right)}{(4\pi)^{\frac{d+1}{2}}d(d-2)\Gamma\left(\frac{d+1}{2}\right)}$$


$$M_\gamma^2 = e^2\frac{4iv\sin(\pi iv)\Gamma\left(\frac{d}{2}+iv\right)\Gamma\left(\frac{d}{2}-iv\right)}{(4\pi)^{\frac{d+1}{2}}d\Gamma\left(\frac{d+1}{2}\right)} + \mathcal{O}(e^4)$$

$$\mathcal{J}_{iz}(\vec{x},z) \underset{z\to 0}{\sim} z^{d-3}\tau_i(\vec{x}) + \mathcal{O}\left(z^{d-2}\right)$$

$$\left(\star^{(D)}\mathcal{J}\right)_{i_1\ldots i_{d-1}}(\vec{x},z) \underset{z\to 0}{\sim} \varepsilon_{i_1\ldots i_d}\tau^{i_d}(\vec{x}) + \mathcal{O}(z)$$

$$\left(\star^{(D)} A\right)_{i_{k+1}\dots i_D} = \frac{\sqrt{|g|}}{(D-k)!}\varepsilon_{i_1\dots i_D}A^{i_1\dots i_k}$$

$$\mathcal{Q}_G = \int_{\Sigma^{(D-2)}} \star^{(D)} \mathcal{J} \underset{\Sigma^{(D-2)}\to\overrightarrow{\Sigma'^{(d-1)}}\subseteq\partial\mathrm{AdS}}{} \int_{\Sigma'^{(d-1)}} \star^{(D)} \mathcal{J} = \int_{\Sigma'^{(d-1)}} \star^{(d)} \tau$$



$$\mathcal{J}_{\mathcal{E}} = \frac{1}{e}F, \mathcal{J}_{\mathcal{M}} = \frac{e}{2\pi}\star^{(D)} F.$$

$$F_{zi} \underset{z\to 0}{\sim} (d-2)j_i(\vec{x}) + \mathcal{O}(z), \left(\star^{(D)} F\right)_{kz} \underset{z\to 0}{\sim} \mathcal{O}(z)$$

$$Q_{\mathcal{E}} = \int_{\Sigma^{(2)}} \star^{(D)} J_{\mathcal{E}} \longrightarrow (d-2)\int_{\Sigma^{(2)}} \star^{(d)} j$$

$$F_{iz} \underset{z\to 0}{\sim} \mathcal{O}(z), \left(\star^{(D)} F\right)_{iz} \underset{z\to 0}{\sim} \varepsilon_{ijk}f_{jk}(\vec{x}) + \mathcal{O}(z).$$

$$Q_{\mathcal{M}} = \int_{\Sigma^{(2)}} \star^{(D)} J_{\mathcal{M}} = \int_{\Sigma^{(2)}} \frac{e}{4\pi} F_{\alpha\beta}\left(\mathrm{d}x^{\alpha} \wedge \mathrm{d}x^{\beta}\right) = \frac{e}{2\pi} \int_{\Sigma^{(2)}} F_{\alpha\beta} \mathrm{d}x^{\alpha} \mathrm{d}x^{\beta} \longrightarrow \frac{e}{2\pi} \int_{\Sigma^{(2)}} f_{ij}\, \mathrm{d}x^{i}\, \mathrm{d}x^{j}$$

$$\nabla_{\mu} (\mathcal{J}_{\mathcal{E}})^{\mu\nu} = J^{\nu}_{\text{mat}}$$

$$\nabla_{\mu} (\mathcal{J}_{\mathcal{E}})^{\mu z} \underset{z\to 0}{\sim} -\frac{1}{e} z^{d+1} (d-2) \partial_i j^i(\vec{x})$$

$$\nabla_{\mu} (\mathcal{J}_{\mathcal{E}})^{\mu\nu} = J^{\nu}_{\text{mat}} \implies \partial_i j^i(\vec{x}) = -\frac{e^{\varphi_1 \partial_z \varphi_2, z^{d+2i\nu-1} \varphi_{[1} \partial_z \varphi_{2]}}}{d-2} \lim_{z\to 0} z^{-(d+1)} J^{z}_{\text{mat}}(\vec{x}, z)$$


$$\varphi_i(\vec{x}, z) \underset{z\to 0}{\sim} z^{\frac{d}{2}+i\nu} \mathcal{O}_i(\vec{x}) \oplus \partial^{\square} \mathfrak{S}_{\text{subleading}} \oslash \partial\tilde{\psi}\hat{\psi}^{\dagger}_{\star}$$
$$\varphi_1 \partial_z \varphi_2(\vec{x}, z) \underset{z\to 0}{\sim} z^{d+2i\nu-1} 2\Delta_{\nu} \mathcal{O}_1 \mathcal{O}_2(\vec{x}) \oplus \partial^{\square} \mathfrak{S}_{\text{subleading}} \oslash \partial\tilde{\psi}\hat{\psi}^{\dagger}_{\star}$$
$$J_{\text{mat},z}(\vec{x}, z) \underset{z\to 0}{\sim} z^{d+2i\nu+1} \partial^i \left(\mathcal{O}_{[1} \partial_i \mathcal{O}_{2]}\right)(\vec{x}) \oplus \partial^{\square} \mathfrak{S}_{\text{subleading}} \oslash \partial\tilde{\psi}\hat{\psi}^{\dagger}_{\star}$$

$$J^{z}_{\text{mat}}(\vec{x}, z) \overset{z\to 0}{\sim} \mathcal{O}\left(z^{d+3+2i\nu}\right)$$

$$Q_{\mathcal{E}} = \int_{\Sigma^{(d-1)} \subseteq \partial \text{AdS}} \star^{(D)} J_{\mathcal{E}} = \int_{\Sigma^{(d-1)} \subseteq \partial \text{AdS}} \star^{(d)} j$$

$$\left(\frac{d}{2} \pm i\nu\right) z^{d-1} \mathcal{O}_1 \mathcal{O}_2$$

$$\nabla_{\mu} (\mathcal{J}_{\mathcal{E}})^{\mu z} = e J^{z}_{\text{mat}} \implies z^{d+1} (d-2) \partial^i j_i(\vec{x}) = z^{d+1} e \underbrace{2i\nu \mathcal{O}_1 \mathcal{O}_2(\vec{x})}_{t(\vec{x})}$$



$$\int_{\Sigma z^{d-1} v^2 t(\vec{x})} \left\langle \mathcal{O}_{\Delta_1}(P_1) W[\gamma(P_1, P_2)] \mathcal{O}_{\Delta_2}(P_2) f_{ij}(P_3) \right\rangle \mathrm{d}x^{i}_{3}\, \mathrm{d}x^{j}_{3}$$

$$\varepsilon_{ijk} \left\langle \mathcal{O}_{\Delta_1}(P_1) \mathcal{O}_{\Delta_2}(P_2) \tilde{j}_k(P_3) \right\rangle = \frac{c_{\mathcal{O}_{\Delta_1} \mathcal{O}_{\Delta_2} f}}{|P_{12}|^{\Delta_{123}} |P_{13}|^{\Delta_{132}} |P_{23}|^{\Delta_{231}}} \varepsilon_{ijk} R_k(P_1, P_2 \mid P_3)$$
$$\Delta_{ijk} := \frac{\Delta_i + \Delta_j - \Delta_k}{2}, R_k(P_1, P_2 \mid P_3) := \frac{\partial P^{M}_{3}}{\partial x^k} \left[\frac{(P_1 \cdot P_3) P_{2,M} - (P_2 \cdot P_3) P_{1,M}}{\sqrt{(P_1 \cdot P_2)(P_1 \cdot P_3)(P_2 \cdot P_3)}} \right]$$

$$\int_{\Sigma^{(d-1)}} \left\langle \mathcal{O}_{\Delta_1}(P_1) W[\gamma(P_1, P_2)] \mathcal{O}_{\Delta_2}(P_2) \left(\star^{(d)} j\right)(P_3) \right\rangle = 0$$

$$\left\langle \varphi_1(P_1) W_\gamma[P_1,P_2] \varphi_2(P_2) \varphi_1(P_3) W_\gamma[P_3,P_4] \varphi_2(P_4) \right\rangle$$

$$\begin{aligned}\Pi^{\mathcal{D}}_{d-1}(X_1,X_2;W_1,W_2) =& \int_{-\infty}^{+\infty} \mathrm{d}\lambda \Pi^{\perp}_{\mathcal{D}}(\lambda) \Omega^{(1)}_\lambda(X_1,X_2;W_1,W_2) \\ &+ \int_{-\infty}^{+\infty} \mathrm{d}\lambda \Pi^{\parallel}_{\mathcal{D}}(\lambda)(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega^{(0)}_\lambda(X_1,X_2)\end{aligned}$$

$$\begin{aligned}-\nabla_1^2 \Omega^{(1)}_\lambda(X_1,X_2;W_1,W_2) &= \left(\lambda^2+\frac{d^2}{4}+1\right)\Omega^{(1)}_\lambda(X_1,X_2;W_1,W_2) \\ (K_1\cdot\nabla_1)\Omega^{(1)}_\lambda(X_1,X_2;W_1,W_2) &= 0\end{aligned}$$

$$\begin{aligned}K_A =& \frac{d-1}{2}\left[\frac{\partial}{\partial W^A} - X_A\left(X\cdot\frac{\partial}{\partial W}\right)\right] + \left(W\cdot\frac{\partial}{\partial W}\right)\frac{\partial}{\partial W^A} \\ &+ X_A\left(W\cdot\frac{\partial}{\partial W}\right)\left(X\cdot\frac{\partial}{\partial W}\right) - \frac{1}{2}W_A\left[\frac{\partial^2}{\partial W\cdot\partial W} + \left(X\cdot\frac{\partial}{\partial W}\right)\left(X\cdot\frac{\partial}{\partial W}\right)\right]\end{aligned}$$

$$H_{A_1\ldots A_J} = \frac{1}{J!\left(\frac{d-1}{2}\right)_J} K_{A_1}\ldots K_{A_J} H(X,W)$$

$$\begin{aligned}&\left[-\nabla_1^2 - d + \frac{1}{\frac{d-1}{2}}\left(1-\frac{1}{\xi}\right)(W_1\cdot\nabla_1)(K_1\cdot\nabla_1)\right]\int_{-\infty}^{+\infty}\mathrm{d}\lambda\Pi^{\perp}_{\mathcal{D}}(\lambda)\Omega^{(1)}_\lambda(X_1,X_2;W_1,W_2) \\ &\qquad = \int_{-\infty}^{+\infty}\mathrm{d}\lambda\left(\lambda^2+\left(\frac{d}{2}-1\right)^2\right)\Pi^{\perp}_{\mathcal{D}}(\lambda)\Omega^{(1)}_\lambda(X_1,X_2;W_1,W_2)\end{aligned}$$

$$\Pi^{\perp}_{\mathcal{D}}(\lambda) = \frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}$$

$$\begin{aligned}\int_{-\infty}^{+\infty}\mathrm{d}\lambda\Omega^{(1)}_\lambda(X_1,X_2;W_1,W_2) =& (W_1\cdot W_2)\delta^{d+1}(X_1,X_2) \\ &-(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{1}{\lambda^2+\frac{d^2}{2}}\Omega^{(0)}_\lambda(X_1,X_2)\end{aligned}$$

$$\begin{aligned}&-\nabla_1^2\left[(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega^{(0)}_\lambda(X_1,X_2)\right] \\ &= (W_1\cdot\nabla_1)(W_2\cdot\nabla_2)[d-\nabla_1^2]\Omega^{(0)}_\lambda(X_1,X_2) \\ &= \left(\lambda^2+\frac{d^2}{4}+d\right)(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega^{(0)}_\lambda(X_1,X_2) \\ &(K_1\cdot\nabla_1)(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega^{(0)}_\lambda(X_1,X_2) \\ &= (W_2\cdot\nabla_2)\frac{d-1}{2}\nabla_1^2\Omega^{(0)}_\lambda(X_1,X_2) \\ &= -\frac{d-1}{2}\left(\lambda^2+\frac{d^2}{4}\right)(W_2\cdot\nabla_2)\Omega^{(0)}_\lambda(X_1,X_2)\end{aligned}$$

$$[\nabla_1^2,(W_1\cdot\nabla_1)] = -d(W_1\cdot\nabla_1)$$

$$0 = -\frac{1}{\lambda^2+\frac{d^2}{4}} + \Pi_{\mathcal{D}}^{\|}\left[\lambda^2+\frac{d^2}{4} - \left(1-\frac{1}{\xi}\right)\left(\lambda^2+\frac{d^2}{4}\right)\right]$$

$$\Pi_{\mathcal{D}}^{\|}(\lambda) = \frac{\xi}{\left(\lambda^2+\frac{d^2}{4}\right)^2}$$

$$\Pi_{d-1}^{\mathcal{D}}(X_1,X_2;W_1,W_2) = \int_{-\infty}^{+\infty} \mathrm{d}\lambda \frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\Omega_{\lambda}^{(1)}(X_1,X_2;W_1,W_2)$$
$$+\int_{-\infty}^{+\infty} \mathrm{d}\lambda \frac{\xi}{\left(\lambda^2+\frac{d^2}{4}\right)^2}(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega_{\lambda}^{(0)}(X_1,X_2)$$

$$u := \frac{(X_1-X_2)^2}{2} = -(1+X_1\cdot X_2)$$

$$u = \frac{(z_1-z_2)^2+|\vec{x}_{12}|^2}{2z_1z_2}$$

$$u \sim \frac{|\vec{x}_{12}|^2+z_2^2}{2z_2}\frac{1}{z_1} \to \infty$$

$$\Pi_{d-1,\mu\nu}^{\mathcal{D}}(x_1,x_2) = -F_0^{\mathcal{D}}\frac{\partial^2 u}{\partial x_1^{\mu}\partial x_2^{\nu}} + F_1^{\mathcal{D}}\frac{\partial u}{\partial x_1^{\mu}}\frac{\partial u}{\partial x_2^{\nu}}$$

$$\Omega_{\lambda}^{(1)}(X_1,X_2;W_1,W_2) = \omega_{0,\lambda}(u)(W_1\cdot W_2) + \omega_{1,\lambda}(u)(X_1\cdot W_2)(X_2\cdot W_1),$$
$$(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega_{\lambda}^{(0)}(X_1,X_2) = -\frac{\partial\Omega_{\lambda}^{(0)}(u)}{\partial u}(W_1\cdot W_2) + \frac{\partial^2\Omega_{\lambda}^{(0)}(u)}{\partial u^2}(X_1\cdot W_2)(X_2\cdot W_1),$$

$$\omega_{0,\lambda}(u) = \frac{\lambda\sinh(\pi\lambda)(d^2+4\lambda^2)\Gamma\left(\frac{d}{2}-1\pm i\lambda\right)}{2^{d+4}\pi^{\frac{d+3}{2}}\Gamma\left(\frac{d+3}{2}\right)}$$
$$\left[(d+1)\,{}_2F_1\left(\frac{d}{2}+i\lambda,\frac{d}{2}-i\lambda,\frac{d+1}{2};-\frac{u}{2}\right)\right.$$
$$\left.-(1+u)\,{}_2F_1\left(\frac{d}{2}+1+i\lambda,\frac{d}{2}+1-i\lambda,\frac{d+3}{2};-\frac{u}{2}\right)\right]$$
$$\omega_{1,\lambda}(u) = \frac{\lambda\sinh(\pi\lambda)(d^2+4\lambda^2)\Gamma\left(\frac{d}{2}-1\pm i\lambda\right)}{2^{d+4}\pi^{\frac{d+3}{2}}\Gamma\left(\frac{d+3}{2}\right)}\frac{1}{u(2+u)}$$
$$\left[(d+1)(1+u)\,{}_2F_1\left(\frac{d}{2}+i\lambda,\frac{d}{2}-i\lambda,\frac{d+1}{2};-\frac{u}{2}\right)\right.$$
$$\left.-(d+(1+u)^2)\,{}_2F_1\left(\frac{d}{2}+1+i\lambda,\frac{d}{2}+1-i\lambda,\frac{d+3}{2};-\frac{u}{2}\right)\right]$$
$$\Omega_{\lambda}^{(0)}(u) = \frac{1}{(4\pi)^{\frac{d+1}{2}}\Gamma\left(\frac{d+1}{2}\right)}\frac{\Gamma\left(\frac{d}{2}\pm i\lambda\right)}{\Gamma(\pm i\lambda)}\,{}_2F_1\left(\frac{d}{2}+i\lambda,\frac{d}{2}-i\lambda,\frac{d+1}{2};-\frac{u}{2}\right)$$

$$(W_1 \cdot W_2) \mapsto -\frac{\partial^2 u}{\partial x_1 \partial x_2}, (X_1 \cdot W_2)(X_2 \cdot W_1) \mapsto \frac{\partial u}{\partial x_1}\frac{\partial u}{\partial x_2}$$

$$F_0^{\mathcal{D}}(u) = \int_{-\infty}^{+\infty} \mathrm{d}\lambda \frac{1}{\lambda^2 + \left(\frac{d}{2}-1\right)^2}\omega_{0,\lambda}(u) - \frac{\xi}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\frac{\partial \Omega_\lambda^{(0)}(u)}{\partial u}$$

$$F_1^{\mathcal{D}}(u) = \int_{-\infty}^{+\infty} \mathrm{d}\lambda \frac{1}{\lambda^2 + \left(\frac{d}{2}-1\right)^2}\omega_{1,\lambda}(u) + \frac{\xi}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\frac{\partial^2 \Omega_\lambda^{(0)}(u)}{\partial u^2}$$

$$F_0^{\mathcal{D}}(u) \underset{u\to\infty}{\sim} -\frac{1}{2\pi^{\frac{d+4}{2}}}\int_{-\infty}^{+\infty} \mathrm{d}\lambda \frac{\Gamma(1+i\lambda)(2\lambda+id)\sinh(\pi\lambda)\Gamma\left(\frac{d}{2}-i\lambda-1\right)}{(d-2)^2+4\lambda^2}\left(\frac{1}{2u}\right)^{\frac{d}{2}-i\lambda} + \frac{\Gamma(1-i\lambda)(2\lambda-id)\sinh(\pi\lambda)\Gamma\left(\frac{d}{2}+i\lambda-1\right)}{(d-2)^2+4\lambda^2}\left(\frac{1}{2u}\right)^{\frac{d}{2}+i\lambda}$$

$$F_0^{\mathcal{D}}(u) \underset{u\to\infty}{\sim} \frac{\Gamma\left(\frac{d+1}{2}\right)}{2\pi^{\frac{d+1}{2}}}\frac{1}{d-2}\frac{1}{u^{d-1}} + \mathcal{O}\left(\frac{1}{u^d}\right)$$

$$\frac{1}{\mathcal{C}_{d-1}^{(1)}}\left\langle A_i^{\mathcal{D}}(x_1,z_1)A_j^{\mathcal{D}}(x_2,z_2)\right\rangle \underset{z_1,z_2\to 0}{\sim} z_1^{d-2}z_2^{d-2}\frac{1}{|\vec{x}_{12}|^{2(d-1)}}\left(\delta_{ij} - 2\frac{\vec{x}_{12,i}\vec{x}_{12,j}}{|\vec{x}_{12}|^2}\right),$$

$$\frac{1}{\mathcal{C}_{d-1}^{(1)}}\left\langle A_z^{\mathcal{D}}(x_1,z_1)A_z^{\mathcal{D}}(x_2,z_2)\right\rangle \underset{z_1,z_2\to 0}{\sim} z_1^{d-1}z_2^{d-1}(-2(d-1))\frac{1}{|\vec{x}_{12}|^{2d}},$$

$$A_i \underset{z\to 0}{\sim} z^{d-2}j_i(\vec{x}), A_z \underset{z\to 0}{\sim} \mathcal{O}\left(z^{d-1}\right)$$

$$\int_{\lambda=\pm i\left(\frac{d}{2}-1\right)}^{\circlearrowleft} \Omega_\lambda^{(1)}(X_1,X_2;W_1,W_2) + (W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\int_{\lambda=\pm i\frac{d}{2}}^{\circlearrowleft}\frac{1}{\lambda^2+\frac{d^2}{4}}\Omega_\lambda^{(0)}(X_1,X_2)$$

$$id\mathrm{Res}_{\lambda=-i\left(\frac{d}{2}-1\right)}\left[\Omega_\lambda^{(1)}(X_1,X_2;W_1,W_2)\right] = (W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega^{(0)}_{-i\frac{d}{2}}(X_1,X_2).$$

$$F_0^{\mathcal{D}}(u) \underset{u\to\infty}{\sim} \frac{\Gamma\left(\frac{d+1}{2}\right)}{2\pi^{\frac{d+1}{2}}}\frac{1}{d-2}\left(\frac{1}{u^{d-1}} - \frac{d-1}{u^d}\right), F_1^{\mathcal{D}}(u) \underset{u\to\infty}{\sim} \frac{\Gamma\left(\frac{d+1}{2}\right)}{2\pi^{\frac{d+1}{2}}}\frac{1}{d-2}\left(\frac{1}{u^d} - \frac{d}{u^{d+1}}\right)$$

$$\Omega_\lambda^{(1)}(X_1,X_2;W_1,W_2) = \frac{i\lambda}{2\pi}\left[\Pi^{(1)}_{\frac{d}{2}+i\lambda}(X_1,X_2;W_1,W_2) - \Pi^{(1)}_{\frac{d}{2}-i\lambda}(X_1,X_2;W_1,W_2)\right]$$

$$F_0^{\mathrm{Hom}}(u)=\int_{\lambda=\pm i\left(\frac{d}{2}-1\right)}^{\circlearrowleft}\mathrm{d}\lambda\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\,\omega_{0,\lambda}(u)-\int_{\lambda=\pm i\frac{d}{2}}^{\circlearrowleft}\mathrm{d}\lambda\frac{\xi}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\,\frac{\partial\Omega_\lambda^{(0)}(u)}{\partial u}\,,$$

$$F_1^{\mathrm{Hom}}(u)=\int_{\lambda=\pm i\left(\frac{d}{2}-1\right)}^{\circlearrowleft}\mathrm{d}\lambda\frac{1}{\lambda^2+\left(\frac{d}{2}-1\right)^2}\,\omega_{1,\lambda}(u)+\int_{\lambda=\pm i\frac{d}{2}}^{\circlearrowleft}\mathrm{d}\lambda\frac{\xi}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\,\frac{\partial^2\Omega_\lambda^{(0)}(u)}{\partial u^2}\,.$$

$$F_0^{\mathrm{Hom}}(u)\underset{u\to\infty}{\sim}-\frac{\Gamma\left(\frac{d+1}{2}\right)}{2\pi^{\frac{d+1}{2}}(d-2)}\frac{1}{u^{d-1}}-\frac{1}{4\pi^{\frac{d}{2}}\Gamma\left(2-\frac{d}{2}\right)}\left[\left(1-\frac{d-2}{d}\xi\right)\log\left(\frac{u}{2}\right)+C\right]\frac{1}{u}$$

$$C:=-\frac{d-1}{d-2}+\left(1-\frac{d-2}{d}\xi\right)\left(\log(4)+\gamma_E+\psi^{(0)}\left(-\frac{d}{2}\right)-\frac{2}{d}\right)$$

$$\langle F^{AB}(X_1)F_{CD}(X_2)\rangle=\nabla^{1,[A}\nabla_{2,[C}K^{1,B]}K_{2,D]}(W_1^MW_2^N\langle A_M(X_1)A_N(X_2)\rangle).$$

$$\mathbb{T}_1^{AB,CD}=G_{12}^{AC}G_{12}^{BD}-G_{12}^{AD}G_{12}^{BC},$$
$$\mathbb{T}_2^{AB,CD}=V_1^{[A}G_{12}^{B][C}V_2^{D]},$$

$$G_{12}^{AC}=\eta^{AC}(X_1\cdot X_2)-X_2^AX_1^C,V_1=X_2+(X_1\cdot X_2)X_1,V_2=X_1+(X_1\cdot X_2)X_2$$

$$\langle F^{AB}(X_1)F^{CD}(X_2)\rangle=\alpha_{FS}(u)\mathbb{T}_1^{AB,CD}+\beta_{FS}(u)\mathbb{T}_2^{AB,CD}$$
$$\alpha_{FS}(u)=-\frac{2}{(1+u)^2}\left(F_1(u)+\frac{\partial F_0(u)}{\partial u}\right)$$
$$\beta_{FS}(u)=\frac{2}{(1+u)^2}\left(F_1(u)+\frac{\partial F_0(u)}{\partial u}\right)+\frac{1}{1+u}\left(\frac{\partial F_1(u)}{\partial u}+\frac{\partial^2F_0(u)}{\partial u^2}\right).$$

$$\alpha_{FS}^{\mathcal{D}}(u)\underset{u\to\infty}{\sim}\frac{\Gamma\left(\frac{d+1}{2}\right)}{\pi^{\frac{d+1}{2}}}\frac{1}{u^{d+2}}$$

$$\beta_{FS}^{\mathcal{D}}(u)\underset{u\to\infty}{\sim}\frac{d-2}{2}\frac{\Gamma\left(\frac{d+1}{2}\right)}{\pi^{\frac{d+1}{2}}}\frac{1}{u^{d+2}}$$

$$\alpha_{FS}^{\mathrm{Hom}}(u)\underset{u\to\infty}{\sim}-\frac{\Gamma\left(\frac{d+1}{2}\right)}{\pi^{\frac{d+1}{2}}}\frac{1}{u^{d+2}}+\frac{\Gamma\left(\frac{d-2}{2}\right)}{2\pi^{\frac{d+2}{2}}}\sin\left(\frac{\pi d}{2}\right)\frac{1}{u^4}$$

$$\beta_{FS}^{\mathrm{Hom}}(u)\underset{u\to\infty}{\sim}-\frac{d-2}{2}\frac{\Gamma\left(\frac{d+1}{2}\right)}{\pi^{\frac{d+1}{2}}}\frac{1}{u^{d+2}}+\frac{3\Gamma\left(\frac{d-4}{2}\right)}{4\pi^{\frac{d+2}{2}}}\sin\left(\frac{\pi d}{2}\right)\frac{1}{u^6}$$

$$\frac{\partial X_1^A}{\partial x_1^\alpha}\frac{\partial X_2^B}{\partial x_2^\beta}\eta_{AB}=\frac{\partial^2}{\partial x_1^\alpha\partial x_2^\beta}(X_1\cdot X_2)=-\frac{\partial^2u}{\partial x_1^\alpha\partial x_2^\beta}$$
$$\frac{\partial X_1^A}{\partial x_1^\alpha}X_{2,A}=\frac{\partial}{\partial x_1^\alpha}(X_1\cdot X_2)=-\frac{\partial u}{\partial x_1^\alpha}\frac{\partial X^A}{\partial x^\alpha}$$

$$\mathbb{J}_1^{0j,0k} = \frac{(\vec{x}_1 - \vec{x}_2)^2 + z_1^2 + z_2^2}{2z_1^3 z_2^3}\delta_{j,k},$$

$$\mathbb{J}_2^{0j,0k} = \frac{(\vec{x}_1 - \vec{x}_2)^2 + z_1^2 + z_2^2}{8z_1^5 z_2^5}\big[((\vec{x}_1 - \vec{x}_2)^4 - (z_1^2 - z_2^2)^2)\delta_{j,k} - 2((\vec{x}_1 - \vec{x}_2)^2 + z_1^2 + z_2^2)\big(x_{1,j} - x_{2,j}\big)\big(x_{1,k} - x_{2,k}\big)\big],$$

$$\lim_{z_1 \to 0} \mathbb{J}_1^{ab,cd} = \mathcal{O}\left(\frac{1}{z_1^4}\right),$$

$$\mathbb{J}_2^{ab,cd} = -\frac{(\vec{x}_1 - \vec{x}_2)^2 + z_1^2 + z_2^2}{2z_1^4 z_2^4}\big[(\vec{x}_{1,d} - \vec{x}_{2,d})(\vec{x}_{1,b} - \vec{x}_{2,b})\delta_{a,c} - (\vec{x}_{1,d} - \vec{x}_{2,d})(\vec{x}_{1,a} - \vec{x}_{2,a})\delta_{b,c} - (\vec{x}_{1,b} - \vec{x}_{2,b})(\vec{x}_{1,c} - \vec{x}_{2,c})\delta_{a,d} + (\vec{x}_{1,a} - \vec{x}_{2,a})(\vec{x}_{1,c} - \vec{x}_{2,c})\delta_{b,d}\big].$$

$$\begin{aligned}\int_{\mathbb{R}^d} \mathrm{d}^d\vec{x}\,\frac{z^{d+2\alpha}}{(z^2+\vec{x})^{d+\alpha}} &= z^{-d}\int_{\mathbb{R}^d} \mathrm{d}^d\vec{x}\,\frac{1}{\left(1+\frac{\vec{x}^2}{z^2}\right)^{d+\alpha}} \\ &= \int_{\mathbb{R}^d} \mathrm{d}^d\vec{x}\,\frac{1}{(1+\vec{x}^2)^{d+\alpha}} \\ &= \Omega_{d-1}\int_0^{+\infty} \mathrm{d}x\,\frac{x^{d-1}}{(1+x^2)^{d+\alpha}} \\ &= \pi^{\frac{d}{2}}\frac{\Gamma\left(\frac{d}{2}+\alpha\right)}{\Gamma(d+\alpha)}\end{aligned}$$

$$\Omega_d = \frac{2\pi^{\frac{d+1}{2}}}{\Gamma\left(\frac{d+1}{2}\right)}$$

$$\lim_{z\to 0}\frac{z^{d+2\alpha}}{(z^2+\vec{x})^{d+\alpha}} = \pi^{\frac{d}{2}}\frac{\Gamma\left(\frac{d}{2}+\alpha\right)}{\Gamma(d+\alpha)}\delta^d(\vec{x}) = \tilde{\kappa}_\nu\delta^d(\vec{x})$$

$$\begin{aligned}\Pi^{(0)}_{\frac{d}{2}+i\nu}(X,P) &= \sqrt{\mathcal{C}_\nu^{(0)}}\left(\frac{z}{z^2+\vec{x}_{ij}^2}\right)^{\frac{d}{2}+i\nu} \\ &= \sqrt{\mathcal{C}_\nu^{(0)}}\,\frac{z^{d+2\left(i\nu-\frac{d}{2}\right)}}{\left(z^2+\vec{x}_{ij}^2\right)^{d+\left(i\nu-\frac{d}{2}\right)}}z^{\frac{d}{2}-i\nu} \\ &\underset{z\to 0}{\sim} \sqrt{\mathcal{C}_\nu^{(0)}}z^{\frac{d}{2}-i\nu}\pi^{\frac{d}{2}}\frac{\Gamma(i\nu)}{\Gamma\left(\frac{d}{2}+i\nu\right)}\delta^d\left(\vec{x}_{ij}\right)\end{aligned}$$

$$\frac{\partial \frac{z^{d+2\alpha}}{(z^2+\vec{x})^{d+\alpha}}}{\partial z}\Pi^{(0)}_{\frac{d}{2}+i\nu}(X,P)=\sqrt{\mathcal{C}_\nu^{(0)}}\left(\frac{d}{2}+i\nu\right)\left[\frac{z^{\frac{d}{2}+i\nu-1}}{\left(z^2+\vec{x}_{ij}^2\right)^{d+\left(i\nu-\frac{d}{2}\right)}}-2\frac{z^{\frac{d}{2}+i\nu+1}}{\left(z^2+\vec{x}_{ij}^2\right)^{d+\left(i\nu+1-\frac{d}{2}\right)}}\right]$$

$$=\sqrt{\mathcal{C}_\nu^{(0)}}\left(\frac{d}{2}+i\nu\right)z^{\frac{d}{2}-i\nu-1}\left[\frac{z^{d+2\left(i\nu-\frac{d}{2}\right)}}{\left(z^2+\vec{x}_{ij}^2\right)^{d+\left(i\nu-\frac{d}{2}\right)}}-2\frac{z^{d+2\left(i\nu+1-\frac{d}{2}\right)}}{\left(z^2+\vec{x}_{ij}^2\right)^{d+\left(i\nu+1-\frac{d}{2}\right)}}\right]$$

$$\sim\sqrt{\mathcal{C}_\nu^{(0)}}\left(\frac{d}{2}+i\nu\right)z^{\frac{d}{2}-i\nu-1}\delta^d(\vec{x}_{ij})\pi^{\frac{d}{2}}\left[\frac{\Gamma(i\nu)}{\Gamma\left(\frac{d}{2}+i\nu\right)}-2\frac{\Gamma(1+i\nu)}{\Gamma\left(1+\frac{d}{2}+i\nu\right)}\right]$$

$$=\sqrt{\mathcal{C}_\nu^{(0)}}z^{\frac{d}{2}-i\nu-1}\delta^d(\vec{x}_{ij})\pi^{\frac{d}{2}}\left(\frac{d}{2}-i\nu\right)\frac{\Gamma(i\nu)}{\Gamma\left(\frac{d}{2}+i\nu\right)}$$

$$T_{12}^z(P_1,P_2,X_1)$$

$$\underset{z\to0}{\sim}z^{d+1}\mathcal{C}_\nu^{(0)}\tilde{\kappa}_\nu\left(\left(\frac{d}{2}+i\nu\right)\frac{\delta^d(\vec{x}_{1\tilde{1}})}{\left(\vec{x}_{2\tilde{1}}^2\right)^{\frac{d}{2}+i\nu}}+\left(\frac{d}{2}-i\nu\right)\frac{\delta^d(\vec{x}_{2\tilde{1}})}{\left(\vec{x}_{1\tilde{1}}^2\right)^{\frac{d}{2}+i\nu}}-(1\leftrightarrow2)+\mathcal{O}(z^2)\right)$$

$$+z^{d+1}z^{2i\nu}\mathcal{C}_\nu^{(0)}\left(\left(\frac{1}{\vec{x}_{1\tilde{1}}^2\vec{x}_{2\tilde{1}}^2}\right)^{\frac{d}{2}+i\nu}-(1\leftrightarrow2)+\mathcal{O}(z^2)\right)$$

$$\underset{z\to0}{\sim}z^{d+1}\left(\frac{\delta^d(\vec{x}_{1\tilde{1}})}{\left(\vec{x}_{2\tilde{1}}^2\right)^{\frac{d}{2}+i\nu}}-\frac{\delta^d(\vec{x}_{2\tilde{1}})}{\left(\vec{x}_{1\tilde{1}}^2\right)^{\frac{d}{2}+i\nu}}+\mathcal{O}(z^2)+\mathcal{O}\left(z^{2(1+i\nu)}\right)\right)$$

$$2i\nu\mathcal{C}_\nu^{(0)}\tilde{\kappa}_\nu=1$$

$$
\begin{aligned}
&\int_{\partial AdS} \frac{\mathrm{d}^d \vec{x}_{\tilde{1}}}{z^{d+1}} T_{12}^A\left(P_1, P_2, X_1\right) \Pi_{\frac{d}{2}+i\lambda}^{(0)}(X_1, P_5)\Big|_{X_1=\left(z, \vec{x}_{\tilde{1}}\right)} \underset{z\to 0}{\sim} \sqrt{\mathcal{C}_\lambda^{(0)}} \int_{\mathbb{R}^d} \mathrm{d}^d \vec{x}_{\tilde{1}} \\
&\left[\tilde{\kappa}_\lambda \delta^d\left(\vec{x}_{5\tilde{1}}\right) z^{\frac{d}{2}-i\lambda}+z^{\frac{d}{2}+i\lambda}\left(\left(\frac{1}{\vec{x}_{5\tilde{1}}^2}\right)^{\frac{d}{2}+i\lambda}+\mathcal{O}\left(z^2\right)\right)\right] \\
&\left[\frac{\delta^d(\vec{x}_{1\tilde{1}})}{\left(\vec{x}_{2\tilde{1}}^2\right)^{\frac{d}{2}+i\nu}}-\frac{\delta^d(\vec{x}_{2\tilde{1}})}{\left(\vec{x}_{1\tilde{1}}^2\right)^{\frac{d}{2}+i\nu}}+\mathcal{O}\left(z^2\right)+\mathcal{O}\left(z^{2(1+i\nu)}\right)\right] \\
&=\sqrt{\mathcal{C}_\lambda^{(0)}} \tilde{\kappa}_\lambda z^{\frac{d}{2}-i\lambda}\left(\frac{\delta^d(\vec{x}_{15})}{\left(\vec{x}_{25}^2\right)^{\frac{d}{2}+i\nu}}-\frac{\delta^d(\vec{x}_{25})}{\left(\vec{x}_{15}^2\right)^{\frac{d}{2}+i\nu}}+\mathcal{O}\left(z^2\right)\right) \\
&+\mathcal{O}\left(z^{\frac{d}{2}-i\lambda+2(1+i\nu)}\right) \\
&+\sqrt{\mathcal{C}_\lambda^{(0)}} z^{\frac{d}{2}+i\lambda} \frac{1}{\left(\vec{x}_{12}^2\right)^{\frac{d}{2}+i\nu}}\left(\frac{1}{\left(\vec{x}_{15}^2\right)^{\frac{d}{2}+i\lambda}}-\frac{1}{\left(\vec{x}_{25}^2\right)^{\frac{d}{2}+i\lambda}}\right) \\
&+\mathcal{O}\left(z^{\frac{d}{2}+i\lambda+2(1+i\nu)}\right)
\end{aligned}
$$

$$
\begin{aligned}
&\int_{\partial AdS} \frac{\mathrm{d}^d \vec{x}_{\tilde{2}}}{z^{d+1}} T_{34}^A\left(P_3, P_4, X_2\right) \Pi_{\frac{d}{2}-i\lambda}^{(0)}(X_2, P_5) \\
&\qquad \underset{z\to 0}{\sim} \sqrt{\mathcal{C}_{-\lambda}^{(0)}} \tilde{\kappa}_{-\lambda} z^{\frac{d}{2}+i\lambda}\left(\frac{\delta^d(\vec{x}_{35})}{\left(\vec{x}_{45}^2\right)^{\frac{d}{2}+i\nu}}-\frac{\delta^d(\vec{x}_{45})}{\left(\vec{x}_{35}^2\right)^{\frac{d}{2}+i\nu}}+\mathcal{O}\left(z^2\right)\right) \\
&\qquad +\mathcal{O}\left(z^{\frac{d}{2}+i\lambda+2(1+i\nu)}\right) \\
&\qquad +\sqrt{\mathcal{C}_{-\lambda}^{(0)}} z^{\frac{d}{2}-i\lambda} \frac{1}{\left(\vec{x}_{34}^2\right)^{\frac{d}{2}+i\nu}}\left(\frac{1}{\left(\vec{x}_{35}^2\right)^{\frac{d}{2}-i\lambda}}-\frac{1}{\left(\vec{x}_{45}^2\right)^{\frac{d}{2}-i\lambda}}\right) \\
&\qquad +\mathcal{O}\left(z^{\frac{d}{2}-i\lambda+2(1+i\nu)}\right)
\end{aligned}
$$

$$
\begin{aligned}
\mathcal{A}_{\mathcal{D}}^{\parallel}(z) \underset{z\to 0}{\sim} (-ie)^2 \frac{\xi}{\pi} \int_{-\infty}^{+\infty} \mathrm{d}\lambda \frac{\lambda^2 \mathcal{C}_\lambda^{(0)} \mathcal{C}_{-\lambda}^{(0)}}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\Big(&\#z^d+\#z^{d+2}+\ldots \\
&+\#z^{d+2+2i\nu}+\#z^{d+4+2i\nu}+\ldots \\
&+\#z^{d+4+4i\nu}+\#z^{d+6+4i\nu}+\ldots \\
&+\#z^{d\pm 2i\lambda}+\#z^{d+2\pm 2i\lambda}+\ldots \\
&+\#z^{d\pm 2i\lambda+2(1+i\nu)}+\#z^{d+2\pm 2i\lambda+2(1+i\nu)}+\ldots \\
&+\#z^{d\pm 2i\lambda+4(1+i\nu)}+\#z^{d+2\pm 2i\lambda+4(1+i\nu)}+\ldots\Big).
\end{aligned}
$$

$$(-ie^{z^{d+2+2i\nu}\otimes z^{d+4+4i\nu}\backslash z^{d\pm 2i\lambda}})^2\frac{\xi}{\pi}z^d\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{\lambda^2\mathcal{C}_{\lambda}^{(0)}\mathcal{C}_{-\lambda}^{(0)}}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\frac{1}{(\vec{x}_{12}^2\vec{x}_{34}^2)^{\frac{d}{2}+i\nu}}\Big[\tilde{\kappa}_{\lambda}\tilde{\kappa}_{-\lambda}\big(\delta^d(\vec{x}_{13})-\delta^d(\vec{x}_{14})-\delta^d(\vec{x}_{23})+\delta^d(\vec{x}_{24})\big)$$
$$+\int_{\mathbb{R}^d}\mathrm{d}^dx_5\left(\frac{1}{(\vec{x}_{15}^2)^{\frac{d}{2}+i\lambda}}-\frac{1}{(\vec{x}_{25}^2)^{\frac{d}{2}+i\lambda}}\right)\left(\frac{1}{(\vec{x}_{35}^2)^{\frac{d}{2}-i\lambda}}-\frac{1}{(\vec{x}_{45}^2)^{\frac{d}{2}-i\lambda}}\right)\big(1+\mathcal{O}(z^2)\big)\Big]$$

$\Delta_\nu\geq\frac{d-2}{2}$; indeed $d+2+2i\nu>0$ is true in any $d>0$ if $i\nu\geq-1$

$$\mathcal{A}_{\mathcal{D}}^{\parallel}\underset{z\to 0}{\sim}(-ie)^2\frac{\xi}{\pi}\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{\lambda^2}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\mathcal{C}_{-\lambda}^{(0)}\mathcal{C}_{\lambda}^{(0)}\frac{1}{\vec{x}_{12}^{2\left(\frac{d}{2}+i\nu\right)}\vec{x}_{34}^{2\left(\frac{d}{2}+i\nu\right)}}$$
$$\left[z^{d-2i\lambda}\tilde{\kappa}_{\lambda}\left(\frac{1}{\vec{x}_{13}^{2\left(\frac{d}{2}-i\lambda\right)}}-\frac{1}{\vec{x}_{14}^{2\left(\frac{d}{2}-i\lambda\right)}}-\frac{1}{\vec{x}_{23}^{2\left(\frac{d}{2}-i\lambda\right)}}+\frac{1}{\vec{x}_{24}^{2\left(\frac{d}{2}-i\lambda\right)}}\right)\right.$$
$$\left.+z^{d+2i\lambda}\tilde{\kappa}_{-\lambda}\left(\frac{1}{\vec{x}_{13}^{2\left(\frac{d}{2}+i\lambda\right)}}-\frac{1}{\vec{x}_{14}^{2\left(\frac{d}{2}+i\lambda\right)}}-\frac{1}{\vec{x}_{23}^{2\left(\frac{d}{2}+i\lambda\right)}}+\frac{1}{\vec{x}_{24}^{2\left(\frac{d}{2}+i\lambda\right)}}\right)\right]$$


$\mathcal{C}_{-\lambda}^{(0)}\tilde{\kappa}_{-\lambda}=\frac{i}{2\lambda}$, and that $\mathcal{C}_{\lambda}^{(0)}=\frac{1}{2\pi^{d/2}}\frac{\Gamma\left(\frac{d}{2}+i\lambda\right)}{\Gamma(1+i\lambda)}$ we have a λ-integral

$$(-ie)^2\frac{\xi}{\pi}\frac{i}{4\pi^{\frac{d}{2}}}\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{\lambda}{\left(\lambda^2+\frac{d^2}{4}\right)^2}\frac{\Gamma\left(\frac{d}{2}+i\lambda\right)}{\Gamma(1+i\lambda)}\frac{1}{\vec{x}_{i,j}^{2\left(\frac{d}{2}+i\lambda\right)}}z^{d+2i\lambda}$$

$z^{2i\lambda}=e^{-2i\lambda|\log z|}$ and $\log z<0$ given that $z\to 0$

$$\lambda=-i\frac{d}{2}\Longrightarrow z^{d+2i\lambda}\Big|_{\lambda=-i\frac{d}{2}\backslash\lambda=\mp i\frac{d}{2}}=z^{2d}\longrightarrow e^{z^{d\pm 2i\lambda+4(1+i\nu)}}\star e^{z^{d\pm 2i\lambda+4(1+i\nu)}}$$

$$\mathcal{A}_{\mathcal{D}}^{\parallel}(z)\supseteq z^{d+2i\lambda+2n(1+i\nu)}\Big|_{\lambda=-i\frac{d}{2}}=z^{2d+2n(1+i\nu)}\leq z^{2d}\longrightarrow 0$$



$$(X_1,X_2;W_1,W_2)=2\Pi_{\frac{d}{2}+i\nu_1}^{(0)}(X_1,X_2)(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Pi_{\frac{d}{2}+i\nu_2}^{(0)}(X_1,X_2)$$
$$2\Pi_{\frac{d}{2}+i\nu_2}^{(0)}(X_1,X_2)(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Pi_{\frac{d}{2}+i\nu_1}^{(0)}(X_1,X_2)$$
$$-(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\left[\Pi_{\frac{d}{2}+i\nu_1}^{(0)}(X_1,X_2)\Pi_{\frac{d}{2}+i\nu_2}^{(0)}(X_1,X_2)\right].$$

$$(X_1,X_2;W_1,W_2)= -\int_{-\infty}^{+\infty} \mathrm{d}\lambda \langle JJ\rangle_{\nu_1,\nu_2}(\lambda)\Omega_\lambda^{(1)}(X_1,X_2;W_1,W_2)$$
$$-\int_{-\infty}^{+\infty} \mathrm{d}\lambda \langle JJ\rangle^{\parallel}_{\nu_1,\nu_2}(\lambda)(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega_\lambda^{(0)}(X_1,X_2)$$

$$\langle JJ\rangle_{\nu_1,\nu_2}(\lambda) * \langle JJ\rangle^{\parallel}_{\nu_1,\nu_2}(\lambda)$$

$$(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega_\lambda^{(0)}(X_1,X_2)$$

$$\langle JJ\rangle^{\parallel}_{\nu_1,\nu_2}(\lambda) \propto \left(\lambda^2+\frac{d^2}{4}\right)^{-1}$$

$$\langle JJ\rangle^{\parallel}_{\nu_1,\nu_2}(\lambda) \triangleq \left(\lambda^2+\frac{d^2}{4}\right)^{-1}$$

$$\langle JJ\rangle_{\nu_1,\nu_2}(\lambda) = -\frac{2\pi^{-\frac{d}{2}}\,(d+i\nu_1+i\nu_2+2)_{-\frac{d}{2}}}{\left(\frac{d}{2}+i\nu_1+i\nu_2+1\right)\left(\frac{d}{2}+i\nu_1+1\right)_{\frac{d}{2}}\left(\frac{d}{2}+i\nu_2+1\right)_{-\frac{d}{2}}}\Bigg[$$
$$\frac{{}_7F_6\left(\begin{matrix}\frac{d+2}{2},\frac{d+2}{2}+i\nu_1,\frac{1+(\nu_1+\nu_2)}{2},1+i\frac{\nu_1+\nu_2}{2},\frac{d+2}{4}+\frac{i\lambda}{2}+i\frac{\nu_1+\nu_2}{2},\frac{d+2}{2}+i\nu_2,\frac{d+2}{2}+i\nu_1+i\nu_2\\ 1+i\nu_1,\frac{d+2}{2}+i\frac{\nu_1+\nu_2}{2},\frac{d+3}{2}+i\frac{\nu_1+\nu_2}{2},\frac{d}{4}+\frac{i\lambda}{2}+i\frac{\nu_1+\nu_2}{2}+\frac{3}{2},1+i\nu_2,1+i\nu_1+i\nu_2\end{matrix};1\right)}{\frac{d}{2}+i\lambda+i\nu_1+i\nu_2+1}$$
$$+(\lambda\to-\lambda)\Bigg].$$


$$Z=\sum_{i=1}^{q+1} a_{\;{}_{q+1}F_q\left(\begin{smallmatrix}a_1,\ldots,a_{q+1}\\ b_1,\ldots,b_q\end{smallmatrix};z\right)} - \sum_{i=1}^{q} b_{\;{}_{q+1}F_q\left(\begin{smallmatrix}a_1,\ldots,a_{q+1}\\ b_1,\ldots,b_q\end{smallmatrix};z\right)}$$

$${}_{q+1}F_q\left(\begin{matrix}a_1,\ldots,a_{q+1}\\ b_1,\ldots,b_q\end{matrix};1\right)$$
$$=-\frac{1}{Z}\sum_{j=1}^{q}\frac{\Pi_{k=1}^{q+1}(b_j-a_k)}{b_j\Pi_{k=1,k\neq j}^{q}(b_j-b_k)^{q+1}}F_q\left(\begin{matrix}a_1,\ldots,a_{q+1}\\ b_1,\ldots,b_j+1,\ldots,b_q\end{matrix};1\right)$$

$$\langle JJ\rangle_{\nu,\nu}(\lambda)=\frac{1}{(d-1)(d-2)(d-3)}\langle\widetilde{J}\widetilde{J}\rangle_{\nu,\nu}(\lambda)$$
$$\langle JJ\rangle_{\nu,-\nu}(\lambda)=\frac{1}{(d-1)(d-2)(d-3)}\langle\widetilde{J}\widetilde{J}\rangle_{\nu,-\nu}(\lambda)$$

$$\langle JJ\rangle_{\nu_1,\nu_2}(\lambda) \underset{d\to3}{\sim} \frac{-\frac{1}{24\pi^2}\lambda^2+c(\nu_1,\nu_2)}{d-3}+\overline{\langle JJ\rangle}_{\nu,\pm\nu}(\lambda)+\mathcal{O}(d-3),$$

$$T_\nu(X_1,X_2;W_1,W_2)=$$



$$T_\nu(X_1,X_2;W_1,W_2)=(W_1\cdot W_2)\mathcal{T}_\nu\delta^{d+1}(X_1,X_2)$$

$$T_\nu(X_1,X_2;W_1,W_2)=\mathcal{T}_\nu\left(\int_{-\infty}^{+\infty}\mathrm{d}\lambda\Omega_\lambda^{(1)}(X_1,X_2;W_1,W_2)+\int_{-\infty}^{+\infty}\mathrm{d}\lambda\frac{1}{\lambda^2+\frac{d^2}{4}}(W_1\cdot\nabla_1)(W_2\cdot\nabla_2)\Omega_\lambda^{(0)}(X_1,X_2)\right)$$

$$\mathcal{T}_\nu=\lim_{X_4\to X_3}\Pi_{\frac{d}{2}+i\nu}^{\left(\lambda^2+\frac{d^2}{4}\right)^{-1}}(X_4,X_3)=\int_{-\infty}^{\infty}d\nu'\frac{1}{\nu'^2-\nu^2}\Omega_{\nu'}^{AdS}(x,x)$$

$$\Omega_\nu^{\left(\lambda^2+\frac{d^2}{4}\right)^{-1}}(x,x)=\frac{\Gamma\left(\frac{d}{2}\right)}{4\pi^{\frac{d}{2}+1}\Gamma(d)}\frac{\Gamma\left(\frac{d}{2}+i\nu\right)\Gamma\left(\frac{d}{2}-i\nu\right)}{\Gamma(i\nu)\Gamma(-i\nu)}$$

$$\nu'=\nu\text{ and }\nu'=i\left(\frac{d}{2}+n\right)_{n\in\mathbb{N}\cdot}$$

$$\mathcal{T}_\nu=\frac{\Gamma\left(\frac{d}{2}\right)}{4\pi^{\frac{d}{2}+1}\Gamma(d)}\Bigg[-i\sinh(\pi\nu)\Gamma\left(\frac{d}{2}-i\nu\right)\Gamma\left(\frac{d}{2}+i\nu\right)$$
$$+\frac{2i\pi}{\nu(d^2+4\nu^2)}\frac{\Gamma(d)}{\Gamma\left(-\frac{d}{2}\right)\Gamma\left(\frac{d}{2}\right)}\Big((d+2i\nu)_3F_2\left(\frac{d}{2}+1,d,\frac{d}{2}-i\nu;\frac{d}{2},\frac{d}{2}-i\nu+1;1\right)$$
$$-(d-2i\nu)_3F_2\left(\frac{d}{2}+1,d,\frac{d}{2}+i\nu;\frac{d}{2},\frac{d}{2}+i\nu+1;1\right)\Big)\Bigg]$$

$$\Pi_{\frac{d}{2}-i\nu}^{\left(\lambda^2+\frac{d^2}{4}\right)^{-1}}(X_4,X_3)=\Pi_{\frac{d}{2}+i\nu}^{\left(\lambda^2+\frac{d^2}{4}\right)^{-1}}(X_4,X_3)+\frac{2i\pi}{\nu}\Omega_\nu^{\left(\lambda^2+\frac{d^2}{4}\right)^{-1}}(x_4,x_3)$$

$$\Longrightarrow\lim_{X_4\to X_3}\Pi_{\frac{d}{2}-i\nu}^{\left(\lambda^2+\frac{d^2}{4}\right)^{-1}}(X_4,X_3)=\mathcal{T}_\nu+\frac{\Gamma\left(\frac{d}{2}\right)}{4\pi^{\frac{d}{2}+1}\Gamma(d)}\left[2i\sinh(\pi\nu)\Gamma\left(\frac{d}{2}\pm i\nu\right)\right]=\mathcal{T}(-\nu)$$

$$\mathcal{T}_\nu\underset{d\to3}{\sim}\frac{c_T(\nu)}{3-d}+\mathcal{O}(d-3)$$

$$c_T(\nu)=\frac{\cosh(\pi\nu)}{8\pi^3}\Gamma\left(\frac{3}{2}\pm i\nu\right)=\frac{\nu^2+\frac{1}{4}}{8\pi^2}.$$

$$\int_{Y_1}\int_{Y_2}\langle A(X_1)A(Y_1)\rangle B_{\nu_1,\nu_2}^{(1)}(Y_1,Y_2)\langle A(Y_2)A(X_2)\rangle,$$

$$\Gamma(a\pm ib)\underset{b\to\infty}{\sim}2\pi e^{-\pi b}b^{2a-1}\text{ valid for }|\mathrm{Arg}(b)|_{\nu'^{d-2}}\leq\pi-\epsilon$$

$$\Psi_L(2,1,4)=\begin{pmatrix}u_1&u_2&u_3&\nu\\d_1&d_2&d_3&e\end{pmatrix}_L,\Psi_R(1,2,4)=\begin{pmatrix}u_1&u_2&u_3&\nu\\d_1&d_2&d_3&e\end{pmatrix}_R$$

$$H_L(2,1,4) = \begin{pmatrix} \chi_1^u & \chi_2^u & \chi_3^u & \chi^\nu \\ \chi_1^d & \chi_2^d & \chi_3^d & \chi^e \end{pmatrix}_L, H_R(1,2,4) = \begin{pmatrix} \chi_1^u & \chi_2^u & \chi_3^u & \chi^\nu \\ \chi_1^d & \chi_2^d & \chi_3^d & \chi^e \end{pmatrix}_R$$

$$\langle \chi_L^\nu \rangle = \kappa_L, \langle \chi_R^\nu \rangle = \kappa_R$$

$$\Sigma_L(1,1,15) = \frac{1}{\sqrt{2}} \begin{pmatrix} \frac{\mathcal{O}_3}{\sqrt{2}} + \frac{\mathcal{O}_8}{\sqrt{6}} + \frac{N}{\sqrt{12}} & \mathcal{O}_{12} & \mathcal{O}_{13} & U_1 \\ \mathcal{O}_{12}^c & -\frac{\mathcal{O}_3}{\sqrt{2}} + \frac{\mathcal{O}_8}{\sqrt{6}} + \frac{N}{\sqrt{12}} & \mathcal{O}_{23} & U_2 \\ \mathcal{O}_{13}^c & \mathcal{O}_{23}^c & -\sqrt{\frac{2}{3}}\mathcal{O}_8 + \frac{N}{\sqrt{12}} & U_3 \\ U_1^c & U_2^c & U_3^c & -\frac{\sqrt{3}}{2}N \end{pmatrix}_L$$

$$\Omega_{L,R}(1,1,10) = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}\mathcal{S}_{11} & \mathcal{S}_{12} & \mathcal{S}_{13} & D_1 \\ \mathcal{S}_{12} & \sqrt{2}\mathcal{S}_{22} & \mathcal{S}_{23} & D_2 \\ \mathcal{S}_{13} & \mathcal{S}_{23} & \sqrt{2}\mathcal{S}_{33} & D_3 \\ D_1 & D_2 & D_3 & \sqrt{2}E^- \end{pmatrix}_{L,R}$$

$$\Omega_{L,R}(1,1,10) = D_{L,R}\left(3,1,-\frac{1}{3}\right) + E_{L,R}^-(1,1,-1) + \mathcal{S}_{L,R}\left(6,1,\frac{1}{3}\right)$$

$$\Psi_L \leftrightarrow \Psi_R, \Omega_L \leftrightarrow \Omega_R, \Sigma_L \leftrightarrow \Sigma_R^c, H_L \leftrightarrow H_R$$

$$\begin{aligned} \mathcal{L}_{\text{Yuk}} = {} & \sqrt{2}Y_{15}^\dagger\{\text{Tr}(\Sigma_L\Psi_L^T H_L^*) + \text{Tr}(\bar{\Sigma}_L\Psi_R^T H_R^*)\} \\ & -\sqrt{2}Y_{10}\{\text{Tr}(\bar{\Psi}_L\tilde{H}_L\Omega_R) + \text{Tr}(\bar{\Psi}_R\tilde{H}_R\Omega_L)\} \\ & +M_{10}\bar{\Omega}_L\Omega_R + M_{15}(\overline{\Sigma^c})_R\Sigma_L + \text{ h.c.} \end{aligned}$$

$$\Psi_L \to V_L\Psi_L V_c^T, \Psi_R \to V_R\Psi_R V_c^T, \Sigma_L \to V_c\Sigma_L V_c^\dagger, \Omega_{L,R} \to V_c\Omega_{L,R}V_c^T$$
$$H_L \to V_L H_L V_c^T, H_R \to V_R H_R V_c^T$$

$$M_{10}^\dagger = M_{10}, M_{15}^\dagger = M_{15}$$

$$\begin{aligned} \mathcal{L}_{\text{Yuk}} = {} & Y_{15}^\dagger\Big(U_L^c u_L\chi_L^{\nu*} + U_L^c d_L\chi_L^{e*} + U_L e_L\chi_L^{d*} + U_L\nu_L\chi_L^{u*} - \frac{\sqrt{3}}{2}N_L\nu_L\chi_L^{\nu*} - \frac{\sqrt{3}}{2}N_L e_L\chi_L^{e*} \\ & + \frac{1}{2\sqrt{3}}N_L d_L\chi_L^{d*} + \frac{1}{2\sqrt{3}}N_L u_L\chi_L^{u*} + \sqrt{2}\mathcal{O}_{L\beta}^\alpha d_{L\alpha}\chi_L^{d*\beta} + \sqrt{2}\mathcal{O}_{L\beta}^\alpha u_{L\alpha}\chi_L^{u*\beta}\Big) \\ & + Y_{15}^\dagger\Big(\bar{U}_L u_R\chi_R^{\nu*} + \bar{U}_L d_R\chi_R^{e*} + \overline{U_L^c}e_R\chi_R^{d*} + \overline{U_L^c}\nu_R\chi_R^{u*} - \frac{\sqrt{3}}{2}\overline{N_L}\nu_R\chi_R^{\nu*} - \frac{\sqrt{3}}{2}\overline{N_L}e_R\chi_R^{e*} \\ & + \frac{1}{2\sqrt{3}}\overline{N_L}d_R\chi_R^{d*} + \frac{1}{2\sqrt{3}}\overline{N_L}u_R\chi_R^{u*} + \sqrt{2}\overline{\mathcal{O}}_{L\beta}^\alpha d_{R\alpha}\chi_R^{d*\beta} + \sqrt{2}\overline{\mathcal{O}}_{L\beta}^\alpha u_{R\alpha}\chi_R^{u*\beta}\Big) \\ & + Y_{10}\big(\bar{d}_L D_R\chi_L^{\nu*} - \bar{u}_L D_R\chi_L^{e*} + \sqrt{2}\bar{e}_L E_R\chi_L^{\nu*} - \sqrt{2}\bar{\nu}_L E_R\chi_L^{e*} + \bar{e}_L D_R\chi_L^{u*} - \bar{\nu}_L D_R\chi_L^{d*} \\ & - \bar{u}_L^\alpha(\mathcal{S}_R)_{\alpha\beta}\chi_L^{d*\beta} + \bar{d}_L^\alpha(\mathcal{S}_R)_{\alpha\beta}\chi_L^{u*\beta}\big) + Y_{10}\big(\bar{d}_R D_L\chi_R^{\nu*} - \bar{u}_R D_L\chi_R^{e*} + \sqrt{2}\bar{e}_R E_L\chi_R^{\nu*} \\ & - \sqrt{2}\bar{\nu}_R E_L\chi_R^{e*} + \bar{e}_R D_L\chi_R^{u*} - \bar{\nu}_R D_L\chi_R^{d*} - \bar{u}_R^\alpha(\mathcal{S}_L)_{\alpha\beta}\chi_R^{d*\beta} + \bar{d}_R^\alpha(\mathcal{S}_L)_{\alpha\beta}\chi_R^{u*\beta}\big) \\ & + M_{10}\big(\bar{D}_L D_R + \bar{E}_L E_R + \overline{\mathcal{S}}_L\mathcal{S}_R\big) + M_{15}\Big(U_L^c U_L + \frac{1}{2}N_L N_L + \frac{1}{2}(\mathcal{O}_L)_\alpha^\beta(\mathcal{O}_L)_\beta^\alpha\Big) + \text{ h.c.} \end{aligned}$$

$$\mathcal{L}_{\text{mass}} = (\bar{f}_L \bar{F}_L)\mathcal{M}_f \begin{pmatrix} f_R \\ F_R \end{pmatrix},$$

$$\mathcal{M}_u = \begin{pmatrix} 0 & Y_{15}\kappa_L \\ Y_{15}^\dagger \kappa_R & M_{15}^\dagger \end{pmatrix}, \mathcal{M}_d = \begin{pmatrix} 0 & Y_{10}\kappa_L \\ Y_{10}^\dagger \kappa_R & M_{10} \end{pmatrix}, \mathcal{M}_e = \begin{pmatrix} 0 & \sqrt{2}Y_{10}\kappa_L \\ \sqrt{2}Y_{10}^\dagger \kappa_R & M_{10} \end{pmatrix}.$$

$$\mathcal{M}_{\mathcal{O}} = M_{15}^\dagger, \mathcal{M}_{\mathcal{S}} = M_{10}$$

mass matrices for $M_{u,d,e}$ given the approximation $\left(Y_{15}^\dagger \kappa_R M_{15}^{-1}\right) \ll 1$ and $\left(Y_{10}^\dagger \kappa_R M_{10}^{-1}\right) \ll 1$

$$\begin{aligned} M_u &\simeq -\left(Y_{15} M_{15}^{-1} Y_{15}^\dagger\right)\kappa_L \kappa_R \\ M_d &\simeq -\left(Y_{10} M_{10}^{-1} Y_{10}^\dagger\right)\kappa_L \kappa_R \\ M_e &\simeq -2\left(Y_{10} M_{10}^{-1} Y_{10}^\dagger\right)\kappa_L \kappa_R \end{aligned}$$

$$\left(Y_{15}^\dagger \kappa_R M_{15}^{-1}\right) \ll 1 \text{ and } \left(Y_{10}^\dagger \kappa_R M_{10}^{-1}\right) \ll 1$$

$$\begin{aligned} V = &-\mu_L^2 \text{Tr}\left(H_L^\dagger H_L\right) - \mu_R^2 \text{Tr}\left(H_R^\dagger H_R\right) + \lambda_1 \left\{\left(\text{Tr}\left(H_L^\dagger H_L\right)\right)^2 + \left(\text{Tr}\left(H_R^\dagger H_R\right)\right)^2\right\} \\ &+\lambda_2 \left\{\text{Tr}\left(H_L^\dagger H_L H_L^\dagger H_L\right) + \text{Tr}\left(H_R^\dagger H_R H_R^\dagger H_R\right)\right\} + \lambda_3 \text{Tr}\left(H_L^\dagger H_L\right)\text{Tr}\left(H_R^\dagger H_R\right) \\ &+\lambda_4 \text{Tr}\left(H_L^\dagger H_L H_R^\dagger H_R\right) + \lambda_5 \left\{\text{Tr}\left(H_L^T \tilde{H}_L^* H_R^\dagger \tilde{H}_R\right) + \text{Tr}\left(H_R^T \tilde{H}_R^* H_L^\dagger \tilde{H}_L\right)\right\} \\ &+\left\{\lambda_6 H_L^{\alpha a} H_L^{\beta b} H_R^{\gamma c} H_R^{\delta d} \epsilon_{\alpha\beta\gamma\delta} \epsilon_{ab} \epsilon_{cd} + \text{ h.c.}\right\} \end{aligned}$$

$$\begin{aligned} \mu_L^2 &= 2(\lambda_1 + \lambda_2)\kappa_L^2 + (\lambda_3 + \lambda_4)\kappa_R^2 \\ \mu_R^2 &= 2(\lambda_1 + \lambda_2)\kappa_R^2 + (\lambda_3 + \lambda_4)\kappa_L^2 \end{aligned}$$

$$W_{L\mu}^\pm, W_{R\mu}^\pm, Z_\mu, Z'_\mu, X_\mu$$

$$\chi_L^\nu = \frac{\sigma_L + i a_L}{\sqrt{2}} + \kappa_L, \chi_R^\nu = \frac{\sigma_R + i a_R}{\sqrt{2}} + \kappa_R$$

$$M_{\chi_{L,R}^u}^2 = \begin{pmatrix} -\lambda_4 \kappa_R^2 & \lambda_4 \kappa_L \kappa_R \\ \lambda_4 \kappa_L \kappa_R & -\lambda_4 \kappa_L^2 \end{pmatrix}$$

$$\chi_1^u = \frac{(\kappa_R \chi_L^u - \kappa_L \chi_R^u)}{\sqrt{\kappa_L^2 + \kappa_R^2}}, \chi_2^u = \frac{(\kappa_L \chi_L^u + \kappa_R \chi_R^u)}{\sqrt{\kappa_L^2 + \kappa_R^2}}$$

$$M_{\chi_1^u}^2 = -\lambda_4(\kappa_L^2 + \kappa_R^2), M_{\chi_2^u}^2 = 0$$

$$M_{\chi_{L,R}^d}^2 = \begin{pmatrix} -2\lambda_2 \kappa_L^2 - \lambda_4 \kappa_R^2 & -2\lambda_5 \kappa_L \kappa_R \\ -2\lambda_5 \kappa_L \kappa_R & -2\lambda_2 \kappa_R^2 - \lambda_4 \kappa_L^2 \end{pmatrix}.$$

$$M_{\sigma_{L,R}}^2 = \begin{pmatrix} 4(\lambda_1 + \lambda_2)\kappa_L^2 & 2(\lambda_3 + \lambda_4)\kappa_L \kappa_R \\ 2(\lambda_3 + \lambda_4)\kappa_L \kappa_R & 4(\lambda_1 + \lambda_2)\kappa_R^2 \end{pmatrix}.$$

$$\mathcal{L}_{\text{kinetic}} = \text{Tr}\left\{\left(D_\mu H_L\right)^\dagger \left(D^\mu H_L\right)\right\} + \text{Tr}\left\{\left(D_\mu H_R\right)^\dagger \left(D^\mu H_R\right)\right\}$$

$$D_\mu H_L = \partial_\mu H_L - i g_{2L}(\vec{T}_L \cdot \vec{W}_{L\mu}) H_L - i g_4 H_L (\vec{\lambda} \cdot \vec{G}_\mu)^T$$
$$D_\mu H_R = \partial_\mu H_R - i g_{2R}(\vec{T}_R \cdot \vec{W}_{R\mu}) H_R - i g_4 H_R (\vec{\lambda} \cdot \vec{G}_\mu)^T$$

$$M^2_{W_L^{\mu\pm}} = \frac{1}{2} g_{2L}^2 \kappa_L^2$$
$$M^2_{W_R^{\mu\pm}} = \frac{1}{2} g_{2R}^2 \kappa_R^2$$
$$M^2_{X^\mu} = \frac{1}{2} g_4^2 (\kappa_L^2 + \kappa_R^2)$$

$$M^2_{Z-Z'} = \frac{1}{2}\begin{pmatrix} (g_Y^2 + g_{2L}^2)\kappa_L^2 & g_Y^2 \sqrt{\dfrac{g_Y^2 + g_{2L}^2}{g_{2R}^2 - g_Y^2}}\,\kappa_L^2 \\ g_Y^2 \sqrt{\dfrac{g_Y^2 + g_{2L}^2}{g_{2R}^2 - g_Y^2}}\,\kappa_L^2 & \dfrac{g_Y^4}{g_{2R}^2 - g_Y^2}\kappa_L^2 + \dfrac{g_{2R}^4}{g_{2R}^2 - g_Y^2}\kappa_R^2 \end{pmatrix}.$$

$$\frac{1}{g_Y^2} = \frac{1}{g_{2R}^2} + \frac{2}{3g_4^2}$$

$$A_\mu = \frac{\sqrt{3} g_4 g_{2R} W^0_{\mu L} + \sqrt{3} g_4 g_{2L} W^0_{\mu R} + \sqrt{2} g_{2L} g_{2R} G^{15}_\mu}{\sqrt{2 g_{2L}^2 g_{2R}^2 + 3 g_4^2 (g_{2L}^2 + g_{2R}^2)}}$$
$$Z_{L\mu} = \frac{g_{2L}(3g_4^2 + 2g_{2R}^2) W^0_{\mu L} - 3 g_4^2 g_{2R} W^0_{\mu R} - \sqrt{6} g_{2R}^2 g_4 G^{15}_\mu}{\sqrt{(3g_4^2 + 2g_{2R}^2)\left(2 g_{2L}^2 g_{2R}^2 + 3 g_4^2 (g_{2L}^2 + g_{2R}^2)\right)}}$$
$$Z_{R\mu} = \frac{-\sqrt{2} g_{2R} W^0_{\mu R} + \sqrt{3} g_4 G^{15}_\mu}{\sqrt{3 g_4^2 + 2 g_{2R}^2}}$$

$$\mathcal{L}_{\text{kin}} = i\bar{\psi}_L \not{D} \psi_L + i\bar{\psi}_R \not{D} \psi_R + i\bar{\Sigma}_L \not{D} \Sigma_L + i\bar{\Omega}_L \not{D} \Omega_L + i\bar{\Omega}_R \not{D} \Omega_R,$$

$$D_\mu \Sigma_L = \partial_\mu \Sigma_L - i g_4 [\vec{\lambda} \cdot \vec{G}_\mu, \Sigma_L]$$
$$D_\mu \Omega_{L,R} = \partial_\mu \Omega_{L,R} - i g_4 \vec{\lambda} \cdot \vec{G}_\mu \Omega_{L,R} - i g_4 \Omega_{L,R} (\vec{\lambda} \cdot \vec{G}_\mu)^T$$


$$\mathcal{L}_{X_\mu} = \frac{g_4}{\sqrt{2}} \Big[(\bar{u}_L \gamma^\mu \nu_L + \bar{d}_L \gamma^\mu e_L) X_\mu + \sqrt{2} \bar{D}_L \gamma^\mu E_L X_\mu + \overline{S}_L^{\alpha\beta} \gamma^\mu D_{L\beta} X_{\mu\alpha} + \text{L} \leftrightarrow \text{R}$$
$$+ \frac{1}{\sqrt{3}} \overline{N_L} \gamma^\mu U_L^c X_\mu + \frac{1}{\sqrt{3}} \overline{N_R^c} \gamma^\mu U_R X_\mu + \frac{1}{\sqrt{2}} \overline{(\mathcal{O}_L)^\alpha_\beta} \gamma^\mu U^c_{L\alpha} X^\beta_\mu + + \frac{1}{\sqrt{2}} \overline{(\mathcal{O}_R^c)^\alpha_\beta} \gamma^\mu U_{R\alpha} X^\beta_\mu \Big] + \text{ h.c.}$$

$$m_t(m_t) = 163.6\text{GeV}, \quad m_c(m_t) = 0.620\text{GeV}, \quad m_u(m_t) = 1.30\text{MeV}$$
$$m_b(m_t) = 2.78\text{GeV}, \quad m_s(m_t) = 55.0\text{MeV}, \quad m_d(m_t) = 2.70\text{MeV}$$
$$m_\tau(m_t) = 1.78\text{GeV}, \quad m_\mu(m_t) = 105.7\text{MeV}, \quad m_e(m_t) = 0.511\text{MeV}$$

$$\frac{m_\tau}{m_b} = 1.47, \frac{m_\mu}{m_s} = 3.97, \frac{m_e}{m_d} = 0.39 \ (\mu = 5.2 \times 10^{13}\text{GeV})$$

$m_f(\mu_P)$	[GeV]

m_t	8.50×10^{1}
m_b	1.19
m_τ	1.76
m_c	2.89×10^{-1}
m_μ	1.04×10^{-1}
m_s	2.63×10^{-2}
m_d	1.29×10^{-3}
m_u	6.06×10^{-4}
m_e	5.05×10^{-4}



$$\Delta M_d \simeq \left(\frac{11}{4}\right)\frac{Y_{15}M_{15}Y_{15}^{\dagger}}{16\pi^2}(2\lambda_5\kappa_L\kappa_R)F\left(M_{153}^2, M_{\chi_1^d}^2, M_{\chi_2^d}^2\right)$$

$$\Delta M_e \simeq (3)\frac{Y_{15}M_{15}Y_{15}^{\dagger}}{16\pi^2}(2\lambda_5\kappa_L\kappa_R)F\left(M_{153}^2, M_{\chi_1^d}^2, M_{\chi_2^d}^2\right)$$

$$F(x,y,z) = \frac{x\log\, x}{(x-y)(x-z)} + \frac{y\log\, y}{(y-x)(y-z)} + \frac{z\log\, z}{(z-x)(z-y)}.$$

$$Y_t(\mu_P) = \frac{|(Y_{15})_{33}|^2\kappa_R}{\sqrt{|(M_{15})_3|^2 + |(Y_{15})_{33}|^2\kappa_R^2}},$$

$$F(x,y,z) \simeq \frac{y\log\left(\frac{x}{y}\right) - z\log\left(\frac{x}{z}\right)}{x(y-z)}, x \gg y,z$$

$$F(x,y,z) \simeq \frac{1}{x}\left(\log\left(\frac{x}{y}\right) - 1\right), x \gg y = z$$

$$M_d = \begin{pmatrix} m_{11} & m_{12} & m_{13} \\ m_{12}^* & m_{22} & m_{23} \\ m_{13}^* & m_{23}^* & m_{33} \end{pmatrix} + \frac{11}{5} m_0 \begin{pmatrix} |\delta_1|^2 & \delta_1\delta_2^* & \delta_1\delta_3^* \\ \delta_1^*\delta_2 & |\delta_2|^2 & \delta_2\delta_3^* \\ \delta_1^*\delta_3 * & \delta_2^*\delta_3 & |\delta_3|^2 \end{pmatrix}$$

$$M_e = 2\begin{pmatrix} m_{11} & m_{12} & m_{13} \\ m_{12}^* & m_{22} & m_{23} \\ m_{13}^* & m_{23}^* & m_{33} \end{pmatrix} + \frac{12}{5} m_0 \begin{pmatrix} |\delta_1|^2 & \delta_1\delta_2^* & \delta_1\delta_3^* \\ \delta_1^*\delta_2 & |\delta_2|^2 & \delta_2\delta_3^* \\ \delta_1^*\delta_3^* & \delta_2^*\delta_3 & |\delta_3|^2 \end{pmatrix}$$

$$\delta_i = (Y_{15})_{i3}$$

$$m_0 = \left(\frac{5}{4}\right)\frac{(M_{15})_3}{16\pi^2}(2\lambda_5\kappa_L\kappa_R)F\left(M_{15_3}^2, M_{\chi_1^d}^2, M_{\chi_2^d}^2\right)$$

$$M_e = \begin{pmatrix} m_e & & \\ & m_\mu & \\ & & m_\tau \end{pmatrix}, M_d = \begin{pmatrix} \frac{m_e}{2} + m_0|\delta_1|^2 & m_0\delta_1\delta_2^* & m_0\delta_1\delta_3^* \\ m_0\delta_1^*\delta_2 & \frac{m_\mu}{2} + m_0|\delta_2|^2 & m_0\delta_2\delta_3^* \\ m_0\delta_1^*\delta_3 & m_0\delta_2^*\delta_3 & \frac{m_\tau}{2} + m_0|\delta_3|^2 \end{pmatrix}$$

$$m_\tau, |\delta_3|^2 m_0 \gg m_\mu, |\delta_2|^2 m_0 \gg m_e, |\delta_1|^2 m_0$$

$$m_b \simeq \frac{m_\tau}{2} + |\delta_3|^2 m_0$$

$$m_s m_b \simeq \frac{m_\mu m_\tau}{4} + \frac{1}{2}|\delta_2|^2 m_0 m_\tau + \frac{1}{2}|\delta_3|^2 m_0 m_\mu$$

$$m_d m_s m_b \simeq \frac{m_e m_\mu m_\tau}{8} + \frac{1}{4}|\delta_3|^2 m_0 m_e m_\mu + \frac{1}{4}|\delta_2|^2 m_0 m_e m_\tau + \frac{1}{4}|\delta_1|^2 m_0 m_\mu m_\tau$$

$$|\delta_3|^2 m_0 \simeq \left(m_b - \frac{m_\tau}{2}\right)$$

$$|\delta_2|^2 m_0 \simeq \frac{2m_b}{m_\tau}\left(m_s - \frac{m_\mu}{2}\right)$$

$$|\delta_1|^2 m_0 \simeq \frac{4m_s m_b}{m_\mu m_\tau}\left(m_d - \frac{m_e}{2}\right)$$

$$|\delta_3|^2 m_0 = 0.313\text{GeV}, |\delta_2|^2 m_0 = 0.035\text{GeV}, |\delta_1|^2 m_0 = 7.09 \times 10^{-4}\text{GeV}$$

$$(Y_{15})_{33} = 1.434, (Y_{15})_{23} = 0.481, (Y_{15})_{13} = 0.068$$

$$|(Y_{15})_{33}/(M_{15})_3| = 0.358$$

$$\frac{1}{2}|(Y_{15})_{33}/(M_{15})_3|^2 \simeq 0.064$$

$$m_c \sim \{(Y_{15})_{23}\}^4 \kappa_L\kappa_R/(\{(Y_{15})_{33}\}^2 (M_{15})_2)$$

$$(M_{15})_2/\kappa_R \geq 15$$

$$|(Y_{15})_{22}|^2 \kappa_L \kappa_R/(M_{15})_2$$

$$(M_{15})_2/\kappa_R \sim 600.$$

$(M_{15})_1$ to ΔM_d and ΔM_e

$$V_d = \begin{pmatrix} 0.991 & 0.132 & -0.024 \\ 0.134 & -0.987 & 0.083 \\ 0.013 & 0.086 & 0.996 \end{pmatrix}$$

$$\Delta M_d \simeq (3)\frac{Y_{10}M_{10}Y_{10}^{\dagger}}{16\pi^2}(2\lambda_4\kappa_L\kappa_R)F\left(M_{10_3}^2, M_{\chi_1^u}^2, M_{\chi_2^u}^2\right)$$

$$\Delta M_e \simeq (5)\frac{Y_{10}M_{10}Y_{10}^{\dagger}}{16\pi^2}(2\lambda_4\kappa_L\kappa_R)F\left(M_{10_3}^2, M_{\chi_1^u}^2, M_{\chi_2^u}^2\right)$$

$$M_d = \begin{pmatrix} m_{11} & m_{12} & m_{13} \\ m_{12}^* & m_{22} & m_{23} \\ m_{13}^* & m_{23}^* & m_{33} \end{pmatrix} + 4m_0 \begin{pmatrix} |\delta_1|^2 & \delta_1\delta_2^* & \delta_1\delta_3^* \\ \delta_1^*\delta_2 & |\delta_2|^2 & \delta_2\delta_3^* \\ \delta_1^*\delta_3 & \delta_2^*\delta_3 & |\delta_3|^2 \end{pmatrix}$$

$$M_e = 2\begin{pmatrix} m_{11} & m_{12} & m_{13} \\ m_{12}^* & m_{22} & m_{23} \\ m_{13}^* & m_{23}^* & m_{33} \end{pmatrix} + 6m_0 \begin{pmatrix} |\delta_1|^2 & \delta_1\delta_2^* & \delta_1\delta_3^* \\ \delta_1^*\delta_2 & |\delta_2|^2 & \delta_2\delta_3^* \\ \delta_1^*\delta_3 & \delta_2^*\delta_3 & |\delta_3|^2 \end{pmatrix}$$

$$\delta_i = (Y_{10})_{i3}$$

$$m_0 = \left(\frac{1}{2}\right)\frac{(M_{10})_3}{16\pi^2}(2\lambda_4\kappa_L\kappa_R)F\left(M_{10_3}^2, M_{\chi_1^u}^2, M_{\chi_2^u}^2\right)$$


$$(Y_{10})_{33} = 1.434, (Y_{10})_{23} = 0.481, (Y_{10})_{13} = 0.068$$


$$m_\tau \simeq 2\frac{|(Y_{10})_{33}|^2\kappa_L\kappa_R}{(M_{10})_3} + 6m_0|\delta_3|^2$$

$$\lambda_4 = -3.14, (M_{10})_3 = 219\kappa_R,$$


$$m_0 = -0.152\text{GeV and } m_\tau = 1.76\text{GeV}$$

mass relation $M_{\chi_1^u} \simeq \sqrt{-\lambda_4}\kappa_R$





$$\mathcal{M}_\nu = \begin{pmatrix} 0 & 0 & -\frac{\sqrt{3}}{2} Y_{15}^* \kappa_L \\ 0 & 0 & -\frac{\sqrt{3}}{2} Y_{15} \kappa_R \\ -\frac{\sqrt{3}}{2} Y_{15}^\dagger \kappa_L - \frac{\sqrt{3}}{2} Y_{15}^T \kappa_R & & M_{15} \end{pmatrix}.$$



(a) (b) (c)




(d) (e)





(f) (g)





(h) (i)

















$$\mathcal{M}_\nu = \begin{pmatrix} m_L & m_\nu^D & -\frac{\sqrt{3}}{2}Y_{15}^*\kappa_L \\ (m_\nu^D)^T & m_R & -\frac{\sqrt{3}}{2}Y_{15}\kappa_R \\ -\frac{\sqrt{3}}{2}Y_{15}^\dagger\kappa_L & -\frac{\sqrt{3}}{2}Y_{15}^T\kappa_R & M_{15} \end{pmatrix}.$$

$$M_\nu^\ell \approx m_L - \left(m_\nu^D - \frac{\sqrt{3}}{2}Y_{15}^*\kappa_L\right)\begin{pmatrix} m_R & -\frac{\sqrt{3}}{2}Y_{15}\kappa_R \\ -\frac{\sqrt{3}}{2}Y_{15}^T\kappa_R & M_{15} \end{pmatrix}^{-1}\begin{pmatrix} (m_\nu^D)^T \\ -\frac{\sqrt{3}}{2}Y_{15}^\dagger\kappa_L \end{pmatrix}$$

$$\approx -\frac{\kappa_L}{\kappa_R}\left(m_\nu^D\left(Y_{15}^T\right)^{-1}Y_{15}^\dagger + Y_{15}^*Y_{15}^{-1}(m_\nu^D)^T\right) + \frac{\kappa_L^2}{\kappa_R^2}\left(Y_{15}^*Y_{15}^{-1}m_R\left(Y_{15}^T\right)^{-1}Y_{15}^\dagger\right) + m_L$$

$$M_{15} \ll \kappa_R m_R^{-1}\kappa_R$$

$$m_L \simeq \frac{3}{4}Y_{15}^\dagger M_{15}\left[\frac{3}{32\pi^2}\frac{M_{Z_L}^2}{M_{15_3}^2 - M_{Z_L}^2}\ln\left(\frac{M_{15_3}^2}{M_{Z_L}^2}\right) + \frac{1}{32\pi^2}\frac{M_{\sigma_L}^2}{M_{15_3}^2 - M_{\sigma_L}^2}\ln\left(\frac{M_{15_3}^2}{M_{\sigma_L}^2}\right)\right]Y_{15}^*$$

$$m_R \simeq \frac{3}{4}Y_{15}^\dagger M_{15}\left[\frac{3}{32\pi^2}\frac{M_{Z_R}^2}{M_{15_3}^2 - M_{Z_R}^2}\ln\left(\frac{M_{15_3}^2}{M_{Z_R}^2}\right) + \frac{1}{32\pi^2}\frac{M_{\sigma_R}^2}{M_{15_3}^2 - M_{\sigma_R}^2}\ln\left(\frac{M_{15_3}^2}{M_{\sigma_R}^2}\right)\right]Y_{15}^*$$

$$m_\nu^D = m_\nu^D(\mathrm{f}+\mathrm{g}) + m_\nu^D(\mathrm{h}) + m_\nu^D(\mathrm{i})$$

$$m_\nu^D(\mathrm{f}+\mathrm{g}) \simeq 3Y_{15}^\dagger M_{15}\left[\frac{3g_4^2}{32\pi^2}\frac{\kappa_L\kappa_R}{M_{15_3}^2 - M_X^2}\ln\left(\frac{M_{15_3}^2}{M_X^2}\right) + \frac{\lambda_4\kappa_L\kappa_R}{16\pi^2}F\left(M_{15_3}^2, M_{\chi_1^u}^2, M_X^2\right)\right]Y_{15}$$

$$m_\nu^D(\mathrm{h}) \simeq \frac{3(2\lambda_5\kappa_L\kappa_R)}{16\pi^2}Y_{10}^\dagger M_{10}Y_{10}F\left(M_{10_3}^2, M_{\chi_1^d}^2, M_{\chi_2^d}^2\right)$$

$$m_\nu^D(\mathrm{i}) \simeq \frac{2(\lambda_3+\lambda_4)\kappa_L\kappa_R}{16\pi^2}Y_{15}^\dagger M_{15}Y_{15}F\left(M_{15_3}^2, M_{\sigma_L}^2, M_{\sigma_R^d}^2\right)$$

$$g_{Z_L\nu\nu} = \sqrt{g_Y^2 + g_{2L}^2}/2 \text{ and } g_{Z_R\nu_R\nu_R} = g_{2R}^2/\left(2\sqrt{g_{2R}^2 - g_Y^2}\right)$$

$$\mathcal{L}_\theta^{\text{QCD}} = \frac{g_s^2}{32\pi^2}\theta_{\text{QCD}} G_{\mu\nu}^a \tilde{G}^{a\mu\nu}$$

$$\tilde{G}^{a\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\alpha\beta} G_{\alpha\beta}^a$$

$$\bar{\theta} = \theta_{\text{QCD}} + \text{ArgDet}(\mathcal{M}_u\mathcal{M}_d) + 5\text{ArgDet}\mathcal{M}_S + 3\text{ArgDet}\mathcal{M}_\mathcal{O}$$

$$\mathcal{M}_i^{1\,\text{L}} \equiv \mathcal{M}_i + \delta\mathcal{M}_i = \mathcal{M}_i(1 + \Sigma_i), \Sigma_i = \mathcal{M}_i^{-1}\delta\mathcal{M}_i$$

$$\bar{\theta}^{(1)} = \text{ImTr}\left[\Sigma_u^{(1)} + \Sigma_d^{(1)} + 5\Sigma_S^{(1)} + 3\Sigma_\mathcal{O}^{(1)}\right]$$

$$\Sigma_q = \mathcal{M}_q^{-1}\delta\mathcal{M}_q$$

$$\bar{\theta}_{(u+d)}^{(1)} = \text{ImTr}\left[\mathcal{M}_u^{-1}\delta\mathcal{M}_u + \mathcal{M}_d^{-1}\delta\mathcal{M}_d\right]$$

$$\delta\mathcal{M}_q = \begin{pmatrix} \delta\mathcal{M}_{LL}^q & \delta\mathcal{M}_{LH}^q \\ \delta\mathcal{M}_{HL}^q & \delta\mathcal{M}_{HH}^q \end{pmatrix}$$

$$\bar{\theta}_d^{(1)} = \text{ImTr}\left[-\frac{1}{\kappa_L\kappa_R}\delta\mathcal{M}_{LL}^d\left(Y_{10}^\dagger\right)^{-1}M_{10}Y_{10}^{-1} + \frac{1}{\kappa_L}\delta\mathcal{M}_{LH}^d Y_{10}^{-1} + \frac{1}{\kappa_R}\delta\mathcal{M}_{HL}^d\left(Y_{10}^\dagger\right)^{-1}\right]$$

$$-\mathcal{L}_{\text{eff}}^{e,\,\text{tree}} = (\bar{e}_R\bar{E}_R)\mathcal{M}_e^\dagger k^2\left(k^2 - \mathcal{M}_e\mathcal{M}_e^\dagger\right)^{-1}\begin{pmatrix} e_L \\ E_L \end{pmatrix}$$

$$\left(\mathcal{M}_e\mathcal{M}_e^\dagger - k^2\right)^{-1} = \begin{pmatrix} P_e(k^2) & Q_e(k^2) \\ Q_e^\dagger(k^2) & R_e(k^2) \end{pmatrix}, \text{ with } P_e(k^2) = P_e^\dagger(k^2), R_e(k^2) = R_e^\dagger(k^2)$$

$$\left(\mathcal{M}_e\mathcal{M}_e^\dagger - k^2\right)\left(\mathcal{M}_e\mathcal{M}_e^\dagger - k^2\right)^{-1} = \mathbb{1}$$

$$\left(2\kappa_R^2 Y_{10}^\dagger Y_{10} + M_{10}M_{10}^\dagger - k^2\right)Q_e^\dagger(k^2) = -\sqrt{2}\kappa_L M_{10}Y_{10}^\dagger P_e(k^2)$$
$$2\kappa_L Y_{10}Y_{10}^\dagger P_e(k^2) + \sqrt{2}Y_{10}M_{10}^\dagger Q_e^\dagger(k^2) = \frac{1}{\kappa_L}\left(\mathbb{1} + k^2 P_e(k^2)\right)$$
$$Q_e(k^2) = -\sqrt{2}\kappa_L H_e(k^2)Y_{10}M_{10}^\dagger R_e(k^2)$$

$$H_e(k^2) = \left(2\kappa_L^2 Y_{10}Y_{10}^\dagger - k^2\right)^{-1} = H_e^\dagger(k^2)$$
$$R_e(k^2) = \left[\left(2\kappa_R^2 Y_{10}^\dagger Y_{10} + M_{10}M_{10}^\dagger - k^2\right) - 2\kappa_L^2 M_{10}Y_{10}^\dagger H_e Y_{10}M_{10}^\dagger\right]^{-1} = R_e^\dagger(k^2)$$

$$-\mathcal{L}_{\text{eff}}^{e,\,\text{tree}} = \bar{E}_R\left[\frac{k^4}{\sqrt{2}\kappa_L}Y_{10}^{-1}Q_e(k^2)\right]E_L + \bar{e}_R\left[\sqrt{2}k^2 Y_{10}\kappa_R R_e(k^2)\right]E_L$$
$$+\bar{E}_R\left[\frac{k^2}{\sqrt{2}\kappa_L}Y_{10}^{-1}\left(\mathbb{1} + k^2 P_e(k^2)\right)\right]e_L + \bar{e}_R\left[\sqrt{2}k^2 Y_{10}\kappa_R Q_e^\dagger(k^2)\right]e_L + \text{ h.c.}$$

$$\mathcal{O}(\kappa_L/\kappa_R)\ \text{ð}\left(\overline{\nu_L}\overline{\nu_L^c}\overline{N_L}\right)\mathcal{M}_\nu'(\nu_R^c\nu_R N_R^c)^T$$

$$U \simeq \begin{pmatrix} \mathbb{1} & -\rho \\ \rho^\dagger & \mathbb{1} \end{pmatrix}, \rho = \frac{\kappa_L}{\kappa_R} Y_{15} (Y_{15}^*)^{-1}$$

$$\nu_{\ell L} = \nu_L - \rho \nu_L^c \text{ vanishes to } \mathcal{O}(\kappa_L^2/\kappa_R^2)$$

$$\nu_{\ell L} = \nu_L - \rho \nu_L^c, \nu_{hL} = \nu_L^c + \rho^\dagger \nu_L$$

$$\mathcal{L}_{\text{mass}}^{\nu_h} = \left(\overline{\nu_{hL}} \overline{N_L}\right) \mathcal{M}_{\nu_h} \begin{pmatrix} \nu_{hR}^c \\ N_R^c \end{pmatrix} + \text{ h.c.}$$

$$\mathcal{M}_{\nu_h} = \begin{pmatrix} 0 & -\frac{\sqrt{3}}{2} \kappa_R Y_{15}^* \\ -\frac{\sqrt{3}}{2} \kappa_R Y_{15}^\dagger & M_{15}^\dagger \end{pmatrix} + \mathcal{O}(\kappa_L^2/\kappa_R^2)$$

$$\left(\mathcal{M}_{\nu_h} \mathcal{M}_{\nu_h}^\dagger - k^2\right)^{-1} = \begin{pmatrix} P_\nu(k^2) & Q_\nu(k^2) \\ Q_\nu^\dagger(k^2) & R_\nu(k^2) \end{pmatrix}, \text{ with } P_\nu(k^2) = P_\nu^\dagger(k^2), R_\nu(k^2) = R_\nu^\dagger(k^2)$$


$$\left(\frac{3}{4}\kappa_R^2 Y_{15}^\dagger Y_{15} + M_{15}^\dagger M_{15} k^2\right) Q_\nu^\dagger(k^2) = \frac{\sqrt{3}}{2} \kappa_R M_{15}^\dagger Y_{15}^T P_\nu(k^2)$$

$$\frac{3}{4}\kappa_R Y_{15}^* Y_{15}^T P_\nu(k^2) - \frac{\sqrt{3}}{2} Y_{15}^* M_{15} Q_\nu^\dagger(k^2) = \frac{1}{\kappa_R}\left(\mathbb{1} + k^2 P_\nu(k^2)\right)$$

$$Q_\nu(k^2) = \frac{\sqrt{3}}{2} \kappa_R H_\nu(k^2) Y_{15}^* M_{15} R_\nu(k^2)$$

$$H_\nu(k^2) = \left(\frac{3}{4}\kappa_R^2 Y_{15}^* Y_{15}^T - k^2\right)^{-1} = H_\nu^\dagger(k^2)$$

$$R_\nu(k^2) = \left[\left(\frac{3}{4}\kappa_R^2 Y_{15}^\dagger Y_{15} + M_{15}^\dagger M_{15} - k^2\right) - \frac{3}{4}\kappa_R^2 M_{15}^\dagger Y_{15}^T H_\nu Y_{15}^* M_{15}\right]^{-1} = R_\nu^\dagger(k^2)$$



$$-\mathcal{L}_{\text{eff}}^{\nu,\text{ tree}} = \overline{N_R^c}\left[-\frac{2k^4}{\sqrt{3}\kappa_R}(Y_{15}^*)^{-1} Q_\nu(k^2)\right] N_L + \overline{\nu_{hR}^c}\left[-\frac{\sqrt{3}k^2}{2} Y_{15} \kappa_R R_\nu(k^2)\right] N_L$$
$$+\overline{N_R^c}\left[-\frac{2k^2}{\sqrt{3}\kappa_R}(Y_{15}^*)^{-1}\left(\mathbb{1} + k^2 P_\nu(k^2)\right)\right] \nu_{hL} + \overline{\nu_{hR}^c}\left[-\frac{\sqrt{3}k^2}{2} Y_{15} \kappa_R Q_\nu^\dagger(k^2)\right] \nu_{hL} + \text{ h.c.}$$

$$\delta \mathcal{M}_{LL}^d = \int \frac{d^4k}{(2\pi)^4} \frac{Y_{15} M_{15} Y_{15}^\dagger (2\lambda_5 \kappa_L \kappa_R)}{(k^2 - M_{15}^2)\left((p-k)^2 - M_{\chi_R^d}^2\right)\left((p-k)^2 - M_{\chi_L^d}^2\right)},$$

$$\mathrm{ImTr}\left[-2\lambda_5 \int \frac{d^4k}{(2\pi)^4} \frac{Y_{15} M_{15} Y_{15}^\dagger}{(k^2 - M_{15}^2)\left((p-k)^2 - M_{\chi_R^d}^2\right)\left((p-k)^2 - M_{\chi_L^d}^2\right)} \left(Y_{10}^\dagger\right)^{-1} M_{10} Y_{10}^{-1}\right].$$

$$Y_{15} M_{15} Y_{15}^\dagger \text{ and } \left(Y_{10}^\dagger\right)^{-1} M_{10} Y_{10}^{-1}$$

$$\mathrm{ImTr}\left[\left(Y_{10} M_{10} Y_{10}^\dagger\right)\left(\left(Y_{10}^\dagger\right)^{-1} M_{10} Y_{10}^{-1}\right)\right] = 0$$

$$\mathrm{ImTr}\left[Q_e Y_{10}^\dagger \left(Y_{10}^\dagger\right)^{-1} M_{10} Y_{10}^{-1}\right] = -\sqrt{2}\kappa_L \mathrm{ImTr}\left[\left(2\kappa_L^2 Y_{10} Y_{10}^\dagger - k^2\right)^{-1}\left(M_{10}^\dagger R_e(k^2) M_{10}\right)\right],$$

$$\mathrm{ImTr}\left[R_e^\dagger Y_{10}^\dagger \left(Y_{10}^\dagger\right)^{-1}\right] = 0$$

$$\mathrm{ImTr}\left[\left(\mathbb{1} + k^2 P_e(k^2)\right)^\dagger (Y_{10}^{-1})^\dagger Y_{10}^{-1}\right] = 0$$

$$\mathrm{ImTr}\left[\left(Y_{15} M_{15} Y_{15}^\dagger\right)\left(\left(Y_{15}^\dagger\right)^{-1} M_{15}^\dagger Y_{15}^{-1}\right)\right] = 0$$

$\left(Y_{15}^\dagger\right)^{-1} M_{15}^\dagger Y_{15}^{-1}$ appears from the $\delta\mathcal{M}_{LL}^u$ insertion

$$\mathrm{ImTr}\left[\left(Y_{10}^\dagger M_{10}^\dagger Y_{10}\right)\left(\left(Y_{15}^\dagger\right)^{-1} M_{15}^\dagger Y_{15}^{-1}\right)\right] = 0$$

$$\mathrm{ImTr}\left[Y_{15}(Y_{15}^*)^{-1} Q_\nu(k^2) Y_{15}^\dagger \left(\left(Y_{15}^\dagger\right)^{-1} M_{15}^\dagger Y_{15}^{-1}\right)\right]$$

$$\mathrm{ImTr}\left[\left(\frac{3}{4}\kappa_R^2 Y_{15}^T Y_{15}^* - k^2\right)^{-1} M_{15}^\dagger R_\nu(k^2) M_{15}^\dagger\right] = 0$$

since $R_\nu(k^2)$ is Hermitian and $\left(\frac{3}{4}\kappa_R^2 Y_{15}^T Y_{15}^* - k^2\right)^{-1}$ is also Hermitian

$$\mathrm{ImTr}\left[\rho Q_\nu(k^2) Y_{15}^\dagger \left(\left(Y_{15}^\dagger\right)^{-1} M_{15}^\dagger Y_{15}^{-1}\right)\right]$$


$$\mathrm{ImTr}\left[\frac{\kappa_L}{\kappa_R}\left(\frac{3}{4}\kappa_R^2 Y_{15}^T Y_{15}^* - k^2\right)^{-1} M_{15}^\dagger R_\nu(k^2) M_{15}^\dagger\right] = 0$$

$$\mathrm{ImTr}\left[R_\nu^\dagger(k^2) Y_{15}^\dagger \left(Y_{15}^\dagger\right)^{-1}\right] = 0$$

$$\mathrm{ImTr}\left[\rho\left(\mathbb{1} + k^2 P_\nu^\dagger(k^2)\right)\left(Y_{15}^T\right)^{-1} Y_{15}^{-1}\right] = \mathrm{ImTr}\left[(Y_{15}^*)^{-1}\left(\mathbb{1} + k^2 P_\nu^\dagger(k^2)\right)\left(Y_{15}^T\right)^{-1}\right] = 0,$$

$$P_\nu(k^2) = P_\nu^\dagger(k^2)$$

$$\mathrm{ImTr}\left[Y_{15}\xi Y_{15}^\dagger \left(\left(Y_{10}^\dagger\right)^{-1} M_{10} Y_{10}^{-1}\right)\right]$$



$$\xi = \frac{\sqrt{3}}{2}\kappa_R\left(k^4\left(M_{15}^{\dagger}\right)^{-1} - k^2 M_{15}^{\dagger} + \frac{9}{16}\kappa^4 Y_{15}^{\dagger}Y_{15}M_{15}^{-1}Y_{15}^{T}Y_{15}^{*} - \frac{3}{4}k^2\kappa_R^2\left(M_{15}^{-1}Y_{15}^{T}Y_{15}^{*} + Y_{15}^{\dagger}Y_{15}M_{15}^{-1}\right)\right)^{-1}$$

$$\bar{\theta} \sim \frac{\lambda_5}{96\pi^2}\frac{m_t}{m_b}M_{15}\kappa_R F\left(M_{15_3}^2, M_{\chi_1^d}^2, M_{\chi_2^d}^2\right),$$

Input: $\lambda_5 = 3, \frac{m_t}{m_b} = 10^2, M_{15} = 10^5\text{GeV},$
$M_{15_3} = 10^{14}\text{GeV}, \kappa_R = 5\times10^{13}\text{GeV}, M_{\chi_1^d} = 8\times10^{13}\text{GeV}, M_{\chi_2^d} = 7\times10^{13}\text{GeV},$

Output: $\bar{\theta} \approx 10^{-10}$.

$$\bar{\theta} \sim \frac{\lambda_5}{96\pi^2}Y_{15}M_{15_3}Y_{15}^{\dagger}\left(\left(Y_{10}^{\dagger}\right)^{-1}M_{10}Y_{10}^{-1}\right)F\left(M_{15_3}^2, M_{\chi_1^d}^2, M_{\chi_2^d}^2\right),$$

Input: $\lambda_5 = 0.1, Y_{15} = 1, Y_{10} = 10^{-2}, M_{153} = 10^{13}\text{GeV},$
$M_{10} = 10^5\text{GeV}, M_{\chi_1^d} = 8\times10^{13}\text{GeV}, M_{\chi_2^d} = 7\times10^{13}\text{GeV},$
Output: $\bar{\theta} \approx 10^{-10}$.

$$\left(H_L^T\tilde{H}_L^*\right)\left(H_R^{\dagger}\tilde{H}_R\right)$$

$$\lambda_5 \sim \frac{3}{16\pi^2}\text{Tr}\left(Y_{10}^{\dagger}Y_{10}Y_{15}^{\dagger}Y_{15}\right)\ln\left(\frac{M_{15_3}^2}{M_{\text{Pl}}^2}\right) \sim 4.4\times10^{-5}$$

$Y_{10} \sim 10^{-2}$ (effective bottom Yukawa), $Y_{15} \sim 1$ (top Yukawa), $M_{15_3} \sim 10^{14}\text{GeV}$, and $M_{\text{Pl}} \sim$

10^{19}GeV

$$\bar{\theta}^{(1)}_{(\mathcal{S}+\mathcal{O})} = \mathrm{Im}\mathrm{Tr}\left[5\mathcal{M}_{\mathcal{S}}^{-1}\delta\mathcal{M}_{\mathcal{S}} + 3\mathcal{M}_{\mathcal{O}}^{-1}\delta\mathcal{M}_{\mathcal{O}}\right]$$

$$5k^2\kappa_R \mathrm{Im}\mathrm{Tr}[M_{10}^{-1}\zeta_a]$$


$$\zeta_a = \left(Y_{10}^{\dagger}Y_{10}R_d^{\dagger}M_{10}Y_{10}^{\dagger}H_dY_{10}\right)$$

$$\zeta_a = -\kappa_L\Big(k^4Y_{10}^{-1}\left(Y_{10}^{\dagger}\right)^{-1}M_{10}^{-1}Y_{10}^{-1}\left(Y_{10}^{\dagger}\right)^{-1} - k^2Y_{10}^{-1}\left(Y_{10}^{\dagger}\right)^{-1}M_{10}^{\dagger}Y_{10}^{-1}\left(Y_{10}^{\dagger}\right)^{-1} + \kappa_L^2\kappa_R^2M_{10}^{-1}$$
$$-k^2\left(\kappa_L^2M_{10}^{-1}Y_{10}^{-1}\left(Y_{10}^{\dagger}\right)^{-1} + \kappa_R^2Y_{10}^{-1}\left(Y_{10}^{\dagger}\right)^{-1}M_{10}^{-1}\right)\Big)^{-1}$$

$$\bar{\theta}^1_{\mathcal{S}(a)} \sim \frac{5\lambda_4}{8\pi^2}Y_{10}^{\dagger}\left(\frac{\kappa_L^2}{M_{10_3}^2 - M_{\chi_1^u}^2}\ln\left(\frac{M_{10_3}^2}{M_{\chi_1^u}^2}\right) - \frac{\kappa_L^2}{M_{10_3}^2 - M_X^2}\ln\left(\frac{M_{10_3}^2}{M_X^2}\right)\right)Y_{10}$$

$$5k^2\kappa_R \mathrm{Im}\mathrm{Tr}\left[M_{10}^{-1}Y_{10}^{\dagger}\zeta_bY_{10}\right]\mathrm{Tr}(M_{10}^{-1}\zeta_a)$$

$$\zeta_b = -\kappa_L\left(k^4\left(Y_{15}^{\dagger}\right)^{-1}\left(M_{15}^{\dagger}\right)^{-1}Y_{15}^{-1} - k^2\left(Y_{15}^{\dagger}\right)^{-1}M_{15}Y_{15}^{-1} + \kappa_L^2\kappa_R^2Y_{15}\left(M_{15}^{\dagger}\right)^{-1}Y_{15}^{\dagger}\right.$$
$$\left.-k^2\left(\kappa_L^2Y_{15}\left(M_{15}^{\dagger}\right)^{-1}Y_{15}^{-1} + \kappa_R^2\left(Y_{15}^{\dagger}\right)^{-1}\left(M_{15}^{\dagger}\right)^{-1}Y_{15}^{\dagger}\right)\right)^{-1}$$



$$\bar{\theta}^1_{\mathcal{S}(b)} \sim \frac{5\lambda_5\kappa_L^2}{8\pi^2M_{10}}Y_{10}^{\dagger}\left(\frac{M_{15}}{M_{15_3}^2 - M_{\chi_1^d}^2}\ln\left(\frac{M_{15_3}^2}{M_{\chi_1^d}^2}\right) - \frac{M_{15}}{M_{15_3}^2 - M_{\chi_2^d}^2}\ln\left(\frac{M_{15_3}^2}{M_{\chi_2^d}^2}\right)\right)Y_{10}.$$

$$\frac{5k^4}{\kappa_L}\mathrm{Im}\mathrm{Tr}[M_{10}^{-1}\zeta_c]$$


$$\zeta_c = -\kappa_L(k^4M_{10}^{-1} - k^2M_{10}^{\dagger} + \kappa_L^2\kappa_R^2Y_{10}^{\dagger}Y_{10}M_{10}^{-1}Y_{10}^{\dagger}Y_{10}$$
$$-k^2\left(\kappa_L^2Y_{10}^{\dagger}Y_{10}M_{10}^{-1} + \kappa_R^2M_{10}^{-1}Y_{10}^{\dagger}Y_{10}\right)\Big)^{-1}$$

$$\frac{1}{\kappa_L}\mathrm{Tr}[\zeta_cM_{10}^{-1}]\langle\chi_L^{\nu}\rangle \neq 0$$

$$\bar{\theta}^1_{\mathcal{S}(c)} \sim \frac{15g_4^2}{32\pi^2}Y_{10}^{\dagger}\frac{\kappa_L^2}{M_{10_3}^2 - M_X^2}\ln\left(\frac{M_{10_3}^2}{M_X^2}\right)Y_{10},$$

factor $(\kappa_L/\kappa_R)^2 \sim 10^{-23}$, or by a factor (M_{10}/κ_R) or (M_{15}/κ_R)

$$V(\lambda_6) = 4\lambda_6 \left(\chi_L^{u\alpha}\chi_L^{d\beta}\chi_R^{d\gamma}\chi_R^{\nu}\epsilon_{\alpha\beta\gamma} + \chi_L^{d\alpha}\chi_R^{u\beta}\chi_R^{d\gamma}\chi_L^{\nu}\epsilon_{\alpha\beta\gamma}\right) + \text{ h.c.}$$



$$A[(\bar{\nu}_L d_R)(\bar{e}_L d_R)(\bar{e}^c{}_L u_R)] = \frac{4\lambda_6\kappa_R}{M_{\chi_L}^4 M_{\chi_R^d}^2}\left(\frac{Y_{10}^2\kappa_R}{M_{10}}\right)^2\left(\frac{Y_{15}^2\kappa_R}{M_{15}}\right).$$

$$n \to e^+e^-\nu$$
$$p \to \pi^+\pi^0 e^+e^-\nu$$

$$\mathcal{L}_{\text{gravity}}^{\text{VEVs}\,\langle\chi_{L,R}^{\nu}\rangle} = \frac{A}{M_{\text{P}}}\text{Tr}(\bar{\Psi}_L H_L)\text{Tr}\left(H_R^{\dagger}\Psi_R\right) + \frac{B}{M_{\text{P}}}\text{Tr}\left(\bar{\Psi}_L H_L H_R^{\dagger}\Psi_R\right)$$

$$\mathcal{L}_{\text{gravity}}^{\text{VEVs}\,\langle\chi_{L,R}^{\mu}\rangle} = \frac{c}{M_{\text{P}}^2}G_{\mu\nu}G_{\alpha\beta}\epsilon^{\mu\nu\alpha\beta}\left(\text{Tr}\left(H_L^{\dagger}H_L\right) - \text{Tr}\left(H_R^{\dagger}H_R\right)\right)$$

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \begin{pmatrix} S^{-1} & -S^{-1}BD^{-1} \\ -D^{-1}CS^{-1} & D^{-1} + D^{-1}CS^{-1}BD^{-1} \end{pmatrix}$$

$$\delta\Phi_{\pm} = i\theta_{\pm}\Phi_{\pm}$$

$$\delta\left(\Phi_+(t_{\text{f}}) - \Phi_-(t_{\text{f}})\right) = i\theta_+\Phi_+(t_{\text{f}}) - i\theta_-\Phi_-(t_{\text{f}}) = i(\theta_+ - \theta_-)\Phi_+(t_{\text{f}}) = 0$$

$$\delta\pi_{\pm} = v_{\mu}^{\pm}x^{\mu}$$

$$\delta\left(\pi_+(t_{\text{f}}) - \pi_-(t_{\text{f}})\right) = (v_0^+ - v_0^-)t_{\text{f}} + (v_i^+ - v_i^-)x^i = 0$$

$$S = -\int \text{d}^4x\left[\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \left(D_\mu\Phi\right)^* D^\mu\Phi + V(\Phi^*\Phi)\right]$$

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu, D_\mu = \partial_\mu - iqA_\mu$$

$$V = m^2|\Phi|^2$$

$$\Phi(x) \to e^{iq\lambda(x)}\Phi(x), \text{ and } A_\mu(x) \to A_\mu(x) + \partial_\mu\lambda(x)$$

$$S_{\mathrm{GF}}[A] \equiv -\frac{1}{2\xi}\int \mathrm{d}^4x\left(\partial_\mu A^\mu\right)^2$$

$$S_{\text{Becchi–Rouet–Stora–Tyutin (BRST)}} = -\int \mathrm{d}^4x\left[\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \left(D_\mu\Phi\right)^* D^\mu\Phi + V(\Phi^*\Phi) + \frac{1}{2\xi}\left(\partial_\mu A^\mu\right)^2 + \bar{c}\Box c\right]$$

$$\begin{aligned} A_\mu(x) &\to A_\mu(x) + \eta\partial_\mu c(x), & c(x) &\to c(x) \\ \Phi(x) &\to \Phi(x) + iq\eta c(x)\Phi(x), & \bar{c}(x) &\to \bar{c}(x) - \frac{\eta}{\xi}\partial_\mu A^\mu(x) \end{aligned}$$

$$Q_{\mathrm{BRST}} = \int \mathrm{d}^{\mathcal{D}}\mathbf{x}\left[\Pi_A^i\partial_i c + iq(\Pi_\Phi\Phi - \Pi_{\Phi^*}\Phi^*)c - \frac{1}{\xi}\Pi_{\bar{c}}\left(\partial_\mu A^\mu\right)\right]$$

where $\Pi_{f^{\mathfrak{a}}} = \partial\mathcal{L}_{\mathrm{BRST}}/\partial\dot{f}^{\mathfrak{a}}$ is the conjugate momentum to $\boldsymbol{f} = \{f^{\mathfrak{a}}\} = (A, c, \bar{c}, \Phi, \Phi^*)$ where $S_{\mathrm{BRST}} =$

$$\int \mathrm{d}^4x\mathcal{L}_{\mathrm{BRST}}$$

Schrödinger-picture operator $\hat{A}_S^\mu(\mathbf{x}), \hat{\Phi}_S(\mathbf{x}), \hat{c}_S(\mathbf{x})$ and $\hat{\bar{c}}_S(\mathbf{x})$

respective field eigenstates:

$$\begin{aligned} \hat{A}_S^\mu(\mathbf{x})|\mathrm{A}\rangle &= \mathrm{A}^\mu(\mathbf{x})|\mathrm{A}\rangle, & \hat{c}_S(\mathbf{x})|\mathrm{c}\rangle &= \Phi(\mathbf{x})|\mathrm{c}\rangle \\ \hat{\Phi}_S(\mathbf{x})|\Phi\rangle &= & \hat{\bar{c}}_S(\mathbf{x})|\bar{\mathrm{c}}\rangle &= \bar{\mathrm{c}}(\mathbf{x})|\bar{\mathrm{c}}\rangle \\ (\mathbf{x})|\Phi\rangle, & & & \end{aligned}$$

$$\hat{A}^\mu(x) = \hat{U}(t, t_{\mathrm{i}})\hat{A}_S^\mu(\mathbf{x})\hat{U}^\dagger(t, t_{\mathrm{i}})$$

$$\begin{aligned} &\langle \mathrm{A_f c_f}\bar{\mathrm{c}}_{\mathrm{f}}\Phi_{\mathrm{f}}\Phi_{\mathrm{f}}^*|\hat{U}(t_{\mathrm{f}}, t_{\mathrm{i}})|\mathrm{A_i c_i}\bar{\mathrm{c}}_{\mathrm{i}}\Phi_{\mathrm{i}}\Phi_{\mathrm{i}}^*\rangle \\ &= \int_{\mathrm{A_i}}^{\mathrm{A_f}}\mathcal{D}[A]\int_{\mathrm{c_i}}^{\mathrm{c_f}}\mathcal{D}[c]\int_{\bar{\mathrm{c}}_{\mathrm{i}}}^{\bar{\mathrm{c}}_{\mathrm{f}}}\mathcal{D}[\bar{c}]\int_{\Phi_{\mathrm{i}}}^{\Phi_{\mathrm{f}}}\mathcal{D}[\Phi]\int_{\Phi_{\mathrm{i}}^*}^{\Phi_{\mathrm{f}}^*}\mathcal{D}[\Phi^*]\mu e^{iS_{\mathrm{BRST}}[A,c,\bar{c},\Phi,\Phi^*;t_{\mathrm{i}},t_{\mathrm{f}}]} \end{aligned}$$

$$Z[J_A, J_\Phi, J_{\Phi^*}, J_c, J_{\bar{c}}] = \int_{\mathrm{vac}}^{\mathrm{vac}}\mathcal{D}[A, c, \bar{c}, \Phi, \Phi^*]\mu e^{iS_{\mathrm{BRST}}[A,c,\bar{c},\Phi,\Phi^*] + i\int \mathrm{d}^4x\left(J_A\cdot A + J_\Phi\Phi + J_{\Phi^*}\Phi^* + J_c c + J_{\bar{c}}\bar{c}\right)}$$


$$(-i)^2\frac{\delta^2 Z}{\delta J_A^\mu(x)\delta J_A^\nu(y)}\bigg|_{J_{\mathfrak{a}}=0} = \langle 0|\mathcal{T}\{\hat{A}_\mu(x)\hat{A}_\nu(y)\}|0\rangle$$


$$\rho(t_{\mathrm{f}}) = \hat{U}(t_{\mathrm{f}}, t_{\mathrm{i}})\rho_{\mathrm{i}}\hat{U}^\dagger(t_{\mathrm{f}}, t_{\mathrm{i}})$$

$$\begin{aligned} \langle \mathrm{A_+ c_+}\bar{\mathrm{c}}_+\Phi_+\Phi_+^*|\rho(t_{\mathrm{f}})|\mathrm{A_- c_-}\bar{\mathrm{c}}_-\Phi_-\Phi_-^*\rangle &=: \langle \mathbf{f}_+|\rho(t_{\mathrm{f}})|\mathbf{f}_-\rangle \\ &= \langle \mathbf{f}_+|\hat{U}(t_{\mathrm{f}}, t_{\mathrm{i}})\rho_{\mathrm{i}}\hat{U}^\dagger(t_{\mathrm{f}}, t_{\mathrm{i}})|\mathbf{f}_-\rangle \\ = \int \mathrm{d}[\mathbf{f}_{+\mathrm{i}}, \mathbf{f}_{-\mathrm{i}}]\langle \mathbf{f}_+|\hat{U}(t_{\mathrm{f}}, t_{\mathrm{i}})|\mathbf{f}_{+\mathrm{i}}\rangle\langle \mathbf{f}_{+\mathrm{i}}|\rho_{\mathrm{i}}|\mathbf{f}_{-\mathrm{i}}\rangle\langle \mathbf{f}_{-\mathrm{i}}|\hat{U}^\dagger(t_{\mathrm{f}}, t_{\mathrm{i}})|\mathbf{f}_-\rangle \end{aligned}$$

compact notation $\boldsymbol{f} = (A, c, \bar{c}, \Phi, \Phi^*)$ for the fields and $\mathbf{f}_j = \left(\mathrm{A}_j, \mathrm{c}_j, \bar{\mathrm{c}}_j, \Phi_j, \Phi_j^*\right)$ for their eigenstates

$$\langle \mathbf{f}_+|\rho(t_\mathrm{f})|\mathbf{f}_-\rangle = \int \mathrm{d}[\mathbf{f}_{+\mathrm{i}},\mathbf{f}_{-\mathrm{i}}]\langle \mathbf{f}_{+\mathrm{i}}|\rho_\mathrm{i}|\mathbf{f}_{-\mathrm{i}}\rangle \int_{\mathbf{f}_{+\mathrm{i}}}^{\mathbf{f}_+} \mathcal{D}[\boldsymbol{f}_+] \int_{\mathbf{f}_{-\mathrm{i}}}^{\mathbf{f}_-} \mathcal{D}[\boldsymbol{f}_-]\mu e^{iS_\mathrm{BRST}[\boldsymbol{f}_+]-iS_\mathrm{BRST}[\boldsymbol{f}_-]}$$





$$Z_{\mathrm{in-in}}[\boldsymbol{J}^+,\boldsymbol{J}^-] = \int \mathrm{d}[\mathbf{f},\mathbf{f}_{+\mathrm{i}},\mathbf{f}_{-\mathrm{i}}]\langle \mathbf{f}_{+\mathrm{i}}|\rho_\mathrm{i}|\mathbf{f}_{-\mathrm{i}}\rangle$$
$$\times \int_{\mathbf{f}_{+\mathrm{i}}}^{\mathbf{f}} \mathcal{D}[\boldsymbol{f}_+] \int_{\mathbf{f}_{-\mathrm{i}}}^{\mathbf{f}} \mathcal{D}[\boldsymbol{f}_-]\mu e^{iS_\mathrm{BRST}[\boldsymbol{f}_+]-iS_\mathrm{BRST}[\boldsymbol{f}_-]+i\int \mathrm{d}^4x(\boldsymbol{J}^+\cdot\boldsymbol{f}_+-\boldsymbol{J}^-\cdot\boldsymbol{f}_-)},$$

$$\boldsymbol{J}^\pm = \{J_\mathrm{a}^\pm\} = \left(J_A^\pm, J_c, J_{\bar{c}}^\pm, J_\Phi^\pm, J_{\Phi^*}^\pm\right)$$

$$\left.\frac{\delta^2 Z_{\mathrm{in-in}}}{\delta J_A^{-\mu}(x)\delta J_A^{+\nu}(y)}\right|_{J_\mathrm{a}^\pm=0} = \mathrm{Tr}\left(\hat{A}_\mu(x)\hat{A}_\nu(y)\rho_\mathrm{i}\right)$$

$$A_{\pm\mu}(x) \rightarrow A_{\pm\mu}(x) + \eta_\pm\partial_\mu c_\pm(x), c_\pm(x) \rightarrow c_\pm(x)$$
$$\Phi_\pm(x) \rightarrow \Phi_\pm(x) + iq\eta_\pm c_\pm(x)\Phi_\pm(x), \bar{c}_\pm(x) \rightarrow \bar{c}_\pm(x) - \frac{\eta_\pm}{\xi}\partial_\mu A_\pm^\mu(x)$$

$$\hat{s}A_{\pm\mu}(x) = \partial_\mu c_\pm(x)$$
$$\hat{s}\Phi_\pm(x) = iqc_\pm(x)\Phi_\pm(x)$$
$$\hat{s}c_\pm(x) = 0$$
$$\hat{s}\bar{c}_\pm(x) = -\frac{1}{\xi}\partial_\mu A_\pm^\mu(x)$$

$$S_\mathrm{BRST} \supset -\int \mathrm{d}^4x\left[\frac{1}{2\xi}\left(\partial_\mu A^\mu\right)^2 + \bar{c}\Box c\right]$$

$$S_\mathrm{BRST} \supset -\int \mathrm{d}^4x\left[\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \left(D_\mu\Phi\right)^* D^\mu\Phi + V(\Phi^*\Phi)\right]$$

$$\boldsymbol{f}_\mathrm{r} = \frac{\boldsymbol{f}_+ + \boldsymbol{f}_-}{2} \text{ and }, \boldsymbol{f}_\mathrm{a} = \boldsymbol{f}_+ - \boldsymbol{f}_-,$$

$$\boldsymbol{J}^\mathrm{r} = \frac{\boldsymbol{J}^+ + \boldsymbol{J}^-}{2} \text{ and } \boldsymbol{J}^\mathrm{a} = \boldsymbol{J}^+ - \boldsymbol{J}^-$$

$$Z_{\mathrm{in-in}}[\boldsymbol{J}^\mathrm{r},\boldsymbol{J}^\mathrm{a}] = \int \mathrm{d}[\mathbf{f},\mathbf{f}_{+\mathrm{i}},\mathbf{f}_{-\mathrm{i}}]\langle \mathbf{f}_{+\mathrm{i}}|\rho_\mathrm{i}|\mathbf{f}_{-\mathrm{i}}\rangle$$
$$\times \int_{(\mathbf{f}_{+\mathrm{i}}+\mathbf{f}_{-\mathrm{i}})/2}^{\mathbf{f}} \mathcal{D}[\boldsymbol{f}_\mathrm{r}] \int_{\mathbf{f}_{+\mathrm{i}}-\mathbf{f}_{-\mathrm{i}}}^{\mathbf{0}} \mathcal{D}[\boldsymbol{f}_\mathrm{a}]\mu e^{iS_\mathrm{BRST}\left[\boldsymbol{f}_\mathrm{r}+\frac{\boldsymbol{f}_\mathrm{a}}{2}\right]-iS_\mathrm{BRST}\left[\boldsymbol{f}_\mathrm{r}-\frac{\boldsymbol{f}_\mathrm{a}}{2}\right]-i\int \mathrm{d}^4x(\boldsymbol{J}^\mathrm{r}\cdot\boldsymbol{f}_\mathrm{a}+\boldsymbol{J}^\mathrm{a}\cdot\boldsymbol{f}_\mathrm{r})}$$

$$S_{\mathrm{BRST}}\left[\boldsymbol{f}_{\mathrm{r}}+\frac{\boldsymbol{f}_{\mathrm{a}}}{2}\right]-S_{\mathrm{BRST}}\left[\boldsymbol{f}_{\mathrm{r}}-\frac{\boldsymbol{f}_{\mathrm{a}}}{2}\right]\simeq\int\ \mathrm{d}^4x\left[\boldsymbol{f}_{\mathrm{a}}\cdot\frac{\delta S[\boldsymbol{f}_{\mathrm{r}}]}{\delta\boldsymbol{f}_{\mathrm{r}}}+(\square::\boldsymbol{f}_{\mathrm{a}})\right]$$

$$-\frac{\delta^2 Z_{\mathrm{in-in}}}{\delta J_A^{\mathrm{a}\mu}(x)\delta J_A^{\mathrm{r}\nu}(y)}\bigg|_{J_{\mathrm{a}}^{\pm}=0}=\theta(x^0-y^0)\mathrm{Tr}\big(\big[\hat{A}_\mu(x),\hat{A}_\nu(y)\big]\rho_{\mathrm{i}}\big),$$

$$A_{\mathrm{r}\mu}\to A_{\mathrm{r}\mu}+\eta\partial_\mu c_{\mathrm{r}},\ \text{and}\ A_{\mathrm{a}\mu}\to A_{\mathrm{a}\mu}+\eta\partial_\mu c_{\mathrm{a}}$$

$$\mathbf{G_0}=\begin{pmatrix}G_{++} & G_{+-}\\ G_{-+} & G_{--}\end{pmatrix}$$

$$\begin{aligned}G_{++}(x,y)&=\mathrm{Tr}[\rho\mathcal{T}\hat{\phi}(x)\hat{\phi}(y)]\\ G_{+-}(x,y)&=\mathrm{Tr}[\rho\hat{\phi}(y)\hat{\phi}(x)]\\ G_{-+}(x,y)&=\mathrm{Tr}[\rho\hat{\phi}(x)\hat{\phi}(y)]\\ G_{--}(x,y)&=\mathrm{Tr}[\rho\hat{\mathcal{T}}\hat{\phi}(x)\hat{\phi}(y)]\end{aligned}$$

$$\hat{\phi}(x)=\int\ \frac{\mathrm{d}^4k}{(2\pi)^4}\theta(k^0)2\pi\delta(k^2+m^2)\big(e^{ik\cdot x}\hat{a}_{\mathbf{k}}+e^{-ik\cdot x}\hat{a}_{\mathbf{k}}^{\dagger}\big)$$

$$\mathbf{G_0}(x,y)=\int\ \frac{\mathrm{d}^4k}{(2\pi)^4}e^{ik\cdot(x-y)}\mathbf{G_0}(k)$$

$$\mathbf{G_0}(k)=\begin{pmatrix}-\dfrac{i}{k^2+m^2-i\epsilon}+2\pi\delta(k^2+m^2)\mathfrak{n}(k) & \theta(-k^0)2\pi\delta(k^2+m^2)+2\pi\delta(k^2+m^2)\mathfrak{n}(k)\\ \theta(k^0)2\pi\delta(k^2+m^2)+2\pi\delta(k^2+m^2)\mathfrak{n}(k) & \dfrac{i}{k^2+m^2+i\epsilon}+2\pi\delta(k^2+m^2)\mathfrak{n}(k)\end{pmatrix},$$

with $\mathfrak{n}(k)=\theta(k^0)n(\mathbf{k})+\theta(-k^0)n(-\mathbf{k})$, and

$$n(\mathbf{k})2\omega_k(2\pi)^3\delta^3(\mathbf{k}-\mathbf{k}')=\mathrm{Tr}\big(\rho\hat{a}_{\mathbf{k}}^{\dagger}\hat{a}_{\mathbf{k}'}\big)$$

where $\omega_k=\sqrt{\mathbf{k}^2+m^2}$

$$\mathbf{G}_{0A}=\mathbf{G}_0-\begin{pmatrix}G_{-+} & G_{-+}\\ G_{-+} & G_{-+}\end{pmatrix}=\begin{pmatrix}G_A & -\Delta\\ 0 & G_R\end{pmatrix},$$

$$\Delta=G_{-+}-G_{+-}=G_R-G_A$$

$$G_R(x,y)=\theta(x^0-y^0)\Delta(x,y),\ G_A(x,y)=-\theta(y^0-x^0)\Delta(x,y)$$

$$\mathbf{G}_{0A}(k)=\begin{pmatrix}-\dfrac{i}{k^2+m^2-i\epsilon\sigma(k^0)} & -\sigma(k^0)2\pi\delta(k^2+m^2)\\ 0 & \dfrac{i}{k^2+m^2+i\epsilon\sigma(k^0)}\end{pmatrix}$$

$$\sigma(k^0)=\theta(k^0)-\theta(-k^0)$$

$$\frac{1}{x\mp i\epsilon}=P\left(\frac{1}{x}\right)\pm i\pi\delta(x)$$

$$(x-i\epsilon)(x+i\epsilon)\pi\delta(x)=\epsilon$$

$$(k^2+m^2-i\epsilon)(k^2+m^2+i\epsilon)\pi\delta(k^2+m^2)=\epsilon$$

$$\det[\mathbf{G_0}(k)]=\frac{1}{(k^2+m^2-i\epsilon)(k^2+m^2+i\epsilon)},$$

$$\mathbf{G_0^{-1}}(k)=\begin{pmatrix} i(k^2+m^2-i\epsilon)+2\epsilon\mathfrak{n}(k) & -\theta(k^0)2\epsilon-2\epsilon\mathfrak{n}(k) \\ -\theta(-k^0)2\epsilon-2\epsilon\mathfrak{n}(k) & -i(k^2+m^2+i\epsilon)+2\epsilon\mathfrak{n}(k)\end{pmatrix}$$

$$\det[\mathbf{G_0}]=\det[\mathbf{G_{0A}}]$$

$$S_{\text{in-in}}=\frac{i}{2}\int\frac{\mathrm{d}^4k}{(2\pi)^4}(\phi_+(-k)\phi_-(-k))\mathbf{G_0^{-1}}(k)\begin{pmatrix}\phi_+(k)\\ \phi_-(k)\end{pmatrix}$$
$$=\int\mathrm{d}^4x\left(-\frac{1}{2}(\partial\phi_+)^2-\frac{1}{2}m^2\phi_+^2+\frac{1}{2}(\partial\phi_-)^2+\frac{1}{2}m^2\phi_-^2\right)+S_{\mathrm{i}\epsilon}$$

$$S_{\mathrm{i}\epsilon}=\int\frac{\mathrm{d}^4k}{(2\pi)^4}i\epsilon\left[\phi_{\mathrm{a}}(-k)\frac{k^0}{|k^0|}\phi_{\mathrm{r}}(k)+\left(\frac{1}{2}+n(\mathbf{k})\right)|\phi_{\mathrm{a}}(k)|^2\right]$$

$$\theta(k^0)=\frac{1}{2}+\frac{1}{2}\frac{k^0}{|k^0|}$$

$$S_{\text{in-in}}=\int\mathrm{d}^4x\phi_{\mathrm{a}}(x)[\Box-m^2-2\epsilon\partial_t]\phi_{\mathrm{r}}(x)+i\epsilon\int\mathrm{d}^4x\int\mathrm{d}^4y\phi_{\mathrm{a}}(x)K_\phi(x,y)\phi_{\mathrm{a}}(y)$$

$$S_{\mathrm{i}\epsilon}=-\epsilon\int\mathrm{d}^4x(\phi_+(x)-\phi_-(x))\partial_t(\phi_+(x)+\phi_-(x))$$
$$+i\epsilon\int\mathrm{d}^4x\int\mathrm{d}^4y(\phi_+(x)-\phi_-(x))K_\phi(x,y)(\phi_+(y)-\phi_-(y))$$

$$K_\phi(x,y)=\int\frac{\mathrm{d}^4k}{(2\pi)^4}e^{ik\cdot(x-y)}(1+2\mathfrak{n}(k))\omega_k$$

$$\omega_k=\sqrt{\mathbf{k}^2+m^2}$$

$$K_\phi^\beta(x,y)=\int\frac{\mathrm{d}^4k}{(2\pi)^4}e^{ik\cdot(x-y)}\coth(\beta\omega_k/2)\omega_k$$

$$\lim_{\substack{t_{\mathrm{i}}\to-\infty\\ t_{\mathrm{f}}\to+\infty}}\int\mathrm{d}[\phi,\phi_{+\mathrm{i}},\phi_{-\mathrm{i}}]\int_{\phi_{+\mathrm{i}}}^{\phi}\mathcal{D}[\phi_+]\int_{\phi_{-\mathrm{i}}}^{\phi}\mathcal{D}[\phi_-]\langle\phi_{+\mathrm{i}}|\rho|\phi_{-\mathrm{i}}\rangle\mu e^{iS_{\text{in-in}}[\phi_+,\phi_-;t_{\mathrm{i}},t_{\mathrm{f}}]}$$
$$\equiv\int\mathcal{D}[\phi_+]\int\mathcal{D}[\phi_-]\mu e^{iS_{\text{in-in}}[\phi_+,\phi_-]+S_{i\epsilon}}$$

$$\hat\Phi(x)=\int\frac{\mathrm{d}^4k}{(2\pi)^4}\theta(k^0)2\pi\delta(k^2+m^2)\left(e^{ik\cdot x}\hat a_{\mathbf{k}}+e^{-ik\cdot x}\hat b_{\mathbf{k}}^\dagger\right)$$

$$n_+(\mathbf{k})2\omega_k(2\pi)^3\delta^3(\mathbf{k}-\mathbf{k}')=\mathrm{Tr}\left(\rho\hat a_{\mathbf{k}}^\dagger\hat a_{\mathbf{k}'}\right)$$
$$n_-(\mathbf{k})2\omega_k(2\pi)^3\delta^3(\mathbf{k}-\mathbf{k}')=\mathrm{Tr}\left(\rho\hat b_{\mathbf{k}}^\dagger\hat b_{\mathbf{k}'}\right)$$

$$\mathrm{Tr}(\rho\hat a_{\mathbf{k}}\hat a_{\mathbf{k}'})=\mathrm{Tr}\left(\rho\hat b_{\mathbf{k}}\hat b_{\mathbf{k}'}\right)=\mathrm{Tr}\left(\rho\hat a_{\mathbf{k}}\hat b_{\mathbf{k}'}^\dagger\right)=0$$

$$G_{++}(x,y) = \mathrm{Tr}\left[\rho\mathcal{T}\hat{\Phi}(x)\hat{\Phi}^{\dagger}(y)\right]$$
$$G_{+-}(x,y) = \mathrm{Tr}\left[\rho\hat{\Phi}^{\dagger}(y)\hat{\Phi}(x)\right]$$
$$G_{-+}(x,y) = \mathrm{Tr}\left[\rho\hat{\Phi}(x)\hat{\Phi}^{\dagger}(y)\right]$$
$$G_{--}(x,y) = \mathrm{Tr}\left[\rho\overline{\mathcal{T}}\hat{\Phi}(x)\hat{\Phi}^{\dagger}(y)\right]$$

$$\mathfrak{n}(k) = \theta(k^0)n_+(\mathbf{k}) + \theta(-k^0)n_-(-\mathbf{k}).$$

$$S_{\mathrm{in-in}} = \int \mathrm{d}^4x(\Phi_{\mathrm{a}}^*(x)[\Box - m^2 - 2\epsilon\partial_t]\Phi_{\mathrm{r}}(x) + \Phi_{\mathrm{r}}^*(x)[\Box - m^2 + 2\epsilon\partial_t]\Phi_{\mathrm{a}}(x))$$
$$+2i\epsilon\int \mathrm{d}^4x\int \mathrm{d}^4y\Phi_{\mathrm{a}}^*(x)K_{\Phi}(x,y)\Phi_{\mathrm{a}}(y)$$

$$K_{\Phi}(x,y) = \int \frac{\mathrm{d}^4k}{(2\pi)^4}e^{ik.(x-y)}(1 + 2\mathfrak{n}(k))\omega_k$$

$$\Phi_-^*(x)\Phi_+(x)$$

$$\Phi_-^*(x)\Phi_+(x) \to e^{iq\lambda_+(x) - iq\lambda_-(x)}\Phi_-^*(x)\Phi_+(x)$$

$$\hat{s}(\Phi_-^*(x)\Phi_+(x)) = iq(c_+(x) - c_-(x))\Phi_-^*(x)\Phi_+(x)$$

$$\lambda_{\pm} = \lambda_r \pm \frac{1}{2}\lambda_{\mathrm{a}} \text{ with } A_{\mu}^{\pm} = A_{\mu}^{\mathrm{r}} \pm \frac{1}{2}A_{\mu}^{\mathrm{a}}$$

$$A_{\mu}^{\mathrm{r,a}} \to A_{\mu}^{\mathrm{r,a}} + \partial_{\mu}\lambda_{\mathrm{r,a}}(x)$$

$$\Phi_{\pm}(x) = e^{\mp\frac{1}{2}iq\int_x^{x_{\mathrm{f}}}\mathrm{d}z^{\mu}A_{\mu}^{\mathrm{a}}(z)}\widetilde{\Phi}_{\pm}(x) = e^{\mp\frac{1}{2}iq(x_{\mathrm{f}}^{\mu} - x^{\mu})\int_0^1 \mathrm{d}sA_{\mu}^{\mathrm{a}}(x + s(x_{\mathrm{f}} - x))}\widetilde{\Phi}_{\pm}(x)$$

$$\Phi_+(t_{\mathrm{f}}) = \Phi_-(t_{\mathrm{f}}) \implies \widetilde{\Phi}_+(t_{\mathrm{f}}) = \widetilde{\Phi}_-(t_{\mathrm{f}})$$

$$\widetilde{\Phi}_{\pm}(x) \to e^{iq\lambda_{\mathrm{r}}(x)}\widetilde{\Phi}_{\pm}(x)$$

$$\Phi_{\pm}(x) \to e^{iq\lambda_{\mathrm{r}}(x)}\Phi_{\pm}(x)$$

$$\widetilde{\Phi}_{\pm}(x) \to e^{\pm i\frac{1}{2}q\lambda_{\mathrm{a}}(t_{\mathrm{f}})}\widetilde{\Phi}_{\pm}(x) = \widetilde{\Phi}_{\pm}(x)$$

$$\lambda_{\mathrm{a}}(t_{\mathrm{f}}) = \lambda_+(t_{\mathrm{f}}) - \lambda_-(t_{\mathrm{f}}) = 0$$

$$\Phi_{\pm}(x) \to e^{\pm i\frac{1}{2}q\lambda_{\mathrm{a}}(x)}\Phi_{\pm}(x)$$

$$\hat{s}A_{\mu}^{\mathrm{r}} = \partial_{\mu}c_{\mathrm{r}}(x), \hat{s}\widetilde{\Phi}_{\pm}(x) = iqc_{\mathrm{r}}(x)\widetilde{\Phi}_{\pm}(x), \hat{s}\widetilde{\Phi}_{\pm}^*(x) = -iqc_{\mathrm{r}}(x)\widetilde{\Phi}_{\pm}^*(x)$$

$$S_{i\epsilon} = -2\epsilon\int \mathrm{d}^4x\left(\widetilde{\Phi}_{\mathrm{a}}^*(x)D_t[A_{\mathrm{r}}]\widetilde{\Phi}_{\mathrm{r}}(x) - \widetilde{\Phi}_{\mathrm{r}}^*(x)D_t[A_r]\widetilde{\Phi}_{\mathrm{a}}(x)\right) + 2i\epsilon\int \mathrm{d}^4x\int \mathrm{d}^4y\widetilde{\Phi}_{\mathrm{a}}^*(x)K_{\widetilde{\Phi}}(x,y)\widetilde{\Phi}_{\mathrm{a}}(y)$$

$$K_{\widetilde{\Phi}}(x,y) \to e^{-iq\lambda_{\mathrm{r}}(x) + iq\lambda_{\mathrm{r}}(y)}K_{\widetilde{\Phi}}(x,y)$$

$$\widetilde{\Phi}_{\pm}(x) = e^{iq\int_{x_{\mathrm{i}}}^{x}\mathrm{d}z^{\mu}A_{\mu}^{\mathrm{r}}(z)}\breve{\Phi}_{\pm}(x) = e^{iq(x^{\mu} - x_{\mathrm{i}}^{\mu})\int_0^1 \mathrm{d}sA_{\mu}^{\mathrm{r}}(x_{\mathrm{i}} + s(x - x_{\mathrm{i}}))}\breve{\Phi}_{\pm}(x)$$

$$\breve{\Phi}_{\pm}(x) \to e^{iq\lambda_{\mathrm{r}}(x_{\mathrm{i}})}\breve{\Phi}_{\pm}(x),$$

$$S_{i\epsilon} = -2\epsilon \int \mathrm{d}^4x\left(\check{\Phi}^*_{\mathrm{a}}(x)\partial_t\check{\Phi}_{\mathrm{r}}(x) - \check{\Phi}^*_{\mathrm{r}}(x)\partial_t\check{\Phi}_{\mathrm{a}}(x)\right) + 2i\epsilon \int \mathrm{d}^4x \int \mathrm{d}^4y\check{\Phi}^*_{\mathrm{a}}(x)K_\Phi(x,y)\check{\Phi}_{\mathrm{a}}(y)$$

$$\tilde{A}^{\mathrm{r}}_\mu(x) = A^{\mathrm{r}}_\mu(x) - \partial_\mu\left((x^\nu - x^\nu_{\mathrm{i}})\int_0^1 \mathrm{d}sA^{\mathrm{r}}_\nu\left(x_{\mathrm{i}} + s(x - x_{\mathrm{i}})\right)\right)$$

$$\tilde{A}^{\mathrm{a}}_\mu(x) = A^{\mathrm{a}}_\mu(x) + \partial_\mu\left((x^\nu_{\mathrm{f}} - x^\nu)\int_0^1 \mathrm{d}sA^{\mathrm{a}}_\nu\left(x + s(x_{\mathrm{f}} - x)\right)\right)$$

$$\begin{aligned} S_{i\epsilon} =& -2\epsilon \int \mathrm{d}^4x\left(A_{\mathrm{a}\mu}(x)\partial_t A^\mu_{\mathrm{r}}(x) - \bar{c}_{\mathrm{a}}(x)\partial_t c_{\mathrm{r}}(x) + \bar{c}_{\mathrm{r}}(x)\partial_t c_{\mathrm{a}}(x)\right) \\ &+ i\epsilon \int \mathrm{d}^4x \int \mathrm{d}^4y\left(A_{\mathrm{a}\mu}(x)K_A^{\mu\nu}(x,y)A_{\mathrm{a}\nu}(y) + c_{\mathrm{a}}(y)K_c(x,y)\bar{c}_{\mathrm{a}}(x)\right) \end{aligned}$$

$$\begin{aligned} \hat{s}S_{i\epsilon} =& -2\epsilon \int \mathrm{d}^4x\left(\partial_\mu c_{\mathrm{a}}(x)\partial_t A^\mu_{\mathrm{r}}(x) + A_{\mathrm{a}\mu}(x)\partial_t\partial^\mu c_{\mathrm{r}}(x) + \partial_\mu A^\mu_{\mathrm{a}}(x)\partial_t c_{\mathrm{r}}(x) - \partial_\mu A^\mu_{\mathrm{r}}(x)\partial_t c_{\mathrm{a}}(x)\right) \\ &+ i\epsilon \int \mathrm{d}^4x \int \mathrm{d}^4y\left(2\partial_\mu c_{\mathrm{a}}(x)K_A^{\mu\nu}(x,y)A_{\nu\mathrm{a}}(y) + c_{\mathrm{a}}(y)K_c(x,y)\partial_\mu A^\mu_{\mathrm{a}}(x)\right) \end{aligned}$$

$$\hat{s}S_{i\epsilon} = -2i\epsilon \int \mathrm{d}^4x \int \mathrm{d}^4y c_{\mathrm{a}}(x)\left(2\partial_{x\mu}K_A^{\mu\nu}(x,y) + \partial^\nu_x K_c(x,y)\right)A_{\nu\mathrm{a}}(y)$$

$$2\partial_{x\mu}K_A^{\mu\nu}(x,y) + \partial^\nu_x K_c(x,y) = 0$$

$$2k_\mu K_A^{\mu\nu}(k) + k^\nu K_c(k) = 0$$

$$K_A^{\mu\nu}(k) = \eta^{\mu\nu} + 2T^{\mu\alpha}(k)\mathfrak{n}_{\alpha\beta}(k)T^{\beta\nu}(k), K_c(k) = -2,$$

$$T^{\mu\nu}(k) = \eta^{\mu\nu} - \frac{k^\mu u^\nu + u^\mu k^\nu}{u.k} + u^2\frac{k^\mu k^\nu}{(u.k)^2},$$

$$k_\mu T^{\mu\nu} = 0 \text{ and } T^{\mu\alpha}T^\nu_\alpha = T^{\mu\nu}$$

$$K_A^{\mu\nu}(k) = \eta^{\mu\nu}\left(1 + 2\mathfrak{n}_\beta(k)\right), K_c(k) = -2\left(1 + 2\mathfrak{n}_\beta(k)\right),$$

$$\mathfrak{n}_\beta(k) = \frac{1}{e^{\beta\omega_k} - 1}.$$

$$S_{\mathrm{BRST}}[A, c, \bar{c}, \Phi, \Phi^*] = S_{\mathcal{S}}[A, c, \bar{c}] + S_{\mathcal{E}}[\Phi, \Phi^*] + S_{\mathrm{int}}[A, \Phi, \Phi^*]$$

Density matrix:

$$\varrho_{\mathcal{S}} \equiv \mathrm{Tr}_{\mathcal{E}}[\rho]$$

$$\langle \mathrm{A}_+\mathrm{c}_+\bar{\mathrm{c}}_+|\varrho_{\mathcal{S}}(t_{\mathrm{f}})|\mathrm{A}_-\mathrm{c}_-\bar{\mathrm{c}}_-\rangle = \int \mathrm{d}[\Phi, \Phi^*]\langle \mathrm{A}_+\mathrm{c}_+\bar{\mathrm{c}}_+\Phi\Phi^*|\rho(t_{\mathrm{f}})|\mathrm{A}_-\mathrm{c}_-\bar{\mathrm{c}}_-\Phi\Phi^*\rangle$$

$$\rho_{\mathrm{i}} = \varrho_{\mathcal{S}\mathrm{i}} \otimes \varrho_{\mathcal{E}\mathrm{i}}$$

$$\langle \mathrm{A}_{+\mathrm{i}}\mathrm{c}_{+\mathrm{i}}\bar{\mathrm{c}}_{+\mathrm{i}}\Phi_{+\mathrm{i}}\Phi^*_{+\mathrm{i}}|\rho_{\mathrm{i}}|\ \mathrm{A}_{-\mathrm{i}}\mathrm{c}_{-\mathrm{i}}\bar{\mathrm{c}}_{-\mathrm{i}}\Phi_{-\mathrm{i}}\Phi^*_{-\mathrm{i}}\rangle = \langle \mathrm{A}_{+\mathrm{i}}\mathrm{c}_{+\mathrm{i}}\bar{\mathrm{c}}_{+\mathrm{i}}|\varrho\varrho_{\mathcal{S}}|\mathrm{A}_{-\mathrm{i}}\mathrm{c}_{-\mathrm{i}}\bar{\mathrm{c}}_{-\mathrm{i}}\rangle \cdot \langle \Phi_{+\mathrm{i}}\Phi^*_{+\mathrm{i}}|\varrho\mathcal{E}_{\mathrm{i}}|\Phi_{-\mathrm{i}}\Phi^*_{-\mathrm{i}}\rangle$$

$$\langle \mathrm{A}_+\mathrm{c}_+\bar{\mathrm{c}}_+|\varrho_{\mathcal{S}}(+\infty)|\mathrm{A}_-\mathrm{c}_-\bar{\mathrm{c}}_-\rangle$$
$$= \int \mathrm{d}[\,\mathrm{A}_{\pm\mathrm{i}}, \mathrm{c}_{\pm\mathrm{i}}, \bar{\mathrm{c}}_{\pm\mathrm{i}}]\langle \mathrm{A}_{+\mathrm{i}}\mathrm{c}_{+\mathrm{i}}\bar{\mathrm{c}}_{+\mathrm{i}}|\varrho_{\mathcal{S}}|\mathrm{A}_{-\mathrm{i}}\mathrm{c}_{-\mathrm{i}}\bar{\mathrm{c}}_{-\mathrm{i}}\rangle$$
$$\times \int_{\mathrm{A}_{+\mathrm{i}}\mathrm{c}_{+\mathrm{i}}\bar{\mathrm{c}}_{+\mathrm{i}}}^{\mathrm{A}_+\mathrm{c}_+\bar{\mathrm{c}}_+} \mathcal{D}[A_+, c_+, \bar{c}_+] \int_{\mathrm{A}_{-\mathrm{i}}\mathrm{c}_{-\mathrm{i}}\bar{\mathrm{c}}_{-\mathrm{i}}}^{\mathrm{A}_-\mathrm{c}_-\bar{\mathrm{c}}_-} \mathcal{D}[A_-, c_-, \bar{c}_-]\tilde{\mu}e^{iS_{\mathcal{S}}[A_+,c_+,\bar{c}_+]-iS_{\mathcal{S}}[A_-,c_-,\bar{c}_-]+iS_{\mathrm{IF}}[A_+,A_-]}$$

$$\tilde{\mu}e^{iS_{\mathrm{IF}}[A_+,A_-]}$$
$$= \int \mathrm{d}\big[\Phi, \Phi^*, \Phi_{\pm\mathrm{i}}, \Phi^*_{\pm\mathrm{i}}\big]\langle \Phi_{+\mathrm{i}}\Phi^*_{+\mathrm{i}}|\varrho\mathcal{E}_{\mathrm{i}}|\Phi_{-\mathrm{i}}\Phi^*_{-\mathrm{i}}\rangle$$
$$\times \int_{\Phi_{+\mathrm{i}}\Phi^*_{+\mathrm{i}}}^{\Phi\Phi^*} \mathcal{D}[\Phi_+, \Phi^*_+] \int_{\Phi_{-\mathrm{i}}\Phi^*_{-\mathrm{i}}}^{\Phi\Phi^*} \mathcal{D}[\Phi_-, \Phi^*_-]\mu e^{iS_{\mathcal{E}}[\Phi_+,\Phi^*_+]+iS_{\mathrm{int}}[A_+,\Phi_+,\Phi^*_+]-iS_{\mathcal{E}}[\Phi_-,\Phi^*_-]-iS_{\mathrm{int}}[A_-,\Phi_-,\Phi^*_-]}$$

$$Z_{\mathcal{S}}[J_A^+, J_c^+, J_{\bar{c}}^+, J_A^-, J_c^-, J_{\bar{c}}^-]$$
$$= \int \mathrm{d}\big[\,\mathrm{A}, \mathrm{c}, \bar{\mathrm{c}}', \mathrm{A}_{\pm\mathrm{i}}, \mathrm{c}_{\pm\mathrm{i}}, \bar{\mathrm{c}}_{\pm\mathrm{i}}\big]\langle \mathrm{A}_{+\mathrm{i}}\mathrm{c}_{+\mathrm{i}}\bar{\mathrm{c}}_{+\mathrm{i}}|\varrho\varrho_i|\mathrm{A}_{-\mathrm{i}}\mathrm{c}_{-\mathrm{i}}\bar{\mathrm{c}}_{-\mathrm{i}}\rangle \int_{\mathrm{A}_{+\mathrm{i}}\mathrm{c}_{+\mathrm{i}}\bar{\mathrm{c}}_{+\mathrm{i}}}^{\mathrm{Ac}\bar{\mathrm{c}}} \mathcal{D}[A_+, c_+, \bar{c}_+] \int_{\mathrm{A}_{-\mathrm{i}}\mathrm{c}_{-\mathrm{i}}\bar{\mathrm{c}}_{-\mathrm{i}}}^{\mathrm{Ac}\bar{\mathrm{c}}} \mathcal{D}[A_-, c_-, \bar{c}_-]$$
$$\times \tilde{\mu}e^{iS_{\mathcal{S}}[A_+,c_+,\bar{c}_+]-iS_{\mathcal{S}}[A_-,c_-,\bar{c}_-]+iS_{\mathrm{IF}}[A_+,c_+,\bar{c}_+,A_-,c_-,\bar{c}_-]-i\int \mathrm{d}^4x\left(J_A^+\cdot A_+ + J_{\bar{c}}^+ c_+ + J_{\bar{c}}^+\bar{c}_+ - J_A^-\cdot A_- - J_{\bar{c}}^- c_- - J_{\bar{c}}^-\bar{c}_-\right)}$$

$$Z_{\mathcal{S}}[J_A^+, J_c^+, J_{\bar{c}}^+, J_A^-, J_c^-, J_{\bar{c}}^-] = Z_{\mathrm{in-in}}[J_A^+, J_c^+, J_{\bar{c}}^+, 0,0, J_A^-, J_c^-, J_{\bar{c}}^-, 0,0]$$

$$\hat{s}S_{\text{influence functional (IF)}}[A_+, A_-, \dots] = 0$$

$$\hat{s} = \hat{s}_{\mathcal{S}} + \hat{s}_{\mathcal{E}}$$

$$\hat{s}_{\mathcal{E}}A_{\pm\mu}(x) = \hat{s}_{\mathcal{E}}c_{\pm}(x) = \hat{s}_{\mathcal{E}}\bar{c}_{\pm}(x) = 0, \text{ and } \hat{s}_{\mathcal{E}}\Phi_{\pm}(x) = iqc_{\pm}(x)\Phi_{\pm}(x)$$

$$e^{iS_{\mathcal{S}}+iS_{\mathrm{IF}}[A_+,A_-,\dots]} = \int \mathcal{D}[\Phi_+, \Phi^*_+, \Phi_-, \Phi^*_-]e^{iS_{\mathrm{BRST}}+iS_{i\epsilon}}$$

$$\hat{s}_{\mathcal{S}}S_{\mathrm{IF}}[A_+, A_-, \dots] = e^{-iS_{\mathcal{S}}-iS_{\mathrm{IF}}[A_+,A_-,\dots]}\int \mathcal{D}[\Phi_+, \Phi^*_+, \Phi_-, \Phi^*_-]\big(\hat{s}_{\mathcal{S}}(S_{\mathrm{BRST}} + iS_{i\epsilon})\big)e^{iS_{\mathrm{BRST}}+iS_{i\epsilon}}$$

$$\int \mathcal{D}[\Phi_+, \Phi^*_+, \Phi_-, \Phi^*_-]\big(\hat{s}_{\mathcal{E}}(S_{\mathrm{BRST}} + iS_{i\epsilon})\big)e^{iS_{\mathrm{BRST}}+iS_{i\epsilon}} = 0$$

$$\hat{s}_{\mathcal{S}}S_{\mathrm{IF}}[A_+, A_-, \dots] = e^{-iS_{\mathcal{S}}-iS_{\mathrm{IF}}[A_+,A_-,\dots]}\int \mathcal{D}[\Phi_+, \Phi^*_+, \Phi_-, \Phi^*_-]\big(\hat{s}(S_{\mathrm{BRST}} + iS_{i\epsilon})\big)e^{iS_{\mathrm{BRST}}+iS_{i\epsilon}}$$

$$\hat{s}(S_{\mathrm{BRST}} + iS_{i\epsilon}) = 0 \Rightarrow \hat{s}_{\mathcal{S}}S_{\mathrm{IF}}[A_+, A_-, \dots] \equiv \hat{s}S_{\mathrm{IF}}[A_+, A_-, \dots] = 0$$

$$\hat{s}S_{\mathrm{IF}}[A_+, A_-] = \int \mathrm{d}^4x\left[\partial_\mu c_+\frac{\delta S_{\mathrm{IF}}}{\delta A_{+\mu}} + \partial_\mu c_-\frac{\delta S_{\mathrm{IF}}}{\delta A_{-\mu}}\right] = -\int \mathrm{d}^4x\left[c_+\partial_\mu\frac{\delta S_{\mathrm{IF}}}{\delta A_{+\mu}} + c_-\partial_\mu\frac{\delta S_{\mathrm{IF}}}{\delta A_{-\mu}}\right] = 0$$

$$\partial_\mu\frac{\delta S_{\mathrm{IF}}}{\delta A_{+\mu}} = 0, \text{ and } \partial_\mu\frac{\delta S_{\mathrm{IF}}}{\delta A_{-\mu}} = 0$$

$$\chi_\pm \to \chi_\pm + \alpha\lambda_\pm(x)$$

$$\hat{s}\chi_\pm = \alpha c_\pm(x)$$

$$\hat{s}S_{\mathrm{IF}}[A_+,A_-,\chi_+,\chi_-] = \int \mathrm{d}^4x\left[\partial_\mu c_+\frac{\delta S_{\mathrm{IF}}}{\delta A_{+\mu}}+\partial_\mu c_-\frac{\delta S_{\mathrm{IF}}}{\delta A_{-\mu}}+\alpha\chi_+\frac{\delta S_{\mathrm{IF}}}{\delta \chi_+}+\alpha\chi_-\frac{\delta S_{\mathrm{IF}}}{\delta \chi_-}\right]$$
$$= -\int \mathrm{d}^4x\left[c_+\left(\partial_\mu\frac{\delta S_{\mathrm{IF}}}{\delta A_{+\mu}}-\alpha\frac{\delta S_{\mathrm{IF}}}{\delta \chi_+}\right)+c_-\left(\partial_\mu\frac{\delta S_{\mathrm{IF}}}{\delta A_{-\mu}}-\alpha\frac{\delta S_{\mathrm{IF}}}{\delta \chi_-}\right)\right]=0$$

$$\partial_\mu\frac{\delta S_{\mathrm{IF}}}{\delta A_{+\mu}}-\alpha\frac{\delta S_{\mathrm{IF}}}{\delta \chi_+}=0 \text{ and } \partial_\mu\frac{\delta S_{\mathrm{IF}}}{\delta A_{-\mu}}-\alpha\frac{\delta S_{\mathrm{IF}}}{\delta \chi_-}=0$$

$$A'_\mu(x)=A_\mu(x)+\partial_\mu\lambda(x),\Phi'(x)=e^{iq\lambda(x)}\Phi(x).$$

$$\lambda(x)=\frac{1}{q}\theta+\xi(x)$$

$$A'_\mu(x)=A_\mu(x)+\partial_\mu\xi(x),\Phi'(x)=e^{i\theta}\Phi(x)$$

$$c_\pm(x)=\frac{1}{q}c_0+C_\pm(x)$$

$$\hat{s}A_{\pm\mu}=\partial_\mu C_\pm,\hat{s}C_\pm=0,\hat{s}\bar{c}_\pm=-\frac{1}{\xi}\partial_\mu A^\mu_\pm,\hat{s}\Phi^\pm=ic_0\Phi^\pm$$

$$\tilde{\mu}e^{i\tilde{S}_{\mathrm{IF}}[A_+,A_-,\chi_+,\chi_-]}=\int \mathrm{d}\left[\mathrm{H},\varphi,\varphi^*,\mathrm{H}_{\pm\mathrm{i}},\varphi_{\pm\mathrm{i}},\varphi^*_{\pm\mathrm{i}}\right]\langle \mathrm{H}_{+\mathrm{i}}\varphi_{+\mathrm{i}}\varphi^*_{+\mathrm{i}}|\varrho_{\mathcal{E}\mathrm{i}}|\mathrm{H}_{-i}\varphi_{-\mathrm{i}}\varphi^*_{-\mathrm{i}}\rangle\int_{\mathrm{H}_{+\mathrm{i}}\varphi_{+\mathrm{i}}\varphi^*_{+\mathrm{i}}}^{\mathrm{H}\varphi\varphi^*}\mathcal{D}[H_+,\varphi_+,\varphi^*_+]$$
$$\int_{\mathrm{H}_{-\mathrm{i}}\varphi_{-\mathrm{i}}\varphi^*_{-\mathrm{i}}}^{\mathrm{H}\varphi\varphi^*}\mathcal{D}[H_-,\varphi_-,\varphi^*_-]\mu e^{iS_{\mathcal{E}}[H_+,\varphi_+,\varphi^*_+]+iS_{\mathrm{int}}[H_+,A_+,\varphi_+,\varphi^*_+]-iS_{\mathcal{E}}[H_-,\varphi_-,\varphi^*_-]-iS_{\mathrm{int}}[H_-,A_-,\varphi_-,\varphi^*_-]}$$

$$\breve{\mu}[A_+,A_-]e^{iS_{\mathrm{IF}}[A_+,A_-]}=\int \mathrm{d}[\chi,\chi_{\pm\mathrm{i}}]\langle\chi_{+\mathrm{i}}|\varrho_{\chi\mathrm{i}}|\chi_{-\mathrm{i}}\rangle$$
$$\int_{\chi_{+\mathrm{i}}}^{\chi}\mathcal{D}[\chi_+]\int_{\chi_{-\mathrm{i}}}^{\chi}\mathcal{D}[\chi_-]\tilde{\mu}[A_+,A_-,\chi_+,\chi_-]e^{i\tilde{S}_{\mathrm{IF}}[A_+,A_-,\chi_+,\chi_-]}$$

$$iS_{\mathrm{IF}}[A_+,A_-]\equiv W[A_1,A_2],$$
$$i\tilde{S}_{\mathrm{IF}}[A_+,A_-,\chi_+,\chi_-]\equiv iI[B_1,B_2],$$

$$\ddot{\phi}+\gamma\dot{\phi}-\nabla^2\phi+m^2\phi=0.$$

$$E=\frac{1}{2}\int \mathrm{d}^{\square}\mathbf{x}\left(\dot{\phi}^2+(\nabla\phi)^2+m^2\phi^2\right)$$

$$\frac{\mathrm{d}E}{\mathrm{d}t}=-\gamma\int \mathrm{d}^{\square}\mathbf{x}\dot{\phi}^2<0$$

$$S=\int \mathrm{d}^4xe^{\gamma t}\left(-\frac{1}{2}(\partial\phi)^2-\frac{1}{2}m^2\phi^2\right)$$

$$S=\int \mathrm{d}^4xa^3(t)\left(\frac{1}{2}\dot{\phi}^2-\frac{1}{2a^2(t)}(\nabla\phi)^2-\frac{1}{2}m^2\phi^2\right)$$

$$\ddot{\phi}+\frac{3\dot{a}(t)}{a(t)}\dot{\phi}-\frac{1}{a^2(t)}\nabla^2\phi+m^2\phi=0$$

$$\ddot{\phi} + \gamma(-\nabla^2)\dot{\phi} - \nabla^2\phi + m^2\phi = 0.$$

$$[-\omega^2 - i\omega\gamma(\mathbf{k}^2) + \mathbf{k}^2 + m^2]\phi(k) = 0,$$

$$S = \frac{1}{2}\int \mathrm{d}^4x\left(-\partial_\mu\phi e^{\gamma(-\nabla^2)t}\partial^\mu\phi - m^2\phi e^{\gamma(-\nabla^2)t}\phi\right)$$

$$= \frac{1}{2}\int \mathrm{d}t \int \frac{\mathrm{d}^{\square}\mathbf{k}}{(2\pi)^3} e^{\gamma(\mathbf{k}^2)t}\left(\left|\dot{\phi}_{\mathbf{k}}(t)\right|^2 - (\mathbf{k}^2 + m^2)|\phi_{\mathbf{k}}(t)|^2\right)$$

$$S = \frac{1}{2}\int \mathrm{d}t \int \frac{\mathrm{d}^{\square}\mathbf{k}}{(2\pi)^3} \sum_{n,m=0}^{\infty} c_{nm}(\mathbf{k}^2)\partial_t^n\phi_{-\mathbf{k}}(t)e^{\gamma(\mathbf{k}^2)t}\partial_t^m\phi_{\mathbf{k}}(t)$$

$$\sum_{n,m=0}^{\infty} c_{nm}(\mathbf{k}^2)\left(-\partial_t - \gamma(\mathbf{k}^2)\right)^n\partial_t^m\phi_{\mathbf{k}}(t) = 0$$

$$\sum_{n,m=0}^{\infty} c_{nm}(\mathbf{k}^2)\left(i\omega - \gamma(\mathbf{k}^2)\right)^n(-i\omega)^m\phi(k) = 0$$

$$S = \int \mathrm{d}^4x e^{\gamma t}\left(-\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + J_\mu A^\mu\right)$$

$$\partial_\mu F^{\mu\nu} + \gamma u_\mu F^{\mu\nu} = -J^\nu$$

$$A_\mu \to A_\mu + \partial_\mu\lambda$$

$$\partial_\mu(e^{\gamma t}J^\mu) = 0 \;\Rightarrow\; \partial_\mu J^\mu + \gamma u_\mu J^\mu = 0$$

$$S_{\mathrm{in-in}} = \int \mathrm{d}^4x e^{\gamma t}\left(-\frac{1}{4}F_{+\mu\nu}^{+}F_{+}^{\mu\nu} + J_\mu^{+}A_{+}^{\mu} + \frac{1}{4}F_{-\mu\nu}^{-}F_{-}^{\mu\nu} - J_\mu^{-}A_{-}^{\mu}\right)$$

$$S_{\mathrm{in-in}} = \int \mathrm{d}^4x e^{\gamma t}\left(-\frac{1}{2}F_{\mu\nu}^{\mathrm{a}}F_{\mathrm{r}}^{\mu\nu} + J_\mu^{\mathrm{r}}A_{\mathrm{a}}^{\mu} + J_\mu^{\mathrm{a}}A_{\mathrm{r}}^{\mu}\right)$$

$$A_\mu^{\mathrm{a}} = e^{-\gamma t}a_\mu, \text{ and } J_\mu^{\mathrm{a}} = e^{-\gamma t}j_\mu^{\mathrm{a}}$$

$$S_{\mathrm{in-in}} = \int \mathrm{d}^4x\left(-\frac{1}{2}(B_{\mu\nu} - 2\gamma u_\mu a_\nu)F_{\mathrm{r}}^{\mu\nu} + J_\mu^{\mathrm{r}}a^\mu + j_\mu^{\mathrm{a}}A^\mu\right)$$

with $B_{\mu\nu} = \partial_\mu a_\nu - \partial_\nu a_\mu$

$$\delta a_\mu = \partial_\mu\lambda - \gamma u_\mu\lambda$$

$$\delta a_\mu(-k) = iv_\mu\lambda(-k),$$

$v^\mu = (i\gamma_2, -\mathbf{k})$ and $\gamma_2 = \gamma - i\omega$

$$\partial_\mu J_{\mathrm{r}}^\mu + \gamma u_\mu J_{\mathrm{r}}^\mu = 0$$

$$\delta A_\mu^{\mathrm{a}} = \partial_\mu(e^{-\gamma t}\lambda),$$

$$\partial_\mu\left(e^{\gamma t}\left(J_r^\mu+\xi^\mu\right)\right)=0$$

$$S_{\text{in-in}}=\int\frac{\mathrm{d}^4k}{(2\pi)^4}\left(a^0(-k)ik_iF_{\mathrm{r}}^{0i}(k)+a_i(-k)\left(\gamma_2(k)F_{\mathrm{r}}^{0i}(k)-\gamma_3(k)ik_jF_{\mathrm{r}}^{ij}(k)+\gamma_4(k)\epsilon^{ijk}F_{jk}^{\mathrm{r}}(k)\right)\right)$$

$$\gamma_2(k)=\hat{\Gamma}(k)-i\omega$$

$$S_{\text{in-in}}=\int\frac{\mathrm{d}^4k}{(2\pi)^4}\left(a_i(-k)\hat{\Gamma}(k)F_{\mathrm{r}}^{0i}(k)-B_{0i}(-k)F_{\mathrm{r}}^{0i}(k)+\frac{1}{2}B_{ij}(-k)\gamma_3(k)F_{\mathrm{r}}^{ij}(k)+\gamma_4(k)\epsilon^{ijk}a_i(-k)F_{jk}^{r}(k)\right)$$

$$\delta a_\mu(k)=ik_\mu\lambda(k)-\hat{\Gamma}(-k)u_\mu\lambda(k)$$

$$\delta B_{\mu\nu}(k)=-i\hat{\Gamma}(-k)\left(k_\mu u_\nu-u_\mu k_\nu\right)\lambda(k)$$

$$\hat{\Gamma}(-k)=\sum_{n=0}^{\infty}(-i\omega)^n\Gamma_n(\mathbf{k}^2)$$

$$\delta a_0(t,\mathbf{k})=\partial_t\lambda(t,\mathbf{k})-\sum_{n=0}^{\infty}\Gamma_n(\mathbf{k}^2)\partial_t^n\lambda(t,\mathbf{k})$$

$$\lambda(t,\mathbf{k})=e^{\alpha(\mathbf{k}^2)t}\Lambda(t,\mathbf{k})$$

$$\alpha(\mathbf{k}^2)-\sum_{n=0}^{\infty}\Gamma_n(\mathbf{k}^2)\alpha^n(\mathbf{k}^2)=0$$

$$\begin{aligned}\delta a_0(t,\mathbf{k})&=e^{\alpha(\mathbf{k}^2)t}\left[\partial_t\Lambda(t,\mathbf{k})-\sum_{n=1}^{\infty}\Gamma_n(\mathbf{k}^2)((\alpha+\partial_t)^n-\alpha^n)\Lambda(t,\mathbf{k})\right]\\&=e^{\alpha(\mathbf{k}^2)t}\left[\partial_t\Lambda(t,\mathbf{k})-\sum_{n=1}^{\infty}\sum_{m=1}^{n}\Gamma_n(\mathbf{k}^2)\frac{n!}{m!\,(n-m)!}\alpha^{n-m}\partial_t^m\Lambda(t,\mathbf{k})\right]\end{aligned}$$

$$a_i(t,\mathbf{k})=e^{\alpha(\mathbf{k}^2)t}A_i^{\mathrm{a}}(t,\mathbf{k})$$

$$a_0(t,\mathbf{k})=e^{\alpha(\mathbf{k}^2)t}\left(1-\sum_{n=1}^{\infty}\sum_{m=1}^{n}\Gamma_n(\mathbf{k}^2)\frac{n!}{m!\,(n-m)!}\alpha^{n-m}\partial_t^{m-1}\right)A_0^{\mathrm{a}}(t,\mathbf{k})$$

$$\delta A_\mu^{\mathrm{a}}=\partial_\mu\Lambda$$

$$S=\int\mathrm{d}^4x\left[-\frac{1}{4}F_{\mu\nu}^2-|D[A]\Phi|^2-m^2|\Phi|^2+\sum_I\left(|D_t[A]\varphi_I|^2-\Gamma_I^2|\varphi_I|^2+g_I(\varphi_I^*\Phi+\Phi^*\varphi_I)\right)\right]$$

$$\Phi\to e^{iq\lambda}\Phi,\varphi_I\to e^{iq\lambda}\varphi_I$$

$$e^{iS_{\mathrm{IF}}[A_+,A_-,\Phi_+,\Phi_-,\Phi_+^*,\Phi_-^*]}=\int\mathcal{D}[\varphi_I^+,\varphi_I^-]$$

$$\frac{\mu}{\tilde{\mu}}e^{i\int\mathrm{d}^4x\sum_I\left(|D_t[A_+]\varphi_I^+|^2-\Gamma_I^2|\varphi_I^+|^2+g_I(\varphi_I^{+*}\Phi^++\Phi^{+*}\varphi_I^+)-|D_t[A_-]\varphi_I^-|^2+\Gamma_I^2|\varphi_I^-|^2-g_I(\varphi_I^{-*}\Phi^-+\Phi^{-*}\varphi_I^-)\right)+iS_{i\epsilon}}$$

$$\partial_t \lambda^{\mathrm{a}}(x) = A_0^{\mathrm{a}}(x)$$

$$\lambda^{\mathrm{a}}(t,\mathbf{x}) = -\int_t^{t_{\mathrm{f}}} \mathrm{d}t' A_0^{\mathrm{a}}(t',\mathbf{x})$$

$$\varphi_I^{\pm}(x) = U_{\pm}(x)\tilde{\varphi}_I^{\pm}(x)$$

$$U_{\pm}(x) = e^{\mp\frac{i}{2}q\int_t^{t_{\mathrm{f}}} \mathrm{d}t' A_0^{\mathrm{a}}(t',\mathbf{x})}$$

$$e^{iS_{\mathrm{IF}}[A_+,A_-,\Phi_+,\Phi_-,\Phi_+^*,\Phi_-^*]} = \int \mathcal{D}[\tilde{\varphi}_I^+,\tilde{\varphi}_I^-]$$

$$\frac{\mu}{\tilde{\mu}} e^{i\int \mathrm{d}^4x \sum_I \left(|D_t[A_{\mathrm{r}}]\tilde{\varphi}_I^+|^2 - \Gamma_I^2|\tilde{\varphi}_I^+|^2 + g_I(\tilde{\varphi}_I^{+*}\tilde{\Phi}^+ + \tilde{\Phi}^{+*}\tilde{\varphi}_I^+) - |D_t[A_{\mathrm{r}}]\tilde{\varphi}_I^-|^2 + \Gamma_I^2|\tilde{\varphi}_I^-|^2 - g_I(\tilde{\varphi}_I^{-*}\tilde{\Phi}^- + \tilde{\Phi}^{-*}\tilde{\varphi}_I^-)\right) + iS_{i\epsilon}}$$

$$\Phi^{\pm}(x) = U_{\pm}(x)\tilde{\Phi}^{\pm}(x)$$

$$S_{\mathrm{IF}}[A_+,A_-,\Phi_+,\Phi_-,\Phi_+^*,\Phi_-^*] = -i\sum_I \mathrm{Trlog}\ \mathbf{J}_I[A_{\mathrm{r}}] + i\sum_I \mathrm{Trlog}\ \mathbf{J}_{IA}[A_{\mathrm{r}}]$$
$$+i\sum_I g_I^2 \int \mathrm{d}^4x \int \mathrm{d}^4y \left(\tilde{\Phi}_+^*(x) - \tilde{\Phi}_-^*(x)\right)\mathbf{J}_I[A_{\mathrm{r}}](x,y)\begin{pmatrix}\tilde{\Phi}_+(y)\\ -\tilde{\Phi}_-(y)\end{pmatrix}$$

$$\mathbf{J}_I(x,y) = \begin{pmatrix} J_I^{++}(x,y) & J_I^{+-}(x,y) \\ J_I^{-+}(x,y) & J_I^{--}(x,y)\end{pmatrix},$$

$$\begin{aligned} J_I^{++}(x,y) &= \mathrm{Tr}\left[\rho\mathcal{T}\hat{\varphi}_I(x)\hat{\varphi}_I^{\dagger}(y)\right] \\ J_I^{+-}(x,y) &= \mathrm{Tr}\left[\rho\hat{\varphi}_I^{\dagger}(y)\hat{\varphi}_I(x)\right] \\ J_I^{-+}(x,y) &= \mathrm{Tr}\left[\rho\hat{\varphi}_I(x)\hat{\varphi}_I^{\dagger}(y)\right] \\ J_I^{--}(x,y) &= \mathrm{Tr}\left[\rho\overline{\mathcal{T}}\hat{\varphi}_I(x)\hat{\varphi}_I^{\dagger}(y)\right] \end{aligned}$$

$$\mathbf{J}_{IA}(x,y) = \mathbf{J}_I(x,y) - \begin{pmatrix} J_I^{-+}(x,y) & J_I^{-+}(x,y) \\ J_I^{-+}(x,y) & J_I^{-+}(x,y)\end{pmatrix}$$

$$D_t[A_{\mathrm{r}}]^2\hat{\tilde{\varphi}}_I(x) = -\Gamma_I^2\hat{\tilde{\varphi}}_I(x)$$

$$\partial_t \lambda^{\mathrm{r}}(x) = A_0^{\mathrm{r}}(x)$$

$$\lambda^{\mathrm{r}}(t,\mathbf{x}) = \int_{t_{\mathrm{i}}}^{t} \mathrm{d}t' A_0^{\mathrm{r}}(t',\mathbf{x}) + \lambda_{\mathrm{i}}^{\mathrm{r}}(\mathbf{x})$$

$$\hat{\tilde{\varphi}}_I(x) = V(x)\frac{1}{\sqrt{2\Gamma_I}}\left(e^{-i\Gamma_I t}\hat{a}_I(\mathbf{x}) + e^{i\Gamma_I t}\hat{b}_I^{\dagger}(\mathbf{x})\right)$$

$$V(x) = e^{iq\int_{t_{\mathrm{i}}}^{t} \mathrm{d}t' A_0^{\mathrm{r}}(t',\mathbf{x})}$$

$$\left[\hat{a}_I(\mathbf{x}),\hat{a}_J^{\dagger}(\mathbf{y})\right] = \delta_{IJ}\delta^3(\mathbf{x}-\mathbf{y}), \left[\hat{b}_I(\mathbf{x}),\hat{b}_J^{\dagger}(\mathbf{y})\right] = \delta_{IJ}\delta^3(\mathbf{x}-\mathbf{y})$$

$$\mathrm{Tr}\left[\rho\hat{a}_I^{\dagger}(\mathbf{x})\hat{a}_J(\mathbf{y})\right] = \delta_{IJ}W(\mathbf{x},\mathbf{y})H_I(\mathbf{x},\mathbf{y})$$

$$W(\mathbf{x},\mathbf{y}) = e^{iq\int_0^1 \mathrm{d}s \frac{\mathrm{d}z^i}{\mathrm{d}s} A_i(t_\mathrm{i},\mathbf{z}(s))}$$

$$z(s) = \left(t_\mathrm{i}, y^i + s\left(x^i - y^i\right)\right)$$

$$\mathbf{J}_I(x,y) = \mathcal{W}^\mathrm{r}_{C_1}(x,y)\mathbf{D}_I(x-y)$$

$$\mathcal{W}^\mathrm{r}_{C_1}(x,y) = e^{iq\int_{C_1} A^\mathrm{r}_\mu(z)\mathrm{d}z^\mu} = V(x)W(\mathbf{x},\mathbf{y})V(y)^*$$

$$\mathbf{U}(x) = \begin{pmatrix} U_+(x) & 0 \\ 0 & U_-(x) \end{pmatrix},$$

$$S_\mathrm{IF}[A_+,A_-,\Phi_+,\Phi_-,\Phi_+^*,\Phi_-^*] = -i\sum_I \mathrm{Trlog}\ \mathbf{D}_I + i\sum_I \mathrm{Trlog}\ \mathbf{D}_{IA}$$
$$+i\sum_I g_I^2 \int \mathrm{d}^4x \int \mathrm{d}^4y (\Phi_+^*(x) - \Phi_-^*(x))\mathcal{W}^\mathrm{r}_{C_1}(x,y)\mathbf{U}(x)\mathbf{D}_I(x,y)\mathbf{U}^\dagger(y)\begin{pmatrix}\Phi_+(y)\\ -\Phi_-(y)\end{pmatrix}$$

$$\mathbf{D}_I(\omega,\mathbf{k}) = \begin{pmatrix} -\frac{i}{-\omega^2+\Gamma_I^2-i\epsilon} + 2\pi\delta(-\omega^2+\Gamma_I^2)n_I(k) & \theta(-\omega)2\pi\delta(-\omega^2+\Gamma_I^2) + 2\pi\delta(-\omega^2+\Gamma_I^2)n_I(k) \\ \theta(\omega)2\pi\delta(-\omega^2+\Gamma_I^2) + 2\pi\delta(-\omega^2+\Gamma_I^2)n_I(k) & \frac{i}{-\omega^2+\Gamma_I^2+i\epsilon} + 2\pi\delta(-\omega^2+\Gamma_I^2)n_I(k) \end{pmatrix}$$

$$n_I(k) = \theta(\omega)n_{I+}(\mathbf{k}) + \theta(-\omega)n_{I-}(-\mathbf{k})$$

$$\lim_{q\to 0} S_\mathrm{IF}[A_+,A_-,\Phi_+,\Phi_-,\Phi_+^*,\Phi_-^*] =$$
$$i\sum_I g_I^2 \int \mathrm{d}^4x \int \mathrm{d}^4y (\Phi_+^*(x) - \Phi_-^*(x))\mathbf{D}_I(x,y)\begin{pmatrix}\Phi_+(y)\\ -\Phi_-(y)\end{pmatrix} - i\sum_I \mathrm{Trlog}\ \mathbf{D}_I + i\sum_I \mathrm{Trlog}\ \mathbf{D}_{IA}$$

$$S[A,\Psi,\bar\Psi] = \int \mathrm{d}^4x \left[-\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + i\bar\Psi \not{D}\Psi - m\bar\Psi\Psi\right]$$

$$e^{iS_\mathrm{IF}[\Psi_+,\bar\Psi_+,\Psi_-,\bar\Psi_-]} = \int \mathcal{D}[A_+,A_-,c_+,c_-,\bar c_+,\bar c_-]$$
$$\frac{\mu}{\tilde\mu} e^{i\int \mathrm{d}^4x\left(\frac{1}{2}A_\mu^+\Box A_+^\mu + \bar\Psi_+ q\not{A}_+\Psi_+ - \frac{1}{2}A_\mu^-\Box A_-^\mu - \bar\Psi_- q\not{A}_-\Psi_- + \bar c_+\Box c_+ - \bar c_-\Box c_-\right) + iS_{\mathrm{i}\epsilon}}$$

$$S_\mathrm{IF}[\Psi_+,\bar\Psi_+,\Psi_-,\bar\Psi_-] = \frac{i}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y \left(J_+^\mu(x) - J_-^\mu(x)\right)\mathbf{D}_{\mu\nu}(x,y)\begin{pmatrix}J_+^\nu(y)\\ -J_-^\nu(y)\end{pmatrix}$$

$$J_\pm^\mu(x) = \bar\Psi_\pm(x) q\gamma^\mu \Psi_\pm(x)$$

$$\mathbf{D}_{\mu\nu}(x,y) = \begin{pmatrix} D_{\mu\nu}^{++}(x,y) & D_{\mu\nu}^{+-}(x,y) \\ D_{\mu\nu}^{-+}(x,y) & D_{\mu\nu}^{--}(x,y) \end{pmatrix}$$

$$\mathbf{D}_{\mu\nu}(k) = \begin{pmatrix} -\frac{i\eta_{\mu\nu}}{k^2-i\epsilon} + \pi\delta(k^2)\left(-\eta_{\mu\nu} + K_{\mu\nu}^A(k)\right) & \pi\delta(k^2)\left(\eta_{\mu\nu}[2\theta(-k^0)-1] + K_{\mu\nu}^A(k)\right) \\ \pi\delta(k^2)\left(\eta_{\mu\nu}[2\theta(k^0)-1] + K_{\mu\nu}^A(k)\right) & \frac{i\eta_{\mu\nu}}{k^2+i\epsilon} + \pi\delta(k^2)\left(-\eta_{\mu\nu} + K_{\mu\nu}^A(k)\right) \end{pmatrix}$$

$$e^{iS_{\mathrm{IF}}[A_+,A_-]} = \int \mathcal{D}[\Psi_+,\bar{\Psi}_+,\Psi_-,\bar{\Psi}_-]$$

$$\frac{\mu}{\tilde{\mu}} e^{i\int \mathrm{d}^4x(i\bar{\Psi}_+\partial\Psi_+ - m\bar{\Psi}_+\Psi_+ - i\bar{\Psi}_-\partial\Psi_- + m\bar{\Psi}_-\Psi_- + \bar{\Psi}_+q\mathbb{A}_+\Psi_+ - \bar{\Psi}_-q\mathbb{A}_-\Psi_-) + iS_{\mathrm{i}\epsilon}}$$

$$\mathbf{S}_{\alpha\beta}(x,y) = \begin{pmatrix} S^{++}_{\alpha\beta}(x,y) & S^{+-}_{\alpha\beta}(x,y) \\ S^{-+}_{\alpha\beta}(x,y) & S^{--}_{\alpha\beta}(x,y) \end{pmatrix}$$

$$\begin{aligned} S^{++}_{\alpha\beta}(x,y) &= \mathrm{Tr}\left[\rho\mathcal{T}\hat{\Psi}_\alpha(x)\hat{\bar{\Psi}}_\beta(y)\right] \\ S^{+-}_{\alpha\beta}(x,y) &= -\mathrm{Tr}\left[\rho\hat{\bar{\Psi}}_\beta(y)\hat{\Psi}_\alpha(x)\right] \\ S^{-+}_{\alpha\beta}(x,y) &= \mathrm{Tr}\left[\rho\hat{\Psi}_\alpha(x)\hat{\bar{\Psi}}_\beta(y)\right] \\ S^{--}_{\alpha\beta}(x,y) &= \mathrm{Tr}\left[\rho\overline{\mathcal{T}}\hat{\Psi}_\alpha(x)\hat{\bar{\Psi}}_\beta(y)\right] \end{aligned}$$

$$[i\not{\partial}[A_{\mathrm{r}}] - m]\hat{\Psi} = 0$$

$$\begin{aligned} S^{++}_{\alpha\beta}(x,y) &= Z^{-1}\mathrm{Tr}\left[\rho\overline{\mathcal{T}}^* e^{i\hat{S}_{\mathrm{int}}[A_{\mathrm{a}}]}\mathcal{T}^* e^{i\hat{S}_{\mathrm{int}}[A_{\mathrm{a}}]}\hat{\Psi}_\alpha(x)\hat{\bar{\Psi}}_\beta(y)\right] \\ S^{+-}_{\alpha\beta}(x,y) &= -Z^{-1}\mathrm{Tr}\left[\rho\overline{\mathcal{T}}^*\left(e^{i\hat{S}_{\mathrm{int}}[A_{\mathrm{a}}]}\hat{\bar{\Psi}}_\beta(y)\right)\mathcal{T}^*\left(e^{i\hat{S}_{\mathrm{int}}[A_{\mathrm{a}}]}\hat{\Psi}_\alpha(x)\right)\right] \\ S^{-+}_{\alpha\beta}(x,y) &= Z^{-1}\mathrm{Tr}\left[\rho\overline{\mathcal{T}}^*\left(e^{i\hat{S}_{\mathrm{int}}[A_{\mathrm{a}}]}\hat{\Psi}_\alpha(x)\right)\mathcal{T}^*\left(e^{i\hat{S}_{\mathrm{int}}[A_{\mathrm{a}}]}\hat{\bar{\Psi}}_\beta(y)\right)\right] \\ S^{--}_{\alpha\beta}(x,y) &= Z^{-1}\mathrm{Tr}\left[\rho\overline{\mathcal{T}}^*\left(e^{i\hat{S}_{\mathrm{int}}[A_{\mathrm{a}}]}\hat{\Psi}_\alpha(x)\right)\mathcal{T}^*\left(e^{i\hat{S}_{\mathrm{int}}[A_{\mathrm{a}}]}\hat{\bar{\Psi}}_\beta(y)\right)\right] \end{aligned}$$

$$\hat{S}_{\mathrm{int}} = \int \mathrm{d}^4x q\hat{\bar{\Psi}}\not{\mathcal{A}}_{\mathrm{a}}\hat{\Psi}$$

$$Z = \mathrm{Tr}\left[\rho\overline{\mathcal{T}}^* e^{i\hat{S}_{\mathrm{int}}[A_{\mathrm{a}}]}\mathcal{T}^* e^{i\hat{S}_{\mathrm{int}}[A_{\mathrm{a}}]}\right]$$

$$\begin{aligned} e^{iS_{\mathrm{IF}}[A_+,A_-]} &= \int \mathcal{D}[\Psi_+,\bar{\Psi}_+,\Psi_-,\bar{\Psi}_-]\frac{\mu}{\tilde{\mu}}\exp\left(-(\bar{\Psi}_+\bar{\Psi}_-)\mathbf{S}^{-1}[A_+,A_-]\begin{pmatrix}\Psi_+\\ \Psi_-\end{pmatrix}\right) \\ &= \frac{\det[\mathbf{S}_A[A_+,A_-]]}{\det[\mathbf{S}[A_+,A_-]]} \end{aligned}$$

$$S_{\mathrm{IF}}[A_+,A_-] = i\mathrm{Trlog}\ \mathbf{S}[A_+,A_-] - i\mathrm{Trlog}\ \mathbf{S}_A[A_+,A_-]$$

$$\mathbf{S}^0(k) = (-\not{k} + m)\begin{pmatrix} \dfrac{-i}{k^2+m^2-i\epsilon} + \mathfrak{n}(k)2\pi\delta(k^2+m^2) & [\theta(-k^0)+\mathfrak{n}(k)]2\pi\delta(k^2+m^2) \\ [\theta(k^0)+\mathfrak{n}(k)]2\pi\delta(k^2+m^2) & \dfrac{i}{k^2+m^2+i\epsilon} + \mathfrak{n}(k)2\pi\delta(k^2+m^2) \end{pmatrix}$$

$$\mathfrak{n}(k) = \theta(k^0)n_+(|\mathbf{k}|) - \theta(-k^0)n_-(|\mathbf{k}|).$$

$$\mathbf{S}^{-1} = \mathbf{S}_0^{-1} - iq\mathbf{A},$$

$$\mathbb{A}(x,y) = \begin{pmatrix} A_+(x) & 0 \\ 0 & -A_-(x) \end{pmatrix}\delta^4(x-y)$$

$$\mathbf{S}[A_+,A_-] = (1 - \mathbf{S}^0 iq\mathbf{A})^{-1}\mathbf{S}^0$$

$$\Psi_\pm(x) = U_\pm(x)\tilde{\Psi}_\pm(x)$$

$$U_{\pm}(x) = e^{\pm\frac{i}{2}q\int_0^1 \mathrm{d}s(x_\mathrm{f}-x)^\mu A_\mu^\mathrm{a}(x+s(x_\mathrm{f}-x))}$$

$$\tilde{A}_\mu^\mathrm{a}(x) = A_\mu^\mathrm{a}(x) - \partial_\mu \int_0^1 \mathrm{d}s(x_\mathrm{f}-x)^\nu A_\nu^\mathrm{a}\big(x+s(x_\mathrm{f}-x)\big)$$

$$\tilde{\Psi}_{\pm}(x) \to e^{iq\lambda_\mathrm{r}(x)}\tilde{\Psi}_{\pm}(x)$$

$$\mathbf{S}[A_+, A_-](x,y) = \mathbf{U}(x)\tilde{\mathbf{S}}\big[A_\mathrm{r}, \tilde{A}_\mathrm{a}\big](x,y)\mathbf{U}(y)^\dagger,$$

where $\tilde{\mathbf{S}}\big[A_\mathrm{r}, \tilde{A}_\mathrm{a}\big](x,y)$ gauge transformations

$$\tilde{\mathbf{S}}\big[A_\mathrm{r}, \tilde{A}_\mathrm{a}\big](x,y) \to e^{iq(\lambda_\mathrm{r}(x)-\lambda_\mathrm{r}(y))}\tilde{\mathbf{S}}\big[A_\mathrm{r}, \tilde{A}_\mathrm{a}\big](x,y),$$

$$\mathbf{U}(x) = \begin{pmatrix} U_+(x) & 0 \\ 0 & U_-(x) \end{pmatrix}.$$

$$\tilde{\mathbf{S}}\big[A_\mathrm{r}, \tilde{A}_\mathrm{a}\big](x,y) = \mathcal{W}_{C_3}(x,y)\tilde{\mathbf{S}}\big[A_\mathrm{r}, \tilde{A}_\mathrm{a}\big](x,y),$$

$$\check{\mathbf{S}}\big[A_\mathrm{r}, \tilde{A}_\mathrm{a}\big](t_\mathrm{i}, \mathbf{x}, t_\mathrm{i}, \mathbf{y}) = \mathbf{S}^0(\mathbf{x}, \mathbf{y})$$

$$\mathbf{S}[A_+, A_-](x,y) = \mathbf{U}(x)\left[\left(1 - \tilde{\mathbf{S}}[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\right)^{-1}\tilde{\mathbf{S}}[A_\mathrm{r}]\right](x,y)\mathbf{U}(y)^\dagger$$

$$\tilde{A}_\mathrm{a}(x,y) = \frac{1}{2}\begin{pmatrix} \tilde{A}_\mathrm{a}(x) & 0 \\ 0 & \tilde{A}_\mathrm{a}(x) \end{pmatrix}\delta^4(x-y)$$

$$S_\mathrm{IF}[A_+, A_-] = i\mathrm{Trlog}\ \tilde{\mathbf{S}}[A_\mathrm{r}] - i\mathrm{Trlog}\left(1 - \tilde{\mathbf{S}}[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\right) - i\mathrm{Trlog}\ \tilde{\mathbf{S}}_A[A_\mathrm{r}] + i\mathrm{Trlog}\left(1 - \tilde{\mathbf{S}}_A[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\right)$$

$$S_\mathrm{IF}[A_+, A_-] = i\mathrm{Trlog}\ \tilde{\mathbf{S}}[A_\mathrm{r}] - i\mathrm{Trlog}\left(1 - \tilde{\mathbf{S}}[A_\mathrm{r}]iqA_\mathrm{a}\right) - i\mathrm{Trlog}\ \tilde{\mathbf{S}}_A[A_\mathrm{r}] + i\mathrm{Trlog}\left(1 - \tilde{\mathbf{S}}_A[A_\mathrm{r}]iqA_\mathrm{a}\right),$$

$$A_\mathrm{a}(x,y) = \frac{1}{2}\begin{pmatrix} A_\mathrm{a}(x) & 0 \\ 0 & A_\mathrm{a}(x) \end{pmatrix}\delta^4(x-y).$$

$$\begin{aligned} S_\mathrm{IF}[A_\mathrm{r}, A_\mathrm{a}] =& i\mathrm{Trlog}\ \tilde{\mathbf{S}}[A_\mathrm{r}] + i\mathrm{Tr}\big(\tilde{\mathbf{S}}[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\big) + \frac{i}{2}\mathrm{Tr}\big(\tilde{\mathbf{S}}[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\big)^2 + \cdots \\ & - i\mathrm{Trlog}\ \tilde{\mathbf{S}}_A[A_\mathrm{r}] - i\mathrm{Tr}\big(\tilde{\mathbf{S}}_A[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\big) - \frac{i}{2}\mathrm{Tr}\big(\tilde{\mathbf{S}}_A[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\big)^2 + \cdots \end{aligned}$$

$$\mathrm{Trlog}\ \tilde{\mathbf{S}}[A_\mathrm{r}] = \mathrm{Trlog}\ \tilde{\mathbf{S}}_A[A_\mathrm{r}]$$

$$\det\tilde{\mathbf{S}}[A_\mathrm{r}] = \det\tilde{\mathbf{S}}_A[A_\mathrm{r}]$$

$$S_\mathrm{IF}[A_\mathrm{r}, A_\mathrm{a}] = i\mathrm{Tr}\big(\tilde{\mathbf{S}}[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\big) + \frac{i}{2}\mathrm{Tr}\big(\tilde{\mathbf{S}}[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\big)^2 - i\mathrm{Tr}\big(\tilde{\mathbf{S}}_A[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\big) - \frac{i}{2}\mathrm{Tr}\big(\tilde{\mathbf{S}}_A[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\big)^2 + \cdots$$

$$S_\mathrm{IF}[A_\mathrm{r}, A_\mathrm{a}] = i\mathrm{Tr}\big(\tilde{\mathbf{S}}[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\big) + \frac{i}{2}\mathrm{Tr}\big(\tilde{\mathbf{S}}[A_\mathrm{r}]iq\tilde{A}_\mathrm{a}\big)^2 + \cdots$$

$$\partial^{\nu}F^{\mathrm{r}}_{\nu\mu}(x) = -\frac{\delta}{\delta A^{\mu}_{\mathrm{a}}(x)}\Big(i\mathrm{Tr}\big(\tilde{\mathbf{S}}[A_{\mathrm{r}}]iq\tilde{A}_{\mathrm{a}}\big)\Big)$$

$$= -J^{p}_{\mu}(x) + \int_0^1 \mathrm{d}s\,\frac{1}{(1-s)}(x_{\mathrm{f}}-x)^{\mu}\big(\partial^{\nu}J^{p}_{\nu}\big)(x_{\mathrm{ret}}(s))$$

$$J^{p}_{\mu}(x) = -\frac{1}{2}q\mathrm{tr}\big[\gamma_{\mu}\big(\tilde{S}^{++}[A_{\mathrm{r}}](x,x) + \tilde{S}^{--}[A_{\mathrm{r}}](x,x)\big)\big]$$

$$= -\frac{1}{2}q\mathrm{tr}\big[\gamma_{\mu}\big(\tilde{S}^{+-}[A_{\mathrm{r}}](x,x) + \tilde{S}^{-+}[A_{\mathrm{r}}](x,x)\big)\big]$$

$$x_{\mathrm{ret}}(s) = x - \frac{s}{1-s}(x_{\mathrm{f}}-x)$$

$$\partial^{\mu}J^{p}_{\mu}(x) = 0$$

$$\partial^{\mu}F^{\mathrm{r}}_{\mu\nu}(x) = -J^{p}_{\nu}(x)$$

$$S_{\mathrm{IF}}[A_{\mathrm{r}},A_{\mathrm{a}}] = \int \mathrm{d}^4x A^{\mu}_{\mathrm{a}}(x)J^{p}_{\mu}[A_{\mathrm{r}}](x) + \mathcal{O}(A^2_{\mathrm{a}})$$

$$W_{\alpha\beta}[A_{\mathrm{r}}](x,p) = \frac{1}{2}\int \mathrm{d}^4y e^{-ip.y}\big(\check{S}^{-+}_{\alpha\beta}[A_{\mathrm{r}}](x-y/2,x+y/2) + \check{S}^{+-}_{\alpha\beta}[A_{\mathrm{r}}](x-y/2,x+y/2)\big),$$

$$\int \frac{\mathrm{d}^4p}{(2\pi)^4}W_{\alpha\beta}[A_{\mathrm{r}}](x,p) = \frac{1}{2}\check{S}^{-+}_{\alpha\beta}[A_{\mathrm{r}}](x,x) + \frac{1}{2}\check{S}^{+-}_{\alpha\beta}[A_{\mathrm{r}}](x,x) = \frac{1}{2}\tilde{S}^{-+}_{\alpha\beta}[A_{\mathrm{r}}](x,x) + \frac{1}{2}\tilde{S}^{+-}_{\alpha\beta}[A_{\mathrm{r}}](x,x)$$

$$\partial^{\mu}F^{\mathrm{r}}_{\mu\nu}(x) = q\int \frac{\mathrm{d}^4p}{(2\pi)^4}\mathrm{tr}(\gamma_{\nu}W[A_{\mathrm{r}}](x,p))$$

$$\hat{\Psi}(x) = \int \mathrm{d}\tilde{k}\sum_{s}\big[u_{k,s}(x)b_{s}(\mathbf{k}) + v_{k,s}(x)d^{\dagger}_{s}(\mathbf{k})\big]$$

$$\int \mathrm{d}\tilde{k} = \int \frac{\mathrm{d}^4k}{(2\pi)^4}\theta(k^0)2\pi\delta(k^2+m^2)$$

$$[i\not{\partial}[A_{\mathrm{r}}] - m]u_{k,s}(x) = 0, [i\not{\partial}[A_{\mathrm{r}}] - m]v_{k,s}(x) = 0,$$

$$u_{k,s}(t_{\mathrm{i}},\mathbf{x}) = u_{s}(\mathbf{k})e^{i\mathbf{k}\cdot\mathbf{x}}, v_{k,s}(t_{\mathrm{i}},\mathbf{x}) = v_{s}(\mathbf{k})e^{-i\mathbf{k}\cdot\mathbf{x}}$$

$$\big\{b_{s}(\mathbf{k}), b^{\dagger}_{s'}(\mathbf{k}')\big\} = \big\{d_{s}(\mathbf{k}), d^{\dagger}_{s'}(\mathbf{k}')\big\} = 2\omega_{k}(2\pi)^3\delta^3(\mathbf{k}-\mathbf{k}')\delta_{ss'}$$

$$J^{p}_{\mu}(x) = q\int \mathrm{d}\tilde{k}\int \mathrm{d}\tilde{k}'\sum_{s,s'}\Big(\frac{1}{2}\bar{u}_{k',s'}(x)\gamma_{\mu}u_{k,s}(x)\mathrm{Tr}\Big[\rho\big[b^{\dagger}_{s'}(\mathbf{k}'),b_{s}(\mathbf{k})\big]\Big] + \bar{u}_{k',s'}(x)\gamma_{\mu}v_{k,s}(x)\mathrm{Tr}\big[\rho b^{\dagger}_{s'}(\mathbf{k}')d^{\dagger}_{s}(\mathbf{k})\big]$$
$$+\bar{v}_{k',s'}(x)\gamma_{\mu}u_{k,s}(x)\mathrm{Tr}[\rho d_{s'}(\mathbf{k}')b_{s}(\mathbf{k})] + \frac{1}{2}\bar{v}_{k',s'}(x)\gamma_{\mu}v_{k,s}(x)\mathrm{Tr}\Big[\rho\big[d_{s'}(\mathbf{k}'),d^{\dagger}_{s}(\mathbf{k})\big]\Big]\Big)$$

$$\mathrm{Tr}[\rho d_{s'}(\mathbf{k}')b_{s}(\mathbf{k})] = \mathrm{Tr}\big[\rho b^{\dagger}_{s'}(\mathbf{k}')d^{\dagger}_{s}(\mathbf{k})\big] = 0$$

$$J^{p,\,\mathrm{finite}}_{\mu}(x) = q\int \mathrm{d}\tilde{k}\int \mathrm{d}\tilde{k}'\sum_{s,s'}\big(\bar{u}_{k',s'}(x)\gamma_{\mu}u_{k,s}(x)\mathrm{Tr}\big[\rho b^{\dagger}_{s'}(\mathbf{k}')b_{s}(\mathbf{k})\big] - \bar{v}_{k',s'}(x)\gamma_{\mu}v_{k,s}(x)\mathrm{Tr}\big[\rho d^{\dagger}_{s'}(\mathbf{k}')d_{s}(\mathbf{k})\big]\big)$$

$$J_{\mu}^{p,\,\text{vacuum}}(x)=\frac{1}{2}q\int \mathrm{d}\tilde{k}\sum_{s}\left(\bar{v}_{k,s}(x)\gamma_{\mu}v_{k,s}(x)-\bar{u}_{k,s}(x)\gamma_{\mu}u_{k,s}(x)\right)$$

$$J_{\mu}^{p,\text{vacuum}}(x)=\frac{\delta S_{1-\text{ loop vac}}\left[A_{\text{r}}\right]}{\delta A_{\text{r}}^{\mu}(x)}$$

$$S_{1\text{-loop vac}}\left[A_{\text{r}}\right]=\frac{i}{2}\text{Trlog}\left[\check{S}_{\text{vac}}^{++}\left[A_{\text{r}}\right]\right]+\frac{i}{2}\text{Trlog}\left[\check{S}_{\text{vac}}^{--}\left[A_{\text{r}}\right]\right]$$
$$=-\frac{i}{2}\text{Trlog}\left[i\not{D}[A_{\text{r}}]-m+i\epsilon\right]-\frac{i}{2}\text{Trlog}\left[i\not{D}[A_{\text{r}}]-m-i\epsilon\right]$$

$$\partial^{\mu}J_{\mu}^{p,\,\text{vacuum}}(x)=0$$

$$[-i\not{D}[A_{\text{r}}]-m][i\not{D}[A_{\text{r}}]-m]u_{k,s}(x)=\left[-D[A_{\text{r}}]^{2}+m^{2}-\frac{1}{2}q\sigma_{\mu\nu}F^{\mu\nu}(x)\right]u_{k,s}(x)=0$$

$$\sigma_{\mu\nu}=\frac{i}{2}\left[\gamma_{\mu},\gamma_{\nu}\right]$$

$u_{k,s}(x)=e^{ik\cdot x}U_{k,s}(x)$ with $k^{2}+m^{2}=0$ so that

$$\left[-2ik.D[A_{\text{r}}]-D[A_{\text{r}}]^{2}-\frac{1}{2}q\sigma_{\mu\nu}F^{\mu\nu}(x)\right]U_{k,s}(x)=0$$

$$\left[-ik.D[A_{\text{r}}]-\frac{1}{4}q\sigma_{\mu\nu}F^{\mu\nu}(x)\right]U_{k,s}(x)\simeq 0$$

$$U_{k,s}(x)=e^{iq\int_{\tau_{\text{i}}}^{0}\mathrm{d}\tau A_{\mu}(z(\tau))\frac{\mathrm{d}z^{\mu}}{\mathrm{d}\tau}}\mathcal{P}e^{-\frac{i}{4}q\int_{\tau_{\text{i}}}^{0}\mathrm{d}\tau\sigma_{\alpha\beta}F^{\alpha\beta}(z(\tau))}u_{s}(\mathbf{k})$$

$$z^{\mu}(\tau)=x^{\mu}+\tau k^{\mu}$$

$$z^{\mu}(0)=x^{\mu}\text{ and }z^{0}(\tau_{\text{i}})=t_{\text{i}}=t+k^{0}\tau_{\text{i}}$$

$$U_{k,s}(x)\simeq e^{iq\int_{\tau_{\text{i}}}^{0}\mathrm{d}\tau A_{\mu}(z(\tau))\frac{\mathrm{d}z^{\mu}}{\mathrm{d}\tau}}u_{s}(\mathbf{k})$$

$$u_{k,s}(x)\simeq e^{-imt}V(x)u_{s}(\mathbf{k})$$

$$\hat{\Psi}(x)\simeq\int \mathrm{d}\tilde{k}\sum_{s}\left[e^{ik\cdot x}e^{iq\int_{\tau_{i}}^{0}\mathrm{d}\tau A_{\mu}(z(\tau))\frac{\mathrm{d}z^{\mu}}{\mathrm{d}\tau}}\mathcal{P}e^{-\frac{i}{4}q\int_{\tau_{\text{i}}}^{0}\mathrm{d}\tau\sigma_{\alpha\beta}F^{\alpha\beta}(z(\tau))}u_{s}(\mathbf{k})b_{s}(\mathbf{k})\right.$$
$$\left.+e^{-ik\cdot x}e^{iq\int_{\tau_{i}}^{0}\mathrm{d}\tau A_{\mu}(z(\tau))\frac{\mathrm{d}z^{\mu}}{\mathrm{d}\tau}}\mathcal{P}e^{\frac{i}{4}q\int_{\tau_{i}}^{0}\mathrm{d}\tau\sigma_{\alpha\beta}F^{\alpha\beta}(z(\tau))}v_{s}(\mathbf{k})d_{s}^{\dagger}(\mathbf{k})\right]$$

$$\tilde{S}^{-+}[A_{\text{r}}](x,y)=\text{Tr}\{\rho\Psi(x)\bar{\Psi}(y)\}$$
$$=\int \mathrm{d}\tilde{k}\int \mathrm{d}\tilde{k}'\sum_{s,s'}\left[u_{k,s}(x)\bar{u}_{k',s'}(y)\text{Tr}\left[\rho b_{s}(\mathbf{k})b_{s'}^{\dagger}(\mathbf{k}')\right]+v_{k,s}(x)\bar{v}_{k',s'}(y)\text{Tr}\left[\rho d_{s}^{\dagger}(\mathbf{k})d_{s'}(\mathbf{k}')\right]\right]$$

$$\tilde{S}^{-+}(t_{\mathrm{i}},\mathbf{x},t_{\mathrm{i}},\mathbf{y}) = \int \mathrm{d}\tilde{k} \int \mathrm{d}\tilde{k}' \sum_{s,s'} \Big[e^{-i(\omega_k-\omega_{k'})t_{\mathrm{i}}} e^{i\mathbf{k}\cdot\mathbf{x}-i\mathbf{k}'\cdot\mathbf{y}} u_s(\mathbf{k})\bar{u}_{s'}(\mathbf{k}')\mathrm{Tr}\big[\rho b_s(\mathbf{k})b_{s'}^{\dagger}(\mathbf{k}')\big]$$
$$+ e^{i(\omega_k-\omega_{k'})t_{\mathrm{i}}} e^{-i\mathbf{k}\cdot\mathbf{x}+i\mathbf{k}'\cdot\mathbf{y}} v_s(\mathbf{k})\bar{v}_{s'}(\mathbf{k}')\mathrm{Tr}\big[\rho d_s^{\dagger}(\mathbf{k})d_{s'}(\mathbf{k}')\big]\Big]$$

$$\tilde{S}^{-+}(t_{\mathrm{i}},\mathbf{x},t_{\mathrm{i}},\mathbf{y}) = \mathcal{W}_{C_3}(t_{\mathrm{i}},\mathbf{x},t_{\mathrm{i}},\mathbf{y})\check{S}^{-+}(\mathbf{x},\mathbf{y})$$

$$\check{S}^{-+}(\mathbf{x},\mathbf{y}) = \int \mathrm{d}\tilde{k} \sum_{s} \Big[e^{i\mathbf{k}\cdot(\mathbf{x}-\mathbf{y})} u_s(\mathbf{k})\bar{u}_s(\mathbf{k})(1-n_{+s}(\mathbf{k})) + e^{-i\mathbf{k}\cdot(\mathbf{x}-\mathbf{y})} v_s(\mathbf{k})\bar{v}_s(\mathbf{k})n_{-s}(\mathbf{k})\Big]$$

$$\check{S}^{-+}(\mathbf{x},\mathbf{y})\mathrm{Tr}\big[\rho b_s(\mathbf{k})b_{s'}^{\dagger}(\mathbf{k}')\big]$$

$$\mathrm{Tr}\big[\rho b_s(\mathbf{k})b_{s'}^{\dagger}(\mathbf{k}')\big] \propto \delta^3(\mathbf{k}-\mathbf{k}')$$

$$\mathrm{Tr}\big[\rho b_{s'}^{\dagger}(\mathbf{k}')b_s(\mathbf{k})\big] = \mathcal{M}_{ss'}^{b}(\mathbf{k}',\mathbf{k})\delta(\omega_k-\omega_{k'})$$
$$\mathrm{Tr}\big[\rho d_{s'}^{\dagger}(\mathbf{k})d_s(\mathbf{k}')\big] = \mathcal{M}_{ss'}^{d}(\mathbf{k}',\mathbf{k})\delta(\omega_k-\omega_{k'})$$

$$\tilde{S}^{-+}(x,y) = \tilde{S}_{\mathrm{vac}}^{-+}(x,y) +$$
$$\int \mathrm{d}\tilde{k} \int \mathrm{d}\tilde{k}' \sum_{s,s'} \big[-u_{k,s}(x)\bar{u}_{k',s'}(y)\mathcal{M}_{ss'}^{b}(\mathbf{k}',\mathbf{k})\delta(\omega_k-\omega_{k'}) + v_{k,s}(x)\bar{v}_{k',s'}(y)\mathcal{M}_{ss'}^{d}(\mathbf{k}',\mathbf{k})\delta(\omega_k-\omega_{k'})\big]$$

$$J_{\mu}^{p,\,\mathrm{finite}}(x) = q\int \mathrm{d}\tilde{k} \int \mathrm{d}\tilde{k}' \sum_{s,s'} \big[\bar{u}_{k',s'}(x)\gamma_{\mu}u_{k,s}(x)\mathcal{M}_{ss'}^{b}(\mathbf{k}',\mathbf{k}) - \bar{v}_{k',s'}(x)\gamma_{\mu}v_{k,s}(x)\mathcal{M}_{ss'}^{d}(\mathbf{k}',\mathbf{k})\big]\delta(\omega_k-\omega_{k'})$$

$$\mathcal{U}_{\pm}(k,x) = e^{iq\int_{-\infty}^{0}\mathrm{d}\tau k^{\mu}A_{\mu}(x+k\tau)}\mathcal{P}e^{\mp\frac{i}{4}q\int_{-\infty}^{0}\mathrm{d}\tau\sigma_{\alpha\beta}F^{\alpha\beta}(x+k\tau)}$$

$$J_{\mu}^{p,\,\mathrm{finite}}(x) = q\int \mathrm{d}\tilde{k} \int \mathrm{d}\tilde{k}' \sum_{s,s'} \big[\bar{u}_{s'}(\mathbf{k}')\mathcal{U}_{+}^{\dagger}(k',x)\gamma_{\mu}\mathcal{U}_{+}(k,x)u_s(\mathbf{k})\mathcal{M}_{ss'}^{b}(\mathbf{k}',\mathbf{k})$$
$$-\bar{v}_{s'}(\mathbf{k}')\mathcal{U}_{-}^{\dagger}(k',x)\gamma_{\mu}\mathcal{U}_{-}(k,x)v_s(\mathbf{k})\mathcal{M}_{ss'}^{d}(\mathbf{k}',\mathbf{k})\big]\delta(\omega_k-\omega_{k'})$$

$$\mathrm{Tr}\big[\rho b_{s'}^{\dagger}(\mathbf{k}')b_s(\mathbf{k})\big] = \delta_{ss'}2\omega_k(2\pi)^3\delta^3(\mathbf{k}-\mathbf{k}')n_{+s}(\mathbf{k}), \mathrm{Tr}\big[\rho d_{s'}^{\dagger}(\mathbf{k}')d_s(\mathbf{k})\big] = \delta_{ss'}2\omega_k(2\pi)^3\delta^3(\mathbf{k}-\mathbf{k}')n_{-s}(\mathbf{k}),$$

$$J_{\mu}^{p,\,\mathrm{finite}}(x) = q\int \mathrm{d}\tilde{k} \sum_{s} \big(\bar{u}_s(\mathbf{k})\mathcal{U}_{+}^{\dagger}(k,x)\gamma_{\mu}\mathcal{U}_{+}(k,x)u_s(\mathbf{k})n_{+s}(\mathbf{k}) - \bar{v}_s(\mathbf{k})\mathcal{U}_{-}^{\dagger}(k,x)\gamma_{\mu}\mathcal{U}_{-}(k,x)v_s(\mathbf{k})n_{-s}(\mathbf{k})\big).$$

$$S_{\mathrm{gravity}}[A_{\mathrm{r}},A_{\mathrm{a}}] = \frac{i}{2}\mathrm{Tr}\big(\tilde{\mathbf{S}}[A_{\mathrm{r}}]iq\tilde{A}_{\mathrm{a}}\big)^2$$

$$S_{\mathrm{gravity}}[A_{\mathrm{r}},A_{\mathrm{a}}] = \frac{i}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y\tilde{A}_{\mathrm{a}}^{\mu}(x)N_{\mu\nu}[A_{\mathrm{r}}](x,y)\tilde{A}_{\mathrm{a}}^{\nu}(y)$$

$$N_{\mu\nu}[A_{\mathrm{r}}](x,y) = -\frac{1}{4}q^2\big(\mathrm{tr}\big[\tilde{S}_{++}[A_{\mathrm{r}}](y,x)\gamma_{\mu}\tilde{S}_{++}[A_{\mathrm{r}}](x,y)\gamma_{\nu}\big] + \mathrm{tr}\big[\tilde{S}_{--}[A_{\mathrm{r}}](y,x)\gamma_{\mu}\tilde{S}_{--}[A_{\mathrm{r}}](x,y)\gamma_{\nu}\big]$$
$$+\mathrm{tr}\big[\tilde{S}_{+-}[A_{\mathrm{r}}](y,x)\gamma_{\mu}\tilde{S}_{-+}[A_{\mathrm{r}}](x,y)\gamma_{\nu}\big] + \mathrm{tr}\big[\tilde{S}_{-+}[A_{\mathrm{r}}](y,x)\gamma_{\mu}\tilde{S}_{+-}[A_{\mathrm{r}}](x,y)\gamma_{\nu}\big]\big),$$

$$N_{\mu\nu}[A_{\mathrm{r}}](x,y) = N_{\nu\mu}[A_{\mathrm{r}}](y,x),$$

$$N_{\mu\nu}[A_{\mathrm{r}}](x,y) = -\frac{q^2}{2}\left(\mathrm{tr}\left[\tilde{S}_{+-}[A_{\mathrm{r}}](y,x)\gamma_\mu \tilde{S}_{-+}[A_{\mathrm{r}}](x,y)\gamma_\nu\right] + \mathrm{tr}\left[\tilde{S}_{-+}[A_{\mathrm{r}}](y,x)\gamma_\mu \tilde{S}_{+-}[A_{\mathrm{r}}](x,y)\gamma_\nu\right]\right)$$

$$\partial_x^\mu N_{\mu\nu}[A_{\mathrm{r}}](x,y) = D_x^\mu[A_{\mathrm{r}}]N_{\mu\nu}[A_{\mathrm{r}}](x,y) = 0.$$

$$[i\not{\partial}_x[A_{\mathrm{r}}] - m]\tilde{S}_{+-}(x,y) = [i\not{\partial}_x[A_{\mathrm{r}}] - m]\tilde{S}_{-+}(x,y) = 0$$
$$\tilde{S}_{+-}(x,y)\left[-i\not{D}_y[A_{\mathrm{r}}] - m\right] = \tilde{S}_{-+}(x,y)\left[-i\not{D}_y[A_{\mathrm{r}}] - m\right] = 0$$

$$S_{\text{gravity}}[A_{\mathrm{r}}, A_{\mathrm{a}}] = \frac{i}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y A_{\mathrm{a}}^\mu(x) N_{\mu\nu}[A_{\mathrm{r}}](x,y) A_{\mathrm{a}}^\nu(y)$$

$$S[A,\Psi,\bar{\Psi},\chi_I] = S_{\mathcal{S}}[A,\Psi,\bar{\Psi}] + S_{\mathcal{E}}[\chi_I] + S_{\mathrm{int}}[A,\Psi,\bar{\Psi},\chi_I]$$

$$S_{\mathcal{S}}[A,\Psi,\bar{\Psi}] = \int \mathrm{d}^4x\left[-\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + i\bar{\Psi}\not{\partial}\Psi - m\bar{\Psi}\Psi\right]$$
$$S_{\mathcal{E}}[\chi_I] = \sum_I \int \mathrm{d}^4x[i\bar{\chi}_I\not{\partial}\chi_I - M_I\bar{\chi}_I\chi_I]$$
$$S_{\mathrm{int}}[A,\Psi,\bar{\Psi},\chi_I] = \sum_I \int \mathrm{d}^4x g_I(\bar{\Psi}\chi_I + \bar{\chi}_I\Psi).$$

$$S_{\mathrm{IF}}[A_+,\Psi_+,\bar{\Psi}_+,A_-,\Psi_-,\bar{\Psi}_-] = \int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I ig_I^2(\bar{\Psi}_+(x) - \bar{\Psi}_-(x))\mathbf{S}_I[A_+,A_-](x,y)\begin{pmatrix}\Psi_+(y)\\-\Psi_-(y)\end{pmatrix}$$
$$+i\sum_I \mathrm{Tr}\left[\log \mathbf{S}_I[A_+,A_-]\right] - i\sum_I \mathrm{Tr}\left[\log \mathbf{S}_{IA}[A_+,A_-]\right]$$

$$S_{\mathrm{IF}} = \int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I ig_I^2\left(\bar{\tilde{\Psi}}_+(x) - \bar{\tilde{\Psi}}_-(x)\right)\left(1 - \tilde{\mathbf{S}}_I[A_{\mathrm{r}}]iq\tilde{A}_{\mathrm{a}}\right)^{-1}\tilde{\mathbf{S}}_I[A_{\mathrm{r}}](x,y)\begin{pmatrix}\tilde{\Psi}_+(y)\\-\tilde{\Psi}_-(y)\end{pmatrix}$$
$$+i\sum_I \mathrm{Tr}\left[\log \tilde{\mathbf{S}}_I[A_{\mathrm{r}}]\right] - i\sum_I \mathrm{Tr}\left[\log\left(1 - \tilde{\mathbf{S}}_I[A_{\mathrm{r}}]iq\tilde{A}_{\mathrm{a}}\right)\right]$$
$$-i\sum_I \mathrm{Tr}\left[\log \tilde{\mathbf{S}}_{IA}[A_{\mathrm{r}}]\right] + i\sum_I \mathrm{Tr}\left[\log\left(1 - \tilde{\mathbf{S}}_{IA}[A_{\mathrm{r}}]iq\tilde{A}_{\mathrm{a}}\right)\right]$$

$$
\begin{aligned}
S_{\mathrm{IF}} =& \frac{1}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I i g_I^2 \left(\overline{\tilde{\Psi}}_{\mathrm{r}}(x) - \overline{\tilde{\Psi}}_{\mathrm{r}}(x)\right) \tilde{\mathbf{S}}_I[A_{\mathrm{r}}](x,y) \begin{pmatrix} \tilde{\Psi}_{\mathrm{a}}(y) \\ \tilde{\Psi}_{\mathrm{a}}(y) \end{pmatrix} \\
&+ \frac{1}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I i g_I^2 \left(\overline{\tilde{\Psi}}_{\mathrm{a}}(x)\overline{\tilde{\Psi}}_{\mathrm{a}}(x)\right) \tilde{\mathbf{S}}_I[A_{\mathrm{r}}](x,y) \begin{pmatrix} \tilde{\Psi}_{\mathrm{r}}(y) \\ -\tilde{\Psi}_{\mathrm{r}}(y) \end{pmatrix} \\
&+ \int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I i g_I^2 \left(\overline{\tilde{\Psi}}_{\mathrm{r}}(x) - \overline{\tilde{\Psi}}_{\mathrm{r}}(x)\right) \tilde{\mathbf{S}}_I[A_{\mathrm{r}}] i q \tilde{A}_{\mathrm{a}} \tilde{\mathbf{S}}_I[A_{\mathrm{r}}](x,y) \begin{pmatrix} \tilde{\Psi}_{\mathrm{r}}(y) \\ -\tilde{\Psi}_{\mathrm{r}}(y) \end{pmatrix} \\
&+ \frac{1}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I i g_I^2 \left(\overline{\tilde{\Psi}}_{\mathrm{a}}(x)\overline{\tilde{\Psi}}_{\mathrm{a}}(x)\right) \tilde{\mathbf{S}}_I[A_{\mathrm{r}}] i q \tilde{A}_{\mathrm{a}} \tilde{\mathbf{S}}_I[A_{\mathrm{r}}](x,y) \begin{pmatrix} \tilde{\Psi}_{\mathrm{r}}(y) \\ -\tilde{\Psi}_{\mathrm{r}}(y) \end{pmatrix} \\
&+ \frac{1}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I i g_I^2 \left(\overline{\tilde{\Psi}}_{\mathrm{r}}(x) - \overline{\tilde{\Psi}}_{\mathrm{r}}(x)\right) \tilde{\mathbf{S}}_I[A_{\mathrm{r}}] i q \tilde{A}_{\mathrm{a}} \tilde{\mathbf{S}}_I[A_{\mathrm{r}}](x,y) \begin{pmatrix} \tilde{\Psi}_{\mathrm{a}}(y) \\ \tilde{\Psi}_{\mathrm{a}}(y) \end{pmatrix} \\
&+ \int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I i g_I^2 \left(\overline{\tilde{\Psi}}_{\mathrm{r}}(x) - \overline{\tilde{\Psi}}_{\mathrm{r}}(x)\right) \left(\tilde{\mathbf{S}}_I[A_{\mathrm{r}}] i q \tilde{A}_{\mathrm{a}}\right)^2 \tilde{\mathbf{S}}_I[A_{\mathrm{r}}](x,y) \begin{pmatrix} \tilde{\Psi}_{\mathrm{r}}(y) \\ -\tilde{\Psi}_{\mathrm{r}}(y) \end{pmatrix} \\
&+ i \sum_I \mathrm{Tr}\left[\tilde{\mathbf{S}}_I[A_{\mathrm{r}}] i q \tilde{A}_{\mathrm{a}}\right] + \frac{i}{2}\sum_I \mathrm{Tr}\left[\left(\tilde{\mathbf{S}}_I[A_{\mathrm{r}}] i q \tilde{A}_{\mathrm{a}}\right)^2\right] + \cdots \\
&- i \sum_I \mathrm{Tr}\left[\tilde{\mathbf{S}}_{IA}[A_{\mathrm{r}}] i q \tilde{A}_{\mathrm{a}}\right] - \frac{i}{2}\sum_I \mathrm{Tr}\left[\left(\tilde{\mathbf{S}}_{IA}[A_{\mathrm{r}}] i q \tilde{A}_{\mathrm{a}}\right)^2\right] + \cdots
\end{aligned}
$$

$$
\lim_{q\to 0} S_{\mathrm{IF}}[A_+, \Psi_+, \bar{\Psi}_+, A_-, \Psi_-, \bar{\Psi}_-] = \int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I i g_I^2 (\bar{\Psi}_+(x) - \bar{\Psi}_-(x)) \tilde{\mathbf{S}}_I[0](x,y) \begin{pmatrix} \Psi_+(y) \\ -\Psi_-(y) \end{pmatrix}
$$

$$
S_{\mathcal{S}}[A, c, \bar{c}] = -\int \mathrm{d}^4x \left[\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{\left(\partial_\mu A^\mu\right)^2}{2\xi} + \bar{c} \Box c\right]
$$

$$
S_{\mathcal{E}}[\Phi, \Phi^*] = -\int \mathrm{d}^4x \left[\partial_\mu \Phi^* \partial^\mu \Phi + m^2 \Phi^* \Phi\right]
$$

$$
S_{\mathrm{int}}[A, \Phi] = -\int \mathrm{d}^4x \left[i q A^\mu \left(\Phi^* \partial_\mu \Phi - \Phi \partial_\mu \Phi^*\right) + q^2 A_\mu A^\mu \Phi^* \Phi\right]
$$


$$
\mathcal{F}^\beta(x,y) \equiv G_{++}(x,y)\big|_{A_+ = A_- = 0, \rho = \varrho_\beta} = \int \frac{\mathrm{d}^4k}{(2\pi)^4} \mathcal{F}^\beta(k) e^{ik\cdot(x-y)}
$$

$$
\mathcal{W}^\beta(x,y) \equiv G_{-+}(x,y)\big|_{A_+ = A_- = 0, \rho = \varrho_\beta} = \int \frac{\mathrm{d}^4k}{(2\pi)^4} \mathcal{W}^\beta(k) e^{ik\cdot(x-y)}
$$

$$
\mathcal{F}^\beta(k) = \frac{-i}{k^2 + m^2 - i\epsilon} + \frac{2\pi\delta(k^2 + m^2)}{e^{\beta|k_0|} - 1} \text{ and } \mathcal{W}^\beta(k) = 2\pi\delta(k^2 + m^2)\left[\theta(k^0) + \frac{1}{e^{\beta|k_0|} - 1}\right].
$$

$$
\begin{aligned}
S_{\mathrm{IF}}[A_+, A_-] \simeq \frac{q^2}{2}\int \frac{\mathrm{d}^4k}{(2\pi)^4} \Big[& -A_+^\mu(k) \Pi_{\mu\nu}^\beta(k) A_+^\nu(-k) - i A_+^\mu(k) \mathcal{N}_{\mu\nu}^\beta(-k) A_-^\nu(-k) \\
& - i A_-^\mu(k) \mathcal{N}_{\mu\nu}^\beta(k) A_+^\nu(-k) + A_-^\mu(k) \Pi_{\mu\nu}^{\beta *}(k) A_-^\nu(-k)\Big] + \mathcal{O}(q^4)
\end{aligned}
$$

$$
\begin{aligned}
\Pi_{\mu\nu}^\beta(k) &\equiv 2\eta_{\mu\nu}\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\left[\mathcal{F}^\beta(\ell)\right] - 2i\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\left[\mathcal{F}^\beta(\ell)\right]\left[\mathcal{F}^\beta(-\ell - k)\right]\left(2\ell_\mu + k_\mu\right)\ell_\nu \\
\mathcal{N}_{\mu\nu}^\beta(k) &\equiv 2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4} \mathcal{W}^\beta(\ell)\mathcal{W}^\beta(-\ell - k)\left(2\ell_\mu + k_\mu\right)\ell_\nu
\end{aligned}
$$



Feynman-Vernon's influence functional:

$$k^{\mu}\Pi^{\beta}_{\mu\nu}(k) = k^{\mu}\mathcal{N}^{\beta}_{\mu\nu}(k) = 0$$

$$\mathcal{P}^{\mathrm{L}}_{\mu\nu}(k) + \frac{k^2}{|\mathbf{k}|^2}\mathcal{P}^{\mathrm{T}}_{\mu\nu}(k) = k^2\eta_{\mu\nu} - k_{\mu}k_{\nu}.$$

$$\mathcal{P}^{\mathrm{L}}_{\mu\nu}(k) = k^2\frac{\left(u_{\mu} - \frac{u\cdot k}{k^2}k_{\mu}\right)\left(u_{\nu} - \frac{u\cdot k}{k^2}k_{\nu}\right)}{\left(u - \frac{u\cdot k}{k^2}k\right)^2},$$

$$\begin{bmatrix}\mathcal{P}^{\mathrm{L}}_{00} & \mathcal{P}^{\mathrm{L}}_{i0}\\ \mathcal{P}^{\mathrm{L}}_{0j} & \mathcal{P}^{\mathrm{L}}_{ij}\end{bmatrix} = \begin{bmatrix}-|\mathbf{k}|^2 & -k_ik_0\\ -k_0k_j & -k_ik_j\cdot\dfrac{k_0^2}{|\mathbf{k}|^2}\end{bmatrix} \text{ and } , \begin{bmatrix}\mathcal{P}^{\mathrm{T}}_{00} & \mathcal{P}^{\mathrm{T}}_{i0}\\ \mathcal{P}^{\mathrm{T}}_{0j} & \mathcal{P}^{\mathrm{T}}_{ij}\end{bmatrix} = \begin{bmatrix}0 & 0\\ 0 & |\mathbf{k}|^2\delta_{ij} - k_ik_j\end{bmatrix}.$$

$$\eta^{\nu\rho}\mathcal{P}^{\mathfrak{p}}_{\mu\nu}(k)\mathcal{P}^{\mathfrak{q}}_{\rho\sigma}(k) = \delta^{\mathfrak{p}\mathfrak{q}}\mathcal{P}^{\mathfrak{p}}_{\mu\sigma}(k), \text{ for } \mathfrak{p},\mathfrak{q} = \mathrm{L},\mathrm{T}.$$

$$S_{\mathrm{IF}}[A_+,A_-] \simeq \frac{q^2}{2}\sum_{\mathfrak{p}=\mathrm{L},\,\mathrm{T}}\int\frac{\mathrm{d}^4k}{(2\pi)^4}\mathcal{P}^{\mathfrak{p}}_{\mu\nu}(k)\Big[-A^{\mu}_{+}(k)\Pi^{\beta}_{\mathfrak{p}}(k)A^{\nu}_{+}(-k) - iA^{\mu}_{+}(k)\mathcal{N}^{\beta}_{\mathfrak{p}}(-k)A^{\nu}_{-}(-k)$$
$$-iA^{\mu}_{-}(k)\mathcal{N}^{\beta}_{\mathfrak{p}}(k)A^{\nu}_{+}(-k) + A^{\mu}_{-}(k)\Pi^{\beta *}_{\mathfrak{p}}(k)A^{\nu}_{-}(-k)\Big] + \mathcal{O}(q^4)$$

$$2\mathrm{Im}\left[\Pi^{\beta}_{\mathrm{L},\,\mathrm{T}}(k)\right] + \mathcal{N}^{\beta}_{\mathrm{L},\,\mathrm{T}}(k) + \mathcal{N}^{\beta}_{\mathrm{L},\,\mathrm{T}}(-k) = 0$$

$$\mathcal{N}^{\beta}_{\mathrm{L}}(k) = -\frac{\theta(-k^2-4m^2)}{8\pi|\mathbf{k}|^3}\int_{\Omega_-}^{\Omega_+}\mathrm{d}\Omega\left(\theta(-k^0)\coth\left(\frac{\beta\Omega}{2}\right)(2\Omega-|k_0|)^2 + \frac{4\Omega^2-2|k_0|\Omega}{[e^{\beta\Omega}-1][e^{\beta(|k_0|-\Omega)}-1]}\right)$$
$$-\frac{\theta(k^2)}{4\pi|\mathbf{k}|^3}\int_{\Omega_+}^{\infty}\mathrm{d}\Omega\left(\frac{\theta(k^0)}{e^{\beta\Omega}-1} + \frac{\theta(-k^0)}{e^{\beta(\Omega-|k_0|)}-1} + \frac{1}{[e^{\beta\Omega}-1][e^{\beta(\Omega-|k_0|)}-1]}\right)(2\Omega-|k_0|)^2$$

$$\Omega_{\pm} \equiv \frac{|k_0|}{2} \pm \frac{|\mathbf{k}|}{2}\sqrt{1+\frac{4m^2}{k^2}}.$$

$$S_{\mathrm{IF}}[A_{\mathrm{r}},A_{\mathrm{a}}] \simeq q^2\sum_{\mathfrak{p}=\mathrm{L},\,\mathrm{T}}\int\frac{\mathrm{d}^4k}{(2\pi)^4}\mathcal{P}^{\mathfrak{p}}_{\mu\nu}(k)\left[-A^{\mu}_{\mathrm{a}}(k)\mathcal{D}^{\beta}_{\mathfrak{p}}(k)A^{\nu}_{\mathrm{r}}(-k) + \frac{i}{2}A^{\mu}_{\mathrm{a}}(k)\mathcal{S}^{\beta}_{\mathfrak{p}}(k)A^{\nu}_{\mathrm{a}}(-k)\right]$$

$$\mathcal{D}^{\beta}_{\mathfrak{p}}(k) \equiv \mathrm{Re}\left[\Pi^{\beta}_{\mathfrak{p}}(k)\right] - \frac{i\left[\mathcal{N}^{\beta}_{\mathfrak{p}}(k) - \mathcal{N}^{\beta}_{\mathfrak{p}}(-k)\right]}{2}$$
$$\mathcal{S}^{\beta}_{\mathfrak{p}}(k) \equiv \frac{1}{2}\mathrm{Im}\left[\Pi^{\beta}_{\mathfrak{p}}(k)\right] - \frac{\mathcal{N}^{\beta}_{\mathfrak{p}}(k) + \mathcal{N}^{\beta}_{\mathfrak{p}}(-k)}{4}$$

$$\mathrm{Re}\left[\Pi_{\mathrm{L}}^{\beta}(k)\right] \simeq -\frac{1}{3|\mathbf{k}|^2\beta^2}\left[\frac{k_0}{2|\mathbf{k}|}\log\left|\frac{k_0/|\mathbf{k}|+1}{k_0/|\mathbf{k}|-1}\right|-1\right]+\mathcal{O}\left(\frac{1}{\beta}\right)$$

$$\mathrm{Im}\left[\Pi_{\mathrm{L}}^{\beta}(k)\right] \simeq \frac{\theta(k^2)\pi}{3|\mathbf{k}|^3\beta^3}-\frac{\theta(k^2)+\theta(-k^2-4m^2)}{2\pi|\mathbf{k}|^2\beta^2}\left[\sqrt{1+\frac{4m^2}{k^2}}-\frac{|k_0|}{2|\mathbf{k}|}\log\left|\frac{\Omega_+}{\Omega_-}\right|\right]+\mathcal{O}\left(\frac{1}{\beta}\right)$$

$$\mathcal{N}_{\mathrm{L}}^{\beta}(k) \simeq -\frac{\theta(k^2)\pi}{3|\mathbf{k}|^3\beta^3}-\frac{\theta(k^2)\pi k_0}{12|\mathbf{k}|^3\beta^2}+\frac{\theta(k^2)+\theta(-k^2-4m^2)}{2\pi|\mathbf{k}|^2\beta^2}\left[\sqrt{1+\frac{4m^2}{k^2}}-\frac{|k_0|}{2|\mathbf{k}|}\log\left|\frac{\Omega_+}{\Omega_-}\right|\right]$$

$$\mathrm{Re}\left[\Pi_{\mathrm{T}}^{\beta}(k)\right] \simeq \frac{1}{6|\mathbf{k}|^2\beta^2}\left[\frac{k_0^2}{|\mathbf{k}|^2}+\left(1-\frac{k_0^2}{|\mathbf{k}|^2}\right)\frac{k_0}{2|\mathbf{k}|}\log\left|\frac{k_0/|\mathbf{k}|+1}{k_0/|\mathbf{k}|-1}\right|-1\right]+\mathcal{O}\left(\frac{1}{\beta}\right)$$

$$\mathrm{Im}\left[\Pi_{\mathrm{T}}^{\beta}(k)\right] \simeq \frac{2k^2}{|\mathbf{k}|^2}\Bigg\{-\frac{\theta(k^2)\pi}{12|\mathbf{k}|^3\beta^3},$$

$$+\frac{\theta(k^2)+\theta(-k^2-4m^2)}{8\pi|\mathbf{k}|^2\beta^2}\left[\sqrt{1+\frac{4m^2}{k^2}}+\left(1+\frac{4m^2}{k^2}-\frac{k_0^2}{|\mathbf{k}|^2}\right)\frac{|\mathbf{k}|}{2|k_0|}\log\left|\frac{\Omega_+}{\Omega_-}\right|\right]\Bigg\}+\mathcal{O}\left(\frac{1}{\beta}\right),$$

$$\mathcal{N}_{\mathrm{T}}^{\beta}(k) \simeq \frac{2k^2}{|\mathbf{k}|^2}\Bigg\{+\frac{\theta(k^2)\pi}{12|\mathbf{k}|^3\beta^3}-\frac{\theta(k^2)\pi k_0}{24|\mathbf{k}|^3\beta^2},$$

$$-\frac{\theta(k^2)+\theta(-k^2-4m^2)}{8\pi|\mathbf{k}|^2\beta^2}\left[\sqrt{1+\frac{4m^2}{k^2}}+\left(1+\frac{4m^2}{k^2}-\frac{k_0^2}{|\mathbf{k}|^2}\right)\frac{|\mathbf{k}|}{2|k_0|}\log\left|\frac{\Omega_+}{\Omega_-}\right|\right]\Bigg\}+\mathcal{O}\left(\frac{1}{\beta}\right),$$

$$\mathrm{Re}\left[\Pi_{\mathrm{L},T}^{\beta}\right] \sim \mathcal{O}(\beta^{-2})$$

$$\frac{1}{e^{\beta\Omega}-1}=\frac{e^{-\beta\Omega}}{1-e^{-\beta\Omega}}=\sum_{n=1}^{\infty}e^{-n\beta\Omega},$$

$$\mathrm{Re}\left[\Pi_{\mathrm{L}}^{\beta}(k)\right] \simeq -\frac{1}{32}\left[-\frac{k^2}{15m^2}+\mathcal{O}(m^{-4})\right]+e^{-\beta m}\frac{\sqrt{m}\left(\frac{k^2}{k_0^2}-1\right)}{\sqrt{2}(\pi\beta)^{3/2}|\mathbf{k}|^2}+\mathcal{O}\left(m^{-\frac{1}{2}}\right)$$

$$\mathrm{Im}\left[\Pi_{\mathrm{L}}^{\beta}(k)\right] \simeq \frac{\theta(k^2)}{8\pi|\mathbf{k}|^3}\left(1+e^{-\beta|k_0|}\right)e^{-\beta m\frac{|\mathbf{k}|}{|k|}}\left[\frac{4|\mathbf{k}|^2m^2}{\beta k^2}+\frac{8m\mathbf{k}}{\beta^2|k|}+\mathcal{O}(m^0)\right]$$

$$\mathrm{Re}\left[\Pi_{\mathrm{T}}^{\beta}(k)\right] \simeq \frac{k^2}{32\pi^2|\mathbf{k}|^2}\left[-\frac{k^2}{15m^2}+\mathcal{O}(m^{-4})\right]+e^{-\beta m}\left[\frac{\sqrt{m}(k^2+5|\mathbf{k}|^2)}{4\sqrt{2}\pi^{3/2}\beta^{3/2}|\mathbf{k}|^4}+\mathcal{O}\left(m^{-\frac{1}{2}}\right)\right]$$

$$\mathrm{Im}\left[\Pi_{\mathrm{T}}^{\beta}(k)\right] \simeq -\frac{\theta(k^2)}{16\pi|\mathbf{k}|^5}\left(1+e^{-\beta|k_0|}\right)e^{-\beta m\frac{|\mathbf{k}|}{|k|}}\left[\frac{8m|\mathbf{k}||k|}{\beta^2}+\mathcal{O}(m^0)\right]$$

$$\mathcal{N}_{\mathrm{L}}^{\beta}(k) \simeq -\frac{\theta(k^2)}{4\pi|\mathbf{k}|^3}e^{-\beta m\frac{|\mathbf{k}|}{|k|}}\left[\theta(k^0)+e^{\beta|k_0|}\theta(-k^0)+1\right]\left[\frac{4|\mathbf{k}|^2m^2}{\beta k^2}+\frac{8|\mathbf{k}|m}{\beta^2|k|}+\mathcal{O}(m^0)\right]$$

$$\mathcal{N}_{\mathrm{T}}^{\beta}(k) \simeq \frac{\theta(k^2)}{8\pi|\mathbf{k}|^5}e^{-\beta m\frac{|\mathbf{k}|}{|k|}}\left[\theta(k^0)+e^{\beta|k_0|}\theta(-k^0)+1\right]\left[\frac{8m|\mathbf{k}||k|}{\beta^2}+\mathcal{O}(m^0)\right]$$

$$V(\Phi^*\Phi)=\lambda(v^2-\Phi^*\Phi)^2$$

$$\Phi=\left(v+\frac{1}{\sqrt{2}}\zeta\right)e^{\frac{i\chi}{\sqrt{2}v}}$$

$$D_\mu \Phi = \left[\frac{\partial_\mu \zeta}{\sqrt{2}} - i\left(qA_\mu - \frac{\partial_\mu \chi}{\sqrt{2}v}\right)\left(v + \frac{\zeta}{\sqrt{2}}\right)\right] e^{\frac{i\chi}{\sqrt{2}v}}.$$

$$\chi(x) \to \chi(x) + \sqrt{2}qv\lambda(x)$$

mass term $q^2v^2A_\mu A^\mu$ as mixing term $-\sqrt{2}qvA_\mu \partial^\mu \chi$.

$$S_{\mathrm{GF}}[A] \to -\frac{1}{2\xi}\int \mathrm{d}^4x\left(\partial_\mu A^\mu - \sqrt{2}qv\xi\chi\right)^2$$

$$S_{\mathrm{BRST}} = -\int \mathrm{d}^4x\left[\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + q^2v^2A_\mu A^\mu + \frac{\left(\partial_\mu A^\mu\right)^2}{2\xi} + \frac{1}{2}(\partial\chi)^2 + \xi q^2v^2\chi^2 + \bar{c}(\Box - 2\xi q^2v^2)c\right.$$
$$\left. + \frac{1}{2}(\partial\zeta)^2 + 2\lambda v^2\zeta^2 + \sqrt{2}v\lambda\zeta^3 + \frac{\lambda}{4}\zeta^4 + \left(\partial_\mu\chi - \sqrt{2}vqA_\mu\right)\left(\partial^\mu\chi - \sqrt{2}vqA^\mu\right)\left(\frac{\zeta}{\sqrt{2}v} + \frac{\zeta^2}{4v^2}\right)\right]$$

$$\hat{s}A_\mu = \partial_\mu c, \hat{s}\chi = \sqrt{2}vqc, \hat{s}\zeta = 0, \hat{s}c = 0, \hat{s}\bar{c} = -\frac{1}{\xi}\left(\partial_\mu A^\mu + \sqrt{2}qv\xi\chi\right)$$

$$S_{\mathrm{BRST}} = -\int \mathrm{d}^4x\left[-\frac{1}{2}A_\mu\Box A^\mu + q^2v^2A_\mu A^\mu + \frac{1}{2}(\partial\chi)^2 + q^2v^2\chi^2 + \bar{c}(\Box - 2q^2v^2)c\right.$$
$$\left. + \frac{1}{2}(\partial\zeta)^2 + 2\lambda v^2\zeta^2 + \sqrt{2}v\lambda\zeta^3 + \frac{\lambda}{4}\zeta^4 + \mathcal{D}_\mu\chi\mathcal{D}^\mu\chi\left(\frac{\zeta}{\sqrt{2}v} + \frac{\zeta^2}{4v^2}\right)\right],$$

$$\mathcal{D}_\mu\chi \equiv \partial_\mu\chi - \sqrt{2}vqA_\mu,$$

$$\zeta \to H = \zeta + \frac{\zeta^2}{2\sqrt{2}v},$$

$$(\partial\zeta)^2 = (\partial H)^2\left(1 - \frac{\sqrt{2}H}{v} + \frac{2H^2}{v^2} + \cdots\right).$$

$$S_{\text{counterterms}} \simeq -\int \mathrm{d}^4x\left[\frac{a_1}{\sqrt{2}v}H + \frac{a_2}{2v^2}H^2 + \frac{b_2}{2v^2}(\partial H)^2\left(1 - \frac{\sqrt{2}H}{v} + \frac{2H^2}{v^2}\right) + \frac{b_4}{v^4}(\mathcal{D}\chi)^4 + \cdots\right],$$

$$\lim_{x\to y}H(x)H(y) = \lim_{x\to y}\left(\zeta(x) + \frac{\zeta^2(x)}{2\sqrt{2}v}\right)\left(\zeta(y) + \frac{\zeta^2(y)}{2\sqrt{2}v}\right) \propto \frac{1}{v^2(x-y)^4}\zeta(x)\zeta(y) \propto (x-y)^{-2}\mathbb{I} + \cdots \text{ as } x \to y$$

$$S_S[A,\chi,c,\bar{c}] = -\int \mathrm{d}^4x\left[-\frac{1}{2}A_\mu\Box A^\mu + q^2v^2A_\mu A^\mu + \frac{1}{2}(\partial\chi)^2 + q^2v^2\chi^2 + \bar{c}(\Box - 2q^2v^2)c + \frac{b_4}{4v^4}\left(\mathcal{D}_\mu\chi\right)^4\right],$$

$$S_{\mathcal{E}}[H] = -\int \mathrm{d}^4x\left[\frac{1}{2}(\partial H)^2 + \frac{1}{2}M^2H^2 - \frac{(\partial H)^2H}{\sqrt{2}v} + \frac{(\partial H)^2H^2}{v^2} - \frac{\sqrt{2}(\partial H)^2H^3}{v^3} + \cdots\right]$$
$$-\int \mathrm{d}^4x\left[\frac{a_1}{\sqrt{2}v}H + \frac{b_2}{2v^2}(\partial H)^2 + \frac{a_2}{2v^2}H^2 - \frac{b_2(\partial H)^2H}{\sqrt{2}v^3} + \cdots\right]$$

$$S_{\text{int}}[A,\chi,H] := -\frac{1}{\sqrt{2}v}\int \mathrm{d}^4x\left(\mathcal{D}_\mu\chi\right)(\mathcal{D}^\mu\chi)H$$

$$M \equiv 2\sqrt{\lambda}v$$

$$e^{iS_{\rm IF}[A_+,\chi_+,A_-,\chi_-]} = \int \mathcal{D}[H_+]\int \mathcal{D}[H_-]\frac{\mu}{\tilde{\mu}}e^{iS_{\mathcal{E}}[H_+]-iS_{\mathcal{E}}[H_-]+iS_{\rm int}[A_+,\chi_+,H_+]-iS_{\rm int}[A_-,\chi_-,H_-]+iS_{i\epsilon}}$$

$$\mathcal{F}(x,y) \equiv G_{++}(x,y)|_{A_+=A_-=0,\rho=\varrho_{\mathcal{E}}\mathrm{i}} = \int \frac{\mathrm{d}^4k}{(2\pi)^4}\mathcal{F}(k)e^{ik\cdot(x-y)}$$
$$\mathcal{W}(x,y) \equiv G_{-+}(x,y)|_{A_+=A_-=0,\rho=\varrho_{\mathcal{E}}\mathrm{i}} = \int \frac{\mathrm{d}^4k}{(2\pi)^4}\mathcal{W}(k)e^{ik\cdot(x-y)}$$

$$\mathcal{F}(k) = \frac{-i}{k^2+M^2-i\epsilon} + 2\pi\mathfrak{n}(k)\delta(k^2+M^2), \text{ and } \mathcal{W}(k) = 2\pi[\theta(k^0)+\mathfrak{n}(k)]\delta(k^2+M^2)$$

$$J_H = -(\mathcal{D}_\mu\chi)(\mathcal{D}^\mu\chi)/(\sqrt{2}v)$$

$$\begin{aligned}
&S_{\rm IF}[A_+,\chi_+,A_-,\chi_-] \\
&\simeq -\frac{1}{\sqrt{2}v}\int \mathrm{d}^4x\left[\left(\mathscr{D}_+\chi_+(x)\right)^2\langle H_+(x)\rangle_{\mathcal{E}}^{\rm conn} - \left(\mathscr{D}_-\chi_-(x)\right)^2\langle H_-(x)\rangle_{\mathcal{E}}^{\rm conn}\right] \\
&\quad +\frac{i}{4v^2}\int \mathrm{d}^4x\int \mathrm{d}^4y\left[\left(\mathscr{D}_+\chi_+(x)\right)^2\left(\mathscr{D}_+\chi_+(y)\right)^2\langle H_+(x)H_+(y)\rangle_{\mathcal{E}}^{\rm conn}\right. \\
&\qquad -\left(\mathscr{D}_+\chi_+(x)\right)^2\left(\mathscr{D}_-\chi_-(y)\right)^2\langle H_+(x)H_-(y)\rangle_{\mathcal{E}}^{\rm conn} \\
&\qquad -\left(\mathscr{D}_-\chi_-(x)\right)^2\left(\mathscr{D}_+\chi_+(y)\right)^2\langle H_-(x)H_+(y)\rangle_{\mathcal{E}}^{\rm conn} \\
&\qquad \left.+\left(\mathscr{D}_-\chi_-(x)\right)^2\left(\mathscr{D}_-\chi_-(y)\right)^2\langle H_-(x)H_-(y)\rangle_{\mathcal{E}}^{\rm conn}\right]
\end{aligned}$$

$$\begin{aligned}
&-\frac{1}{12\sqrt{2}v^3}\int \mathrm{d}^4x\int \mathrm{d}^4y\int \mathrm{d}^4z\left[\left(\mathscr{D}_+\chi_+(x)\right)^2\left(\mathscr{D}_+\chi_+(y)\right)^2\left(\mathscr{D}_+\chi_+(z)\right)^2\langle H_+(x)H_+(y)H_+(z)\rangle_{\mathcal{E}}^{\rm conn}\right. \\
&\qquad -3\left(\mathscr{D}_+\chi_+(x)\right)^2\left(\mathscr{D}_+\chi_+(y)\right)^2\left(\mathscr{D}_-\chi_-(z)\right)^2\langle H_+(x)H_+(y)H_-(z)\rangle_{\mathcal{E}}^{\rm conn} \\
&\qquad +3\left(\mathscr{D}_+\chi_+(x)\right)^2\left(\mathscr{D}_-\chi_-(y)\right)^2\left(\mathscr{D}_-\chi_-(z)\right)^2\langle H_+(x)H_-(y)H_-(z)\rangle_{\mathcal{E}}^{\rm conn} \\
&\qquad \left.-\left(\mathscr{D}_-\chi_-(x)\right)^2\left(\mathscr{D}_-\chi_-(y)\right)^2\left(\mathscr{D}_-\chi_-(z)\right)^2\langle H_-(x)H_-(y)H_-(z)\rangle_{\mathcal{E}}^{\rm conn}\right]
\end{aligned}$$

$$\mathcal{D}_\pm\chi_\pm \equiv \partial\chi_\pm - \sqrt{2}veA_\pm$$

$$\langle f[H_+,H_-]\rangle_{\mathcal{E}} = \int \mathcal{D}[H_+]\int \mathcal{D}[H_-]f[H_+,H_-]e^{iS_{\mathcal{E}}[H_+]-iS_{\mathcal{E}}[H_-]+S_{i\epsilon}}$$

$$\langle H_+(x)H_+(y)\rangle_{\mathcal{E}}^{\rm conn} \simeq \mathcal{F}(x,y)+\mathcal{O}(v^{-2}), \text{ and } \langle H_-(x)H_+(y)\rangle_{\mathcal{E}}^{\rm conn} \simeq \mathcal{W}(x,y)+\mathcal{O}(v^{-2})$$

$$\begin{aligned}
S_{\rm IF}[A_+,\chi_+,A_-,\chi_-] \simeq \frac{i}{4v^2}\int \mathrm{d}^4x\int \mathrm{d}^4y[&(\mathcal{D}_+\chi_+(x))^2\mathcal{F}(x,y)(\mathcal{D}_+\chi_+(y))^2 \\
&-(\mathcal{D}_+\chi_+(x))^2\mathcal{W}^*(x,y)(\mathcal{D}_-\chi_-(y))^2 \\
&-(\mathcal{D}_-\chi_-(x))^2\mathcal{W}(x,y)(\mathcal{D}_+\chi_+(y))^2 \\
&+(\mathcal{D}_-\chi_-(x))^2\mathcal{F}^*(x,y)(\mathcal{D}_-\chi_-(y))^2] + \mathcal{O}(v^{-4})
\end{aligned}$$

$$\mathcal{F}(x,y) \simeq -\frac{i}{M^2}\delta(x-y) \text{ and }, \mathcal{W}(x,y) \simeq 0$$

$$\lim_{M^2\gg\Box} S_{\rm IF}[A_+,\chi_+,A_-,\chi_-] \simeq \frac{1}{4v^2M^2}\int \mathrm{d}^4x[(\mathcal{D}_+\chi_+(x))^4-(\mathcal{D}_-\chi_-(x))^4]+\mathcal{O}\left(\frac{\Box^2}{M^4}\right),$$

$$\begin{aligned}
S_{\rm IF}[A_+,\chi_+,A_-,\chi_-] \;\simeq\; & \frac{i}{4v^2}\int \mathrm{d}^4x\int \mathrm{d}^4y\,\Big[\,(\mathscr{D}_+\chi_+(x))^2(\mathscr{D}_+\chi_+(y))^2\left(\mathcal{F}(x,y)+\frac{\mathcal{Q}_{\mathcal{F}}(x,y)}{v^2}\right)\\
&-(\mathscr{D}_+\chi_+(x))^2(\mathscr{D}_-\chi_-(y))^2\left(\mathcal{W}^*(x,y)+\frac{\mathcal{Q}^*_{\mathcal{W}}(x,y)}{v^2}\right)\\
&-(\mathscr{D}_-\chi_-(x))^2(\mathscr{D}_+\chi_+(y))^2\left(\mathcal{W}(x,y)+\frac{\mathcal{Q}_{\mathcal{W}}(x,y)}{v^2}\right)\\
&+(\mathscr{D}_-\chi_-(x))^2(\mathscr{D}_-\chi_-(y))^2\left(\mathcal{F}^*(x,y)+\frac{\mathcal{Q}^*_{\mathcal{F}}(x,y)}{v^2}\right)\,\Big]\\
-\frac{i}{24v^4}\int \mathrm{d}^4x\int \mathrm{d}^4y\int \mathrm{d}^4z\,\Big[\,&(\mathscr{D}_+\chi_+(x))^2(\mathscr{D}_+\chi_+(y))^2(\mathscr{D}_+\chi_+(z))^2\,\Gamma_{\rm u}(x,y,z)\\
&-3(\mathscr{D}_+\chi_+(x))^2(\mathscr{D}_+\chi_+(y))^2(\mathscr{D}_-\chi_-(z))^2\,\Gamma^*_{\rm n}(x,y,z)\\
&-3(\mathscr{D}_+\chi_+(x))^2(\mathscr{D}_-\chi_-(y))^2(\mathscr{D}_-\chi_-(z))^2\,\Gamma_{\rm n}(x,y,z)\\
&+(\mathscr{D}_-\chi_-(x))^2(\mathscr{D}_-\chi_-(y))^2(\mathscr{D}_-\chi_-(z))^2\,\Gamma^*_{\rm u}(x,y,z)\,\Big]+\mathcal{O}(v^{-6})
\end{aligned}$$


$$\Gamma_{\rm u,n}(x,y,z)=\int\frac{\mathrm{d}^4k}{(2\pi)^4}\int\frac{\mathrm{d}^4p}{(2\pi)^4}\int\frac{\mathrm{d}^4q}{(2\pi)^4}(2\pi)^4\delta^{(4)}(k+p+q)\gamma_{\rm u,n}(k,p,q)e^{ik\cdot x+ip\cdot y+iq\cdot z}$$


$$\gamma_{\rm u}(k,p,q)=(k^2+p^2+q^2)(\mathcal{F}(k)\mathcal{F}(p)\mathcal{F}(q)-\mathcal{W}(-k)\mathcal{W}(-p)\mathcal{W}(-q))$$
$$\gamma_{\rm n}(k,p,q)=(k^2+p^2+q^2)(\mathcal{F}^*(k)\mathcal{F}^*(p)\mathcal{W}(-q)-\mathcal{W}(k)\mathcal{W}(p)\mathcal{F}(q))$$

$$\mathcal{Q}_{\mathcal{F},\mathcal{W}}(x,y)=\int\frac{\mathrm{d}^4k}{(2\pi)^4}\mathcal{Q}_{\mathcal{F},\mathcal{W}}(k)e^{ik\cdot(x-y)}$$

$$\begin{bmatrix}\mathcal{Q}_{\mathcal{F}}(k) & \mathcal{Q}_{\mathcal{W}}(-k)\\ \mathcal{Q}_{\mathcal{W}}(k) & \mathcal{Q}^*_{\mathcal{F}}(k)\end{bmatrix}=\begin{bmatrix}\mathcal{F}(k) & \mathcal{W}(-k)\\ \mathcal{W}(k) & \mathcal{F}^*(k)\end{bmatrix}\begin{bmatrix}\Sigma_{\mathcal{F}}(k) & \Sigma_{\mathcal{W}}(-k)\\ \Sigma_{\mathcal{W}}(k) & \Sigma^*_{\mathcal{F}}(k)\end{bmatrix}\begin{bmatrix}\mathcal{F}(k) & \mathcal{W}(-k)\\ \mathcal{W}(k) & \mathcal{F}^*(k)\end{bmatrix}$$

$$\Sigma_{\mathcal{F}}(k)=-\frac{1}{4}\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\mathcal{F}(\ell)\mathcal{F}(k-\ell)$$
$$-2i\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+\ell^2)\mathcal{F}(\ell)-i[a_2+b_2k^2+b_4(k^2+M^2)^2]$$
$$\Sigma_{\mathcal{W}}(k)=\frac{1}{4}\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\mathcal{W}(\ell)\mathcal{W}(k-\ell)$$





$$S_{\mathrm{IF}}[A_{\mathrm{r}},\chi_{\mathrm{r}},A_{\mathrm{a}},\chi_{\mathrm{a}}] \simeq \frac{i}{2v^2}\int \mathrm{d}^4x \int \mathrm{d}^4y \Big[(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(x)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(x))(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(y))^2\left(\mathcal{R}(x,y)+\frac{\mathcal{Q}_{\mathcal{R}}(x,y)}{v^2}\right)$$
$$+(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(x))^2(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(y)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(y))\left(\mathcal{A}(x,y)+\frac{\mathcal{Q}_{\mathcal{A}}(x,y)}{v^2}\right)$$
$$+8(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(x)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(x))(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(y)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(y))\left(\mathcal{K}(x,y)+\frac{\mathcal{Q}_{\mathcal{K}}(x,y)}{v^2}\right)\Big]$$
$$-\frac{i}{24v^4}\int \mathrm{d}^4x \int \mathrm{d}^4y \int \mathrm{d}^4z[(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(x)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(x))(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(y))^2(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(z))^2(\Gamma_{\mathrm{u}}-\Gamma_{\mathrm{u}}^*-3\Gamma_{\mathrm{n}}-3\Gamma_{\mathrm{n}}^*)(x,y,z)$$
$$+(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(x))^2(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(y)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(y))(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(z))^2(\Gamma_{\mathrm{u}}-\Gamma_{\mathrm{u}}^*+3\Gamma_{\mathrm{n}}-3\Gamma_{\mathrm{n}}^*)(x,y,z)$$
$$+(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(x))^2(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(y))^2(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(z)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(z))(\Gamma_{\mathrm{u}}-\Gamma_{\mathrm{u}}^*+3\Gamma_{\mathrm{n}}+3\Gamma_{\mathrm{n}}^*)(x,y,z)$$
$$+(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(x)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(x))(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(y)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(y))(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(z))^2(\Gamma_{\mathrm{u}}+\Gamma_{\mathrm{u}}^*+3\Gamma_{\mathrm{n}}-3\Gamma_{\mathrm{n}}^*)(x,y,z)$$
$$+(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(x)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(x))(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(y))^2(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(z)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(z))(\Gamma_{\mathrm{u}}+\Gamma_{\mathrm{u}}^*+3\Gamma_{\mathrm{n}}+3\Gamma_{\mathrm{n}}^*)(x,y,z)$$
$$+(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(x))^2(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(y)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(y))(\mathcal{D}_{\mathrm{a}}\chi_{\mathrm{a}}(z)\cdot\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(z))(\Gamma_{\mathrm{u}}+\Gamma_{\mathrm{u}}^*-3\Gamma_{\mathrm{n}}+3\Gamma_{\mathrm{n}}^*)(x,y,z)$$

$$\mathcal{D}_{a,r}\chi_{a,r} \equiv \partial\chi_{a,r} - \sqrt{2}veA_{a,r}$$

$$\mathcal{R} = \tfrac{i}{2}\mathrm{Im}[\mathcal{F}+\mathcal{W}], \mathcal{A} = \tfrac{i}{2}\mathrm{Im}[\mathcal{F}-\mathcal{W}] \text{ and } \mathcal{K} = \tfrac{1}{2}\mathrm{Re}[\mathcal{F}+\mathcal{W}]$$

$$\mathcal{Q}_{\mathcal{R}} \equiv \frac{i}{2}\mathrm{Im}[\mathcal{Q}_{\mathcal{F}}+\mathcal{Q}_{\mathcal{W}}], \mathcal{Q}_{\mathcal{A}} \equiv \frac{i}{2}\mathrm{Im}[\mathcal{Q}_{\mathcal{F}}-\mathcal{Q}_{\mathcal{W}}] \text{ and }, \mathcal{Q}_{\mathcal{K}} \equiv \frac{1}{2}\mathrm{Re}[\mathcal{Q}_{\mathcal{F}}+\mathcal{Q}_{\mathcal{W}}]$$

$$[\mathcal{F}+\mathcal{F}^*-\mathcal{W}-\mathcal{W}^*](x,y) = 0$$
$$[\mathcal{Q}_{\mathcal{F}}+\mathcal{Q}_{\mathcal{F}}^*-\mathcal{Q}_{\mathcal{W}}-\mathcal{Q}_{\mathcal{W}}^*](x,y) = 0$$

$$[\Gamma_{\mathrm{u}}+\Gamma_{\mathcal{U}}^*-3\Gamma_{\mathcal{N}}-3\Gamma_{\mathcal{N}}^*](x,y,z) \to 0$$

$$\mathfrak{n}(k) = \theta(k^0)n(|\mathbf{k}|)+\theta(-k^0)n(|\mathbf{k}|) = n(|\mathbf{k}|)$$

$$\Sigma_{\mathcal{F}}(k) = i\frac{M^2(M^2-k^2)}{8\pi^2}\log\left(\frac{M^2}{\mu^2}\right)$$
$$-i\int_{-1}^{+1}\mathrm{d}y\frac{y^2(5y^2+3)(k^2)^2-12(2y^2+1)k^2M^2+24M^4}{256\pi^2}\log\left|\frac{(1-y^2)k^2+4M^2}{\mu^2}\right|$$
$$-\frac{ik^2}{\pi^2}\int_M^\infty \mathrm{d}\Omega\sqrt{\Omega^2-M^2}n\left(\sqrt{\Omega^2-M^2}\right)$$
$$-\frac{i(k^2-2M^2)^2}{32\pi^2|\mathbf{k}|}\int_M^\infty \mathrm{d}\Omega n\left(\sqrt{\Omega^2-M^2}\right)\log\left|\frac{\left(k^2-2|\mathbf{k}|\sqrt{\Omega^2-M^2}\right)^2-4k_0^2\Omega^2}{\left(k^2+2|\mathbf{k}|\sqrt{\Omega^2-M^2}\right)^2-4k_0^2\Omega^2}\right|$$
$$-\theta(-k^2-4M^2)\frac{(k^2-2M^2)^2}{64\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+}\mathrm{d}\Omega\left[1+4n\left(\sqrt{\Omega^2-M^2}\right)\right.$$
$$\left.+2n\left(\sqrt{\Omega^2-M^2}\right)n\left(\sqrt{(\Omega-|k_0|)^2-M^2}\right)\right]$$
$$-\theta(k^2)\frac{(k^2-2M^2)^2}{32\pi|\mathbf{k}|}\int_{\Omega_+}^\infty \mathrm{d}\Omega\left[n\left(\sqrt{\Omega^2-M^2}\right)+n\left(\sqrt{(\Omega-|k_0|)^2-M^2}\right)\right.$$
$$\left.+2n\left(\sqrt{\Omega^2-M^2}\right)n\left(\sqrt{(\Omega-|k_0|)^2-M^2}\right)\right]$$
$$\Sigma_{\mathcal{W}}(k) = \theta(-k^2-4M^2)\frac{(k^2-2M^2)^2}{32\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+}\mathrm{d}\Omega\left[\theta(k^0)+2\theta(k^0)n\left(\sqrt{\Omega^2-M^2}\right)\right.$$
$$\left.+n\left(\sqrt{\Omega^2-M^2}\right)n\left(\sqrt{(\Omega-|k_0|)^2-M^2}\right)\right]$$
$$+\theta(k^2)\frac{(k^2-2M^2)^2}{16\pi|\mathbf{k}|}\int_{\Omega_+}^\infty \mathrm{d}\Omega\left[\theta(-k^0)n\left(\sqrt{\Omega^2-M^2}\right)+\theta(k^0)n\left(\sqrt{(\Omega-|k_0|)^2-M^2}\right)\right.$$
$$\left.+n\left(\sqrt{\Omega^2-M^2}\right)n\left(\sqrt{(\Omega-|k_0|)^2-M^2}\right)\right]$$

$$\Omega_\pm = \frac{|k_0|}{2}\pm\frac{|\mathbf{k}|}{2}\sqrt{1+\frac{4M^2}{k^2}}$$

$$-\Sigma_{\mathcal{F}}(k)\simeq\Sigma_{\mathcal{W}}(k)\simeq\frac{\theta(-k^2-4M^2)(-k^2+2M^2)^2}{32\pi|\mathbf{k}|}\left[\frac{\log\left(\frac{\Omega_+}{\Omega_-}\right)-\log\left(\frac{|k_0|-\Omega_+}{|k_0|-\Omega_-}\right)}{|k_0|\beta^2}+\mathcal{O}(\beta^{-1})\right]$$
$$+\frac{\theta(k^2)(-k^2+2M^2)^2}{16\pi|\mathbf{k}|}\left[\frac{\log\left(\frac{\Omega_+}{\Omega_+-|k_0|}\right)}{|k_0|\beta^2}+\mathcal{O}(\beta^{-1})\right]$$

$$S=-\int \mathrm{d}^4x\left[\frac{1}{4}F_{\mu\nu}F^{\mu\nu}+|D^\mu[A]\Phi|^2+\lambda(v^2-\Phi^*\Phi)^2\right.$$
$$\left.+\sum_I\left(-|D_0[A]\varphi_I|^2+\Gamma_I^2|\varphi_I|^2-g_I(\Phi^*\varphi_I+\varphi_I^*\Phi)\right)\right]$$

$$\langle\Phi\rangle=v_0=\sqrt{v^2+\frac{1}{2\lambda}\sum_I\frac{g_I^2}{\Gamma_I^2}},\langle\varphi_I\rangle=v_I=\frac{g_I}{\Gamma_I^2}v_0.$$

$\Phi=\left(v_0+\zeta/\sqrt{2}\right)$ with ζ real and $\varphi_I=v_I+\frac{1}{\sqrt{2}}(\alpha_I+i\beta_I)$

$$\Phi \to \left(v_0 + \frac{1}{\sqrt{2}}\zeta\right) e^{\frac{i\chi}{\sqrt{2}v_0}}, \varphi_I \to \left(v_I + \frac{1}{\sqrt{2}}(\alpha_I + i\beta_I)\right) e^{\frac{i\chi}{\sqrt{2}v_0}}$$

$$S = -\int \mathrm{d}^4x \left(\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}(\partial_\mu\zeta)^2 + \frac{\left(v_0 + \frac{1}{\sqrt{2}}\zeta\right)^2}{2v_0^2}(\mathcal{D}_\mu\chi)^2 \right.$$

$$+\sum_I \left[-\frac{1}{2}(\partial_0\alpha_I)^2 - \frac{1}{2}(\partial_0\beta_I)^2 - \frac{\left(v_I + \frac{1}{\sqrt{2}}\alpha_I\right)^2 + \frac{1}{2}\beta_I^2}{2v_0^2}(\mathcal{D}_0\chi)^2 + \Gamma_I^2\left(v_I^2 + \frac{1}{2}\alpha_I^2 + \frac{1}{2}\beta_I^2\right) \right.$$

$$\left. -\frac{1}{v_0}\left(v_I\partial_0\beta_I + \frac{\alpha_I\partial_0\beta_I - \beta_I\partial_0\alpha_I}{\sqrt{2}}\right)\mathcal{D}^0\chi - 2g_I\left(v_0v_I + \frac{1}{2}\zeta\alpha_I\right)\right]$$

$$\left. +\lambda\left[(v^2 - v_0^2)^2 + (3v_0^2 - v^2)\zeta^2 + \sqrt{2}v_0\zeta^3 + \frac{1}{4}\zeta^4\right]\right)$$

$$\mathcal{D}_\mu\chi \equiv \partial_\mu\chi - \sqrt{2}qv_0A_\mu$$

$$(\alpha_I + i\beta_I) \to e^{i\theta}(\alpha_I + i\beta_I)$$

$$S_{\text{in-in}} = -\int \mathrm{d}^4x \left[\frac{1}{4}F_{\mu\nu}^+F_+^{\mu\nu} + \frac{1}{2}(\partial_\mu\zeta_+)^2 + \lambda\left((3v_0^2 - v^2)\zeta_+^2 + \sqrt{2}v_0\zeta_+^3 + \frac{1}{4}\zeta_+^4\right) \right.$$

$$\left. + \frac{\left(v_0 + \frac{1}{\sqrt{2}}\zeta_+\right)^2}{2v_0^2}(\mathcal{D}_\mu^+\chi_+)^2 - \sum_I \frac{v_I^2}{2v_0^2}(\mathcal{D}_0^+\chi_+)^2 \right]$$

$$+\int \mathrm{d}^4x \left[\frac{1}{4}F_{\mu\nu}^-F_-^{\mu\nu} + \frac{1}{2}(\partial_\mu\zeta_-)^2 + \lambda\left((3v_0^2 - v^2)\zeta_-^2 + \sqrt{2}v_0\zeta_-^3 + \frac{1}{4}\zeta_-^4\right) \right.$$

$$\left. + \frac{\left(v_0 + \frac{1}{\sqrt{2}}\zeta_-\right)^2}{2v_0^2}(\mathcal{D}_\mu^-\chi_-)^2 - \sum_I \frac{v_I^2}{2v_0^2}(\mathcal{D}_0^-\chi_-)^2 \right]$$

$$+S_{\text{IF}}[A_+, \zeta_+, \chi_+, A_-, \zeta_-, \chi_-]$$

$$e^{iS_{\text{IF}}[A_+,\zeta_+,\chi_+,A_-,\zeta_-,\chi_-]} =$$

$$\int \mathcal{D}[\alpha_+, \beta_+] \int \mathcal{D}[\alpha_-, \beta_-] e^{iS_{\mathcal{E}}[\alpha_+,\beta_+] - iS_{\mathcal{E}}[\alpha_-,\beta_-] + iS_{\text{int}}[\alpha_+,\beta_+,A_+,\zeta_+,\chi_+] - iS_{\text{int}}[\alpha_-,\beta_-,\chi_-,A_-,\zeta_-,\chi_-] + iS_{i\epsilon}},$$

$$S_{\mathcal{E}}[\alpha, \beta] = \int \mathrm{d}^4x \sum_I \left[\frac{1}{2}(\partial_0\alpha_I)^2 + \frac{1}{2}(\partial_0\beta_I)^2 - \frac{1}{2}\Gamma_I^2(\alpha_I^2 + \beta_I^2) \right]$$

$$S_{\text{int}}[\alpha, \beta, A, \zeta, \chi] =$$

$$\int \mathrm{d}^4x \sum_I \left[\frac{\sqrt{2}v_I\alpha_I + \frac{1}{2}(\alpha_I^2 + \beta_I^2)}{2v_0^2}(\mathcal{D}_0\chi)^2 + \frac{1}{v_0}\left(v_I\partial_0\beta_I + \frac{\alpha_I\partial_0\beta_I - \beta_I\partial_0\alpha_I}{\sqrt{2}}\right)\mathcal{D}^0\chi + g_I\zeta\alpha_I \right]$$

$$S_{\text{int}}[\alpha, \beta, A, \zeta, \chi] \simeq \int \mathrm{d}^4x \sum_I g_I \left[\frac{\alpha_I}{\sqrt{2}\Gamma_I^2 v_0}(\mathcal{D}_0\chi)^2 - \frac{1}{\Gamma_I^2}\beta_I\partial_0\mathcal{D}^0\chi + \zeta\alpha_I \right]$$

$$v_I/v_0 = g_I/\Gamma_I^2$$

$$S_{\mathrm{IF}}[A_+,\chi_+,\zeta_+,A_-,\chi_-,\zeta_-] = \frac{1}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I ig_I^2\left(\tilde{\zeta}_I^+(x) - \tilde{\zeta}_I^-(x)\right)\mathbf{D}_I(x-y)\begin{pmatrix}\tilde{\zeta}_I^+(y)\\ -\tilde{\zeta}_I^-(y)\end{pmatrix}$$
$$+\frac{1}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I \frac{ig_I^2}{\Gamma_I^4}(\mathcal{D}_0^+\chi^+(x) - \mathcal{D}_0^-\chi^-(x))\partial_0^2\mathbf{D}_I(x-y)\begin{pmatrix}\mathcal{D}_0^+\chi^+(y)\\ -\mathcal{D}_0^-\chi^-(y)\end{pmatrix}$$

$$\tilde{\zeta}_I^{\pm}(x) = \zeta^{\pm}(x) + \frac{1}{\sqrt{2}\Gamma_I^2 v_0}(\mathcal{D}_0\chi^{\pm})^2$$

$$\mathbf{D}_I(k) = \begin{pmatrix} -\frac{i}{-\omega^2+\Gamma_I^2-i\epsilon} + 2\pi\delta(-\omega^2+\Gamma_I^2)\mathfrak{n}_I(k) & \theta(-\omega)2\pi\delta(-\omega^2+\Gamma_I^2) + 2\pi\delta(-\omega^2+\Gamma_I^2)\mathfrak{n}_I(k) \\ \theta(\omega)2\pi\delta(-\omega^2+\Gamma_I^2) + 2\pi\delta(-\omega^2+\Gamma_I^2)\mathfrak{n}_I(k) & \frac{i}{-\omega^2+\Gamma_I^2+i\epsilon} + 2\pi\delta(-\omega^2+\Gamma_I^2)\mathfrak{n}_I(k) \end{pmatrix}$$

$$\partial_0^2\mathbf{D}_I(x-y) + \Gamma_I^2\mathbf{D}_I(x-y) = -i\delta^4(x-y)\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}$$

$$S_{\mathrm{IF}}[A_+,\chi_+,\zeta_+,A_-,\chi_-,\zeta_-] = -\frac{C}{2}\int \mathrm{d}^4x[(\mathcal{D}_0^+\chi^+(x))^2 - (\mathcal{D}_0^-\chi^-(x))^2]$$
$$-\frac{1}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y(\mathcal{D}_0^+\chi^+(x) - \mathcal{D}_0^-\chi^-(x))\mathbf{d}(x-y)\begin{pmatrix}\mathcal{D}_0^+\chi^+(y)\\ -\mathcal{D}_0^-\chi^-(y)\end{pmatrix}$$
$$+\frac{1}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y \sum_I ig_I^2\left(\tilde{\zeta}_I^+(x) - \tilde{\zeta}_I^-(x)\right)\mathbf{D}_I(x-y)\begin{pmatrix}\tilde{\zeta}_I^+(y)\\ -\tilde{\zeta}_I^-(y)\end{pmatrix}$$

$$C = \sum_I \frac{g_I^2}{\Gamma_I^4}, \text{ and } \mathbf{d}(x-y) = \sum_I \frac{ig_I^2}{\Gamma_I^2}\mathbf{D}_I(x-y)$$

$$S_{\mathrm{in-in}}[A_+,\chi_+,A_-,\chi_-] = \int d^4x\left(-\frac{1}{4}F_{\mu\nu}^+F^{+\mu\nu} - \frac{1}{2}\left(\mathcal{D}_\mu^+\chi^+\right)^2 + \frac{1}{4}F_{\mu\nu}^-F^{-\mu\nu} + \frac{1}{2}\left(\mathcal{D}_\mu^-\chi^-\right)^2\right)$$
$$-C\int d^4x[(\mathcal{D}_0^+\chi^+(x))^2 - (\mathcal{D}_0^-\chi^-(x))^2]$$
$$-\frac{1}{2}\int d^4x \int d^4y(\mathcal{D}_0^+\chi^+(x) - \mathcal{D}_0^-\chi^-(x))\mathbf{d}(x-y)\begin{pmatrix}\mathcal{D}_0^+\chi^+(y)\\ -\mathcal{D}_0^-\chi^-(y)\end{pmatrix} + \cdots$$

$$S_{\text{in-in}}[A^{\mathrm{r}},A^{\mathrm{a}},\chi^{\mathrm{r}},\chi^{\mathrm{a}}] = \int d^4x\left[-\frac{1}{2}F_{\mu\nu}^{\mathrm{r}}F^{\mathrm{a}\mu\nu} - \mathcal{D}_\mu^{\mathrm{r}}\chi^{\mathrm{r}}\mathcal{D}^{\mathrm{a}\mu}\chi^{\mathrm{a}} - 2C\mathcal{D}_0^{\mathrm{r}}\chi^{\mathrm{r}}\mathcal{D}_0^{\mathrm{a}}\chi^{\mathrm{a}}\right]$$
$$-\int \frac{d^4k}{(2\pi)^4}\left[\mathcal{D}_0^{\mathrm{a}}\chi^{\mathrm{a}}(-k)d^D(\omega,\mathbf{k})\mathcal{D}_0^{\mathrm{r}}\chi^{\mathrm{r}}(k) + \frac{i}{2}\mathcal{D}_0^{\mathrm{a}}\chi^{\mathrm{a}}(-k)d^N(\omega,\mathbf{k})\mathcal{D}_0^{\mathrm{a}}\chi^{\mathrm{a}}(k)\right] + \cdots,$$

$$d^N(\omega,\mathbf{k}) = \sum_I \frac{g_I^2}{\Gamma_I^2}2\pi\delta(-\omega^2+\Gamma_I^2)\left(\frac{1}{2} + \mathfrak{n}_I(\mathbf{k})\right)$$

$$d^D(\omega,\mathbf{k}) = \sum_I \frac{g_I^2}{\Gamma_I^2}\frac{1}{-(\omega+i\epsilon)^2+\Gamma_I^2}$$

$$\partial^\mu F_{\mu\nu} = -J_\nu^{\text{medium}} - J_\nu^{\text{free}}$$

$$J^{\mu}_{\text{medium}} = \partial_{\nu} P^{\mu\nu},$$

$$\partial^{\mu}\left(F_{\mu\nu} + P_{\mu\nu}\right) = -J^{\text{free}}_{\nu}$$

$$P_{\mu\nu} = P_{\mu\nu}\left[F_{\alpha\beta}\right]$$

$$\frac{\delta P_{\mu\nu}(x)}{\delta F_{\alpha\beta}(y)} = 0, \text{ past lightcone of } x.$$

$$S_{\text{in-in}} = \int \mathrm{d}^4x \left[-\frac{1}{2}\left(F^{\text{r}}_{\mu\nu} + P_{\mu\nu}\left[F^{\text{r}}_{\alpha\beta}\right]\right)F^{\mu\nu}_{\text{a}} + J^{\text{r}}_{\mu}A^{\mu}_{\text{a}} + J^{\text{a}}_{\mu}A^{\mu}_{\text{r}}\right]$$

$$\frac{\delta S_{\text{in-in}}}{\delta A^{\text{a}}_{\nu}(x)} = \partial^{\mu}\left(F^{\text{r}}_{\mu\nu} + P_{\mu\nu}\left[F^{\text{r}}_{\alpha\beta}\right]\right) + J^{\text{r}}_{\nu} = 0$$

$$S_{\text{in-in}} = \int \mathrm{d}^4x \left[-\frac{1}{2}\left(F^{\text{r}}_{\mu\nu} + P_{\mu\nu}\left[F^{\text{r}}_{\alpha\beta}\right]\right)F^{\mu\nu}_{\text{a}}\right] + \int \mathrm{d}^4x \int \mathrm{d}^4y F^{\mu\nu}_{\text{a}}(x) K_{\mu\nu;\alpha\beta}\left[F^{\text{r}}_{\alpha\beta}\right](x,y) F^{\alpha\beta}_{\text{a}}(y) + \cdots$$
$$+ \int \mathrm{d}^4x \left[J^{\text{r}}_{\mu}(x)A^{\mu}_{\text{a}}(x) + J^{\text{a}}_{\mu}(x)A^{\mu}_{\text{r}}(x)\right]$$

$$K_{\mu\nu;\alpha\beta}\left[F^{\text{r}}_{\alpha\beta}\right](x,y) \star P^{\mu\nu}\left[F^{\text{r}}_{\alpha\beta}\right](x) = -\frac{\delta S_{\text{EH}}\left[F^{\text{r}}_{\alpha\beta}\right]}{\delta F^{\text{r}}_{\mu\nu}(x)} + \cdots$$

$$H_{\mu\nu} = F_{\mu\nu} + P_{\mu\nu}$$

$$H_{\rho\sigma}(x) \equiv \int \mathrm{d}^4y \chi_{\alpha\beta\rho\sigma}(x,y) F^{\alpha\beta}(y)$$

$$\chi^{\mu\nu\alpha\beta} = -\chi^{\nu\mu\alpha\beta} = -\chi^{\mu\nu\beta\alpha}$$

$$S_{\text{in-in}} \equiv -\frac{1}{2}\int \mathrm{d}^4x \int \mathrm{d}^4y F^{\alpha\beta}_{\text{r}}(x) \chi_{\alpha\beta\rho\sigma}(x,y) F^{\rho\sigma}_{\text{a}}(y) + \int \mathrm{d}^4x \int \mathrm{d}^4y F^{\mu\nu}_{\text{a}}(x) K_{\mu\nu;\alpha\beta}\left[F^{\text{r}}_{\alpha\beta}\right](x,y) F^{\alpha\beta}_{\text{a}}(y)$$
$$+ \int \mathrm{d}^4x \left[J^{\text{r}}_{\mu}(x)A^{\mu}_{\text{a}}(x) + J^{\text{a}}_{\mu}(x)A^{\mu}_{\text{r}}(x)\right]$$

$$f_n \equiv f_n(x-y) \text{ and } g_n \equiv g_n(x-y)$$

$$\tilde{F}_{\mu\nu} \equiv \frac{1}{2}\epsilon_{\mu\nu\rho\sigma}F^{\rho\sigma}$$

$$S_{\text{in-in}} = \int \mathrm{d}^4x \left[-\frac{1}{2}F^{\text{r}}_{\alpha\beta}F^{\alpha\beta}_{\text{a}} + \theta F^{\alpha\beta}_{\text{r}}\tilde{F}^{\text{a}}_{\alpha\beta}\right] + S_{\text{IF}}$$

$$\partial_{[\alpha}F_{\beta\gamma]} = 0$$

$$\partial_i B_i = 0, \text{ and } \partial_0 B_k = -\epsilon_{ijk}\partial_i E_j,$$

$$S_{\text{supergravity}} = \int \mathrm{d}^{\square}x \int \mathrm{d}^{\square}y \Big\{ f_1(x-y) F^{\mathrm{r}}_{\alpha\beta}(x) F^{\alpha\beta}_{\mathrm{a}}(y) + f_2(x-y) F^{\mathrm{r}\gamma}_{\alpha}(x) F^{\mathrm{a}}_{\gamma\beta}(y) n^{\alpha} n^{\beta}$$
$$+ f_3(x-y) F^{\mathrm{r}}_{\alpha\beta}(x) \tilde{F}^{\alpha\beta}_{\mathrm{a}}(y) + f_4(x-y) F^{\mathrm{r}\gamma}_{\alpha}(x) \tilde{F}^{\mathrm{a}}_{\beta\gamma}(y) n^{\alpha} n^{\beta} + f_5(x-y) \tilde{F}^{\mathrm{r}}_{\alpha\beta}(x) A^{\beta}_{\mathrm{a}}(y) n^{\alpha}$$
$$+ f_6(x-y) F^{\gamma\alpha}_{\mathrm{r}}(x) \partial_{\gamma} \tilde{F}^{\mathrm{a}}_{\beta\alpha}(y) n^{\beta} + f_7(x-y) \tilde{F}^{\mathrm{r}}_{\alpha\beta}(x) \partial_{\gamma} F^{\gamma\alpha}_{\mathrm{a}}(y) n^{\beta}$$
$$+ f_8(x-y) \partial_{\alpha} F^{\alpha\beta}_{\mathrm{r}}(x) \partial_{\gamma} F^{\mathrm{a}\gamma}_{\beta}(y) + f_9(x-y) \partial_{\alpha} F^{\alpha}_{\mathrm{r}\sigma}(x) \partial_{\nu} F^{\nu}_{\mathrm{a}\rho}(y) n^{\sigma} n^{\rho} \Big\}$$

$$S_{\text{gravity}} = \frac{i}{2} \int \mathrm{d}^4 x \int \mathrm{d}^4 y \Big\{ g_1(x-y) F^{\mathrm{a}}_{\alpha\beta}(x) F^{\alpha\beta}_{\mathrm{a}}(y) + g_2(x-y) F^{\mathrm{a}}_{\alpha\gamma}(x) F^{\mathrm{a}\gamma}_{\beta}(y) n^{\alpha} n^{\beta}$$
$$+ g_3(x-y) F^{\mathrm{a}}_{\alpha\beta}(x) \tilde{F}^{\alpha\beta}_{\mathrm{a}}(y) + g_4(x-y) F^{\mathrm{a}\gamma}_{\alpha}(x) \tilde{F}^{\mathrm{a}}_{\beta\gamma}(y) n^{\alpha} n^{\beta} + g_5(x-y) \tilde{F}^{\mathrm{a}}_{\alpha\beta}(x) A^{\alpha}_{\mathrm{a}}(y) n^{\beta}$$
$$+ g_6(x-y) F^{\gamma\alpha}_{\mathrm{a}} \partial_{\gamma} \tilde{F}^{\mathrm{a}}_{\beta\alpha} n^{\beta} + g_7(x-y) \partial_{\alpha} F^{\alpha\beta}_{\mathrm{a}}(x) \partial_{\gamma} F^{\mathrm{a}\gamma}_{\beta}(y) + g_8(x-y) \partial_{\alpha} F^{\alpha}_{\mathrm{a}\sigma}(x) \partial_{\nu} F^{\nu}_{\mathrm{a}\rho} n^{\sigma} n^{\rho} \Big\}$$

$\partial_{\nu} F^{\nu}{}_{\rho}(y)$ which in S -matrix

$$n^{\alpha} = (1, \mathbf{0}), \left(n_{\lambda} n^{\lambda} = -1\right)$$

$F_{0i} = -E_i, F_{ij} = \epsilon_{ijk} B^k$, and using that $\epsilon^{ijk} \epsilon_{jkl} = 2\delta^i_l$ and $\epsilon_{0ijk} = -\epsilon_{ijk}$

$$S_{\text{supergravity}} = \int \mathrm{d}^{\square}x \int \mathrm{d}^{\square}y \{ f_1(x-y) [E^{\mathrm{r}}_i(x) E^{\mathrm{a}}_i(y) - B^{\mathrm{r}}_i(x) B^{\mathrm{a}}_i(y)] + f_2(x-y) E^{\mathrm{r}}_i(x) E^{\mathrm{a}}_i(y)$$
$$+ f_3(x-y) [E^{\mathrm{r}}_i(x) B^{\mathrm{a}}_i(y) + B^{\mathrm{r}}_i(x) E^{\mathrm{a}}_i(y)] + f_4(x-y) E^{\mathrm{r}}_i(x) B^{\mathrm{a}}_i(y) - f_5(x-y) B^{\mathrm{r}}_i(x) A^{\mathrm{a}}_i(y)$$
$$- f_6(x-y) \epsilon_{ijk} B^{\mathrm{r}}_k(x) \partial_i B^{\mathrm{a}}_j(y) + f_8(x-y) \partial_i E^{\mathrm{r}}_i(x) \partial_j E^{\mathrm{a}}_j(y) \}$$

$$S_{\text{gravity}} = \frac{i}{2} \int \mathrm{d}^4 x \int \mathrm{d}^4 y \{ g_1(x-y) [E^{\mathrm{a}}_i(x) E^{\mathrm{a}}_i(y) - B^{\mathrm{a}}_i(x) B^{\mathrm{a}}_i(y)] + g_2(x-y) E^{\mathrm{a}}_i(x) E^{\mathrm{a}}_i(y)$$
$$+ g_3(x-y) [E^{\mathrm{a}}_i(x) B^{\mathrm{a}}_i(y) + B^{\mathrm{a}}_i(x) E^{\mathrm{a}}_i(y)] + g_4(x-y) B^{\mathrm{a}}_i(x) E^{\mathrm{a}}_i(y) - g_5(x-y) B^{\mathrm{a}}_i(x) A^{\mathrm{a}}_i(y)$$
$$- g_6(x-y) \epsilon_{ijk} B^{\mathrm{a}}_k(x) \partial_i B^{\mathrm{a}}_j(y) + g_7(x-y) \partial_i E^{\mathrm{a}}_i(x) \partial_j E^{\mathrm{a}}_j(y) \}$$

$$f_n(x-y) = \sum_{r=0}^{p} \frac{1}{r!} c_{nr} \partial^r_{x^0} \delta(x-y), g_n(x-y) = \sum_{r=0}^{p} \frac{1}{r!} g_{nr} \partial^r_{x^0} \delta(x-y)$$

$$S_{\text{supergravity}} = \int \frac{\mathrm{d}^{\square}k}{(2\pi)^4} \Big\{ 2 f_1(-k) A^{\beta}_{\mathrm{r}}(k) \big(k^2 \eta_{\alpha\beta} - k_{\alpha} k_{\beta}\big) A^{\alpha}_{\mathrm{a}}(-k) + f_2(-k) \big[A^{\mathrm{r}}_{\gamma}(k) (k \cdot n)^2 A^{\gamma}_{\mathrm{a}}(-k)$$
$$+ A^{\beta}_{\mathrm{r}}(k) \big(k^2 n_{\alpha} n_{\beta} - (k \cdot n) \big[k_{\alpha} n_{\beta} + k_{\beta} n_{\alpha}\big]\big) A^{\alpha}_{\mathrm{a}}(-k) \big]$$
$$- 2 f_4(-k) \epsilon_{\alpha\beta\delta\sigma} A^{\beta}_{\mathrm{r}}(k) (k \cdot n) k^{\delta} n^{\sigma} A^{\alpha}_{\mathrm{a}}(-k) - i 2 f_5(-k) \epsilon_{\alpha\beta\gamma\delta} A^{\beta}_{\mathrm{r}}(k) k^{\gamma} n^{\delta} A^{\alpha}_{\mathrm{a}}(-k)$$
$$+ 2 f_6(-k) \epsilon_{\beta\gamma\alpha\delta} A^{\gamma}_{\mathrm{r}}(k) k^2 k^{\alpha} n^{\delta} A^{\beta}_{\mathrm{a}}(-k) + A^{\beta}_{\mathrm{r}}(k) [f_8(-k) k^2 + f_9(-k) (k \cdot n)^2] k_{\alpha} k_{\beta} A^{\alpha}_{\mathrm{a}}(-k) \Big\},$$

$$S_{\text{gravity}} = \frac{i}{2}\int \frac{\mathrm{d}^4k}{(2\pi)^4}\Big\{2g_1(-k)A_{\mathrm{a}}^{\beta}(k)\big(k^2\eta_{\alpha\beta}-k_\alpha k_\beta\big)A_{\mathrm{a}}^{\alpha}(-k)+g_2\big[(-k)A_{\gamma}^{\mathrm{a}}(k)(k\cdot n)^2A_{\mathrm{a}}^{\gamma}(-k)$$
$$+A_{\mathrm{a}}^{\beta}(k)\big(k^2n_\alpha n_\beta-(k\cdot n)\big[k_\alpha n_\beta+k_\beta n_\alpha\big]\big)A_{\mathrm{a}}^{\alpha}(-k)\big]$$
$$-2g_4(-k)\epsilon_{\alpha\beta\delta\sigma}A_{\mathrm{a}}^{\beta}(k)(k\cdot n)k^{\delta}n^{\sigma}A_{\mathrm{a}}^{\alpha}(-k)-i2g_5(-k)\epsilon_{\alpha\beta\gamma\delta}A_{\mathrm{a}}^{\beta}(k)k^{\gamma}n^{\delta}A_{\mathrm{a}}^{\alpha}(-k)$$
$$+2g_6(-k)\epsilon_{\beta\gamma\alpha\delta}A_{\mathrm{a}}^{\gamma}(k)k^2k^{\alpha}n^{\delta}A_{\mathrm{a}}^{\beta}(-k)+A_{\mathrm{a}}^{\beta}(k)[g_7(-k)k^2+g_8(-k)(k\cdot n)^2]k_\alpha k_\beta A_{\mathrm{a}}^{\alpha}(-k)\Big\}.$$

$$S_{\text{IF}} = q^2\int \frac{\mathrm{d}^4k}{(2\pi)^4}\left[-A_{\mathrm{a}}^{\alpha}(-k)\mathcal{D}_{\alpha\beta}(k)A_{\mathrm{r}}^{\beta}(k)+\frac{i}{2}A_{\mathrm{a}}^{\alpha}(-k)\mathcal{S}_{\alpha\beta}(k)A_{\mathrm{a}}^{\beta}(k)\right]$$

$$-q^2\mathcal{D}_{\alpha\beta}(k) = 2f_1(-k)\big(k^2\eta_{\alpha\beta}-k_\alpha k_\beta\big)+f_2(-k)\big((n\cdot k)^2\eta_{\alpha\beta}+k^2n_\alpha n_\beta-(k\cdot n)\big[k_\alpha n_\beta+k_\beta n_\alpha\big]\big)$$
$$q^2\mathcal{S}_{\alpha\beta}(k) = 2g_1(-k)\big(k^2\eta_{\alpha\beta}-k_\alpha k_\beta\big)+g_2(-k)\big((n\cdot k)^2\eta_{\alpha\beta}+k^2n_\alpha n_\beta-(k\cdot n)\big[k_\alpha n_\beta+k_\beta n_\alpha\big]\big)$$

$$k^{\mu}\mathcal{D}_{\mu\nu} = k^{\mu}\mathcal{S}_{\mu\nu} = 0$$

$$\mathcal{D}_{\alpha\beta} = \mathcal{D}_{\mathrm{L}}(k)\mathcal{P}_{\alpha\beta}^{\mathrm{L}}+\mathcal{D}_{\mathrm{T}}(k)\mathcal{P}_{\alpha\beta}^{\mathrm{T}}$$
$$\mathcal{S}_{\alpha\beta} = \mathcal{S}_{\mathrm{L}}(k)\mathcal{P}_{\alpha\beta}^{\mathrm{L}}+\mathcal{S}_{\mathrm{T}}(k)\mathcal{P}_{\alpha\beta}^{\mathrm{T}}$$

scalar kernels $\mathcal{D}_{L,T}(k)$ and $\mathcal{S}_{L,T}(k)$

$$-q^2\mathcal{D}_{\mathrm{L}}(k) = 2f_1(-k)-f_2(-k), \qquad q^2\mathcal{S}_{\mathrm{L}}(k) = 2g_1(-k)-g_2(-k),$$
$$-q^2\mathcal{D}_{\mathrm{T}}(k) = \frac{2k^2}{|\mathbf{k}|^2}f_1(-k)+\frac{k_0^2}{|\mathbf{k}|^2}f_2(-k), \qquad q^2\mathcal{S}_{\mathrm{T}}(k) = \frac{2k^2}{|\mathbf{k}|^2}g_1(-k)+\frac{k_0^2}{|\mathbf{k}|^2}g_2(-k)$$

$$f_n(x-y) = 0 \text{ unless } (x-y)^2 \le 0 \text{ and } x^0 > y^0.$$

$$S_{\text{in-in}} = \int \mathrm{d}^4x\left[-\frac{1}{2}F_{\alpha\beta}^{\mathrm{r}}F_{\mathrm{a}}^{\alpha\beta}+\theta F_{\mathrm{r}}^{\alpha\beta}\tilde{F}_{\mathrm{a}}^{\alpha\beta}-\mathcal{D}_{\mu}^{\mathrm{r}}\chi_{\mathrm{r}}(x)\mathcal{D}_{\mathrm{a}}^{\mu}\chi_{\mathrm{a}}(x)\right]+S_{\text{IF}}$$

$$\Delta S^{\text{supergravity}} = \int \mathrm{d}^{\square}x\int \mathrm{d}^{\square}y\{f_{10}(x-y)\mathcal{D}_0^{\mathrm{r}}\chi_{\mathrm{r}}(x)\mathcal{D}_0^{\mathrm{a}}\chi_{\mathrm{a}}(y)+f_{11}(x-y)\eta^{ij}\mathcal{D}_i^{\mathrm{r}}\chi_{\mathrm{r}}(x)\mathcal{D}_j^{\mathrm{a}}\chi_{\mathrm{a}}(y)$$
$$+h_1(x-y)F_{0i}^{\mathrm{r}}(x)\mathcal{D}_i^{\mathrm{a}}\chi_{\mathrm{a}}(y)+h_2(x-y)\mathcal{D}_i^{\mathrm{r}}\chi_{\mathrm{r}}(x)F_{0i}^{\mathrm{a}}(y)+h_3(x-y)\epsilon_{ijk}F_{ij}^{\mathrm{r}}(x)\mathcal{D}_k^{\mathrm{a}}\chi_{\mathrm{a}}(y)$$
$$+h_4(x-y)\epsilon_{ijk}\mathcal{D}_k^{\mathrm{r}}\chi_{\mathrm{r}}(x)F_{ij}^{\mathrm{a}}(y)\}$$

$$\Delta S^{\text{gravity}} = \int \mathrm{d}^4x\int \mathrm{d}^4y\{g_9(x-y)\mathcal{D}_0^{\mathrm{a}}\chi_{\mathrm{a}}(x)\mathcal{D}_0^{\mathrm{a}}\chi_{\mathrm{a}}(y)+g_{10}(x-y)\eta^{ij}\mathcal{D}_i^{\mathrm{a}}\chi_{\mathrm{a}}(x)\mathcal{D}_j^{\mathrm{a}}\chi_{\mathrm{a}}(y)$$
$$+c_1(x-y)F_{0i}^{\mathrm{a}}(x)\mathcal{D}_i^{\mathrm{a}}\chi_{\mathrm{a}}(y)+c_2(x-y)\epsilon_{ijk}F_{ij}^{\mathrm{a}}(x)\mathcal{D}_k^{\mathrm{a}}\chi_{\mathrm{a}}(y)\}$$

$$\mathcal{T}e^{-i\int \mathrm{d}^4x\hat{\mathcal{H}}_{\text{int}}(x)} \equiv \mathcal{T}^*e^{i\int \mathrm{d}^4x\hat{\mathcal{L}}_{\text{int}}(x)}$$

$$Z = \int \mathcal{D}[\phi_I]\int \mathcal{D}[\pi_I]e^{i\int \mathrm{d}^4x\sum_I \pi_I\partial_t\phi_I-\mathcal{H}[\phi_I,\pi_I]} = \int \mathcal{D}[\phi_I]e^{iS[\phi_I]}$$

$$\hat{\mathcal{H}}_{\text{int}}(x) \neq -\hat{\mathcal{L}}_{\text{int}}(x).$$

$$\int \mathcal{D}[\pi_I] e^{i\int \mathrm{d}^4x \sum_I \pi_I \partial_t \phi_I - \mathcal{H}[\phi_\jmath, \pi_\jmath]} = \mu[\phi_I] e^{iS[\phi_I]}$$

$$Z = \int \mathcal{D}[\phi] \sqrt{\det[G_A[\phi]]} e^{iS[\phi]}$$

$$\int \mathrm{d}^4 z \frac{\delta^2 S[\phi]}{\delta\phi(x)\delta\phi(z)} G_A[\phi](z,y) = i\delta^4(x,y)$$

$$\Gamma[\phi] = S[\phi] - \frac{i}{2}\mathrm{Trlog}\ G_F[\phi] + \frac{i}{2}\mathrm{Trlog}\ G_A[\phi]$$

$$G_A(x,y)G_A(y,z)G_A(z,x)$$

$\mathcal{T}^* e^{i\int \mathrm{d}^4x \hat{\mathcal{L}}_{\mathrm{int}}(x)}$ is the canonical path integral, computing matrix elements of $\mathcal{T} e^{-i\int \mathrm{d}^4x \hat{\mathcal{H}}_{\mathrm{int}}(x)}$ be

Matthews' theorem.

$$\Gamma[\phi] = S[\phi] - \frac{i}{2}\log\left(\frac{\det G_F[\phi]}{\det G_A[\phi]}\right)$$

$$\Gamma[\phi_+, \phi_-] = S[\phi_+] - S[\phi_-] - \frac{i}{2}\mathrm{Trlog}\ \mathbf{G}[\phi_+, \phi_-] + \frac{i}{2}\mathrm{Trlog}\ \mathbf{G}_A[\phi_+, \phi_-]$$

$$\det \mathbf{G}_A[\phi_+, \phi_-] = \det G_A[\phi_+] \det G_R[\phi_-]$$

$$\det \mathbf{G}[\phi_{\mathrm{r}}, \phi_{\mathrm{r}}] = \det G_A[\phi_{\mathrm{r}}] \det G_R[\phi_{\mathrm{r}}]$$

$$\Gamma[\phi_{\mathrm{r}}, \phi_{\mathrm{r}}] = 0$$

$$S_{\mathrm{IF}}[\phi_{\mathrm{r}}, \phi_{\mathrm{r}}] = 0$$

$$\tilde{\mu}[\phi_+, \phi_-] e^{iS_{\mathrm{IF}}[\phi_+, \phi_-]} = \int \mathcal{D}[H_+, H_-] \mu[\phi_+, \phi_-, H_+, H_-] e^{iS[\phi_+, \phi_-, H_+, H_-] + iS_{i\epsilon}}$$

$$\sum_{\beta=\pm} \int \mathrm{d}^4 z \begin{pmatrix} \dfrac{\delta^2 S[\phi]}{\delta\phi_\alpha(x)\delta\phi_\beta(z)} & \dfrac{\delta^2 S[\phi]}{\delta\phi_\alpha(x)\delta H_\beta(z)} \\ \dfrac{\delta^2 S[\phi]}{\delta H_\alpha(x)\delta\phi_\beta(z)} & \dfrac{\delta^2 S[\phi]}{\delta H_\alpha(x)\delta H_\beta(z)} \end{pmatrix} \mathbf{G}_{A\beta\gamma}[\phi_+, \phi_-, H_+, H_-](x,y) = i\delta^4(x,y)\delta_{\alpha\gamma} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mu[\phi_+, \phi_-, H_+, H_-] = \sqrt{\det \mathbf{G}_{A\beta\gamma}[\phi_+, \phi_-, H_+, H_-]}$$

$$S[\phi_+, \phi_-, H_+, H_-], H_\pm = 0, \mu[\phi_+, \phi_-, 0,0] \frac{\delta^2 S[\phi]}{\delta H_\alpha(x)\delta\phi_\beta(z)} = 0$$

$$\mu[\phi_+, \phi_-, 0,0] = \sqrt{\det \mathbf{G}_{\phi A}[\phi_+, \phi_-]} \sqrt{\det \mathbf{G}_{HA}[\phi_+, \phi_-]}$$

$$\tilde{\mu}[\phi_+, \phi_-] = \sqrt{\det \mathbf{G}_{\phi A}[\phi_+, \phi_-]}$$

$$S_{\mathrm{IF}}[\phi_+,\phi_-] = -\frac{i}{2}\mathrm{Trlog}\ \mathbf{G}_H[\phi_+,\phi_-] + \frac{i}{2}\mathrm{Trlog}\ \mathbf{G}_{HA}[\phi_+,\phi_-]$$

$$S_{\mathrm{IF}}[\phi_{\mathrm{r}},\phi_{\mathrm{r}}] = -\frac{i}{2}\mathrm{Trlog}\ \mathbf{G}_H[\phi_{\mathrm{r}},\phi_{\mathrm{r}}] + \frac{i}{2}\mathrm{Trlog}\ \mathbf{G}_{HA}[\phi_{\mathrm{r}},\phi_{\mathrm{r}}] = 0$$

$$\mathrm{Trlog}\,\mathbf{G}_{\phi A}[\phi_+,\phi_-] \text{ or } \mathrm{Trlog}\,\mathbf{G}_{HA}[\phi_+,\phi_-]$$

$$S[\Phi,\Phi^*,A] = \int_{t_{\mathrm{i}}}^{t_{\mathrm{f}}} \mathrm{d}t(|D_0[A]\Phi|^2 - \Gamma^2|\Phi|^2)$$

$$D_0[A] = \partial_t - iqA_0(t)$$

$$e^{iW[J_+,J_-,A_+,A_-]} = \int \mathrm{d}\left[\Phi_{\mathrm{f}},\Phi_{\mathrm{f}}^*,\Phi_{\pm\mathrm{i}},\Phi_{\pm\mathrm{i}}^*\right]\langle\Phi_+\Phi_+^*|\rho|\Phi_-\Phi_-^*\rangle \int_{\Phi_{+\mathrm{i}}\Phi_{+\mathrm{i}}^*}^{\Phi_{\mathrm{f}}\Phi_{\mathrm{f}}^*} \mathcal{D}[\Phi_+,\Phi_+^*] \int_{\Phi_{-\mathrm{i}}\Phi_{-\mathrm{i}}^*}^{\Phi_{\mathrm{f}}\Phi_{\mathrm{f}}^*} \mathcal{D}[\Phi_-,\Phi_-^*]$$

$$\times e^{iS[\Phi_+,\Phi_+^*,A_+]-iS[\Phi_-,\Phi_-^*,A_-]+i\int_{t_{\mathrm{i}}}^{t_{\mathrm{f}}}\mathrm{d}t(J_+^*(t)\Phi_+(t)+\Phi_+^*(t)J_+(t)-J_-^*(t)\Phi_-(t)-\Phi_-^*(t)J_-(t))}$$

$$\langle\Phi_+\Phi_+^*|\rho|\Phi_-\Phi_-^*\rangle = Ae^{-\kappa|\Phi_+|^2-\kappa|\Phi_-|^2}$$

$$\begin{aligned} S_{\text{in-in}} =& S[\Phi_+,\Phi_+^*,A_+] - S[\Phi_-,\Phi_-^*,A_-] + \int_{t_{\mathrm{i}}}^{t_{\mathrm{f}}} \mathrm{d}t(J_+^*\Phi_+(t) + \Phi_+^*J_+(t) - J_-^*\Phi_-(t) - \Phi_-^*J_-(t)) \\ &+\lambda^*\big(\Phi_+(t_{\mathrm{f}}) - \Phi_-(t_{\mathrm{f}})\big) + \big(\Phi_+^*(t_{\mathrm{f}}) - \Phi_-^*(t_{\mathrm{f}})\big)\lambda + i\kappa|\Phi_+(t_{\mathrm{i}})|^2 + i\kappa|\Phi_-(t_{\mathrm{i}})|^2 \end{aligned}$$

$$\begin{aligned} D_0[A_+]^2\Phi_+ + \Gamma^2\Phi_+ &= J_+(t) \\ D_0[A_-]^2\Phi_- + \Gamma^2\Phi_- &= J_-(t) \end{aligned}$$

$$\begin{aligned} D_0[A_+]\Phi_+(t_{\mathrm{i}}) &= i\kappa\Phi_+(t_{\mathrm{i}}), & D_0[A_-]\Phi_-(t_{\mathrm{i}}) &= -i\kappa\Phi_-(t_{\mathrm{i}}), \\ \Phi_+(t_{\mathrm{f}}) &= \Phi_-(t_{\mathrm{f}}), & \dot{\Phi}_+(t_{\mathrm{f}}) &= \dot{\Phi}_-(t_{\mathrm{f}}), \end{aligned}$$

$$A_0^+(t_{\mathrm{f}}) = A_0^-(t_{\mathrm{f}})$$

$$\Phi_\pm(t) = e^{iq\int_{t_{\mathrm{i}}}^{t}\mathrm{d}\tau A_0^\pm(\tau)}\sigma_\pm(t), \text{ and } J_\pm(t) = e^{iq\int_{t_{\mathrm{i}}}^{t}\mathrm{d}\tau A_0^\pm(\tau)}j_\pm(t)$$

$$\partial_t\sigma_+(t_{\mathrm{i}}) = i\kappa\sigma_+(t_{\mathrm{i}}), \partial_t\sigma_-(t_{\mathrm{i}}) = -i\kappa\sigma_-(t_{\mathrm{i}})$$

$$\begin{aligned} \partial_t^2\sigma_+ + \Gamma^2\sigma_+ &= j_+(t) \\ \partial_t^2\sigma_- + \Gamma^2\sigma_- &= j_-(t) \end{aligned}$$

$$\sigma_+(t_{\mathrm{f}}) = e^{-iq\int_{t_{\mathrm{i}}}^{t_{\mathrm{f}}}\mathrm{d}\tau A_0^{\mathrm{a}}(\tau)}\sigma_-(t_{\mathrm{f}}), \dot{\sigma}_+(t_{\mathrm{f}}) = e^{-iq\int_{t_{\mathrm{i}}}^{t_{\mathrm{f}}}\mathrm{d}\tau A_0^{\mathrm{a}}(\tau)}\dot{\sigma}_-(t_{\mathrm{f}})$$

$$\sigma_\pm(t) = e^{\mp\frac{i}{2}q\int_{t_{\mathrm{i}}}^{t_{\mathrm{f}}}\mathrm{d}\tau A_0^{\mathrm{a}}(\tau)}\tilde{\sigma}_\pm(t), j_\pm(t) = e^{\mp\frac{i}{2}q\int_{t_{\mathrm{i}}}^{t_{\mathrm{f}}}\mathrm{d}\tau A_0^{\mathrm{a}}(\tau)\tilde{j}_\pm(t)}$$

$$\begin{aligned} \partial_t\tilde{\sigma}_+(t_{\mathrm{i}}) &= i\kappa\tilde{\sigma}_+(t_{\mathrm{i}}), & \partial_t\tilde{\sigma}_-(t_{\mathrm{i}}) &= -i\kappa\tilde{\sigma}_-(t_{\mathrm{i}}), \\ \partial_t^2\tilde{\sigma}_+ + \Gamma^2\tilde{\sigma}_+ &= \tilde{j}_+(t), & \partial_t^2\tilde{\sigma}_- + \Gamma^2\tilde{\sigma}_- &= \tilde{j}_-(t), \\ \tilde{\sigma}_+(t_{\mathrm{f}}) &= \tilde{\sigma}_-(t_{\mathrm{f}}), & \dot{\tilde{\sigma}}_+(t_{\mathrm{f}}) &= \dot{\tilde{\sigma}}_-(t_{\mathrm{f}}). \end{aligned}$$

$$W[J_+,J_-,A_+,A_-] = i\int_{t_\mathrm{i}}^{t_\mathrm{f}} \mathrm{d}t \int_{t_\mathrm{i}}^{t_\mathrm{f}} \mathrm{d}t' (\tilde{j}_+^*(t) - \tilde{j}_-^*(t))\mathbf{D}(t,t') \begin{pmatrix} \tilde{J}_+(t') \\ -\tilde{J}_-(t') \end{pmatrix}$$

$$W[J_+,J_-,A_+,A_-] = i\int_{t_\mathrm{i}}^{t_\mathrm{f}} \mathrm{d}t \int_{t_\mathrm{i}}^{t_\mathrm{f}} \mathrm{d}t' (J_+^*(t) - J_-^*(t))V(t)\mathbf{U}(t)\mathbf{D}(t,t')\mathbf{U}^\dagger(t')V^*(t') \begin{pmatrix} J_+(t') \\ -J_-(t') \end{pmatrix}$$

$$V(t) = e^{\,i\int_{t_\mathrm{i}}^{t} \mathrm{d}t' A_0^\mathrm{r}(t')}$$

$$\mathbf{U}(t) = \begin{pmatrix} U_+(t) & 0 \\ 0 & U_-(t) \end{pmatrix}$$

$$U_\pm(t) = e^{\mp\frac{i}{2}\int_t^{t_\mathrm{f}} \mathrm{d}t' A_0^\mathrm{a}(t')}$$

$$e^{iS_\mathrm{IF}[A_+,A_-]} = \int \mathcal{D}[\Psi_+,\bar{\Psi}_+,\Psi_-,\bar{\Psi}_-]$$
$$e^{i\int \mathrm{d}^4x (i\bar{\Psi}_+ \partial\!\!\!/ \Psi_+ - m\bar{\Psi}_+\Psi_+ - i\bar{\Psi}_- \partial\!\!\!/ \Psi_- + m\bar{\Psi}_-\Psi_- + \bar{\Psi}_+ q\mathbb{A}\!\!\!/_+ \Psi_+ - \bar{\Psi}_- q\mathbb{A}\!\!\!/_- \Psi_-)}$$

$$\Psi_\pm(x) \to \Psi_\pm(x) + i\lambda_\pm(x)\Psi_\pm(x)$$

$$\int \mathcal{D}[\Psi_+,\bar{\Psi}_+,\Psi_-,\bar{\Psi}_-] \int \mathrm{d}^4x \partial_\mu \lambda_\pm(x)\Psi_\pm(x)\gamma^\mu\Psi_\pm(x) e^{i\int \mathrm{d}^4y \cdots} =$$
$$\int \mathcal{D}[\Psi_+,\bar{\Psi}_+,\Psi_-,\bar{\Psi}_-] \left(-\int \mathrm{d}^4x \lambda_\pm(x)\partial_\mu(\Psi_\pm(x)\gamma^\mu\Psi_\pm(x))\right) e^{i\int \mathrm{d}^4y \cdots} = 0$$

$$\partial_\mu \frac{\delta S_\mathrm{IF}[A_+,A_-]}{\delta A_\mu^\pm(x)} = 0$$

$$S_\mathrm{IF}[A_+,A_-] \simeq q^2 \int \mathrm{d}^4x \big(-A_{+\mu}(x)A_+^\mu(x) + A_{-\mu}(x)A_-^\mu(x)\big)\mathcal{F}(x,x)$$
$$+iq^2 \int \mathrm{d}^4x \int \mathrm{d}^4y \left(A_+^\mu(x)A_+^\nu(y)\left[\mathcal{F}(x,y)\frac{\partial^2\mathcal{F}(x,y)}{\partial x^\mu \partial y^\nu} - \frac{\partial\mathcal{F}(x,y)}{\partial x^\mu}\frac{\partial\mathcal{F}(x,y)}{\partial y^\mu}\right]\right.$$
$$-A_+^\mu(x)A_-^\nu(y)\left[\mathcal{W}^*(x,y)\frac{\partial^2\mathcal{W}^*(x,y)}{\partial x^\mu \partial y^\nu} - \frac{\partial\mathcal{W}^*(x,y)}{\partial x^\mu}\frac{\partial\mathcal{W}^*(x,y)}{\partial y^\mu}\right]$$
$$-A_-^\mu(x)A_+^\nu(y)\left[\mathcal{W}(x,y)\frac{\partial^2\mathcal{W}(x,y)}{\partial x^\mu \partial y^\nu} - \frac{\partial\mathcal{W}(x,y)}{\partial x^\mu}\frac{\partial\mathcal{W}(x,y)}{\partial y^\mu}\right]$$
$$\left.+A_-^\mu(x)A_-^\nu(y)\left[\mathcal{F}^*(x,y)\frac{\partial^2\mathcal{F}^*(x,y)}{\partial x^\mu \partial y^\nu} - \frac{\partial\mathcal{F}^*(x,y)}{\partial x^\mu}\frac{\partial\mathcal{F}^*(x,y)}{\partial y^\mu}\right]\right) + \mathcal{O}(q^4)$$

$$\Pi_{\mu\nu}^\beta \to \Pi_{\mu\nu} \text{ and } \mathcal{N}_{\mu\nu}^\beta \to \mathcal{N}_{\mu\nu}$$

$$\mathcal{F}^\mathrm{vac}(k) = \frac{-i}{k^2 + m^2 - i\epsilon} \text{ and } \mathcal{W}^\mathrm{vac}(k) = 2\pi\delta(k^2+m^2)\theta(k^0)$$

$$\Pi_{\mu\nu}(k) \equiv 2\eta_{\mu\nu}\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{F}^{\mathrm{vac}}(\ell) - 2i\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{F}^{\mathrm{vac}}(\ell)\mathcal{F}^{\mathrm{vac}}(\ell-k)\big(2\ell_\mu - k_\mu\big)\ell_\nu$$

$$= -2i\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\left[\frac{\eta_{\mu\nu}}{\ell^2+m^2-i\epsilon} - \frac{\big(2\ell_\mu - k_\mu\big)\ell_\nu}{[\ell^2+m^2-i\epsilon][(\ell-k)^2+m^2-i\epsilon]}\right]$$

$$= -2i\int_{-1}^{+1}\mathrm{d}y\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\frac{\eta_{\mu\nu}[\ell^2 - 2\ell\cdot k + k^2 + m^2] - \big(2\ell_\mu - k_\mu\big)\ell_\nu}{2\left(\ell^2 + (y-1)\ell\cdot k - \frac{1}{2}(y-1)k^2 + m^2 - i\epsilon\right)^2}$$

$$\frac{1}{AB} = 2\int_{-1}^{+1}\frac{\mathrm{d}y}{[(1+y)A + (1-y)B]^2}$$

$$\ell \to p = \ell + \frac{1}{2}(y-1)k$$

$$\Pi_{\mu\nu}(k) = -i\int_{-1}^{+1}\mathrm{d}y\int \frac{\mathrm{d}^4p}{(2\pi)^4}\frac{\eta_{\mu\nu}\left[p^2 + \frac{1}{4}(y^2+1)k^2 + m^2\right] - 2p_\mu p_\nu - \frac{1}{2}y^2k_\mu k_\nu}{\left(p^2 + \frac{1}{4}\Sigma - i\epsilon\right)^2}$$

$$\Sigma \equiv (1-y^2)k^2 + 4m^2$$

$$\int \frac{\mathrm{d}^Dp}{(2\pi)^D}\frac{p_\mu p_\nu}{\left(p^2 + \frac{1}{4}\Sigma - i\epsilon\right)^2} \to \frac{\eta_{\mu\nu}}{D}\int \frac{\mathrm{d}^Dp}{(2\pi)^D}\frac{p^2}{\left(p^2 + \frac{1}{4}\Sigma - i\epsilon\right)^2}$$

D dimensions:

$$\Pi_{\mu\nu}(k) = \int_{-1}^{+1}\mathrm{d}y\left(\left[\frac{1}{4}(y^2+1)\eta_{\mu\nu}k^2 + \eta_{\mu\nu}m^2 - \frac{1}{2}y^2k_\mu k_\nu\right]\mathcal{L}_{(0,2)}\left(\frac{1}{4}\Sigma\right) + \eta_{\mu\nu}\left(1 - \frac{2}{D}\right)\mathcal{L}_{(1,2)}\left(\frac{1}{4}\Sigma\right)\right)$$

$$\mathcal{L}_{(n,N)}(m^2) \equiv \mu^{4-D}\int \frac{\mathrm{d}^Dp}{(2\pi)^D}\frac{-i(p^2)^n}{(p^2+m^2-i\epsilon)^N}$$

$$p_0 \simeq \pm\sqrt{|\mathbf{p}|^2 + m^2} \mp i\epsilon$$

$$\mathcal{L}_{(n,N)}(m^2) = \mu^{4+2n-2N}\frac{\Gamma\left(\frac{D}{2}+n\right)\Gamma\left(-\frac{D}{2}-n+N\right)}{(4\pi)^{D/2}\Gamma\left(\frac{D}{2}\right)\Gamma(N)}\left(\frac{m^2-i\epsilon}{\mu^2}\right)^{\frac{D}{2}+n-N}$$

$\Sigma(y) = (1-y^2)k^2 + 4m^2$ factors out a $k^2\eta_{\mu\nu} - k_\mu k_\nu$ structure

$$\Pi_{\mu\nu}(k) = \left[k^2\eta_{\mu\nu} - k_\mu k_\nu\right]\int_{-1}^{+1}\mathrm{d}y\frac{y^2}{2}\cdot\frac{\Gamma\left(\frac{4-D}{2}\right)}{(4\pi)^{D/2}}\left(\frac{\mu^2}{\Sigma(y)/4 - i\epsilon}\right)^{\frac{4-D}{2}}$$

Expanding for $4 - D \ll 1$ gives:

$$\Pi_{\mu\nu}(k) \simeq \big(k^2\eta_{\mu\nu} - k_\mu k_\nu\big)\left[\frac{1}{24\pi^2(4-D)} - \frac{1}{32\pi^2}\int_{-1}^{+1}\mathrm{d}y y^2\log\left(\frac{(1-y^2)k^2 + 4m^2 - i\epsilon}{16\pi e^{-\gamma}\mu^2}\right)\right]$$

$$\mathcal{N}_{\mu\nu}(k) = 2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{W}^{\mathrm{vac}}(\ell)\mathcal{W}^{\mathrm{vac}}(-k-\ell)\big(2\ell_\mu + k_\mu\big)\ell_\nu$$

$$k^\mu\mathcal{N}_{\mu\nu}(k) = 0$$

$$\mathcal{N}_{\mu\nu}(k) = \big(k^2\eta_{\mu\nu} - k_\mu k_\nu\big)\mathcal{N}(k) \text{ with } \mathcal{N}(k) \equiv \frac{2}{3k^2}\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{W}^{\mathrm{vac}}(\ell)\mathcal{W}^{\mathrm{vac}}(-k-\ell)(2\ell^2 + k\cdot\ell)$$

$$\begin{aligned}\mathcal{N}(k) &= \frac{1}{6\pi^2k^2}\int \mathrm{d}^4\ell\theta(\ell_0)\theta(-k_0-\ell_0)\delta(\ell^2+m^2)\delta(\ell^2+2\ell\cdot k+k^2+m^2)(2\ell^2+\ell\cdot k)\\ &= \frac{\theta(-k^0)}{6\pi^2k^2}\int_0^{-k_0}\mathrm{d}\ell_0\int \mathrm{d}^3\ell\delta(\ell^2+m^2)\delta(\ell^2+2\ell\cdot k+k^2+m^2)(2\ell^2+\ell\cdot k)\\ &= \frac{\theta(-k^0)}{3\pi k^2}\int_0^{-k_0}\mathrm{d}\ell_0\int_m^\infty \mathrm{d}\Omega\int_{-1}^{+1}\mathrm{d}\mu\delta(-\ell_0^2+\Omega^2)\delta\left(-\ell_0^2+\Omega^2-2\ell_0k_0+2\mu\sqrt{\Omega^2-m^2}|\mathbf{k}|+k^2\right)\\ &\quad\times\Omega\sqrt{\Omega^2-m^2}(-2\ell_0^2+2\Omega^2-2m^2-\ell_0k_0+\mu L|\mathbf{k}|)\end{aligned}$$

$$|\boldsymbol{\ell}| \to \Omega = \sqrt{|\boldsymbol{\ell}|+m^2} \text{ and } \theta \to \mu = \cos\,\theta$$

$$\theta(-k^0) \to \theta(-k_0-m)$$

$$\mathcal{N}(k) = \frac{\theta(-k_0-m)}{12\pi k^2|\mathbf{k}|}\int_m^{-k_0}\mathrm{d}\Omega\int_{-1}^{+1}\mathrm{d}\mu\left(-2m^2-\Omega k_0+\mu\sqrt{\Omega^2-m^2}|\mathbf{k}|\right)\delta\left(\mu-\frac{2\Omega k_0-k^2}{2\sqrt{\Omega^2-m^2}|\mathbf{k}|}\right)$$

$$-1 < \frac{2\Omega k_0-k^2}{2\sqrt{\Omega^2-m^2}|\mathbf{k}|} < 1$$

$$|\mathbf{k}| > 0 \text{ and } -k_0 > \Omega > m > 0$$

$$0 < 4m^2 < -k^2 \text{ and } -\frac{k_0}{2}-\frac{|\mathbf{k}|}{2}\sqrt{1+\frac{4m^2}{k^2}} < \Omega < -\frac{k_0}{2}+\frac{|\mathbf{k}|}{2}\sqrt{1+\frac{4m^2}{k^2}}$$

$$\begin{aligned}\mathcal{N}(k) &= \frac{\theta(-k^0)\theta(-k^2-4m^2)}{12\pi k^2|\mathbf{k}|}\int_{-\frac{k_0}{2}-\frac{|\mathbf{k}|}{2}\sqrt{1+\frac{4m^2}{k^2}}}^{-\frac{k_0}{2}+\frac{|\mathbf{k}|}{1+\frac{4m^2}{k^2}}}\mathrm{d}\Omega\frac{-k^2-4m^2}{2}\\ &= -\frac{\theta(-k^0)\theta(-k^2-4m^2)}{24\pi}\left(1+\frac{4m^2}{k^2}\right)^{3/2}\end{aligned}$$

$$\mp\frac{q^2}{2}\int \frac{\mathrm{d}^4k}{(2\pi)^4}A_\pm^\mu(k)\Pi_{\mu\nu}(k)A_\pm^\nu(-k) \supset \mp\frac{q^2}{2}\int \frac{\mathrm{d}^4k}{(2\pi)^4}A_\pm^\mu(k)\frac{k^2\eta_{\mu\nu}-k_\mu k_\nu}{24\pi^2(4-D)}A_\pm^\nu(-k)$$

$$-\frac{q^2c}{4}\int \mathrm{d}^4xF_{\mu\nu}F^{\mu\nu} = -\frac{q^2c}{2}\int \mathrm{d}^4xA^\mu(k)\big(k^2\eta_{\mu\nu}-k_\mu k_\nu\big)A^\nu(-k)$$

$$c = -\frac{1}{24\pi^2(4-D)}+\frac{1}{48\pi^2}\log\left(\frac{m^2}{4\pi e^{-\gamma}\mu^2}\right)$$

$$\lim_{\beta\to\infty} S_{\mathrm{IF}}[A_+,A_-] \simeq \frac{q^2}{2}\int \frac{\mathrm{d}^4k}{(2\pi)^4}\big(k^2\eta_{\mu\nu}-k_\mu k_\nu\big)[\,-A_+^\mu(k)\Pi(k)A_+^\nu(-k)-iA_+^\mu(k)\mathcal{N}(-k)A_-^\nu(-k)$$
$$-iA_-^\mu(k)\mathcal{N}(k)A_+^\nu(-k)+A_-^\mu(k)\Pi^*(k)A_-^\nu(-k)]+\mathcal{O}(q^4)$$

$$\Pi(k)=\frac{1}{32\pi^2}\int_{-1}^{+1}\mathrm{d}y\,y^2\log\left(\frac{4m^2}{(1-y^2)k^2+4m^2-i\epsilon}\right)$$

$$\log\left(\tfrac{1}{z-i\epsilon}\right)=\log\left|\tfrac{1}{z}\right|+i\pi\theta(-z)\text{ for real }z\xrightarrow{yields}\text{real and imaginary parts}$$

$$\mathrm{Re}[\Pi(k)]\;=\frac{1}{32\pi^2}\int_{-1}^{+1}\mathrm{d}y\,y^2\log\left|\frac{4m^2}{(1-y^2)k^2+4m^2}\right|$$
$$\mathrm{Im}[\Pi(k)]\;=\frac{\theta(-k^2-4m^2)}{48\pi}\left(1+\frac{4m^2}{k^2}\right)^{3/2}$$

$$2\mathrm{Im}[\Pi(k^2)]+\mathcal{N}(k)+\mathcal{N}(-k)=0$$

$$\mathcal{F}^\beta(k)=\mathcal{F}^{\mathrm{vac}}(k)+\mathcal{C}^\beta(k)\;\text{ and }\;\mathcal{W}^\beta(k)=\mathcal{W}^{\mathrm{vac}}(k)+\mathcal{C}^\beta(k)$$

$$\mathcal{C}^\beta(k)\equiv\frac{2\pi\delta(k^2+m^2)}{e^{\beta|k_0|}-1}$$

$$\Pi_{\mu\nu}^\beta(k)=\Pi_{\mu\nu}(k)+\bar{\Pi}_{\mu\nu}^\beta(k)\;\text{ and }\;\mathcal{N}_{\mu\nu}^\beta(k)=\mathcal{N}_{\mu\nu}+\overline{\mathcal{N}}_{\mu\nu}^\beta(k)$$

$$\bar{\Pi}_{\mu\nu}^\beta(k)=\eta_{\mu\nu}\mathcal{T}^\beta-2i\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}\big(\mathcal{C}^\beta(\ell)\mathcal{F}^{\mathrm{vac}}(\ell-k)\big(2\ell_\mu-k_\mu\big)(2\ell_\nu-k_\nu)+\mathcal{C}^\beta(\ell)\mathcal{C}^\beta(\ell-k)\big(2\ell_\mu-k_\mu\big)\ell_\nu\big)$$
$$\overline{\mathcal{N}}_{\mu\nu}^\beta(k)=2\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}\big(\mathcal{C}^\beta(\ell)\mathcal{W}^{\mathrm{vac}}(\ell-k)\big(2\ell_\mu-k_\mu\big)(2\ell_\nu-k_\nu)+\mathcal{C}^\beta(\ell)\mathcal{C}^\beta(\ell-k)\big(2\ell_\mu-k_\mu\big)\ell_\nu\big)$$

$$\mathcal{T}^\beta:=2\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{C}^\beta(\ell)$$

$$\eta^{\mu\nu}\bar{\Pi}_{\mu\nu}^\beta(k)\;=4\mathcal{T}^\beta-\mathcal{E}^{\beta\mathcal{F}}(k)+i\mathcal{E}^{\beta\beta}(k)+\mathcal{K}^{\beta\mathcal{F}}(k)-i\mathcal{K}^{\beta\beta}(k)$$
$$n^\mu n^\nu\bar{\Pi}_{\mu\nu}^\beta(k)\;=-\mathcal{T}^\beta+\mathcal{E}^{\beta\mathcal{F}}(k)-i\mathcal{E}^{\beta\beta}(k)$$
$$\eta^{\mu\nu}\overline{\mathcal{N}}_{\mu\nu}^\beta(k)\;=-\mathcal{E}^{\beta\mathcal{W}}(k)-\mathcal{E}^{\beta\beta}(k)+\mathcal{K}^{\beta\mathcal{W}}(k)+\mathcal{K}^{\beta\beta}(k)$$
$$n^\mu n^\nu\overline{\mathcal{N}}_{\mu\nu}^\beta(k)\;=\mathcal{E}^{\beta\mathcal{W}}(k)+\mathcal{E}^{\beta\beta}(k)$$

$$\mathcal{K}^{\beta\mathcal{F}}(k) \equiv -2i\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{C}^{\beta}(\ell)\mathcal{F}^{\mathrm{vac}}(\ell-k)|2\ell-\mathbf{k}|^2$$

$$\mathcal{E}^{\beta\mathcal{F}}(k) \equiv -2i\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{C}^{\beta}(\ell)\mathcal{F}^{\mathrm{vac}}(\ell-k)(2\ell_0-k_0)^2$$

$$\mathcal{K}^{\beta\mathcal{W}}(k) \equiv 2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{C}^{\beta}(\ell)\mathcal{W}^{\mathrm{vac}}(\ell-k)|2\ell-\mathbf{k}|^2$$

$$\mathcal{E}^{\beta\mathcal{W}}(k) \equiv 2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{C}^{\beta}(\ell)\mathcal{W}^{\mathrm{vac}}(\ell-k)(2\ell_0-k_0)^2$$

$$\mathcal{K}^{\beta\beta}(k) \equiv 2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{C}^{\beta}(\ell)\mathcal{C}^{\beta}(\ell-k)(2|\ell|^2-\mathbf{k}\cdot\ell)$$

$$\mathcal{E}^{\beta\beta}(k) \equiv 2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\mathcal{C}^{\beta}(\ell)\mathcal{C}^{\beta}(\ell-k)(2\ell_0^2-k_0\ell_0)$$

$$\mathcal{T}^{\beta} = 2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\frac{2\pi\delta(\ell^2+m^2)}{e^{\beta|\ell_0|}-1} = \frac{1}{\pi^2}\int_m^{\infty}\mathrm{d}\Omega\frac{\sqrt{\Omega^2-m^2}}{e^{\beta\Omega}-1}$$

energy variable $\Omega = \sqrt{|\boldsymbol{\ell}|^2+m^2}$

$$\mathcal{T}^{\beta} \simeq \frac{1}{6\beta^2}\ (m\beta \ll 1)$$

$$\mathcal{K}^{\beta\mathcal{F}}(k) = -2i\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\frac{2\pi\delta(\ell^2+m^2)}{e^{\beta|\ell_0|}-1}\cdot\frac{-i}{(\ell-k)^2+m^2-i\epsilon}\cdot(|\mathbf{k}|^2-4\mathbf{k}\cdot\ell+4|\ell|^2)$$

$$= \frac{1}{8\pi^2|\mathbf{k}|}\int_m^{\infty}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}\int_{-1}^{+1}\mathrm{d}\mu\left[\frac{|\mathbf{k}|^2-4|\mathbf{k}|\sqrt{\Omega^2-m^2}\mu+4\Omega^2-4m^2}{\mu-\frac{k^2+2k_0\Omega}{2|\mathbf{k}|\sqrt{\Omega^2-m^2}}+i\epsilon}+\frac{|\mathbf{k}|^2-4|\mathbf{k}|\sqrt{\Omega^2-m^2}\mu+4\Omega^2-4m^2}{\mu-\frac{k^2-2k_0\Omega}{2|\mathbf{k}|\sqrt{\Omega^2-m^2}}+i\epsilon}\right]$$

spherical coordinates $(|\boldsymbol{\ell}|,\theta,\varphi)$, integrated over φ and swapped $|\boldsymbol{\ell}| \to \Omega = \sqrt{|\boldsymbol{\ell}|^2+m^2}$ and $\theta \to \mu =$

$$\cos\,\theta$$

simplify this by using the identity $\frac{a-b\mu}{\mu-x}+\frac{a-b\mu}{\mu-y} = -2b+\frac{a-bx}{\mu-x}+\frac{a-by}{\mu-y}$

$$\mathcal{K}^{\beta\mathcal{F}}(k) = \frac{1}{8\pi^2|\mathbf{k}|}\int_m^{\infty}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}\Big[-8|\mathbf{k}|\sqrt{\Omega^2-m^2}+(2k_0^2-|\mathbf{k}|^2-4k_0\Omega+4\Omega^2-4m^2)g\left(\frac{k^2+2k_0\Omega}{2|\mathbf{k}|\sqrt{\Omega^2-m^2}}\right)$$
$$+(2k_0^2-|\mathbf{k}|^2+4k_0\Omega+4\Omega^2-4m^2)g\left(\frac{k^2-2k_0\Omega}{2|\mathbf{k}|\sqrt{\Omega^2-m^2}}\right)\Big].$$

$$g(x) \equiv \int_{-1}^{+1}\frac{\mathrm{d}\mu}{\mu-x+i\epsilon} = \log\left|\frac{x-1}{x+1}\right| - i\pi\theta(1-x^2)\ \text{ for } x\in\mathbb{R}$$

$$\mathrm{Re}\left[\mathcal{K}^{\beta\mathcal{F}}(k)\right] = -\mathcal{T}^{\beta}+\frac{1}{8\pi^2|\mathbf{k}|}\int_m^{\infty}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}(2k_0^2-|\mathbf{k}|^2+4\Omega^2-4m^2)\log\left|\frac{\left(k^2-2|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2-4k_0^2\Omega^2}{\left(k^2+2|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2-4k_0^2\Omega^2}\right|$$
$$+\frac{k_0}{2\pi^2|\mathbf{k}|}\int_m^{\infty}\mathrm{d}\Omega\frac{\Omega}{e^{\beta\Omega}-1}\log\left|\frac{(k^2)^2-4\left(k_0\Omega+|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2}{(k^2)^2-4\left(k_0\Omega-|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2}\right|$$

$$\mathrm{Im}\left[\mathcal{K}^{\beta\mathcal{F}}(k)\right]=-\frac{1}{8\pi|\mathbf{k}|}\int_m^\infty\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}\left[(2k_0^2-|\mathbf{k}|^2-4k_0\Omega+4\Omega^2-4m^2)\theta\left(1-\frac{(k^2+2k_0\Omega)^2}{4|\mathbf{k}|^2(\Omega^2-m^2)}\right)\right.$$
$$\left.+(2k_0^2-|\mathbf{k}|^2+4k_0\Omega+4\Omega^2-4m^2)\theta\left(1-\frac{(k^2-2k_0\Omega)^2}{4|\mathbf{k}|^2(\Omega^2-m^2)}\right)\right]$$

$$\mathrm{Im}\left[\mathcal{K}^{\beta\mathcal{F}}(k)\right]=-\frac{\theta(-k^2-4m^2)}{8\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+}\frac{\mathrm{d}\Omega(4\Omega^2-4m^2+2k_0^2-|\mathbf{k}|^2-4|k_0|\Omega)}{e^{\beta\Omega}-1}$$
$$-\frac{\theta(k^2)}{8\pi|\mathbf{k}|}\left[\int_{\Omega_+}^\infty\frac{\mathrm{d}\Omega(4\Omega^2-4m^2+2k_0^2-|\mathbf{k}|^2-4|k_0|\Omega)}{e^{\beta\Omega}-1}+\int_{-\Omega_-}^\infty\frac{\mathrm{d}\Omega(4\Omega^2-4m^2+2k_0^2-|\mathbf{k}|^2+4|k_0|\Omega)}{e^{\beta\Omega}-1}\right]$$

$$-\Omega_-=|\Omega_-|,-\Omega_-+|k_0|=\Omega_+,\Omega\to\Omega+|k_0|,\Omega>\Omega_+,\theta(k^2)\text{ terms}$$

$$\mathrm{Im}\left[\mathcal{K}^{\beta\mathcal{F}}(k)\right]=-\frac{\theta(-k^2-4m^2)}{8\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+}\frac{\mathrm{d}\Omega(4\Omega^2-4m^2+2k_0^2-|\mathbf{k}|^2-4|k_0|\Omega)}{e^{\beta\Omega}-1}$$
$$-\frac{\theta(k^2)}{8\pi|\mathbf{k}|}\int_{\Omega_+}^\infty\mathrm{d}\Omega\left[\frac{1}{e^{\beta\Omega}-1}+\frac{1}{e^{\beta(\Omega-|k_0|)}-1}\right](4\Omega^2-4m^2+2k_0^2-|\mathbf{k}|^2-4|k_0|\Omega)$$

$$\mathrm{Re}\left[\mathcal{E}^{\beta\mathcal{F}}(k)\right]=\frac{1}{8\pi^2|\mathbf{k}|}\int_m^\infty\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}(k_0^2+4\Omega^2)\log\left|\frac{\left(k^2-2|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2-4k_0^2\Omega^2}{\left(k^2+2|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2-4k_0^2\Omega^2}\right|$$
$$+\frac{k_0}{2\pi^2|\mathbf{k}|}\int_m^\infty\mathrm{d}\Omega\frac{\Omega}{e^{\beta\Omega}-1}\log\left|\frac{(k^2)^2-4\left(k_0\Omega+|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2}{(k^2)^2-4\left(k_0\Omega-|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2}\right|$$
$$\mathrm{Im}\left[\mathcal{E}^{\beta\mathcal{F}}(k)\right]=-\frac{\theta(-k^2-4m^2)}{8\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}(k_0^2-4|k_0|\Omega+4\Omega^2)$$
$$-\frac{\theta(k^2)}{8\pi|\mathbf{k}|}\int_{\Omega_+}^\infty\mathrm{d}\Omega\left[\frac{1}{e^{\beta\Omega}-1}+\frac{1}{e^{\beta(\Omega-|k_0|)}-1}\right](k_0^2-4|k_0|\Omega+4\Omega^2)$$

$$\mathcal{L}_f(k)=2\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}2\pi\delta(\ell^2+m^2)\cdot2\pi\delta((k-\ell)^2+m^2)\cdot f(\ell_0,|\ell|,\mathbf{k}\cdot\ell,k_0)$$

using spherical coordinates $(|\ell|,\theta,\varphi)$, integrating over φ and swapping $|\boldsymbol{\ell}|\to\Omega=\sqrt{|\boldsymbol{\ell}|^2+m^2}$ and

$$\theta\to\mu=\cos\,\theta$$

$$\mathcal{L}_f(k)=\frac{1}{4\pi|\mathbf{k}|}\int_m^\infty\mathrm{d}\Omega\int_{-1}^{+1}\mathrm{d}\mu\delta\left(\mu-\frac{k^2+2k_0\Omega}{2|\mathbf{k}|\sqrt{\Omega^2-m^2}}\right)f\left(\Omega,\sqrt{\Omega^2-m^2},|\mathbf{k}|\sqrt{\Omega^2-m^2}\mu,k_0\right)$$
$$+\frac{1}{4\pi|\mathbf{k}|}\int_m^\infty\mathrm{d}\Omega\int_{-1}^{+1}\mathrm{d}\mu\delta\left(\mu-\frac{k^2-2k_0\Omega}{2|\mathbf{k}|\sqrt{\Omega^2-m^2}}\right)f\left(-\Omega,\sqrt{\Omega^2-m^2},|\mathbf{k}|\sqrt{\Omega^2-m^2}\mu,k_0\right)$$

$$-1<\frac{k^2\pm2k_0\Omega}{2|\mathbf{k}|\sqrt{\Omega^2-m^2}}<+1$$

$$\mathcal{L}_f(k) = \frac{\theta(-k^2-4m^2)}{4\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+} \mathrm{d}\Omega\left[\theta(k^0) f\left(\Omega,\sqrt{\Omega^2-m^2},\frac{k^2+2|k_0|\Omega}{2},|k_0|\right)+\theta(-k^0) f\left(-\Omega,\sqrt{\Omega^2-m^2},\frac{k^2+2|k_0|\Omega}{2},-|k_0|\right)\right]$$
$$+\frac{\theta(k^2)}{4\pi|\mathbf{k}|}\int_{\Omega_+}^{\infty} \mathrm{d}\Omega\left[\theta(k^0) f\left(\Omega,\sqrt{\Omega^2-m^2},\frac{k^2+2|k_0|\Omega}{2},|k_0|\right)+\theta(-k^0) f\left(-\Omega,\sqrt{\Omega^2-m^2},\frac{k^2+2|k_0|\Omega}{2},-|k_0|\right)\right]$$
$$+\frac{\theta(k^2)}{4\pi|\mathbf{k}|}\int_{-\Omega_-}^{\infty} \mathrm{d}\Omega\left[\theta(k^0) f\left(-\Omega,\sqrt{\Omega^2-m^2},\frac{k^2-2|k_0|\Omega}{2},|k_0|\right)+\theta(-k^0) f\left(\Omega,\sqrt{\Omega^2-m^2},\frac{k^2-2|k_0|\Omega}{2},-|k_0|\right)\right]$$

$$\mathcal{K}^{\beta\beta}(k) = 2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\frac{2\pi\delta(\ell^2+m^2)}{e^{\beta|\ell_0|}-1}\frac{2\pi\delta((k-\ell)^2+m^2)}{e^{\beta|k_0-\ell_0|}-1}(2|\ell|^2-\mathbf{k}\cdot\ell)$$

$$f(\ell_0,|\ell|,\mathbf{k}\cdot\ell,k_0)\to\frac{1}{e^{\beta|\ell_0|}-1}\cdot\frac{2|\ell|^2-\mathbf{k}\cdot\ell}{e^{\beta|k_0-\ell_0|}-1}$$

$$-i\mathcal{K}^{\beta\beta}(k) = -\frac{i\theta(-k^2-4m^2)}{8\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}\cdot\frac{4\Omega^2-4m^2+k_0^2-|\mathbf{k}|^2-2|k_0|\Omega}{e^{\beta(|k_0|-\Omega)}-1}$$
$$-\frac{i\theta(k^2)}{8\pi|\mathbf{k}|}\int_{\Omega_+}^{\infty}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}\cdot\frac{2(4\Omega^2-4m^2+2k_0^2-|\mathbf{k}|^2-4|k_0|\Omega)}{e^{\beta(\Omega-|k_0|)}-1}$$

$$\mathcal{E}^{\beta\beta}(k) = \frac{\theta(-k^2-4m^2)}{8\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}\cdot\frac{4\Omega^2-2|k_0|\Omega}{e^{\beta(|k_0|-\Omega)}-1}+\frac{\theta(k^2)}{16\pi|\mathbf{k}|}\int_{\Omega_+}^{\infty}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}\cdot\frac{2(2\Omega-|k_0|)^2}{e^{\beta(\Omega-|k_0|)}-1}$$

$$\mathcal{K}^{\beta\mathcal{W}}(k) = \frac{\theta(-k^2-4m^2)}{4\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}\theta(-k^0)[(2\Omega-|k_0|)^2-k^2-4m^2]$$
$$+\frac{\theta(k^2)}{4\pi|\mathbf{k}|}\int_{\Omega_+}^{\infty}\mathrm{d}\Omega\left[\frac{\theta(k^0)}{e^{\beta\Omega}-1}+\frac{\theta(-k^0)}{e^{\beta(\Omega-|k_0|)}-1}\right][(2\Omega-|k_0|)^2-k^2-4m^2]$$

$$\mathcal{E}^{\beta\mathcal{W}}(k) = \frac{\theta(-k^2-4m^2)}{4\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}\theta(-k^0)(2\Omega-|k_0|)^2$$
$$+\frac{\theta(k^2)}{4\pi|\mathbf{k}|}\left[\int_{\Omega_+}^{\infty}\mathrm{d}\Omega\left[\frac{\theta(k^0)}{e^{\beta\Omega}-1}+\frac{\theta(-k^0)}{e^{\beta(\Omega-|k_0|)}-1}\right](2\Omega-|k_0|)^2\right.$$

$$\Pi^{\beta}_{\mu\nu}(k)+c\left(k^2\eta_{\mu\nu}-k_\mu k_\nu\right) = \left(k^2\eta_{\mu\nu}-k_\mu k_\nu\right)\Pi(k)+\bar{\Pi}^{\beta}_{\mu\nu}(k)$$
$$\mathcal{N}^{\beta}_{\mu\nu}(k) = \left(k^2\eta_{\mu\nu}-k_\mu k_\nu\right)\mathcal{N}(k)+\overline{\mathcal{N}}^{\beta}_{\mu\nu}(k)$$

$$\left(k^2\eta_{\mu\nu}-k_\mu k_\nu\right)\Pi(k)+\bar{\Pi}^{\beta}_{\mu\nu}(k) = \Pi^{\beta}_{\mathrm{L}}(k)\mathcal{P}^{\mathrm{L}}_{\mu\nu}(k)+\Pi^{\beta}_{\mathrm{T}}(k)\mathcal{P}^{\mathrm{T}}_{\mu\nu}(k)$$
$$\left(k^2\eta_{\mu\nu}-k_\mu k_\nu\right)\mathcal{N}(k)+\overline{\mathcal{N}}^{\beta}_{\mu\nu}(k) = \mathcal{N}^{\beta}_{\mathrm{L}}(k)\mathcal{P}^{\mathrm{L}}_{\mu\nu}(k)+\mathcal{N}^{\beta}_{\mathrm{T}}(k)\mathcal{P}^{\mathrm{T}}_{\mu\nu}(k)$$

$$\eta^{\mu\nu}\mathcal{P}^{\mathrm{L}}_{\mu\nu}(k) = k^2\eta^{\mu\nu}\mathcal{P}^{\mathrm{T}}_{\mu\nu}(k) = 2|\mathbf{k}|^2$$
$$n_\mu n_\nu\mathcal{P}^{\mathrm{L}}_{\mu\nu}(k) = -|\mathbf{k}|^2 n_\mu n_\nu\mathcal{P}^{\mathrm{T}}_{\mu\nu}(k) = 0$$

$$\Pi^{\beta}_{\mathrm{L}}(k) = \Pi(k)+\frac{1}{|\mathbf{k}|^2}\left(\mathcal{T}^{\beta}-\mathcal{E}^{\beta\mathcal{F}}(k)+i\mathcal{E}^{\beta\beta}(k)\right)$$
$$\Pi^{\beta}_{\mathrm{T}}(k) = \frac{k^2}{|\mathbf{k}|^2}\Pi(k)+\frac{1}{2|\mathbf{k}|^4}\left((k_0^2+3|\mathbf{k}|^2)\mathcal{T}^{\beta}-k_0^2\left[\mathcal{E}^{\beta\mathcal{F}}(k)-i\mathcal{E}^{\beta\beta}(k)\right]+|\mathbf{k}|^2\left[\mathcal{K}^{\beta\mathcal{F}}-i\mathcal{K}^{\beta\beta}\right]\right)$$

$$\mathcal{N}^{\beta}_{\mathrm{L}}(k) = \mathcal{N}(k)-\frac{1}{|\mathbf{k}|^2}\left(\mathcal{E}^{\beta\mathcal{W}}(k)+\mathcal{E}^{\beta\beta}(k)\right)$$
$$\mathcal{N}^{\beta}_{\mathrm{T}}(k) = \frac{k^2}{|\mathbf{k}|^2}\mathcal{N}(k)+\frac{1}{2|\mathbf{k}|^4}\left(-k_0^2\left[\mathcal{E}^{\beta\mathcal{W}}(k)+\mathcal{E}^{\beta\beta}(k)\right]+|\mathbf{k}|^2\left[\mathcal{K}^{\beta\mathcal{W}}+\mathcal{K}^{\beta\beta}\right]\right)$$

$$\mathrm{Re}\left[\Pi_{\mathrm{L}}^{\beta}(k)\right]=\frac{1}{32\pi^2}\int_{-1}^{+1}\mathrm{d}yy^2\log\left|\frac{4m^2}{(1-y^2)k^2+4m^2}\right|+\frac{1}{\pi^2|\mathbf{k}|^2}\int_m^{\infty}\mathrm{d}\Omega\frac{\sqrt{\Omega^2-m^2}}{e^{\beta\Omega}-1}$$
$$-\frac{1}{8\pi^2|\mathbf{k}|^3}\int_m^{\infty}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}(k_0^2+4\Omega^2)\log\left|\frac{\left(k^2-2|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2-4k_0^2\Omega^2}{\left(k^2+2|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2-4k_0^2\Omega^2}\right|$$
$$-\frac{k_0}{2\pi^2|\mathbf{k}|^3}\int_m^{\infty}\mathrm{d}\Omega\frac{\Omega}{e^{\beta\Omega}-1}\log\left|\frac{(k^2)^2-4\left(k_0\Omega+|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2}{(k^2)^2-4\left(k_0\Omega-|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2}\right|$$
$$\mathrm{Im}\left[\Pi_{\mathrm{L}}^{\beta}(k)\right]=\frac{\theta(-k^2-4m^2)}{8\pi|\mathbf{k}|^3}\int_{\Omega_-}^{\Omega_+}\mathrm{d}\Omega\left(\frac{1}{2}\coth\left(\frac{\beta\Omega}{2}\right)(2\Omega-|k_0|)^2+\frac{4\Omega^2-2|k_0|\Omega}{[e^{\beta\Omega}-1][e^{\beta(|k_0|-\Omega)}-1]}\right)$$
$$+\frac{\theta(k^2)}{8\pi|\mathbf{k}|^3}\int_{\Omega_+}^{\infty}\mathrm{d}\Omega\frac{1+e^{\beta|k_0|}}{[e^{\beta\Omega}-1][1-e^{-\beta(\Omega-|k_0|)}]}(2\Omega-|k_0|)^2$$
$$\mathrm{Re}\left[\Pi_{\mathrm{T}}^{\beta}(k)\right]=\frac{k^2}{32\pi^2|\mathbf{k}|^2}\int_{-1}^{+1}\mathrm{d}yy^2\log\left|\frac{4m^2}{(1-y^2)k^2+4m^2}\right|+\frac{k_0^2+2|\mathbf{k}|^2}{2\pi^2|\mathbf{k}|^4}\int_m^{\infty}\mathrm{d}\Omega\frac{\sqrt{\Omega^2-m^2}}{e^{\beta\Omega}-1}$$
$$+\frac{1}{16\pi^2|\mathbf{k}|^5}\int_m^{\infty}\frac{\mathrm{d}\Omega}{e^{\beta\Omega}-1}[k^2(4\Omega^2-k^2)-4|\mathbf{k}|^2m^2]\log\left|\frac{\left(k^2-2|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2-4k_0^2\Omega^2}{\left(k^2+2|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2-4k_0^2\Omega^2}\right|$$
$$+\frac{k^2\cdot k_0}{4\pi^2|\mathbf{k}|^5}\int_m^{\infty}\mathrm{d}\Omega\frac{\Omega}{e^{\beta\Omega}-1}\log\left|\frac{(k^2)^2-4\left(k_0\Omega+|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2}{(k^2)^2-4\left(k_0\Omega-|\mathbf{k}|\sqrt{\Omega^2-m^2}\right)^2}\right|$$
$$\mathrm{Im}\left[\Pi_{\mathrm{T}}^{\beta}(k)\right]=-\frac{\theta(-k^2-4m^2)}{16\pi|\mathbf{k}|^5}\int_{\Omega_-}^{\Omega_+}\mathrm{d}\Omega\left(\frac{1}{2}\coth\left(\frac{\beta\Omega}{2}\right)(k^2(2\Omega-|k_0|)^2-(k^2+4m^2)|\mathbf{k}|^2)+\frac{k^2(4\Omega^2-2|k_0|\Omega)-(k^2+4m^2)|\mathbf{k}|^2}{[e^{\beta\Omega}-1][e^{\beta(|k0|-\Omega)}-1]}\right)$$
$$-\frac{\theta(k^2)}{16\pi|\mathbf{k}|^5}\int_{\Omega_+}^{\infty}\mathrm{d}\Omega\frac{1+e^{\beta|k_0|}}{[e^{\beta\Omega}-1][1-e^{-\beta(\Omega-|k_0|)}]}(k^2(2\Omega-|k_0|)^2-(k^2+4m^2)|\mathbf{k}|^2)$$

$$\mathcal{N}_{\mathrm{L}}^{\beta}(k)=-\frac{\theta(-k^2-4m^2)}{8\pi|\mathbf{k}|^3}\int_{\Omega_-}^{\Omega_+}\mathrm{d}\Omega\left(\theta(-k^0)\coth\left(\frac{\beta\Omega}{2}\right)(2\Omega-|k_0|)^2+\frac{4\Omega^2-2|k_0|\Omega}{[e^{\beta\Omega}-1][e^{\beta(|k_0|-\Omega)}-1]}\right)$$
$$-\frac{\theta(k^2)}{4\pi|\mathbf{k}|^3}\int_{\Omega_+}^{\infty}\mathrm{d}\Omega\left(\frac{\theta(k^0)}{e^{\beta\Omega}-1}+\frac{\theta(-k^0)}{e^{\beta(\Omega-|k_0|)}-1}+\frac{1}{[e^{\beta\Omega}-1][e^{\beta(\Omega-|k_0|)}-1]}\right)(2\Omega-|k_0|)^2$$
$$\mathcal{N}_{\mathrm{T}}^{\beta}(k)=\frac{\theta(-k^2-4m^2)}{16\pi|\mathbf{k}|^5}\int_{\Omega_-}^{\Omega_+}\mathrm{d}\Omega\left(\theta(-k^0)\coth\left(\frac{\beta\Omega}{2}\right)(k^2(2\Omega-|k_0|)^2-(k^2+4m^2)|\mathbf{k}|^2)+\frac{k^2(4\Omega^2-2|k_0|\Omega)-(k^2+4m^2)|\mathbf{k}|^2}{[e^{\beta\Omega}-1][e^{\beta(|k_0|-\Omega)}-1]}\right)$$
$$+\frac{\theta(k^2)}{8\pi|\mathbf{k}|^5}\int_{\Omega_+}^{\infty}\mathrm{d}\Omega\left(\frac{\theta(k^0)}{e^{\beta\Omega}-1}+\frac{\theta(-k^0)}{e^{\beta(\Omega-|k_0|)}-1}+\frac{1}{[e^{\beta\Omega}-1][e^{\beta(\Omega-|k_0|)}-1]}\right)(k^2(2\Omega-|k_0|)^2-(k^2+4m^2)|\mathbf{k}|^2$$

$$\langle H_+(x)\rangle_{\mathcal{E}}\simeq\frac{1}{\sqrt{2}vM^2}\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}\ell^2\mathcal{F}(\ell)-\frac{a_1}{\sqrt{2}vM^2}+\mathcal{O}(v^{-3})$$

$$\mathcal{F}(0)=-iM^{-2}\ \text{ and }\ \mathcal{W}(0)=0$$

$$\langle H_-(x)\rangle_{\mathcal{E}}=\langle H_+(x)\rangle_{\mathcal{E}}$$

$$a_1\simeq\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}\ell^2\mathcal{F}(\ell)+\mathcal{O}(v^{-2})$$

$$\mathcal{Q}_{\mathcal{F}}(k) = -2i(\mathcal{F}(k)^2 - \mathcal{W}(k)\mathcal{W}(-k))\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+\ell^2)\mathcal{F}(\ell)$$
$$+i(\mathcal{F}(k)^2 - \mathcal{W}(k)\mathcal{W}(-k))\cdot\frac{k^2}{M^2}\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\ell^2\mathcal{F}(\ell)$$
$$-\frac{1}{4}\mathcal{F}(k)^2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\mathcal{F}(\ell)\mathcal{F}(k-\ell)$$
$$+\frac{1}{4}\mathcal{F}(k)\mathcal{W}(k)\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k+\ell)^2+\ell^2)\mathcal{W}(\ell)\mathcal{W}(-k-\ell)$$
$$+\frac{1}{4}\mathcal{F}(k)\mathcal{W}(-k)\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)\mathcal{W}(\ell)\mathcal{W}(k-\ell)$$
$$-\frac{1}{4}\mathcal{W}(k)\mathcal{W}(-k)\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+\ell^2+(k-\ell)^2)^2\mathcal{F}^*(\ell)\mathcal{F}^*(k-\ell)$$
$$-i\left(a_2+\left[b_2+\frac{a_1}{M^2}\right]k^2\right)(\mathcal{F}(k)^2 - \mathcal{W}(k)\mathcal{W}(-k))$$

$$\mathcal{Q}_{\mathcal{W}}(k) = -2i\mathcal{W}(k)(\mathcal{F}(k) - \mathcal{F}^*(k))\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+\ell^2)\mathcal{F}(\ell)$$
$$+i\mathcal{W}(k)(\mathcal{F}(k) - \mathcal{F}^*(k))\cdot\frac{k^2}{M^2}\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}\ell^2\mathcal{F}(\ell)$$
$$-\frac{1}{4}\mathcal{W}(k)\mathcal{F}(k)\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\mathcal{F}(\ell)\mathcal{F}(k-\ell)$$
$$+\frac{1}{4}\mathcal{F}^*(k)\mathcal{F}(k)\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\mathcal{W}(\ell)\mathcal{W}(k-\ell)$$
$$+\frac{1}{4}\mathcal{W}(k)\mathcal{W}(k)\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k+\ell)^2+\ell^2)^2\mathcal{W}(\ell)\mathcal{W}(-k-\ell)$$
$$-\frac{1}{4}\mathcal{W}(k)\mathcal{F}^*(k)\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\mathcal{F}^*(\ell)\mathcal{F}^*(k-\ell)$$
$$-i\left(a_2+\left[b_2+\frac{a_1}{M^2}\right]k^2\right)\mathcal{W}(k)(\mathcal{F}(k) - \mathcal{F}^*(k))$$

$$I(k) := -2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+\ell^2)\mathcal{F}(\ell)$$
$$L_{\mathcal{F}(k)} \equiv \int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\mathcal{F}(\ell)\mathcal{F}(k-\ell)$$
$$L_{\mathcal{W}}(k) \equiv \int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\mathcal{W}(\ell)\mathcal{W}(k-\ell)$$

$$\mathcal{Q}_{\mathcal{F}}(k) = i(\mathcal{F}(k)^2 - \mathcal{W}(k)\mathcal{W}(-k))\left(I(k) - \frac{1}{4}\mathrm{Im}[L_{\mathcal{F}}(k)] - a_2 - b_2k^2\right)$$
$$-\frac{1}{4}(\mathcal{F}(k)^2 + \mathcal{W}(k)\mathcal{W}(-k))\mathrm{Re}[L_{\mathcal{F}}(k)]$$
$$+\frac{1}{4}\mathcal{F}(k)(\mathcal{W}(k)L_{\mathcal{W}}(-k) + \mathcal{W}(-k)L_{\mathcal{W}}(k))$$

$$\mathcal{Q}_{\mathcal{W}}(k) = i\mathcal{W}(k)(\mathcal{F}(k) - \mathcal{F}^*(k))\left(I(k) - \frac{1}{4}\mathrm{Im}[L_{\mathcal{F}}(k)] - a_2 - b_2k^2\right)$$
$$-\frac{1}{4}\mathcal{W}(k)(\mathcal{F}(k) + \mathcal{F}^*(k))\mathrm{Re}[L_{\mathcal{F}}(k)]$$
$$+\frac{1}{4}\mathcal{F}^*(k)\mathcal{F}(k)L_{\mathcal{W}}(k) + \frac{1}{4}\mathcal{W}(k)\mathcal{W}(k)L_{\mathcal{W}}(-k)$$

$$I(k) = I^{\mathrm{vac}}(k) + \delta I(k) \text{ and } L_{\mathcal{F},\mathcal{W}}(k) = L^{\mathrm{vac}}_{\mathcal{F},\mathcal{W}}(k) + \delta L_{\mathcal{F},\mathcal{W}}(k)$$

$$I^{\mathrm{vac}}(k) := -2\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2 + \ell^2)\mathcal{F}^{\mathrm{vac}}(\ell)$$
$$L^{\mathrm{vac}}_{\mathcal{F}}(k) \equiv \int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2 + (k-\ell)^2 + \ell^2)^2\mathcal{F}^{\mathrm{vac}}(\ell)\mathcal{F}^{\mathrm{vac}}(k-\ell)$$
$$L^{\mathrm{vac}}_{\mathcal{W}}(k) \equiv \int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2 + (k-\ell)^2 + \ell^2)^2\mathcal{W}^{\mathrm{vac}}(\ell)\mathcal{W}^{\mathrm{vac}}(k-\ell)$$

$$I^{\mathrm{vac}}(k) = -2k^2\mathcal{L}_{(0,1)}(M^2) - 2\mathcal{L}_{(1,1)}(M^2)$$
$$= -\frac{2M^2(k^2 - M^2)\Gamma\left(1 - \frac{D}{2}\right)}{(4\pi)^{D/2}}\left(\frac{M^2}{\mu^2}\right)^{\frac{D-4}{2}}$$
$$\simeq \frac{M^2(k^2 - M^2)}{4\pi^2(4-D)} - \frac{M^2(k^2 - M^2)}{8\pi^2}\log\left(\frac{M^2}{4\pi e^{1-\gamma}\mu^2}\right) + \mathcal{O}(4-D)$$

$$L^{\mathrm{vac}}_{\mathcal{F}}(k) = -\frac{1}{2}\int \frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2 + (k-\ell)^2 + \ell^2)^2\int_{-1}^{+1}\frac{\mathrm{d}y}{\left[\ell^2 + M^2 + (y-1)\left(k\cdot\ell - \frac{1}{2}k^2\right) - i\epsilon\right]^2}$$
$$= -2\int_{-1}^{+1}\mathrm{d}y\int \frac{\mathrm{d}^4p}{(2\pi)^4}\frac{\left[p^2 - yp\cdot k + \frac{1}{4}(3+y^2)k^2\right]^2}{\left(p^2 + \frac{1}{4}\Sigma(y) - i\epsilon\right)^2}$$

$$\Sigma(y) = (1-y^2)k^2 + 4M^2 \propto (p\cdot k)^2 = k^\mu k^\nu p_\mu p_\nu$$

$$L^{\mathrm{vac}}_{\mathcal{F}}(k) = -2\int_{-1}^{+1}\mathrm{d}y\int \frac{\mathrm{d}^Dp}{(2\pi)^D}\frac{(p^2)^2 + \left[\left(\frac{1}{D} + \frac{1}{2}\right)y^2 + \frac{3}{2}\right]k^2p^2 + \frac{1}{16}(3+y^2)^2(k^2)^2}{\left(p^2 + \frac{1}{4}\Sigma(y) - i\epsilon\right)^2}$$
$$= -2i\int_{-1}^{+1}\mathrm{d}y\left[\mathcal{L}_{(2,2)}\left(\frac{1}{4}\Sigma\right) + \frac{\left(\frac{2}{D} + 1\right)y^2 + 3}{2}k^2\mathcal{L}_{(1,2)}\left(\frac{1}{4}\Sigma\right) + \frac{(3+y^2)^2(k^2)^2}{16}\mathcal{L}_{(0,2)}\left(\frac{1}{4}\Sigma\right)\right]$$

$$L^{\mathrm{vac}}_{\mathcal{F}}(k) = 4i\int_{-1}^{+1}\mathrm{d}y\frac{\Gamma\left(1 - \frac{D}{2}\right)}{(16\pi)^{D/2}}\left(\frac{(1-y^2)k^2 + 4M^2 - i\epsilon}{\mu^2}\right)^{\frac{D-4}{2}}$$
$$\times[(k^2)^2((D+1)y^4 + (2D-5)y^2 + D - 4) - 4k^2M^2((D+2)y^2 + D - 1) + 4(D+2)M^4]$$
$$\simeq -i\frac{(k^2)^2 - 10k^2M^2 + 12M^4}{8\pi^2(4-D)} + i\frac{11(k^2)^2 - 10k^2M^2 - 6M^4}{120\pi^2}$$
$$+i\int_{-1}^{+1}\mathrm{d}y\frac{y^2(5y^2+3)(k^2)^2 - 12(2y^2+1)k^2M^2 + 24M^4}{64\pi^2}\log\left(\frac{(1-y^2)k^2 + 4M^2 - i\epsilon}{16\pi e^{3/5-\gamma}\mu^2}\right)$$

$$\log(x - i\epsilon) = \log|x| - i\pi\theta(-x)$$

$$\mathrm{Re}[L_{\mathcal{F}}^{\mathrm{vac}}(k)] = \frac{\theta(-k^2-4M^2)}{16\pi}(k^2-2M^2)^2\sqrt{1+\frac{4m^2}{k^2}}$$

$$\mathrm{Im}[L_{\mathcal{F}}^{\mathrm{vac}}(k)] = -\frac{(k^2)^2-10k^2M^2+12M^4}{8\pi^2(4-D)}+\frac{11(k^2)^2-10k^2M^2-6M^4}{120\pi^2}$$
$$+\int_{-1}^{+1}\mathrm{d}y\,\frac{y^2(5y^2+3)(k^2)^2-12(2y^2+1)k^2M^2+24M^4}{64\pi^2}\log\left|\frac{(1-y^2)k^2+4M^2}{16\pi e^{3/5-\gamma}\mu^2}\right|$$

$$L_{\mathcal{W}}^{\mathrm{vac}}(k) = \frac{\theta(k^0)\theta(-k^2-4M^2)}{8\pi}(k^2-2M^2)^2\sqrt{1+\frac{4m^2}{k^2}}$$

$$\left(I(k)-\frac{1}{4}\mathrm{Im}[L_{\mathcal{F}}(k)]\right)_{\text{divergence}} \simeq \frac{(k^2+M^2)^2-4k^2M^2+3M^4}{32\pi^2(4-D)}+\cdots$$

$$S_{\mathcal{S}}[A_+,\chi_+,c_+,\bar{c}_+]-S_{\mathcal{S}}[A_-,\chi_-,c_-,\bar{c}_-]+S_{\mathrm{IF}}[A_+,\chi_+,A_-,\chi_-]$$
$$\supset -\frac{b_4}{4v^4}\int \mathrm{d}^4x[(\mathcal{D}_+\chi_+(x))^4-(\mathcal{D}_-\chi_-(x))^4]$$
$$+\frac{i}{4v^4}\int \mathrm{d}^4x\int \mathrm{d}^4y[(\mathcal{D}_+\chi_+(x))^2\mathcal{Q}_{\mathcal{F}}(x,y)(\mathcal{D}_+\chi_+(y))^2+(\mathcal{D}_-\chi_-(x))^2\mathcal{Q}_{\mathcal{F}}^*(x,y)(\mathcal{D}_-\chi_-(y))^2]$$
$$\supset -\frac{1}{4v^4}\int \mathrm{d}^4x\int \mathrm{d}^4y[(\mathcal{D}_+\chi_+(x))^2(\mathcal{D}_+\chi_+(y))^2-(\mathcal{D}_-\chi_-(x))^2(\mathcal{D}_-\chi_-(y))^2]\left[b_4\delta(x-y)+\mathrm{Im}[\mathcal{Q}_F(x,y)]\right]$$

$$\mathcal{Q}_{\mathcal{F}}(k)\to\mathcal{Q}_{\mathcal{F}}(k)+ib_4$$
$$= i(\mathcal{F}(k)^2-\mathcal{W}(k)\mathcal{W}(-k))\left(I(k)-\frac{1}{4}\mathrm{Im}[L_{\mathcal{F}}(k)]-a_2-b_2k^2\right)$$
$$-\frac{1}{4}(\mathcal{F}(k)^2+\mathcal{W}(k)\mathcal{W}(-k))\mathrm{Re}[L_{\mathcal{F}}(k)]$$
$$+\frac{1}{4}\mathcal{F}(k)(\mathcal{W}(k)L_{\mathcal{W}}(-k)+\mathcal{W}(-k)L_{\mathcal{W}}(k))+ib_4$$

$$\mathcal{F}(k)=\frac{-i}{k^2+M^2}+2\pi\left(\frac{1}{2}+\mathfrak{n}(k)\right)\delta(k^2+M^2)\ \text{ and }\ \mathcal{W}(k)=2\pi(\theta(k_0)+\mathfrak{n}(k))\delta(k^2+M^2)$$

$$i(\mathcal{F}(k)^2-\mathcal{W}(k)\mathcal{W}(-k))=\frac{-i}{(k^2+M^2)^2}-2\pi\left(\frac{1}{2}+\mathfrak{n}(k)\right)\delta'(k^2+M^2)$$

$$\frac{\delta(x)}{x+i\epsilon}=-\frac{1}{2}\delta'(x)-i\pi[\delta(x)]^2$$

$(x-i\epsilon)^{-2}=x^{-2}-i\pi\delta'(x)$ where formally $x^{-2}\to\mathcal{H}(x^{-2})$ with $\mathcal{H}$ the Hadamard finite part.

using the identity $x^2\delta'(x)=0$, we have

$$(k^2+M^2)^2\cdot i(\mathcal{F}(k)^2-\mathcal{W}(k)\mathcal{W}(-k))=-i$$

$$\mathcal{Q}_{\mathcal{F}}(k)=i(\mathcal{F}(k)^2-\mathcal{W}(k)\mathcal{W}(-k))\left(I(k)-\frac{1}{4}\mathrm{Im}[L_{\mathcal{F}}(k)]-a_2-b_2k^2-b_4(k^2+M^2)^2\right)$$
$$-\frac{1}{4}(\mathcal{F}(k)^2+\mathcal{W}(k)\mathcal{W}(-k))\mathrm{Re}[L_{\mathcal{F}}(k)]$$
$$+\frac{1}{4}\mathcal{F}(k)(\mathcal{W}(k)L_{\mathcal{W}}(-k)+\mathcal{W}(-k)L_{\mathcal{W}}(k))$$

$$i\mathcal{W}(k)(\mathcal{F}(k)-\mathcal{F}^*(k)) = -2\pi[\theta(k^0)+\mathfrak{n}(k)]\delta'(k^2+M^2)$$

$$(k^2+M^2)^2\cdot i\mathcal{W}(k)(\mathcal{F}(k)-\mathcal{F}^*(k)) = 0$$

$$\begin{aligned}\mathcal{Q}_{\mathcal{W}}(k) =& i\mathcal{W}(k)(\mathcal{F}(k)-\mathcal{F}^*(k))\left(I(k)-\frac{1}{4}\mathrm{Im}[L_{\mathcal{F}}(k)]-a_2-b_2k^2-b_4(k^2+M^2)^2\right)\\ &-\frac{1}{4}\mathcal{W}(k)(\mathcal{F}(k)+\mathcal{F}^*(k))\mathrm{Re}[L_{\mathcal{F}}(k)]\\ &+\frac{1}{4}\mathcal{F}^*(k)\mathcal{F}(k)L_{\mathcal{W}}(k)+\frac{1}{4}\mathcal{W}(k)\mathcal{W}(k)L_{\mathcal{W}}(-k)\end{aligned}$$

$$\begin{bmatrix}\mathcal{Q}_{\mathcal{F}}(k) & \mathcal{Q}_{\mathcal{W}}(-k)\\ \mathcal{Q}_{\mathcal{W}}(k) & \mathcal{Q}^*_{\mathcal{F}}(k)\end{bmatrix} = \begin{bmatrix}\mathcal{F}(k) & \mathcal{W}(-k)\\ \mathcal{W}(k) & \mathcal{F}^*(k)\end{bmatrix}\begin{bmatrix}\Sigma_{\mathcal{F}}(k) & \Sigma_{\mathcal{W}}(-k)\\ \Sigma_{\mathcal{W}}(k) & \Sigma^*_{\mathcal{F}}(k)\end{bmatrix}\begin{bmatrix}\mathcal{F}(k) & \mathcal{W}(-k)\\ \mathcal{W}(k) & \mathcal{F}^*(k)\end{bmatrix}.$$

$$\begin{aligned}\mathrm{Re}[\Sigma_{\mathcal{F}}(k)] &= -\frac{1}{4}\mathrm{Re}[L_{\mathcal{F}}(k)]\\ \mathrm{Im}[\Sigma_{\mathcal{F}}(k)] &= I(k)-\frac{1}{4}\mathrm{Im}[L_{\mathcal{F}}(k)]-a_2-b_2k^2-b_4(k^2+M^2)^2\\ \Sigma_{\mathcal{W}}(k) &= \frac{1}{4}L_{\mathcal{W}}(k)\end{aligned}$$

$$\begin{aligned}\mathrm{Im}[\Sigma^{\mathrm{vac}}_{\mathcal{F}}(k)]:&= I^{\mathrm{vac}}(k)-\frac{1}{4}\mathrm{Im}[L^{\mathrm{vac}}_{\mathcal{F}}(k)]-a_2-b_2k^2-b_4(k^2+M^2)^2\\ &=\frac{(k^2+M^2)^2-4k^2M^2+3M^4}{32\pi^2(4-D)}-a_2-b_2k^2-b_4(k^2+M^2)^2\\ &-\frac{11(k^2)^2-10k^2M^2-6M^4}{480\pi^2}-\frac{M^2(k^2-M^2)}{8\pi^2}\log\left(\frac{M^2}{4\pi e^{1-\gamma}\mu^2}\right)\\ &-\int_{-1}^{+1}\mathrm{d}y\frac{y^2(5y^2+3)(k^2)^2-12(2y^2+1)k^2M^2+24M^4}{256\pi^2}\log\left|\frac{(1-y^2)k^2+4M^2}{16\pi e^{3/5-\gamma}\mu^2}\right|\end{aligned}$$

$$\begin{aligned}\mathrm{Im}[\Sigma^{\mathrm{vac}}_{\mathcal{F}}(k)] =& -\frac{M^2(k^2-M^2)}{8\pi^2}\log\left(\frac{M^2}{\mu^2}\right)\\ &-\int_{-1}^{+1}\mathrm{d}y\frac{y^2(5y^2+3)(k^2)^2-12(2y^2+1)k^2M^2+24M^4}{256\pi^2}\log\left|\frac{(1-y^2)k^2+4M^2}{\mu^2}\right|\end{aligned}$$

$$\delta I(k) = -2\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+\ell^2)\cdot 2\pi\mathfrak{n}(\ell)\delta(\ell^2+M^2)$$

$$\delta L_{\mathcal{F}}(k) = 2L_{\mathcal{F}\mathfrak{n}}(k)+L_{\mathfrak{n}\mathfrak{n}}(k)\ \text{ and }\ \delta L_{\mathcal{W}}(k) = 2L_{\mathcal{W}_{\mathfrak{n}}}(k)+L_{\mathfrak{n}\mathfrak{n}}(k)$$

$$\begin{aligned}L_{\mathcal{F}\mathfrak{n}}(k) &\equiv \int\frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\cdot 2\pi\mathfrak{n}(\ell)\delta(\ell^2+M^2)\cdot\frac{-i}{(k-\ell)^2+M^2-i\epsilon}\\ L_{\mathcal{W}\mathfrak{n}}(k) &\equiv \int\frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\cdot 2\pi\mathfrak{n}(\ell)\delta(\ell^2+M^2)\cdot 2\pi\theta(k^0-\ell^0)\delta((k-\ell)^2+M^2)\\ L_{\mathfrak{n}\mathfrak{n}}(k) &\equiv \int\frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2\cdot 2\pi\mathfrak{n}(\ell)\delta(\ell^2+M^2)\cdot 2\pi\mathfrak{n}(k-\ell)\delta((k-\ell)^2+M^2)\end{aligned}$$

$$n(\mathbf{k})\to n(|\mathbf{k}|)$$

$$\mathfrak{n}(k) = \theta(k^0)n(|\mathbf{k}|)+\theta(-k^0)n(|\mathbf{k}|) = n(|\mathbf{k}|)$$

$$\delta I(k) = \frac{M^2 - k^2}{\pi^2}\int_M^\infty \mathrm{d}\Omega\sqrt{\Omega^2 - M^2}\, n\left(\sqrt{\Omega^2 - M^2}\right)$$

$$\Omega = \sqrt{|\boldsymbol{\ell}|^2 + M^2}$$

$$L_{\mathcal{F}\mathfrak{n}}(k) = \frac{i(-3k^2 + 4M^2)}{2\pi^2}\int_M^\infty \mathrm{d}\Omega\sqrt{\Omega^2 - M^2}\, n\left(\sqrt{\Omega^2 - M^2}\right)$$
$$+\frac{i(k^2 - 2M^2)^2}{16\pi^2|\mathbf{k}|}\int_M^\infty \mathrm{d}\Omega n\left(\sqrt{\Omega^2 - M^2}\right)\left[g\left(\frac{k^2/2 + k_0\Omega}{|\mathbf{k}|\sqrt{\Omega^2 - M^2}}\right) + g\left(\frac{k^2/2 - k_0\Omega}{|\mathbf{k}|\sqrt{\Omega^2 - M^2}}\right)\right]$$

$$\mathrm{Re}[L_{\mathcal{F}\mathfrak{n}}(k)] = \frac{\theta(k^2)(k^2 - 2M^2)^2}{16\pi|\mathbf{k}|}\int_{\Omega_+}^\infty \mathrm{d}\Omega\left[n\left(\sqrt{\Omega^2 - M^2} + n\left(\sqrt{(\Omega - |k_0|)^2 - M^2}\right)\right]\right.$$
$$+\frac{\theta(-k^2 - 4M^2)(k^2 - 2M^2)^2}{8\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+} \mathrm{d}\Omega n\left(\sqrt{\Omega^2 - M^2}\right)$$

$$\mathrm{Im}[L_{\mathcal{F}\mathfrak{n}}(k)] = \frac{(-3k^2 + 4M^2)}{2\pi^2}\int_M^\infty \mathrm{d}\Omega\sqrt{\Omega^2 - M^2}\, n\left(\sqrt{\Omega^2 - M^2}\right)$$
$$+\frac{(k^2 - 2M^2)^2}{16\pi^2|\mathbf{k}|}\int_M^\infty \mathrm{d}\Omega n\left(\sqrt{\Omega^2 - M^2}\right)\log\left|\frac{\left(k^2 - 2|\mathbf{k}|\sqrt{\Omega^2 - M^2}\right)^2 - 4k_0^2\Omega^2}{\left(k^2 + 2|\mathbf{k}|\sqrt{\Omega^2 - M^2}\right)^2 - 4k_0^2\Omega^2}\right|$$

$$L_{\mathcal{W}\mathfrak{n}}(k) = \frac{\theta(k^0)\theta(-k^2 - 4M^2)(k^2 - 2M^2)^2}{8\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+} \mathrm{d}\Omega n\left(\sqrt{\Omega^2 - M^2}\right)$$
$$+\frac{\theta(k^2)(k^2 - 2M^2)^2}{8\pi|\mathbf{k}|}\int_{\Omega_+}^\infty \mathrm{d}\Omega\left[\theta(-k^0)n\left(\sqrt{\Omega^2 - M^2}\right) + \theta(k^0)n\left(\sqrt{(\Omega - |k_0|)^2 - M^2}\right)\right]$$

$$L_{\mathfrak{n}\mathfrak{n}}(k) = \frac{\theta(-k^2 - 4M^2)(k^2 - 2M^2)^2}{8\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+} \mathrm{d}\Omega n\left(\sqrt{\Omega^2 - M^2}\right)n\left(\sqrt{(\Omega - |k_0|)^2 - M^2}\right)$$
$$+\frac{\theta(k^2)(k^2 - 2M^2)^2}{4\pi|\mathbf{k}|}\int_{\Omega_+}^\infty \mathrm{d}\Omega n\left(\sqrt{\Omega^2 - M^2}\right)n\left(\sqrt{(\Omega - |k_0|)^2 - M^2}\right)$$

$$n(|\mathbf{k}|) = \frac{1}{e^{\beta\sqrt{|\mathbf{k}|^2 + M^2}} + 1}$$

$$L_{\mathfrak{n}\mathfrak{n}}(k) = \frac{\theta(-k^2 - 4M^2)(-k^2 + 2M^2)^2}{8\pi|\mathbf{k}|}\int_{\Omega_-}^{\Omega_+} \frac{\mathrm{d}\Omega}{[e^{\beta\Omega} - 1][e^{\beta(|k_0| - \Omega)} - 1]}$$
$$+\frac{\theta(k^2)(-k^2 + 2M^2)^2}{4\pi|\mathbf{k}|}\int_{\Omega_+}^\infty \frac{\mathrm{d}\Omega}{[e^{\beta\Omega} - 1][e^{\beta(\Omega - |k_0|)} - 1]}$$

$$L_{\mathfrak{n}\mathfrak{n}}(k) = \frac{\theta(-k^2 - 4M^2)(-k^2 + 2M^2)^2}{8\pi|\mathbf{k}|}\left[\frac{\log\left(e^{\beta\Omega} - 1\right) - \log\left(e^{\beta(|k_0| - \Omega)} - 1\right) - \beta\Omega}{\beta(e^{\beta|k_0|} - 1)}\right]\Bigg|_{\Omega_-}^{\Omega_+}$$
$$+\frac{\theta(k^2)(-k^2 + 2M^2)^2}{4\pi|\mathbf{k}|}\left[\frac{\log\left(e^{\beta\Omega} - 1\right)}{\beta(e^{-\beta|k_0|} - 1)} + \frac{\log\left(e^{\beta(\Omega - |k_0|)} - 1\right)}{\beta(e^{\beta|k_0|} - 1)} + \Omega\right]\Bigg|_{\Omega_+}^\infty$$

$$\Sigma_{\mathcal{F}}(k) \simeq -\tfrac{1}{4}L_{\mathfrak{n}\mathfrak{n}}(k) \text{ and } \Sigma_{\mathcal{W}}(k) \simeq +\tfrac{1}{4}L_{\mathfrak{n}\mathfrak{n}}(k)$$

$$\mathcal{F}(k) = \frac{-i}{k^2+M^2} + f(k)\delta(k^2+M^2) \text{ with } f(k) = 2\pi\left(\frac{1}{2} + \mathfrak{n}(k)\right)$$
$$\mathcal{W}(k) = w(k)\delta(k^2+M^2) \text{ with } w(k) = 2\pi(\theta(k_0) + \mathfrak{n}(k))$$

$$\mathfrak{n}(k) = \theta(k^0)n(\mathbf{k}) + \theta(-k^0)n(-\mathbf{k})$$

$$f(k) = f(-k) \text{ however } w(k) \neq w(-k)$$

$$\Gamma_{\mathrm{rrr}}(x,y,z) \equiv \Gamma_{\mathcal{U}}(x,y,z) + \Gamma^*_{\mathcal{U}}(x,y,z) - 3\Gamma_{\mathcal{N}}(x,y,z) - 3\Gamma^*_{\mathcal{N}}(x,y,z)$$

$$\Gamma_{\mathrm{rrr}}(x,y,z) = \int \frac{\mathrm{d}^4k}{(2\pi)^4}\int \frac{\mathrm{d}^4p}{(2\pi)^4}\int \frac{\mathrm{d}^4q}{(2\pi)^4}(2\pi)^4\delta^{(4)}(k+p+q)\bar{\Gamma}_{\mathrm{rrr}}(k,p,q)e^{ik\cdot x+ip\cdot y+iq\cdot z}$$

$$\frac{\bar{\Gamma}_{\mathrm{rrr}}(k,p,q)}{k^2+p^2+q^2} = \frac{\Gamma_{\mathcal{U}}(k,p,q) + \Gamma^*_{\mathcal{U}}(-k,-p,-q) - 3\Gamma_{\mathcal{N}}(k,p,q) - 3\Gamma^*_{\mathcal{N}}(-k,-p,-q)}{k^2+p^2+q^2}$$
$$= -2\left(\frac{f(k)\delta(k^2+M^2)}{(M^2+p^2)(M^2+q^2)} + \frac{f(p)\delta(M^2+p^2)}{(k^2+M^2)(M^2+q^2)} + \frac{\left(f(q) - \frac{3}{2}(w(-q)+w(q))\right)\delta(M^2+q^2)}{(k^2+M^2)(M^2+p^2)}\right)$$
$$-3i\left(\frac{f(p)(w(-q)-w(q))\delta(M^2+p^2)\delta(M^2+q^2)}{k^2+M^2} + \frac{f(k)(w(-q)-w(q))\delta(k^2+M^2)\delta(M^2+q^2)}{M^2+p^2}\right.$$
$$\left. + \frac{\delta(k^2+M^2)(w(k)w(p)-w(-k)w(-p))\delta(M^2+p^2)}{M^2+q^2}\right)$$
$$+[3f(q)w(-k)w(-p) + 3f(q)w(k)w(p) - 3f(k)f(p)w(-q) - 3f(k)f(p)w(q)$$
$$+2f(k)f(p)f(q) - w(-k)w(-p)w(-q) - w(k)w(p)w(q)]\delta(k^2+M^2)\delta(M^2+p^2)\delta(M^2+q^2)$$

$w(k) + w(-k) = 2f(k)$, and replace $w(-k) = 2f(k) - w(k)$

$$\frac{\bar{\Gamma}_{\mathrm{rrr}}(k,p,q)}{k^2+p^2+q^2} = -2\left(\frac{f(k)\delta(k^2+M^2)}{(M^2+p^2)(M^2+q^2)} + \frac{f(p)\delta(M^2+p^2)}{(k^2+M^2)(M^2+q^2)} - \frac{2f(q)\delta(M^2+q^2)}{(k^2+M^2)(M^2+p^2)}\right)$$
$$-6i\left[\frac{\delta(k^2+M^2)\delta(M^2+p^2)(f(p)w(k) + f(k)w(p) - 2f(k)f(p))}{M^2+q^2}\right.$$
$$\left. + \frac{\delta(M^2+p^2)\delta(M^2+q^2)(f(p)f(q) - f(p)w(q))}{k^2+M^2} + \frac{\delta(k^2+M^2)\delta(M^2+q^2)(f(k)f(q) - f(k)w(q))}{M^2+p^2}\right]$$
$$-2[f(p)f(q)w(k) + f(k)f(q)w(p) - 2f(k)f(p)w(q)$$
$$+f(p)w(k)w(q) - 2f(q)w(k)w(p) + f(k)w(p)w(q)]\delta(k^2+M^2)\delta(M^2+p^2)\delta(M^2+q^2)$$

$$e^{ik\cdot x+ip\cdot y+iq\cdot z}\int \mathrm{d}^{\square}x\int \mathrm{d}^{\square}y\int \mathrm{d}^{\square}z$$

$$\int \mathrm{d}^4x(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(x))^2\int \mathrm{d}^4y(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(y))^2\int \mathrm{d}^4z(\mathcal{D}_{\mathrm{r}}\chi_{\mathrm{r}}(z))^2\bar{\Gamma}_{\mathrm{rrr}}(x,y,z) = 0$$

$$\mathcal{Q}_{\mathrm{rr}}(x,y) \equiv \int \frac{\mathrm{d}^4k}{(2\pi)^4}\overline{\mathcal{Q}}_{\mathrm{rr}}(k)e^{ik\cdot(x-y))} \text{ with } \overline{\mathcal{Q}}_{\mathrm{rr}}(k) \equiv \mathcal{Q}_{\mathcal{F}(k)} + \mathcal{Q}^*_{\mathcal{F}}(k) - \mathcal{Q}_{\mathcal{W}}(k) - \mathcal{Q}_{\mathcal{F}}(-k)$$

$$\mathcal{Q}_{\mathcal{F}}(-k) = \mathcal{Q}_{\mathcal{F}}(k) \text{ and } \mathcal{Q}^*_{\mathcal{W}}(k) = \mathcal{Q}_{\mathcal{W}}(k)$$

$$\overline{\mathcal{Q}}_{\mathrm{rr}}(k)=i\left(I(k)-\frac{1}{4}\mathrm{Im}\left[L_{\mathcal{F}(k)}\right]\right)\{\mathcal{F}(k)^2-\mathcal{F}^*(k)^2-(\mathcal{W}(k)+\mathcal{W}(-k))(\mathcal{F}(k)-\mathcal{F}^*(k))\}$$
$$-\frac{1}{4}\mathrm{Re}\left[L_{\mathcal{F}(k)}\right]\{\mathcal{F}(k)^2+\mathcal{F}^*(k)^2+2\mathcal{W}(k)\mathcal{W}(-k)$$
$$+(\mathcal{W}(k)+\mathcal{W}(-k))(\mathcal{F}(k)+\mathcal{F}^*(k))\}$$
$$+\frac{1}{4}\{(\mathcal{F}(k)+\mathcal{F}^*(k))(\mathcal{W}(k)L_{\mathcal{W}}(-k)+\mathcal{W}(-k)L_{\mathcal{W}}(k))$$
$$-\mathcal{F}^*(k)\mathcal{F}(k)L_{\mathcal{W}}(k)-\mathcal{W}(k)\mathcal{W}(k)L_{\mathcal{W}}(-k)$$
$$-\mathcal{F}^*(k)\mathcal{F}(k)L_{\mathcal{W}}(-k)-\mathcal{W}(-k)\mathcal{W}(-k)L_{\mathcal{W}}(k)\}$$

$w(-k)=2f(k)-w(k)$, and use $f(k)=2\pi\left(\frac{1}{2}+\mathfrak{n}(k)\right)$ and $w(k)=2\pi(\theta(k^0)+\mathfrak{n}(k))$

$$\overline{\mathcal{Q}}_{\mathrm{rr}}(k)=-\frac{1}{4}\mathrm{Re}\left[L_{\mathcal{F}(k)}\right]\left\{-\frac{2}{(k^2+M^2)^2}-2\pi^2[\delta(k^2+M^2)]^2\right\}$$
$$+\frac{1}{4}\left\{-(L_{\mathcal{W}}(k)+L_{\mathcal{W}}(-k))\left(\frac{1}{(k^2+M^2)^2}+\pi^2[\delta(k^2+M^2)]^2\right)\right\}$$
$$=+\frac{1}{4}\left(2\mathrm{Re}\left[L_{\mathcal{F}(k)}\right]-L_{\mathcal{W}}(k)-L_{\mathcal{W}}(-k)\right)\mathcal{F}^{\mathrm{vac}}(k)\mathcal{F}^{\mathrm{vac}*}(k)$$

$$2\mathrm{Re}[L_{\mathcal{F}}(k)]-L_{\mathcal{W}}(k)-L_{\mathcal{W}}(-k)$$
$$=\int\frac{\mathrm{d}^4\ell}{(2\pi)^4}(k^2+(k-\ell)^2+\ell^2)^2[\mathcal{F}(\ell)\mathcal{F}(k-\ell)+\mathcal{F}^*(\ell)\mathcal{F}^*(k-\ell)$$
$$-\mathcal{W}(\ell)\mathcal{W}(k-\ell)-\mathcal{W}(-\ell)\mathcal{W}(-k+\ell)]=0$$

$$Z_{\text{in-in}}\left[\boldsymbol{J}^+,\boldsymbol{J}^-\right]=\mathrm{Tr}\left[\mathcal{U}(\boldsymbol{J}^+)\rho_{\mathrm{i}}\mathcal{U}^\dagger(\boldsymbol{J}^-)\right]$$

$$\mathcal{U}(\mathbf{0})=U(t_{\mathrm{f}},t_{\mathrm{i}})$$

$$Z_{\text{in-in}}\left[\boldsymbol{J},\boldsymbol{J}\right]=1$$

$$\frac{\delta P_{\mu\nu}(x)}{\delta F_{\alpha\beta}(y)}=0\text{ when }y^0>x^0$$

$$V^\star=\Omega^{m-k}(M)\times\Omega^{m-k-1}(\partial M)$$

$$\langle(\alpha,\alpha_\partial),\varphi\rangle=\int_M\varphi\wedge\alpha+\int_{\partial M}\imath_{\partial M}^*\varphi\wedge\alpha_\partial,\varphi\in V,(\alpha,\alpha_\partial)\in V^\star$$

$$L_\nabla\colon TV\to\mathbb{R}$$

$$L_\nabla(\varphi,\dot{\varphi})=\int_M\mathcal{L}\left(\varphi(x),\dot{\varphi}(x),\mathrm{d}^\nabla\varphi(x)\right)$$

$$L\colon T\mathcal{C}(P)\to\mathbb{R}$$

$$L_\nabla(A,\dot{A})=\int_M\mathcal{L}\left(A(x),\dot{A}(x),\mathrm{d}^A A(x)\right)$$

$$L\colon TQ\to\mathbb{R},Q=\mathcal{C}(P)\times\Omega^0(M,\tilde{V})$$

$$L\left(A,\dot{A},\ \mathrm{d}^A A,\varphi,\dot{\varphi},\ \mathrm{d}^A\varphi\right)=\int_M\mathcal{L}_{\mathrm{gau}}\left(A(x),\dot{A}(x),\mathrm{d}^A A(x)\right)+\mathcal{L}_{\mathrm{mat}}\left(\varphi(x),\dot{\varphi}(x),\mathrm{d}^A\varphi(x)\right)$$

$\Omega^b_{T^*Q}: T(T^*Q) \to T^*(T^*Q)$, namely, $D_{T^*Q} = \text{graph}\left(\Omega^b_{T^*Q}\right) \subset T(T^*Q) \oplus T^*(T^*Q)$

$$D_{T^*Q}(p_q) = \text{graph}\left(\Omega^b_{T^*Q}(p_q)\right)$$
$$= \left\{\left(v_{p_q}, \alpha_{p_q}\right) \in T_{p_q}(T^*Q) \times T^*_{p_q}(T^*Q) \mid \alpha_{p_q} = \Omega^b_{T^*Q}(p_q)\left(v_{p_q}\right)\right\}$$

$\mathrm{d}_D L = \gamma_Q \circ \mathrm{d}L: TQ \to T^*(T^*Q)$, where $\gamma_Q: T^*(TQ) \to T^*(T^*Q)$ is the canonical isomorphism given

by $\gamma_Q(q, \delta q, \delta p, p) = (q, p, -\delta p, \delta q)$

$$\mathrm{d}_D L(q, v) = \left(q, \frac{\partial L}{\partial v}, -\frac{\partial L}{\partial q}, v\right)$$

map $\tilde{F}: TQ \to T^*(T^*Q)$

$$\langle \tilde{F}(q, v), W\rangle = \left\langle F(q, v), T_{\mathbb{F}L(q,v)}\pi_Q(W)\right\rangle$$

for $(q, v) \in TQ$ and $W \in T_{\mathbb{F}L(q,v)}(T^*Q)$

$\mathbb{F}L(q, v) = \left(q, \frac{\partial L}{\partial v}(q, v)\right)$, and $\pi_Q: T^*Q \to Q$ is the projection map

$$\tilde{F}(q, v) = \left(q, \frac{\partial L}{\partial v}(q, v), F(q, v), 0\right)$$

$$(q, v, p): [t_0, t_1] \to TQ \oplus T^*Q$$

$$\left((q, p, \dot{q}, \dot{p}), \mathrm{d}_D L(q, v) - \tilde{F}(q, v)\right) \in D_{T^*Q}(q, p)$$

$$\begin{cases} \dot{q} = v \\ p = \dfrac{\partial L}{\partial v}(q, v) \\ \dot{p} = \dfrac{\partial L}{\partial q}(q, v) + F(q, \dot{q}) \end{cases}$$

for the curve $(q, v, p): [t_0, t_1] \to TQ \oplus T^*Q$ imply Euler-Lagrange equations

$$\frac{d}{dt}\frac{\partial L}{\partial \dot{q}}(q, \dot{q}) - \frac{\partial L}{\partial q}(q, \dot{q}) = F(q, \dot{q})$$

$$\delta \int_{t_0}^{t_1} (L(q, v) + \langle p, \dot{q} - v\rangle) dt + \int_{t_0}^{t_1} \langle F(q, \dot{q}), \delta q\rangle dt = 0$$

$$\delta \int_{t_0}^{t_1} L(q, \dot{q}) dt + \int_{t_0}^{t_1} \langle F(q, \dot{q}), \delta q\rangle dt = 0$$

$$V^\dagger = \Omega^{m-k}(M, E^*) \times \Omega^{m-k-1}(\partial M, E^*)$$

$$\left((\varphi, \alpha, \alpha_\partial, \dot{\varphi}, \dot{\alpha}, \dot{\alpha}_\partial), \mathrm{d}^\dagger_D L_\nabla(\varphi, \nu) - \tilde{F}^\dagger(\varphi, \nu)\right) \in D_{T^\dagger V}(\varphi, \alpha, \alpha_\partial)$$

for a curve $(\varphi, \nu, \alpha, \alpha_\partial): [t_0, t_1] \to TV \oplus T^\dagger V$, where $F^\dagger: TV \to T^\dagger V$

$$\begin{cases}\dfrac{\partial}{\partial t}\dfrac{\partial \mathcal{L}}{\partial \dot{\varphi}} = \dfrac{\partial \mathcal{L}}{\partial \varphi} - (-1)^k \, \mathrm{d}^* \dfrac{\partial \mathcal{L}}{\partial \zeta} + \mathcal{F}^\dagger \\ \mathcal{F}_\partial^\dagger = -\iota_{\partial M}^* \dfrac{\partial \mathcal{L}}{\partial \zeta}\end{cases}$$

$$\frac{d}{dt}\int_M \mathcal{E} = \underbrace{\int_M \dot{\varphi} \wedge \mathcal{F}^\dagger}_{\text{spatially distributed contribution}} + \underbrace{\int_{\partial M} \iota_{\partial M}^* \dot{\varphi} \wedge \mathcal{F}_\partial^\dagger}_{\text{boundary contribution}}$$

$$\mathcal{C}(P)^\dagger = \Omega^{m-1}(M, \tilde{\mathfrak{g}}^*) \times \Omega^{m-2}(\partial M, \tilde{\mathfrak{g}}^*)$$

$$\left((A, \varsigma, \varsigma_\partial, \dot{A}, \dot{\varsigma}, \dot{\varsigma}_\partial), \mathrm{d}_D^\dagger L(A, \varepsilon) - \tilde{F}^\dagger(A, \varepsilon)\right) \in D_{T^\dagger \mathcal{C}(P)}(A, \varsigma, \varsigma_\partial),$$

for a curve $(A, \varepsilon, \varsigma, \varsigma_\partial)\colon [t_0, t_1] \to T\mathcal{C}(P) \oplus T^\dagger \mathcal{C}(P)$, where $F^\dagger\colon T\mathcal{C}(P) \to T^\dagger \mathcal{C}(P)$ is a given force

$$\begin{cases}\dot{E} - \delta^A B_A = -J, \\ \star_\partial \left(\iota_{\partial M}^*(\star\, B_A)\right) = j,\end{cases}$$

$$\mathrm{d}^A B_A = 0, \dot{B}_A = -\mathrm{d}^A E$$

$$\left((q, p, \dot{q}, \dot{p}), \mathrm{d}_D^\dagger L(q, v) - \tilde{F}^\dagger(q, v)\right) \in D_{T^\dagger V}(q, p)$$

with $q = (A, \varphi), v = (v, \epsilon)$, and $p = (\varsigma, \varsigma_\partial, \alpha, \alpha_\partial)$

$$\begin{cases}\dot{E} - \delta^A B_A + \tilde{\varrho}_\varphi^*((\mathrm{d}^A \varphi)^{b_\kappa})^{\sharp_\partial} = -J, \\ \star_\partial \left(\iota_{\partial M}^*(\star\, B_A)\right) = j,\end{cases} \quad \begin{cases}\ddot{\varphi} + \delta^A(\, \mathrm{d}^A \varphi) + \mathrm{grad}_\kappa \mathbf{V} = \text{ב}, \\ \star_\partial \left(\iota_{\partial M}^*(\star\, \mathrm{d}^A \varphi)\right) = \text{ב}.\end{cases}$$

$\iota_{\partial M}\colon \partial M \to M$. Let $\pi_{E,M}\colon E \to M$

$$\pi_{E^*,M}\colon E^* \to M$$

$$\pi_{E,M}\colon E \to M$$

$$\eta \cdot \xi \in C^\infty(M) = \Omega^0(M).$$

$$\pi_{E',M}\colon E' \to M,$$

$\pi_{E,M}\colon E \to M$ and $\pi_{E',M}\colon E' \to M$ is denoted by $\pi_{E \otimes E',M}\colon E \otimes E' \to M$.

$$V = \Omega^k(M, E) = \Gamma\left(\textstyle\bigwedge^k T^*M \otimes E\right), 0 \le k \le m$$

$$\Lambda_r^s(M, E^*) = \Gamma(\textstyle\bigwedge^s TM \otimes \bigwedge^r T^*M \otimes E^*), 0 \le r, s \le m$$

$\Lambda_r^s(M) = \Gamma(\bigwedge^s TM \otimes \bigwedge^r T^*M)$ for the space of (r, s) symmetric tensor fields on M.

$$\begin{aligned}\chi \cdot \varphi &= (\tilde{\chi} \cdot \tilde{\varphi})(\eta \cdot \xi), \quad \varphi = \tilde{\varphi} \otimes \xi \in \Omega^k(M, E), \chi = \tilde{\chi} \otimes \eta \in \Lambda_m^k(M, E^*), \\ \chi \wedge \varphi &= (\tilde{\chi} \wedge \tilde{\varphi})(\eta \cdot \xi), \quad \varphi = \tilde{\varphi} \otimes \xi \in \Omega^k(M, E), \chi = \tilde{\chi} \otimes \eta \in \Omega^{m-k}(M, E^*).\end{aligned}$$

$$\wedge\colon \Omega^k(M, E) \times \Omega^{m-k}(M, E^*) \to \Omega^m(M)$$

$\pi_{\iota_{\partial M}^* E^*, \partial M}\colon \iota_{\partial M}^* E^* \to \partial M$, where $\iota_{\partial M}\colon \partial M \to M$ is the inclusion $E^* \equiv \iota_{\partial M}^* E^*$

$$V^{\star} = \Lambda_m^k(M, E^*) \times \Lambda_{m-1}^k(\partial M, E^*)$$

$$\langle(\alpha, \alpha_\partial), \varphi\rangle_{\star} = \int_M \alpha \cdot \varphi + \int_{\partial M} \alpha_\partial \cdot \iota_{\partial M}^* \varphi, (\alpha, \alpha_\partial) \in V^{\star}, \varphi \in V$$

$$V^{\dagger} = \Omega^{m-k}(M, E^*) \times \Omega^{m-k-1}(\partial M, E^*)$$

$$\langle(\alpha, \alpha_\partial), \varphi\rangle_{\dagger} = \int_M \varphi \wedge \alpha + \int_{\partial M} \iota_{\partial M}^* \varphi \wedge \alpha_\partial, (\alpha, \alpha_\partial) \in V^{\dagger}, \varphi \in V$$

$$V = \Omega^0(M, E) = \Gamma\left(\pi_{E,M}\right)$$

$$\Psi_{\star}: V^{\star} \to V' \text{ and } \Psi_{\dagger}: V^{\dagger} \to V'$$

$$\Psi_{\star}(\alpha, \alpha_\partial)(\varphi) = \langle(\alpha, \alpha_\partial), \varphi\rangle_{\star} \text{ and } \Psi_{\dagger}(\beta, \beta_\partial)(\varphi) = \langle(\beta, \beta_\partial), \varphi\rangle_{\dagger},$$

$$\varphi \in V, (\alpha, \alpha_\partial) \in V^{\star}, \text{and } (\beta, \beta_\partial) \in V^{\dagger}$$

$$U = U_1 \wedge \cdots \wedge U_k \in \bigwedge^k T_x M$$

$$i_U: \bigwedge^m T_x^* M \to \bigwedge^{m-k} T_x^* M, \mu \mapsto i_U \mu$$

where $(i_U\mu)(u_1, \ldots, u_{m-k}) = \mu(U_1, \ldots, U_k, u_1, \ldots, u_{m-k})$ for each $u_1, \ldots, u_{m-k} \in T_x M$

$$\alpha = \alpha_1 \wedge \cdots \wedge \alpha_k \in \bigwedge^k T_x^* M$$

$$i_\alpha: \bigwedge^m T_x M \to \bigwedge^{m-k} T_x M, U \mapsto i_\alpha U$$

where $(i_\alpha U)(\beta_1, \ldots, \beta_{m-k}) = U(\alpha_1, \ldots, \alpha_k, \beta_1, \ldots, \beta_{m-k})$ for each $\beta_1, \ldots, \beta_{m-k} \in T_x^* M$.

$$\Phi_E: \bigwedge^k TM \otimes \bigwedge^m T^*M \otimes E^* \to \bigwedge^{m-k} T^*M \otimes E^*, U \otimes \mu \otimes \eta \mapsto i_U\mu \otimes \eta.$$

$$\Phi_E^{-1}: \bigwedge^{m-k} T^*M \otimes E^* \to \bigwedge^k TM \otimes \bigwedge^m T^*M \otimes E^*, \alpha \otimes \eta \mapsto (-1)^{k(m-k)} (\star\, \alpha)^{\sharp_\partial} \otimes \mu_g \otimes \eta,$$

$\sharp: T^*M \to TM$ and $b: TM \to T^*M$ isomorphisms by the Riemannian metric, the Riemannian volume form $\mu_g \in \Omega^m(M)$ and $\star: \Omega^k(M) \to \Omega^{m-k}(M)$ is the Hodge star operator $g(\alpha, \beta)\mu_g = \alpha \wedge \star\, \beta$ for each $\alpha, \beta \in \Omega^k(M)$.

$\star\, \mu_g = (-1)^{m-1}$, yield $i_U \mu_g = (-1)^{(m-1)} \star U^{\mathrm{b}}$ for $U \in TM$

multivectors $i_U \mu_g = (-1)^{m-k} \star U^b$ for $U \in \bigwedge^k TM$

(i) $(\Phi_E^{-1} \circ \Phi_E)\left(U \otimes \mu_g \otimes \eta\right) = (-1)^{k(m-k)} \left(\star \left(i_U \mu_g\right)\right)^{\sharp_\partial} \otimes \mu_g \otimes \eta = (-1)^k \left(\star \left(\star\, U^b\right)\right)^{\sharp_\partial} \otimes$

$\mu_g \otimes \eta = \left(U^{\mathrm{b}}\right)^{\sharp_\partial} \otimes \mu_g \otimes \eta = U \otimes \mu_g \otimes \eta$, have used $\star\star = (-1)^{k(m-k)}$.

(ii) $(\Phi_E \circ \Phi_E^{-1})(\alpha \otimes \eta) = (-1)^{k(m-k)} i_{(\star\alpha)^{\sharp_\partial}} \mu_g \otimes \eta = \alpha \otimes \eta$ for each $\beta \in \bigwedge^{m-k} T^*M$

$$g\left(i_{(\star\alpha)^{\sharp_\partial}}\mu_g,\beta\right)=g\left(\mu_g,\star\,\alpha\wedge\beta\right)=(-1)^{k(m-k)}g\left(\mu_g,\alpha\wedge\star\,\beta\right)=g\left(\mu_g,\star\,\beta\wedge\alpha\right)$$
$$=g\left(i_{(\star\beta)^{\sharp_\partial}}\mu_g,\alpha\right)=(-1)^{m-k}g(\star\star\,\beta,\alpha)=(-1)^{k(m-k)}g(\alpha,\beta).$$

$$I=(i_1,\dots,i_k)\in\mathbb{N}^k,\text{ where }1\le i_1<\cdots<i_k\le m$$

$$\partial_I=\partial_{i_1}\wedge\cdots\wedge\partial_{i_k},dx^I=dx^{i_1}\wedge\cdots\wedge dx^{i_k},d^mx=dx^1\wedge\cdots\wedge dx^m,\boldsymbol{\partial}_m=\partial_1\wedge\cdots\wedge\partial_m\text{ and}$$

$$d_I^{m-k}x=i_{\partial_I}d^mx$$

$$\Phi_E^{-1}\colon\bigwedge^{m-k}T^*M\otimes E^*\to\bigwedge^k TM\otimes\bigwedge^m T^*M\otimes E^*,\alpha\otimes\eta\mapsto(-1)^{k(m-k)}(i_\alpha\boldsymbol{\partial}_m)\otimes d^mx\otimes\eta,$$

$$g_\partial=\imath_{\partial M}^*g$$

$$\Phi_{E,\partial}\colon\bigwedge^kT\partial M\otimes\bigwedge^{m-1}T^*\partial M\otimes E^*\to\bigwedge^{m-k-1}T^*\partial M\otimes E^*,U_\partial\otimes\mu_\partial\otimes\eta\mapsto i_{U_\partial}\mu_\partial\otimes\eta$$

$$\Phi_{E,\partial}^{-1}\colon\bigwedge^{m-k-1}T^*\partial M\otimes E^*\to\bigwedge^kT\partial M\otimes\bigwedge^{m-1}T^*\partial M\otimes E^*$$

$$\alpha_\partial\otimes\eta\mapsto(-1)^{k(m-k-1)}(\star_\partial\,\alpha_\partial)^{\sharp_\partial}\otimes\mu_g^\partial\otimes\eta,$$

$$\sharp_\partial\colon T^*\partial M\to T\partial M\text{ and }b_\partial\colon T\partial M\to T^*\partial M$$

$$\star_\partial\colon\Omega^k(\partial M)\to\Omega^{m-k-1}(\partial M)$$

$$\mu_g^\partial=\imath_{\partial M}^*\left(i_n\mu_g\right)\in\Omega^{m-1}(\partial M)$$

$$n\in\mathfrak{X}(M)|_{\partial M}$$

$$V^\star\to V^\dagger,(\alpha,\alpha_\partial)\mapsto\left(\Phi_E(\alpha),\Phi_{E,\partial}(\alpha_\partial)\right)$$
$$V^\dagger\to V^\star,(\beta,\beta_\partial)\mapsto\left(\Phi_E^{-1}(\beta),\Phi_{E,\partial}^{-1}(\beta_\partial)\right)$$

$$\Psi_\dagger\circ\left(\Phi_E,\Phi_{E,\partial}\right)=\Psi_\star\text{ and }\Psi_\star\circ\left(\Phi_E^{-1},\Phi_{E,\partial}^{-1}\right)=\Psi_\dagger$$

$$T^\star V=V\times V^\star=\Omega^k(M,E)\times\Lambda_m^k(M,E^*)\times\Lambda_{m-1}^k(\partial M,E^*)$$
$$T^\dagger V=V\times V^\dagger=\Omega^k(M,E)\times\Omega^{m-k}(M,E^*)\times\Omega^{m-k-1}(\partial M,E^*)$$

$$T^\star(TV)=V\times V\times V^\star\times V^\star,\quad T^\dagger(TV)=V\times V\times V^\dagger\times V^\dagger,$$
$$T(T^\star V)=V\times V^\star\times V\times V^\star,\quad T\left(T^\dagger V\right)=V\times V^\dagger\times V\times V^\dagger,$$
$$T^\star(T^\star V)=V\times V^\star\times V^\star\times V,\quad T^\dagger\left(T^\dagger V\right)=V\times V^\dagger\times V^\dagger\times V.$$

$$T(T^\star V)\oplus T^\star(T^\star V)=V\times V^\star\times(V\times V^\star\times V^\star\times V)\left(\subset T(T^\star V)\oplus T'(T^\star V)\right)$$
$$T\left(T^\dagger V\right)\oplus T^\dagger\left(T^\dagger V\right)=V\times V^\dagger\times\left(V\times V^\dagger\times V^\dagger\times V\right)\left(\subset T\left(T^\dagger V\right)\oplus T'\left(T^\dagger V\right)\right)$$

$V^\star=\Lambda_m^k(M,E^*)\times\Lambda_{m-1}^k(\partial M,E^*)$, the canonical form $\Theta_{T^\star V}\in\Omega^1\left(T^\star\Omega^k(M,E)\right)$, is defined

$$\Theta_{T^\star V}(z)\cdot\delta z=\langle z,T_z\pi_V^\star(\delta z)\rangle_\star,z\in T^\star V,\delta z\in T_z(T^\star V)$$

where $\pi_V^\star: T^\star V \to V$

$$\Omega_{T^\star V} = -\mathrm{d}\Theta_{T^\star V} \in \Omega^2\big(T^\star\Omega^k(M,E)\big)$$

$V^\dagger = \Omega^{m-k}(M,E^*) \times \Omega^{m-k-1}(\partial M, E^*)$, the canonical form $\Theta_{T^\dagger V} \in \Omega^1\big(T^\dagger\Omega^k(M,E)\big)$, is defined

$$\Theta_{T^\dagger V}(z)\cdot\delta z = \big\langle z, T_z\pi_V^\dagger(\delta z)\big\rangle_\dagger, z \in T^\dagger V, \delta z \in T_z\big(T^\dagger V\big)$$

where $\pi_V^\dagger: T^\dagger V \to V$

$$\Omega_{T^\dagger V} = -\mathrm{d}\Theta_{T^\dagger V} \in \Omega^2\big(T^\dagger\Omega^k(M,E)\big)$$

$$\Theta_{T^\star V}(\varphi,\alpha,\alpha_\partial)\cdot(\delta\varphi,\delta\alpha,\delta\alpha_\partial) = \langle(\varphi,\alpha,\alpha_\partial),\delta\varphi\rangle_\star = \int_M \alpha\cdot\delta\varphi + \int_{\partial M}\alpha_\partial\cdot\iota_{\partial M}^*\delta\varphi$$

$(\varphi,\alpha,\alpha_\partial) \in T^\star V$ and $(\delta\varphi,\delta\alpha,\delta\alpha_\partial) \in T_{(\varphi,\alpha,\alpha_\partial)}(T^\star V)$, and

$$\Theta_{T^\dagger V}(\varphi,\alpha,\alpha_\partial)\cdot(\delta\varphi,\delta\alpha,\delta\alpha_\partial) = \langle(\varphi,\alpha,\alpha_\partial),\delta\varphi\rangle_\dagger = \int_M \delta\varphi\wedge\alpha + \int_{\partial M}\iota_{\partial M}^*\delta\varphi\wedge\alpha_\partial$$

$(\varphi,\alpha,\alpha_\partial) \in T^\dagger V$ and $(\delta\varphi,\delta\alpha,\delta\alpha_\partial) \in T_{(\varphi,\alpha,\alpha_\partial)}\big(T^\dagger V\big)$

$$\Omega_{T^\star V}(\varphi,\alpha,\alpha_\partial)\big((\dot\varphi,\dot\alpha,\dot\alpha_\partial),(\delta\varphi,\delta\alpha,\delta\alpha_\partial)\big) = \langle(\delta\alpha,\delta\alpha_\partial),\dot\varphi\rangle_\star - \langle(\dot\alpha,\dot\alpha_\partial),\delta\varphi\rangle_\star,$$

$(\varphi,\alpha,\alpha_\partial) \in T^\star V$ and $(\delta\varphi,\delta\alpha,\delta\alpha_\partial),(\dot\varphi,\dot\alpha,\dot\alpha_\partial) \in T_{(\varphi,\alpha,\alpha_\partial)}(T^\star V) \simeq V\times V^\star$

$$\Omega_{T^\dagger V}(\varphi,\alpha,\alpha_\partial)\big((\dot\varphi,\dot\alpha,\dot\alpha_\partial),(\delta\varphi,\delta\alpha,\delta\alpha_\partial)\big) = \big\langle(\delta\alpha,\delta\alpha_\partial),\dot\varphi_\dagger - \langle(\dot\alpha,\dot\alpha_\partial),\delta\varphi\rangle_\dagger,$$

for each $(\varphi,\alpha,\alpha_\partial) \in T^\dagger V$ and $(\delta\varphi,\delta\alpha,\delta\alpha_\partial),(\dot\varphi,\dot\alpha,\dot\alpha_\partial) \in T_{\left(\varphi,\alpha,\alpha_\partial\right)}\big(T^\dagger V\big) \simeq V\times V^\dagger$

$$\Omega_{T^\star V}^\natural: T(T^\star V) \to T^\star(T^\star V), (\varphi,\alpha,\alpha_\partial,\dot\varphi,\dot\alpha,\dot\alpha_\partial) \mapsto (\varphi,\alpha,\alpha_\partial,-\dot\alpha,-\dot\alpha_\partial,\dot\varphi)$$

$$\Omega_{T^\dagger V}^\natural: T\big(T^\dagger V\big) \to T^\dagger\big(T^\dagger V\big), (\varphi,\alpha,\alpha_\partial,\dot\varphi,\dot\alpha,\dot\alpha_\partial) \mapsto (\varphi,\alpha,\alpha_\partial,-\dot\alpha,-\dot\alpha_\partial,\dot\varphi)$$

$V^\star = \Lambda_m^k(M,E^*) \times \Lambda_{m-1}^k(\partial M, E^*)$, the corresponding Tulczyjew

$$\begin{array}{ccccc} & & \gamma_{T^\star V} = \Omega_{T^\star V}^\flat \circ \kappa_{T^\star V}^{-1} & & \\ & \overset{\kappa_{T^\star V}}{\longleftarrow} & & \overset{\Omega_{T^\star V}^\flat}{\longrightarrow} & \\ T^\star(T\Omega^k(M,E)) & & T(T^\star\Omega^k(M,E)) & & T^\star(T^\star\Omega^k(M,E)) \\ (\varphi,\dot\varphi,\dot\alpha,\dot\alpha_\partial,\alpha,\alpha_\partial) & \longmapsto & (\varphi,\alpha,\alpha_\partial,\dot\varphi,\dot\alpha,\dot\alpha_\partial) & \longmapsto & (\varphi,\alpha,\alpha_\partial,-\dot\alpha,-\dot\alpha_\partial,\dot\varphi)\,. \end{array}$$

For $V^\dagger = \Omega^{m-k}(M,E^*) \times \Omega^{m-k-1}(\partial M, E^*)$

$T^\star V = \Omega^k(M,E) \times \Lambda_m^k(M,E^*) \times \Lambda_{m-1}^k(\partial M,E^*)$ is the subbundle $D_{T^\star V} = \text{graph}\Omega^b_{T^\star V}$. For each

$(\varphi,\alpha,\alpha_\partial) \in T^\star V$, it reads

$$D_{T^\star V}(\varphi,\alpha,\alpha_\partial) = \{(\dot\varphi,\dot\alpha,\dot\alpha_\partial,\delta\alpha,\delta\alpha_\partial,\delta\varphi) \in T_{(\varphi,\alpha,\alpha_\partial)}(T^\star V) \times T^\star_{(\varphi,\alpha,\alpha_\partial)}(T^\star V) \big| \\ -\dot\alpha = \delta\alpha, -\dot\alpha_\partial = \delta\alpha_\partial, \dot\varphi = \delta\varphi\}.$$

canonical Dirac $T^\dagger V = \Omega^k(M,E) \times \Omega^{m-k}(M,E^*) \times \Omega^{m-k-1}(\partial M,E^*)$ is the subbundle $D_{T^\dagger V} =$

$\text{graph}\Omega^b_{T^\dagger\Omega^k(M,E)}$. For each $(\varphi,\alpha,\alpha_\partial) \in T^\dagger V$, it reads

$$D_{T^\dagger V}(\varphi,\alpha,\alpha_\partial) = \left\{(\dot\varphi,\dot\alpha,\dot\alpha_\partial,\delta\alpha,\delta\alpha_\partial,\delta\varphi) \in T_{\left(\varphi,\alpha,\alpha_\partial\right)}\left(T^\dagger V\right) \times T^\dagger_{\left(\varphi,\alpha,\alpha_\partial\right)}\left(T^\dagger V\right) \right| \\ -\dot\alpha = \delta\alpha, -\dot\alpha_\partial = \delta\alpha_\partial, \dot\varphi = \delta\varphi\}$$

$$\text{tr}\colon \Lambda^{k+1}TM \otimes \Lambda^m T^*M \otimes E^* \to \Lambda^k TM \otimes \Lambda^{m-1}T^*M \otimes E^*, U \otimes \mu \otimes \eta \mapsto \sum_{i=1}^{k+1} \hat U^i \otimes i_{U_i}\mu \otimes \eta,$$

$U = U_1 \wedge \cdots \wedge U_{k+1} \in \Lambda^{k+1} TM, \mu \in T^*M$ and $\hat U^r = U_1 \wedge \cdots \wedge U_{r-1} \wedge U_{r+1} \wedge \cdots \wedge U_{k+1} \in \Lambda^k TM.$

In local coordinates, for $\chi \otimes \eta \in \Lambda_m^{k+1}(M,E^*)$ with $\chi = \chi^J \partial_J \otimes d^m x$, it reads

$$\text{tr}(\chi \otimes \eta) = \sum_{r=1}^{k+1} \chi^J \partial_{J_r} \otimes d^{m-1}_{(j_r)} x \otimes \eta$$

$$\chi = \chi^i \partial_i \otimes d^m x$$

$$\text{tr}(\chi \otimes \eta) = \chi^i d_i^{m-1} x \otimes \eta = i_{\partial_i}\chi\left(dx^i, \dots\right) \otimes \eta$$

$$\text{d}^\nabla\colon \Omega^s(M,E) \to \Omega^{s+1}(M,E), \text{d}^{\nabla^*}\colon \Omega^s(M,E^*) \to \Omega^{s+1}(M,E^*)$$

$$\Lambda_s^k(M,E^*) \to \Lambda_{s+1}^k(M,E^*)$$

$$\text{d}^{\nabla^*}(U \otimes \alpha) = U \otimes \ \text{d}^{\nabla^*}\alpha, U \in \Gamma\left(\Lambda^k TM\right), \alpha \in \Omega^s(M,E^*)$$

$$\text{div}^{\nabla^*}\colon \Lambda_m^{k+1}(M,E^*) \mapsto \Lambda_m^k(M,E^*), \chi \mapsto \text{d}^{\nabla^*}(\text{tr}\chi)$$

$$\text{div}^{\nabla^*}(\xi) = \sum_{i=1}^{k+1} \hat U^i \otimes \ \text{d}^{\nabla^*}\left(i_{U_i}\mu \otimes \eta\right), \xi = U \otimes \mu \otimes \eta \in \Lambda_m^{k+1}(M,E^*)$$

$$U = U_1 \wedge \dots \wedge U_{k+1}$$

$$\text{div}\colon \Lambda_m^{k+1}(M) \to \Lambda_m^k(M), \chi \mapsto \text{d}(\text{tr}\chi)$$

$$\alpha = \alpha^I \partial_I \otimes d^m x \in \Lambda_m^k(M) \text{ and } \varphi = \varphi_I dx^I \in \Omega^k(M)$$

$$\alpha \cdot \varphi = \alpha^I \varphi_I (\partial_I \cdot dx^I) \otimes d^m x = \alpha^I \varphi_I d^m x$$

$$\chi = \chi^J \partial_J \otimes d^m x \in \Lambda_m^{k+1}(M)$$

$$\chi \cdot \varphi = \chi^J \varphi_I \left(\partial_J \cdot dx^I\right) \otimes d^m x = (-1)^k \sum_{r=1}^{k+1} \chi^J \varphi_{J_r} \partial_{j_r} \otimes d^m x$$

$$\partial_J \cdot dx^I = (-1)^k \partial_{j_r}$$

$$\delta\varphi \in \Omega^k(M, E) \text{ and } \chi \in \Lambda_m^{k+1}(M, E^*)$$

$$\mathrm{div}(\chi \cdot \delta\varphi) = \chi \cdot \mathrm{d}^\nabla \delta\varphi + (-1)^k \left(\mathrm{div}^{\nabla^*} \chi\right) \cdot \delta\varphi$$

$$\int_M \chi \cdot \ \mathrm{d}^\nabla \delta\varphi = -(-1)^k \int_M \left(\mathrm{div}^{\nabla^*} \chi\right) \cdot \delta\varphi + \int_{\partial M} \iota_{\partial M}^* (\mathrm{tr}\chi) \cdot \iota_{\partial M}^* \delta\varphi$$

$$\delta\varphi = (\delta\varphi)_I^a dx^I \otimes B_a, \chi = \chi_a^J \partial_J \otimes d^m x \otimes B^a$$

$$(\delta\varphi)_I^a, \chi_a^J \in C^\infty(M), \text{ where } |I| = k, |J| = k+1 \text{ and } 1 \le a \le n$$

$$\mathrm{d}^\nabla \delta\varphi = \left(\partial_i (\delta\varphi)_I^a + \Gamma_{i,b}^a (\delta\varphi)_I^b\right) dx^i \wedge dx^I \otimes B_a$$

$$\Gamma_{i,b}^a \in C^\infty(M), 1 \le i \le m, 1 \le a, b \le n$$

$$\mathrm{div}^{\nabla^*} \chi = \mathrm{d}^{\nabla^*} \left(\sum_{r=1}^{k+1} \chi_a^J \partial_{J_r} \otimes d_{(j_r)}^{m-1} x \otimes B^a \right) = \sum_{r=1}^{k+1} \left(\partial_{j_r} \chi_a^J - \Gamma_{j_r,a}^b \chi_b^J\right) \partial_{J_r} \otimes d^m x \otimes B^a$$

$$-\Gamma_{i,a}^b, 1 \le i \le m, 1 \le a, b \le m$$

$$\chi \cdot \mathrm{d}^\nabla \delta\varphi = (-1)^k \sum_{r=1}^{k+1} \chi_a^J \left(\partial_{j_r} (\delta\varphi)_{J_r}^a + \Gamma_{j_r,b}^a (\delta\varphi)_{J_r}^b\right) d^m x$$

$$(\mathrm{div}\nabla^* \chi) \cdot \delta\varphi = \sum_{r=1}^{k+1} \left(\partial_{j_r} \chi_a^J - \Gamma_{j_r,a}^b \chi_b^J\right) (\delta\varphi)_{J_r}^a d^m x$$

$$\chi \cdot \delta\varphi = (-1)^k \sum_{r=1}^{k+1} \chi_a^J (\delta\varphi)_{J_r}^a \partial_{j_r} \otimes d^m x, \mathrm{tr}(\chi \cdot \delta\varphi) = \sum_{r=1}^{k+1} \chi_a^J (\delta\varphi)_{J_r}^a d_{(j_r)}^{m-1} x$$

$$\begin{aligned}(-1)^k \chi \cdot \ \mathrm{d}^\nabla \delta\varphi + \left(\mathrm{div}^{\nabla^*} \chi\right) \cdot \delta\varphi &= \sum_{r=1}^{k+1} \left(\chi_a^J \partial_{j_r} (\delta\varphi)_{J_r}^a + \left(\partial_{j_r} \chi_a^J\right) (\delta\varphi)_{J_r}^a\right) d^m x \\ &= \sum_{r=1}^{k+1} \partial_{j_r} \left(\chi_a^J (\delta\varphi)_{J_r}^a\right) d^m x \\ &= (-1)^k \mathrm{div}(\chi \cdot \delta\varphi)\end{aligned}$$

$$\mathrm{tr}(\chi \cdot \delta\varphi) = \sum_{r=1}^{k+1} \chi_a^J (\delta\varphi)_{J_r}^a d_{(j_r)}^{m-1} x = (\mathrm{tr}\chi) \cdot \delta\varphi$$

$$\iota_{\partial M}^* ((\mathrm{tr}\chi) \cdot \delta\varphi) = \iota_{\partial M}^* (\mathrm{tr}\chi) \cdot \iota_{\partial M}^* \delta\varphi$$

$$\delta\varphi \in \Omega^k(M,E), \zeta \in \Lambda_m^k(M,E^*) \text{ and } \chi \in \Lambda_m^{k+1}(M,E^*)$$

$$\zeta \cdot \delta\varphi = \delta\varphi \wedge \Phi_E(\zeta), \left(\mathrm{div}^{\nabla^*}\chi\right) \cdot \delta\varphi = \delta\varphi \wedge \mathrm{d}^{\nabla^*}\Phi_E(\chi)$$
$$(\mathrm{tr}\chi) \cdot \delta\varphi = \delta\varphi \wedge \Phi_E(\chi).$$

$$\delta\varphi = (\delta\varphi)_I^a dx^I \otimes B_a, \zeta = \zeta_a^I \partial_I \otimes d^m x \otimes B^a, \chi = \chi_a^J \partial_J \otimes d^m x \otimes B^a$$

$$(\delta\varphi)_I^a, \zeta_a^I, \chi_a^J \in C^\infty(M), \text{ where } |I| = k, |J| = k+1, \text{ and } B^a \mid 1 \le a \le n\}$$

$$\mathrm{div}\nabla^*\chi = \sum_{r=1}^{k+1} \left(\partial_{j_r}\chi_a^J - \Gamma_{j_r,a}^b \chi_b^J\right)\partial_{J_r} \otimes d^m x \otimes B^a, \qquad \Phi_E(\chi) = \chi_a^J d_J^{m-k-1} x \otimes B^a$$

$$\mathrm{d}^{\nabla^*}\Phi_E(\chi) = \sum_{r=1}^{k+1} \left(\partial_{j_r}\chi_a^J - \Gamma_{j_r,a}^b \chi_b^J\right) d_{J_r}^{m-k} x \otimes B^a, \quad \mathrm{tr}\chi = \sum_{r=1}^{k+1} \chi_a^J \partial_{J_r} \otimes d_{(j_r)}^{m-1} x \otimes B^a$$

$\Gamma_{i,b}^a \in C^\infty(M), 1 \le i \le m, 1 \le a, b \le n$ are the Christoffel symbols of ∇

$$\delta\varphi \wedge \Phi_E(\zeta) = (\delta\varphi)_{I'}^a \zeta_a^I dx^{I'} \wedge d_I^{m-k} x = (\delta\varphi)_I^a \zeta_a^I d^m x = \zeta \cdot \delta\varphi$$

$$\delta\varphi \wedge \mathrm{d}^{\nabla^*}\Phi_E(\chi) = \sum_{j=1}^{k+1} \left(\partial_{j_r}\chi_a^J - \Gamma_{j_r,a}^b \chi_b^J\right)(\delta\varphi)_I^a dx^I \wedge d_{J_r}^{m-k} x$$
$$= \sum_{r=1}^{k+1} \left(\partial_{j_r}\chi_a^J - \Gamma_{j_r,a}^b \chi_b^J\right)(\delta\varphi)_{J_r}^a d^m x = \left(\mathrm{div}^{\nabla^*}\chi\right) \cdot \delta\varphi$$

$$\delta\varphi \wedge \Phi_E(\chi) = \chi_a^J (\delta\varphi)_I^a dx^I \wedge d_J^{m-k-1} x$$
$$= \sum_{r=1}^{k+1} \chi_a^J (\delta\varphi)_{J_r}^a d_{(j_r)}^{m-1} x = (\mathrm{tr}\chi) \cdot \delta\varphi$$

$$L_\nabla : TV \to \mathbb{R}, L_\nabla(\varphi, \nu) = \int_M \mathcal{L}\left(\varphi, \nu, \mathrm{d}^\nabla\varphi\right)$$

$$\mathcal{L} : W^k(M,E) := \left(\Lambda^k T^*M \otimes E\right) \times_M \left(\Lambda^k T^*M \otimes E\right) \times_M \left(\Lambda^{k+1} T^*M \otimes E\right) \to \Lambda^{m+1} T^*M$$

$$\frac{\partial\mathcal{L}}{\partial\varphi} : W^k(M,E) \to \Lambda^k TM \otimes \Lambda^m T^*M \otimes E^*,$$
$$\frac{\partial\mathcal{L}}{\partial\nu} : W^k(M,E) \to \Lambda^k TM \otimes \Lambda^m T^*M \otimes E^*,$$
$$\frac{\partial\mathcal{L}}{\partial\zeta} : W^k(M,E) \to \Lambda^{k+1} TM \otimes \Lambda^m T^*M \otimes E^*,$$

$$x \in M \text{ and } (\varphi_x, \nu_x, \zeta_x) \in W^k(M,E)_x$$

$$\frac{\partial\mathcal{L}}{\partial\varphi}(\varphi_x, \nu_x, \zeta_x) \cdot \delta\varphi_x = \frac{d}{d\epsilon}\Big|_{\epsilon=0} \mathcal{L}(\varphi_x + \epsilon\delta\varphi_x, \nu_x, \zeta_x), \quad \delta\varphi_x \in \Lambda^k T_x^*M \otimes E_x$$
$$\frac{\partial\mathcal{L}}{\partial\nu}(\varphi_x, \nu_x, \zeta_x) \cdot \delta\nu_x = \frac{d}{d\epsilon}\Big|_{\epsilon=0} \mathcal{L}(\varphi_x, \nu_x + \epsilon\delta\nu_x, \zeta_x), \quad \delta\nu_x \in \Lambda^k T_x^*M \otimes E_x$$
$$\frac{\partial\mathcal{L}}{\partial\zeta}(\varphi_x, \nu_x, \zeta_x) \cdot \delta\zeta_x = \frac{d}{d\epsilon}\Big|_{\epsilon=0} \mathcal{L}(\varphi_x, \nu_x, \zeta_x + \epsilon\delta\zeta_x), \quad \delta\zeta_x \in \Lambda^{k+1} T_x^*M \otimes E_x$$

$$\frac{\partial \mathcal{L}}{\partial \varphi}: W^k(M,E) \to \Lambda^{m-k} T^*M \otimes E^*,$$
$$\frac{\partial \mathcal{L}}{\partial \nu}: W^k(M,E) \to \Lambda^{m-k} T^*M \otimes E^*,$$
$$\frac{\partial \mathcal{L}}{\partial \zeta}: W^k(M,E) \to \Lambda^{m-k-1} T^*M \otimes E^*.$$

$$\delta\varphi_x \wedge \frac{\partial \mathcal{L}}{\partial \varphi}(\varphi_x, \nu_x, \zeta_x) = \frac{d}{d\epsilon}\Big|_{\epsilon=0} \mathcal{L}(\varphi_x + \epsilon\delta\varphi_x, \nu_x, \zeta_x), \delta\varphi_x \in \Lambda^k T_x^*M \otimes E_x$$
$$\delta\nu_x \wedge \frac{\partial \mathcal{L}}{\partial \nu}(\varphi_x, \nu_x, \zeta_x) = \frac{d}{d\epsilon}\Big|_{\epsilon=0} \mathcal{L}(\varphi_x, \nu_x + \epsilon\delta\nu_x, \zeta_x), \delta\nu_x \in \Lambda^k T_x^*M \otimes E_x$$
$$\delta\zeta_x \wedge \frac{\partial \mathcal{L}}{\partial \zeta}(\varphi_x, \nu_x, \zeta_x) = \frac{d}{d\epsilon}\Big|_{\epsilon=0} \mathcal{L}(\varphi_x, \nu_x, \zeta_x + \epsilon\delta\zeta_x), \delta\zeta_x \in \Lambda^{k+1} T_x^*M \otimes E_x$$

$$\frac{\partial \mathcal{L}}{\partial \varphi} = \Phi_E \circ \frac{\partial \mathcal{L}}{\partial \varphi}, \frac{\partial \mathcal{L}}{\partial \nu} = \Phi_E \circ \frac{\partial \mathcal{L}}{\partial \nu}, \frac{\partial \mathcal{L}}{\partial \zeta} = \Phi_E \circ \frac{\partial \mathcal{L}}{\partial \zeta}.$$

$$\frac{\delta L}{\delta \varphi}, \frac{\delta L}{\delta \nu}: TV = T\Omega^k(M,E) \to V'$$

$$\frac{\delta L}{\delta \varphi}(\varphi,\nu)(\delta\varphi) = \frac{d}{d\epsilon}\Big|_{\epsilon=0} L(\varphi + \epsilon\delta\varphi, \nu), \frac{\delta L}{\delta \nu}(\varphi,\nu)(\delta\nu) = \frac{d}{d\epsilon}\Big|_{\epsilon=0} L(\varphi, \nu + \epsilon\delta\nu)$$

$$\mathrm{d}L: TV \to T'(TV), (\varphi,\nu) \mapsto \left(\varphi, \nu, \frac{\delta L}{\delta \varphi}(\varphi,\nu), \frac{\delta L}{\delta \nu}(\varphi,\nu)\right)$$

$$\frac{\delta L_\nabla}{\delta \varphi}(\varphi,\nu) = \left(\frac{\partial \mathcal{L}}{\partial \varphi}(\varphi,\nu,\mathrm{d}^\nabla\varphi) - (-1)^k \mathrm{div}^{\nabla^*}\left(\frac{\partial \mathcal{L}}{\partial \zeta}(\varphi,\nu,\ \mathrm{d}^\nabla\varphi)\right), \iota_{\partial M}^*\left(\mathrm{tr}\frac{\partial \mathcal{L}}{\partial \zeta}(\varphi,\nu,\ \mathrm{d}^\nabla\varphi)\right)\right)$$
$$\frac{\delta L_\nabla}{\delta \nu}(\varphi,\nu) = \left(\frac{\partial \mathcal{L}}{\partial \nu}(\varphi,\nu,\mathrm{d}^\nabla\varphi), 0\right)$$

$$\frac{\delta L_\nabla}{\delta \varphi}(\varphi,\nu) = \left(\frac{\partial \mathcal{L}}{\partial \varphi}(\varphi,\nu,\mathrm{d}^\nabla\varphi) - (-1)^k\ \mathrm{d}^{\nabla^*}\left(\frac{\partial \mathcal{L}}{\partial \zeta}(\varphi,\nu,\ \mathrm{d}^\nabla\varphi)\right), \iota_{\partial M}^*\left(\frac{\partial \mathcal{L}}{\partial \zeta}(\varphi,\nu,\ \mathrm{d}^\nabla\varphi)\right)\right)$$
$$\frac{\delta L_\nabla}{\delta \nu}(\varphi,\nu) = \left(\frac{\partial \mathcal{L}}{\partial \nu}(\varphi,\nu,\mathrm{d}^\nabla\varphi), 0\right)$$

$$\begin{aligned}
\frac{\delta L_\nabla}{\delta \varphi}(\varphi,\nu)(\delta\varphi) &= \frac{d}{d\epsilon}\Big|_{\epsilon=0} \int_M \mathcal{L}\big(\varphi + \epsilon\delta\varphi, \nu, \mathrm{d}^\nabla(\varphi + \epsilon\delta\varphi)\big) \\
&= \int_M \left(\frac{\partial \mathcal{L}}{\partial \varphi}(\varphi,\nu,\ \mathrm{d}^\nabla\varphi)\cdot\delta\varphi + \frac{\partial \mathcal{L}}{\partial \zeta}(\varphi,\nu,\mathrm{d}^\nabla\varphi)\cdot\mathrm{d}^\nabla\delta\varphi\right) \\
&= \int_M \left(\frac{\partial \mathcal{L}}{\partial \varphi}(\varphi,\nu,\ \mathrm{d}^\nabla\varphi) - (-1)^k \mathrm{div}^\nabla\left(\frac{\partial \mathcal{L}}{\partial \zeta}(\varphi,\nu,\mathrm{d}^\nabla\varphi)\right)\right)\cdot\delta\varphi \\
&\quad + \int_{\partial M} \iota_{\partial M}^*\left(\mathrm{tr}\frac{\partial \mathcal{L}}{\partial \zeta}(\varphi,\nu,\ \mathrm{d}^\nabla\varphi)\right)\cdot\iota_{\partial M}^*\delta\varphi \\
\frac{\delta L_\nabla}{\delta \nu}(\varphi,\nu)(\delta\nu) &= \frac{d}{d\epsilon}\Big|_{\epsilon=0} \int_M \mathcal{L}\big(\varphi, \nu + \epsilon\delta\nu, \mathrm{d}^\nabla\varphi\big) = \int_M \frac{\partial \mathcal{L}}{\partial \nu}(\varphi,\nu,\ \mathrm{d}^\nabla\varphi)\cdot\delta\nu
\end{aligned}$$

$$\frac{\delta L_\nabla}{\delta \varphi}(\varphi,\nu)(\delta\varphi)=\int_M \delta\varphi\wedge\left(\frac{\partial \mathcal{L}}{\partial \varphi}\left(\varphi,\nu,\ \mathrm{d}^\nabla\varphi\right)-(-1)^k\ \mathrm{d}^{\nabla^*}\left(\frac{\partial \mathcal{L}}{\partial \zeta}\left(\varphi,\nu,\ \mathrm{d}^\nabla\varphi\right)\right)\right)$$

$$+\int_{\partial M}\iota_{\partial M}^*\delta\varphi\wedge\iota_{\partial M}^*\left(\frac{\partial \mathcal{L}}{\partial \zeta}\left(\varphi,\nu,\ \mathrm{d}^\nabla\varphi\right)\right)$$

$$\frac{\delta L_\nabla}{\delta \nu}(\varphi,\nu)(\delta\nu)=\int_M \delta\nu\wedge\frac{\partial \mathcal{L}}{\partial \nu}\left(\varphi,\nu,\ \mathrm{d}^\nabla\varphi\right)$$

$$\mathrm{d}L_\nabla: TV\to T'(TV),(\varphi,\nu)\mapsto\left(\varphi,\nu,\frac{\delta L_\nabla}{\delta\varphi}(\varphi,\nu),\frac{\delta L_\nabla}{\delta\nu}(\varphi,\nu)\right)$$

$$\mathrm{d}_D^\star L_\nabla=\gamma_{T^\star V}\circ\ \mathrm{d}L_\nabla: TV\to T^\star(T^\star V),(\varphi,\nu)\mapsto\left(\varphi,\frac{\delta L_\nabla}{\delta\nu}(\varphi,\nu),-\frac{\delta L_\nabla}{\delta\varphi}(\varphi,\nu),\nu\right)$$

$$\mathrm{d}_D^\dagger L_\nabla=\gamma_{T^\dagger V}\circ\ \mathrm{d}L_\nabla: TV\to T^\dagger\left(T^\dagger V\right),(\varphi,\nu)\mapsto\left(\varphi,\frac{\delta L_\nabla}{\delta\nu}(\varphi,\nu),-\frac{\delta L_\nabla}{\delta\varphi}(\varphi,\nu),\nu\right)$$

$$\mathbb{F}L_\nabla: TV\to T'V,(\varphi,\nu)\mapsto\mathbb{F}L_\nabla(\varphi,\nu)=\left(\varphi,\frac{\delta L_\nabla}{\delta\nu}(\varphi,\nu)\right)$$

$$V^\star=\Lambda_m^k(M,E^*)\times\Lambda_{m-1}^k(\partial M,E^*)$$

$$\mathbb{F}^\star L_\nabla: TV\to T^\star V,(\varphi,\nu)\mapsto\left(\varphi,\frac{\partial \mathcal{L}}{\partial\nu}\left(\varphi,\nu,\mathrm{d}^\nabla\varphi\right),0\right)$$

$$V^\dagger=\Omega^{m-k}(M,E^*)\times\Omega^{m-k-1}(\partial M,E^*)$$

$$\mathbb{F}^\dagger L_\nabla: TV\to T^\dagger V,(\varphi,\nu)\mapsto\left(\varphi,\frac{\partial \mathcal{L}}{\partial\nu}\left(\varphi,\nu,\mathrm{d}^\nabla\varphi\right),0\right)$$

$$V^\star=\Lambda_m^k(M,E^*)\times\Lambda_{m-1}^k(\partial M,E^*)$$

$$F^\star: TV\to T^\star V,(\varphi,\nu)\mapsto F^\star(\varphi,\nu)=(\varphi,\mathcal{F}^\star(\varphi,\nu),\mathcal{F}_\partial^\star(\varphi,\nu)),$$

$$\mathcal{F}^\star: TV\to\Lambda_m^k(M,E^*)\text{ and }\mathcal{F}_\partial^\star: TV\to\Lambda_{m-1}^k(\partial M,E^*)$$

$$\tilde{F}^\star: TV\to T^\star(T^\star V)$$

$$\left\langle\tilde{F}^\star(\varphi,\nu),W\right\rangle_\star=\left\langle F^\star(\varphi,\nu),T_{\mathbb{F}^\star L_\nabla(\varphi,\nu)}\pi_V^\star(W)\right\rangle_\star,(\varphi,\nu)\in TV,W\in T_{\mathbb{F}^\star L(\varphi,\nu)}(T^\star V)$$

$\pi_V^\star: T^\star V\to V$ is the natural projection and $T\pi_V^\star: T(T^\star V)\to TV$ is the tangent map

$$\tilde{F}^\star(\varphi,\nu)=\left(\varphi,\frac{\partial \mathcal{L}}{\partial\nu}\left(\varphi,\nu,\mathrm{d}^\nabla\varphi\right),0,\mathcal{F}^\star(\varphi,\nu),\mathcal{F}_\partial^\star(\varphi,\nu),0\right).$$

$$V^\dagger=\Omega^{m-k}(M,E^*)\times\Omega^{m-k-1}(\partial M,E^*)$$

$$F^\dagger: TV\to T^\dagger V,(\varphi,\nu)\mapsto F^\dagger(\varphi,\nu)=\left(\varphi,\mathcal{F}^\dagger(\varphi,\nu),\mathcal{F}_\partial^\dagger(\varphi,\nu)\right),$$

$$\mathcal{F}^\dagger: TV\to\Omega^{m-k}(M,E^*)\text{ and }\mathcal{F}_\partial^\dagger: TV\to\Omega^{m-k-1}(\partial M,E^*)$$

$$\tilde{F}^{\dagger}: TV \to T^{\dagger}\left(T^{\dagger}V\right)$$

$$\left\langle \tilde{F}^{\dagger}(\varphi,\nu), W\right\rangle_{\dagger} = \left\langle F^{\dagger}(\varphi,\nu), T_{\mathbb{F}^{\dagger}L_{\nabla}(\varphi,\nu)}\pi_V^{\dagger}(W)\right\rangle_{\dagger}, (\varphi,\nu) \in TV, W \in T_{\mathbb{F}^{\dagger}L(\varphi,\nu)}\left(T^{\dagger}V\right)$$

where $\pi_V^{\dagger}: T^{\dagger}V \to V$ is the natural projection and $T\pi_V^{\dagger}: T\left(T^{\dagger}V\right) \to TV$ denotes the tangent map.

$$\tilde{F}^{\dagger}(\varphi,\nu) = \left(\varphi, \frac{\partial \mathcal{L}}{\partial \nu}\left(\varphi,\nu,\mathrm{d}^{\nabla}\varphi\right), 0, \mathcal{F}^{\dagger}(\varphi,\nu), \mathcal{F}_{\partial}^{\dagger}(\varphi,\nu), 0\right).$$

$$V^{\star} = \Lambda_m^k(M,E^*) \times \Lambda_{m-1}^k(\partial M, E^*)$$

$$V = \Omega^k(M,E), D_{T^{\star}V}, L_{\nabla}, F^{\star})$$

$$(\varphi,\nu,\alpha,\alpha_{\partial}): [t_0,t_1] \to TV \oplus T^{\star}V$$

$$\left((\varphi,\alpha,\alpha_{\partial},\dot{\varphi},\dot{\alpha},\dot{\alpha}_{\partial}), \mathrm{d}_D^{\star}L_{\nabla}(\varphi,\nu) - \tilde{F}^{\star}(\varphi,\nu)\right) \in D_{T^{\star}V}(\varphi,\alpha,\alpha_{\partial}).$$

$$V^{\dagger} = \Omega^{m-k}(M,E^*) \times \Omega^{m-k-1}(\partial M, E^*)$$

$$V = \Omega^k(M,E), D_{T^{\dagger}V}, L_{\nabla}, F^{\dagger}$$

$$(\varphi,\nu,\alpha,\alpha_{\partial}): [t_0,t_1] \to TV \oplus T^{\dagger}V$$

$$\left((\varphi,\alpha,\alpha_{\partial},\dot{\varphi},\dot{\alpha},\dot{\alpha}_{\partial}), \mathrm{d}_D^{\dagger}L_{\nabla}(\varphi,\nu) - \tilde{F}^{\dagger}(\varphi,\nu)\right) \in D_{T^{\dagger}V}(\varphi,\alpha,\alpha_{\partial})$$

$V^{\star} = \Lambda_m^k(M,E^*) \times \Lambda_{m-1}^k(\partial M, E^*)$, a curve $(\varphi,\nu,\alpha,\alpha_{\partial}): [t_0,t_1] \to TV \oplus T^{\star}V$

$$\begin{cases} \dot{\varphi} = \nu, \\ \alpha = \dfrac{\partial \mathcal{L}}{\partial \nu}\left(\varphi,\nu,\mathrm{d}^{\nabla}\varphi\right), \quad \dot{\alpha} = \dfrac{\partial \mathcal{L}}{\partial \varphi}\left(\varphi,\nu,\mathrm{d}^{\nabla}\varphi\right) - (-1)^k \mathrm{div}^{\nabla^*}\left(\dfrac{\partial \mathcal{L}}{\partial \zeta}\left(\varphi,\nu,\ \mathrm{d}^{\nabla}\varphi\right)\right) + \mathcal{F}^{\star}(\varphi,\nu) \\ \alpha_{\partial} = 0, \qquad\qquad \dot{\alpha}_{\partial} = \iota_{\partial M}^{*}\left(\mathrm{tr}\dfrac{\partial \mathcal{L}}{\partial \zeta}\left(\varphi,\nu,\ \mathrm{d}^{\nabla}\varphi\right)\right) + \mathcal{F}_{\partial}^{\star}(\varphi,\nu) \end{cases}$$

$V^{\dagger} = \Omega^{m-k}(M,E^*) \times \Omega^{m-k-1}(\partial M, E^*)$, a curve $(\varphi,\nu,\alpha,\alpha_{\partial}): [t_0,t_1] \to TV \oplus T^{\dagger}V$

$$\begin{cases} \dot{\varphi} = \nu \\ \alpha = \dfrac{\partial \mathcal{L}}{\partial \nu}\left(\varphi,\nu,\mathrm{d}^{\nabla}\varphi\right), \quad \dot{\alpha} = \dfrac{\partial \mathcal{L}}{\partial \varphi}\left(\varphi,\nu,\mathrm{d}^{\nabla}\varphi\right) - (-1)^k\ \mathrm{d}^{*}\left(\dfrac{\partial \mathcal{L}}{\partial \zeta}\left(\varphi,\nu,\ \mathrm{d}^{\nabla}\varphi\right)\right) + \mathcal{F}^{\dagger}(\varphi,\nu) \\ \alpha_{\partial} = 0, \qquad\qquad \dot{\alpha}_{\partial} = \iota_{\partial M}^{*}\left(\dfrac{\partial \mathcal{L}}{\partial \zeta}\left(\varphi,\nu,\ \mathrm{d}^{\nabla}\varphi\right)\right) + \mathcal{F}_{\partial}^{\dagger}(\varphi,\nu) \end{cases}$$

$$\partial M = \{(x^1,\dots,x^m) \in \mathbb{R}^m \mid x^m = 0\}.$$

$$\Gamma_{i,b}^{a} \in C^{\infty}(M), 1 \le i \le m, 1 \le a,b \le m$$

$$\mathcal{I} = \{(J,r) \in \mathbb{N}^{k+1} \times \{1,\dots,k+1\} \mid 1 \le j_1 < \cdots < j_{k+1} \le m, J_r = I\}.$$

$$\mathcal{I}_m = \{I \in \mathbb{N}^k \mid 1 \le i_1 < \cdots < i_k < m\}$$

$$I^m = (i_1, \dots, i_k, m) \in \mathbb{N}^{k+1} \text{ for each } I \in \mathcal{I}_m$$

$$\alpha = \alpha_a^I \partial_I \otimes d^m m \otimes B^a \in \Lambda_m^k(M, E^*), \alpha = \alpha_a^I d_I^{m-k} x \otimes B^a \in \Omega^{m-k}(M, E^*).$$

$$\begin{cases} \dot{\varphi}_I^a = v_I^a, \\ \alpha_a^I = \dfrac{\partial \mathcal{L}}{\partial v_I^a}, \quad \dot{\alpha}_a^I = \dfrac{\partial \mathcal{L}}{\partial \varphi_I^a} - (-1)^k \left(\displaystyle\sum_{(J,r)\in\mathcal{J}} \partial_{j_r} \frac{\partial \mathcal{L}}{\partial \zeta_J^a} - \Gamma_{j_r,a}^b \frac{\partial \mathcal{L}}{\partial \zeta_J^b} \right) + \mathcal{F}_a^I, \\ (\alpha_\partial)_a^I = 0, \quad (\dot{\alpha}_\partial)_a^I = \left. \dfrac{\partial \mathcal{L}}{\partial \zeta_{I^m}^a} \right|_{x^m=0} + (\mathcal{F}_\partial)_a^I, I \in \mathcal{I}_m. \end{cases}$$

$$\begin{cases} \dot{\varphi}_i^a = v_i^a, \\ \alpha_a^i = \dfrac{\partial \mathcal{L}}{\partial v_i^a}, \quad \dot{\alpha}_a^i = \dfrac{\partial \mathcal{L}}{\partial \varphi_i^a} + \displaystyle\sum_{j=1}^m \epsilon(i,j) \left(\partial_j \frac{\partial \mathcal{L}}{\partial \zeta_{(i,j)}^a} - \Gamma_{j,a}^b \frac{\partial \mathcal{L}}{\partial \zeta_{(i,j)}^b} \right) + \mathcal{F}_a^i, \\ (\alpha_\partial)_a^i = 0, \quad (\dot{\alpha}_\partial)_a^i = \left. \dfrac{\partial \mathcal{L}}{\partial \zeta_{(i,m)}^a} \right|_{x^m=0} + (\mathcal{F}_\partial)_a^i, 1 \le i < m, \end{cases}$$

$$\zeta \in \Omega^{k+1}(M, E) \text{ and } \zeta_{(i,j)} = -\zeta_{(j,i)} \text{ for each } 1 \le i < j \le m$$

$$\epsilon(i,j) = \begin{cases} -1, & 1 \le j \le i-1 \\ 0, & i = j \\ +1, & i+1 \le j \le m \end{cases}$$

$$\mathcal{E}: W^k(M, E) \to \Lambda^m T^* M$$

$$\begin{aligned} \mathcal{E}(\varphi_x, v_x, \zeta_x) &= \frac{\partial \mathcal{L}}{\partial v}(\varphi_x, v_x, \zeta_x) \cdot v_x - \mathcal{L}(\varphi_x, v_x, \zeta_x) \\ &= v_x \wedge \frac{\partial \mathcal{L}}{\partial v}(\varphi_x, v_x, \zeta_x) - \mathcal{L}(\varphi_x, v_x, \zeta_x) \end{aligned}$$

$$(\varphi_x, v_x, \zeta_x) \in W^k(M, E)_x$$

$$V^\star = \Lambda_m^k(M, E^*) \times \Lambda_{m-1}^k(\partial M, E^*), \text{ let } (\varphi, v, \alpha, \alpha_\partial): [t_0, t_1] \to TV \oplus T^\star V$$

$$\frac{\partial}{\partial t} \mathcal{E}(\varphi, v, \mathrm{d}^\nabla \varphi) = -\mathrm{div}\left(\frac{\partial \mathcal{L}}{\partial \zeta}(\varphi, v, \mathrm{d}^\nabla \varphi) \cdot v \right) + \mathcal{F}^\star(\varphi, v) \cdot v$$

$$\frac{d}{dt} \int_M \mathcal{E}(\varphi, v, \mathrm{d}^\nabla \varphi) = \underbrace{\int_M \mathcal{F}^\star(\varphi, v) \cdot v}_{\text{spatially distributed contribution}} + \underbrace{\int_{\partial M} \mathcal{F}_\partial^\star(\varphi, v) \cdot \iota_{\partial M}^* v}_{\text{boundary contribution}}$$

$$V^\dagger = \Omega^{m-k}(M, E^*) \times \Omega^{m-k-1}(\partial M, E^*), \text{ let } (\varphi, v, \alpha, \alpha_\partial): [t_0, t_1] \to TV \oplus T^\dagger V$$

$$\frac{\partial}{\partial t} \mathcal{E}(\varphi, v, \mathrm{d}^\nabla \varphi) = -\mathrm{d}\left(v \wedge \frac{\partial \mathcal{L}}{\partial \zeta}(\varphi, v, \mathrm{d}^\nabla \varphi) \right) + v \wedge \mathcal{F}^\dagger(\varphi, v)$$

$$\frac{d}{dt} \int_M \mathcal{E}(\varphi, v, \mathrm{d}^\nabla \varphi) = \underbrace{\int_M v \wedge \mathcal{F}^\dagger(\varphi, v)}_{\text{spatially distributed contribution}} + \underbrace{\int_{\partial M} \iota_{\partial M}^* v \wedge \mathcal{F}_\partial^\dagger(\varphi, v)}_{\text{boundary contribution}}$$

$$V^\star = \Lambda_m^k(M, E^*) \times \Lambda_{m-1}^k(\partial M, E^*)$$

$$\begin{aligned}\frac{\partial}{\partial t}\mathcal{E}(\varphi, \nu, \mathrm{d}^\nabla\varphi) &= \left(\frac{\partial}{\partial t}\frac{\partial \mathcal{L}}{\partial \nu}\right)\cdot \nu + \frac{\partial \mathcal{L}}{\partial \nu}\cdot \dot{\nu} - \left(\frac{\partial \mathcal{L}}{\partial \varphi}\cdot \dot{\varphi} + \frac{\partial \mathcal{L}}{\partial \nu}\cdot \dot{\nu} + \frac{\partial \mathcal{L}}{\partial \zeta}\cdot \mathrm{d}^\nabla \nu\right) \\ &= \left(-(-1)^k \mathrm{div}^\nabla \frac{\partial \mathcal{L}}{\partial \zeta} + \mathcal{F}^\star(\varphi, \nu)\right)\cdot \nu - \frac{\partial \mathcal{L}}{\partial \zeta}\cdot \mathrm{d}^\nabla \nu \\ &= -\mathrm{div}\left(\frac{\partial \mathcal{L}}{\partial \zeta}\cdot \nu\right) + \mathcal{F}^\star(\varphi, \nu)\cdot \nu\end{aligned}$$

$$\begin{aligned}\frac{d}{dt}\int_M \mathcal{E}(\varphi, \nu, \mathrm{d}^\nabla\varphi) &= \int_M \left[-\mathrm{div}\left(\frac{\partial \mathcal{L}}{\partial \zeta}\cdot \nu\right) + \mathcal{F}^\star(\varphi, \nu)\cdot \nu\right] \\ &= \int_M \mathcal{F}^\star(\varphi, \nu)\cdot \nu - \int_{\partial M} \iota_{\partial M}^*\left(\mathrm{tr}\frac{\partial \mathcal{L}}{\partial \zeta}\right)\cdot \iota_{\partial M}^* \nu \\ &= \int_M \mathcal{F}^\star(\varphi, \nu)\cdot \nu + \int_{\partial M} \mathcal{F}_\partial^\star(\varphi, \nu)\cdot \iota_{\partial M}^* \nu\end{aligned}$$

$$V^\dagger = \Omega^{m-k}(M, E^*) \times \Omega^{m-k-1}(\partial M, E^*)$$

$$\begin{aligned}\frac{\partial}{\partial t}\mathcal{E}(\varphi, \nu, \mathrm{d}^\nabla\varphi) &= \nu \wedge \frac{\partial}{\partial t}\frac{\partial \mathcal{L}}{\partial \nu} + \dot{\nu} \wedge \frac{\partial \mathcal{L}}{\partial \nu} - \left(\dot{\varphi} \wedge \frac{\partial \mathcal{L}}{\partial \varphi} + \dot{\nu} \wedge \frac{\partial \mathcal{L}}{\partial \nu} + \mathrm{d}^\nabla \nu \wedge \frac{\partial \mathcal{L}}{\partial \zeta}\right) \\ &= \nu \wedge \left(-(-1)^k\, \mathrm{d}^\nabla \frac{\partial \mathcal{L}}{\partial \zeta} + \mathcal{F}^\dagger(\varphi, \nu)\right) - \mathrm{d}^\nabla \nu \wedge \frac{\partial \mathcal{L}}{\partial \zeta} \\ &= -\mathrm{d}\left(\nu \wedge \frac{\partial \mathcal{L}}{\partial \zeta}\right) + \nu \wedge \mathcal{F}^\dagger(\varphi, \nu)\end{aligned}$$

$$\mathrm{d}\left(\nu \wedge \frac{\partial \mathcal{L}}{\partial \zeta}\right) = \mathrm{d}^\nabla \nu \wedge \frac{\partial \mathcal{L}}{\partial \zeta} + (-1)^k \nu \wedge \left(\mathrm{d}^{\nabla^*}\frac{\partial \mathcal{L}}{\partial \zeta}\right).$$

$$\begin{aligned}\frac{d}{dt}\int_M \mathcal{E}(\varphi, \nu, \mathrm{d}^\nabla\varphi) &= \int_M \left[-\mathrm{d}\left(\nu \wedge \frac{\partial \mathcal{L}}{\partial \zeta}\right) + \nu \wedge \mathcal{F}^\dagger(\varphi, \nu)\right] \\ &= \int_M \nu \wedge \mathcal{F}^\dagger(\varphi, \nu) - \int_{\partial M} \iota_{\partial M}^* \nu \wedge \iota_{\partial M}^*\left(\frac{\partial \mathcal{L}}{\partial \zeta}\right) \\ &= \int_M \nu \wedge \mathcal{F}^\dagger(\varphi, \nu) + \int_{\partial M} \iota_{\partial M}^* \nu \wedge \mathcal{F}_\partial^\dagger(\varphi, \nu)\end{aligned}$$

$$S = \frac{\partial \mathcal{L}}{\partial \zeta}\cdot \nu \text{ or } \mathrm{S} = \nu \wedge \frac{\partial \mathcal{L}}{\partial \zeta}$$

$$\mathcal{F}_\partial^\star \in \Lambda_{m-1}^k(\partial M, E^*) \text{ and } \iota_{\partial M}^* \nu \in \Omega^k(\partial M, E)$$

$$\mathcal{F}^\star \in \Lambda_m^k(M, E^*) \text{ and } \nu \in \Omega^k(M, E)$$

$$\mathcal{F}^\star \cdot \nu \in \Omega^m(M) \text{ and } \mathcal{F}_\partial^\star \cdot \iota_{\partial M}^* \nu \in \Omega^{m-1}(\partial M)$$

$$\mathcal{F}_\partial^\dagger \in \Omega^{m-k-1}(\partial M, E^*) \text{ and } \mathcal{F}^\dagger \in \Omega^{m-k}(M, E^*)$$

$$\nu \wedge \mathcal{F}^\dagger \in \Omega^m(M) \text{ and } \iota_{\partial M}^* \nu \wedge \mathcal{F}_\partial^\dagger \in \Omega^{m-1}(\partial M)$$

$(q(t), v(t), p(t)) \in TQ \oplus T^*Q$ of the Lagrange-Dirac dynamical system

$$(\varphi(t), \nu(t), \alpha(t), \alpha_\partial(t)) \in TV \oplus T^\star V \text{ or } TV \oplus T^\dagger V$$

Lagrange-d'Alembert density:

$$\begin{cases} \dfrac{\partial}{\partial t}\dfrac{\partial \mathcal{L}}{\partial \nu}\left(\varphi, \dot{\varphi}, \mathrm{d}^\nabla \varphi\right) = \dfrac{\partial \mathcal{L}}{\partial \varphi}\left(\varphi, \dot{\varphi}, \mathrm{d}^\nabla \varphi\right) - (-1)^k \mathrm{div}^\nabla \dfrac{\partial \mathcal{L}}{\partial \zeta}\left(\varphi, \dot{\varphi}, \mathrm{d}^\nabla \varphi\right) + \mathcal{F}^\star(\varphi, \dot{\varphi}) \\ \mathcal{F}_\partial^\star(\varphi, \dot{\varphi}) = -\iota_{\partial M}^* \left(\mathrm{tr} \dfrac{\partial \mathcal{L}}{\partial \zeta}\left(\varphi, \dot{\varphi}, \mathrm{d}^\nabla \varphi\right) \right) \end{cases}$$

$$\begin{cases} \dfrac{\partial}{\partial t}\dfrac{\partial \mathcal{L}}{\partial \nu}\left(\varphi, \dot{\varphi}, \mathrm{d}^\nabla \varphi\right) = \dfrac{\partial \mathcal{L}}{\partial \varphi}\left(\varphi, \dot{\varphi}, \mathrm{d}^\nabla \varphi\right) - (-1)^k \, \mathrm{d}^* \left(\dfrac{\partial \mathcal{L}}{\partial \zeta}\left(\varphi, \dot{\varphi}, \mathrm{d}^\nabla \varphi\right) \right) + \mathcal{F}^\dagger(\varphi, \dot{\varphi}) \\ \mathcal{F}_\partial^\dagger(\varphi, \dot{\varphi}) = -\iota_{\partial M}^* \left(\dfrac{\partial \mathcal{L}}{\partial \zeta}\left(\varphi, \dot{\varphi}, \mathrm{d}^\nabla \varphi\right) \right) \end{cases}$$

$V^\star = \Lambda_m^k(M, E^*) \times \Lambda_{m-1}^k(\partial M, E^*)$, a curve $\varphi\colon [t_0, t_1] \to \Omega^k(M, E)$

$$\begin{aligned} &\delta \int_{t_0}^{t_1} L_\nabla(\varphi, \dot{\varphi}) dt + \int_{t_0}^{t_1} \langle F^\star(\varphi, \dot{\varphi}), \delta\varphi \rangle_\star dt \\ &= \delta \int_{t_0}^{t_1} \left(\int_M \mathcal{L}\left(\varphi, \dot{\varphi}, \mathrm{d}^\nabla \varphi\right) \right) dt + \int_{t_0}^{t_1} \left(\int_M \mathcal{F}^\star(\varphi, \dot{\varphi}) \cdot \delta\varphi + \int_{\partial M} \mathcal{F}_\partial^\star(\varphi, \dot{\varphi}) \cdot \iota_{\partial M}^* \delta\varphi \right) dt = 0 \end{aligned}$$

$V^\dagger = \Omega^{m-k}(M, E^*) \times \Omega^{m-k-1}(\partial M, E^*)$, a curve $\varphi\colon [t_0, t_1] \to \Omega^k(M, E)$

$$\begin{aligned} &\delta \int_{t_0}^{t_1} L_\nabla(\varphi, \dot{\varphi}) dt + \int_{t_0}^{t_1} \left\langle F^\dagger(\varphi, \dot{\varphi}), \delta\varphi \right\rangle_\dagger dt \\ &= \delta \int_{t_0}^{t_1} \left(\int_M \mathcal{L}\left(\varphi, \dot{\varphi}, \mathrm{d}^\nabla \varphi\right) \right) dt + \int_{t_0}^{t_1} \left(\int_M \delta\varphi \wedge \mathcal{F}^\dagger(\varphi, \dot{\varphi}) + \int_{\partial M} \delta\varphi \wedge \mathcal{F}_\partial^\dagger(\varphi, \dot{\varphi}) \right) dt = 0 \end{aligned}$$

$$\begin{aligned} 0 &= \int_{t_0}^{t_1} \int_M \left(\frac{\partial \mathcal{L}}{\partial \varphi} \cdot \delta\varphi + \frac{\partial \mathcal{L}}{\partial \nu} \cdot \delta\dot{\varphi} + \frac{\partial \mathcal{L}}{\partial \zeta} \cdot \delta \mathrm{d}^\nabla \varphi \right) dt + \int_{t_0}^{t_1} \langle F^\star(\varphi, \dot{\varphi}), \delta\varphi \rangle_\star \\ &= \int_{t_0}^{t_1} \left[\int_M \left(\frac{\partial \mathcal{L}}{\partial \varphi} - \frac{\partial}{\partial t}\frac{\partial \mathcal{L}}{\partial \nu} - (-1)^k \mathrm{div}\nabla^* \frac{\partial \mathcal{L}}{\partial \zeta} + \mathcal{F}^\star(\varphi, \dot{\varphi}) \right) \cdot \delta\varphi \right. \\ &\quad \left. + \int_{\partial M} \left(\iota_{\partial M}^* \left(\mathrm{tr} \frac{\partial \mathcal{L}}{\partial \zeta} \right) + \mathcal{F}_\partial^\star(\varphi, \dot{\varphi}) \right) \cdot \iota_{\partial M}^* \delta\varphi \right] dt + \left[\int_M \frac{\partial \mathcal{L}}{\partial \nu} \cdot \delta\varphi \right]_{t=t_0}^{t=t_1} \end{aligned}$$

For $V^\star = \Lambda_m^k(M, E^*) \times \Lambda_{m-1}^k(\partial M, E^*)$, a curve $(\varphi, \nu, \alpha, \alpha_\partial)\colon [t_0, t_1] \to TV \oplus T^\star V$

$$\begin{aligned} &\delta \int_{t_0}^{t_1} (L_\nabla(\varphi, \nu) + \langle (\alpha, \alpha_\partial), \dot{\varphi} - \nu \rangle_\star) dt + \int_{t_0}^{t_1} \langle F^\star(\varphi, \dot{\varphi}), \delta\varphi \rangle_\star dt \\ &= \delta \int_{t_0}^{t_1} \left(\int_M \mathcal{L}\left(\varphi, \nu, \mathrm{d}^\nabla \varphi\right) + \int_M \alpha \cdot (\dot{\varphi} - \nu) + \int_{\partial M} \alpha_\partial \cdot (\dot{\varphi} - \nu) \right) dt \\ &+ \int_{t_0}^{t_1} \left(\int_M \mathcal{F}^\star(\varphi, \dot{\varphi}) \cdot \delta\varphi + \int_{\partial M} \mathcal{F}_\partial^\star(\varphi, \dot{\varphi}) \cdot \iota_{\partial M}^* \delta\varphi \right) dt = 0 \end{aligned}$$

$V^\dagger = \Omega^{m-k}(M, E^*) \times \Omega^{m-k-1}(\partial M, E^*)$, a curve $(\varphi, \nu, \alpha, \alpha_\partial)\colon [t_0, t_1] \to TV \oplus T^\dagger V$

$$\delta \int_{t_0}^{t_1} \left(L_\nabla(\varphi, \nu) + \langle (\alpha, \alpha_\partial), \dot{\varphi} - \nu \rangle_\dagger\right) dt + \int_{t_0}^{t_1} \left\langle F^\dagger(\varphi, \dot{\varphi}), \delta\varphi \right\rangle_\dagger dt$$
$$= \delta \int_{t_0}^{t_1} \left(\int_M \mathcal{L}\left(\varphi, \nu, \mathrm{d}^\nabla \varphi\right) + \int_M (\dot{\varphi} - \nu) \wedge \alpha + \int_{\partial M} (\dot{\varphi} - \nu) \wedge \alpha_\partial \right) dt$$
$$+ \int_{t_0}^{t_1} \left(\int_M \delta\varphi \wedge \mathcal{F}^\dagger(\varphi, \dot{\varphi}) + \int_{\partial M} \iota_{\partial M}^* \delta\varphi \wedge \mathcal{F}_\partial^\dagger(\varphi, \dot{\varphi}) \right) dt = 0$$

$$V^\star = \Lambda_m^k(M) \times \Lambda_{m-1}^k(\partial M) \text{ and } V^\dagger = \Omega^{m-k}(M) \times \Omega^{m-k-1}(\partial M)$$

$$\langle (\alpha, \alpha_\partial), \varphi \rangle_\star = \int_M \alpha \cdot \varphi + \int_{\partial M} \alpha_\partial \cdot \iota_{\partial M}^* \varphi, (\alpha, \alpha_\partial) \in V^\star, \varphi \in V$$

$$\langle (\alpha, \alpha_\partial), \varphi \rangle_\dagger = \int_M \varphi \wedge \alpha + \int_{\partial M} \iota_{\partial M}^* \varphi \wedge \alpha_\partial, (\alpha, \alpha_\partial) \in V^\dagger, \varphi \in V,$$

$$\mathrm{div}(\chi \cdot \delta\varphi) = \chi \cdot \mathrm{d}\delta\varphi + (-1)^k \mathrm{div}(\chi) \cdot \delta\varphi$$

$\varphi \in \Omega^k(M)$ and $\chi \in \Lambda_m^{k+1}(M)$

$$L\colon T\Omega^k(M) \to \mathbb{R}, (\varphi, \nu) \mapsto L(\varphi, \nu) = \int_M \mathcal{L}(\varphi, \nu, \mathrm{d}\varphi)$$

$$\mathbf{g}\colon \left(\Lambda^k T^*M \otimes E\right) \times_M \left(\Lambda^k T^*M \otimes E\right) \to \mathbb{R}$$

$$\mathbf{g}_x(\alpha_1 \otimes \varphi_1, \alpha_2 \otimes \varphi_2) = g_x(\alpha_1, \alpha_2)\kappa_x(\varphi_1, \varphi_2)$$

$x \in M, \alpha_i \in \Lambda^k T_x^*M$ and $\varphi_i \in E_x, i = 1,2$

$\sharp\colon T^*M \to TM$ and $b\colon TM \to T^*M$

$\sharp_\kappa\colon E^* \to E$ and $b_\kappa\colon E \to E^*$, where $\pi_{E^*,M}\colon E^* \to M$ is the dual bundle of $\pi_{E,M}$

$$\sharp\colon \Lambda^k T^*M \otimes E \to \Lambda^k TM \otimes E^*, \quad \alpha \otimes \varphi \mapsto \alpha^\sharp \otimes \varphi^{b_\kappa},$$
$$\natural\colon \Lambda^k TM \otimes E^* \to \Lambda^k T^*M \otimes E, \quad U \otimes \eta \mapsto U^\natural \otimes \eta^{\sharp_\kappa}.$$

$\star\,(\alpha \otimes \varphi) = (\star\,\alpha) \otimes \varphi$, where $\star\colon \Lambda^k T^*M \to \Lambda^{m-k} T^*M$ denotes the standard Hodge star operator

$\mathbf{g}(\sigma_1, \sigma_2)\mu_g = \sigma_1^{\natural_\kappa} \wedge \star\, \sigma_2$ for each $x \in M$ and $\sigma_1, \sigma_2 \in \Lambda^k T_x^*M \otimes E_x$, where $\mu_g \in \Omega^m(M)$ is the

Riemannian volume $\sigma_1^{b_\kappa} \in \Lambda^k T_x^*M \otimes E_x^*$.

$$g_\partial = \iota_{\partial M}^* g$$

isomorphisms are denoted by $\sharp_\partial\colon T^*\partial M \to T\partial M$ and $b_\partial\colon T\partial M \to T^*\partial M$

$\star_\partial\colon \Omega^k(\partial M) \to \Omega^{m-k-1}(\partial M)$ and $\mu_g^\partial = \iota_{\partial M}^*\left(i_n \mu_g\right) \in \Omega^{m-1}(\partial M)$ where $n \in \mathfrak{X}(M)|_{\partial M}$ is the vector

field on ∂M, and $i_U\colon \Omega^k(M) \to \Omega^{k-1}(M)$ is the left interior by $U \in \mathfrak{X}(M)$

$$\bar{\pi}_{E,M}\colon \bar{E} \to \bar{M}, \text{ where } \bar{M} = [t_0, t_1] \times M \text{ and } \bar{E} = [t_0, t_1] \times E$$

$$\eta = -dt \otimes dt + g$$

$$\boldsymbol{\eta}\colon \left(\Lambda^k T^*\bar{M} \otimes \bar{E}\right) \times \bar{M}\left(\Lambda^k T^*\bar{M} \otimes \bar{E}\right) \to \mathbb{R}$$

$$\boldsymbol{\eta}_{(t,x)}(\bar{\alpha}_1 \otimes \bar{\varphi}_1, \bar{\alpha}_2 \otimes \bar{\varphi}_2) = \eta_{(t,x)}(\bar{\alpha}_1, \bar{\alpha}_2)\kappa_x(\varphi_1, \varphi_2)$$

$$(t, x) \in \bar{M}, \bar{\alpha}_i \in \Lambda^k T^*_{(t,x)}\bar{M} \text{ and } \bar{\varphi}_i = (t, \varphi_i) \in \bar{E}_{(t,x)} = \{t\} \times E_x, i = 1,2$$

1. If $\bar{\alpha}_1 = dt$, then

$$\boldsymbol{\eta}_{(t,x)}(\bar{\alpha}_1 \otimes \bar{\varphi}_1, \bar{\alpha}_2 \otimes \bar{\varphi}_2) = \underbrace{\eta_{(t,x)}(dt, dt)}_{-1}\kappa_x(\varphi_1, \varphi_2) = -\mathbf{g}_x(\varphi_1, \varphi_2)$$

2. If $\bar{\alpha}_i = \alpha_i \in T^*_x M, i = 1,2$, then

$$\boldsymbol{\eta}_{(t,x)}(\bar{\alpha}_1 \otimes \bar{\varphi}_1, \bar{\alpha}_2 \otimes \bar{\varphi}_2) = \underbrace{\eta_{(t,x)}(\alpha_1, \alpha_2)}_{g_x(\alpha_1,\alpha_2)}\kappa_x(\varphi_1, \varphi_2) = \mathbf{g}_x(\alpha_1 \otimes \varphi_1, \alpha_2 \otimes \varphi_2)$$

$$\bar{\pi}_{E,M}\colon \bar{E} \to \bar{M}$$

$$\bar{\nabla}\bar{\varphi}(t, x) = (t, \nabla\varphi(t, x) + \dot{\varphi}(t, x)dt), \bar{\varphi}(t, x) = (t, \varphi(t, x)) \in \Gamma(\bar{E} \to \bar{M}) = \Omega^0(\bar{M}, \bar{E})$$

$$\mathcal{A}\colon \Omega^0(\bar{M}, \bar{E}) \to \mathbb{R}, \bar{\varphi} \mapsto \mathcal{A}(\bar{\varphi}) = \int_{\bar{M}} \mathfrak{L}\left(\bar{\varphi},\ \mathrm{d}^{\bar{\nabla}}\bar{\varphi}\right)$$

$$\mathfrak{L}\colon (\Lambda^0 T^*\bar{M} \otimes \bar{E}) \times_{\bar{M}} (\Lambda^1 T^*\bar{M} \otimes \bar{E}) \to \Lambda^{m+1} T^*\bar{M}$$

$$\mathcal{L}\colon (\Lambda^0 T^*M \otimes E) \times_M (\Lambda^0 T^*M \otimes E) \times_M (\Lambda^1 T^*M \otimes E) \to \Lambda^m T^*M$$

$$\mathfrak{L}\left(\bar{\varphi}, \mathrm{d}^{\bar{\nabla}}\bar{\varphi}\right) = \mathcal{L}\left(\varphi, \dot{\varphi}, \mathrm{d}^{\nabla}\varphi\right)dt$$

$$\mathcal{L}\left(\varphi, \dot{\varphi}, \mathrm{d}^{\nabla}\varphi\right) = i_{\partial_t}\left(\mathfrak{L}\left(\bar{\varphi},\ \mathrm{d}^{\bar{\nabla}}\bar{\varphi}\right)\right)$$

$$\mathcal{A}(\bar{\varphi}) = \int_{t_0}^{t_1} L_{\nabla}(\varphi, \dot{\varphi})dt = \int_{t_0}^{t_1}\int_M \mathcal{L}\left(\varphi, \dot{\varphi},\ \mathrm{d}^{\nabla}\varphi\right)dt$$

$$\mathfrak{L}\left(\bar{\varphi}, \mathrm{d}^{\bar{\nabla}}\bar{\varphi}\right) = -\left(\frac{1}{2}\boldsymbol{\eta}\left(\ \mathrm{d}^{\bar{\nabla}}\bar{\varphi}, \mathrm{d}^{\bar{\nabla}}\bar{\varphi}\right) + \mathbf{V}(\bar{\varphi})\right)dt \wedge \mu_g$$

$$\mathbf{V}(t, \varphi_x) = \mathbf{V}(\varphi_x) \text{ for each } (t, x) \in \bar{M} \text{ and } \varphi_x \in E_x$$

$$\boldsymbol{\eta}\left(\mathrm{d}^{\bar{\nabla}}\bar{\varphi}, \mathrm{d}^{\bar{\nabla}}\bar{\varphi}\right) = -\mathbf{g}(\dot{\varphi}, \dot{\varphi}) + \mathbf{g}\left(\mathrm{d}^{\nabla}\varphi, \mathrm{d}^{\nabla}\varphi\right)$$

$$\mathrm{d}^{\bar{\nabla}}\bar{\varphi} = \bar{\nabla}\bar{\varphi} \text{ and } \mathrm{d}^{\nabla}\varphi = \nabla\varphi$$

$$\mathcal{L}(\varphi_x, \nu_x, \zeta_x) = \left(\frac{1}{2}\mathbf{g}(\nu_x, \nu_x) - \frac{1}{2}\mathbf{g}(\zeta_x, \zeta_x) - \mathbf{V}(\varphi_x)\right)\mu_g$$

$$= \frac{1}{2}\left(\nu_x^{b_\kappa} \wedge\star\, \nu_x - \zeta_x^{b_\kappa} \wedge\star\, \zeta_x\right) -\star\, \mathbf{V}(\varphi_x)$$

$$(\varphi_x, v_x, \zeta_x) \in E_x \times E_x \times T_x^* M \otimes E_x \text{ and } x \in M$$

$$\begin{cases} \dot{\varphi}^a = v^a \\ \alpha_a = \dfrac{\partial \mathcal{L}}{\partial v^a}, \dot{\alpha}_a = \dfrac{\partial \mathcal{L}}{\partial \varphi^a} - \left(\partial_j \dfrac{\partial \mathcal{L}}{\partial \zeta_j^a} - \Gamma_{j,a}^b \dfrac{\partial \mathcal{L}}{\partial \zeta_j^b} \right) + \mathcal{F}_a \\ (\alpha_\partial)_a = 0, (\dot{\alpha}_\partial)_a = \left. \dfrac{\partial \mathcal{L}}{\partial \zeta_m^a} \right|_{x^m = 0} + (\mathcal{F}_\partial)_a \end{cases}$$

$$\partial \mathbf{V} / \partial \varphi : E \to E^*$$

$$\frac{\partial \mathbf{V}}{\partial \varphi}(\varphi_x) \cdot \delta \varphi_x = \left. \frac{d}{d\epsilon} \right|_{\epsilon = 0} \mathbf{V}(\varphi_x + \epsilon \delta \varphi_x), \varphi_x, \delta \varphi_x \in E_x, x \in M$$

$$\operatorname{grad}_\kappa \mathbf{V} = \sharp_\kappa \circ \partial \mathbf{V} / \partial \varphi : E \to E$$

$$V = \Omega^0(M, E) \text{ and } (\varphi, \dot{\varphi}) \in TV$$

$$\zeta = \mathrm{d}^\nabla \varphi \in \Omega^1(M, E)$$

$$\begin{aligned} &\frac{\partial \mathcal{L}}{\partial \varphi} = -\frac{\partial \mathbf{V}}{\partial \varphi} \otimes \mu_g, \quad \frac{\partial \mathcal{L}}{\partial v} = \dot{\varphi}^{\sharp_\partial} \otimes \mu_g, \quad \frac{\partial \mathcal{L}}{\partial \zeta} = -\left(\mathrm{d}^\nabla \varphi\right)^{\sharp_\partial} \otimes \mu_g, \\ &\frac{\partial \mathcal{L}}{\partial \varphi} = -\star \frac{\partial \mathbf{V}}{\partial \varphi}, \qquad \frac{\partial \mathcal{L}}{\partial v} = \star\, \dot{\varphi}^{\mathfrak{G}_\kappa}, \qquad \frac{\partial \mathcal{L}}{\partial \zeta} = -\star \left(\mathrm{d}^\nabla \varphi\right)^{\mathfrak{G}_\kappa}. \end{aligned}$$

$$\mathcal{F}^\star : TV \to \Lambda_m^0(M, E^*) = \Omega^m(M, E^*) \text{ and } \mathcal{F}_\partial^\star : TV \to \Lambda_{m-1}^0(\partial M, E^*) = \Omega^{m-1}(\partial M, E^*)$$

$$\begin{cases} \dot{\varphi} = v, \\ \alpha = \dot{\varphi}^{\sharp_\partial} \otimes \mu_g, \quad \dot{\alpha} = -\dfrac{\partial \mathbf{V}}{\partial \varphi} \otimes \mu_g + \operatorname{div}^{\nabla^*} \left(\left(\mathrm{d}^\nabla \varphi \right)^{\sharp_\partial} \otimes \mu_g \right) + \mathcal{F}^\star, \\ \alpha_\partial = 0, \qquad \dot{\alpha}_\partial = -\iota_{\partial M}^* \left(\operatorname{tr}\left(\mathrm{d}^\nabla \varphi \right)^{\sharp_\partial} \otimes \mu_g \right) + \mathcal{F}_\partial^\star. \end{cases}$$

$$\begin{cases} \ddot{\varphi}^{\sharp_\partial} \otimes \mu_g - \operatorname{div}^{\nabla^*} \left(\left(\mathrm{d}^\nabla \varphi \right)^{\sharp_\partial} \otimes \mu_g \right) + \dfrac{\partial \mathbf{V}}{\partial \varphi} \otimes \mu_g = \mathcal{F}^\star, \\ \iota_{\partial M}^* \left(\operatorname{tr}\left(\mathrm{d}^\nabla \varphi \right)^{\sharp_\partial} \otimes \mu_g \right) = \mathcal{F}_\partial^\star. \end{cases}$$

$$\mathcal{F}^\dagger : TV \to \Omega^m(M, E^*) \text{ and } \mathcal{F}_\partial^\dagger : TV \to \Omega^{m-1}(\partial M, E^*)$$

$$\begin{cases} \dot{\varphi} = v, \\ \alpha = \star\, \dot{\varphi}^{\mathfrak{G}_\kappa}, \quad \dot{\alpha} = -\star \dfrac{\partial \mathbf{V}}{\partial \varphi} + \mathrm{d}^{\nabla^*} \left(\star \left(\mathrm{d}^\nabla \varphi \right)^{\mathfrak{G}_\kappa} \right) + \mathcal{F}^\dagger \\ \alpha_\partial = 0, \qquad \dot{\alpha}_\partial = -\iota_{\partial M}^* \left(\star \left(\mathrm{d}^\nabla \varphi \right)^{\mathfrak{G}_\kappa} \right) + \mathcal{F}_\partial^\dagger \end{cases}$$

$$\begin{cases} \star\, \ddot{\varphi}^{\mathfrak{G}_\kappa} - \mathrm{d}^{\nabla^*} \left(\star \left(\mathrm{d}^\nabla \varphi \right)^{\mathfrak{G}_\kappa} \right) + \star \dfrac{\partial \mathbf{V}}{\partial \varphi} = \mathcal{F}^\dagger \\ \iota_{\partial M}^* \left(\star \left(\mathrm{d}^\nabla \varphi \right)^{\mathfrak{G}_\kappa} \right) = \mathcal{F}_\partial^\dagger \end{cases}$$

$\mathrm{d}\kappa(\varphi_1, \varphi_2) = \kappa(\nabla \varphi_1, \varphi_2) + \kappa(\varphi_1, \nabla \varphi_2)$ for each $\varphi_1, \varphi_1 \in V$, we have $\mathfrak{G}_\kappa \circ \mathrm{d}^\nabla = \mathrm{d}^{\nabla^*} \circ \mathfrak{G}_\kappa$

$\mathcal{F}^{\dagger} = \star\, ב^{\flat_\kappa}$ and $\mathcal{F}_{\partial}^{\dagger} = \star_{\partial}\, ב^{\flat_\kappa}$ for some $ב: TV \to \Omega^0(M,E)$ and $ב: TV \to \Omega^0(\partial M,E)$

$$\begin{cases} \ddot{\varphi} + \delta^{\nabla}\left(\mathrm{d}^{\nabla}\varphi\right) + \mathrm{grad}_{\kappa}\mathbf{V} = ב \\ \star_{\partial}\left(\iota_{\partial M}^{*}\left(\star\ \mathrm{d}^{\nabla}\varphi\right)\right) = I \end{cases}$$

$$\delta^{\nabla} = -\star \circ \mathrm{d}^{\nabla} \circ \star : \Omega^1(M,E) \to \Omega^0(M,E)$$

In local coordinates, the Riemannian metric $\pi_{E,M}: E \to M$ given $g = g_{ij}dx^i \otimes dx^j$ and $\kappa =$

$\kappa_{ab}B^a \otimes B^b$ for some $g_{ij}, \kappa_{ab} \in C^{\infty}(M), 1 \le i,j \le m, 1 \le a,b \le n$

$\left(g^{ij}\right)_{1\le i,j\le m}$ and $\left(\kappa^{ab}\right)_{1\le a,b\le n}$ the inverse matrices of $\left(g_{ij}\right)_{1\le i,j\le m}$ and $(\kappa_{ab})_{1\le a,b\le n}$

$$\begin{cases} \ddot{\varphi}^a - \partial_j\left(g^{ij}\partial_i\varphi^a\right) + \kappa^{ab}\dfrac{\partial \mathbf{V}}{\partial \varphi^b} = ב^a \\ g^{im}\partial_i\varphi^a\big|_{x^m=0} = ב^a \end{cases}$$

where we have denoted $ב = ב^a B_a$ and $ב = ב^a B_a$, for some $ב^a \in C^{\infty}(M)$ and $ב^a \in C^{\infty}(\partial M)$, $1 \le a \le$

n.

Energy considerations:

$$\begin{aligned} \mathcal{E} &= \left(\frac{1}{2}\mathbf{g}(\dot{\varphi},\dot{\varphi}) + \frac{1}{2}\mathbf{g}\left(\mathrm{d}^{\nabla}\varphi, \mathrm{d}^{\nabla}\varphi\right) + \mathbf{V}(\varphi)\right)\mu_g \\ &= \frac{1}{2}\left(\dot{\varphi}^{\flat_\kappa} \wedge \star\, \dot{\varphi} + \left(\mathrm{d}^{\nabla}\varphi\right)^{\flat_\kappa} \wedge \star\, \mathrm{d}^{\nabla}\varphi\right) + \mathbf{V}(\varphi)\mu_g \end{aligned}$$

$$\frac{\partial \mathcal{E}}{\partial t} = -\mathrm{div}S + \mathcal{F}^{\star} \cdot \dot{\varphi}$$

$$\frac{d}{dt}\int_M \mathcal{E} = \underbrace{\int_M \mathcal{F}^{\star} \cdot \dot{\varphi}}_{\text{spatially distributed contribution}} + \underbrace{\int_{\partial M} \mathcal{F}_{\partial}^{\star} \cdot \iota_{\partial M}^{*}\dot{\varphi}}_{\text{boundary contribution}}$$

$$S = -\dot{\varphi} \cdot \left(\mathrm{d}^{\nabla}\varphi\right)^{\sharp_\partial} \otimes \mu_g \in \Lambda_m^1(M)$$

$$\frac{\partial \mathcal{E}}{\partial t} = -\mathrm{dS} + \dot{\varphi} \wedge \star\, ב^{\flat_K}$$

$$\frac{d}{dt}\int_M \mathcal{E} = \underbrace{\int_M \dot{\varphi} \wedge \star\, ב^{\flat_\kappa}}_{\text{spatially distributed contribution}} + \underbrace{\int_{\partial M} (\iota_{\partial M}^{*}\dot{\varphi}) \wedge \star_{\partial}\, ב^{\flat_\kappa}}_{\text{boundary contribution}}$$

$$S = -\dot{\varphi} \wedge \star \left(\mathrm{d}^{\nabla}\varphi\right)^{\flat_\kappa} \in \Omega^{m-1}(M)$$

$\tilde{\mathfrak{g}} = (P \times \mathfrak{g})/G \to M$ by $\mathrm{Conn}(P) = (J^1P)/G \to M$

$$\mathcal{C}(P) = \Gamma(\mathrm{Conn}(P) \to M)$$

$$T^*M \otimes \tilde{\mathfrak{g}} \to M \text{ and } \pi_{P,M}: P \to M$$

$$\mathcal{C}(P) \ni A \overset{\blacksquare:\blacksquare}{\leftrightarrow} \mathbf{A} \in \Omega^1(P, \mathfrak{g})$$

$$T\mathcal{C}(P) \simeq \mathcal{C}(P) \times \Omega^1(M, \tilde{\mathfrak{g}}), v_A \mapsto (A, \dot{A})$$

where $\dot{A} = \dot{\bar{\gamma}}(0)$, for $\gamma: (-\epsilon, \epsilon) \to \mathcal{C}(P)$ a curve such that $\gamma(0) = A$ and $\dot{\gamma}(0) = v_A$, and $\bar{\gamma}: (-\epsilon, \epsilon) \to$

$\Omega^1(M, \tilde{\mathfrak{g}})$ defined as $\bar{\gamma}(t) = \gamma(t) - A$ for each $t \in (-\epsilon, \epsilon)$

$T^\star\mathcal{C}(P) = \mathcal{C}(P) \times \Lambda^1_m(M, \tilde{\mathfrak{g}}^*) \times \Lambda^1_{m-1}(\partial M, \tilde{\mathfrak{g}}^*)$ given by

$$\langle (\varsigma, \varsigma_\partial), \dot{A} \rangle_\star = \int_M \varsigma \cdot \dot{A} + \int_{\partial M} \varsigma_\partial \cdot \imath^*_{\partial M} \dot{A}$$

$(\varsigma, \varsigma_\partial) \in \Lambda^1_m(M, \tilde{\mathfrak{g}}^*) \times \Lambda^1_{m-1}(\partial M, \tilde{\mathfrak{g}}^*)$ and $\dot{A} \in \Omega^1(M, \tilde{\mathfrak{g}})$

$$T^\dagger\mathcal{C}(P) = \mathcal{C}(P) \times \Omega^{m-1}(M, \tilde{\mathfrak{g}}^*) \times \Omega^{m-2}(\partial M, \tilde{\mathfrak{g}}^*)$$

$$\langle (\varsigma, \varsigma_\partial), \dot{A} \rangle_\dagger = \int_M \dot{A} \wedge \varsigma + \int_{\partial M} \imath^*_{\partial M} \dot{A} \wedge \varsigma_\partial$$

$(\varsigma, \varsigma_\partial) \in \Omega^{m-1}(M, \tilde{\mathfrak{g}}^*) \times \Omega^{m-2}(\partial M, \tilde{\mathfrak{g}}^*)$ and $\dot{A} \in \Omega^1(M, \tilde{\mathfrak{g}})$

$$\mathrm{d}^{\mathbf{A}}: \Omega^k(P, \mathfrak{g}) \to \Omega^{k+1}(P, \mathfrak{g})$$

$$\mathrm{d}^{\mathbf{A}}\eta = \mathrm{d}\eta + [\mathbf{A}, \eta] + [\eta, \mathbf{A}], \eta \in \bar{\Omega}^1(P, \mathfrak{g})$$

$$\mathrm{d}^{A}: \Omega^k(M, \tilde{\mathfrak{g}}) \to \Omega^{k+1}(M, \tilde{\mathfrak{g}})$$

$$\mathrm{d}^{A*}: \Omega^k(M, \tilde{\mathfrak{g}}^*) \to \Omega^{k+1}(M, \tilde{\mathfrak{g}}^*)$$

$$\mathrm{div}^{A*}: \Lambda^{k+1}_m(M, \tilde{\mathfrak{g}}^*) \to \Lambda^k_m(M, \tilde{\mathfrak{g}}^*)$$

the curvature $\mathbf{A} \in \Omega^1(P, \mathfrak{g})$ is given by $B_{\mathbf{A}} = \mathrm{d}^{\mathbf{A}}\mathbf{A} \in \bar{\Omega}^2(P, \mathfrak{g})$ and $B_A \in \Omega^2(M, \tilde{\mathfrak{g}})$

$$\mathcal{L}_{\mathrm{gau}}: \mathrm{Conn}(P) \times_M (T^*M \otimes \tilde{\mathfrak{g}}) \times_M (\wedge^2 T^*M \otimes \tilde{\mathfrak{g}}) \to \wedge^m T^*M$$

$$\frac{\partial \mathcal{L}_{\mathrm{gau}}}{\partial A}: \mathrm{Conn}(P) \times_M (T^*M \otimes \tilde{\mathfrak{g}}) \times_M (\wedge^2 T^*M \otimes \tilde{\mathfrak{g}}) \to TM \otimes \wedge^m T^*M \otimes \tilde{\mathfrak{g}}^*$$

$$\frac{\partial \mathcal{L}_{\mathrm{gau}}}{\partial A}(A_x, \varepsilon_x, \beta_x) \cdot \delta A_x = \frac{d}{d\epsilon}\Big|_{\epsilon=0} \mathcal{L}_{\mathrm{gau}}(A_x + \epsilon \delta A_x, \varepsilon_x, \beta_x), \delta A_x \in T^*_x M \otimes \tilde{\mathfrak{g}}_x$$

$x \in M$ and $(A_x, \varepsilon_x, \beta_x) \in \mathrm{Conn}(P)_x \times (T^*M \otimes \tilde{\mathfrak{g}})_x \times (\wedge^2 T^*M \otimes \tilde{\mathfrak{g}})_x$

$\partial \mathcal{L}_{\mathrm{gau}} / \partial \varepsilon$ and $\partial \mathcal{L}_{\mathrm{gau}} / \partial \beta$

$$\frac{\partial \mathcal{L}}{\partial A}: \mathrm{Conn}(P) \times_M (T^*M \otimes \tilde{\mathfrak{g}}) \times_M (\wedge^2 T^*M \otimes \tilde{\mathfrak{g}}) \to \wedge^{m-1} T^*M \otimes \tilde{\mathfrak{g}}^*$$

$$\delta A_x \wedge \frac{\partial \mathcal{L}_{\text{gau}}}{\partial A}(A_x, \varepsilon_x, \beta_x) = \frac{d}{d\epsilon}\Big|_{\epsilon=0} \mathcal{L}_{\text{gau}}(A_x + \epsilon\delta A_x, \varepsilon_x, \beta_x), \delta A_x \in T_x^* M \otimes \tilde{\mathfrak{g}}_x$$

$\partial \mathcal{L}_{\text{gau}} / \partial \varepsilon$ and $\partial \mathcal{L}_{\text{gau}} / \partial \beta$ are defined analogously

$$L_{\text{gau}} : T\mathcal{C}(P) \simeq \mathcal{C}(P) \times \Omega^1(M, \tilde{\mathfrak{g}}) \to \mathbb{R}, (A, \varepsilon) \mapsto \int_M \mathcal{L}_{\text{gau}}(A, \varepsilon, B_A)$$

$$\frac{\delta L_{\text{gau}}}{\delta A}, \frac{\delta L_{\text{gau}}}{\delta \varepsilon} : T\mathcal{C}(P) \to \Omega^1(M, \tilde{\mathfrak{g}})'$$

$$\frac{\delta L_{\text{gau}}}{\delta A}(A, \varepsilon)(\delta A) = \frac{d}{d\epsilon}\Big|_{\epsilon=0} L_{\text{gau}}(A + \epsilon\delta A, \varepsilon), \frac{\delta L_{\text{gau}}}{\delta \varepsilon}(A, \varepsilon)(\delta \varepsilon) = \frac{d}{d\epsilon}\Big|_{\epsilon=0} L_{\text{gau}}(A, \varepsilon + \epsilon\delta\varepsilon)$$

$$(A, \varepsilon) \in T\mathcal{C}(P) \text{ and } \delta A, \delta\varepsilon \in \Omega^1(M, \tilde{\mathfrak{g}})$$

$$(A, \varepsilon) \in T\mathcal{C}(P) \simeq \mathcal{C}(P) \times \Omega^1(M, \tilde{\mathfrak{g}})$$

$$\frac{\delta L_{\text{gau}}}{\delta A}(A, \varepsilon) = \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial A}(A, \varepsilon, B_A) + \text{div}^{A*}\left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A, \varepsilon, B_A) \right), \iota_{\partial M}^* \left(\text{tr} \frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A, \varepsilon, B_A) \right) \right)$$

$$\frac{\delta L_{\text{gau}}}{\delta \varepsilon}(A, \varepsilon) = \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \varepsilon}(A, \varepsilon, B_A), 0 \right)$$

$$\frac{\delta L_{\text{gau}}}{\delta A}(A, \varepsilon) = \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial A}(A, \varepsilon, B_A) + \text{d}^{A*}\left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A, \varepsilon, B_A) \right), \iota_{\partial M}^* \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A, \varepsilon, B_A) \right) \right)$$

$$\frac{\delta L_{\text{gau}}}{\delta \varepsilon}(A, \varepsilon) = \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \varepsilon}(A, \varepsilon, B_A), 0 \right)$$

the curvature is $B_{\mathbf{A}} = \text{d}\mathbf{A} + [\mathbf{A}, \mathbf{A}]$, where $[\mathbf{A}, \mathbf{A}](u, v) = [\mathbf{A}(u), \mathbf{A}(v)]$ for each $u, v \in \mathfrak{X}(P)$

$$\begin{aligned} \frac{d}{d\epsilon}\Big|_{\epsilon=0} B_{\mathbf{A}+\epsilon\delta\mathbf{A}} &= \frac{d}{d\epsilon}\Big|_{\epsilon=0} \text{d}(\mathbf{A} + \epsilon\delta\mathbf{A}) + [\mathbf{A} + \epsilon\delta\mathbf{A}, \mathbf{A} + \epsilon\delta\mathbf{A}] \\ &= \text{d}(\delta\mathbf{A}) + [\mathbf{A}, \delta\mathbf{A}] + [\delta\mathbf{A}, \mathbf{A}] \\ &= \text{d}^{\mathbf{A}}(\delta\mathbf{A}) \end{aligned}$$

$$d/d\epsilon|_{\epsilon=0} B_{A+\epsilon\delta A} = \text{d}^A(\delta A)$$

$$\begin{aligned} \frac{\delta L_{\text{gau}}}{\delta A}(A, \varepsilon)(\delta A) &= \frac{d}{d\epsilon}\Big|_{\epsilon=0} \int_M \mathcal{L}_{\text{gau}}(A + \epsilon\delta A, \varepsilon, B_{A+\epsilon\delta A}) \\ &= \int_M \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial A} \cdot \delta A + \frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta} \cdot \text{d}^A(\delta A) \right) \\ &= \int_M \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial A} + \text{div}^{A*}\left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta} \right) \right) \cdot \delta A + \int_{\partial M} \iota_{\partial M}^* \left(\text{tr} \frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta} \right) \cdot \iota_{\partial M}^* \delta A \end{aligned}$$

$$\text{d}L_{\text{gau}} : T\mathcal{C}(P) \to T'(T\mathcal{C}(P)), (A, \varepsilon) \mapsto \left(A, \varepsilon, \frac{\delta L_{\text{gau}}}{\delta A}(A, \varepsilon), \frac{\delta L_{\text{gau}}}{\delta \varepsilon}(A, \nu) \right)$$

$$\hat{F}^{\star}: T\mathcal{C}(P) \to T^{\star}\mathcal{C}(P), (A,\varepsilon) \mapsto \left(A, \hat{\mathcal{F}}^{\star}(A,\varepsilon), \hat{\mathcal{F}}_{\partial}^{\star}(A,\varepsilon)\right)$$

$$\hat{F}^{\dagger}: T\mathcal{C}(P) \to T^{\dagger}\mathcal{C}(P), (A,\varepsilon) \mapsto \left(A, \hat{\mathcal{F}}^{\dagger}(A,\varepsilon), \hat{\mathcal{F}}_{\partial}^{\dagger}(A,\varepsilon)\right)$$

$$L_{\text{gau}}: T\mathcal{C}(P) \simeq \mathcal{C}(P) \times \Omega^1(M, \tilde{\mathfrak{g}}) \to \mathbb{R}$$

for a curve $(A, \varepsilon, \varsigma, \varsigma_{\partial}): [t_0, t_1] \to T\mathcal{C}(P) \oplus T^{\star}\mathcal{C}(P)$

$$\begin{cases} \dot{A} = \varepsilon \\ \varsigma = \dfrac{\partial \mathcal{L}_{\text{gau}}}{\partial \varepsilon}(A,\varepsilon,B_A), \quad \dot{\varsigma} = \dfrac{\partial \mathcal{L}_{\text{gau}}}{\partial A}(A,\varepsilon,B_A) + \operatorname{div}^{A*}\left(\dfrac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A,\varepsilon,B_A)\right) + \hat{\mathcal{F}}^{\star}(A,\varepsilon) \\ \varsigma_{\partial} = 0, \qquad\qquad \dot{\varsigma}\partial = \iota_{\partial M}^{*}\left(\operatorname{tr}\dfrac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A,\varepsilon,B_A)\right) + \hat{\mathcal{F}}_{\partial}^{\star}(A,\varepsilon) \end{cases}$$

for a curve $(A, \varepsilon, \varsigma, \varsigma_{\partial}): [t_0, t_1] \to T\mathcal{C}(P) \oplus T^{\dagger}\mathcal{C}(P)$

$$\begin{cases} \dot{A} = \varepsilon, \\ \varsigma = \dfrac{\partial \mathcal{L}_{\text{gau}}}{\partial \varepsilon}(A,\varepsilon,B_A), \quad \dot{\varsigma} = \dfrac{\partial \mathcal{L}_{\text{gau}}}{\partial A}(A,\varepsilon,B_A) + \mathrm{d}^{A*}\left(\dfrac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A,\varepsilon,B_A)\right) + \hat{\mathcal{F}}^{\dagger}(A,\varepsilon) \\ \varsigma_{\partial} = 0, \qquad\qquad \dot{\varsigma_{\partial}} = \iota_{\partial M}^{*}\left(\dfrac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A,\varepsilon,B_A)\right) + \hat{\mathcal{F}}_{\partial}^{\dagger}(A,\varepsilon) \end{cases}$$

$$dA/dt \simeq (A, \dot{A}): [t_0, t_1] \to T\mathcal{C}(P) \simeq \mathcal{C}(P) \times \Omega^1(M, \tilde{\mathfrak{g}})$$

$$\begin{aligned} \mathcal{E}_{\text{gau}}(A_x, \varepsilon_x, \beta_x) &= \frac{\partial \mathcal{L}_{\text{gau}}}{\partial \varepsilon}(A_x, \varepsilon_x, \beta_x) \cdot \varepsilon_x - \mathcal{L}_{\text{gau}}(A_x, \varepsilon_x, \beta_x) \\ &= \varepsilon_x \wedge \frac{\partial \mathcal{L}_{\text{gau}}}{\partial \varepsilon}(A_x, \varepsilon_x, \beta_x) - \mathcal{L}_{\text{gau}}(A_x, \varepsilon_x, \beta_x) \end{aligned}$$

$$(A_x, \varepsilon_x, \beta_x) \in \operatorname{Conn}(P)_x \times (T^*M \otimes \tilde{\mathfrak{g}})_x \times (\Lambda^2 T^*M \otimes \tilde{\mathfrak{g}})_x$$

$$T^{\star}\mathcal{C}(P) = \mathcal{C}(P) \times \Lambda_m^1(M, \tilde{\mathfrak{g}}^*) \times \Lambda_{m-1}^1(\partial M, \tilde{\mathfrak{g}}^*), \text{ let } (A, \varepsilon, \varsigma, \varsigma_{\partial}): [t_0, t_1] \to T\mathcal{C}(P) \oplus T^{\star}\mathcal{C}(P)$$

$$\frac{\partial}{\partial t}\mathcal{E}_{\text{gau}}(A,\varepsilon,B_A) = -\operatorname{div}\left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A,\varepsilon,B_A) \cdot \varepsilon\right) + \hat{\mathcal{F}}^{\star}(A,\varepsilon) \cdot \varepsilon$$

$$\frac{d}{dt}\int_M \mathcal{E}_{\text{gau}}(A,\varepsilon,B_A) = \underbrace{\int_M \hat{\mathcal{F}}^{\star}(A,\varepsilon) \cdot \varepsilon}_{\text{spatially distributed contribution}} + \underbrace{\int_{\partial M} \hat{\mathcal{F}}_{\partial}^{\star}(A,\varepsilon) \cdot \iota_{\partial M}^{*}\varepsilon}_{\text{boundary contribution}}$$

$$T^{\dagger}\mathcal{C}(P) = \mathcal{C}(P) \times \Omega^{m-1}(M, \tilde{\mathfrak{g}}^*) \times \Omega^{m-2}(\partial M, \tilde{\mathfrak{g}}^*), \text{ let } (A, \varepsilon, \varsigma, \varsigma_{\partial}): [t_0, t_1] \to T\mathcal{C}(P) \oplus T^{\dagger}\mathcal{C}(P)$$

$$\frac{\partial}{\partial t}\mathcal{E}_{\text{gau}}(A,\varepsilon,B_A) = -\mathrm{d}\left(\varepsilon \wedge \frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A,\varepsilon,B_A)\right) + \varepsilon \wedge \hat{\mathcal{F}}^{\dagger}(A,\varepsilon)$$

$$\frac{d}{dt}\int_M \mathcal{E}_{\text{gau}}(A,\varepsilon,B_A) = \underbrace{\int_M \varepsilon \wedge \hat{\mathcal{F}}^{\dagger}(A,\varepsilon)}_{\text{spatially distributed contribution}} + \underbrace{\int_{\partial M} \iota_{\partial M}^{*}\varepsilon \wedge \hat{\mathcal{F}}_{\partial}^{\dagger}(A,\varepsilon)}_{\text{boundary contribution}}$$

vector field density $S = \frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta} \cdot \varepsilon = \frac{\partial \mathcal{L}_{\text{gau}}}{\partial B_A} \cdot \dot{A}$ or the $(m-1)$-form $\mathrm{S} = \varepsilon \wedge \frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta} = \dot{A} \wedge \frac{\partial \mathcal{L}_{\text{gau}}}{\partial B_A}$

$$\hat{\mathcal{F}}_{\partial}^{\star} \in \Lambda_{m-1}^{1}(\partial M, \tilde{\mathfrak{g}}^{*}) \text{ and } \iota_{\partial M}^{*}\varepsilon = \iota_{\partial M}^{*}\dot{A} \in \Omega^{1}(\partial M, \tilde{\mathfrak{g}})$$

$$\hat{\mathcal{F}}^{\star} \in \Lambda_{m}^{1}(M, E^{*}) \text{ and } \varepsilon = \dot{A} \in \Omega^{1}(M, E)$$

$$\hat{\mathcal{F}}_{\partial}^{\dagger} \in \Omega^{m-2}(\partial M, \tilde{\mathfrak{g}}^{*}) \text{ and } \hat{\mathcal{F}}^{\dagger} \in \Omega^{m-2}(\partial M, \tilde{\mathfrak{g}}^{*})$$

$$\begin{cases} \frac{\partial}{\partial t}\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \varepsilon}(A, \dot{A}, B_A) = \frac{\partial \mathcal{L}_{\text{gau}}}{\partial A}(A, \dot{A}, B_A) + \operatorname{div}^{A_*}\left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A, \dot{A}, B_A)\right) + \hat{\mathcal{F}}^{\star}(A, \dot{A}), \\ \hat{\mathcal{F}}_{\partial}^{\star}(A, \dot{A}) = -\iota_{\partial M}^{*}\left(\operatorname{tr}\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A, \dot{A}, B_A)\right), \end{cases}$$

$$\begin{cases} \frac{\partial}{\partial t}\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \varepsilon}(A, \dot{A}, B_A) = \frac{\partial \mathcal{L}_{\text{gau}}}{\partial A}(A, \dot{A}, B_A) + \mathrm{d}^{A_*}\left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A, \dot{A}, B_A)\right) + \hat{\mathcal{F}}^{\dagger}(A, \dot{A}) \\ \hat{\mathcal{F}}_{\partial}^{\dagger}(A, \dot{A}) = -\iota_{\partial M}^{*}\left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}(A, \dot{A}, B_A)\right) \end{cases}$$

$$\mathbf{g}\colon \left(\Lambda^{k} T^{*}M \otimes \tilde{\mathfrak{g}}\right) \times_{M} \left(\Lambda^{m} T^{*}M \otimes \tilde{\mathfrak{g}}\right) \to \mathbb{R},$$

$$\mathbf{g}_x(\alpha_1 \otimes \sigma_1, \alpha_2 \otimes \sigma_2) = -g_x(\alpha_1, \alpha_2)K(\sigma_1, \sigma_2), x \in M, \alpha_1, \alpha_2 \in \Lambda^{k} T_x^{*}M, \sigma_1, \sigma_2 \in \tilde{\mathfrak{g}}_x,$$

$$\begin{aligned} \sharp_k &\colon \Lambda^{k} T^{*}M \otimes \tilde{\mathfrak{g}} \to \Lambda^{k} TM \otimes \tilde{\mathfrak{g}}^{*}, \quad \alpha \otimes \sigma \mapsto \alpha^{\sharp_k} \otimes \sigma^{\mathfrak{6}_k}, \\ \mathfrak{6} &\colon \Lambda^{k} TM \otimes \tilde{\mathfrak{g}}^{*} \to \Lambda^{k} T^{*}M \otimes \tilde{\mathfrak{g}}, \quad U \otimes \eta \mapsto U^{\mathfrak{6}} \otimes \eta^{\sharp_\kappa}. \end{aligned}$$

$E = \tilde{\mathfrak{g}}$ or $E = \tilde{\mathfrak{g}}^{*}$ for each $\alpha \in \Lambda^{k} T^{*}M$ and $\sigma \in E$ as $\star\,(\alpha \otimes \sigma) = (\star\,\alpha) \otimes \sigma$, where $\star\colon \Lambda^{k} T^{*}M \to$

$\Lambda^{m-k} T^{*}M$ is the standard Hodge star operator.

$\bar{\pi}_{P,M}\colon \bar{P} \to \bar{M}$, where $\bar{P} = [t_0, t_1] \times P$ and $\bar{M} = [t_0, t_1] \times M$

$\eta = -dt \otimes dt + g$, where $t \in [t_0, t_1]$ is the coordinate system.

$$\boldsymbol{\eta}\colon \left(\Lambda^{k} T^{*}\bar{M} \otimes ([t_0, t_1] \times \tilde{\mathfrak{g}})\right) \times \bar{M}\left(\Lambda^{m} T^{*}\bar{M} \otimes ([t_0, t_1] \times \tilde{\mathfrak{g}})\right) \to \mathbb{R}$$

$\bar{\pi}_{P,M}\colon \bar{P} \to \bar{M}$ is given by $(\bar{P} \times \mathfrak{g}) / G = [t_0, t_1] \times \tilde{\mathfrak{g}}$

$$\boldsymbol{\eta}_{(t,x)}(\bar{\alpha}_1 \otimes \bar{\sigma}_1, \bar{\alpha}_2 \otimes \bar{\sigma}_2) = \eta_{(t,x)}(\bar{\alpha}_1, \bar{\alpha}_2)K(\sigma_1, \sigma_2)$$

$$(t, x) \in \bar{M}, \bar{\alpha}_i \in \Lambda^{k} T_{(t,x)}^{*}\bar{M} \text{ and } \bar{\sigma}_i = (t, \sigma_i) \in ([t_0, t_1] \times \tilde{\mathfrak{g}})_{(t,x)}, i = 1,2$$

$$\eta_{(t,x)}(\bar{\alpha}_1, \bar{\alpha}_2) = \det\begin{pmatrix} \eta_{(t,x)}(dt, dt) & \eta_{(t,x)}(dt, \alpha_2) \\ \eta_{(t,x)}(\alpha_1, dt) & \eta_{(t,x)}(\alpha_1, \alpha_2) \end{pmatrix} = \det\begin{pmatrix} -1 & 0 \\ 0 & g_x(\alpha_1, \alpha_2) \end{pmatrix} = -g_x(\alpha_1, \alpha_2)$$

$$\boldsymbol{\eta}_{(t,x)}(\bar{\alpha}_1 \otimes \bar{\sigma}_1, \bar{\alpha}_2 \otimes \bar{\sigma}_2) = \mathbf{g}_x(\alpha_1 \otimes \sigma_1, \alpha_2 \otimes \sigma_2)$$

$$\bar{\alpha}_i = \alpha_i^1 \wedge \alpha_i^2 \in \Lambda^2 T_x^{*}M, i = 1,2$$

$$\eta_{(t,x)}(\bar{\alpha}_1,\bar{\alpha}_2)=\det\begin{pmatrix}\eta_{(t,x)}(\alpha_1^1,\alpha_2^1) & \eta_{(t,x)}(\alpha_1^1,\alpha_2^2)\\ \eta_{(t,x)}(\alpha_1^2,\alpha_2^1) & \eta_{(t,x)}(\alpha_1^2,\alpha_2^2)\end{pmatrix}$$
$$=\det\begin{pmatrix}g_x(\alpha_1^1,\alpha_2^1) & g_x(\alpha_1^1,\alpha_2^2)\\ g_x(\alpha_1^2,\alpha_2^1) & g_x(\alpha_1^2,\alpha_2^2)\end{pmatrix}$$
$$=g_x(\alpha_1,\alpha_2)$$

$$\boldsymbol{\eta}_{(t,x)}(\bar{\alpha}_1\otimes\bar{\sigma}_1,\bar{\alpha}_2\otimes\bar{\sigma}_2)=-\mathbf{g}_x(\alpha_1\otimes\sigma_1,\alpha_2\otimes\sigma_2)$$

$\bar{\pi}_{P,M}\colon \bar{P}\to\bar{M}$ is denoted by $\mathrm{Conn}(\bar{P})=(J^1\bar{P})/G\to\bar{M}$

$$\mathcal{C}(\bar{P})\ni \mathrm{A}\overset{\blacksquare:\blacksquare}{\leftrightarrow}\mathcal{A}\in\Omega^1(\bar{P},\mathfrak{g})$$

where $\mathcal{C}(\bar{P})=\Gamma(\mathrm{Conn}(\bar{P})\to\bar{M})$.

$$\Omega^1(\bar{M},[t_0,t_1]\times\tilde{\mathfrak{g}})$$

$$\mathrm{d}^{\mathrm{A}}\colon\Omega^k(\bar{M},[t_0,t_1]\times\tilde{\mathfrak{g}})\to\Omega^{k+1}(\bar{M},[t_0,t_1]\times\tilde{\mathfrak{g}})$$

$$\delta^{\mathrm{A}}\colon\Omega^k(\bar{M},[t_0,t_1]\times\tilde{\mathfrak{g}})\to\Omega^{k-1}(\bar{M},[t_0,t_1]\times\tilde{\mathfrak{g}})$$

$$\bar{\star}\colon\bigwedge^k T^*\bar{M}\to\bigwedge^{m+1-k}T^*\bar{M}$$

$$\eta=-dt\otimes dt+g \text{ on } \bar{M}$$

$$\mathrm{d}^{\mathcal{A}}\colon\Omega^k(\bar{P},\mathfrak{g})\to\Omega^{k+1}(\bar{P},\mathfrak{g})$$

$$\mathcal{L}_{\mathrm{YM}}\colon\mathrm{Conn}(\bar{P})\times_{\bar{M}}\left(\bigwedge^2 T^*\bar{M}\otimes([t_0,t_1]\times\tilde{\mathfrak{g}})\right)\to\bigwedge^{m+1}T^*\bar{M}$$

$$\mathcal{L}_{\mathrm{YM}}\left(\mathrm{A}_{(t,x)},\mathrm{B}_{(t,x)}\right)=\frac{1}{2}\boldsymbol{\eta}\left(\mathrm{B}_{(t,x)},\mathrm{B}_{(t,x)}\right)dt\wedge\mu_g$$

$\mathcal{A}(t,x)=\mathbf{A}(t,x)+\mathbf{A}_0(t,x)dt$ for each $(t,x)\in\bar{M}$, where $\mathbf{A}(t,\cdot)\in\Omega^1(P,\mathfrak{g})$

$$\mathbf{A}_0(t,\cdot)\in C_G^\infty(P,\mathfrak{g})$$

$$\mathrm{B}_{\mathcal{A}}=\mathrm{d}^{\mathcal{A}}\mathcal{A}=\mathrm{d}^{\mathbf{A}}\mathbf{A}+\left(-\dot{\mathbf{A}}+\mathrm{d}^{\mathbf{A}}\mathbf{A}_0\right)\wedge dt\in\bar{\Omega}^2(\bar{P},\mathfrak{g})$$

$$\mathcal{A}_{\mathrm{YM}}(\mathrm{A})=\frac{1}{2}\int_{\bar{M}}\boldsymbol{\eta}\left(\mathrm{B}_{\mathrm{A}},\mathrm{B}_{\mathrm{A}}\right)dt\wedge\mu_g=\frac{1}{2}\int_{t_0}^{t_1}\int_M\left(\mathbf{g}(\dot{A},\dot{A})-\mathbf{g}(B_A,B_A)\right)\mu_g dt$$

$$\mathcal{L}_{\mathrm{YM}}(A_x,\varepsilon_x,\beta_x)=\frac{1}{2}\left(\mathbf{g}(\varepsilon_x,\varepsilon_x)-\mathbf{g}(\beta_x,\beta_x)\right)\mu_g.$$

$E(t)=-\dot{A}(t,\blacksquare)\in\Omega^1(M,\tilde{\mathfrak{g}})$ and $B_{A(t,\blacksquare)}\in\Omega^2(M,\tilde{\mathfrak{g}})$

$$b_K\circ\mathrm{d}^A=\mathrm{d}^{A^*}\circ b_K$$

$A=A_i^a dx^i\otimes\hat{\xi}_a$ and $K=K_{ab}\hat{\xi}^a\otimes\hat{\xi}^b$

$A_i^a,K_{ab}\in C^\infty(M)$ such that $K_{ab}=K_{ba},1\le a,b\le n,1\le i\le m$

$$K_{ab} = \epsilon_a \delta_{ab}, 1 \leq a, b \leq n, \text{ where } \epsilon_a \in \{-1,1\}.$$

$$-f_{bc}^a = K\big(\hat{\xi}_a, f_{bc}^d \hat{\xi}_d\big) = K\big(\hat{\xi}_a, [\hat{\xi}_b, \hat{\xi}_c]\big) = K\big([\hat{\xi}_a, \hat{\xi}_b], \hat{\xi}_c\big) = K\big(f_{ab}^d \hat{\xi}_d, \hat{\xi}_c\big) = -f_{ab}^c$$

$$f_{bc}^a \in \mathbb{R}, 1 \leq a, b, c \leq n$$

$$f_{bc}^a = -f_{ba}^c \text{ for each } 1 \leq a, b, c \leq n$$

$$\xi_1 = \xi_1^a \hat{\xi}_a, \xi_2 = \xi_2^a \hat{\xi}_a \in \Gamma(\tilde{\mathfrak{g}} \to M)$$

$$\nabla^A \xi_1 = \big(\partial_i \xi_1^a + f_{bc}^a A_i^b \xi_1^c\big) dx^i \otimes \hat{\xi}_a$$

$$\begin{aligned}
\mathrm{d}\big(K(\xi_1, \xi_2)\big) &= -\mathrm{d}(\xi_1^a \xi_2^a) = -\partial_i (\xi_1^a \xi_2^a) dx^i \\
&= -\big((\partial_i \xi_1^a)\xi_2^a + \xi_1^a (\partial_i \xi_2^a)\big) dx^i \\
&= -\Big(\big(\partial_i \xi_1^a - f_{bc}^a A_i^b \xi_1^a\big)\xi_2^c + \xi_1^a \big(\partial_i \xi_2^a + f_{bc}^a A_i^b \xi_2^c\big)\Big) dx^i \\
&= -\Big(\big(\partial_i \xi_1^a + f_{bc}^a A_i^b \xi_1^c\big)\xi_2^a + \xi_1^a \big(\partial_i \xi_2^a + f_{bc}^a A_i^b \xi_2^c\big)\Big) dx^i \\
&= K(\nabla^A \xi_1, \xi_2) + K(\xi_1, \nabla^A \xi_2)
\end{aligned}$$

$$(A, \varepsilon) \in T\mathcal{C}(P) \simeq \mathcal{C}(P) \times \Omega^1(M, \tilde{\mathfrak{g}})$$

$$\begin{cases}
\dfrac{\partial \mathcal{L}_{\mathrm{YM}}}{\partial A} = 0, & \dfrac{\partial \mathcal{L}_{\mathrm{YM}}}{\partial \varepsilon} = -E^{\sharp_\partial} \otimes \mu_g, & \dfrac{\partial \mathcal{L}_{\mathrm{YM}}}{\partial \beta} = -B_A^{\sharp_\partial} \otimes \mu_g, \\
\dfrac{\partial \mathcal{L}_{\mathrm{YM}}}{\partial A} = 0, & \dfrac{\partial \mathcal{L}_{\mathrm{YM}}}{\partial \varepsilon} = - \star E^{\mathfrak{G}_k}, & \dfrac{\partial \mathcal{L}_{\mathrm{YM}}}{\partial \beta} = - \star B_A^{\mathfrak{G}_k}.
\end{cases}$$

$$A = A_i^a dx^i \otimes \hat{\xi}_a \text{ and } E = -\varepsilon = -\varepsilon_i^a dx^i \otimes \hat{\xi}_a$$

$$A_i^a, \varepsilon_i^a \in C^\infty(M), 1 \leq i \leq m, 1 \leq a \leq n$$

$$B_A = \big(\partial_i A_j^a + \Gamma_{i,b}^a A_j^b\big) dx^i \wedge dx^j \otimes \hat{\xi}_a$$

$\Gamma_{i,b}^a \in C^\infty(M), 1 \leq i \leq m, 1 \leq a, b \leq n$, are the Christoffel symbols of ∇^A, i.e., $\Gamma_{i,b}^a = f_{cb}^a A_i^c$.

Riemannian metric and the Killing form $g = g_{ij} dx^i \otimes dx^j$ and $K = -\delta_{ab} \hat{\xi}^a \otimes \hat{\xi}^b$

$$\mu_g = \sqrt{|g|} d^m x \in \Omega^m(M)$$

$$\star\big(dx^i \otimes \hat{\xi}_a\big) = \sqrt{|g|} g^{ij} d_{(j)}^{m-1} x \otimes \hat{\xi}_a, \star\big(dx^i \wedge dx^j \otimes \hat{\xi}_a\big) = \sqrt{|g|} g^{ik} g^{jl} d_{(k,l)}^{m-2} x \otimes \hat{\xi}_a$$

$$E^{\sharp_\partial} = -g^{ij} \varepsilon_j^a \partial_i \otimes \hat{\xi}^a, B_A^\sharp = g^{ik} g^{jl} \big(\partial_i A_j^a + \Gamma_{i,c}^a A_j^c\big) \partial_k \wedge \partial_l \otimes \hat{\xi}^a.$$

$$\begin{cases}
\dfrac{\partial \mathcal{L}_{\mathrm{YM}}}{\partial A} = \Phi_E \circ \dfrac{\partial \mathcal{L}_{\mathrm{YM}}}{\partial A} = \Phi_E \\
\dfrac{\partial \mathcal{L}_{\mathrm{YM}}}{\partial \varepsilon} = \Phi_E \circ \big(-E^{\sharp_\partial} \otimes \mu_g\big) = \sqrt{|g|} g^{ij} \varepsilon_j^a d_{(i)}^{m-1} x \otimes \hat{\xi}^a = -(\star E)^{\mathfrak{G}_k}, \\
\dfrac{\partial \mathcal{L}_{\mathrm{YM}}}{\partial \beta} = \Phi_E \circ \Big(-B_A^{\sharp_\partial} \otimes \mu_g\Big) = -\sqrt{|g|} g^{ik} g^{jl} \big(\partial_i A_j^a + \Gamma_{i,c}^a A_j^c\big) d_{(k,l)}^{m-2} x \otimes \hat{\xi}^a = -(\star B_A)^{\mathfrak{G}_k}.
\end{cases}$$

$$\begin{cases} \dot{A} = -E, \\ \varsigma = -E^{\sharp_\partial} \otimes \mu_g, & \dot{\varsigma} = -\mathrm{div}^{A*}\left(B_A^{\sharp_\partial} \otimes \mu_g\right) + \hat{\mathcal{F}}^\star, \\ \varsigma_\partial = 0, & \dot{\varsigma}_\partial = -\iota_{\partial M}^*\left(\mathrm{tr}\left(B_A^{\sharp_\partial} \otimes \mu_g\right)\right) + \hat{\mathcal{F}}_\partial^\star. \end{cases}$$

$$\begin{cases} \dot{A} = -E, \\ \varsigma = - \star E^{\mathfrak{G}_K}, & \dot{\varsigma} = -\mathrm{d}^{A*}\left(\star B_A^{\mathfrak{G}_K}\right) + \hat{\mathcal{F}}^\dagger, \\ \varsigma_\partial = 0, & \dot{\varsigma}_\partial = -\iota_{\partial M}^*\left(\star B_A^{\mathfrak{G}_K}\right) + \hat{\mathcal{F}}_\partial^\dagger. \end{cases}$$

$\star\, \hat{\mathcal{F}}^\dagger = -J^{b_K}$ and $\star_\partial\, \hat{\mathcal{F}}_\partial^\dagger = j^{b_K}$, where $J: T\mathcal{C}(P) \to \Omega^1(M, \tilde{\mathfrak{g}})$ and $j: T\mathcal{C}(P) \to \Omega^1(\partial M, \tilde{\mathfrak{g}})$

$$\dot{E} - \delta^A B_A = -J, \star_\partial \left(\iota_{\partial M}^*(\star\, B_A)\right) = j,$$

$$\delta^A = (-1)^{m+1} \star \circ\, \mathrm{d}^A \circ \star : \Omega^2(M, \tilde{\mathfrak{g}}) \to \Omega^1(M, \tilde{\mathfrak{g}})$$

$$\mathcal{E}_{\mathrm{YM}}(A, E, B_A) = \frac{1}{2}\big(\mathbf{g}(E, E) + \mathbf{g}(B_A, B_A)\big)\mu_g.$$

$$\frac{\partial \mathcal{E}_{\mathrm{YM}}}{\partial t} = -\mathrm{dS} + E \wedge \star\, J^{\mathfrak{G}_K}$$

$$\frac{d}{dt}\int_M \mathcal{E}_{\mathrm{YM}} = \int_M E \wedge \star\, J^{\mathfrak{G}_K} + \int_{\partial M} \left(\star_\partial\, j^{\mathfrak{G}_K}\right) \wedge \iota_{\partial M}^* E$$

$$\mathrm{S} = E \wedge \star\, B_A^{b_K} \in \Omega^{m-1}(M)$$

$$\mathbf{S} = (\star\, \mathrm{S})^{\sharp_\partial} \in \mathfrak{X}(M)$$

$$\mathrm{J} = J + \rho dt \in \Omega^1(\bar{M}, [t_0, t_1] \times \tilde{\mathfrak{g}})$$

$$\boldsymbol{\rho} \in C_G^\infty(\bar{P}, \mathfrak{g}) \text{ and } \mathbf{J} \in \bar{\Omega}^1(\bar{P}, \mathfrak{g})$$

$$\delta^{\mathrm{A}}\, \mathrm{B_A} = \mathrm{J},\ \mathrm{d}^{\mathrm{A}}\, \mathrm{B_A} = 0, \delta^{\mathrm{A}}\, \mathrm{J} = 0$$

$$\delta^{\mathrm{A}}\, \mathrm{B_A} = \varpi_1 dt + \varpi_2 \in \Omega^1(\bar{M}, [t_0, t_1] \times \tilde{\mathfrak{g}})$$

where $\varpi_1 \in \Omega^0(\bar{M}, [t_0, t_1] \times \tilde{\mathfrak{g}})$ and $\varpi_2(t, \cdot) \in \Omega^1(M, \tilde{\mathfrak{g}})$ for each $t \in [t_0, t_1]$

$$\mathrm{B_A}(t, \cdot) = B_{A(t,\cdot)} + E(t, \cdot) \wedge dt, t \in [t_0, t_1]$$

$$\alpha \in \Omega^0(M, \tilde{\mathfrak{g}}) = C^\infty(M, \tilde{\mathfrak{g}})\mathrm{cl}(\mathrm{supp}\alpha) \subset \mathrm{int}M = M - \partial M$$

$$\begin{aligned} \boldsymbol{\eta}(E \wedge dt,\ \mathrm{d}^A\alpha \wedge dt) &= \det\begin{pmatrix} \boldsymbol{\eta}(dt, dt) & \boldsymbol{\eta}(dt,\ \mathrm{d}^A\alpha) \\ \boldsymbol{\eta}(E, dt) & \boldsymbol{\eta}(E,\ \mathrm{d}^A\alpha) \end{pmatrix} \\ &= \det\begin{pmatrix} -1 & 0 \\ 0 & \boldsymbol{\eta}(E,\ \mathrm{d}^A\alpha) \end{pmatrix} = -\boldsymbol{\eta}(E,\ \mathrm{d}^A\alpha) \end{aligned}$$

$$\mathrm{d}^{\mathrm{A}}(\alpha dt) = \mathrm{d}^A\alpha \wedge dt$$

$$\int_{\bar{M}} \boldsymbol{\eta}(\varpi_1 dt, \alpha dt) dt \wedge \mu_g = \int_{\bar{M}} \boldsymbol{\eta}\big(\delta^{\mathrm{A}}\, \mathrm{B_A}, \alpha dt\big) dt \wedge \mu_g = \int_{\bar{M}} \boldsymbol{\eta}\big(\, \mathrm{B_A},\, \mathrm{d}^{\mathrm{A}}(\alpha dt)\big) dt \wedge \mu_g$$
$$= \int_{\bar{M}} \boldsymbol{\eta}(E \wedge dt,\, \mathrm{d}^A \alpha \wedge dt) dt \wedge \mu_g = -\int_{\bar{M}} \boldsymbol{\eta}(E,\, \mathrm{d}^A \alpha) dt \wedge \mu_g$$
$$= -\int_{\bar{M}} \boldsymbol{\eta}(\delta^A E, \alpha) dt \wedge \mu_g = -\int_{\bar{M}} \boldsymbol{\eta}(\rho, \alpha) dt \wedge \mu_g$$
$$= \int_{\bar{M}} \boldsymbol{\eta}(\rho dt, \alpha dt) dt \wedge \mu_g$$

$\beta(t,\cdot) \in \Omega^1(M, \tilde{\mathfrak{g}})$ for each $t \in [t_0, t_1]\mathrm{cl}\{(t,x) \in \bar{M} \mid \beta(t,x) \neq 0\} \subset \mathrm{int}\bar{M} = \bar{M} - \partial\bar{M}$

$$\int_{\bar{M}} \boldsymbol{\eta}(\varpi_2, \beta) dt \wedge \mu_g = \int_{\bar{M}} \boldsymbol{\eta}\big(\delta^{\mathrm{A}}\, \mathrm{B_A}, \beta\big) dt \wedge \mu_g = \int_{\bar{M}} \boldsymbol{\eta}\big(\, \mathrm{B_A},\, \mathrm{d}^{\mathrm{A}}\beta\big) dt \wedge \mu_g$$
$$= \int_{\bar{M}} \boldsymbol{\eta}\big(B_A + E \wedge dt,\, \mathrm{d}^A \beta - \dot{\beta} \wedge dt\big) dt \wedge \mu_g$$
$$= \int_{\bar{M}} \big(\boldsymbol{\eta}(B_A,\, \mathrm{d}^A \beta) + \boldsymbol{\eta}(E, \dot{\beta})\big) dt \wedge \mu_g$$
$$= \int_{\bar{M}} \boldsymbol{\eta}(\delta^A B_A - \dot{E}, \beta) dt \wedge \mu_g = \int_{\bar{M}} \boldsymbol{\eta}(J, \beta) dt \wedge \mu_g$$

$$\delta^{\mathrm{A}}\, \mathrm{B_A} = \rho dt = J = \mathrm{J}$$

$$\mathrm{d}^{\mathrm{A}}\, \mathrm{B_A} = \mathrm{d}^{\mathrm{A}}(B_A + E \wedge dt) = \mathrm{d}^A B_A - \dot{B}_A \wedge dt - \mathrm{d}^A E \wedge dt$$
$$= \mathrm{d}^A E \wedge dt - \mathrm{d}^A E \wedge dt = 0$$

$$Q = \mathcal{C}(P) \times \Omega^0(M, \tilde{V})$$

$$\mathbf{g}_\kappa : \big(\textstyle\bigwedge^k T^*M \otimes \tilde{V}\big) \times_M \big(\bigwedge^m T^*M \otimes \tilde{V}\big) \to \mathbb{R}$$

$$\sharp_\kappa : \textstyle\bigwedge^k T^*M \otimes \tilde{V} \to \bigwedge^k TM \otimes \tilde{V}^*, \quad \alpha \otimes \sigma \mapsto \alpha^{\sharp_\partial} \otimes \sigma^{\flat_\kappa},$$
$$\flat_\kappa : \textstyle\bigwedge^k TM \otimes \tilde{V}^* \to \bigwedge^k T^*M \otimes \tilde{V}, \quad U \otimes \eta \mapsto U^{\flat} \otimes \eta^{\sharp_\kappa}.$$

$\alpha \in \bigwedge^k T^*M$ and $\sigma \in \tilde{V}$ as $\star\,(\alpha \otimes \sigma) = (\star\,\alpha) \otimes \sigma$, where $\star : \bigwedge^k T^*M \to \bigwedge^{m-k} T^*M$

$$\mathrm{d}^{\mathbf{A}} : \Omega^k(P, V) \to \Omega^{k+1}(P, V)$$

$$\mathrm{d}^{\mathbf{A}}\eta(u, v) = \mathrm{d}\eta(u, v) + \tilde{\varrho}(\mathbf{A}(u), \eta(v)) - \tilde{\varrho}(\mathbf{A}(v), \eta(u)), \eta \in \bar{\Omega}^1(P, V), u, v \in \mathfrak{X}(P)$$

$$\tilde{\varrho}(\sigma, v) = \frac{d}{dt}\Big|_{t=0} \varrho(\exp(\sigma t), v) = (d\varrho_v)_e(\sigma), \sigma \in \mathfrak{g}, v \in V$$

$$\mathrm{d}^A : \Omega^k(M, \tilde{V}) \to \Omega^{k+1}(M, \tilde{V})$$

$$\mathrm{d}^{A*} : \Omega^k\big(M, \tilde{V}^*\big) \to \Omega^{k+1}\big(M, \tilde{V}^*\big) \mathrm{div}^{A*} : \Lambda_m^{k+1}\big(M, \tilde{V}^*\big) \to \Lambda_m^k\big(M, \tilde{V}^*\big)$$

$$\tilde{\varrho}_\varphi : \Omega^0(M, \tilde{\mathfrak{g}}) \to \Omega^0(M, \tilde{V}), \xi \mapsto \tilde{\varrho}(\xi, \varphi),$$

$$\tilde{\varrho}_\varphi^* : \Omega^0\big(M, \tilde{V}^*\big) \to \Omega^0(M, \tilde{\mathfrak{g}}^*)$$

$\eta \cdot \tilde{\varrho}_\varphi(\xi) = \tilde{\varrho}_\varphi^*(\eta) \cdot \xi$ for each $\xi \in \Omega^0(M, \tilde{\mathfrak{g}})$ and $\eta \in \Omega^0\big(M, \tilde{V}^*\big)$

$$\tilde{\varrho}_{\varphi}: \Lambda_r^s(M, \tilde{\mathfrak{g}}) \to \Lambda_r^s(M, \tilde{V}), \tilde{\varrho}_{\varphi}^*: \Lambda_r^s\left(M, \tilde{V}^*\right) \to \Lambda_r^s(M, \tilde{\mathfrak{g}}^*)$$

$$\varphi \in \Omega^0(M, \tilde{V}), \delta A \in \Omega^1(M, \tilde{\mathfrak{g}}) \text{ and } \chi \in \Lambda_m^1\left(M, \tilde{V}^*\right)$$

$$\tilde{\varrho}_{\varphi}^*(\chi) \cdot \delta A = \delta A \wedge \tilde{\varrho}_{\varphi}^*(\Phi_{\tilde{V}}(\chi))$$

$$\delta A = \alpha_i dx^i \otimes \xi \text{ and } \chi = U^i \partial_i \otimes d^m x \otimes \eta$$

$$\alpha_i, U^i \in C^{\infty}(M), \xi \in \Omega^0(M, \tilde{\mathfrak{g}}) \text{ and } \eta \in \Omega^0\left(M, \tilde{V}^*\right)$$

$$\begin{aligned}\tilde{\varrho}_{\varphi}^*(\chi) \cdot \delta A &= U^i \alpha_i \left(\tilde{\varrho}_{\varphi}^*(\eta) \cdot \xi\right) d^m x = \alpha_i U^i \left(\xi \cdot \tilde{\varrho}_{\varphi}^*(\eta)\right)\left(dx^i \wedge d_i^{m-1} x\right) \\ &= \delta A \wedge \tilde{\varrho}_{\varphi}^*\left(U^i d_i^{m-1} x \otimes \eta\right) = \delta A \wedge \tilde{\varrho}_{\varphi}^*(\Phi_{\tilde{V}}(\chi))\end{aligned}$$

$$\mathcal{L}_{\text{int}} : \text{Conn}(P) \times_M (T^*M \otimes \tilde{\mathfrak{g}}) \times_M (\wedge^2 T^*M \otimes \tilde{\mathfrak{g}}) \times_M \tilde{V} \times_M \tilde{V} \times_M \left(T^*M \otimes \tilde{V}\right) \to \wedge^m T^*M$$

$$\mathcal{L}_{\text{int}}(A_x, \varepsilon_x, \beta_x, \varphi_x, \nu_x, \zeta_x) = \mathcal{L}_{\text{gau}}(A_x, \varepsilon_x, \beta_x) + \mathcal{L}_{\text{mat}}(\varphi_x, \nu_x, \zeta_x)$$

$$\begin{aligned}L_{\text{int}}(A, \varepsilon, \varphi, \nu) &= \int_M \mathcal{L}_{\text{int}}(A, \varepsilon, B_A, \varphi, \nu, \ \mathrm{d}^A \varphi) \\ &= \int_M \left(\mathcal{L}_{\text{gau}}(A, \varepsilon, B_A) + \mathcal{L}_{\text{mat}}(\varphi, \nu, \ \mathrm{d}^A \varphi)\right)\end{aligned}$$

$$(A, \varepsilon, \varphi, \nu) \in TQ \simeq \mathcal{C}(P) \times \Omega^1(M, \tilde{\mathfrak{g}}) \times T\Omega^0(M, \tilde{V})$$

$$\frac{\delta L_{\text{int}}}{\delta A} = \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial A} + \text{div}^{A*}\left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}\right) + \tilde{\varrho}_{\varphi}^*\left(\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \zeta}\right), \iota_{\partial M}^*\left(\text{tr}\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}\right)\right)$$

$$\frac{\delta L_{\text{int}}}{\delta \varepsilon} = \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \varepsilon}, 0\right)$$

$$\frac{\delta L_{\text{int}}}{\delta \varphi} = \left(\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \varphi} - \text{div}^{A*}\left(\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \zeta}\right), \iota_{\partial M}^*\left(\text{tr}\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \zeta}\right)\right)$$

$$\frac{\delta L_{\text{int}}}{\delta \nu} = \left(\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \nu}, 0\right)$$

$$\frac{\delta L_{\text{int}}}{\delta A} = \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial A} + \mathrm{d}^{A*}\left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}\right) + \tilde{\varrho}_{\varphi}^*\left(\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \zeta}\right), \iota_{\partial M}^*\left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta}\right)\right)$$

$$\frac{\delta L_{\text{int}}}{\delta \varepsilon} = \left(\frac{\partial \mathcal{L}_{\text{gau}}}{\partial \varepsilon}, 0\right)$$

$$\frac{\delta L_{\text{int}}}{\delta \varphi} = \left(\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \varphi} - \mathrm{d}^{A*}\left(\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \zeta}\right), \iota_{\partial M}^*\left(\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \zeta}\right)\right)$$

$$\frac{\delta L_{\text{int}}}{\delta \nu} = \left(\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \nu}, 0\right)$$

$$\left.\frac{d}{d\epsilon}\right|_{\epsilon=0} \mathrm{d}^{\mathbf{A}+\epsilon\delta\mathbf{A}}\boldsymbol{\varphi} = \left.\frac{d}{d\epsilon}\right|_{\epsilon=0} \mathrm{d}\boldsymbol{\varphi} + \tilde{\varrho}(\mathbf{A} + \epsilon\delta\mathbf{A}, \boldsymbol{\varphi}) = \tilde{\varrho}(\delta\mathbf{A}, \boldsymbol{\varphi})$$

$$d/d\epsilon|_{\epsilon=0}\ \mathrm{d}^{A+\epsilon\delta A}\varphi = \tilde{\varrho}_{\varphi}(\delta A)$$

$$\left.\frac{d}{d\epsilon}\right|_{\epsilon=0}\int_M \mathcal{L}_{\mathrm{mat}}\left(\varphi,\nu,\ \mathrm{d}^{A+\epsilon\delta A}\varphi\right) = \int_M \frac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\cdot\tilde{\varrho}_{\varphi}(\delta A) = \int_M \tilde{\varrho}_{\varphi}^{*}\left(\frac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)\cdot\delta A$$

$$(A,\varepsilon,\varsigma,\varsigma_{\partial},\varphi,\nu,\alpha,\alpha_{\partial})\colon[t_0,t_1]\to TQ\oplus T^{\star}Q$$

$$\begin{cases}\dot{A}=-E, & \dot{\varphi}=\nu,\\ \varsigma=\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\varepsilon}, & \dot{\varsigma}=\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial A}+\mathrm{div}^{A*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\beta}\right)+\tilde{\varrho}_{\varphi}^{*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\hat{\mathcal{F}}^{\star}\\ \varsigma_{\partial}=0, & \dot{\varsigma_{\partial}}=\iota_{\partial M}^{*}\left(\mathrm{tr}\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\beta}\right)+\hat{\mathcal{F}}_{\partial}^{\star}\\ \alpha=\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\nu}, & \dot{\alpha}=\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\varphi}-\mathrm{div}^{A*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\mathcal{F}^{\star}\\ \alpha_{\partial}=0, & \dot{\alpha}_{\partial}=\iota_{\partial M}^{*}\left(\mathrm{tr}\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\mathcal{F}_{\partial}^{\star}\end{cases}$$

$$(A,\varepsilon,\varsigma,\varsigma_{\partial},\varphi,\nu,\alpha,\alpha_{\partial})\colon[t_0,t_1]\to TQ\oplus T^{\dagger}Q$$

$$\begin{cases}\dot{A}=-E, & \dot{\varphi}=\nu\\ \varsigma=\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\varepsilon}, & \dot{\varsigma}=\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial A}+\mathrm{d}^{A*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\beta}\right)+\tilde{\varrho}_{\varphi}^{*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\hat{\mathcal{F}}^{\dagger}\\ \varsigma_{\partial}=0, & \dot{\varsigma}_{\partial}=\iota_{\partial M}^{*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\beta}\right)+\hat{\mathcal{F}}_{\partial}^{\dagger}\\ \alpha=\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\nu}, & \dot{\alpha}=\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\varphi}-\mathrm{d}^{A*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\mathcal{F}^{\dagger},\\ \alpha_{\partial}=0, & \dot{\alpha}_{\partial}=\iota_{\partial M}^{*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\mathcal{F}_{\partial}^{\dagger}.\end{cases}$$

$$\begin{cases}\dfrac{\partial}{\partial t}\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\varepsilon}=\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial A}+\mathrm{div}^{A_*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\beta}\right)+\tilde{\varrho}_{\varphi}^{*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\hat{\mathcal{F}}^{\star}\\ \dfrac{\partial}{\partial t}\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\nu}=\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\varphi}-\mathrm{div}^{A_*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\mathcal{F}^{\star}\\ \hat{\mathcal{F}}_{\partial}^{\star}=-\iota_{\partial M}^{*}\left(\mathrm{tr}\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\beta}\right),\mathcal{F}_{\partial}^{\star}=-\iota_{\partial M}^{*}\left(\mathrm{tr}\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)\end{cases}$$

$$\begin{cases}\dfrac{\partial}{\partial t}\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\varepsilon}=\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial A}+\mathrm{d}^{A_*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\beta}\right)+\tilde{\varrho}_{\varphi}^{*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\hat{\mathcal{F}}^{\dagger}\\ \dfrac{\partial}{\partial t}\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\nu}=\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\varphi}-\mathrm{d}^{A_*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\mathcal{F}^{\dagger}\\ \hat{\mathcal{F}}_{\partial}^{\dagger}=-\iota_{\partial M}^{*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\beta}\right),\mathcal{F}_{\partial}^{\dagger}=-\iota_{\partial M}^{*}\left(\dfrac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)\end{cases}$$

$$\frac{\partial}{\partial t}\mathcal{E}_{\mathrm{gau}}=-\mathrm{d}\left(\varepsilon\wedge\frac{\partial\mathcal{L}_{\mathrm{gau}}}{\partial\beta}\right)+\dot{A}\wedge\tilde{\varrho}_{\varphi}^{*}\left(\frac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)+\varepsilon\wedge\hat{\mathcal{F}}^{\dagger}$$
$$\frac{\partial}{\partial t}\mathcal{E}_{\mathrm{mat}}=-\mathrm{d}\left(\nu\wedge\frac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}\right)-\tilde{\varrho}_{\varphi}(\dot{A})\wedge\frac{\partial\mathcal{L}_{\mathrm{mat}}}{\partial\zeta}+\nu\wedge\mathcal{F}^{\dagger}$$

$S_{\text{tot}} = \frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta} \cdot \dot{A} + \frac{\partial \mathcal{L}_{\text{mat}}}{\partial \zeta} \cdot \dot{\varphi}$, or as the $(m-1)$-form $\mathrm{S}_{\text{tot}} = \dot{A} \wedge \frac{\partial \mathcal{L}_{\text{gau}}}{\partial \beta} + \dot{\varphi} \wedge \frac{\partial \mathcal{L}_{\text{mat}}}{\partial \zeta}$

$$\mathcal{L}_{\text{gau}}(A_x, \varepsilon_x, \beta_x) = \mathcal{L}_{\text{YM}}(A_x, \varepsilon_x, \beta_x) = \frac{1}{2}\big(\mathbf{g}(\varepsilon_x, \varepsilon_x) - \mathbf{g}(\beta_x, \beta_x)\big)\mu_g$$

$\mathcal{L}_{\text{mat}}(\varphi_x, \nu_x, \zeta_x) = \left(\frac{1}{2}\mathbf{g}_\kappa(\nu_x, \nu_x) - \frac{1}{2}\mathbf{g}_\kappa(\zeta_x, \zeta_x) - \mathbf{V}(\varphi_x)\right)\mu_g$ for some potential $\mathbf{V}\colon \tilde{V} \to \mathbb{R}$.

$$\mathbf{V}(\varphi_x) = m\mathbf{g}_\kappa(\varphi_x, \varphi_x)$$

Klein-Gordon field of mass $m > 0$, whilst $\mathbf{V}(\varphi_x) = \lambda \mathbf{g}_\kappa(\varphi_x, \varphi_x)^2 - \mu_H \mathbf{g}_\kappa(\varphi_x, \varphi_x)$ yields the Higgs

field of mass $m_H = \sqrt{2\mu_H} > 0$.

bundle morphism $\partial \mathbf{V}/\partial \varphi\colon \tilde{V} \to \tilde{V}^*$

$$\frac{\partial \mathbf{V}}{\partial \varphi}(\varphi_x) \cdot \delta\varphi_x = \frac{d}{d\epsilon}\Big|_{\epsilon=0} \mathbf{V}(\varphi_x + \epsilon\delta\varphi_x), \varphi_x, \delta\varphi_x \in \tilde{V}_x, x \in M$$

$$\operatorname{grad}_\kappa \mathbf{V} = \sharp_\kappa \circ \partial \mathbf{V}/\partial \varphi\colon \tilde{V} \to \tilde{V}$$

$$\mathcal{L}_{\text{mat}}(\varphi_x, \nu_x, \zeta_x) = \frac{1}{2}\nu_x^{b_\kappa} \wedge\!\star\, \nu_x - \frac{1}{2}\zeta_x^{b_\kappa} \wedge\!\star\, \zeta_x - \star\, \mathbf{V}(\varphi_x)$$

$(\varphi, \dot{\varphi}) \in T\Omega^0(M, \tilde{V})$, by $\zeta = \mathrm{d}^A\varphi \in \Omega^1(M, \tilde{V})$

$$\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \varphi} = -\frac{\partial \mathbf{V}}{\partial \varphi} \otimes \mu_g, \quad \frac{\partial \mathcal{L}_{\text{mat}}}{\partial \nu} = \dot{\varphi}^{\sharp_\kappa} \otimes \mu_g, \quad \frac{\partial \mathcal{L}_{\text{mat}}}{\partial \zeta} = -(\mathrm{d}^A\varphi)^{\sharp_\kappa}$$
$$\frac{\partial \mathcal{L}_{\text{mat}}}{\partial \varphi} = -\star\frac{\partial \mathbf{V}}{\partial \varphi}, \quad \frac{\partial \mathcal{L}_{\text{mat}}}{\partial \nu} = \star\, \dot{\varphi}^{\flat_\kappa}, \quad \frac{\partial \mathcal{L}_{\text{mat}}}{\partial \zeta} = -\star\,(\mathrm{d}^A\varphi)^{\flat_\kappa}$$

$$\begin{cases} \dot{A} = -E, & \dot{\varphi} = \nu, \\ \varsigma = -E^{\sharp_\partial} \otimes \mu_g, & \dot{\varsigma} = -\operatorname{div}^{A*}\left(B_A^{\sharp_\partial} \otimes \mu_g\right) - \tilde{\varrho}_\varphi^*\big((\mathrm{d}^A\varphi)^{\sharp_\kappa} \otimes \mu_g\big) + \hat{\mathcal{F}}^\star, \\ \varsigma_\partial = 0, & \dot{\varsigma_\partial} = -\iota_{\partial M}^*\left(\operatorname{tr}\left(B_A^{\sharp_\partial} \otimes \mu_g\right)\right) + \hat{\mathcal{F}}_\partial^\star, \\ \alpha = \dot{\varphi}^{\sharp_\partial} \otimes \mu_g, & \dot{\alpha} = -\frac{\partial \mathbf{V}}{\partial \varphi} \otimes \mu_g + \operatorname{div}^{A*}\big((\mathrm{d}^A\varphi)^{\sharp_\kappa} \otimes \mu_g\big) + \mathcal{F}^\star, \\ \alpha_\partial = 0, & \dot{\alpha}_\partial = -\iota_{\partial M}^*\big(\operatorname{tr}(\mathrm{d}^A\varphi)^{\sharp_\kappa} \otimes \mu_g\big) + \mathcal{F}_\partial^\star. \end{cases}$$

$$\begin{cases} \dot{A} = -E, & \dot{\varphi} = \nu \\ \varsigma = -\star\, E^{b_K}, & \dot{\varsigma} = -\mathrm{d}^{A*}\big(\star\, B_A^{b_K}\big) - \tilde{\varrho}_\varphi^*\big(\star\,(\mathrm{d}^A\varphi)^{b_\kappa}\big) + \hat{\mathcal{F}}^\dagger, \\ \varsigma_\partial = 0, & \dot{\varsigma_\partial} = -\iota_{\partial M}^*\big(\star\, B_A^{b_K}\big) + \hat{\mathcal{F}}_\partial^\dagger, \\ \alpha = \star\, \dot{\varphi}^{b_\kappa}, & \dot{\alpha} = -\star\frac{\partial \mathbf{V}}{\partial \varphi} + \mathrm{d}^{A*}\big(\star\,(\mathrm{d}^A\varphi)^{b_\kappa}\big) + \mathcal{F}^\dagger, \\ \alpha_\partial = 0, & \dot{\alpha}_\partial = -\iota_{\partial M}^*\big(\star\,(\mathrm{d}^A\varphi)^{b_\kappa}\big) + \mathcal{F}_\partial^\dagger. \end{cases}$$

$J\colon T\mathcal{C}(P) \to \Omega^1(M, \tilde{\mathfrak{g}}), j\colon T\mathcal{C}(P) \to \Omega^1(\partial M, \tilde{\mathfrak{g}}), \exists\colon T\Omega^0(M, \tilde{\mathfrak{g}}) \to \Omega^0(M, \tilde{\mathfrak{g}})$ and $\beth\colon T\Omega^0(M, \tilde{V}) \to$

$\Omega^0(\partial M, \tilde{V})$ the maps defined as $\star\, \hat{\mathcal{F}}^\dagger = -J^{b_K}, \star_\partial\, \hat{\mathcal{F}}_\partial^\dagger = j^{b_K}, \mathcal{F}^\dagger = \star\, \beth^{b_\kappa}$ and $\mathcal{F}_\partial^\dagger = \star_\partial\, \beth^{b_\kappa}$

$$\begin{cases}\dot{E}-\delta^{A}B_{A}+\tilde{\varrho}_{\varphi}^{*}((\mathrm{d}^{A}\varphi)^{b_{\kappa}})^{\sharp_{\kappa}}=-J,\\ \star_{\partial}\left(\iota_{\partial M}^{*}(\star\, B_{A})\right)=j,\end{cases}\quad\begin{cases}\ddot{\varphi}+\delta^{A}(\,\mathrm{d}^{A}\varphi)+\mathrm{grad}_{\kappa}\mathbf{V}=\beth,\\ \star_{\partial}\left(\iota_{\partial M}^{*}(\star\;\mathrm{d}^{A}\varphi)\right)=I.\end{cases}$$

$$\star\left(\mathrm{d}^{A*}\left(\star\,(\,\mathrm{d}^{A}\varphi)^{b_{\kappa}}\right)\right)=\star\left(\mathrm{d}^{A}\left(\star\,(\,\mathrm{d}^{A}\varphi)\right)\right)^{\mathfrak{G}_{\kappa}}=\left(\delta^{A}(\,\mathrm{d}^{A}\varphi)\right)^{\mathfrak{G}_{\kappa}}$$

$$b_{\kappa}\circ\mathrm{d}^{A}=\mathrm{d}^{A*}\circ b_{\kappa}$$

$$\ddot{\varphi}+\left(\delta^{A*}(\,\mathrm{d}^{A}\varphi)^{b_{\kappa}}\right)^{\sharp_{\kappa}}+\mathrm{grad}_{\kappa}\mathbf{V}=\beth$$

$$\iota_{\partial M}^{*}(i_{n}B_{A})=(-1)^{m}\frac{|g|}{|g_{\partial}|}j,\iota_{\partial M}^{*}(i_{n}\,\mathrm{d}^{A}\varphi)=\frac{|g|}{|g_{\partial}|}\beth$$

$$\mathcal{E}_{\mathrm{int}}=\mathcal{E}_{\mathrm{YM}}+\mathcal{E}_{\mathrm{mat}}=\left(\frac{1}{2}\mathbf{g}(E,E)+\frac{1}{2}\mathbf{g}(B_{A},B_{A})+\frac{1}{2}\mathbf{g}_{\kappa}(\dot{\varphi},\dot{\varphi})+\frac{1}{2}\mathbf{g}_{\kappa}(\mathrm{d}^{A}\varphi,\,\mathrm{d}^{A}\varphi)+\mathbf{V}(\varphi)\right)\mu_{g}$$
$$=\frac{1}{2}\left(E^{\mathfrak{G}_{K}}\wedge\star E+B_{A}^{\mathfrak{G}_{k}}\wedge\star B_{A}+\dot{\varphi}^{\mathfrak{G}_{\kappa}}\wedge\star\dot{\varphi}+(\mathrm{d}^{A}\varphi)^{\mathfrak{G}_{\kappa}}\wedge\star\,\mathrm{d}^{A}\varphi\right)+\mathbf{V}(\varphi)\mu_{g}$$

$$\frac{\partial\mathcal{E}_{\mathrm{tot}}}{\partial t}=-\mathrm{dS}_{\mathrm{gau}}+E\wedge\star J^{\mathfrak{G}_{k}}-\mathrm{dS}_{\mathrm{mat}}+\dot{\varphi}\wedge\star\beth^{\mathfrak{G}_{\kappa}}$$

$$\frac{d}{dt}\int_{M}\mathcal{E}_{\mathrm{tot}}=\underbrace{\int_{M}\left(E\wedge\star J^{\mathfrak{G}_{K}}+\dot{\varphi}\wedge\star\beth^{\mathfrak{G}_{\kappa}}\right)}_{\text{spatially distributed contribution}}+\underbrace{\int_{\partial M}\left(\left(\star_{\partial}\,j^{\mathfrak{G}_{K}}\right)\wedge\iota_{\partial M}^{*}E+(\iota_{\partial M}^{*}\dot{\varphi})\wedge\star_{\partial}\,\beth^{\mathfrak{G}_{\kappa}}\right)}_{\text{boundary contribution}}$$

$$\mathrm{S}_{\mathrm{gau}}=\mathrm{S}_{\mathrm{YM}}=E\wedge\star B_{A}^{\mathfrak{G}_{k}}\text{ and }\mathrm{S}_{\mathrm{mat}}=-\dot{\varphi}\wedge\star(\mathrm{d}^{A}\varphi)^{\mathfrak{G}_{\kappa}}$$

$$\dot{\rho}+\delta^{A}\left(J-\tilde{\varrho}_{\varphi}^{*}\left((\,\mathrm{d}^{A}\varphi)^{b_{\kappa}}\right)^{\sharp_{\kappa}}\right)=0.$$

$$S=\int\; d^{4}x\sqrt{-g}\left[-\frac{M_{P}^{2}}{2}\left(1+\frac{\xi_{\Phi}\Phi^{\dagger}\Phi}{M_{P}^{2}}+\frac{\xi_{H}H^{\dagger}H}{M_{P}^{2}}\right)R+\left(\tilde{D}_{\mu}\Phi\right)^{\dagger}\left(\tilde{D}^{\mu}\Phi\right)+\left(D_{\mu}H\right)^{\dagger}(D^{\mu}H)\right.$$
$$\left.-V(\Phi,H)-\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu}-\frac{\epsilon}{2}F'_{\mu\nu}B_{Y}^{\mu\nu}+\bar{\chi}i\not{D}\chi-m_{\chi}\bar{\chi}\chi\right]$$

$$g_{\mu\nu}\mathrm{d}x^{\mu}\mathrm{d}x^{\nu}=\mathrm{d}t^{2}-a^{2}(t)\mathrm{d}x^{i}\,\mathrm{d}x^{i}=a^{2}(\tau)\left[\mathrm{d}\tau^{2}-\mathrm{d}x^{i}\,\mathrm{d}x^{i}\right]$$

$$\tilde{D}_{\mu}\Phi=\partial_{\mu}\Phi-ig_{D}Z'_{\mu}\Phi\text{ and }D_{\mu}\chi=\partial_{\mu}\chi-ig_{D}Q_{\chi}Z'_{\mu}\chi$$

gauge coupling constant; $F'_{\mu\nu}=\partial_{\mu}Z'_{\nu}-\partial_{\nu}Z'_{\mu}$

$$V(\Phi,H)=\frac{\lambda_{H}}{2}\left(H^{\dagger}H-\frac{v_{H}^{2}}{2}\right)^{2}+\frac{\lambda_{\Phi}}{2}\left(\Phi^{\dagger}\Phi-\frac{v_{D}^{2}}{2}\right)^{2}+\lambda_{\Phi H}\left(H^{\dagger}H-\frac{v_{H}^{2}}{2}\right)\left(\Phi^{\dagger}\Phi-\frac{v_{D}^{2}}{2}\right).$$

$$\sin\,\theta=\frac{\lambda_{\Phi H}v_{H}v_{D}}{\sqrt{(\lambda_{\Phi}v_{D}^{2}-\lambda_{H}v_{H}^{2})^{2}+(\lambda_{\Phi H}v_{H}v_{D})^{2}}},$$

$$N(k_*) \equiv \int_{t_{end}}^{t_*} H[\eta(t)]\mathrm{d}t$$

$$\simeq \Delta N + \frac{1}{4}\log\frac{\rho_{\mathrm{eq}}}{\rho_{\mathrm{reh}}} - \log\frac{k_*}{a_0 H_0} + \log\frac{H_{k_*}}{H_{\mathrm{eq}}} + \log\left(219h\Omega_0\right),$$

$$\Delta N \simeq \frac{1}{4}\log\frac{\rho_{\mathrm{end}}}{\rho_{\mathrm{m.d.}}} + \frac{1}{3}\log\frac{\rho_{\mathrm{m.d}}}{\rho_{\mathrm{reh}}}$$

$$\sin\psi \simeq \sqrt{\frac{-\lambda_{\Phi H}}{\lambda_H}}$$

$$\sin\psi \sim \sqrt{\frac{\lambda_\Phi}{\lambda_H}} \text{ for } |\lambda_{\Phi H}| \ll \lambda_\Phi$$

$$\beta_{\lambda_\Phi} = \frac{1}{16\pi^2}\left[5\lambda_\Phi^2 + 2\lambda_{\Phi H}^2 - 6g_D^2\lambda_\Phi + 6g_D^4\right]$$

$$\xi_H \lesssim \xi_\Phi \lesssim 1, \lambda_{\Phi H}, g_D^2 \ll \lambda_\Phi \lesssim 4.2\times 10^{-10}, v_D < 10^{13}\mathrm{GeV}.$$

$$\begin{aligned}
\Phi(t,\mathbf{x}) &= \frac{1}{\sqrt{2}}\rho(t,\mathbf{x})e^{ig_D\theta(t,\mathbf{x})}, && \text{with} && \langle\rho(t,\mathbf{x})\rangle = \varrho(t) \neq 0;\\
G_\mu(t,\mathbf{x}) &= Z'_\mu(t,\mathbf{x}) + \partial_\mu\theta(t,\mathbf{x}), && \text{with} && \langle G_\mu\rangle = 0;\\
&= \left[G_0(t,\mathbf{x}), \partial_i G^{\parallel}(t,\mathbf{x}) + G_i^{\perp}(t,\mathbf{x})\right], && \text{with} && \partial_i G^{\perp i} = 0, G_i^L \equiv \partial_i G^{\parallel};
\end{aligned}$$


$$\mathbf{G}_\mathbf{k}^L = \epsilon_\mathbf{k}^L G_\mathbf{k}^L \; \mathbf{G}_\mathbf{k}^\perp = \sum_{\lambda=\pm}\epsilon_\mathbf{k}^{T\pm}G_\mathbf{k}^{T\lambda}$$

$$\begin{aligned}
&\epsilon_\mathbf{k}^L = \epsilon_{-\mathbf{k}}^{L*}, && \epsilon_\mathbf{k}^{L*}\cdot\epsilon_\mathbf{k}^L = 1, && i\mathbf{k}\cdot\epsilon_\mathbf{k}^L = k,\\
&\epsilon_\mathbf{k}^{T\pm} = \epsilon_{-\mathbf{k}}^{T\pm *}, && \epsilon_\mathbf{k}^{T\lambda *}\cdot\epsilon_\mathbf{k}^{T\lambda'} = \delta^{\lambda\lambda'}, && i\mathbf{k}\cdot\epsilon_\mathbf{k}^{T\pm} = 0,\\
& && && i\mathbf{k}\times\epsilon_\mathbf{k}^{T\pm} = \pm k\epsilon^{T\pm}
\end{aligned}$$

$$\Phi(t,\mathbf{x}) = \frac{\varphi(t) + \delta\phi_R(t,\mathbf{x}) + i\delta\phi_I(t,\mathbf{x})}{\sqrt{2}}$$

$$|\varphi(t)| = |\langle\mathrm{Re}\Phi\rangle| = \varrho(t)$$

$$\delta\phi_{R\mathbf{k}} = \delta\rho_\mathbf{k}, \delta\phi_{I\mathbf{k}} = -\frac{g_D\varrho}{k}G_\mathbf{k}^L, Z_\mathbf{k}^{T\pm} = G_\mathbf{k}^{T\pm} \text{ and } Z_\mathbf{k}^{LL} = 0.$$



$$\delta\tilde{\phi}_{I\mathbf{k}} = \frac{ak}{\sqrt{k^2 + g_D^2\tilde{\varphi}^2}}\delta\phi_{I\mathbf{k}} \text{ with } \tilde{\varphi} = a\varphi \text{ and } k = |\mathbf{k}|$$

$$S^{(2)}_{\phi_I} = \frac{1}{2}\int \frac{\mathrm{d}^{\square}\mathbf{k}\,\mathrm{d}\tau}{(2\pi)^3}\left[\left|\delta\tilde{\phi}'_{I\mathbf{k}}\right|^2 - \left(k^2 + m^2_{I,\mathrm{eff}}\right)\left|\delta\tilde{\phi}_{I\mathbf{k}}\right|^2\right],$$

$$m^2_{I,\mathrm{eff}} = m^2_{T,\mathrm{eff}} - \frac{k^2}{k^2 + m^2_{T,\mathrm{eff}}}\left[\frac{m''_{T,\mathrm{eff}}}{m_{T,\mathrm{eff}}} - \frac{3\left(m'_{T,\mathrm{eff}}\right)^2}{k^2 + m^2_{T,\mathrm{eff}}}\right] \text{ with } m_{T,\mathrm{eff}} = g_D\tilde{\varphi},$$

$$\tilde{\varphi}'' + \underbrace{\left[\lambda_\Phi(\tilde{\varphi}^2 - a^2v_D^2) + a^2\left(\xi_\Phi + \frac{1}{6}\right)R\right]}_{-m''_{T,\mathrm{eff}}/m_{T,\mathrm{eff}}}\tilde{\varphi} = 0 \text{ with } R = \frac{-6a''}{a^3}$$

$$X_{0\mathbf{k}} = \frac{g_D}{2}\frac{[\varphi'\delta\phi_{I\mathbf{k}} - \varphi\delta\phi'_{I\mathbf{k}}]}{(k/a)^2 + (g_D\varphi/2)^2}$$

$$m^2_{I,\mathrm{eff}} \simeq \lambda_\Phi(1+q)\tilde{\varphi}^2 + a^2\left(\xi_\Phi + \frac{1}{6}\right)R$$

$$\widehat{\delta\tilde{\phi}}_{I\mathbf{k}}(\tau) = \chi_k(\tau)\hat{a}_{\mathbf{k}} + \chi^*_k(\tau)\hat{a}^\dagger_{-\mathbf{k}} \text{ and require } \chi_k\chi^*_k - \chi'_k\chi^*_k = i$$

$$\left\langle\delta\tilde{\phi}_I^2\right\rangle = \lim_{\mathbf{x}'\to\mathbf{x}}\left\langle\widehat{\delta\tilde{\phi}}_I(\tau,\mathbf{x}')\widehat{\delta\tilde{\phi}}_I(\tau,\mathbf{x})\right\rangle = \int \frac{dk}{k}\frac{k^3}{2\pi^2}|\chi_k(\tau)|^2 = \int \frac{dk}{k}\Delta^2_{\delta\tilde{\phi}_I}(k,\tau)$$

$$\Delta^2_{\delta\phi_I}(k,\tau) = \frac{k^2 + g_D^2\tilde{\varphi}^2}{a^2k^2}\Delta^2_{\delta\tilde{\phi}_I}(k,\tau) \text{ and } \Delta^2_{G^L_{\mathbf{k}}} = \frac{a^2k^2}{g_D^2\tilde{\varphi}^2}\Delta^2_{\delta\phi_I}(k,\tau).$$

$$\delta\chi''_k + \underbrace{\left(k^2 + m^2_{I,\mathrm{eff}}\right)}_{\omega^2_{Ik}}\delta\chi_k = 0$$

$$\chi_k(\tau) \longrightarrow \frac{e^{-ik\tau}}{\sqrt{2k}}\left(k \gg m_{I,\,\mathrm{eff}}, \sqrt{\lambda_\Phi}\tilde{\varphi}\right) \star \tau \simeq \frac{-1}{aH}$$

$$\Delta^2_{G^L_{\mathbf{k}}}(k, t_{\rm end}) \simeq \left(\frac{kH_k}{2\pi m_{T,\rm eff}(t_{\rm end})}\right)^2 \quad \text{when} \quad k < a_{\rm end}H_{\rm end}$$

$$\langle \rho_G \rangle = \frac{1}{2a^4}\int \frac{\mathrm{d}^{\square}\mathbf{k}}{(2\pi)^3}\left[|\chi_k'|^2 + \left(k^2 + m^2_{I,\rm eff}\right)|\chi_k|^2\right]$$

$$\phi_k''(\tau) + \underbrace{\left[k^2 + 3\lambda_\Phi \tilde{\varphi}^2(\tau)\right]}_{\omega^2_{k\phi}}\phi_k(\tau) = 0 \quad \text{and} \quad \chi_k''(\tau) + \underbrace{\left[k^2 + \lambda_\Phi \tilde{\varphi}^2(\tau)\right]}_{\omega^2_{k\chi}}\chi_k(\tau) = 0$$

$$k \gg m_{T,\,\rm eff}, \sqrt{3q\tilde{\varphi}'} \propto \exp\left(\mu_k^I \tau\right)$$


$$\kappa \equiv \frac{k}{\sqrt{\lambda}\tilde{\varphi}_{\rm end}}$$

$$\lambda_\Phi = 2.5 \times 10^{-10} (\xi_\Phi = 0.5), \lambda_{H\Phi} = 10^{-11}, \lambda_H = 0.25, \varphi_{\rm end} = 1.33 M_P$$

$$k_{\rm IR} = 0.25\sqrt{\lambda_\Phi}\varphi_{\rm end}$$


$$d\tau = 10^{-3}\sqrt{\lambda_\Phi}\varphi_{\rm end}$$

$$\rho_{bg} = \frac{1}{2}\dot{\varphi}^2 + \frac{\lambda_\Phi}{4}(\varphi^2 - v_D^2)^2$$



$$\delta\rho_X \sim \left(\delta\dot{\phi}_X\right)^2$$

$$m_T < k < k_{\text{tail}} \sim O(10)\sqrt{\lambda_\Phi}\tilde{\varphi}_{\text{end}} :: n_k \propto k^{-3/2}$$



$$v_D = 3 \times 10^{16}\text{GeV}, k_{IR} = 0.08\sqrt{\lambda_\Phi}\varphi_{\text{end}}, N = 128, dt = 10^{-3}\sqrt{\lambda_\Phi}\varphi_{\text{end}}$$

$$n_k(\tau) = \left(\frac{\tau}{\tau_{\text{turb}}}\right)^{-q} n_{\tilde{k}}(\tau_{\text{turb}}), \text{ with } \tilde{k} = \left(\frac{\tau}{\tau_{\text{turb}}}\right)^{-p} k,$$

$$\left\langle a^2 \delta\phi_{R,I}^2 \right\rangle \propto \tau^{-2p}$$

$$\omega_k^{\text{peak}}(\tau) \sim (\tau/\tau_{\text{turb}})^p \omega_k^{\text{peak}}(\tau_{\text{turb}})$$

$$w = \frac{\rho_G}{3(\rho_G+\rho_V)}, \text{ assuming } \rho_K = \rho_G + \rho_V$$

$$\frac{a_d}{a_{p.t.}} \lesssim 90\left(\frac{2.5\times10^{-10}}{\lambda_\Phi}\right)^{2/3}\left(\frac{v_D}{10^9\text{GeV}}\right)^{2/3}$$

$$\frac{\mathrm{d}\rho_{\delta\phi}}{\mathrm{d}N} + 3(1+w_{\delta\phi})\rho_{\delta\phi} = -\frac{\left(\gamma_\phi\Gamma_{\delta\phi\to hh} + \gamma_\phi\Gamma_{\delta\phi\to Z'Z'}\right)}{H}(1+w_{\delta\phi})\rho_{\delta\phi}$$

$$\frac{\mathrm{d}\rho_R}{\mathrm{d}N} + 4\rho_R = \frac{\gamma_\phi\Gamma_{\delta\phi\to hh}}{H}(1+w_{\delta\phi})\rho_{\delta\phi} + \frac{\gamma_{Z'}\Gamma_{Z'\to\text{SMSM}}}{H}(1+w_{Z'})\rho_{Z'}$$

$$\frac{\mathrm{d}\rho_{Z'}}{\mathrm{d}N} + 3(1+w_{Z'})\rho_{Z'} = \frac{\gamma_\phi\Gamma_{\delta\phi\to Z'Z'}}{H}(1+w_{\delta\phi})\rho_{\delta\phi} - \left(\frac{\gamma_{Z'}\Gamma_{Z'\to\text{SMSM}}}{H} + \frac{\gamma_{Z'}\Gamma_{Z'\to\chi\bar{\chi}}}{H}\right)(1+w_{Z'})\rho_{Z'}$$

$$\frac{\mathrm{d}\rho_\chi}{\mathrm{d}N} + 3(1+w_\chi)\rho_\chi = \frac{\gamma_{Z'}\Gamma_{Z'\to\chi\bar{\chi}}}{H}(1+w_{Z'})\rho_{Z'}$$

$$H^2 = \frac{\rho_{\delta\phi} + \rho_{Z'} + \rho_R + \rho_\chi}{3M_P^2}$$

$$h^2\Omega_{\text{GW}}^{\text{inf,RD}}(f) = \frac{1}{\rho_c}\frac{d\ln\rho_{\text{GW}}}{d\ln k} \simeq 1\times10^{-16}\times[g_{*s}(T_{\text{hc}})]^{-4/3}g_{*\rho}(T_{\text{hc}})\left[\frac{H_*(f)}{3\times10^{13}\text{GeV}}\right]^2$$

$$\text{where } f \simeq 1\times10^{-5}\text{ Hz}\times[g_{*s}(T_{\text{hc}})]^{-2/3}\left[g_{*\rho}(T_{\text{hc}})\right]^{1/2}\left[\frac{T_{\text{hc}}}{1\text{TeV}}\right]$$

$$h^2\Omega_{\rm GW}^{\rm inf,pre-RD}(f) \simeq \left(\frac{k_{\rm rh}}{k}\right)^{\frac{2(1-3w)}{1+3w}} \times 10^{-16} \times [g_{*s}(T_{\rm rh})]^{-4/3} g_{*\rho}(T_{\rm rh}) \left[\frac{H_*(f)}{3\times 10^{13}{\rm GeV}}\right]^2.$$

$$h^2\Omega_{\rm GW}^{\rm preh}(f) \simeq 7.7\times 10^{-5} \times [g_{*s}(T_{\rm rh})]^{-4/3} g_{*\rho}(T_{\rm rh}) e^{-N_{\rm m.d}} \left\langle \frac{1}{\rho(\tau_{\rm turb})} \frac{{\rm d}\rho_{\rm GW}}{{\rm dlog}\ k}(\tau_{\rm turb}) \right\rangle \Big|_{k=\frac{f}{2\pi a_o}}$$

$$G\mu \simeq 2\pi G v_D^2 \log\left(\frac{m_r}{\Lambda}\right) \text{ where } \Lambda = \max\{g_D v_D, H\}$$

$$G\mu < 1.1\times 10^{-10}\left(\frac{v_D}{10^{13}{\rm GeV}}\right)^2 \left\{1+\frac{1}{26}\log\left[\left(\frac{\lambda_\Phi}{2.5\times 10^{-10}}\right)^{\frac{1}{2}}\left(\frac{v_D}{10^{13}{\rm GeV}}\right)\left(\frac{m_{Z',0}}{2m_e}\right)\right]\right\}$$

$$\Gamma_{\delta\phi\to hh} \simeq \frac{\lambda_{\Phi H}^2 v_D^2}{512\pi m_\phi}\sqrt{1-\frac{4m_h^2}{m_\phi^2}}$$

$$\Gamma_{\delta\phi\to Z'Z'} \simeq \frac{g_D^4 v_D^2}{192\pi}\frac{m_\phi^3}{m_{Z'}^4}\left[1-4\frac{m_{Z'}^2}{m_\phi^2}+12\frac{m_{Z'}^4}{m_\phi^4}\right]\sqrt{1-\frac{4m_{Z'}^2}{m_\phi^2}},$$

$$\Gamma_{Z'\to W^+W^-} = \frac{\epsilon^2 e^2 c_w^2}{16\pi} m_{Z'}\left[7-\frac{5m_{Z'}^2}{m_W^2}-\frac{12m_W^2}{m_{Z'}^2}+\frac{m_{Z'}^4}{m_W^4}\right]\sqrt{1-\frac{4m_W^2}{m_{Z'}^2}}$$

$$\Gamma_{Z'\to f\bar f} = \frac{\epsilon^2 e^2 c_w^2 Q_f^2 N_c}{4\pi} m_{Z'}\left(1+\frac{2m_f^2}{m_{Z'}^2}\right)\sqrt{1-\frac{4m_f^2}{m_{Z'}^2}}$$

$$\Gamma_{Z'\to\chi\bar\chi} = \frac{g_D^2}{4\pi} Q_\chi^2 m_{Z'}\left(1+\frac{2m_\chi^2}{m_{Z'}^2}\right)\sqrt{1-\frac{4m_\chi^2}{m_{Z'}^2}}$$

$$\langle\rho^2\rangle \simeq \max\left\{\langle\rho^2\rangle|_{\tau=\tau_s}\left(\frac{a_s}{a}\right)^{2(1+p)}, v_D^2\right\}$$

$$m_\phi^2 \simeq \frac{\lambda_\Phi}{2}(3\langle\rho^2\rangle - v_D^2)$$

$$m_{Z'}^2 \simeq g_D^2\langle\rho^2\rangle + \lambda_\Phi(\langle\rho^2\rangle - v_D^2)$$

$$m_h^2 \simeq \lambda_H v_H^2 + \frac{\lambda_{\Phi H}}{2}(\langle\rho^2\rangle - v_D^2)$$

$$\gamma_I = \left\langle\frac{m_I}{E_I}\right\rangle_{\Gamma_{\delta\phi\to Z'Z'}=H}$$

$$\langle E^{-1}\rangle \sim a_s/\left[a k_{\rm peak}(\tau)\right] \text{ where } k_{\rm peak}(\tau) = k_{\rm peak}(\tau_s)\left(\frac{a}{a_s}\right)^p$$

$$\gamma_\phi = \min\left\{1, \frac{m_\phi}{k_{\text{peak},\phi}(\tau_s)}\left(\frac{a_s}{a}\right)^{p+1}\right\}$$

$$\gamma_{Z'} = \begin{cases} \min\left\{1, \frac{m_{Z'}}{k_{\text{peak},\phi}(\tau_s)}\left(\frac{a_s}{a}\right)^{p+1}\right\}, & a \le a_d \\ \min\left\{1, \frac{g_D}{\sqrt{\lambda_\Phi}}\frac{a}{a_d}\right\} & a > a_d \end{cases}$$

$$w_{\delta\phi} = \begin{cases} \frac{1}{3} & , a \le a_{p.t.} \\ \frac{(\rho_G)_f}{3\left[(\rho_G)_f + \frac{a}{a_f}(\rho_V)_f\right]} & , a > a_{p.t} \end{cases}$$

$$w_{Z'} = \frac{1}{3}\theta(1-\gamma_{Z'})$$

$$\rho_{\delta\phi}(N_s) \simeq \left\langle \dot{\phi}_R^2(\tau_s)\right\rangle, \rho_{Z'}(N_s) \simeq \left\langle \delta\dot{\phi}_I^2(\tau_s)\right\rangle, \rho_R(N_s) \simeq \rho_h(N_S)$$

$$\left\langle \sigma_{f\bar{f}\to Z'Z'} v\right\rangle \simeq \begin{cases} 4\pi\epsilon^4\alpha_{\text{EM}}^2\cos^4\theta_W Q_f^4 \frac{\left(m_f^2 - m_{Z',0}^2\right)}{\left(2m_f^2 - m_{Z',0}^2\right)^2}\sqrt{1-\frac{m_{Z',0}^2}{m_f^2}} & \text{for } m_f > T \\ 4\pi\epsilon^4\alpha_{\text{EM}}^2\cos^4\theta_W Q_f^4 \frac{75}{s} & \text{for } m_f < T \text{ and } s \simeq 9T^2 \end{cases}$$

$$\langle \sigma_W + W^- \to Z'Z' v\rangle \simeq$$
$$\begin{cases} \frac{\pi\epsilon^4\alpha_{\text{EM}}^2\cos^4\theta_W}{9m_W^2}\frac{\left(m_W^2 - m_{Z',0}^2\right)^2}{\left(2m_W^2 - m_{Z',0}^2\right)^2}\sqrt{1-\frac{m_{Z',0}^2}{m_W^2}}\left(\frac{33m_W^4 + 58m_W^2 m_{Z',0}^2 + 3m_{Z',0}^4}{m_{Z',0}^4}\right) & \text{for } m_W > T \\ \frac{\pi\epsilon^4\alpha_{\text{EM}}^2\cos^4\theta_W}{4320m_{Z',0}^2}\frac{s^2\left(7s - 50m_{Z',0}^2 + 64m_W^2\right)}{m_W^4 m_{Z',0}^2} & \text{for } m_W < T \text{ and } s \simeq 9T^2 \end{cases}$$

$$\sigma_{f\bar{f}\to\chi\bar{\chi}}(s) = \frac{\epsilon^2\alpha_{\text{EM}}Q_f^2 Q_\chi^2 g_D^2}{3s}\sqrt{\frac{s-4m_\chi^2}{s-4m_f^2}}\frac{\left(s+2m_f^2\right)\left(s+2m_\chi^2\right)}{\left(s-m_{Z',0}^2\right)^2 + m_{Z',0}^2\Gamma_{Z',0}^2}$$

$$\sigma_{W^+W^-\to\chi\bar{\chi}}(s) = \frac{\epsilon^2\alpha_{\text{EM}}\cos^2\theta_W Q_\chi^2 g_D^2}{108m_{Z',0}^4}\sqrt{\frac{s-4m_\chi^2}{s-4m_W^2}}\left(1+\frac{2m_\chi^2}{s}\right)\frac{1}{\left(s-m_{Z',0}^2\right)^2 + m_{Z',0}^2\Gamma_{Z'}^2}$$
$$\times\left(4m_W^4\left(5s+12m_{Z',0}^2\right) - 16m_W^6 + s\left(s^2 + 16m_{Z',0}^2 s - 34m_{Z',0}^4\right)\right.$$
$$\left.-2m_W^2\left(4s^2 + 11m_{Z',0}^2 s + 40m_{Z',0}^4\right)\right)$$

$$Y_\chi^{\text{SM}} = \frac{2025}{\pi^4}\sqrt{\frac{2\pi^2}{45}}\frac{M_P}{g_{*,S}\sqrt{g_{*,\rho}}}\int_T^{T_R}\frac{dT}{T^6}\langle\sigma v\rangle_{\text{SMSM}\to\chi\bar{\chi}} n_{\text{SM}}^2(T)$$

$$\langle\sigma v\rangle_{\text{SMSM}\to\chi\bar{\chi}} = \frac{1}{8m_{\text{SM}}^4 T}\int_{4m_\chi^2}^{\infty} ds s^{1/2}\left(s-4m_{\text{SM}}^2\right)\sigma_{\text{SMSM}\to\chi\bar{\chi}}(s)\frac{K_1(\sqrt{s}/T)}{K_2^2(m_{\text{SM}}/T)}$$

$$\sigma_{Z'Z'\to Z'Z'}v=\frac{g_D^4m_{Z',0}^4}{4608\pi s^{3/2}}\sqrt{s-4m_{Z',0}^2}\left[4+\frac{\left(s-2m_{Z',0}^2\right)^2}{m_{Z',0}^4}+\frac{\left(s-2m_{Z',0}^2\right)^4}{16m_{Z',0}^8}\times\right]$$

$$\left[\frac{2}{m_\phi^2\left(s+m_\phi^2-4m_{Z',0}^2\right)}+\frac{1}{\left(s-m_\phi^2\right)^2+m_\phi^2\Gamma_\phi^2}\right.$$

$$\left.+\frac{4\left(3m_\phi^2-4m_{Z',0}^2\right)\left(s-m_\phi^2\right)-m_\phi^2\Gamma_\phi^2}{\left(s-4m_{Z',0}^2\right)\left(s+2m_\phi^2-4m_{Z',0}^2\right)\left[\left(s-m_\phi^2\right)^2+m_\phi^2\Gamma_\phi^2\right]}\log\left(\frac{m_\phi^2}{s+m_\phi^2-4m_{Z',0}^2}\right)\right]$$

$$\frac{\partial f_\chi}{\partial t}-Hp\frac{\partial f_\chi}{\partial p}=\frac{2\pi^2}{p^2}n_{Z'}\Gamma_{Z'\to\chi\bar{\chi}}\delta\left(p-\frac{m_{Z',0}}{2}\right)$$

$$f_\chi(t,p)=\frac{16\pi^2\Gamma_{Z'\to\chi\bar{\chi}}}{m_{Z',0}^3}\int_{t_{\mathrm{m.d}}}^{t_{\mathrm{reh}}}\mathrm{d}t'\frac{n_{Z'}}{H}\delta(t'-t_0),\ \text{where}\ a(t_0)=\frac{2p}{m_{Z',0}}a(t)$$

$$n_{Z'}(t_0)=n_{Z'}(t)\left(\frac{m_{Z',0}}{2p}\right)^3,H(t_0)=H(t>t_{\mathrm{m.d.}})\left(\frac{m_{Z',0}}{2p}\right)^{3/2},\rho_{Z'}\sim m_{Z',0}n_{Z'}$$

$$f_\chi(t_{\mathrm{m.d.}}<t\le t_{\mathrm{reh}},p)\simeq 16\pi^2\frac{\rho_{Z'}}{m_{Z',0}^4}\frac{\Gamma_{Z'\to\chi\bar{\chi}}}{H}\left(\frac{m_{Z',0}}{2p}\right)^{3/2}\theta\left(m_{Z',0}-2p\right)$$

$$f_\chi(t>t_{\mathrm{reh}},p)\simeq 16\pi^2\left[\frac{\rho_{Z'}}{m_{Z',0}^4}\frac{\Gamma_{Z'\to\chi\bar{\chi}}}{H}\right]_{t=t_{\mathrm{rh}}}\left(\frac{m_{Z',0}}{2p}\right)^{3/2}\left(\frac{a_{\mathrm{rh}}}{a}\right)^{3/2}\theta\left(\frac{a_{\mathrm{rh}}}{a}m_{Z',0}-2p\right)$$

$$\boldsymbol{\theta}=(\theta_e)_{e\in E(\Lambda)}\subset[-\pi,\pi]^{E(\Lambda)}$$

$$\boldsymbol{\theta}\mapsto\beta\sum_{p\in P(\Lambda)}\cos\left(\mathrm{d}\theta_p\right)+\alpha\sum_{e\in E(\Lambda)}\cos\left(\theta_e\right).$$

$$\mathrm{d}\theta_p=\theta_{e_1}+\theta_{e_2}-\theta_{e_3}-\theta_{e_4}$$

$$\mathcal{H}_\Lambda(\boldsymbol{\theta})=\sum_{p\in P(\Lambda)}\left[1-\cos\left(\mathrm{d}\theta_p\right)\right]+m\sum_{e\in E(\Lambda)}\left[1-\cos\left(\theta_e\right)\right].$$

$$\mathrm{d}\mu_\Lambda^{\text{Yang–Mills–Higgs (YMH)}}(\boldsymbol{\theta})=\frac{1}{Z_\Lambda^{\mathrm{YMH}}}\exp\left(-\beta\mathcal{H}_\Lambda(\boldsymbol{\theta})\right)\prod_{e\in E(\Lambda)}\mathbf{1}_{\{|\theta_e|\le\pi\}}\mathrm{d}\theta_e,$$

$$\left|\mathbb{E}_{\Lambda_n}^{\mathrm{YMH}}\left[\theta_x\theta_y\right]-\mathbb{E}_{\Lambda_n}^{\mathrm{YMH}}[\theta_x]\cdot\mathbb{E}_{\Lambda_n}^{\mathrm{YMH}}\left[\theta_y\right]\right|\le Ce^{-c\mathrm{dist}(x,y)}.$$

$$m\sum_{e\in E(\Lambda)}\left[1-\cos\left(\theta_e\right)\right]$$

$$\cos\left(\theta\right)\approx 1-\frac{\theta^2}{2}$$

$$\mathcal{H}_\Lambda(\boldsymbol{\theta})\approx\frac{1}{2}\sum_{p\in P(\Lambda)}\left(\mathrm{d}\theta_p\right)^2+\frac{m}{2}\sum_{e\in E(\Lambda)}\left(\theta_e\right)^2$$

$$L \geq L_0(m,d)\exp\left(-c_d m\mathrm{dist}(x,y)\right)$$

$$\lim_{\beta\to\infty}\frac{\psi(\beta)}{\beta}=0$$

$$\Lambda_n \subset \bigcup_{v\in\Lambda'} Q(v)$$

$$\tilde{Q}(v)=Q(v)\cap\Lambda_n,$$

$$\mathrm{F}(u,v)=\left\{(x,y)\in E\left(\mathbb{Z}^d\right)\colon x\in Q(u), y\in Q(v)\right\}$$

$$\mathcal{E}(\mathcal{S})=E\left(\bigcup_{v\in\mathcal{S}} Q(v)\right)$$

$$X(\Gamma)=\{v\in L\mathbb{Z}^d\colon \exists u\in L\mathbb{Z}^d\colon \mathrm{F}(u,v)\in\Gamma\}$$

$$\partial A=\left\{(u,v)\in E\left(\mathbb{Z}^d\right)\colon u\in V(A), v\in V\left(\mathbb{Z}^d\setminus A\right)\right\}$$

$$V_{\mathrm{s}}\left(t_x,t_y;\boldsymbol{\theta}\right)=V_{\mathrm{s}}(\boldsymbol{\theta})=t_x\theta_x+t_y\theta_y$$

$$\mathcal{H}_{\Lambda,\mathrm{s}}(\boldsymbol{\theta})=\mathcal{H}_{\Lambda}(\boldsymbol{\theta})+V_{\mathrm{s}}(\boldsymbol{\theta})$$

$$Z_{\Lambda}^{\mathrm{YMH,s}}=\int_{[-\pi,\pi]^{E(\Lambda)}}\exp\left(-\beta\mathcal{H}_{\Lambda,\mathrm{s}}(\boldsymbol{\theta})\right)\prod_{e\in E(\Lambda)}\mathrm{d}\theta_e$$

$$\frac{\partial}{\partial t_x}\frac{\partial}{\partial t_x}\Big|_{t_x=t_y=0}\log\, Z_{\Lambda}^{\mathrm{YMH,\,s}}=\mathbb{E}_{\Lambda}^{\mathrm{YMH}}\left[\theta_x\theta_y\right]-\mathbb{E}_{\Lambda}^{\mathrm{YMH}}\left[\theta_x\right]\cdot\mathbb{E}_{\Lambda}^{\mathrm{YMH}}\left[\theta_y\right]$$

$$Z_{\Lambda}^{\mathrm{YMH,\,s}}=Z_{\Lambda,f}^{\mathrm{G}}\cdot\Xi_{\Lambda}(\mathbf{0})\cdot\left(\sum_{n\geq 1}\sum_{\substack{\mathrm{P}_1,\ldots,\mathrm{P}_n\subset\Lambda'\\ \mathrm{P}_i\text{ connected}}}\prod_{j=1}^{n}\mathrm{w}\left(\mathrm{P}_j\right)\cdot\prod_{i<j}\delta\left(\mathrm{P}_i,\mathrm{P}_j\right)\right)$$

$$S_{\Lambda}(\boldsymbol{\theta})=\frac{1}{2}\sum_{p\in P(\Lambda)}\left(\mathrm{d}\theta_p\right)^2+\frac{m}{2}\sum_{e\in E(\Lambda)}(\theta_e)^2$$

$$\mathrm{d}\mu_{\Lambda,\eta}^{\mathrm{G}}(\boldsymbol{\theta})=\frac{1}{Z_{\Lambda,\eta}^{\mathrm{G}}}\exp\left(-\beta S_{\Lambda}(\boldsymbol{\theta})\right)\prod_{e\in E(\Lambda)\setminus\partial\Lambda}\mathrm{d}\theta_e\prod_{e\in\partial\Lambda}\delta_{\eta_e}(\theta_e)$$

$$g(t)=1-\cos(t)-\frac{t^2}{2}$$

$$V_{\Lambda}(\boldsymbol{\theta})=\sum_{p\in P(\Lambda)}g\left(\mathrm{d}\theta_p\right)+m\sum_{e\in E(\Lambda)}g(\theta_e)+V_{\mathrm{s}}(\boldsymbol{\theta}),$$

$$\mathrm{d}\mu_{\Lambda}^{\mathrm{YMH}}(\boldsymbol{\theta}) \propto e^{-\beta V_{\Lambda}(\boldsymbol{\theta})-\beta S_{\Lambda}(\boldsymbol{\theta})} \prod_{e\in E(\Lambda)} \mathbf{1}_{\{|\theta_e|\le\pi\}}\mathrm{d}\theta_e$$

$$\propto e^{-\beta V_{\Lambda}(\boldsymbol{\theta})} \cdot \left(\prod_{e\in E(\Lambda)} \mathbf{1}_{\{|\theta_e|\le\pi\}}\right)\mathrm{d}\mu_{\Lambda,f}^{\mathcal{G}}(\boldsymbol{\theta})$$

$$|\theta_e| \approx m^{-1/2}\beta^{-1/2}$$

$$g(\theta_e) \lesssim (\theta_e)^4 \lesssim m^{-2}\beta^{-2}$$

$$T_{\beta} = \frac{\log^{d+2}(\beta)}{\sqrt{\beta}}$$

$\chi(t) = \chi(-t)$ for all $t \in \mathbb{R}$;

$\chi(t) = 1$ if $|t| \le 1$ and $\chi(t) = 0$ if $|t| \ge 2$;

For all $k \ge 1$, we have $\sup_{t\in\mathbb{R}}\left|\chi^{(k)}(t)\right| \le C(k!)^2$ for some $C > 0$.

Setting $\zeta(t) = 1 - \chi(t)$:

$$1 = \prod_{e\in E(\Lambda)} \left(\chi\left(\frac{|\theta_e|}{T_{\beta}}\right) + \zeta\left(\frac{|\theta_e|}{T_{\beta}}\right)\right)$$

$$= \sum_{\mathcal{L}\subset E(\Lambda)} \left(\prod_{e\in E(\Lambda)\setminus\mathcal{L}} \chi\left(\frac{|\theta_e|}{T_{\beta}}\right) \cdot \prod_{e\in\mathcal{L}} \zeta\left(\frac{|\theta_e|}{T_{\beta}}\right)\right)$$

$$1 = \sum_{\mathcal{B}\subset\Lambda'} \chi_{\mathcal{G}} \cdot \zeta_{\mathcal{B}},$$

$$\chi_{\mathcal{G}} = \prod_{v\in\mathcal{G}} \left(\prod_{e\in E(Q(v))} \chi\left(\frac{|\theta_e|}{T_{\beta}}\right)\right)$$

$$\zeta_{\mathcal{B}} = \mathbf{1}_{\square} \cdot \sum_{\mathcal{L}\subset\mathcal{E}(\mathcal{B})} \left(\prod_{e\in\mathcal{L}} \zeta\left(\frac{|\theta_e|}{T_{\beta}}\right) \cdot \prod_{\substack{e\in E(\Lambda)\setminus\mathcal{L} \\ e\in\mathcal{E}(\mathcal{B})}} \chi\left(\frac{|\theta_e|}{T_{\beta}}\right)\right)$$

$$S_{\Lambda,\mathcal{G}}(\boldsymbol{\theta}) = \frac{1}{2} \sum_{\substack{p\in P(\Lambda) \\ E(p)\cap\mathcal{E}(\mathcal{G})\neq\emptyset}} \left(\mathrm{d}\theta_p\right)^2 + \frac{m}{2} \sum_{e\in\mathcal{E}(\mathcal{G})} (\theta_e)^2$$

$$S_{\Lambda}^{\mathcal{B}}(\boldsymbol{\theta}) = S_{\Lambda}(\boldsymbol{\theta}) - S_{\Lambda,\mathcal{G}}(\boldsymbol{\theta}), \text{ and } V_{\Lambda}^{\mathcal{B}}(\boldsymbol{\theta}) = V_{\Lambda}(\boldsymbol{\theta}) - V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta}),$$

$$Z_{\Lambda}^{\mathrm{YMH,\,s}} = \int_{[-\pi,\pi]^{E(\Lambda)}} \exp\left(-\beta \mathcal{H}_{\Lambda,\mathrm{s}}(\boldsymbol{\theta})\right) \prod_{e\in E(\Lambda)} \mathrm{d}\theta_e$$

$$= \sum_{\mathcal{B}\subset\Lambda'} \int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} e^{-\beta V_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta}) - \beta S_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta})} \cdot \zeta_{\mathcal{B}} \cdot \left(\int_{\mathbb{R}^{\mathcal{E}(\mathcal{G})}} e^{-\beta V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta}) - \beta S_{\Lambda,\mathcal{G}}(\boldsymbol{\theta})} \cdot \chi_{\mathcal{G}} \mathrm{d}\boldsymbol{\theta}_{\mathcal{G}} \right) \mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}$$

$$\mathrm{d}\boldsymbol{\theta}_{\mathcal{G}} = \prod_{e\in\mathcal{E}(\mathcal{G})} \mathrm{d}\theta_e$$

$$\int_{\mathbb{R}^{\mathcal{E}(\mathcal{G})}} e^{-\beta V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta}) - \beta S_{\Lambda,\mathcal{G}}(\boldsymbol{\theta})} \cdot \chi_{\mathcal{G}} \mathrm{d}\boldsymbol{\theta}_{\mathcal{G}} = Z_{\mathcal{G},\eta}^{\mathrm{G}} \cdot \mathbb{E}_{\mathcal{G},\eta}^{\mathrm{G}}\left[e^{-\beta V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta})} \chi_{\mathcal{G}}\right]$$

$$\Xi_{\mathcal{G},\eta} = \mathbb{E}_{\mathcal{G},\eta}^{\mathrm{G}}\left[e^{-\beta V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta})} \chi_{\mathcal{G}}\right]$$

$$Z_{\Lambda}^{\mathrm{YMH,\,s}} = \sum_{\mathcal{B}\subset\Lambda'} \int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} e^{-\beta V_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta}) - \beta S_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta})} \cdot \zeta_{\mathcal{B}} \cdot Z_{\mathcal{G},\eta}^{\mathrm{G}} \cdot \Xi_{\mathcal{G},\eta} \mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}$$

$$\mathcal{B}_1 = \left\{v \in \Lambda' : \mathrm{dist}(v,\mathcal{B}) \le r_\beta\right\}$$

$$\mathcal{G}_1 = \Lambda' \setminus \mathcal{B}_1,\ r_\beta^d = o\left(\beta T_\beta^2\right),\ \mathbf{s} = (s_{\mathrm{F}})_{\mathrm{F}\in P'(\mathcal{G}_1)}$$

$$S_{\Lambda,\mathcal{G}}(\boldsymbol{\theta};\mathbf{s}) = \frac{1}{2} \sum_{\substack{p\in P(\Lambda) \\ E(p)\cap\mathcal{E}(\mathcal{G})\neq\emptyset}} \sigma_p(\mathbf{s})\left(\mathrm{d}\theta_p\right)^2 + \frac{m}{2} \sum_{e\in\mathcal{E}(\mathcal{G})} (\theta_e)^2$$

$$\mathrm{d}\mu_{\mathcal{G},\eta,\mathbf{s}}^{\mathrm{G}}(\boldsymbol{\theta}) = \frac{1}{Z_{\mathcal{G},\eta,\mathbf{s}}^{\mathrm{G}}} \exp\left(-\beta S_{\Lambda,\mathcal{G}}(\boldsymbol{\theta};\mathbf{s})\right) \prod_{e\in\mathcal{E}(\mathcal{G})\setminus\partial\mathcal{E}(\mathcal{G})} \mathrm{d}\theta_e \prod_{e\in\partial\mathcal{E}(\mathcal{G})} \delta_{\eta_e}(\theta_e)$$

$$F(\mathbf{1}) = \sum_{\Gamma\subset S} \left(\int_{[0,1]^\Gamma} \partial^\Gamma F(\mathbf{s}_\Gamma) \mathrm{d}\mathbf{s}_\Gamma \right)$$

$$\partial^\Gamma = \prod_{\gamma\in\Gamma} \frac{\partial}{\partial s_\gamma},\ (\mathbf{s}_\Gamma)_\gamma = \begin{cases} s_\gamma & \gamma \in \Gamma \\ 0 & \text{otherwise} \end{cases}$$

$$F(\mathbf{1},s) \doteq \sum_{\Gamma\subset S'} \left(\int_{[0,1]^\Gamma} \partial^\Gamma F(\mathbf{s}_\Gamma, s) \mathrm{d}\mathbf{s}_\Gamma \right)$$

$$F(\mathbf{s}) = \log Z_{\mathcal{G},\eta}^{\mathrm{G}}(\mathbf{s})$$

$$\log Z_{\mathcal{G},\eta}^{\mathrm{G}} = \log Z_{\mathcal{G},\eta,\mathbf{1}}^{\mathrm{G}} = \sum_{\Gamma\subset P'(\mathcal{G}_1)} W_1(\Gamma;\mathcal{G}),$$

$$W_1(\Gamma;\mathcal{G}) = \int_{[0,1]^\Gamma} \partial^\Gamma \log Z_{\mathcal{G},\eta}^{\mathrm{G}}(\mathbf{s}_\Gamma) \mathrm{d}s_\Gamma$$

$$Z_{\mathcal{G},\eta}^{\mathrm{G}} = Z_{\Lambda,f}^{\mathrm{G}} \cdot \frac{Z_{\mathcal{G},\eta}^{\mathrm{G}}(\mathbf{0})}{Z_{\Lambda,f}^{\mathrm{G}}(\mathbf{0})} \cdot \exp\left(\sum_{\Gamma\neq\emptyset} W_2(\Gamma;\mathcal{G}) \right)$$

$$W_2(\Gamma;\mathcal{G}) = W_1(\Gamma;\mathcal{G}) - W_1(\Gamma;\Lambda')$$

$$Z_\Lambda^{\mathrm{YMH,\,s}} = Z_{\Lambda,f}^{\mathrm{G}} \cdot \sum_{\mathcal{B}\subset E(\Lambda)} \int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} e^{-\beta V_\Lambda^{\mathcal{B}(\boldsymbol{\eta})} - \beta S_\Lambda^{\mathcal{B}}(\boldsymbol{\eta})} \cdot \zeta_{\mathcal{B}} \cdot \frac{Z_{\mathcal{G},\eta}^{\mathrm{G}}(\mathbf{0})}{Z_{\Lambda,f}^{\mathrm{G}}(\mathbf{0})} \cdot \exp\left(\sum_{\Gamma\neq\emptyset} W_2(\Gamma;\mathcal{G})\right) \cdot \Xi_{\mathcal{G},\eta}\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}$$

$$\Xi_{\mathcal{G},\eta} = \mathbb{E}_{\mathcal{G},\eta}^{\mathrm{G}}\left[e^{-\beta V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta})}\chi_{\mathcal{G}}\right] = \int_{\mathbb{R}^{\mathcal{E}(\mathcal{G})}} e^{-\beta V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta})}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu_{\mathcal{G},\eta}^{\mathrm{G}}$$

$$V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta};\mathbf{s}) = \sum_{\substack{p\in P(\Lambda)\\ E(p)\cap\mathcal{E}(\mathcal{G})\neq\emptyset}} \sigma_p(\mathbf{s}) g\left(\mathrm{d}\theta_p\right) + m\sum_{e\in E(\mathcal{G})} g(\theta_e) + V_{\mathbf{s}}(\boldsymbol{\theta})$$

$$\Xi_{\mathcal{G},\eta}(\mathbf{s}) = \int_{\mathbb{R}^{\mathcal{E}(\mathcal{G})}} e^{-\beta V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta};\mathbf{s})}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu_{\mathcal{G},\eta,\mathbf{s}}^{\mathrm{G}}$$

$$\Xi_{\mathcal{G},\eta} = \Xi_{\mathcal{G},\eta}(\boldsymbol{\Gamma}) = \sum_{\Delta\subset P'(\mathcal{G}_1)} \int_{[0,1]^\Delta} \partial^\Delta \Xi_{\mathcal{G},\eta}(\mathbf{s}_\Delta)\mathrm{d}\mathbf{s}_\Delta$$

$$S_{\Lambda,\mathcal{G}}(\boldsymbol{\theta};\mathbf{0},+) = \frac{1}{2}\sum_{\substack{p\in P(\Lambda)\\ E(p)\cap\mathcal{E}(\mathcal{G})\neq\emptyset}} \tilde{\sigma}_p\left(\mathrm{d}\theta_p\right)^2 + \frac{m}{2}\sum_{e\in\mathcal{E}(\mathcal{G})} (\theta_e)^2$$

$$V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta};\mathbf{0},+) = \sum_{\substack{p\in P(\Lambda)\\ E(p)\cap\mathcal{E}(\mathcal{G})\neq\emptyset}} \tilde{\sigma}_p g\left(\mathrm{d}\theta_p\right) + m\sum_{e\in\mathcal{E}(\mathcal{G})} g(\theta_e) + V_{\mathbf{s}}(\boldsymbol{\theta})$$

$$\Xi_{\mathcal{G},\eta}(\mathbf{0},+) = \int_{\mathbb{R}^{\mathcal{G}}} e^{-\beta V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta};\mathbf{0},+)}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu_{\mathcal{G},\eta,\mathbf{0},+}^{G}$$

$$K_1(\Delta;\mathcal{G}) = \frac{1}{\Xi_{\mathcal{G},\eta}(\mathbf{0},+)}\int_{[0,1]^\Delta} \partial^\Delta \Xi_{\mathcal{G},\eta}(\mathbf{s}_\Delta)\mathrm{d}\mathbf{s}_\Delta$$

$$\Xi_{\mathcal{G},\eta} = \Xi_{\mathcal{G},\eta}(\mathbf{0},+)\sum_{\Delta\subset P'(\mathcal{G}_1)} K_1(\Delta;\mathcal{G})$$

$$Z_\Lambda^{\mathrm{YMH,\,s}} = Z_{\Lambda,f}^{\mathrm{G}} \cdot \Xi_\Lambda(\mathbf{0}) \cdot \sum_{\mathcal{B}\subset\Lambda'} \int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} \rho_{\mathcal{B}}(\boldsymbol{\eta}) \cdot \exp\left\{\sum_{\Gamma\neq\emptyset} W_2(\Gamma;\mathcal{G})\right\} \cdot \left(\sum_{\Delta} K_1(\Delta;\mathcal{G})\right)\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}} \quad (41)$$

$$\rho_{\mathcal{B}}(\boldsymbol{\eta}) = e^{-\beta V_\Lambda^{\mathcal{B}}(\boldsymbol{\eta}) - \beta S_\Lambda^{\mathcal{B}}(\boldsymbol{\eta})} \cdot \zeta_{\mathcal{B}}(\boldsymbol{\eta}) \cdot \frac{Z_{\mathcal{G},\eta}^{\mathrm{G}}(\mathbf{0})}{Z_{\Lambda,f}^{\mathrm{G}}(\mathbf{0})} \cdot \frac{\Xi_{\mathcal{G},\eta}(\mathbf{0},+)}{\Xi_\Lambda(\mathbf{0})}.$$

$$\exp\left(\sum_{\Gamma\neq\emptyset} W_2(\Gamma;\mathcal{G})\right)=\sum_{n\geq 0}\frac{1}{n!}\left(\sum_{\Gamma_1\neq\emptyset}\cdots\sum_{\Gamma_n\neq\emptyset}\prod_{j=1}^{n} W_2(\Gamma_j;\mathcal{G})\right)$$
$$=\sum_{\mathbf{\Gamma}\in\mathbf{T}}\frac{1}{|\mathbf{\Gamma}|!}\prod_{\Gamma\in\mathbf{\Gamma}} W_2(\Gamma;\mathcal{G})$$

$$Z_{\Lambda}^{\mathrm{YMH,\,s}}=Z_{\Lambda,f}^{\mathrm{G}}\cdot\Xi_{\Lambda}(\mathbf{0})\sum_{\mathcal{B}\subset\Lambda'}\sum_{\mathbf{\Gamma}\in\mathbf{T}}\sum_{\Delta\subset P(\mathcal{G}_1)}\frac{1}{|\mathbf{\Gamma}|!}\int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}}\rho_{\mathcal{B}}(\boldsymbol{\eta})$$
$$\cdot\prod_{\Gamma\in\mathbf{\Gamma}} W_2(\Gamma;\mathcal{G})\cdot K_1(\Delta;\mathcal{G})\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}\quad(44)$$

$$\mathrm{P}=\mathcal{B}\cup\left(\bigcup_{\Gamma\in\mathbf{\Gamma}} X(\Gamma)\right)\cup X(\Delta)$$

$$\mathrm{w}(\mathrm{P})=\sum_{(\mathcal{B},\mathbf{\Gamma},\Delta)\text{ is a polymer on P}}\frac{1}{|\mathbf{\Gamma}|!}\int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}}\rho_{\mathcal{B}}(\boldsymbol{\eta})\cdot\prod_{\Gamma\in\mathbf{\Gamma}} W_2(\Gamma;\mathcal{G})\cdot K_1(\Delta;\mathcal{G})\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}$$

$$K_1(\Delta;\mathcal{G})=\prod_{\delta\text{ is a maximal }\mathcal{G}_1\text{-connected component}} K_1(\delta;\mathcal{G})$$

$$Z_{\Lambda}^{\mathrm{YMH,\,s}}=Z_{\Lambda,f}^{\mathrm{G}}\cdot\Xi_{\Lambda}(\mathbf{0})\cdot\sum_{\mathbf{P}}\prod_{\mathrm{P}\in\mathbf{P}}\mathrm{w}(\mathrm{P}).$$

$$\Psi(\mathbf{X})=\left(\prod_{\substack{\mathrm{P}\subset\Lambda'\\ \mathrm{P}\text{ connected}}}\frac{1}{n_{\mathbf{X}}(\mathrm{P})!}\right)\cdot\left(\prod_{\mathrm{P}\in\mathbf{X}}\mathrm{w}(\mathrm{P})\right)$$
$$\cdot\left(\sum_{G\in\{\text{ connected graphs on }\mathbf{X}\}}\prod_{(\mathrm{P}_i,\mathrm{P}_j)\in E(G)}\left(\delta(\mathrm{P}_i,\mathrm{P}_j)-1\right)\right)$$

$$\bar{\Psi}(X)=\sum_{\substack{\mathbf{X}=(\mathrm{P}_1,\ldots,\mathrm{P}_n)\\ \cup_i\mathrm{P}_i=X}}\Psi(\mathbf{X})$$

$$\log Z_{\Lambda}^{\mathrm{YMH,\,s}}=\log\left(Z_{\Lambda,f}^{\mathrm{G}}\cdot\Xi_{\Lambda}(\mathbf{0})\right)+\sum_{\mathbf{X}}\Psi(\mathbf{X})$$
$$=\log\left(Z_{\Lambda,f}^{\mathrm{G}}\cdot\Xi_{\Lambda}(\mathbf{0})\right)+\sum_{X\subset\Lambda'}\bar{\Psi}(X)$$

$$\|f\|_r=\sup_{x\in L\mathbb{Z}^d}\sum_{\substack{\mathrm{P}\text{ connected}\\ \mathrm{P}\,\mathrm{Pouch}\,x}} f(\mathrm{P})e^{r|\mathrm{P}|}$$

$$\|f\|_r=\sup_{x\in L\mathbb{Z}^d}\sum_{\substack{\Gamma\text{connected}\\ \Gamma\text{ touch }x}} f(\Gamma)e^{r|X(\Gamma)|}$$

$$\sum_{\substack{\widetilde{\mathrm{P}}\subset\Lambda' \text{ connected}\\ \mathrm{P},\widetilde{\mathrm{P}} \text{ touch}}} |\mathrm{w}(\widetilde{\mathrm{P}})|e^{|\widetilde{\mathrm{P}}|} \le |\mathrm{P}|$$

$$\sum_{\substack{\widetilde{\mathrm{P}}\subset\Lambda' \text{ connected}\\ \mathrm{P},\widetilde{\mathrm{P}} \text{ touch}}} \mathrm{w}(\widetilde{\mathrm{P}})e^{|\widetilde{\mathrm{P}}|} \le |\mathrm{P}| \cdot \sup_{x\in\mathrm{P}^2} \sum_{\mathrm{P_P} \text{ connected}} \mathrm{w}(\mathrm{P})e^{|\mathrm{P}|} \le |\mathrm{P}| \cdot \|\mathrm{w}\|_1$$

$$\|\bar{\Psi}\|_r \le \sum_{n\ge 1} (2\|\mathrm{w}\|_{r+2})^n$$

$$\log Z_\Lambda^{\mathrm{YMH,\,s}} = \log\left(Z_{\Lambda,f}^{\mathrm{G}} \cdot \Xi_\Lambda(\mathbf{0})\right) + \sum_{X\subset\Lambda'} \bar{\Psi}(X)$$

$$(t_x, t_y) \mapsto \log Z_\Lambda^{\mathrm{YMH,\,s}}$$

$$\left|\mathbb{E}_\Lambda^{\mathrm{YMH}}[\theta_x\theta_y] - \mathbb{E}_\Lambda^{\mathrm{YMH}}[\theta_x]\cdot\mathbb{E}_\Lambda^{\mathrm{YMH}}[\theta_y]\right|$$

$$= \left|\frac{\partial}{\partial t_x}\frac{\partial}{\partial t_x}\right|_{t_x=t_y=0} \log Z_\Lambda^{\mathrm{YMH,\,s}}\Bigg|$$

$$= \left|\sum_{\substack{X\subset\Lambda'\\ \{x,y\}\subset\mathcal{E}(X)}} \frac{\partial}{\partial t_x}\frac{\partial}{\partial t_x}\right|_{t_x=t_y=0} \bar{\Psi}(X)\Bigg| \le C_\beta \sum_{\substack{X\subset\Lambda'\\ \{x,y\}\subset\mathcal{E}(X)}} \sup_{|t_x|,|t_y|\le\beta^{-10}} \bar{\Psi}(X).$$

$$|X| \ge cL^{-1}\cdot \mathrm{dist}(x,y)$$

$$\left|\mathbb{E}_\Lambda^{\mathrm{YMH}}[\theta_x\theta_y] - \mathbb{E}_\Lambda^{\mathrm{YMH}}[\theta_x]\cdot\mathbb{E}_\Lambda^{\mathrm{YMH}}[\theta_y]\right|$$
$$\le C_\beta \sum_{\substack{X\subset\Lambda'\\ \{x,y\}\subset\mathcal{E}(X)}} \bar{\Psi}(X)e^{|X|}e^{-|X|} \le C_\beta e^{-cL^{-1}\mathrm{dist}(x,y)} \sum_{X\subset\Lambda'} \bar{\Psi}(X)e^{|X|}$$
$$\le C_\beta e^{-cL^{-1}\mathrm{dist}(x,y)}\|\bar{\Psi}\|_1 \dot{\le} C_\beta e^{-cL^{-1}\mathrm{dist}(x,y)} \sum_{n\ge 1} (2\|\mathrm{w}\|_3)^n$$

$$\left|\mathbb{E}_\Lambda^{\mathrm{YMH}}[\theta_x\theta_y] - \mathbb{E}_\Lambda^{\mathrm{YMH}}[\theta_x]\cdot\mathbb{E}_\Lambda^{\mathrm{YMH}}[\theta_y]\right| \le C_\beta e^{-c_L\mathrm{dist}(x,y)},$$

$$\tilde{\rho}(X) = \sum_{\substack{\mathcal{B}\subset\Lambda'\\ \mathcal{B}_1=X}} \left(\int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} \rho_{\mathcal{B}}(\boldsymbol{\eta})\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}\right)$$

$$\lim_{\beta\to\infty} \|\tilde{\rho}\|_r = 0$$

$$\|W_2(\cdot,\mathcal{G})\|_r \le C_L$$

$\mathcal{C}(\mathcal{B}_1) = \{\Delta \subset P'(\mathcal{G}_1)$: all connected components of Δ touch $\mathcal{B}_1\}$.

$$\lim_{\beta\to\infty}\left(e^{-C_L|\mathcal{B}_1|}\sum_{\Delta\in\mathcal{C}(\mathcal{B}_1)} |K_1(\Delta;\mathcal{G})|e^{r|X(\Delta)|}\right) = 0$$

$$\lim_{\beta\to\infty}\|K_1(\cdot,\Lambda')\|_r = 0$$

$$K_1(\emptyset,\mathcal{G}) \le e^{C_L|\mathcal{B}_1|}$$

$$\#\{X\subset\mathbb{Z}^d : 0\in X, |X| = k, \text{and } X \text{ is connected}\} \le \exp(Ck)$$

$$F(\mathbf{X}) = \prod_{j=1}^{n} f(\mathrm{P}_j)$$

$$\sum_{n\ge1}\frac{1}{n!}\left(\sum_{\substack{\mathbf{X}=(\mathrm{P}_1,\dots,\mathrm{P}_n)\\ \mathrm{P}_j \text{ and } \widetilde{\mathrm{P}} \text{ touch}}} F(\mathbf{X})\right) \le \frac{1}{A}\exp\left(A|\widetilde{\mathrm{P}}|\cdot\|f\|_0\right)$$

$$A\sum_{n\ge1}\frac{1}{n!}\left(\sum_{\substack{\mathbf{X}=(\mathrm{P}_1,\dots,\mathrm{P}_n)\\ \mathrm{P}_j \text{ and } \widetilde{\mathrm{P}} \text{ touch}}} F(\mathbf{X})\right)$$
$$= A\sum_{n\ge1}\frac{1}{n!}\left(\sum_{\mathrm{P} \text{ and } \widetilde{\mathrm{P}} \text{ touch}} f(\mathrm{P})\right)^n$$
$$\le \sum_{n\ge1}\frac{1}{n!}\left(A\sum_{\mathrm{P} \text{ and } \widetilde{\mathrm{P}} \text{ touch}} f(\mathrm{P})\right)^n \le \exp\left(A\sum_{\mathrm{P} \text{ and } \widetilde{\mathrm{P}} \text{ touch}} f(\mathrm{P})\right) \le \exp\left(A|\widetilde{\mathrm{P}}|\cdot\|f\|_0\right)$$

$$\mathrm{w}(\mathrm{P}) = \sum_{(\mathcal{B},\boldsymbol{\Gamma},\Delta) \text{ is a polymer on } \mathrm{P}} \frac{1}{|\boldsymbol{\Gamma}|!}\int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} \rho_{\mathcal{B}}(\boldsymbol{\eta})\cdot\prod_{\Gamma\in\boldsymbol{\Gamma}} W_2(\Gamma;\mathcal{G})\cdot K_1(\Delta;\mathcal{G})\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}$$

$$|\mathrm{P}| \le |\mathcal{B}| + \sum_{\Gamma\in\Gamma} |X(\Gamma)| + |X(\Delta)|$$

$$|\mathrm{w}(\mathrm{P})|e^{R|\mathrm{P}|} \le \sum_{(\mathcal{B},\boldsymbol{\Gamma},\Delta) \text{ is a polymer on } \mathrm{P}} \frac{1}{|\boldsymbol{\Gamma}|!}\int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} \rho_{\mathcal{B}}(\boldsymbol{\eta})e^{R|\mathcal{B}|}$$
$$\times\left(\prod_{\Gamma\in\boldsymbol{\Gamma}} W_2(\Gamma;\mathcal{G})e^{R|X(\Gamma)|}\right)\cdot K_1(\Delta;\mathcal{G})e^{R|X(\Delta)|}\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}$$

$$\mathcal{N}_1 = \sum_{\substack{(\mathcal{B},\Gamma,\Delta) \text{ is a polymer on } \mathrm{P}\\ \mathcal{B}\neq\emptyset}} (\cdots), \text{ and } \mathcal{N}_2 = \sum_{(\emptyset,\Gamma,\Delta) \text{ is a polymer on } \mathrm{P}} (\cdots),$$

$$\sum_{\boldsymbol{\Gamma}}\frac{1}{|\boldsymbol{\Gamma}|!}\left(\prod_{\Gamma\in\boldsymbol{\Gamma}} W_2(\Gamma;\mathcal{G})e^{R|X(\Gamma)|}\right) \dot{\le} \exp\left(\|W_2(\cdot;\mathcal{G})\|_R\cdot|\mathcal{B}_1|\right) \dot{\le} \exp\left(C_L|\mathcal{B}_1|\right)$$

$$\mathcal{N}_1 \le \sum_{\substack{(\mathcal{B},\Delta)\\ \mathcal{B}\neq\emptyset,\mathcal{B}\cup X(\Delta)\subset\mathrm{P}}} e^{C_L|\mathcal{B}_1|}\int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} \rho_{\mathcal{B}}(\boldsymbol{\eta})\cdot K_1(\Delta;\mathcal{G})e^{R|X(\Delta)|}\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}$$

$$\exp\left(\|K_1(\cdot,\Lambda')\|_R \cdot |\mathrm{P}|\right) \dot{\leq} \exp\left(|\mathrm{P}|\right)$$

$$\sum_{\Delta\in\mathcal{C}(\mathcal{B}_1)} |K_1(\Delta;\mathcal{G})| e^{R|X(\Delta)|} \leq e^{C_L|\mathcal{B}_1|}$$

$$|K_1(\emptyset;\mathcal{G})| \leq e^{C_L|\mathcal{B}_1|}$$

$$\mathcal{N}_1 \leq e^{|\mathrm{P}|} \sum_{\substack{\mathcal{B}\neq\emptyset \\ \mathcal{B}\subset\mathrm{P}}} e^{C_L|\mathcal{B}_1|} \int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} \rho_{\mathcal{B}}(\boldsymbol{\eta})\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}$$

$$\sum_{\substack{\mathcal{B}\neq\emptyset \\ \mathcal{B}\subset\mathrm{P}}} e^{C_L|\mathcal{B}_1|} \int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} \rho_{\mathcal{B}}(\boldsymbol{\eta})\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}} = \sum_{\emptyset\neq X\subset\mathrm{P}} \prod_{x\subset X} \left(\tilde{\rho}(x) e^{C_L|x|}\right)$$

$$A = \|\tilde{\rho}\|_{C_L}^{-1} \text{ and } \widetilde{\mathrm{P}} = \mathrm{P}$$

$$\sum_{\substack{\mathcal{B}\neq\emptyset \\ \mathcal{B}\subset\mathbf{P}}} e^{C_L|\mathcal{B}_1|} \int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} \rho_{\mathcal{B}}(\boldsymbol{\eta})\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}} \leq \|\tilde{\rho}\|_{C_L} \cdot e^{|\mathbf{P}|}$$

$$\mathcal{N}_1 \leq e^{2|\mathbf{P}|} \|\tilde{\rho}\|_{C_L}$$

$$\mathcal{N}_2 = \sum_{\substack{\Delta \text{ such that} \\ X(\Delta)=\mathrm{P}}} |K_1(\Delta;\Lambda')| e^{R|X(\Delta)|} = e^{R|\mathrm{P}|} \sum_{\substack{\Delta \text{ such that} \\ X(\Delta)=\mathrm{P}}} |K_1(\Delta;\Lambda')|.$$

$$|\mathrm{w}(\mathrm{P})| e^{R|\mathrm{P}|} \leq \mathcal{N}_1 + \mathcal{N}_2 \leq e^{R|\mathrm{P}|} \sum_{\substack{\Delta \text{ such that} \\ X(\Delta)=\mathrm{P}}} |K_1(\Delta;\Lambda')| + e^{2|\mathrm{P}|} \cdot \|\tilde{\rho}\|_{C_L}$$

$$|\mathrm{w}(\mathrm{P})| \leq \sum_{\substack{\Delta \text{ such that} \\ X(\Delta)=\mathrm{P}}} |K_1(\Delta;\Lambda')| + e^{(2-R)|\mathrm{P}|} \cdot \|\tilde{\rho}\|_{C_L}$$

$$\sum_{\substack{\mathrm{P}_{\text{connected}} \\ \mathrm{P}_{\text{touch } x}}} e^{(2-R)|\mathrm{P}|} e^{r|\mathrm{P}|} = \sum_{\substack{\mathrm{P}_{\text{connected}} \\ \mathrm{P}_{\text{touch } x}}} e^{(2-C_d)|\mathrm{P}|} < \infty,$$

$$\sum_{\substack{\mathrm{P}_{\text{connected}} \\ \mathrm{P}_{\text{touch } x}}} e^{r|\mathrm{P}|} \left(\sum_{\substack{\Delta \text{ such that} \\ X(\Delta)=\mathrm{P}}} |K_1(\Delta;\Lambda')| \right) = \sum_{\substack{\Delta_{\text{connected}} \\ \Delta_{\text{touch } x}}} e^{r|X(\Delta)|} |K_1(\Delta;\Lambda')| \leq \|K_1(\cdot,\Lambda')\|_r.$$

$$\|\mathrm{w}\|_r \leq \|K_1(\bullet,\Lambda')\|_r + C\|\tilde{\rho}\|_{C_L},$$

$$\mathcal{H}_{\Lambda,\mathrm{s}}(\boldsymbol{\theta}) = \mathcal{H}_{\Lambda}(\boldsymbol{\theta}) + V_{\mathrm{s}}(\boldsymbol{\theta})$$

$$\mathcal{H}_{\Lambda}(\boldsymbol{\theta}) = \sum_{p\in P(\Lambda)} \left[1 - \cos\left(\mathrm{d}\theta_p\right)\right] + m \sum_{e\in E(\Lambda)} \left[1 - \cos\left(\theta_e\right)\right]$$

$$\mathcal{H}_{\Lambda}(\boldsymbol{\theta}) \geq \frac{1}{10} \sum_{p \in P(\Lambda)} \left(\mathrm{d}\theta_p\right)^2 + \frac{m}{10} \sum_{e \in E(\Lambda)} (\theta_e) \dot{=} \frac{(14)}{5} \cdot S_{\Lambda}(\boldsymbol{\theta})$$

$$V_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta}) + S_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta}) \geq \frac{1}{5} S_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta}) + V_{\mathrm{s}}(\boldsymbol{\eta})$$

$$\boldsymbol{\eta} \in [-\pi, \pi]^{\mathcal{E}(\mathcal{B})} - \cos(\theta) \geq \frac{\theta^2}{10}$$

$$\rho_{\mathcal{B}}(\boldsymbol{\eta}) = e^{-\beta V_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta}) - \beta S_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta})} \cdot \zeta_{\mathcal{B}}(\boldsymbol{\eta}) \cdot \frac{Z_{\mathcal{G},\eta}^{\mathrm{G}}(\mathbf{0})}{Z_{\Lambda,f}^{\mathrm{G}}(\mathbf{0})} \cdot \frac{\Xi_{\mathcal{G},\eta}(\mathbf{0},+)}{\Xi_{\Lambda}(\mathbf{0})}$$

$$e^{-C_L|\mathcal{B}_1|} \leq \frac{\Xi_{\mathcal{G},\eta}(\mathbf{0},+)}{\Xi_{\Lambda}(\mathbf{0})} \leq e^{C_L|\mathcal{B}_1|}$$

$$\frac{\Xi_{\mathcal{G},\eta}(\mathbf{0},+)}{\Xi_{\Lambda}(\mathbf{0})}$$

$$\mathbf{1}_{\{|\theta| \leq T_\beta\}} \leq \chi\left(\frac{|\theta|}{T_\beta}\right) e^{-\beta g(\theta)} \leq 2,$$

$$-\beta g(\theta) = \beta \cdot (\cos(\theta) - 1 + \theta^2/2) \lesssim \beta T_\beta^{-4} \lesssim \frac{\log^{O(1)}(\beta)}{\beta}$$

$$\chi\left(\frac{T_\beta}{T^2}\right) \equiv 1 \text{ for } |\theta| \leq T_\beta$$

$$\chi_Q(\boldsymbol{\theta}) = \prod_{e \in E(Q)} \chi\left(\frac{|\theta_e|}{T_\beta}\right)$$

$$\frac{1}{C_L} \leq \int_{\mathbb{R}^{E(Q)}} \chi_Q(\boldsymbol{\theta}) e^{-\beta V_Q(\boldsymbol{\theta})} \mathrm{d}\mu_{Q,f}^{\mathrm{G}}(\boldsymbol{\theta}) \leq C_L$$

$$\left(\frac{1}{C_L}\right)^{|\mathcal{B}_1|} \leq \frac{\Xi_{\mathcal{G},\eta}(\mathbf{0},+)}{\Xi_{\Lambda}(\mathbf{0})} \leq C_L^{|\mathcal{B}_1|}$$

$$\int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} \rho_{\mathcal{B}}(\boldsymbol{\eta}) \mathrm{d}\boldsymbol{\eta}_{\mathcal{B}} \leq \exp\left(C_L|\mathcal{B}_1| \cdot (\log \beta) - c_L \beta T_\beta^2 |\mathcal{B}|\right)$$

$$\int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} e^{-\beta V_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta}) - \beta S_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta})} \zeta_{\mathcal{B}}(\boldsymbol{\eta}) \mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}$$
$$\dot{\leq} \int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} e^{-\frac{\beta}{5} S_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta}) - \beta V_{\mathrm{s}}(\boldsymbol{\eta})} \zeta_{\mathcal{B}}(\boldsymbol{\eta}) \mathrm{d}\boldsymbol{\eta}_{\mathcal{B}} \leq e^{-c_L \beta T_\beta^2 |\mathcal{B}|}$$

$$\frac{Z_{\mathcal{G},\eta}^{\mathrm{G}}(\mathbf{0})}{Z_{\Lambda,f}^{\mathrm{G}}(\mathbf{0})} \leq \beta^{C_L|\mathcal{B}_1|}$$

$$\int_{\mathbb{R}^{E(Q)}} e^{-\beta S_Q(\boldsymbol{\theta})} \prod_{e \in E(Q)} \mathrm{d}\theta_e \gtrsim \beta^{-|E(Q)|/2} = \beta^{-C_L}$$

$$\int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} \rho_{\mathcal{B}}(\boldsymbol{\eta})\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}} = \int_{[-\pi,\pi]^{\mathcal{E}(\mathcal{B})}} e^{-\beta V_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta})-\beta S_{\Lambda}^{\mathcal{B}}(\boldsymbol{\eta})}\zeta_{\mathcal{B}}(\boldsymbol{\eta}) \cdot \frac{Z_{\Lambda,\eta}^{\mathrm{G}}(\mathbf{0})}{Z_{\Lambda,f}^{\mathrm{G}}(\mathbf{0})} \cdot \frac{\Xi_{\mathcal{G},\eta}(\mathbf{0},+)}{\Xi_{\Lambda}(\mathbf{0})}\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}$$
$$\leq \beta^{C_L|\mathcal{B}_1|}\exp\left(-c_L\beta T_{\beta}^2|\mathcal{B}|\right)$$
$$= \exp\left(C_L|\mathcal{B}_1| \cdot (\log\,\beta) - c_L\beta T_{\beta}^2|\mathcal{B}|\right)$$

$$K_1(\emptyset,\mathcal{G}) = \frac{\Xi_{\mathcal{G},\eta}(\mathbf{0})}{\Xi_{\mathcal{G},\eta}(\mathbf{0},+)}$$

$$K_1(\emptyset,\mathcal{G}) = \frac{\Xi_{\Lambda}(\mathbf{0})}{\Xi_{\mathcal{G},\eta}(\mathbf{0},+)} \cdot \frac{\Xi_{\mathcal{G},\eta}(\mathbf{0})}{\Xi_{\Lambda}(\mathbf{0})} \leq e^{C_L|\mathcal{B}_1|} \cdot \frac{\Xi_{\mathcal{G},\eta}(\mathbf{0})}{\Xi_{\Lambda}(\mathbf{0})}$$

$$\frac{\Xi_{\mathcal{G},\eta}(\mathbf{0})}{\Xi_{\Lambda}(\mathbf{0})} \leq e^{C_L|\mathcal{B}_1|}$$

$$T_{\beta} = \log^{d+2}(\beta)/\sqrt{\beta}$$

$$\tilde{\rho}(X) = \sum_{\substack{\mathcal{B}\subset\Lambda' \\ \mathcal{B}_1=X}} \left(\int_{[-\pi,\pi]} \mathcal{E}_{\mathcal{B}}\rho_{\mathcal{B}}(\boldsymbol{\eta})\mathrm{d}\boldsymbol{\eta}_{\mathcal{B}}\right)$$
$$\leq \sum_{\substack{\mathcal{B}\subset\Lambda' \\ \mathcal{B}_1=X}} \exp\left(C_L|\mathcal{B}_1| \cdot (\log\,\beta) - c_L\beta T_{\beta}^2|\mathcal{B}|\right)$$
$$= e^{-\log(\beta)|X|} \sum_{\substack{\mathcal{B}\subset\Lambda' \\ \mathcal{B}_1=X}} \exp\left((C_L+1)|\mathcal{B}_1| \cdot (\log\,\beta) - c_L\log^{2d+4}(\beta) \cdot |\mathcal{B}|\right)$$

$$|\mathcal{B}_1| \lesssim r_{\beta}^d \cdot |\mathcal{B}| = \log^{2d}(\beta) \cdot |\mathcal{B}|$$

$$\tilde{\rho}(X) \leq e^{-\log(\beta)|X|} \cdot \sum_{\mathcal{B}\subset X} \varepsilon_{\beta}^{|\mathcal{B}|},$$

$$\varepsilon_{\beta} = \exp\left(\log^{2d+2}(\beta) - c_L\log^{2d+4}(\beta)\right)$$

$$\tilde{f}(\mathcal{B}) = \varepsilon_{\beta}^{|\mathcal{B}|}$$

$$\sum_{\mathcal{B}\subset X} \varepsilon_{\beta}^{|\mathcal{B}|} \dot{\leq} \exp\left(|X| \cdot \|\tilde{f}\|_0\right) \dot{\leq} \exp(|X|)$$

$$\tilde{\rho}(X) \leq e^{-(\log(\beta)-1)|X|}$$

$$W_2(\Gamma;\mathcal{G}) = W_1(\Gamma;\mathcal{G}) - W_1(\Gamma;\Lambda)$$

$$W_1(\Gamma;\mathcal{G}) = \int_{[0,1]^{\Gamma}} \partial^{\Gamma}\log\, Z_{\mathcal{G},\eta}^{\mathrm{G}}(\mathbf{s}_{\Gamma})\mathrm{d}s_{\Gamma}$$

$$\mathrm{Cov}_{\mathbf{s}}(\theta_x,\theta_y) = \mathbb{E}_{\mathbf{s}}^{\mathrm{G}}[\theta_x\theta_y] - \mathbb{E}_{\mathbf{s}}^{\mathrm{G}}[\theta_x] \cdot \mathbb{E}_{\mathbf{s}}^{\mathrm{G}}[\theta_y]$$

$$\left|\mathrm{Cov}_{\mathbf{s}}(\theta_x,\theta_y)\right| \leq C\beta^{-1}e^{-cm\mathrm{dist}(x,y)}$$

$$R^{-1}(x,y) = \mathrm{Cov}_{\mathbf{s}}\big(\theta_x, \theta_y\big)$$

$$S_{\Lambda,\mathcal{G}}(\boldsymbol{\theta};\mathbf{s}) = \frac{1}{2} \sum_{\substack{p \in P(\Lambda) \\ E(p)\cap\mathcal{E}(\mathcal{G})\neq\emptyset}} \sigma_p(\mathbf{s})\big(\mathbf{d}\theta_p\big)^2 + \frac{m}{2} \sum_{e\in\mathcal{E}(\mathcal{G})} (\theta_e)^2 = \frac{1}{2}\langle \boldsymbol{\theta}, R\boldsymbol{\theta}\rangle$$

$$\psi \in \ell^2(\mathcal{E}(\mathcal{G}))$$

$$\langle \psi, R\psi\rangle \geq m\|\psi\|^2$$

$$\langle \psi, R\psi\rangle \leq m\|\psi\|^2 + C_d\|\psi\|^2$$

$$S = I - \frac{1}{m + C_d} R$$

$$(m + C_d)R^{-1} = (I - S)^{-1} = \sum_{k=0}^{\infty} S^k$$

$$\begin{aligned}
\big|\mathrm{Cov}_{\mathbf{s}}\big(\theta_x, \theta_y\big)\big| &\leq \frac{1}{m + C_d} \sum_{k=0}^{\infty} |S(x,y)|^k \\
&= \frac{1}{m + C_d} \sum_{k=\lfloor c\mathrm{dist}(x,y)\rfloor}^{\infty} |S(x,y)|^k \\
&\leq \frac{1}{m + C_d} \sum_{k=\lfloor c\mathrm{dist}(x,y)\rfloor}^{\infty} \|S\|^k \\
&\leq \frac{1}{m + C_d} \sum_{k=\lfloor c\mathrm{dist}(x,y)\rfloor}^{\infty} \left(1 - \frac{m}{m + C_d}\right)^k
\end{aligned}$$

$$\partial^{\Gamma} = \prod_{\gamma\in\Gamma} \frac{\partial}{\partial s_\gamma}$$

$$\sup_{\mathbf{s}} \big|\partial^{\Gamma}\mathbb{E}^{\mathrm{G}}_{\mathbf{s}}[\theta_x]\big| \leq c_1\beta^{-1}\big(c_2L^d\big)^{|\Gamma|} e^{-c_3 m d(x,\mathcal{B}_1;\Gamma)}$$

$$\sup_{\mathbf{s}} \big|\partial^{\Gamma}\mathrm{Cov}_{\mathbf{s}}\big(\theta_x, \theta_y\big)\big| \leq c_1\beta^{-1}\big(c_2L^d\big)^{|\Gamma|} e^{-c_3 m d(x,y;\Gamma)}$$

$$\langle \boldsymbol{\theta}, \mathrm{d}^*\,\mathrm{d}(\mathbf{s})\boldsymbol{\theta}\rangle = \sum_{\substack{p \in P(\Lambda) \\ E(p)\cap\mathcal{E}(\mathcal{G})\neq\emptyset}} \sigma_p(\mathbf{s})\big(\mathrm{d}\theta_p\big)^2$$

$$\beta\mathrm{Cov}_{\mathbf{s}}\big(\theta_x, \theta_y\big) = [mI + \mathrm{d}^*\,\mathrm{d}(\mathbf{s})]^{-1}(x,y)$$

$$\frac{\partial}{\partial t}A(t)^{-1} = -A(t)^{-1} \cdot \left(\frac{\partial A(t)}{\partial t}\right) \cdot A(t)^{-1}$$

$$\frac{\partial}{\partial s_{\mathrm{F}}}[mI + \mathrm{d}^*\,\mathrm{d}(\mathbf{s})]^{-1} = -[mI + \mathrm{d}^*\,\mathrm{d}(\mathbf{s})]^{-1} \cdot \left(\frac{\partial\, \mathrm{d}^*\,\mathrm{d}(\mathbf{s})}{\partial s_{\mathrm{F}}}\right) \cdot [mI + \mathrm{d}^*\,\mathrm{d}(\mathbf{s})]^{-1}$$

$$A_{\mathrm{F}} = \frac{\partial}{\partial s_{\mathrm{F}}} \, \mathrm{d}^{*} \, \mathrm{d}(\mathbf{s})$$

$$\partial^{\Gamma}[mI + \mathrm{d}^{*} \, \mathrm{d}(\mathrm{s})]^{-1} = (-1)^{n+1} \sum_{\sigma \in S_n} \left(\prod_{j=1}^{n} [mI + \mathrm{d}^{*} \, \mathrm{d}(\mathrm{s})]^{-1} \cdot A_{\gamma_{\sigma(j)}} \right) \cdot [mI + \mathrm{d}^{*} \, \mathrm{d}(\mathrm{s})]^{-1}$$

$$\left| \partial^{\Gamma}[mI + \mathrm{d}^{*} \, \mathrm{d}(\mathrm{s})]^{-1}(x, y) \right| \lesssim C^{n} \sum_{\substack{w: x \to y \\ w = \{q_1, \ldots, q_n\}}} \prod_{j=1}^{n} \left| [mI + \mathrm{d}^{*} \, \mathrm{d}(\mathrm{s})]^{-1}(q_j, q_{j+1}) \right|$$

$$\beta \left| \mathrm{Cov}_{\mathbf{s}}(\theta_x, \theta_y) \right| \lesssim \beta^{-1} \left(cL^{d} \right)^{|\Gamma|} \sup_{\substack{w: x \to y \\ w = \{q_1, \ldots, q_n\}}} \prod_{j=1}^{n} \left| [mI + \mathrm{d}^{*} \, \mathrm{d}(\mathbf{s})]^{-1}(q_j, q_{j+1}) \right| \dot{\lesssim}$$

$$\beta^{-1} \left(cL^{d} \right)^{|\Gamma|} \sup_{\substack{w: x \to y \\ w = \{q_1, \ldots, q_n\}}} \prod_{j=1}^{n} C e^{-cm \mathrm{dist}(q_j, q_{j+1})} \lesssim \beta^{-1} \left(CL^{d} \right)^{|\Gamma|} e^{-cm d(x, y; \Gamma)}$$

$$\mathbb{E}_{\mathbf{s}}^{\mathrm{G}}[\theta_x] = C_d \sum_{y \in \partial \mathcal{B}} \mathrm{Cov}_{\mathbf{s}}(\theta_x, \theta_y) \theta_y$$

$$\partial^{\Gamma} \mathbb{E}_{\mathbf{s}}^{\mathrm{G}}[F(\boldsymbol{\theta})] = \sum_{\Pi \in \mathcal{P}(\Gamma)} \mathbb{E}_{\mathbf{s}}^{\mathrm{G}} \left[\left(\prod_{\pi \in \Pi} D^{\pi} \right) F(\boldsymbol{\theta}) \right]$$

$$D^{\pi} F = \frac{1}{2} \sum_{x, y \in \mathcal{E}(\mathcal{G})} \left(\partial^{\pi} \mathrm{Cov}_{\mathbf{s}}(\theta_x, \theta_y) \right) \frac{\partial^2 F}{\partial \theta_x \partial \theta_y} + \sum_{x \in \mathcal{E}(\mathcal{G})} (\partial^{\pi} \mathbb{E}_{\mathbf{s}}^{G}[\theta_x]) \frac{\partial F}{\partial \theta_x}$$

$$\Sigma_t = (t\Sigma_1^{-1} + (1-t)\Sigma_0^{-1})^{-1}, t \in [0,1]$$

$$\frac{\partial}{\partial t} \mathbb{E}[F(X_t)] = \frac{1}{2} \sum_{i,j=1}^{N} \frac{\partial \Sigma_t}{\partial t}(i,j) \cdot \mathbb{E} \left[\frac{\partial}{\partial x_i} \frac{\partial}{\partial x_j} F(X_t) \right]$$

$$A = \Sigma_1^{-1} - \Sigma_0^{-1} = \frac{\partial}{\partial t} \Sigma_t^{-1}$$

$$\log \, p_t(x) = C_N + \frac{1}{2} \log \, \det(\Sigma_t^{-1}) - \frac{1}{2} \langle x, \Sigma_t^{-1} x \rangle,$$

$$\frac{\partial}{\partial t} \log \, p_t(x) = \frac{1}{2} (\mathrm{Tr}(\Sigma_t \cdot A) - \langle x, Ax \rangle) = \frac{1}{2} \sum_{\ell, k=1}^{N} A(k, \ell)(\Sigma_t(k, \ell) - x_k x_\ell)$$

$$\frac{\partial}{\partial t}\mathbb{E}[F(X_t)]=\int_{\mathbb{R}^\eta} F(x)\frac{\partial}{\partial t}p_t(x)\mathrm{d}x$$
$$=\int_{\mathbb{R}^\eta} F(x)p_t(x)\frac{\partial}{\partial t}\log\, p_t(x)\mathrm{d}x$$
$$=\frac{1}{2}\sum_{\ell,k=1}^{N} A(k,\ell)\cdot\mathbb{E}[F(X_t)(\Sigma_t(k,\ell)-(X_t)_k(X_t)_\ell)]$$

$$\mathbb{E}[F(X_t)(X_t)_k]=\sum_{j=1}^{N}\Sigma_t(k,j)\cdot\mathbb{E}\left[\frac{\partial}{\partial x_j}F(X_t)\right]$$

$$\mathbb{E}[F(X_t)(X_t)_k(X_t)_\ell]=\sum_{j=1}^{N}\Sigma_t(k,j)\cdot\mathbb{E}\left[\frac{\partial}{\partial x_j}(F(X_t)(X_t)_\ell)\right]$$
$$=\sum_{j=1}^{N}\Sigma_t(k,j)\cdot\left(\sum_{i=1}^{N}\Sigma_t(i,\ell)\cdot\mathbb{E}\left[\frac{\partial^2}{\partial x_j\partial x_i}F(X_t)\right]+\delta_{j=\ell}\mathbb{E}[F(X_t)]\right)$$
$$=\sum_{i,j=1}^{N}\Sigma_t(k,j)\Sigma_t(i,\ell)\mathbb{E}\left[\frac{\partial^2}{\partial x_j\partial x_i}F(X_t)\right]+\Sigma_t(k,\ell)\mathbb{E}[F(X_t)]$$

$$\frac{\partial}{\partial t}\mathbb{E}[F(X_t)]=-\frac{1}{2}\sum_{k,\ell=1}^{N}A(k,\ell)\cdot\left(\sum_{i,j=1}^{N}\Sigma_t(k,j)\Sigma_t(i,\ell)\mathbb{E}\left[\frac{\partial^2}{\partial x_j\partial x_i}F(X_t)\right]\right)$$
$$=-\frac{1}{2}\sum_{i,j=1}^{N}(\Sigma_t\cdot A\cdot\Sigma_t)(i,j)\cdot\mathbb{E}\left[\frac{\partial^2}{\partial x_j\partial x_i}F(X_t)\right]$$
$$\dot{=}-\frac{1}{2}\sum_{i,j=1}^{N}\left(\Sigma_t\cdot\frac{\partial\Sigma_t^{-1}}{\partial t}\cdot\Sigma_t\right)(i,j)\cdot\mathbb{E}\left[\frac{\partial^2}{\partial x_j\partial x_i}F(X_t)\right]$$
$$\dot{=}\frac{1}{2}\sum_{i,j=1}^{N}\frac{\partial\Sigma_t}{\partial t}(i,j)\cdot\mathbb{E}\left[\frac{\partial^2}{\partial x_j\partial x_i}F(X_t)\right]$$

$$W_1(\Gamma;\mathcal{G})=\int_{[0,1]^\Gamma}\partial^\Gamma\log\, Z^{\mathrm{G}}_{\mathcal{G},\eta}(\mathbf{s}_\Gamma)\mathrm{d}s_\Gamma$$

$$\frac{\partial}{\partial s_\gamma}\log\, Z^{\mathrm{G}}_{\mathcal{G},\eta}(\mathbf{s}_\Gamma)=-\mathbb{E}^{\mathrm{G}}_{\mathbf{s}}\left[\frac{\partial}{\partial s_\gamma}S(\boldsymbol{\theta};\mathbf{s})\right]$$

$$\partial^{\widetilde{\Gamma}}\mathbb{E}^{\mathrm{G}}_{\mathbf{s}}\left[\frac{\partial}{\partial s_p}S(\boldsymbol{\theta};\mathbf{s})\right]=\sum_{\Pi\in\mathcal{P}(\widetilde{\Gamma})}\mathbb{E}^{\mathrm{G}}_{\mathbf{s}}\left[\left(\prod_{\pi\in\Pi}D^\pi\right)\frac{\partial S(\boldsymbol{\theta},\mathbf{s})}{\partial s_\gamma}\right]$$

$$\left|\partial^\Gamma\log\, Z^{\mathrm{G}}_{\mathcal{G},\eta}(\mathbf{s}_\Gamma)\right|\leq 2^{|\Gamma|}\sup_{\Pi\in\mathcal{P}(\widetilde{\Gamma})}\left|\mathbb{E}^{\mathrm{G}}_{\mathbf{s}}\left[\left(\prod_{\pi\in\Pi}D^\pi\right)\frac{\partial S(\boldsymbol{\theta},\mathbf{s})}{\partial s_\gamma}\right]\right|$$

$$\left|\partial^\Gamma\log\, Z^{\mathrm{G}}_{\mathcal{G},\eta}(\mathbf{s}_\Gamma)\right|\lesssim\beta^{-1}\left(CL^d\right)^{|\Gamma|}e^{-cmd(\Gamma)}$$

$$\partial^{\pi}\mathrm{Cov}_{\mathbf{s}}(\theta_x,\theta_y) \text{ and } \partial^{\pi}\mathbb{E}^{\mathrm{G}}_{\mathbf{s}}[\theta_x]$$

$$d(\Gamma) \geq \frac{L}{2d}|\Gamma|$$

$$|W_1(\Gamma;\mathcal{G})| \lesssim \left(CL^d\right)^{|\Gamma|} e^{-cmL|\Gamma|}$$

$$CL^d e^{-cmL}$$

$$\|W_2(\bullet,\mathcal{G})\|_r \lesssim \|W_1(\bullet,\mathcal{G})\|_r \leq C_L$$

$$K_1(\Delta;\mathcal{G}) = \frac{1}{\Xi_{\mathcal{G},\eta}(\mathbf{0},+)}\int_{[0,1]^{\Delta}} \partial^{\Delta}\Xi_{\mathcal{G},\eta}(\mathbf{s}_{\Delta})\mathrm{d}\mathbf{s}_{\Delta}$$

$$\Xi_{\mathcal{G},\eta}(\mathbf{s}) = \int_{\mathbb{R}^{\mathcal{E}(\mathcal{G})}} e^{-\beta V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta};\mathbf{s})}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu^{\mathrm{G}}_{\mathcal{G},\eta,\mathbf{s}} = \mathbb{E}^{\mathrm{G}}_{\mathcal{G},\eta,\mathbf{s}}\left[e^{-\beta V_{\Lambda,\mathcal{G}}(\boldsymbol{\theta};\mathbf{s})}\chi_{\mathcal{G}}(\boldsymbol{\theta})\right]$$

$$(\mathbf{s}_{\Delta})_{\mathrm{F}} = \begin{cases} s_{\mathrm{F}} & \text{if } \mathrm{F} \in \Delta \\ 0 & \text{otherwise} \end{cases}$$

$$\mathcal{X} = X(\Delta) \cup (\mathcal{B}_1 \setminus \mathcal{B})$$

$$\left|\partial^{\Delta}\int_{\mathbb{R}^{\mathcal{E}(\mathcal{X})}} e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_{\Delta})}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_{\Delta}}\right| \leq \nu(\beta)\exp\left(-R|X(\Delta)| + C_L|\mathcal{B}_1|\right),$$

$$\lim_{\beta\to\infty}\nu(\beta) = 0$$

$$\begin{aligned}\int_{\mathbb{R}^{E(Q)}} \chi_Q(\boldsymbol{\theta})e^{-\beta V_Q(\boldsymbol{\theta})}\mathrm{d}\mu^{\mathrm{G}}_{Q,f}(\boldsymbol{\theta}) &\geq \int_{\mathbb{R}^{E(Q)}} 1\{\forall e \in Q, |\theta_e| \leq T_\beta\}\mathrm{d}\mu^{\mathrm{G}}_{Q,f}(\boldsymbol{\theta}) \\ &= 1 - \int_{\mathbb{R}^{E(Q)}} 1\{\exists e \in Q, |\theta_e| \geq T_\beta\}\mathrm{d}\mu^{\mathrm{G}}_{Q,f}(\boldsymbol{\theta})\end{aligned}$$

$$\int_{\mathbb{R}^{E(Q)}} 1\{\exists e \in Q, |\theta_e| \geq T_\beta\}\mathrm{d}\mu^{\mathrm{G}}_{Q,f}(\boldsymbol{\theta}) \lesssim |E(Q)|\int_{\beta T_\beta}^{\infty} e^{-x^2}\,\mathrm{d}x \leq C_L e^{-\log^{2d}(\beta)} \leq \frac{1}{4}$$

$$\int_{\mathbb{R}^{E(Q)}} \chi_Q(\boldsymbol{\theta})e^{-\beta V_Q(\boldsymbol{\theta})}\mathrm{d}\mu^{\mathrm{G}}_{Q,f}(\boldsymbol{\theta}) \geq \frac{1}{2}$$

$$\begin{aligned}&|K_1(\Delta;\mathcal{G})| \\ &\leq 2^{|\mathcal{B}_1|+|X(\Delta)|}\left|\partial^{\Delta}\int_{\mathbb{R}^{\mathcal{E}(\mathcal{X})}} e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_{\Delta})}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_{\Delta}}\right| \dot{\leq} \nu(\beta)e^{-(R-2)|X(\Delta)|+C_L|\mathcal{B}_1|}.\end{aligned}$$

$$\begin{aligned}&e^{-C'_L|\mathcal{B}_1|}\sum_{\Delta\in\mathcal{C}(\mathcal{B}_1)} |K_1(\Delta;\mathcal{G})|e^{r|X(\Delta)|} \\ &\qquad \leq \nu(\beta)e^{-|\mathcal{B}_1|}\sum_{\Delta\in\mathcal{C}(\mathcal{B}_1)} e^{-(R-2-r)|X(\Delta)|} \dot{\lesssim} \nu(\beta)e^{-|\mathcal{B}_1|}|\mathcal{B}_1| \leq \nu(\beta) \overset{\beta\to\infty}{\to} 0\end{aligned}$$

$$\left|\partial^{\Delta}\int_{\mathbb{R}^{\mathcal{E}(\mathcal{X})}} e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_{\Delta})}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu^{G}_{\mathcal{X},f,\mathbf{s}_{\Delta}}\right| \leq \nu(\beta)e^{-R|X(\Delta)|}$$

$$|K_1(\Delta;\Lambda')| \le \nu(\beta)2^{|X(\Delta)|}e^{-R|X(\Delta)|}$$

$$\sum_{\substack{\Delta_{\text{connected}}\\ \Delta_{\text{touch }x}}} K_1(\Delta;\Lambda')e^{r|X(\Gamma)|} \le \nu(\beta)\sum_{\substack{\Delta_{\text{connected}}\\ \Delta_{\text{touch }x}}} e^{-(R-r-\log 2)|X(\Delta)|} \dot{\lesssim} \nu(\beta)$$

$$\lim_{\beta\to\infty}\|K_1(\cdot;\Lambda')\|_r = 0$$

$$\partial^{\Delta}\int_{\mathbb{R}^{\mathcal{E}(\mathcal{X})}} e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_\Delta)}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_\Delta}$$

$$= \sum_{\Upsilon\subset\Delta}\partial^{\Upsilon}_{\mathrm{G}}\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_\Delta}\left[\left(\prod_{\mathrm{F}\in\Delta\setminus\Upsilon}\sum_{\substack{p\in P(\Lambda)\\ E(p)\cap\mathrm{F}\neq\emptyset}}\beta g(\,\mathrm{d}\theta_p)\right)\cdot e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_\Delta)}\chi_{\mathcal{G}}(\boldsymbol{\theta})\right]$$

$$\left|\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_\Delta}\left[\left(\prod_{\pi\in\Pi}D^{\pi}\right)e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_\Delta)}\chi_{\mathcal{G}}(\boldsymbol{\theta})\right]\right| \le C^{|\mathcal{X}|}\prod_{\pi\in\Pi}\left(\nu(\beta)\left(L^d\right)^{|\pi|}e^{-cd(\pi)}\right),$$

$$\lim_{\beta\to\infty}\nu(\beta) = 0$$

$$\beta|g(\,\mathrm{d}\theta_p)| \lesssim \beta\left(T_\beta\right)^4 \le \frac{\log^{4d+8}(\beta)}{\beta}$$

$$|\,\partial^{\Delta}\int_{\mathbb{R}^{\mathcal{E}(\mathcal{X})}} e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_\Delta)}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_\Delta}\,|$$

$$\le C^{|\Delta|}\sup_{\Upsilon\subset\Delta}\left[\left(L^{d-1}\frac{\log^{4d+8}(\beta)}{\beta}\right)^{|\Delta\setminus\Upsilon|}\sum_{\Pi\in\mathcal{P}(\Upsilon)}\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_\Delta}\left[\left(\prod_{\pi\in\Pi}D^{\pi}\right)e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_\Delta)}\chi_{\mathcal{G}}(\boldsymbol{\theta})\right]\right]$$

$$\left|\sum_{\Pi\in\mathcal{P}(\Upsilon)}\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_\Delta}\left[\left(\prod_{\pi\in\Pi}D^{\pi}\right)e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_\Delta)}\chi_{\mathcal{G}}(\boldsymbol{\theta})\right]\right| \le C^{|\mathcal{X}|}\sum_{\Pi\in\mathcal{P}(\Upsilon)}\prod_{\pi\in\Pi}\left(\nu(\beta)\left(L^d\right)^{|\pi|}e^{-cd(\pi)}\right).$$

$$\left|\partial^{\Delta}\int_{\mathbb{R}^{\mathcal{E}(\mathcal{X})}} e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_\Delta)}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu^{G}_{\mathcal{X},\eta,\mathbf{s}_\Delta}\right|$$

$$\le C^{|X(\Delta)|+|\mathcal{B}_1|}\sup_{\Upsilon\subset\Delta}\left[\left(L^{d-1}\frac{\log^{4d+8}(\beta)}{\beta}\right)^{|\Delta\setminus\Upsilon|}\sum_{\Pi\in\mathcal{P}(\Upsilon)}\prod_{\pi\in\Pi}\left(\nu(\beta)\left(L^d\right)^{|\pi|}e^{-cd(\pi)}\right)\right]$$

$$\sum_{\Pi\in\mathcal{P}(\Upsilon)}\prod_{\pi\in\Pi}\left(\nu(\beta)\left(L^d\right)^{|\pi|}e^{-cd(\pi)}\right)$$

$$= e^{-R|\Upsilon|}\sum_{\Pi\in\mathcal{P}(\Upsilon)}\prod_{\pi\in\Pi}\left(\nu(\beta)e^{R|\pi|}\left(L^d\right)^{|\pi|}e^{-cd(\pi)}\right)$$

$$\dot{\le} e^{-R|\Upsilon|}\nu(\beta)\exp\left(|\Upsilon|\cdot\left\|\left(L^d\right)^{|\cdot|}e^{-cd(\cdot)}\right\|_R\right)$$

$$d(\pi) \ge cL|\pi| \triangle \left\|\left(L^d\right)^{|\cdot|}e^{-cd(\cdot)}\right\|_R \le 1$$

$$\sum_{\Pi\in\mathcal{P}(\Upsilon)}\prod_{\pi\in\Pi}\left(\nu(\beta)\left(L^{d}\right)^{|\pi|}e^{-cd(\pi)}\right)\leq e^{-(R-1)|\Upsilon|}\nu(\beta)$$

$$L^{d-1}\frac{\log^{4d+8}(\beta)}{\beta}\leq e^{-2R}$$

$$\left|\partial^{\Delta}\int_{\mathbb{R}^{\mathcal{E}(\mathcal{X})}}e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_{\Delta})}\chi_{\mathcal{G}}(\boldsymbol{\theta})\mathrm{d}\mu^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_{\Delta}}\right|\leq\nu(\beta)C^{|X(\Delta)|+|\mathcal{B}_1|}e^{-R|\Delta|}$$

$$\mathrm{C}(x,y;\pi)=\frac{1}{2}\partial^{\pi}\mathrm{Cov}_{\mathbf{s}}\left(\theta_x,\theta_y\right),\mathrm{m}(z;\pi)=\partial^{\pi}\mathbb{E}^{\mathrm{G}}_{\mathbf{s}}[\theta_z]$$

$$\left(\prod_{\pi\in\Pi}D^{\pi}\right)\left(e^{-\beta V_{\Lambda,\mathcal{X}}}\chi_{\mathcal{G}}\right)=\sum_{\mathbf{x},\mathbf{y},\mathbf{z}\in\mathcal{E}(\mathcal{X})^{N}}\sum_{\varepsilon\in\{0,1\}^{N}}$$
$$\left(\left[\prod_{\ell=1}^{N}\left(\left(\mathrm{C}(x_\ell,y_\ell;\pi_\ell)\right)^{\varepsilon_\ell}\left(\mathrm{m}(z_\ell;\pi_\ell)\right)^{1-\varepsilon_\ell}\right)\right]\times\left[\prod_{\ell=1}^{N}\left(\frac{\partial^2}{\partial\theta_{x_\ell}\partial\theta_{y_\ell}}\right)^{\varepsilon_\ell}\cdot\left(\frac{\partial}{\partial\theta_{z_\ell}}\right)^{1-\varepsilon_\ell}\right]\left(e^{-\beta V_{\Lambda,\mathcal{X}}}\chi_{\mathcal{G}}\right)\right)$$

$$\left|\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_{\Delta}}\left[\left(\prod_{\pi\in\Pi}D^{\pi}\right)e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_{\Delta})}\chi_{\mathcal{G}}(\boldsymbol{\theta})\right]\right|\leq$$
$$C^{|\mathcal{X}|}\sum_{\mathbf{x},\mathbf{y},\mathbf{z}\in\mathcal{E}(\mathcal{X})^{N}}\sum_{\boldsymbol{\varepsilon}\in\{0,1\}^{N}}\beta^{-N}\cdot\left(\prod_{\ell=1}^{N}\left(c_2L^{d}\right)^{|\pi_\ell|}\right)e^{-c_3m\sum_{\ell=1}^{N}\left[\varepsilon_\ell d(x_\ell,y_\ell;\pi_\ell)+(1-\varepsilon_\ell)d(z_\ell,\mathcal{B}_1;\pi_\ell)\right]}$$
$$\times\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_{\Delta}}\left|\mathcal{D}(\mathbf{x},\mathbf{y},\mathbf{z},\boldsymbol{\varepsilon})\left(e^{-\beta V_{\Lambda,\mathcal{X}}}\chi_{\mathcal{G}}\right)\right|$$

$$\mathcal{D}(\mathbf{x},\mathbf{y},\mathbf{z},\varepsilon)=\prod_{\ell=1}^{N}\left(\frac{\partial^2}{\partial\theta_{x_\ell}\partial\theta_{y_\ell}}\right)^{\varepsilon_\ell}\cdot\left(\frac{\partial}{\partial\theta_{z_\ell}}\right)^{1-\varepsilon_\ell}$$

$$\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_{\Delta}}\left|\mathcal{D}(\mathbf{x},\mathbf{y},\mathbf{z},\boldsymbol{\varepsilon})\left(e^{-\beta V_{\Lambda,\mathcal{X}}}\chi_{\mathcal{G}}\right)\right|\leq C^{|\mathcal{X}|}\left(\frac{\beta}{\log\beta}\right)^{N}\prod_{(x,y)\in\mathcal{E}(\mathcal{X})^2}(\mathrm{n}(x,y)!)^2\prod_{z\in\mathcal{E}(\mathcal{X})}(\mathrm{n}(z)!)^2,$$

$$\prod_{(x,y)\in\mathcal{E}(\mathcal{X})^2}(\mathrm{n}(x,y)!)^2\prod_{z\in\mathcal{E}(\mathcal{X})}(\mathrm{n}(z)!)^2\leq C_{\eta}^{|\mathcal{X}|}\exp\left(\eta\sum_{\ell=1}^{N}\left[\varepsilon_\ell d(x_\ell,y_\ell;\pi_\ell)+(1-\varepsilon_\ell)d(z_\ell,\mathcal{B}_1;\pi_\ell)\right]\right)$$

$$\left|\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_{\Delta}}\left[\left(\prod_{\pi\in\Pi}D^{\pi}\right)e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_{\Delta})}\chi_{\mathcal{G}}(\boldsymbol{\theta})\right]\right|$$
$$\leq C^{|\mathcal{X}|}\sum_{\mathbf{x},\mathbf{y},\mathbf{z}\in\mathcal{E}(\mathcal{X})^{N}}\sum_{\varepsilon\in\{0,1\}^{N}}(\log(\beta))^{-N}\cdot\left(\prod_{\ell=1}^{N}\left(c_2L^{d}\right)^{|\pi_\ell|}\right)e^{-cm\sum_{\ell=1}^{N}\left[\varepsilon_\ell d(x_\ell,y_\ell;\pi_\ell)+(1-\varepsilon_\ell)d(z_\ell,\mathcal{B}_1;\pi_\ell)\right]}$$
$$\leq C^{|\mathcal{X}|}\frac{1}{(\log(\beta))^{|\Pi|}}\prod_{\pi\in\Pi}\left(\sum_{x,y\in\mathcal{E}(\mathcal{X})}\left(c_2L^{d}\right)^{|\pi|}e^{-cmd(x,y;\pi)}+\sum_{z\in\mathcal{E}(\mathcal{X})}\left(c_2L^{d}\right)^{|\pi|}e^{-cmd(z,\mathcal{B}_1;\pi)}\right)$$

$$\max\left\{\sum_{x,y\in\mathcal{E}(\mathcal{X})}\left(c_2L^{d}\right)^{|\pi|}e^{-cmd(x,y;\pi)},\sum_{z\in\mathcal{E}(\mathcal{X})}\left(c_2L^{d}\right)^{|\pi|}e^{-cmd(z,\mathcal{B}_1;\pi)}\right\}\lesssim\left(L^{d}\right)^{|\pi|}e^{-cd(\pi)}$$

$$\left|\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_\Delta}\left[\left(\prod_{\pi\in\Pi} D^\pi\right)e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta};\mathbf{s}_\Delta)}\chi_{\mathcal{G}}(\boldsymbol{\theta})\right]\right| \leq C^{|\mathcal{X}|}\prod_{\pi\in\Pi}\left(\frac{1}{\log(\beta)}\left(L^d\right)^{|\pi|}e^{-cd(\pi)}\right)$$

$$\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_\Delta}\left|\mathcal{D}(\mathbf{x},\mathbf{y},\mathbf{z},\boldsymbol{\varepsilon})\left(e^{-\beta V_{\Lambda,\mathcal{X}}}\chi_{\mathcal{G}}\right)\right| \leq 2^{\tilde{N}}\max_{\mathbf{w}=\{\mathbf{w}^1,\mathbf{w}^2\}}\sup_{\boldsymbol{\theta}}\left|\mathcal{D}(\mathbf{w}^1)\left(e^{-\beta V_{\Lambda,\mathcal{X}}}\right)\right|\cdot\sup_{\boldsymbol{\theta}}\left|\mathcal{D}(\mathbf{w}^2)\left(\chi_{\mathcal{G}}\right)\right|,$$

$$\sup_{\boldsymbol{\theta}}\left|\mathcal{D}(\mathbf{w}^1)\left(e^{-\beta V_{\Lambda,\mathcal{X}}}\right)\right| \leq \left(\frac{\sqrt{\beta}}{\log\beta}\right)^{N_1}\cdot\left(\prod_{w\in\mathbf{w}^1}\mathrm{n}(w)!\right)\cdot\sup_{\boldsymbol{\theta}:|\theta_e|\leq\frac{\log\beta}{\sqrt{\beta}}} e^{-\beta V_{\Lambda,\mathcal{X}}(\boldsymbol{\theta})}$$

$$e^{-\beta g(\theta)} \leq e^{C\beta\frac{\log^4(\beta)}{\beta^2}} \leq 2$$

$$\sup_{\boldsymbol{\theta}}\left|\mathcal{D}(\mathbf{w}^1)\left(e^{-\beta V_{\Lambda,\mathcal{X}}}\right)\right| \leq C^{|\mathcal{X}|}\left(\frac{\sqrt{\beta}}{\log\beta}\right)^{N_1}\cdot\left(\prod_{w\in\mathbf{w}^1}\mathrm{n}(w)!\right).$$

$$\sup_{\boldsymbol{\theta}}\left|\mathcal{D}(\mathbf{w}^2)\left(\chi_{\mathcal{G}}\right)\right| \leq C^{|\mathcal{X}|}T_\beta^{-N_2}\prod_{w\in\mathbf{w}^2}(\mathrm{n}(w)!)^2 = C^{|\mathcal{X}|}\left(\frac{\sqrt{\beta}}{\log^{2d+2}(\beta)}\right)^{N_2}\prod_{w\in\mathbf{w}^2}(\mathrm{n}(w)!)^2.$$

$$\mathbb{E}^{\mathrm{G}}_{\mathcal{X},\eta,\mathbf{s}_\Delta}\left|\mathcal{D}(\mathbf{x},\mathbf{y},\mathbf{z},\boldsymbol{\varepsilon})\left(e^{-\beta V_{\Lambda,\mathcal{X}}}\chi_{\mathcal{G}}\right)\right| \leq C^{|\mathcal{X}|}\left(\frac{\beta}{\log^2(\beta)}\right)^{N_1/2+N_2/2}\prod_{(x,y)\in\mathcal{X}^2}(\mathrm{n}(x,y)!)^2\prod_{z\in\mathcal{X}}(\mathrm{n}(z)!)^2$$

$$\mathrm{n}(x,y) = \{\pi\in\Pi\colon \mathrm{p}(\pi) = (x,y)\}$$

$$\#\{\pi\in\Pi\colon \mathrm{p}(\pi) = (x,y) \text{ and } d(x,y;\pi)\leq r\} \lesssim r^d$$

$$\mathrm{n}(x,y)^{1/d} \leq Cd(x,y;\pi)$$

$$\mathrm{n}(x,y)^{1+1/d} \leq C\sum_{\pi:\mathrm{p}(\pi)=(x,y)} d(x,y;\pi)$$

$$\begin{aligned}\prod_{(x,y)\in\mathcal{E}(\mathcal{X})^2}(\mathrm{n}(x,y)!)^2 &\leq \exp\left(C\sum_{(x,y)\in\mathcal{E}(\mathcal{X})^2}\mathrm{n}(x,y)\log\,\mathrm{n}(x,y)\right)\\ &\leq C_\eta^{|\mathcal{X}|}\exp\left(\eta\sum_{(x,y)\in\mathcal{E}(\mathcal{X})^2}(\mathrm{n}(x,y))^{1+1/d}\mathbf{1}_{\{\mathrm{n}(x,y)\geq 1\}}\right)\\ &\dot{\leq} C_\eta^{|\mathcal{X}|}\exp\left(\eta\sum_{\ell=1}^{N} d(x_\ell,y_\ell;\pi_\ell)\right)\end{aligned}$$

$$\chi(t) = \frac{\eta(t/2)}{\eta(t/2)+\eta(1-|t|/2)}$$

$$\mathcal{L} = -\frac{1}{2}\left|\mathcal{D}_\mu \vec{\Phi}\right|^2 - V(\vec{\Phi}) - \frac{1}{4}F_{\mu\nu}^{\square} - \frac{1}{4}\vec{G}_{\mu\nu}^{\square}$$
$$\mathcal{D}_\mu \vec{\Phi} = \left(D_\mu - igA_\mu\right)\vec{\Phi}, D_\mu \vec{\Phi} = \left(\partial_\mu + g'\vec{B}_\mu \times\right)\vec{\Phi}$$
$$V(\Phi) = \frac{\lambda}{2}\left(|\vec{\Phi}|^2 - \frac{\mu^2}{\lambda}\right)^2$$

$$\vec{B}_\mu = \hat{B}_\mu + \vec{W}_\mu$$
$$\vec{W}_\mu = W_\mu^1 \hat{n}_1 + W_\mu^2 \hat{n}_2, \hat{n} \cdot \vec{W}_\mu = 0$$
$$\hat{B}_\mu = B_\mu \hat{n} - \frac{1}{g'}\hat{n} \times \partial_\mu \hat{n} = \tilde{B}_\mu + \tilde{C}_\mu$$
$$\tilde{B}_\mu = B_\mu \hat{n}\ \left(B_\mu = \hat{n} \cdot \vec{B}_\mu\right)$$
$$\tilde{C}_\mu = -\frac{1}{g'}\hat{n} \times \partial_\mu \hat{n}$$

$$\delta\vec{B}_\mu = \frac{1}{g'}D_\mu \vec{\alpha}, \delta\hat{n} = -\vec{\alpha} \times \hat{n}$$

$$\delta B_\mu = \frac{1}{g'}\hat{n} \cdot \partial_\mu \vec{\alpha},$$
$$\delta\hat{B}_\mu = \frac{1}{g'}\hat{D}_\mu \vec{\alpha}, \delta\vec{W}_\mu = -\vec{\alpha} \times \vec{W}_\mu$$

$$\vec{G}_{\mu\nu} = \hat{G}_{\mu\nu} + \hat{D}_\mu \vec{W}_\nu - \hat{D}_\nu \vec{W}_\mu + g'\vec{W}_\mu \times \vec{W}_\nu$$
$$\hat{D}_\mu = \partial_\mu + g'\hat{B}_\mu \times$$
$$\hat{G}_{\mu\nu} = \partial_\mu \hat{B}_\nu - \partial_\nu \hat{B}_\mu + g'\hat{B}_\mu \times \hat{B}_\nu = G'_{\mu\nu}\hat{n}$$
$$G'_{\mu\nu} = G_{\mu\nu} + H_{\mu\nu} = \partial_\mu B'_\nu - \partial_\nu B'_\mu$$
$$G_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$$
$$H_{\mu\nu} = -\frac{1}{g'}\hat{n} \cdot \left(\partial_\mu \hat{n} \times \partial_\nu \hat{n}\right) = \partial_\mu C_\nu - \partial_\nu C_\mu$$
$$B'_\mu = B_\mu + C_\mu, C_\mu = -\frac{1}{g'}\hat{n}_1 \cdot \partial_\mu \hat{n}_2$$

$$\hat{n} = \begin{pmatrix} \sin\alpha\cos\beta \\ \sin\alpha\sin\beta \\ \cos\alpha \end{pmatrix}$$

$$\tilde{C}_\mu = -\frac{1}{g'}\hat{n} \times \partial_\mu \hat{n} = \frac{1}{g'}\left(\hat{n}_1 \sin\alpha\partial_\mu\beta - \hat{n}_2\partial_\mu\alpha\right)$$

$$\hat{n}_1 = \begin{pmatrix} \cos\alpha\cos\beta \\ \cos\alpha\sin\beta \\ -\sin\alpha \end{pmatrix}, \hat{n}_2 = \begin{pmatrix} -\sin\beta \\ \cos\beta \\ 0 \end{pmatrix},$$
$$C_\mu = -\frac{1}{g'}\hat{n}_1 \cdot \partial_\mu \hat{n}_2 = -\frac{1}{g'}(1 - \cos\alpha)\partial_\mu\beta,$$

$$\mathcal{L} = -\frac{1}{2}\left|\mathcal{D}_\mu\vec{\Phi}\right|^2 - V(\vec{\Phi}) - \frac{1}{4}F_{\mu\nu}^{\square} - \frac{1}{4}\hat{G}_{\mu\nu}^{\square} - \frac{1}{4}\left(\hat{D}_\mu\vec{W}_\nu - \hat{D}_\nu\vec{W}_\mu\right)^2 - \frac{g'}{2}\hat{G}_{\mu\nu}\cdot\left(\vec{W}_\mu\times\vec{W}_\nu\right)$$

$$-\frac{g'^2}{4}\left(\vec{W}_\mu\times\vec{W}_\nu\right)^2$$

$$\hat{D}_\mu = \partial_\mu + g'\hat{B}_\mu\times D_\mu\vec{\Phi} = \left(\hat{D}_\mu + g'\vec{W}_\mu\times\right)\vec{\Phi}$$

$$\vec{\Phi} = \rho\exp(-i\theta)\hat{n}, \rho = |\vec{\Phi}|, \hat{n}^2 = 1$$

$$\mathcal{D}_\mu\vec{\Phi} = \left[\left(\partial_\mu\rho - ig\rho\bar{A}_\mu\right)\hat{n} + g'\rho\vec{W}_\mu\times\hat{n}\right]\exp(-i\theta)$$

$$\bar{A}_\mu = A_\mu + \frac{1}{g}\partial_\mu\theta$$

$$\mathcal{L} = -\frac{1}{2}\left(\partial_\mu\rho\right)^2 - \frac{\lambda}{2}(\rho^2-\rho_0^2)^2 - \frac{1}{4}\bar{F}_{\mu\nu}^{\square} - \frac{g^2}{2}\rho^2\bar{A}_\mu^{\square}$$
$$-\frac{1}{4}G_{\mu\nu}'^{\square} - \frac{1}{4}\left(\hat{D}_\mu\vec{W}_\nu - \hat{D}_\nu\vec{W}_\mu\right)^2 - \frac{g'^2}{2}\rho^2\vec{W}_\mu^2$$
$$-\frac{g'}{2}G_{\mu\nu}'\hat{n}\cdot\left(\vec{W}_\mu\times\vec{W}_\nu\right) - \frac{g'^2}{4}\left(\vec{W}_\mu\times\vec{W}_\nu\right)^2$$

$$\mathcal{L} = -\frac{1}{2}\left(\partial_\mu\rho\right)^2 - \frac{\lambda}{2}(\rho^2-\rho_0^2)^2 - \frac{1}{4}\bar{F}_{\mu\nu}^{\square} - \frac{g^2}{2}\rho^2\bar{A}_\mu^{\square}$$
$$-\frac{1}{4}G_{\mu\nu}'^{\square} - \frac{1}{2}\left|D_\mu'W_\nu - D_\nu'W_\mu\right|^2 - g'^2\rho^2W_\mu^*W_\mu$$
$$+ig'G_{\mu\nu}'W_\mu^*W_\nu + \frac{g'^2}{4}\left(W_\mu^*W_\nu - W_\nu^*W_\mu\right)^2$$
$$D_\mu' = \partial_\mu + ig'B_\mu', W_\mu = \frac{1}{\sqrt{2}}\left(W_\mu^1 + iW_\mu^2\right)$$

$$\partial_\mu^2\rho - g^2\rho\bar{A}_\mu^2 - 2g'^2\rho W_\mu^*W_\mu = 2\lambda(\rho^2-\rho_0^2)\rho$$
$$D_\mu'\left(D_\mu'W_\nu - D_\nu'W_\mu\right) = g'^2\rho^2W_\nu + igG_{\mu\nu}'W_\mu - g'^2\left(\left(W_\mu^*W_\nu - W_\nu^*W_\mu\right)W_\mu\right)$$
$$\partial_\mu\bar{F}_{\mu\nu} = J_\nu^{(e)}$$
$$\partial_\mu G_{\mu\nu}' = J_\nu^{(s)}$$

$$J_\nu^{(e)} = g^2\rho^2\bar{A}_\nu$$
$$J_\nu^{(s)} = ig'\left(W_\mu^*\left(D_\mu'W_\nu - D_\nu'W_\mu\right) - W_\mu\left(D_\mu'W_\nu - D_\nu'W_\mu\right)^*\right)$$

$$\vec{\Phi} = \rho(r)\exp(-im\varphi)\hat{n}, \hat{n} = \begin{pmatrix}\sin\alpha(r)\cos(n\varphi)\\ \sin\alpha(r)\sin(n\varphi)\\ \cos\alpha(r)\end{pmatrix},$$
$$A_\mu = \frac{m}{g}A(r)\partial_\mu\varphi$$
$$\hat{B}_\mu = \frac{n}{g'}B(r)\partial_\mu\varphi\hat{n} - \frac{1}{g'}\hat{n}\times\partial_\mu\hat{n}$$
$$\vec{W}_\mu = \frac{1}{g'}f(r)\hat{n}\times\partial_\mu\hat{n}$$

$$\rho = \rho(r), \bar{A}_\mu = \frac{m}{g}(A+1)\partial_\mu\varphi = \frac{m}{g}\bar{A}\partial_\mu\varphi,$$
$$B'_\mu = -\frac{n}{g'}(1-\cos\ \alpha - B)\partial_\mu\varphi = -\frac{n}{g'}B'\partial_\mu\varphi,$$
$$\vec{W_\mu} = \frac{f}{g'}\hat{n}\times\partial_\mu\hat{n}.$$

$$\ddot{\rho} + \frac{\dot{\rho}}{r} - \frac{m^2\bar{A}^2 + n^2 f^2 \sin^2\ \alpha}{r^2}\rho = 2\lambda(\rho^2 - \rho_0^2)\rho,$$
$$\ddot{\bar{A}} - \frac{\dot{\bar{A}}}{r} = g^2\rho^2\bar{A},$$
$$\ddot{B}' - \frac{\dot{B}'}{r} = 0,$$
$$\sin\ \alpha\left(\ddot{f} + \frac{\dot{f}}{r} - g'^2\rho^2 f\right) = 0,$$
$$n^2\sin\ \alpha\left(\dot{f}(B'-1) - f\dot{B}'\right) = 0.$$

$$\ddot{\rho} + \frac{\dot{\rho}}{r} - \frac{m^2\bar{A}^2}{r^2}\rho = 2\lambda(\rho^2 - \rho_0^2)\rho$$
$$\ddot{\bar{A}} - \frac{\dot{\bar{A}}}{r} - g^2\rho^2\bar{A} = 0$$
$$\ddot{B}' - \frac{\dot{B}'}{r} = 0$$

$$\rho(0) = 0, \rho(\infty) = \rho_0, \bar{A}(0) = 1, \bar{A}(\infty) = 0,$$

$$\bar{A}_\mu = \frac{m}{g}\bar{A}\partial_\mu\varphi$$
$$\Phi = \oint_{r=0}^{r=\infty} \bar{A}_\mu dx^\mu = -\frac{2\pi m}{g}$$

$$B'_\mu = -\frac{n}{g'}\partial_\mu\varphi$$
$$\Phi = \oint B'_\mu dx^\mu = -\frac{2\pi n}{g'}$$

$$\ddot{\rho} + \frac{\dot{\rho}}{r} - \frac{m^2\bar{A}^2 + n^2 f^2}{r^2}\rho = 2\lambda(\rho^2 - \rho_0^2)\rho,$$
$$\ddot{\bar{A}} - \frac{\dot{\bar{A}}}{r} = g^2\rho^2\bar{A},$$
$$\ddot{f} - \frac{\dot{f}}{r} = g'^2\rho^2 f.$$

$$\rho(0) = 0, \rho(\infty) = \rho_0, f(0) = 1, f(\infty) = 0,$$

$$\rho(0) = 0, \quad \bar{A}(0) = 1, f(0) = 1,$$
$$\rho(\infty) = \rho_0, \quad \bar{A}(\infty) = 0,$$

$$\vec{\Phi} = \rho(r)\hat{r}, \hat{r} = \begin{pmatrix} \sin\,\theta\cos\,\varphi \\ \sin\,\theta\sin\,\varphi \\ \cos\,\theta \end{pmatrix},$$
$$A_\mu = 0, B'_\mu = -\frac{1}{g}(1-\cos\,\theta)\partial_\mu\varphi,$$
$$\vec{W_\mu} = \frac{1}{g'}f(r)\hat{r}\times\partial_\mu\hat{r}$$

$$\ddot{\rho} + \frac{2}{r}\dot{\rho} - 2\frac{f^2}{r^2}\rho = 2\lambda(\rho^2-\rho_0^2)\rho,$$
$$\ddot{f} - \frac{f^2-1}{r^2}\dot{f} - g'^2\rho^2 f = 0.$$

$$\rho = \rho_0 = \sqrt{\mu^2/\lambda}, f = 0$$
$$B'_\mu = -\frac{1}{g'}(1-\cos\,\theta)\partial_\mu\varphi$$

$$\rho(0) = 0, \quad \rho(\infty) = \rho_0,$$
$$f(0) = 1, \quad f(\infty) = 0.$$

$$\rho = \frac{1}{\tanh(r)} - \frac{1}{r}, f = \frac{r}{\sinh(r)}.$$

$$\mathcal{L} = -\left|\mathcal{D}_\mu\phi\right|^2 - \frac{\lambda}{2}\left(|\phi|^2 - \frac{\mu^2}{\lambda}\right)^2 - \frac{1}{4}F^{\square}_{\mu\nu} - \frac{1}{4}\vec{G}^{\square}_{\mu\nu}$$
$$\mathcal{D}_\mu\phi = \left(D_\mu - i\frac{g}{2}A_\mu\right)\phi$$
$$D_\mu\phi = \left(\partial_\mu - i\frac{g'}{2}\vec{\sigma}\cdot\vec{B}_\mu\right)\phi$$

$$\phi = \frac{1}{\sqrt{2}}\rho\xi, \left(\xi^\dagger\xi = 1\right)$$

$$\mathcal{D}_\mu\xi = \left[\partial_\mu - i\frac{g}{2}A_\mu - i\frac{g'}{2}\left(B'_\mu\hat{n} + \vec{W_\mu}\right)\cdot\vec{\sigma}\right]\xi$$
$$\left|\mathcal{D}_\mu\xi\right|^2 = \frac{1}{8}\left(-gA_\mu + g'B'_\mu\right)^2 + \frac{g'^2}{4}\vec{W_\mu}^2$$

$$\mathcal{L} = -\frac{1}{2}\left(\partial_\mu\rho\right)^2 - \frac{\lambda}{8}(\rho^2-\rho_0^2)^2 - \frac{1}{4}F^{\square}_{\mu\nu} - \frac{1}{4}G^{\square}_{\mu\nu} - \frac{1}{2}\left|D'_\mu W_\nu - D'_\nu W_\mu\right|^2$$
$$-\frac{\rho^2}{8}\left(\left(-gA_\mu + g'B'_\mu\right)^2 + 2g'^2W^*_\mu W_\mu\right) + ig'G'_{\mu\nu}W^*_\mu W_\nu + \frac{g'^2}{4}\left(W^*_\mu W_\nu - W^*_\nu W_\mu\right)^2$$

$$D'_\mu = \partial_\mu + ig'B'_\mu$$

$$W_\mu = \frac{1}{\sqrt{2}}\left(W^1_\mu + iW^2_\mu\right)$$

$$\begin{pmatrix} \bar{A}_\mu \\ Z_\mu \end{pmatrix} = \frac{1}{\sqrt{g^2+g'^2}}\begin{pmatrix} g' & g \\ -g & g' \end{pmatrix}\begin{pmatrix} A_\mu \\ B'_\mu \end{pmatrix} = \begin{pmatrix} \cos\,\omega & \sin\,\omega \\ -\sin\,\omega & \cos\,\omega \end{pmatrix}\begin{pmatrix} A_\mu \\ B'_\mu \end{pmatrix}$$

$$\mathcal{L} = -\frac{1}{2}\left(\partial_\mu \rho\right)^2 - \frac{\lambda}{8}(\rho^2 - \rho_0^2)^2 - \frac{1}{4}\bar{F}_{\mu\nu}^{\square} - \frac{1}{4}Z_{\mu\nu}^{\square} - \frac{1}{2}\left|\left(\bar{D}_\mu + i\bar{e}\frac{g'}{g}Z_\mu\right)W_\nu - \left(\bar{D}_\nu + i\bar{e}\frac{g'}{g}Z_\nu\right)W_\mu\right|^2$$

$$-\frac{\rho^2}{4}\left(g'^2 W_\mu^* W_\mu + \frac{g^2 + g'^2}{2}Z_\mu^2\right) + i\bar{e}\left(\bar{F}_{\mu\nu} + \frac{g'}{g}Z_{\mu\nu}\right)W_\mu^* W_\nu$$

$$+\frac{g'^2}{4}\left(W_\mu^* W_\nu - W_\nu^* W_\mu\right)^2$$

$$\bar{F}_{\mu\nu} = \partial_\mu \bar{A}_\nu - \partial_\nu \bar{A}_\mu, Z_{\mu\nu} = \partial_\mu Z_\nu - \partial_\nu Z_\mu$$

$$\bar{D}_\mu = \partial_\mu + i\bar{e}\bar{A}_\mu$$

$$\bar{e} = \frac{gg'}{\sqrt{g^2 + g'^2}} = g'\sin\,\omega = g\cos\,\omega$$

$$M_H = \sqrt{\lambda}\rho_0$$

$$M_W = \frac{g'}{2}\rho_0, M_Z = \frac{\sqrt{g^2 + g'^2}}{2}\rho_0$$

$$\bar{e}\frac{g'}{g} = 2eg \otimes 4e, g' \setminus \sqrt{2 + 2\sqrt{17}}e\tan\,\omega = \frac{2\sqrt{2}}{\sqrt{1 + \sqrt{17}}}$$

$$\begin{pmatrix} \cos\,\omega & \sin\,\omega \\ -\sin\,\omega & \cos\,\omega \end{pmatrix} = \frac{1}{\sqrt{9 + \sqrt{17}}}\begin{pmatrix} \sqrt{1 + \sqrt{17}} & 2\sqrt{2} \\ -2\sqrt{2} & \sqrt{1 + \sqrt{17}} \end{pmatrix}$$

$$\bar{e} = \frac{4\sqrt{2}}{\sqrt{1 + \sqrt{17}}}e \simeq 2.5e$$

$$M_W = \frac{\sqrt{1 + \sqrt{17}}}{\sqrt{2}}e\rho_0$$

$$M_Z = \frac{\sqrt{9 + \sqrt{17}}}{\sqrt{2}}e\rho_0 \simeq 1{,}6M_W.$$

$$\mathcal{L} = -\frac{1}{2}\left(\partial_\mu \rho\right)^2 - \frac{\lambda}{8}(\rho^2 - \rho_0^2)^2 - \frac{1}{4}\bar{F}_{\mu\nu}^{\square} - \frac{1}{4}Z_{\mu\nu}^{\square} - \frac{M_Z^{\square}}{2\rho_0^{\square}}\rho^{\square}Z_\mu^{\square}$$

$$-\frac{1}{2}\left|\left(\bar{D}_\mu + 2ieZ_\mu\right)W_\nu - \left(\bar{D}_\nu + 2ieZ_\nu\right)W_\mu\right|^2 - \frac{M_W^{\square}}{\rho_0^{\square}}\rho^{\square}W_\mu^* W_\mu$$

$$+ i\left(\bar{e}\bar{F}_{\mu\nu} + 2eZ_{\mu\nu}\right)W_\mu^* W_\nu + \frac{M_W^{\square}}{\rho_0^{\square}}\left(W_\mu^* W_\nu - W_\nu^* W_\mu\right)^2$$

$$\partial^2\rho-\left(2\frac{M_W^2}{\rho_0^2}W_\mu^*W_\mu+\frac{M_Z^2}{\rho_0^2}Z_\mu^2\right)\rho$$

$$=\frac{\lambda}{2}(\rho^2-\rho_0^2)\rho\left(\bar{D}_\mu+2ieZ_\mu\right)\left[\left(\bar{D}_\mu+2ieZ_\mu\right)W_\nu-\left(\bar{D}_\nu+2ieZ_\nu\right)W_\mu\right]-\frac{M_W^2}{\rho_0^2}\rho^2W_\nu$$

$$=iW_\mu\left(\bar{e}\bar{F}_{\mu\nu}+2eZ_{\mu\nu}\right)+4\frac{M_W^2}{\rho_0^2}W_\mu\left(W_\mu^*W_\nu-W_\nu^*W_\mu\right)\partial_\mu\left[\bar{F}_{\mu\nu}-i\bar{e}\left(W_\mu^*W_\nu-W_\nu^*W_\mu\right)\right]$$

$$=4e\bar{e}W_\mu^*W_\mu Z_\nu+i\bar{e}\left[W_\mu^*\left(\bar{D}_\mu W_\nu-\bar{D}_\nu W_\mu\right)-W_\mu\left(\bar{D}_\mu W_\nu-\bar{D}_\nu W_\mu\right)^*\right]$$

$$-2e\bar{e}Z_\mu\left(W_\mu^*W_\nu+W_\nu^*W_\mu\right)\partial_\mu\left[Z_{\mu\nu}-2ie\left(W_\mu^*W_\nu-W_\nu^*W_\mu\right)\right]$$

$$=\left(\frac{M_Z^2}{\rho_0^2}\rho^2+8e^2W_\mu^*W_\mu\right)Z_\nu+2ie\left[W_\mu^*\left(\bar{D}_\mu W_\nu-\bar{D}_\nu W_\mu\right)-W_\mu\left(\bar{D}_\mu W_\nu-\bar{D}_\nu W_\mu\right)^*\right]$$

$$-4e^2Z_\mu\left(W_\mu^*W_\nu+W_\nu^*W_\mu\right)$$

$$\partial_\mu\bar{F}_{\mu\nu}=J_\nu^{(s)},\partial_\mu Z_{\mu\nu}=J_\nu^{(e)}$$

$$J_\nu^{(s)}=4e\bar{e}W_\mu^*W_\mu Z_\nu+i\bar{e}\left[\partial_\mu\left(W_\mu^*W_\nu-W_\nu^*W_\mu\right)+\left(W_\mu^*\left(\bar{D}_\mu W_\nu-\bar{D}_\nu W_\mu\right)-\left(\bar{D}_\mu W_\nu-\bar{D}_\nu W_\mu\right)^*W_\mu\right)\right]$$

$$-2e\bar{e}Z_\mu\left(W_\mu^*W_\nu+W_\nu^*W_\mu\right)$$

$$J_\nu^{(e)}=\left(\frac{M_Z^2}{\rho_0^2}\rho^2+8e^2W_\mu^*W_\mu\right)Z_\nu$$

$$+2ie\left[\partial_\mu\left(W_\mu^*W_\nu-W_\nu^*W_\mu\right)+\left(W_\mu^*\left(\bar{D}_\mu W_\nu-\bar{D}_\nu W_\mu\right)-W_\mu\left(\bar{D}_\mu W_\nu-\bar{D}_\nu W_\mu\right)^*\right)\right]$$

$$-4e^2Z_\mu\left(W_\mu^*W_\nu+W_\nu^*W_\mu\right)$$

$$j_\nu^{(e)}=-2e\frac{M_Z}{\rho_0}\rho^2Z_\nu$$

$$j_\nu^{(s)}=\frac{M_W}{\rho_0}\left(\frac{M_Z}{\rho_0}\rho^2+8eW_\mu^*W_\mu\right)Z_\nu$$

$$+2i\frac{M_W}{\rho_0}\left[\partial_\mu\left(W_\mu^*W_\nu-W_\nu^*W_\mu\right)+\left(W_\mu^*\left(\bar{D}_\mu W_\nu-\bar{D}_\nu W_\mu\right)-W_\mu\left(\bar{D}_\mu W_\nu-\bar{D}_\nu W_\mu\right)^*\right)\right]$$

$$-2e\frac{M_W}{\rho_0}Z_\mu\left(W_\mu^*W_\nu+W_\nu^*W_\mu\right)$$

$$\begin{pmatrix}J_\mu^{(s)}\\J_\mu^{(e)}\end{pmatrix}=\begin{pmatrix}M_W/M_Z & \sqrt{1-(M_W/M_Z)^2}\\-\sqrt{1-(M_W/M_Z)^2} & M_W/M_Z\end{pmatrix}\times\begin{pmatrix}j_\mu^{(e)}\\j_\mu^{(s)}\end{pmatrix}$$

$$\phi = \frac{1}{\sqrt{2}}\rho(r)\xi, \xi = \begin{pmatrix} -\sin\frac{\alpha(r)}{2}\exp(-in\varphi) \\ \cos\frac{\alpha(r)}{2} \end{pmatrix},$$

$$A_\mu = \frac{m}{g}A(r)\partial_\mu\varphi$$

$$\hat{B}_\mu = n\frac{1}{g'}(B(r) + 1 - \cos\alpha(r))\partial_\mu\varphi\hat{n} - \frac{1}{g'}\hat{n}\times\partial_\mu\hat{n},$$

$$\vec{W_\mu} = \frac{1}{g'}f(r)\hat{n}\times\partial_\mu\hat{n},$$

$$\hat{n} = -\xi^\dagger\vec{\sigma}\xi = \begin{pmatrix} \sin\alpha(r)\cos n\varphi \\ \sin\alpha(r)\sin n\varphi \\ \cos\alpha(r) \end{pmatrix},$$

$$\rho = \rho(r), \bar{A}_\mu = \bar{e}\left(\frac{m}{g^2}A + \frac{n}{g'^2}B\right)\partial_\mu\varphi$$

$$Z_\mu = -\frac{1}{\sqrt{g^2 + g'^2}}(mA - nB)\partial_\mu\varphi$$

$$\vec{W_\mu} = \frac{1}{g'}f\hat{n}\times\partial_\mu\hat{n}$$

$$\bar{A}_\mu = \frac{m}{\bar{e}}A\partial_\mu\varphi, Z_\mu = 0$$

$$\bar{A}_\mu = 0$$

$$Z_\mu = n\frac{\sqrt{g^2 + g'^2}}{g'^2}B\partial_\mu\varphi = \frac{n}{e}Z\partial_\mu\varphi$$

$$\ddot{\rho} + \frac{\dot{\rho}}{r} - \frac{n^2f^2\sin^2\alpha + (mA - nB)^2}{4r^2}\rho = \frac{\lambda}{2}\rho(\rho^2 - \rho_0^2),$$

$$nf\rho(mA + n)\sin\alpha = 0,$$

$$m\left(\ddot{A} - \frac{\dot{A}}{r}\right) = -\frac{g^2}{4}(mA - nB)\rho^2,$$

$$n\left(\ddot{B} - \frac{\dot{B}}{r}\right) = -\frac{g'^2}{4}(mA - nB)\rho^2,$$

$$\sin\alpha\left(\ddot{f} - \frac{\dot{f}}{r} - \frac{g'^2}{4}\rho^2 f\right) = 0,$$

$$n^2((B + 1)\dot{f} - f\dot{B})\sin\alpha = 0.$$

$$\ddot{\rho} + \frac{\dot{\rho}}{r} - n^2\frac{(g^2 + g'^2)^2}{g'^4}\frac{Z^2}{r^2}\rho = \frac{\lambda}{2}\rho(\rho^2 - \rho_0^2),$$

$$\ddot{Z} - \frac{\dot{Z}}{r} - \frac{g^2 + g'^2}{4}\rho^2 Z = 0.$$

$$\rho(0) = 0, \rho(\infty) = \rho_0, Z(0) = 1, Z(\infty) = 0$$

$$Z_\mu = -\frac{n}{e} Z \partial_\mu \varphi$$
$$\Phi = \oint_{r=0}^{r=\infty} Z_\mu dx^\mu = \frac{2\pi n}{e}$$

$$\ddot{\rho} + \frac{\dot{\rho}}{r} - \frac{n^2}{4}\frac{f^2}{r^2}\rho = \frac{\lambda}{2}(\rho^2 - \rho_0^2)\rho,$$
$$\ddot{f} - \frac{\dot{f}}{r} - \frac{g'^2}{4}\rho^2 f = 0.$$

$$\rho = \rho_0, f = 0, A = -\frac{n}{m} = 1,$$

$$\bar{A}_\mu = \frac{m}{\bar{e}}\partial_\mu \varphi = -\frac{n}{\bar{e}}\partial_\mu \varphi$$
$$\Phi = \oint \bar{A}_\mu dx^\mu = \frac{2\pi m}{\bar{e}}$$

$$\rho(0) = 0, \quad \rho(\infty) = \rho_0,$$
$$f(0) = 1, \quad f(\infty) = 0,$$

$$\rho = \rho(r), \xi = i\begin{pmatrix} \sin(\theta/2)e^{-i\varphi} \\ -\cos(\theta/2) \end{pmatrix},$$
$$A_\mu = -\frac{1}{g}(1 - \cos\theta)\partial_\mu \varphi,$$
$$\vec{B}_\mu = \frac{1}{g'}(f(r) - 1)\hat{r} \times \partial_\mu \hat{r},$$

$$\rho = \rho(r), \bar{A}_\mu = -\frac{1}{\bar{e}}(1 - \cos\theta)\partial_\mu \varphi$$
$$Z_\mu = 0, \vec{W}_\mu = \frac{1}{\bar{e}} f(r)\hat{r} \times \partial_\mu \hat{r}$$

$$\ddot{\rho} + \frac{2}{r}\dot{\rho} - \frac{f^2}{2r^2}\rho = \frac{\lambda}{2}(\rho^2 - \rho_0^2)\rho$$
$$\ddot{f} - \frac{f^2 - 1}{r^2} f - \frac{g'^2}{4}\rho^2 f = 0$$

$$\rho = \rho_0 = \sqrt{2\mu^2/\lambda}, f = 0$$
$$\bar{A}_\mu = -\frac{1}{\bar{e}}(1 - \cos\theta)\partial_\mu \varphi$$

$$\rho(0) = 0, \quad \rho(\infty) = \rho_0,$$
$$f(0) = 1, \quad f(\infty) = 0.$$

$$g = 2e, g' = \frac{\sqrt{1+\sqrt{17}}}{\sqrt{2}}e$$

$$\tan\,\omega = \frac{2\sqrt{2}}{\sqrt{1+\sqrt{17}}}$$

$$M_W = \frac{\sqrt{1+\sqrt{17}}}{2\sqrt{2}}e\rho_0$$

$$M_Z = \frac{\sqrt{9+\sqrt{17}}}{2\sqrt{2}}e\rho_0 \simeq 1.6M_W$$

$$\mathcal{L} = \bar{\psi}\left[i\gamma^\mu\partial_\mu - g(\sigma + i\vec{\tau}\cdot\vec{\pi}\gamma_5)\right]\psi + \frac{1}{2}\partial_\mu\sigma\partial^\mu\sigma + \frac{1}{2}\partial_\mu\vec{\pi}\cdot\partial^\mu\vec{\pi} - U(\sigma,\vec{\pi})$$

$$U(\sigma,\vec{\pi}) = \frac{\lambda}{4}(\sigma^2 + \vec{\pi}^2 - \nu^2)^2 - H\sigma - \frac{m_\pi^4}{4\lambda} + f_\pi^2 m_\pi^2$$

$$\partial_\mu\frac{\partial\mathcal{L}}{\partial\partial_\mu\phi} - \frac{\partial\mathcal{L}}{\partial\phi} = 0,$$

$$\sigma(\vec{r},t) = \sigma(\vec{r}), \vec{\pi}(\vec{r},t) = \vec{\pi}(\vec{r}), \psi(\vec{r},t) = e^{-i\varepsilon t}\sum_{i=1}^{N}\Psi_i(\vec{r}),$$

$$\begin{pmatrix} g\sigma & -i\vec{\sigma}\cdot\vec{\nabla} + ig\vec{\tau}\cdot\vec{\pi} \\ -i\vec{\sigma}\cdot\vec{\nabla} - ig\vec{\tau}\cdot\vec{\pi} & -g\sigma \end{pmatrix}\Psi = \varepsilon\Psi,$$

$$-\vec{\nabla}^2\sigma + \frac{\partial U(\sigma,\vec{\pi})}{\partial\sigma} + gN\bar{\Psi}\Psi = 0,$$

$$-\vec{\nabla}^2\vec{\pi} + \frac{\partial U(\sigma,\vec{\pi})}{\partial\vec{\pi}} + gN\bar{\Psi}i\gamma_5\vec{\tau}\Psi = 0.$$

$$\sigma(\vec{r}) = \sigma(r), \vec{\pi}(\vec{r}) = \hat{\vec{r}}\pi(r),$$

$$\Psi_{G,\Pi=+}(\vec{r}) = \begin{pmatrix} ig_1(r)y_{G+\frac{1}{2},G}(\hat{\vec{r}}) \\ f_1(r)y_{G+\frac{1}{2},G+1}(\hat{\vec{r}}) \end{pmatrix} + \begin{pmatrix} ig_2(r)y_{G-\frac{1}{2},G}(\hat{\vec{r}}) \\ -f_2(r)y_{G-\frac{1}{2},G-1}(\hat{\vec{r}}) \end{pmatrix},$$

$$\Psi_{G,\Pi=-}(\vec{r}) = \begin{pmatrix} ig_1(r)y_{G+\frac{1}{2},G+1}(\hat{\vec{r}}) \\ -f_1(r)y_{G+\frac{1}{2},G}(\hat{\vec{r}}) \end{pmatrix} + \begin{pmatrix} ig_2(r)y_{G-\frac{1}{2},G-1}(\hat{\vec{r}}) \\ f_2(r)y_{G-\frac{1}{2},G}(\hat{\vec{r}}) \end{pmatrix}$$

$$\begin{cases} f_1' + \frac{G+2}{r}f_1 - (\varepsilon - g\sigma)g_1 + \frac{g\pi}{2G+1}\left[f_1 + 2\sqrt{G(G+1)}f_2\right] = 0 \\ f_2' - \frac{G-1}{r}f_2 + (\varepsilon - g\sigma)g_2 - \frac{g\pi}{2G+1}\left[f_2 - 2\sqrt{G(G+1)}f_1\right] = 0 \\ g_1' - \frac{G}{r}g_1 + (\varepsilon + g\sigma)f_1 - \frac{g\pi}{2G+1}\left[g_1 - 2\sqrt{G(G+1)}g_2\right] = 0 \\ g_2' + \frac{G+1}{r}g_2 - (\varepsilon + g\sigma)f_2 + \frac{g\pi}{2G+1}\left[g_2 + 2\sqrt{G(G+1)}g_1\right] = 0 \end{cases}$$

$$\begin{cases} g_1' + \dfrac{G+2}{r} g_1 - (\varepsilon + g\sigma) f_1 - \dfrac{g\pi}{2G+1}\left[g_1 - 2\sqrt{G(G+1)}g_2\right] = 0 \\ g_2' - \dfrac{G-1}{r} g_2 + (\varepsilon + g\sigma) f_2 + \dfrac{g\pi}{2G+1}\left[g_2 + 2\sqrt{G(G+1)}g_1\right] = 0 \\ f_1' - \dfrac{G}{r} f_1 + (\varepsilon - g\sigma) g_1 + \dfrac{g\pi}{2G+1}\left[f_1 + 2\sqrt{G(G+1)}f_2\right] = 0 \\ f_2' + \dfrac{G+1}{r} f_2 - (\varepsilon - g\sigma) g_2 - \dfrac{g\pi}{2G+1}\left[f_2 - 2\sqrt{G(G+1)}f_1\right] = 0 \end{cases}$$

$$\Psi_{G=0,\Pi=+}(\vec{r}) = \begin{pmatrix} i g_1(r) y_{\frac{1}{2},0}(\hat{\vec{r}}) \\ f_1(r) y_{\frac{1}{2},1}(\hat{\vec{r}}) \end{pmatrix}, \Psi_{G=0,\Pi=-}(\vec{r}) = \begin{pmatrix} i g_1(r) y_{\frac{1}{2},1}(\hat{\vec{r}}) \\ -f_1(r) y_{\frac{1}{2},0}(\hat{\vec{r}}) \end{pmatrix}.$$

$$\begin{cases} f_1' + \dfrac{2}{r} f_1 - (\varepsilon - g\sigma) g_1 + g\pi f_1 = 0 \\ g_1' + (\varepsilon + g\sigma) f_1 - g\pi g_1 = 0 \end{cases}$$

$$\begin{cases} g_1' + \dfrac{2}{r} g_1 - (\varepsilon + g\sigma) f_1 - g\pi g_1 = 0 \\ f_1' + (\varepsilon - g\sigma) g_1 + g\pi f_1 = 0 \end{cases}$$

$$\begin{aligned} \frac{\mathrm{d}f_1(r)}{\mathrm{d}r} &= -\frac{2}{r} f_1(r) + (\varepsilon - g\sigma(r)) g_1(r) - g\pi(r) f_1(r), \\ \frac{\mathrm{d}g_1(r)}{\mathrm{d}r} &= -(\varepsilon + g\sigma(r)) f_1(r) + g\pi(r) g_1(r), \\ \frac{\partial U}{\partial \sigma} &= \frac{\mathrm{d}^{\square}\sigma(r)}{\mathrm{d}r^2} + \frac{2}{r}\frac{\mathrm{d}\sigma(r)}{\mathrm{d}r} - Ng(g_1^2(r) - f_1^2(r)), \\ \frac{\partial U}{\partial \pi} &= \frac{\mathrm{d}^{\square}\pi(r)}{\mathrm{d}r^2} + \frac{2}{r}\frac{\mathrm{d}\pi(r)}{\mathrm{d}r} - \frac{2\pi(r)}{r^2} - 2Ngf_1(r)g_1(r). \end{aligned}$$

$$\begin{gathered} f_1(0) = 0, \frac{\mathrm{d}\sigma(0)}{\mathrm{d}r} = 0, \pi(0) = 0 \\ g_1(\infty) = 0, \sigma(\infty) = f_\pi, \pi(\infty) = 0 \end{gathered}$$

$$4\pi \int r^2 (f_1^2(r) + g_1^2(r)) \mathrm{d}r = 1$$

$$E_{\mathrm{cl}} = 3\varepsilon + 4\pi \int_0^\infty \mathrm{d}r r^2 \left[\frac{1}{2}\left(\frac{\mathrm{d}\sigma}{\mathrm{d}r}\right)^2 + \frac{1}{2}\left(\frac{\mathrm{d}\pi}{\mathrm{d}r}\right)^2 + \frac{\pi^2}{r^2} + U(\sigma,\pi)\right],$$

$$\sigma(r) = \sigma_v + \tilde{\sigma}(r), \pi(r) = \pi_v + \tilde{\pi}(r)$$

$$S_\psi[\tilde{\sigma}(r), \tilde{\pi}(r)] = \int \mathrm{d}^{\square}x \left[\bar{\psi}\left(i\gamma_\mu \partial^\mu - m_q - g\left(\tilde{\sigma} + i\gamma_5 \vec{\tau}\cdot\hat{\vec{r}}\tilde{\pi}\right)\right)\psi + \mathcal{L}_{\mathrm{ct}}\right]$$

$$\mathcal{L}_{\mathrm{ct}} = 2a\left(\left(\partial_\mu \sigma\right)^2 + \left(\partial_\mu \pi\right)^2\right) - 2b(\sigma^2 + \vec{\pi}^2 - \sigma_v^2) - 4c(\sigma^2 + \vec{\pi}^2 - \sigma_v^2)^2.$$

$$a= -\frac{g^2}{(4\pi)^2}\left\{\frac{1}{\epsilon}-\gamma-\frac{2}{3}+\ln\left(\frac{4\pi\mu^2}{m_q^2}\right)-6\int_0^1 dx x(1-x)\ln\left[1-x(1-x)\frac{m_s^2}{m_q^2}\right]\right\}$$

$$b=-\frac{g^2 m_q^2}{(4\pi)^2}\left\{\frac{1}{\epsilon}-\gamma+1+\ln\left(\frac{4\pi\mu^2}{m_q^2}\right)\right\}$$

$$c=-\frac{g^4}{(4\pi)^2}\left\{\frac{1}{\epsilon}-\gamma+\ln\left(\frac{4\pi\mu^2}{m_q^2}\right)-\frac{m_s^2}{4m_q^2}-\frac{3}{2}\int_0^1 dx\ln\left[1-x(1-x)\frac{m_s^2}{m_q^2}\right]\right\}$$

$$e^{iS_{\text{eff}}[\tilde{\sigma},\tilde{\pi}]}=\frac{\int [D\psi]\left[iD\psi^\dagger\right]e^{iS_\psi[\tilde{\sigma},\tilde{\pi}]}}{\int [D\psi][iD\psi^\dagger]e^{iS_\psi[\tilde{\sigma}]_{\tilde{\sigma}=\tilde{\pi}=0}}}$$

$$S_{\text{eff}}[\tilde{\sigma},\tilde{\pi}]=\text{Trlog}\,\frac{h_D[\tilde{\sigma},\tilde{\pi}]}{h_D[\tilde{\sigma}]|_{\tilde{\sigma}=\tilde{\pi}=0}}+S_{\text{ct}}$$

$$h_D=i\gamma_\mu\partial^\mu-m_q-g(\tilde{\sigma}+i\vec{\tau}\cdot\hat{\vec{r}}\tilde{\pi})$$

$$E_{\text{vac}}=-S_{\text{eff}}\int \text{d}t$$

$$E_{\text{vac}}[\tilde{\sigma},\tilde{\pi}]=E_{\text{vac}}^{\psi}[\tilde{\sigma},\tilde{\pi}]+E_{\text{ct}}[\tilde{\sigma},\tilde{\pi}]$$

$$E_{\text{vac}}^{\psi}[\tilde{\sigma},\tilde{\pi}]=-\frac{1}{2}\left[\sum_n (2G_n+1)|E_n|+\sum_{\varepsilon_\eta}\left|E_\eta(k)\right|-\sum_{\varepsilon_k}\left|E_q(k)\right|\right],$$

$$E_q(k)=\omega\sqrt{k^2+m_q^2}$$

$$H_{\text{eff}}(\tilde{\sigma},\tilde{\pi})\psi_\alpha=\left[-i\gamma_0\vec{\gamma}\cdot\nabla+\gamma_0 m_q+\gamma_0 g\left(\tilde{\sigma}+i\gamma_5\vec{\tau}\cdot\hat{\vec{r}}\tilde{\pi}\right)\right]\psi_\alpha=E_\alpha\psi_\alpha$$

$$\sum_{\varepsilon_k}\left|E_q(k)\right|=\sum_{G,\omega,\Pi}(2G+1)\int \text{d}k\rho_{G,\omega,\Pi}^{\text{free}}(k)\left|E_q(k)\right|$$

$$\sum_{\varepsilon_\eta}\left|E_\eta(k)\right|=\sum_{G,\omega,\Pi}(2G+1)\int \text{d}k\rho_{G,\omega,\Pi}(k)\left|E_q(k)\right|$$

$$\rho_{G,\omega,\Pi}(k)-\rho_{G,\omega,\Pi}^{\text{free}}(k)=\frac{1}{\pi}\frac{\text{d}\delta_{G,\omega,\Pi}(k)}{\text{d}k}$$

$$E_{\text{vac}}^{\psi}[\tilde{\sigma},\tilde{\pi}]=-\frac{1}{2}\sum_n (2G_n+1)|E_n|-\frac{1}{2}\sum_{G,\omega,\Pi}(2G+1)\int \text{d}k\frac{1}{\pi}\frac{\text{d}\delta_{G,\omega,\Pi}(k)}{\text{d}k}\left|E_q(k)\right|.$$

$$\bar{\delta}_{G,\omega,\Pi}(k)\equiv\delta_{G,\omega,\Pi}(k)-\sum_{\ell=1}^{n}\delta_{G,\omega,\Pi}^{(\ell)}(k)$$

$$E_{\text{vac}}^{\text{ren}}[\tilde{\sigma},\tilde{\pi}]=-\frac{1}{2}\sum_n (2G_n+1)|E_n|-\frac{1}{2}\sum_{G,\omega,\Pi}(2G+1)\int \text{d}k\frac{1}{\pi}\frac{\text{d}\bar{\delta}_{G,\omega,\Pi}(k)}{\text{d}k}\left|E_q(k)\right|+\Gamma_2+\Gamma_4$$

$$\Gamma_2=\frac{g^2}{\pi^2}\int_0^\infty dqq^2[s^2(q)+p^2(q)]\left\{q^2+m_s^2-6\int_0^1 dx\left[m_q^2+x(1-x)q^2\right]\ln\frac{m_q^2+x(1-x)q^2}{m_q^2-x(1-x)m_s^2}\right\}$$
$$-\frac{g^2}{\pi^2}\int_0^\infty dqq^2p^2(q)\left\{m_s^2+2m_q^2\int_0^1 dx\left[3\ln\left(1-x(1-x)\frac{m_s^2}{m_q^2}\right)-2\ln\left(1+x(1-x)\frac{q^2}{m_q^2}\right)\right]\right\}$$

$$s(q)=\int_0^\infty drr^2\frac{\sin\ qr}{qr}\tilde{\sigma}(r),p(q)=\int_0^\infty drr^2\left[\frac{\cos\ qr}{qr}-\frac{\sin\ qr}{(qr)^2}\right]\tilde{\pi}(r)$$

$$\Gamma_4=-c_B\int_0^\infty\frac{dqq^2}{2\pi^2}\tilde{V}_B(q)\tilde{V}_B(-q)\left[\int_0^1 dx\frac{x(1-x)}{m_B^2-x(1-x)m_f^2}\left(q^2+m_f^2\right)+\int_0^1 dx\ln\frac{m_B^2+x(1-x)q^2}{m_B^2-x(1-x)m_f^2}\right]$$

$$\tilde{V}_B(q)=\int_0^\infty drr^2\frac{\sin\ qr}{qr}V_B(r)$$

$$\begin{cases}g_1'+\dfrac{2}{r}g_1-(\varepsilon+g\sigma)f_1-g\pi g_1=0\\ f_1'+(\varepsilon-g\sigma)g_1+g\pi f_1=0\end{cases}$$

$$\begin{cases}g_1'+\dfrac{2}{r}g_1-(\varepsilon+g\sigma_v)f_1=0\\ f_1'+(\varepsilon-g\sigma_v)g_1=0\end{cases}$$

$$f_1(r)=u(r)h_0(kr),u(r)\to k/(\varepsilon+g\sigma_v)(r\to\infty),$$
$$g_1(r)=v(r)h_1(kr),v(r)\to 1(r\to\infty),$$

$$k=\sqrt{\varepsilon^2-g^2\sigma_v^2}$$

$$\begin{cases}v'h_1+v\left(h_1'+\dfrac{2}{r}h_1-g\pi h_1\right)-u(g\sigma+\varepsilon)h_0=0\\ u'h_0+u(h_0'+g\pi h_0)-v(g\sigma-\varepsilon)h_1=0\end{cases}$$

$$\delta_0(k)=\frac{1}{2i}\lim_{r\to 0}\ln\left(u^*(r)u^{-1}(r)\right)=\frac{1}{2i}\lim_{r\to 0}\ln\left(v^*(r)v^{-1}(r)\right)$$

$$\mathcal{G}(r)\equiv\begin{pmatrix}g_1^{(1)}(r) & g_1^{(2)}(r)\\ g_2^{(1)}(r) & g_2^{(2)}(r)\end{pmatrix}=H_v(kr)V(r),\mathcal{F}(r)\equiv\begin{pmatrix}f_1^{(1)}(r) & f_1^{(2)}(r)\\ f_2^{(1)}(r) & f_2^{(2)}(r)\end{pmatrix}=H_u(kr)U(r),$$

$$V(r)\equiv\begin{pmatrix}v_1^{(1)}(r) & v_1^{(2)}(r)\\ v_2^{(1)}(r) & v_2^{(2)}(r)\end{pmatrix},H_v(kr)\equiv\begin{pmatrix}h_{G+1}(kr) & 0\\ 0 & h_{G-1}(kr)\end{pmatrix},$$
$$U(r)\equiv\begin{pmatrix}u_1^{(1)}(r) & u_1^{(2)}(r)\\ u_2^{(1)}(r) & u_2^{(2)}(r)\end{pmatrix},H_u(kr)\equiv\begin{pmatrix}h_G(kr) & 0\\ 0 & h_G(kr)\end{pmatrix},$$

$$\begin{cases} v_1^{(1)\prime}+v_1^{(1)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(1)}-(\varepsilon+g\sigma)\dfrac{h_G}{h_{G+1}}u_1^{(1)}-\dfrac{g\pi}{1+2G}\left(v_1^{(1)}-2\sqrt{G(1+G)}\dfrac{h_{G-1}}{h_{G+1}}v_2^{(1)}\right)=0 \\ v_2^{(1)\prime}+v_2^{(1)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(1)}+(\varepsilon+g\sigma)\dfrac{h_G}{h_{G-1}}u_2^{(1)}+\dfrac{g\pi}{1+2G}\left(v_2^{(1)}+2\sqrt{G(1+G)}\dfrac{h_{G+1}}{h_{G-1}}v_1^{(1)}\right)=0 \\ u_1^{(1)\prime}+u_1^{(1)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(1)}+(\varepsilon-g\sigma)\dfrac{h_{G+1}}{h_G}v_1^{(1)}+\dfrac{g\pi}{1+2G}\left(u_1^{(1)}+2\sqrt{G(1+G)}u_2^{(1)}\right)=0 \\ u_2^{(1)\prime}+u_2^{(1)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(1)}-(\varepsilon-g\sigma)\dfrac{h_{G-1}}{h_G}v_2^{(1)}-\dfrac{g\pi}{1+2G}\left(u_2^{(1)}-2\sqrt{G(1+G)}u_1^{(1)}\right)=0 \end{cases}$$

$$\begin{cases} v_1^{(2)\prime}+v_1^{(2)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(2)}-(\varepsilon+g\sigma)\dfrac{h_G}{h_{G+1}}u_1^{(2)}-\dfrac{g\pi}{1+2G}\left(v_1^{(2)}-2\sqrt{G(1+G)}\dfrac{h_{G-1}}{h_{G+1}}v_2^{(2)}\right)=0 \\ v_2^{(2)\prime}+v_2^{(2)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(2)}+(\varepsilon+g\sigma)\dfrac{h_G}{h_{G-1}}u_2^{(2)}+\dfrac{g\pi}{1+2G}\left(v_2^{(2)}+2\sqrt{G(1+G)}\dfrac{h_{G+1}}{h_{G-1}}v_1^{(2)}\right)=0 \\ u_1^{(2)\prime}+u_1^{(2)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(2)}+(\varepsilon-g\sigma)\dfrac{h_{G+1}}{h_G}v_1^{(2)}+\dfrac{g\pi}{1+2G}\left(u_1^{(2)}+2\sqrt{G(1+G)}u_2^{(2)}\right)=0 \\ u_2^{(2)\prime}+u_2^{(2)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(2)}-(\varepsilon-g\sigma)\dfrac{h_{G-1}}{h_G}v_2^{(2)}-\dfrac{g\pi}{1+2G}\left(u_2^{(2)}-2\sqrt{G(1+G)}u_1^{(2)}\right)=0 \end{cases}$$

$$V(r)\to\begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix}, U(r)\to\begin{pmatrix}k/(\varepsilon+g\sigma_v) & 0\\ 0 & k/(\varepsilon+g\sigma_v)\end{pmatrix}$$

$$\Psi_{SC}(r)=-\mathcal{G}(r)^*+\mathcal{G}(r)S$$

$$S=\lim_{r\to 0}H_v^{-1}(kr)V^{-1}(r)V^*(r)H_v^*(kr)$$

$$\delta(k)=\frac{1}{2i}\mathrm{trln}\ S=\frac{1}{2i}\lim_{r\to 0}\mathrm{trln}\,(V^{-1}(r)V^*(r)).$$

$$\delta(k)=\frac{1}{2i}\lim_{r\to 0}\mathrm{trln}\,(U^{-1}(r)U^*(r))$$

$$\begin{aligned}V(r)=&1+g_sV^{(1,0)}(r)+g_pV^{(0,1)}(r)+g_s^2V^{(2,0)}(r)+g_p^2V^{(0,2)}(r)\\&+g_sg_pV^{(1,1)}(r)+\cdots,\end{aligned}$$

$$\sigma(r)=\sigma_v+g_s\tilde{\sigma}(r),\pi(r)=\pi_v+g_p\tilde{\pi}(r)$$

$$E_{G,\omega,\Pi}=(2G+1)\int_0^\infty\frac{\mathrm{d}k}{\pi}\frac{d\bar{\delta}_{G,\omega,\Pi}(k)}{dk}\left|E_q(k)\right|$$

$$E_{\mathrm{vac}}^{\mathrm{ren}}[\tilde{\sigma},\tilde{\pi}]=-\frac{1}{2}\sum_n(2G_n+1)|E_n|-\frac{1}{2}\sum_{G,\omega,\Pi}E_{G,\omega,\Pi}+\Gamma_2+\Gamma_4,$$

$$V_B(r)=m_B^2\frac{r}{\mathrm{w}}e^{-3\frac{r}{\mathrm{w}}},$$

$$-\frac{d^\square}{dr^\square}u_\ell(r)+\left[\frac{\ell(\ell+1)}{r^2}+V_B(r)\right]u_\ell(r)=k^\square u_\ell(r).$$

$$u_\ell(r)=\exp(2i\beta_\ell(k,r))rh_\ell(kr)$$

$$\beta_\ell(k,r) = g_B\beta_\ell^{(1)}(k,r) + g_B^2\beta_\ell^{(2)}(k,r)$$

$$-i\beta_\ell^{(1)\prime\prime} - 2ikp_\ell(kr)\beta_\ell^{(1)\prime} + \frac{1}{2}V_B(r) = 0$$
$$-i\beta_\ell^{(2)\prime\prime} - 2ikp_\ell(kr)\beta_\ell^{(2)\prime} + 2\left(\beta_\ell^{(1)\prime}\right)^2 = 0$$

$$\delta_{\ell,B}^{(2)}(k) = 2\mathrm{Re}\left[\beta_\ell^{(2)}(k,0)\right].$$

$$E_B^{(2)}(q) = \int_0^q \frac{d^\square k}{\pi}\frac{d\delta_B^{(2)}(k)}{dk}\sqrt{k^2+m_B^2}$$
$$E_F^{(3,4)}(q) = \int_0^q \frac{d^\square k}{\pi}\frac{d\delta_F^{(3,4)}(k)}{dk}\sqrt{k^2+m_q^2}$$

$$\delta_B^{(2)}(k) = 2\sum_\ell (2\ell+1)\delta_{\ell,B}^{(2)}(k)$$
$$\delta_F^{(3,4)}(k) = \sum_{G,\omega,\Pi}(2G+1)\left(\delta_{G,\omega,\Pi}^{(3)}(k) + \delta_{G,\omega,\Pi}^{(4)}(k)\right)$$

$$\Pi_{ren}(q) = -\int_0^1 dx\frac{x(1-x)}{m_B^2 - x(1-x)m_f^2}\left(q^2+m_f^2\right) - \int_0^1 dx\ln\frac{m_B^2 + x(1-x)q^2}{m_B^2 - x(1-x)m_f^2}$$

$$\Gamma_{2B} = \int_0^\infty \frac{d^\square q q^2}{2\pi^2}\tilde{V}_B(q)\tilde{V}_B(-q)\Pi_{ren}(q)$$

$$c_B = \frac{\Gamma_4}{\Gamma_{2B}}, c_B \equiv \lim_{q\to\infty} C_B(q) = \frac{E_F^{(3,4)}\Big|_{q\to\infty}}{E_B^{(2)}\Big|_{q\to\infty}}$$

$$\Gamma_{\mathrm{loc}}^{(4)} = \frac{g^4}{8\pi}\left(\frac{m_s^2}{m_q^2} + 6\int_0^1 dx\ln\left[1 - x(1-x)\frac{m_s^2}{m_q^2}\right]\right) \times \int_0^\infty drr^2[(\sigma(r)^2 + \pi(r)^2 - \sigma_v^2)^2 - 4\sigma_v^2(\sigma(r)-\sigma_v)^2]$$

$$\Gamma_{4,\mathrm{loc}} = \frac{c_B}{4\pi}\left(-\int_0^1 dx\frac{x(1-x)m_f^2}{m_B^2 - x(1-x)m_f^2} + \int_0^1 dx\ln\left[1 - x(1-x)\frac{m_f^2}{m_B^2}\right]\right)\int_0^\infty drr^2V_B(r)^2$$

$$\begin{cases} v_1^{(1)(1,0)\prime} + v_1^{(1)(1,0)}\dfrac{h_{G+1}'}{h_{G+1}} + \dfrac{G+2}{r}v_1^{(1)(1,0)} - (\varepsilon - g\sigma_v)\dfrac{h_G}{h_{G+1}}u_1^{(1)(1,0)} + \beta\sigma_s = 0; \\ v_2^{(1)(1,0)\prime} + v_2^{(1)(1,0)}\dfrac{h_{G-1}'}{h_{G-1}} - \dfrac{G-1}{r}v_2^{(1)(1,0)} + (\varepsilon - g\sigma_v)\dfrac{h_G}{h_{G-1}}u_2^{(1)(1,0)} = 0; \\ u_1^{(1)(1,0)\prime} + u_1^{(1)(1,0)}\dfrac{h_G'}{h_G} - \dfrac{G}{r}u_1^{(1)(1,0)} + (\varepsilon + g\sigma_v)\dfrac{h_{G+1}}{h_G}v_1^{(1)(1,0)} + \sigma_s = 0; \\ u_2^{(1)(1,0)\prime} + u_2^{(1)(1,0)}\dfrac{h_G'}{h_G} + \dfrac{G+1}{r}u_2^{(1)(1,0)} - (\varepsilon + g\sigma_v)\dfrac{h_{G-1}}{h_G}v_2^{(1)(1,0)} = 0; \end{cases}$$

$$\begin{cases} v_1^{(2)(1,0)\prime}+v_1^{(2)(1,0)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(2)(1,0)}-(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G+1}}u_1^{(2)(1,0)}=0; \\ v_2^{(2)(1,0)\prime}+v_2^{(2)(1,0)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(2)(1,0)}+(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G-1}}u_2^{(2)(1,0)}-\beta\sigma_s=0; \\ u_1^{(2)(1,0)\prime}+u_1^{(2)(1,0)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(2)(1,0)}+(\varepsilon+g\sigma_v)\dfrac{h_{G+1}}{h_G}v_1^{(2)(1,0)}=0; \\ u_2^{(2)(1,0)\prime}+u_2^{(2)(1,0)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(2)(1,0)}-(\varepsilon+g\sigma_v)\dfrac{h_{G-1}}{h_G}v_2^{(2)(1,0)}-\sigma_s=0; \end{cases}$$

$$\begin{cases} v_1^{(1)(2,0)\prime}+v_1^{(1)(2,0)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(1)(2,0)}-(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G+1}}u_1^{(1)(2,0)}+\sigma_s u_1^{(1)(1,0)}=0 \\ v_2^{(1)(2,0)\prime}+v_2^{(1)(2,0)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(1)(2,0)}+(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G-1}}u_2^{(1)(2,0)}-\sigma_s u_2^{(1)(1,0)}=0 \\ u_1^{(1)(2,0)\prime}+u_1^{(1)(2,0)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(1)(2,0)}+(\varepsilon+g\sigma_v)\dfrac{h_{G+1}}{h_G}v_1^{(1)(2,0)}+\sigma_s v_1^{(1)(1,0)}=0 \\ u_2^{(1)(2,0)\prime}+u_2^{(1)(2,0)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(1)(2,0)}-(\varepsilon+g\sigma_v)\dfrac{h_{G-1}}{h_G}v_2^{(1)(2,0)}-\sigma_s v_2^{(1)(1,0)}=0 \end{cases}$$

$$\begin{cases} v_1^{(2)(2,0)\prime}+v_1^{(2)(2,0)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(2)(2,0)}-(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G+1}}u_1^{(2)(2,0)}+\sigma_s u_1^{(2)(1,0)}=0 \\ v_2^{(2)(2,0)\prime}+v_2^{(2)(2,0)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(2)(2,0)}+(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G-1}}u_2^{(2)(2,0)}-\sigma_s u_2^{(2)(1,0)}=0 \\ u_1^{(2)(2,0)\prime}+u_1^{(2)(2,0)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(2)(2,0)}+(\varepsilon+g\sigma_v)\dfrac{h_{G+1}}{h_G}v_1^{(2)(2,0)}+\sigma_s v_1^{(2)(1,0)}=0 \\ u_2^{(2)(2,0)\prime}+u_2^{(2)(2,0)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(2)(2,0)}-(\varepsilon+g\sigma_v)\dfrac{h_{G-1}}{h_G}v_2^{(2)(2,0)}-\sigma_s v_2^{(2)(1,0)}=0 \end{cases}$$

$$\begin{cases} v_1^{(1)(0,1)\prime}+v_1^{(1)(0,1)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(1)(0,1)}-(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G+1}}u_1^{(1)(0,1)}+\dfrac{\pi}{1+2G}=0 \\ v_2^{(1)(0,1)\prime}+v_2^{(1)(0,1)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(1)(0,1)}+(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G-1}}u_2^{(1)(0,1)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}=0 \\ u_1^{(1)(0,1)\prime}+u_1^{(1)(0,1)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(1)(0,1)}+(\varepsilon+g\sigma_v)\dfrac{h_{G+1}}{h_G}v_1^{(1)(0,1)}+\dfrac{\beta\pi}{1+2G}=0 \\ u_2^{(1)(0,1)\prime}+u_2^{(1)(0,1)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(1)(0,1)}-(\varepsilon+g\sigma_v)\dfrac{h_{G-1}}{h_G}v_2^{(1)(0,1)}+\dfrac{2\sqrt{G(1+G)}\beta\pi}{1+2G}=0 \end{cases}$$

$$\begin{cases} v_1^{(2)(0,1)\prime}+v_1^{(2)(0,1)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(2)(0,1)}-(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G+1}}u_1^{(2)(0,1)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}=0 \\ v_2^{(2)(0,1)\prime}+v_2^{(2)(0,1)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(2)(0,1)}+(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G-1}}u_2^{(2)(0,1)}+\dfrac{\pi}{1+2G}=0 \\ u_1^{(2)(0,1)\prime}+u_1^{(2)(0,1)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(2)(0,1)}+(\varepsilon+g\sigma_v)\dfrac{h_{G+1}}{h_G}v_1^{(2)(0,1)}+\dfrac{2\sqrt{G(1+G)}\beta\pi}{1+2G}=0 \\ u_2^{(2)(0,1)\prime}+u_2^{(2)(0,1)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(2)(0,1)}-(\varepsilon+g\sigma_v)\dfrac{h_{G-1}}{h_G}v_2^{(2)(0,1)}-\dfrac{\beta\pi}{1+2G}=0 \end{cases}$$

$$\begin{cases} v_1^{(1)(0,2)\prime}+v_1^{(1)(0,2)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(1)(0,2)}-(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G+1}}u_1^{(1)(0,2)}+\dfrac{\pi}{1+2G}v_1^{(1)(0,1)}-\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}v_2^{(1)(0,1)}=0;\\ v_2^{(1)(0,2)\prime}+v_2^{(1)(0,2)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(1)(0,2)}+(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G-1}}u_2^{(1)(0,2)}+\dfrac{\pi}{1+2G}v_2^{(1)(0,1)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}v_1^{(1)(0,1)}=0;\\ u_1^{(1)(0,2)\prime}+u_1^{(1)(0,2)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(1)(0,2)}+(\varepsilon+g\sigma_v)\dfrac{h_{G+1}}{h_G}v_1^{(1)(0,2)}+\dfrac{\pi}{1+2G}u_1^{(1)(0,1)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}u_2^{(1)(0,1)}=0;\\ u_2^{(1)(0,2)\prime}+u_2^{(1)(0,2)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(1)(0,2)}-(\varepsilon+g\sigma_v)\dfrac{h_{G-1}}{h_G}v_2^{(1)(0,2)}-\dfrac{\pi}{1+2G}u_2^{(1)(0,1)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}u_1^{(1)(0,1)}=0; \end{cases}$$

$$\begin{cases} v_1^{(2)(0,2)\prime}+v_1^{(2)(0,2)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(2)(0,2)}-(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G+1}}u_1^{(2)(0,2)}+\dfrac{\pi}{1+2G}v_1^{(2)(0,1)}-\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}v_2^{(2)(0,1)}=0\\ v_2^{(2)(0,2)\prime}+v_2^{(2)(0,2)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(2)(0,2)}+(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G-1}}u_2^{(2)(0,2)}+\dfrac{\pi}{1+2G}v_2^{(2)(0,1)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}v_1^{(2)(0,1)}=0\\ u_1^{(2)(0,2)\prime}+u_1^{(2)(0,2)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(2)(0,2)}+(\varepsilon+g\sigma_v)\dfrac{h_{G+1}}{h_G}v_1^{(2)(0,2)}+\dfrac{\pi}{1+2G}u_1^{(2)(0,1)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}u_2^{(2)(0,1)}=0\\ u_2^{(2)(0,2)\prime}+u_2^{(2)(0,2)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(2)(0,2)}-(\varepsilon+g\sigma_v)\dfrac{h_{G-1}}{h_G}v_2^{(2)(0,2)}-\dfrac{\pi}{1+2G}u_2^{(2)(0,1)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}u_1^{(2)(0,1)}=0 \end{cases}$$

$$\begin{cases} v_1^{(1)(1,1)\prime}+v_1^{(1)(1,1)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(1)(1,1)}-(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G+1}}u_1^{(1)(1,1)}+\sigma_s u_1^{(1)(0,1)}+\dfrac{\pi}{1+2G}v_1^{(1)(1,0)}-\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}v_2^{(1)(1,0)}=0;\\ v_2^{(1)(1,1)\prime}+v_2^{(1)(1,1)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(1)(1,1)}+(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G-1}}u_2^{(1)(1,1)}-\sigma_s u_2^{(1)(0,1)}+\dfrac{\pi}{1+2G}v_2^{(1)(1,0)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}v_1^{(1)(1,0)}=0;\\ u_1^{(1)(1,1)\prime}+u_1^{(1)(1,1)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(1)(1,1)}+(\varepsilon+g\sigma_v)\dfrac{h_{G+1}}{h_G}v_1^{(1)(1,1)}+\sigma_s v_1^{(1)(0,1)}+\dfrac{\pi}{1+2G}u_1^{(1)(1,0)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}u_2^{(1)(1,0)}=0;\\ u_2^{(1)(1,1)\prime}+u_2^{(1)(1,1)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(1)(1,1)}-(\varepsilon+g\sigma_v)\dfrac{h_{G-1}}{h_G}v_2^{(1)(1,1)}-\sigma_s v_2^{(1)(0,1)}-\dfrac{\pi}{1+2G}u_2^{(1)(1,0)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}u_1^{(1)(1,0)}=0; \end{cases}$$

$$\begin{cases} v_1^{(2)(1,1)\prime}+v_1^{(2)(1,1)}\dfrac{h'_{G+1}}{h_{G+1}}+\dfrac{G+2}{r}v_1^{(2)(1,1)}-(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G+1}}u_1^{(2)(1,1)}+\sigma_s u_1^{(2)(0,1)}+\dfrac{\pi}{1+2G}v_1^{(2)(1,0)}-\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}v_2^{(2)(1,0)}=0;\\ v_2^{(2)(1,1)\prime}+v_2^{(2)(1,1)}\dfrac{h'_{G-1}}{h_{G-1}}-\dfrac{G-1}{r}v_2^{(2)(1,1)}+(\varepsilon-g\sigma_v)\dfrac{h_G}{h_{G-1}}u_2^{(2)(1,1)}-\sigma_s u_2^{(2)(0,1)}+\dfrac{\pi}{1+2G}v_2^{(2)(1,0)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}v_1^{(2)(1,0)}=0;\\ u_1^{(2)(1,1)\prime}+u_1^{(2)(1,1)}\dfrac{h'_G}{h_G}-\dfrac{G}{r}u_1^{(2)(1,1)}+(\varepsilon+g\sigma_v)\dfrac{h_{G+1}}{h_G}v_1^{(2)(1,1)}+\sigma_s v_1^{(2)(0,1)}+\dfrac{\pi}{1+2G}u_1^{(2)(1,0)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}u_2^{(2)(1,0)}=0;\\ u_2^{(2)(1,1)\prime}+u_2^{(2)(1,1)}\dfrac{h'_G}{h_G}+\dfrac{G+1}{r}u_2^{(2)(1,1)}-(\varepsilon+g\sigma_v)\dfrac{h_{G-1}}{h_G}v_2^{(2)(1,1)}-\sigma_s v_2^{(2)(0,1)}-\dfrac{\pi}{1+2G}u_2^{(2)(1,0)}+\dfrac{2\sqrt{G(1+G)}\pi}{1+2G}u_1^{(2)(1,0)}=0; \end{cases}$$

$$\beta=k/(\varepsilon+g\sigma_v),\sigma_s=\tilde{\sigma}(r)\text{ and }\pi=\tilde{\pi}(r)$$

$$\Pi=-(-1)^G*v_i^{(j)(n_s,n_p)}$$

$$V_G^{(n_s,n_p)}\equiv\begin{pmatrix} v_1^{(1)(n_s,n_p)} & v_1^{(2)(n_s,n_p)}\\ v_2^{(1)(n_s,n_p)} & v_2^{(2)(n_s,n_p)}\end{pmatrix}$$

$$V_G^{(1)}(r)=V_G^{(1,0)}(r)+V_G^{(0,1)}(r)$$
$$V_G^{(2)}(r)=V_G^{(2,0)}(r)+V_G^{(0,2)}(r)+V_G^{(1,1)}(r)$$

$$\delta_G^{(1)}(k)=\frac{1}{2i}\lim_{r\to 0}\mathrm{tr}\left[V_G^{(1)*}(r)-V_G^{(1)}(r)\right]$$
$$\delta_G^{(2)}(k)=\frac{1}{2i}\lim_{r\to 0}\mathrm{tr}\left[V_G^{(2)*}(r)-V_G^{(2)}(r)-\frac{1}{2}\left[V_G^{(1)}(r)\right]^2+\frac{1}{2}\left[V_G^{(1)*}(r)\right]^2\right]$$

$$\delta_0^{(1)}(k) = \frac{1}{2i}\lim_{r\to 0}\mathrm{tr}[v^*(r) - v(r)]$$
$$\delta_0^{(2)}(k) = \frac{1}{2i}\lim_{r\to 0}\mathrm{tr}\left[v^*(r) - v(r) - \frac{1}{2}v(r)^2 + \frac{1}{2}v^*(r)^2\right].$$

$$\begin{cases} g_1'' + \left(\dfrac{2}{r} - \dfrac{\varepsilon_1'}{\varepsilon_1}\right)g_1' + \left[-\dfrac{2\sqrt{G(G+1)}}{2G+1}\dfrac{g\pi}{r} + \dfrac{2\sqrt{G(G+1)}}{2G+1}g\pi' - \dfrac{\varepsilon_1'}{\varepsilon_1}\dfrac{2\sqrt{G(G+1)}}{2G+1}g\pi\right]g_2 \\ + \left[-\dfrac{(G+1)(G+2)}{r^2} + \dfrac{2(G+1)}{2G+1}\dfrac{g\pi}{r} - \dfrac{\varepsilon_1'}{\varepsilon_1}\dfrac{G+2}{r} + \dfrac{\varepsilon_1'}{\varepsilon_1}\dfrac{g\pi}{2G+1} - \dfrac{g\pi'}{2G+1} + \varepsilon^2 - g^2(\sigma^2+\pi^2)\right]g_1 = 0 \\ g_2'' + \left(\dfrac{2}{r} - \dfrac{\varepsilon_1'}{\varepsilon_1}\right)g_2' + \left[-\dfrac{2\sqrt{G(G+1)}}{2G+1}\dfrac{g\pi}{r} + \dfrac{2\sqrt{G(G+1)}}{2G+1}g\pi' - \dfrac{\varepsilon_1'}{\varepsilon_1}\dfrac{2\sqrt{G(G+1)}}{2G+1}g\pi\right]g_1 \\ + \left[-\dfrac{G(G-1)}{r^2} + \dfrac{2G}{2G+1}\dfrac{g\pi}{r} + \dfrac{\varepsilon_1'}{\varepsilon_1}\dfrac{G-1}{r} - \dfrac{\varepsilon_1'}{\varepsilon_1}\dfrac{g\pi}{2G+1} + \dfrac{g\pi'}{2G+1} + \varepsilon^2 - g^2(\sigma^2+\pi^2)\right]g_2 = 0, \end{cases}$$

$$\begin{cases} v_1^{(1)\prime\prime} + \dfrac{2}{r}v_1^{(1)\prime}\left(1 + \dfrac{rh_{G+1}'}{h_{G+1}}\right) - \dfrac{\varepsilon_1'}{\varepsilon_1}\left(v_1^{(1)\prime} + v_1^{(1)}\dfrac{h_{G+1}'}{h_{G+1}}\right) + w_{11}v_1^{(1)} + w_{12}v_2^{(1)} = 0 \\ v_2^{(1)\prime\prime} + \dfrac{2}{r}v_2^{(1)\prime}\left(1 + \dfrac{rh_{G+1}'}{h_{G+1}}\right) - \dfrac{\varepsilon_1'}{\varepsilon_1}\left(v_2^{(1)\prime} + v_2^{(1)}\dfrac{h_{G+1}'}{h_{G+1}}\right) + w_{21}v_1^{(1)} + w_{22}v_2^{(1)} \\ \qquad + \dfrac{2(2G+1)}{r^2}v_2^{(1)} = 0 \end{cases}$$

$$\begin{cases} v_1^{(2)\prime\prime} + \dfrac{2}{r}v_1^{(2)\prime}\left(1 + \dfrac{rh_{G-1}'}{h_{G-1}}\right) - \dfrac{\varepsilon_1'}{\varepsilon_1}\left(v_1^{(2)\prime} + v_1^{(2)}\dfrac{h_{G-1}'}{h_{G-1}}\right) + w_{11}v_1^{(2)} + w_{12}v_2^{(2)} \\ \qquad - \dfrac{2(2G+1)}{r^2}v_1^{(2)} = 0 \\ v_2^{(2)\prime\prime} + \dfrac{2}{r}v_2^{(2)\prime}\left(1 + \dfrac{rh_{G-1}'}{h_{G-1}}\right) - \dfrac{\varepsilon_1'}{\varepsilon_1}\left(v_2^{(2)\prime} + v_2^{(2)}\dfrac{h_{G-1}'}{h_{G-1}}\right) + w_{21}v_1^{(2)} + w_{22}v_2^{(2)} = 0, \end{cases}$$

$$\begin{aligned} w_{11} =& -g^2(\sigma^2+\pi^2-\sigma_v^2) - \frac{G+2}{r}\frac{\sigma'}{\sigma+\varepsilon/g} - \frac{g\pi'}{2G+1} \\ &+ \frac{2}{r}\frac{G+1}{2G+1}g\pi + \frac{g\pi}{2G+1}\frac{\sigma'}{\sigma+\varepsilon/g}, \\ w_{22} =& -g^2(\sigma^2+\pi^2-\sigma_v^2) + \frac{G-1}{r}\frac{\sigma'}{\sigma+\varepsilon/g} + \frac{g\pi'}{2G+1} \\ &+ \frac{2}{r}\frac{G}{2G+1}g\pi - \frac{g\pi}{2G+1}\frac{\sigma'}{\sigma+\varepsilon/g}, \\ w_{12} =& w_{21} = \frac{2\sqrt{G(G+1)}}{2G+1}\left[g\pi' - g\pi\left(\frac{1}{r} + \frac{\sigma'}{\sigma+\varepsilon/g}\right)\right]. \end{aligned}$$

$$\begin{cases} v_1^{(1)(1,0)\prime\prime}+\dfrac{2}{r}v_1^{(1)(1,0)\prime}\left(1+\dfrac{rh'_{G+1}}{h_{G+1}}\right)+\left(-\dfrac{\sigma'_s}{\varepsilon_v}\right)\dfrac{h'_{G+1}}{h_{G+1}} \\ \qquad +\left(-\dfrac{\sigma'_s}{\varepsilon_v}\right)\dfrac{G+2}{r}=0 \\ v_2^{(1)(1,0)\prime\prime}+\dfrac{2}{r}v_2^{(1)(1,0)\prime}\left(1+\dfrac{rh'_{G+1}}{h_{G+1}}\right)+\dfrac{2(2G+1)}{r^2}v_2^{(1)(1,0)}=0 \end{cases}$$

$$\begin{cases} v_1^{(2)(1,0)\prime\prime}+\dfrac{2}{r}v_1^{(2)(1,0)\prime}\left(1+\dfrac{rh'_{G-1}}{h_{G-1}}\right)-\dfrac{2(2G+1)}{r^2}v_1^{(2)(1,0)}=0 \\ v_2^{(2)(1,0)\prime\prime}+\dfrac{2}{r}v_2^{(2)(1,0)\prime}\left(1+\dfrac{rh'_{G-1}}{h_{G-1}}\right)+\left(-\dfrac{\sigma'_s}{\varepsilon_v}\right)\dfrac{h'_{G-1}}{h_{G-1}}+\dfrac{\sigma'_s}{\varepsilon_v}\dfrac{G-1}{r}=0 \end{cases}$$

$$\begin{cases} -\dfrac{\sigma'_s}{\varepsilon_v}\left(v_1^{(1)(1,0)\prime}+v_1^{(1)(1,0)}\dfrac{h'_{G+1}}{h_{G+1}}\right)+\left(-\dfrac{\sigma'_s}{\varepsilon_v}\right)\dfrac{G+2}{r}v_1^{(1)(1,0)}+\dfrac{\sigma'_s\sigma_s}{\varepsilon_v^2}\dfrac{h'_{G+1}}{h_{G+1}} \\ \qquad +\dfrac{\sigma'_s\sigma_s}{\varepsilon_v^2}\dfrac{G+2}{r}-(\sigma_s^2+2\sigma_v\sigma_s)+v_1^{(1)(2,0)\prime\prime} \\ \qquad +\dfrac{2}{r}v_1^{(1)(2,0)\prime}\left(1+\dfrac{rh'_{G+1}}{h_{G+1}}\right)=0 \\ -\dfrac{\sigma'_s}{\varepsilon_v}\left(v_2^{(1)(1,0)\prime}+v_2^{(1)(1,0)}\dfrac{h'_{G+1}}{h_{G+1}}\right)+\left(\dfrac{\sigma'_s}{\varepsilon_v}\right)\dfrac{G-1}{r}v_2^{(1)(1,0)}+v_2^{(1)(2,0)\prime\prime} \\ \qquad +\dfrac{2}{r}v_2^{(1)(2,0)\prime}\left(1+\dfrac{rh'_{G+1}}{h_{G+1}}\right)+\dfrac{2(2G+1)}{r^2}v_2^{(1)(2,0)}=0 \end{cases}$$

$$\begin{cases} -\dfrac{\sigma'_s}{\varepsilon_v}\left(v_1^{(2)(1,0)\prime}+v_1^{(2)(1,0)}\dfrac{h'_{G-1}}{h_{G-1}}\right)+\left(-\dfrac{\sigma'_s}{\varepsilon_v}\right)\dfrac{G+2}{r}v_1^{(2)(1,0)}+v_1^{(2)(2,0)\prime\prime} \\ \qquad +\dfrac{2}{r}v_1^{(2)(2,0)\prime}\left(1+\dfrac{rh'_{G-1}}{h_{G-1}}\right)-\dfrac{2(2G+1)}{r^2}v_1^{(2)(2,0)}=0 \\ -\dfrac{\sigma'_s}{\varepsilon_v}\left(v_2^{(2)(1,0)\prime}+v_2^{(2)(1,0)}\dfrac{h'_{G-1}}{h_{G-1}}\right)+\dfrac{\sigma'_s}{\varepsilon_v}\dfrac{G-1}{r}v_2^{(2)(1,0)}+\dfrac{\sigma'_s\sigma_s}{\varepsilon_v^2}\dfrac{h'_{G-1}}{h_{G-1}} \\ \qquad -\dfrac{\sigma'_s\sigma_s}{\varepsilon_v^2}\dfrac{G-1}{r}-(\sigma_s^2+2\sigma_v\sigma_s)+v_2^{(2)(2,0)\prime\prime} \\ \qquad +\dfrac{2}{r}v_2^{(2)(2,0)\prime}\left(1+\dfrac{rh'_{G-1}}{h_{G-1}}\right)=0 \end{cases}$$

$$\begin{cases} v_1^{(1)(0,1)''}+\dfrac{2}{r}v_1^{(1)(0,1)'}\left(1+\dfrac{rh'_{G+1}}{h_{G+1}}\right)+\dfrac{G+1}{2G+1}\dfrac{2}{r}\pi-\dfrac{\pi'}{2G+1}=0 \\ v_2^{(1)(0,1)''}+\dfrac{2}{r}v_2^{(1)(0,1)'}\left(1+\dfrac{rh'_{G+1}}{h_{G+1}}\right)+\dfrac{2(2G+1)}{r^2}v_2^{(1)(0,1)} \\ -\dfrac{\pi}{r}\dfrac{2\sqrt{G(G+1)}}{2G+1}+\dfrac{2\sqrt{G(G+1)}}{2G+1}\pi'=0 \end{cases}$$

$$\begin{cases} v_1^{(2)(0,1)''}+\dfrac{2}{r}v_1^{(2)(0,1)'}\left(1+\dfrac{rh'_{G-1}}{h_{G-1}}\right)-\dfrac{2(2G+1)}{r^2}v_1^{(2)(0,1)} \\ -\dfrac{\pi}{r}\dfrac{2\sqrt{G(G+1)}}{2G+1}+\dfrac{2\sqrt{G(G+1)}}{2G+1}\pi'=0 \\ v_2^{(2)(0,1)''}+\dfrac{2}{r}v_2^{(2)(0,1)'}\left(1+\dfrac{rh'_{G-1}}{h_{G-1}}\right)+\dfrac{G}{2G+1}\dfrac{2}{r}\pi+\dfrac{\pi'}{2G+1}=0 \end{cases}$$

$$\begin{cases} \left(-\dfrac{\pi'}{2G+1}+\dfrac{2}{r}\dfrac{G+1}{2G+1}\pi\right)v_1^{(1)(0,1)}+\dfrac{2\sqrt{G(G+1)}}{2G+1}\left(\pi'-\dfrac{\pi}{r}\right)v_2^{(1)(0,1)} \\ -\pi^2+v_1^{(1)(0,2)''}+\dfrac{2}{r}v_1^{(1)(0,2)'}\left(1+\dfrac{rh'_{G+1}}{h_{G+1}}\right)=0 \\ \dfrac{2\sqrt{G(G+1)}}{2G+1}\left(\pi'-\dfrac{\pi}{r}\right)v_1^{(1)(0,1)}+\left(\dfrac{\pi'}{2G+1}+\dfrac{2}{r}\dfrac{G}{2G+1}\pi\right)v_2^{(1)(0,1)} \\ +v_2^{(1)(0,2)''}+\dfrac{2}{r}v_2^{(1)(0,2)'}\left(1+\dfrac{rh'_{G+1}}{h_{G+1}}\right)+\dfrac{2(2G+1)}{r^2}v_2^{(1)(0,2)}=0 \end{cases}$$

$$\begin{cases} \left(-\dfrac{\pi'}{2G+1}+\dfrac{2}{r}\dfrac{G+1}{2G+1}\pi\right)v_1^{(2)(0,1)}+\dfrac{2\sqrt{G(G+1)}}{2G+1}\left(\pi'-\dfrac{\pi}{r}\right)v_2^{(2)(0,1)} \\ +v_1^{(2)(0,2)''}+\dfrac{2}{r}v_1^{(2)(0,2)'}\left(1+\dfrac{rh'_{G-1}}{h_{G-1}}\right)-\dfrac{2(2G+1)}{r^2}v_1^{(2)(0,2)}=0 \\ \dfrac{2\sqrt{G(G+1)}}{2G+1}\left(\pi'-\dfrac{\pi}{r}\right)v_1^{(2)(0,1)}+\left(\dfrac{\pi'}{2G+1}+\dfrac{2}{r}\dfrac{G}{2G+1}\pi\right)v_2^{(2)(0,1)}-\pi^2 \\ +v_2^{(2)(0,2)''}+\dfrac{2}{r}v_2^{(2)(0,2)'}\left(1+\dfrac{rh'_{G-1}}{h_{G-1}}\right)=0 \end{cases}$$

$$\begin{cases} \left(-\frac{\pi'}{2G+1}+\frac{2}{r}\frac{G+1}{2G+1}\pi\right)v_1^{(1)(0,1)}+\frac{2\sqrt{G(G+1)}}{2G+1}\left(\pi'-\frac{\pi}{r}\right)v_2^{(1)(1,0)} \\ +\frac{\sigma_s'}{\varepsilon_v}\frac{\pi}{2G+1}-\frac{\sigma_s'}{\varepsilon_v}\left(v_1^{(1)(0,1)\prime}+v_1^{(1)(0,1)}\frac{h_{G+1}'}{h_{G+1}}\right)-\frac{\sigma_s'}{\varepsilon_v}\frac{G+2}{r}v_1^{(1)(0,1)} \\ +v_1^{(1)(1,1)\prime\prime}+\frac{2}{r}v_1^{(1)(1,1)\prime}\left(1+\frac{rh_{G+1}'}{h_{G+1}}\right)=0, \\ \frac{2\sqrt{G(G+1)}}{2G+1}\left(\pi'-\frac{\pi}{r}\right)v_1^{(1)(1,0)}+\left(\frac{\pi'}{2G+1}+\frac{2}{r}\frac{G}{2G+1}\pi\right)v_2^{(1)(1,0)} \\ -\frac{\sigma_s'\pi}{\varepsilon_v}\frac{2\sqrt{G(G+1)}}{2G+1}-\frac{\sigma_s'}{\varepsilon_v}\left(v_2^{(1)(0,1)\prime}+v_2^{(1)(0,1)}\frac{h_{G+1}'}{h_{G+1}}\right) \\ +\frac{\sigma_s'}{\varepsilon_v}\frac{G-1}{r}v_2^{(1)(0,1)}+v_2^{(1)(1,1)\prime\prime}+\frac{2}{r}v_2^{(1)(1,1)\prime}\left(1+\frac{rh_{G+1}'}{h_{G+1}}\right) \\ +\frac{2(2G+1)}{r^2}v_2^{(1)(1,1)}=0, \end{cases}$$

$$\begin{cases} \left(-\frac{\pi'}{2G+1}+\frac{2}{r}\frac{G+1}{2G+1}\pi\right)v_1^{(2)(1,0)}+\frac{2\sqrt{G(G+1)}}{2G+1}\left(\pi'-\frac{\pi}{r}\right)v_2^{(2)(1,0)} \\ -\frac{\sigma_s'\pi}{\varepsilon_v}\frac{2\sqrt{G(G+1)}}{2G+1}-\frac{\sigma_s'}{\varepsilon_v}\left(v_1^{(2)(0,1)\prime}+v_1^{(2)(0,1)}\frac{h_{G-1}'}{h_{G-1}}\right) \\ -\frac{\sigma_s'}{\varepsilon_v}\frac{G+2}{r}v_1^{(2)(0,1)}+v_1^{(2)(1,1)\prime\prime}+\frac{2}{r}v_1^{(2)(1,1)\prime}\left(1+\frac{rh_{G-1}'}{h_{G-1}}\right) \\ -\frac{2(2G+1)}{r^2}v_1^{(2)(1,1)}=0, \\ \frac{2\sqrt{G(G+1)}}{2G+1}\left(\pi'-\frac{\pi}{r}\right)v_1^{(2)(1,0)}+\left(\frac{\pi'}{2G+1}+\frac{2}{r}\frac{G}{2G+1}\pi\right)v_2^{(2)(1,0)} \\ -\frac{\sigma_s'}{\varepsilon_v}\frac{\pi}{2G+1}-\frac{\sigma_s'}{\varepsilon_v}\left(v_2^{(2)(0,1)\prime}+v_2^{(2)(0,1)}\frac{h_{G-1}'}{h_{G-1}}\right)+\frac{\sigma_s'}{\varepsilon_v}\frac{G-1}{r}v_2^{(2)(0,1)} \\ +v_2^{(2)(1,1)\prime\prime}+\frac{2}{r}v_2^{(2)(1,1)\prime}\left(1+\frac{rh_{G-1}'}{h_{G-1}}\right)=0, \end{cases}$$

$$\frac{1}{m}E=\mathcal{E}_{\mathrm{cl}}(w,b_1,b_2,c)+(1-Q_{\mathrm{vac}})\epsilon_1(w,b_1,b_2,c)+\mathcal{E}_{\mathrm{vac}}(w,b_1,b_2,c)$$

$$s+i\vec{\tau}\cdot\vec{p}=\rho(\xi)\exp\left(i\vec{\tau}\cdot\hat{\vec{r}}\Theta(\xi)\right)$$

$$\rho(\xi)=1+b_1\left[1+b_2^2\frac{\xi}{w}\right]\exp\left(-b_2^2\frac{\xi}{w}\right)$$

$$\Theta(\xi)=-\pi\frac{e^{c^2}-1}{e^{c^2}-3+2e^{c^2\xi/w}}$$


$$\begin{aligned} Q_{\mathrm{vac}}&=\sum_{G,\Pi}(2G+1)\left[\frac{1}{\pi}\delta_{G,+,\Pi}(0)-n_{G,\Pi}^{(+)}\right] \\ &=-\sum_{G,\Pi}(2G+1)\left[\frac{1}{\pi}\delta_{G,-,\Pi}(0)-n_{G,\Pi}^{(-)}\right] \end{aligned}$$

$$\mathcal{L} = -\frac{1}{4}\left(F^a_{\mu\nu}\right)^2 + \bar{\psi}(i\not{D} - m)\psi - \frac{1}{2\xi}\left(\partial_\mu A^\mu\right)^2.$$


$$F^a_{\mu\nu} = \partial_\mu A^a_\nu - \partial_\nu A^a_\mu + g f^{abc} A^b_\mu A^c_\nu.$$

$$D_\mu = \partial_\mu - igA^a_\mu T^a$$


$$\left[\Gamma^{\text{OS}}_{\text{tree}}\right]^{a\mu} = \frac{igT^a}{2m}[(p+p')^\mu - \sigma^{\mu\alpha}k_\alpha],$$





$ig\gamma^\mu T^a$




$gf^{abc}\left[g^{\mu\nu}(k-p)^\rho + g^{\nu\rho}(p-q)^\mu + g^{\rho\mu}(q-k)^\nu\right]$








$$-i\frac{\delta^{ab}}{q^2}\left(g^{\mu\nu} - \xi\frac{q^\mu q^\nu}{q^2}\right)$$




$$\frac{i}{\not{p} - m}$$

$$S(p+k)(igT^a\gamma^\alpha) = \left(i\frac{\not{p} + \not{k} + m}{(p+k)^2 - m^2}\right)(igT^a\gamma^\alpha) = -gT^a\frac{A^\alpha_{p,k}}{D_{p+k}}$$


$$A^\alpha_{p,k} = B^\alpha_{p,k} - C^\alpha_p, \qquad B^\alpha_{p,k} = 2p^\alpha + k^\alpha - \sigma^{\alpha\beta}k_\beta$$
$$C^\alpha_p = \gamma^\alpha(\not{p} - m), \qquad D_q = q^2 - m^2.$$




$igT^a(p+p')^\mu$




$-igT^a\sigma^{\mu\alpha}k_\alpha$



$-ig^2T^aT^bg^{\mu\nu}$



$-ig^2T^aT^b\sigma^{\mu\nu}$



$$M^{\mu\nu,ab}(p,k_1,k_2) = -g^2T^aT^bS(p+k_1+k_2)\gamma^\mu S(p+k_1)\gamma^\nu$$

$$M^{\mu\nu,ab}(p,k_1,k_2) = g^2T^aT^b\frac{A^\mu_{p+k_1,k_2}A^\nu_{p+k_1}}{D_{p+k_1+k_2}D_{p+k_1}}.$$

$$C^\alpha_{p+k}A^\beta_{p,k} = D_{p+k}\gamma^\alpha\gamma^\beta = D_{p+k}\left(\sigma^{\alpha\beta}+g^{\alpha\beta}\right)$$

$$M^{\mu\nu,ab}(p,k_1,k_2) = g^2T^aT^b\left[\frac{B^\mu_{p+k_1,k_2}A^\nu_{p+k_1}}{D_{p+k_1+k_2}D_{p+k_1}} - \frac{g^{\mu\nu}+\sigma^{\mu\nu}}{D_{p+k_1+k_2}}\right].$$

$$J^\mu(a_1,a_2,a_3) = \int \frac{d^Dl}{i\pi^{\frac{D}{2}}}\frac{l^\mu}{\prod_{i=1}^3 D_i^{a_i}}$$

$$J^{\mu\nu}(a_1,a_2,a_3) = \int \frac{d^Dl}{i\pi^{\frac{D}{2}}}\frac{l^\mu l^\nu}{\prod_{i=1}^3 D_i^{a_i}}$$

$$D_i = (p_i+l)^2 - m_i^2.$$

$$J_3^D(a_1,a_2,a_3) = \int \frac{d^Dl}{i\pi^{\frac{D}{2}}}\frac{1}{\prod_{i=1}^3 D_i^{a_i}},$$

$$J_3^D(a_1,a_2,a_3) = \int \mathcal{D}x\frac{1}{c^{D/2}}\mathrm{e}^{\frac{P}{c}-f}$$

$$\int \mathcal{D}x = \prod_{i=1}^3 \frac{(-1)^{a_i}}{\Gamma(a_i)}\int_0^\infty x_i^{a_i-1}dx_i$$

$$c = \sum_{i=1}^3 x_i, f = \sum_{i=1}^3 x_im_i^2$$

$$P = -\sum_{i,j=1}^3 x_ix_jp_i\cdot p_j + c\sum_{i=1}^3 x_ip_i^2$$

$$J^{\mu}(a_1,a_2,a_3) = -\int \mathcal{D}x \frac{\sum_{i=1}^{3} x_i p_i^{\mu}}{c^{(D+2)/2}} \mathrm{e}^{\frac{P}{c}-f}$$

$$J^{\mu}(a_1,a_2,a_3) = \sum_{i=1}^{3} a_i p_i^{\mu} \mathbf{i}^{+} J_3^{D+2}(a_1,a_2,a_3),$$

$$\mathbf{i}^{+} J_3^{D+2}(\dots,a_i,\dots) = J_3^{D+2}(\dots,a_i+1,\dots).$$

$$J^{\mu\nu}(a_1,a_2,a_3) = -\frac{1}{2} g^{\mu\nu} J_3^{D+2}(a_1,a_2,a_3) + \sum_{i,j=1}^{3} c_{ij} p_i^{\mu} p_j^{\nu} \mathbf{i}^{+}\mathbf{j}^{+} J_3^{D+4}(a_1,a_2,a_3)$$

$$c_{ij} = \begin{cases} a_i(a_i+1) & \text{for } i=j \\ a_i a_j & \text{for } i \neq j. \end{cases}$$

$$\sum_{i=1}^{3} \frac{\partial}{\partial m_i^2} J_3^{D}(a_1,a_2,a_3) = -\int \mathcal{D}x \frac{\sum_{i=1}^{3} x_i}{c^{D/2}} \mathrm{e}^{\frac{P}{c}-f}$$

$$J_3^{D-2}(a_1,a_2,a_3) = -\sum_{i=1}^{3} a_i \mathbf{i}^{+} J_3^{D}(a_1,a_2,a_3).$$

$$J_1^{D}(a_1,a_2,a_3) = \int \frac{d^D l}{i\pi^{\frac{D}{2}}} \frac{1}{(p'+l)^{2a_1}(p+l)^{2a_2}(l^2-m^2)^{a_3}}$$

$$J_2^{D}(a_1,a_2,a_3) = \int \frac{d^D l}{i\pi^{\frac{D}{2}}} \frac{1}{D_{p'+l}^{a_1} D_{p+l}^{a_2} (l^2)^{a_3}}$$

$$\int \frac{d^D l}{i\pi^{\frac{D}{2}}} \frac{\partial}{\partial_{\mu}} \left[\left(x_3^{(i)} l_{\mu} + x_1^{(i)} p'_{\mu} + x_2^{(i)} p_{\mu} \right) A^{(i)} \right] = 0$$

$$A^{(1)} = \frac{1}{(p'+l)^{2a_1}(p+l)^{2a_2}(l^2-m^2)^{a_3}}$$

$$A^{(2)} = \frac{1}{D_{p'+l}^{a_1} D_{p+l}^{a_2} (l^2)^{a_3}}$$

$$x_1^{(1)} = \frac{2}{Y^{(1)}}(p^2 p' \cdot k - m^2 p \cdot k),$$
$$x_2^{(1)} = -\frac{2}{Y^{(1)}}(p'^2 p \cdot k - m^2 p' \cdot k),$$
$$x_1^{(2)} = \frac{2}{Y^{(2)}}(p^2 p' \cdot k + m^2 p \cdot k),$$
$$x_2^{(2)} = -\frac{2}{Y^{(2)}}(p'^2 p \cdot k + m^2 p' \cdot k),$$
$$x_3^{(i)} = \frac{4G}{Y^{(i)}}, i = 1,2,$$

$$G = p^2 p^2 - (p \cdot p')^2,$$

$$Y^{(1)} = -2k^2(m^4 - 2m^2 p \cdot p' + p^2 p'^2)$$

$$Y^{(2)} = -2\left[k^2(m^4 + p^2 p'^2) + 2m^2\left(p'^2 p \cdot p' + p^2(p \cdot p' - 2p'^2)\right)\right]$$

$$J_i^{D+2}(a_1, a_2, a_3) = -\frac{1}{x_3^{(i)}\left(d + 1 - \sum_{j=1}^{3} a_j\right)}\left[J_i^D(a_1, a_2, a_3) + x_1^{(i)} J_i^D(a_1 - 1, a_2, a_3)\right.$$
$$\left. + x_2^{(i)} J_i^D(a_1, a_2 - 1, a_3) + \left(x_3^{(i)} - x_1^{(i)} - x_2^{(i)}\right) J_i^D(a_1, a_2, a_3 - 1)\right]$$

$$S_1 = \{J_1^D(0,0,1), J_1^D(0,1,1), J_1^D(1,0,1) J_1^D(1,1,0), J_1^D(1,1,1)\}$$

$$S_2 = \{J_2^D(0,1,0), J_2^D(0,1,1), J_2^D(1,0,1) J_2^D(1,1,0), J_2^D(1,1,1)\}$$

$$i\left[\Gamma^{(A)}\right]^{\mu a} = g^3 T^b T^a T^b \int \frac{d^D l}{(2\pi)^D} \gamma^\alpha \frac{\not{p}' + \not{l} + m}{(p' + l)^2 - m^2} \gamma^\mu \frac{\not{p} + \not{l} + m}{(p + l)^2 - m^2} \gamma^\beta \Delta_{\alpha\beta}(l)$$

$$\Delta_{\alpha\beta}(l) = \frac{1}{l^2}\left(g_{\alpha\beta} - \xi \frac{l_\alpha l_\beta}{l^2}\right)$$

$$\left[\Gamma^{(A)}\right]^{\mu a} = g^3 \bar{C} T^a \int \frac{d^D l}{i(2\pi)^D} \gamma^\alpha \left(\frac{B^\mu_{p+l,k} B^\beta_{p,l}}{D_{p'+l} D_{p+l}} - \frac{B^\mu_{p+l,k} C^\beta_p}{D_{p'+l} D_{p+l}} - \frac{\gamma^\mu \gamma^\beta}{D_{p'+l}}\right) \Delta_{\alpha\beta}(l)$$

$$C_F = \frac{N^2 - 1}{2N}, C_A = N$$

$$\gamma^\alpha B^\mu_{p+l,k} = B^\mu_{p+l,k} \gamma^\alpha + 2(\gamma^\mu k^\alpha - g^{\mu\alpha} \not{k}),$$
$$= (p + p' + 2l)^\mu \gamma^\alpha - \sigma^{\mu\beta} k_\beta \gamma^\alpha + 2(\gamma^\mu k^\alpha - g^{\mu\alpha} \not{k}).$$

$$\left[\Gamma^{(A)}\right]^{\mu a} = \frac{g^3 \bar{C} T^a}{(4\pi)^{\frac{D}{2}}}\left(\left[\Gamma_L^{(A)}\right]^\mu + \left[\Gamma_T^{(A)}\right]^\mu\right)$$

$$\left[\Gamma_L^{(A)}\right]^\mu = \int \frac{d^D l}{i\pi^{\frac{D}{2}}}\left\{\frac{(p+p'+2l)^\mu}{D_{p'+l}D_{p+l}l^2}\left[2\not{p}+\left(2-D-\xi-2\xi\frac{l\cdot p}{l^2}\right)\not{l}+(\xi-D)(\not{p}-m)\right]\right.$$

$$\left.+\frac{(D-2-\xi)}{D_{p'+l}l^2}\gamma^\mu+\frac{2\xi l^\mu}{D_{p'+l}l^4}\not{l}\right\}$$

$$\left[\Gamma_T^{(A)}\right]^\mu = \int \frac{d^D l}{i\pi^{\frac{D}{2}}}\frac{1}{D_{p'+l}D_{p+l}l^2}\left\{\sigma^{\mu\nu}k_\nu\left[\left(D-2+\xi+2\xi\frac{l\cdot p}{l^2}\right)\not{l}-2\not{p}+(D-4-\xi)(\not{p}-m)\right]\right.$$

$$\left.+\frac{2\xi}{l^2}(k\cdot l\gamma^\mu-l^\mu\not{k})\not{l}(\not{p}-m)+4-2\xi\left(1+2\frac{l\cdot p}{l^2}\right)(k\cdot l\gamma^\mu-l^\mu\not{k})\right\}$$

$$\begin{aligned}\left[\Gamma_L^{(A)}\right]^\mu =&(p+p')^\mu\{[(2-D)J_2^{D+2}(2,1,1)+\xi(p^2-m^2)J_2^{D+2}(2,1,2)-\xi J_2^{D+2}(2,0,2)]\not{p}\\&+[(2-D)J_2^{D+2}(1,2,1)+\xi(p^2-m^2)J_2^{D+2}(1,2,2)]\not{p}'+[m(D-\xi)+(2-D+\xi)\not{p}]J_2^D(1,1,1)\}\\&+2[m(D-\xi)+(2-D+\xi)\not{p}][p^\mu J_2^{D+2}(1,2,1)+p'^\mu J_2^{D+2}(2,1,1)]+(D-2)[\gamma^\mu J_2^{D+2}(1,1,1)\\&-2\left((2p^\mu J_2^{D+4}(1,3,1)+p'^\mu J_2^{D+4}(2,2,1))\not{p}+(2p'^\mu J_2^{D+4}(3,1,1)+p^\mu J_2^{D+4}(2,2,1))\not{p}'\right)]\\&+(D-\xi-2)J_2^D(1,0,1)\gamma^\mu+\xi(m^2-p^2)\left[J_2^{D+2}(1,1,2)\gamma^\mu-2\left((2p^\mu J_2^{D+4}(1,3,2)+p'^\mu J_2^{D+4}(2,2,2))\not{p}\right.\right.\\&\left.\left.+(2p'^\mu J_2^{D+4}(3,1,2)+p^\mu J_2^{D+4}(2,2,2))\not{p}'\right)\right]\end{aligned}$$

$$\begin{aligned}\left[\Gamma_T^{(A)}\right]^\mu =&\sigma^{\mu\nu}k_\nu\{[(4-D+\xi)m+(D-6-\xi)\not{p}]J_2^D(1,1,1)+(D-2)\left[\not{p}J_2^{D+2}(1,2,1)+\not{p}'J_2^{D+2}(2,1,1)\right]\\&+\xi(m^2-p^2)\left[\not{p}J_2^{D+2}(1,2,2)+\not{p}'J_2^{D+2}(2,1,2)\right]+\xi\not{p}'J_2^{D+2}(2,0,2)\}+4J_2^D(1,1,1)(k\cdot p\gamma^\mu-p^\mu\not{k})\\&+2\xi\{\gamma^\mu\not{p}[2k\cdot pJ_2^{D+4}(1,3,2)+k\cdot p'J_2^{D+4}(2,2,2)]-\not{k}p[2p^\mu J_2^{D+4}(1,3,2)+p'^\mu J_2^{D+4}(2,2,2)]\\&+\gamma^\mu\not{p}'[2k\cdot p'J_2^{D+4}(3,1,2)+k\cdot pJ_2^{D+4}(2,2,2)]-\not{k}\not{p}'[2p^\mu J_2^{D+4}(1,3,2)+p'^\mu J_2^{D+4}(2,2,2)]\\&J_2^{D+2}(1,1,2)(k^\mu-\gamma^\mu\not{k})\}(\not{p}-m)+4[\gamma^\mu(k\cdot pJ_2^{D+2}(1,2,1)+k\cdot p'J_2^{D+2}(2,1,1))\\&-\not{k}(p^\mu J_2^{D+2}(1,2,1)+p'^\mu J_2^{D+2}(2,1,1))]+4(k^\mu-\gamma^\mu\not{k})\left[\not{p}J_2^{D+2}(1,2,1)+\not{p}'J_2^{D+2}(2,1,1)\right]\\&-2\xi J_2^{D+2}(2,0,2)(k\cdot p'\gamma^\mu-p'^\mu\not{k})-2\xi(m^2-p^2)[\gamma^\mu(k\cdot pJ_2^{D+2}(1,2,2)+k\cdot p'J_2^{D+2}(2,1,2))\\&-\not{k}(p^\mu J_2^{D+2}(1,2,2)+p'^\mu J_2^{D+2}(2,1,2))].\end{aligned}$$

$$\left[\Gamma^{(NA)}\right]^{\mu a} = g^3 f^{abc}T^cT^b\int\frac{d^Dl}{(2\pi)^D}\gamma^{\nu'}\left(\frac{-\not{l}+m}{l^2-m^2}\right)\gamma^{\rho'}\Delta_{\rho\rho'}(p+l)\Delta_{\nu\nu'}(p'+l)V_{\rho\nu\mu}(p+l,-p'-l,k),$$

$$V_{\alpha\beta\mu}(r,s,t) = g^{\alpha\mu}(t-r)^\beta + g^{\alpha\beta}(r-s)^\mu + g^{\beta\mu}(s-t)^\alpha$$

$$\begin{aligned}\left[\Gamma^{(NA)}\right]^{\mu a} =&\frac{1}{2}g^3C_AT^a\int\frac{d^Dl}{i(2\pi)^D}\gamma^{\nu'}\left(\frac{-\not{=}m}{q'^2q^2D_l}\right)\gamma^{\rho'}\{g^{\mu\rho'}(k-q)^{\nu'}-g^{\mu\nu'}(k+q')^{\rho'}+g^{\nu'\rho'}(q+q')^\mu\\&+\frac{\xi q^{\rho'}}{q^2}\left[g^{\mu\nu'}q\cdot(k+q')-k^{\nu'}q^\mu-q^{\nu'}q^\mu\right]+\frac{\xi q'^{\nu'}}{q'^2}\left[g^{\mu\rho'}q'\cdot(q-k)+k^{\rho'}q'^\mu-q^\mu q'^{\rho'}\right]\\&\left.+\frac{\xi^2q^{\rho'}q'^{\nu'}}{q'^2q^2}(k\cdot q'q^\mu-k\cdot qq'^\mu)\right\}\end{aligned}$$

$$\begin{aligned}\gamma^{\nu'}(-\not{=}+m)\gamma^{\rho'} &= \gamma^{\nu'}A^{\rho'}_{-l,0} = \gamma^{\nu'}B^{\rho'}_{-l,0}-\gamma^{\nu'}C^{\rho'}_{-l}\\&= -2\gamma^{\nu'}l^{\rho'}+\left(\sigma^{\nu'\rho'}+g^{\nu'\rho'}\right)(\not{l}+m)\end{aligned}$$

$$\left[\Gamma^{(NA)}\right]^{\mu a} = \frac{1}{2(4\pi)^{\frac{D}{2}}}g^3C_AT^a\left(\left[\Gamma_L^{(NA)}\right]^\mu+\left[\Gamma_T^{(NA)}\right]^\mu\right)$$

$$\left[\Gamma_L^{(NA)}\right]^\mu = \int \frac{d^D l}{i\pi^{\frac{D}{2}}} \frac{1}{q'^2 q^2 D_l} \{(q+q')^\mu [(D-3-\xi)l + m(D-1-\xi)] - 2(\not{k} - \not{q})l^\mu + 2\gamma^\mu l \cdot (k+q')$$
$$+ \frac{\xi}{q'^2} \not{q}' \left[2(q' \cdot (k-q) l^\mu - l \cdot k q'^\mu + l \cdot q' q^\mu) + \left((q^2 - k^2)\gamma^\mu + q'^\mu \not{k}\right)(l+m)\right]$$
$$- \frac{\xi}{q^2} \left[2l \cdot q (q \cdot (k+q')\gamma^\mu - \not{k} q^\mu - \not{q} q'^\mu) + \left((k^2 - q'^2)\gamma^\mu + q^\mu \not{k}\right)\not{q}(l+m)\right]\Big\}$$

$$\left[\Gamma_T^{(NA)}\right]^\mu = -\int \frac{d^D l}{i\pi^{\frac{D}{2}}} \frac{1}{q'^2 q^2 D_l} \left\{3\sigma^{\mu\nu} k_\nu (l+m) + \frac{\xi^2}{q'^2 q^2} q' [2l \cdot q - \not{q}(l+m)](q^\mu k \cdot q' - q'^\mu k \cdot q)\right\}.$$

$$\left[\Gamma_L^{(NA)}\right]^\mu = \sum_{i=1}^{12} \alpha_i h_i^\mu,$$

$$\left[\Gamma_T^{(NA)}\right]^\mu = \sum_{i=1}^{8} \beta_i T_i^\mu,$$

$$T_1^\mu = p' \cdot k p^\mu - p \cdot k p'^\mu,$$
$$T_2^\mu = (p' \cdot k p^\mu - p \cdot k p'^\mu)(\not{p} + \not{p}'),$$
$$T_3^\mu = k^2 \gamma^\mu - k^\mu \not{k},$$
$$T_4^\mu = -(p' \cdot k p^\mu - p \cdot k p^\mu)\sigma_{\nu\lambda} p^\nu p^\lambda,$$
$$T_5^\mu = \sigma^{\mu\nu} k_\nu,$$
$$T_6^\mu = \gamma^\mu (p'^2 - p^2) - (p+p')^\mu \not{k},$$
$$T_7^\mu = \frac{1}{2}(p^2 - p'^2)\left[(p+p')^\mu - \gamma^\mu (\not{p} + \not{p}')\right]$$
$$-(p+p')^\mu \sigma^{\nu\lambda} p^\nu p^\lambda,$$
$$T_8^\mu = \gamma^\mu \sigma_{\nu\lambda} p^\nu p^\lambda - p^\mu \not{p} + p^\mu \not{p}'.$$

$$\Gamma^\mu = \sum_{i=1}^{4} \lambda_i L_i^\mu + \sum_{i=1}^{8} \tau_i T_i^\mu,$$

$$\bar{u}_{s'}(p')\left[\Gamma^{(A)}\right]^{\mu a} u_s(p) = \bar{u}_{s'}(p')\left[g^3 \bar{C} T^a \int \frac{d^D l}{i(2\pi)^D} \gamma^\alpha \right.$$
$$\left. \times \frac{\not{p}' + \not{l} + m}{(p'+l)^2 - m^2} \gamma^\mu \frac{\not{p} + \not{l} + m}{l^2[(p+l)^2 - m^2]} \gamma_\alpha\right] u_s(p), \quad (53)$$

$$(\not{p} - m) u_s(p) = 0, \bar{u}_{s'}(p')(\not{p}' - m) = 0.$$

$$\bar{u}_{s'}(p') = \bar{u}_{s'}(p') \frac{(\not{p}' + m)}{2m}.$$

$$\bar{u}_{s'}[\Gamma^A]^\mu u_s = \bar{C} \bar{u}_{s'} \left[\frac{g^3}{2m} \int \frac{d^D l}{i(2\pi)^D} (\not{p}' + m) \times \gamma^\alpha \frac{\not{p}' + \not{l} + m}{(p'+l)^2 - m^2} \gamma^\mu \frac{\not{p} + \not{l} + m}{l^2[(p+l)^2 - m^2]} \gamma_\alpha\right] u_s,$$

$$\bar{u}_{s'}[\Gamma^A]^\mu u_s = \bar{C} \bar{u}_{s'} \left[\frac{g^3}{2m} \int \frac{d^D l}{i(2\pi)^D} \frac{1}{l^2 D_{p'+l} D_{p+l}} \times A^\alpha_{p'+l,-l} A^\mu_{p',l} A_{\alpha p,l}\right] u_s$$

$$\bar{u}_{s'}[\Gamma^A]^\mu u_s = \bar{C}\bar{u}_{s'}\left\{\frac{g^3}{2m}\int \frac{d^D l}{i(2\pi)^D}\frac{1}{l^2 D_{p'+l}D_{p+l}}\right.$$

$$\left.\times\left[B^\alpha_{p'+l,-l}B^\mu_{p+l,k}B_{\alpha p,l} - D_{p+l}B_{\alpha p'+l,-l}(\sigma^{\mu\alpha}+g^{\mu\alpha}) - D_{p'+l}(\sigma^{\alpha\mu}+g^{\mu\alpha})B_{\alpha p,l}\right]\right\}u_s$$

$$\bar{u}_{s'}[\Gamma^A]^\mu u_s = \frac{g\bar{C}}{2m}\bar{u}_{s'}([\Gamma^A_{\mathrm{sc-sc-sc}}]^\mu + \left[\Gamma^A_{\mathrm{sp-sc-sc}}\right]^\mu + \left[\Gamma^A_{\mathrm{sp-sp-sc}}\right]^\mu + \left[\Gamma^A_{\mathrm{sc-sc-sp}}\right]^\mu + \left[\Gamma^A_{\mathrm{sp-sc-sp}}\right]^\mu$$

$$+\left[\Gamma^A_{\mathrm{sp-sp-sp}}\right]^\mu + [\Gamma^A_{\mathrm{sc-sc}}]^\mu + \left[\Gamma^A_{\mathrm{sp-sc}}\right]^\mu + \left[\Gamma^A_{\mathrm{sc-sp}}\right]^\mu + \left[\Gamma^A_{\mathrm{sp-sp}}\right]^\mu)u_s$$

$$\bar{u}_{s'}(p')[\Gamma^A]^\mu u_s(p) = e\bar{u}_{s'}\left[\frac{F_1^A(k^2)}{2m}(p+p')^\mu - \frac{F^A_{\mathrm{sp}}(k^2)}{2m}\sigma^{\mu\nu}k_\nu\right]$$

$$F^A_{\mathrm{sp}}(k^2) = F_1^A(k^2) + F_2^A(k^2)$$

$$F_1^A(k^2) = \frac{e^2}{(4\pi)^{\frac{D}{2}}}\{(D-2)[J_2^{D+2}(1,1,1) + J_2^D(1,1,0)$$
$$-4m^2(J_2^{D+4}(2,2,1) + 2J_2^{D+4}(3,1,1))]$$
$$+4p\cdot p'[J_2^D(1,1,1) - 2J_2^{D+2}(2,1,1)]\}$$

$$F^A_{\mathrm{sp}}(k^2) = \frac{e^2}{(4\pi)^{\frac{D}{2}}}\{4p\cdot p'J_2^D(1,1,1)$$
$$-8(m^2+p\cdot p')J_2^{D+2}(2,1,1) + (D-2)[J_2^D(1,1,0)$$
$$+J_2^{D+2}(1,1,1)]\}$$

$$F_2^A(k^2) = \frac{4m^2e^2}{(4\pi)^{\frac{D}{2}}}\{(D-2)[J_2^{D+4}(2,2,1)$$
$$+2J_2^{D+4}(3,1,1)] - 2J_2^{D+2}(2,1,1)\}$$

$$g = 2F^A_{\mathrm{sp}}(0) = 2 + 2F_2^A(0)$$

$$F_1^A(0) = \frac{\alpha}{4\pi}\left[\frac{1}{\epsilon} + \ln\left(\frac{\bar{\mu}^2}{m^2}\right) - 2\ln\left(\frac{m^2}{m_\gamma^2}\right) + 4\right]$$

$$\bar{u}_{s'}\left[\Gamma^{(NA)}\right]^{\mu a}u_s = C_A\bar{u}_{s'}\left\{\frac{g^3}{4m}\int \frac{d^D l}{i(2\pi)^D}\frac{1}{q'^2q^2D_l}(\not{p}'+m)\gamma^\nu(-\not{\neq}+m)\gamma^\rho V_{\rho\nu\mu}(q,-q',k)\right\}u_s,$$

$$\bar{u}_{s'}\left[\Gamma^{(NA)}\right]^{\mu a}u_s = C_A\bar{u}_{s'}\left\{\frac{g^3}{4m}\int \frac{d^D l}{i(2\pi)^D}\frac{1}{q'^2q^2D_l}A^\nu_{-l,p'+l}A^\rho_{p,-p-l}V_{\rho\nu\mu}(q,-q',k)\right\}u_s$$

$$= C_A\bar{u}_{s'}\left\{\frac{g^3}{4m}\int \frac{d^D l}{i(2\pi)^D}\frac{1}{q'^2q^2D_l}\left[B^\nu_{-l,q'}B^\rho_{p,-q} - D_l(\sigma^{\nu\rho}+g^{\nu\rho})\right]V_{\rho\nu\mu}(q,-q',k)\right\}u_s$$

$$\bar{u}_{s'}\left[\Gamma^{(NA)}\right]^{\mu a}u_s = \frac{gC_A}{4m}\bar{u}_{s'}([\Gamma^{NA}_{\mathrm{sc-sc}}]^\mu + \left[\Gamma^{NA}_{\mathrm{sc-sp}}\right]^\mu + \left[\Gamma^{NA}_{\mathrm{sp-sp}}\right]^\mu + [\Gamma^{NA}_{\mathrm{sc}}]^\mu + \left[\Gamma^{NA}_{\mathrm{sp}}\right]^\mu)u_s$$

$$\bar{u}_{s'}\left[\Gamma^{(NA)}\right]^{\mu a}u_s = g\bar{u}_{s'}\left[\frac{F_1^{NA}(k^2)}{2m}(p+p')^\mu - \frac{F^{NA}_{\mathrm{sp}}(k^2)}{2m}\sigma^{\mu\nu}k_\nu\right]$$

$$F_2^{NA}(k^2) = F_{\text{sp}}^{NA}(k^2) - F_1^{NA}(k^2).$$

$$F_1^{NA}(0) = \frac{g^2 C_A}{2(4\pi)^2}\left[\frac{3}{\epsilon} + 4 + 3\ln\left(\frac{\bar{\mu}^2}{m^2}\right)\right].$$

$$G = \frac{g^2}{(4\pi)^2}\left[\Gamma_{\text{sp}-\text{sp}}^{NA}\right]^{\mu}$$

$$F_2^{NA}(0) = \frac{C_A g^2}{(4\pi)^2}\left[3 - \ln\left(\frac{m^2}{m_g^2}\right)\right],$$

$$\alpha_1 = (2-D)J_1^{D+2}(1,1,1) - \xi m^2 J_1^D(2,0,1) + \left(4p\cdot p' - \frac{m^2\xi}{2} - 2p^2\right)J_1^{D+2}(1,2,1) + 2J_1^D(1,1,0)$$
$$+m^2\xi J_1^D(2,1,1)(p^2 - 2p\cdot p' + p'^2) + J_1^D(1,1,1)(2m^2 + \xi[p^2 - 2p\cdot p' + p'^2)]$$
$$+J_1^{D+2}(2,1,1)\left(-\frac{m^2\xi}{2} - 2p\cdot p' + 4p'^2\right) - \xi J_1^D(0,1,1) - \xi J_1^D(2,0,0) - \frac{1}{2}\xi J_1^{D+2}(1,2,0)$$
$$-\frac{1}{2}\xi J_1^{D+2}(2,1,0) + \xi p^2 J_1^D(0,2,1) + 2\xi p^2 J_1^{D+2}(0,3,1) - \xi p^2 J_1^D(1,2,1)(p^2 - 2p\cdot p' + p'^2)$$
$$+\xi J_1^D(2,1,0)(p^2 - 2p\cdot p' + p'^2) - 2\xi p^2 J_1^{D+2}(1,3,1)(p^2 - 2p\cdot p' + p'^2) - 2\xi p'^2 J_1^{D+2}(3,0,1)$$
$$+2\xi p\cdot p' J_1^{D+2}(2,2,1)(-p^2 + 2p\cdot p' - p'^2) + 2\xi p'^2 J_1^{D+2}(3,1,1)(p^2 - 2p\cdot p' + p'^2),$$
$$\alpha_2 = m(D-\xi-1)J_1^D(1,1,1) + 2m(D-\xi-1)J_1^{D+2}(1,2,1) - m\xi J_1^{D+2}(2,1,1) + m\xi p^2 J_1^D(1,2,1)$$
$$+4m\xi p^2 J_1^{D+2}(1,3,1) + 6m\xi p^2 J_1^{D+4}(1,4,1) - m\xi J_1^{D+2}(2,2,1)(p^2 - 6p\cdot p' + 3p'^2)$$
$$+2m\xi J_1^{D+4}(2,3,1)(p^2 + 2p\cdot p' - p^2) + mJ_1^{D+4}(3,2,1)(4\xi p\cdot p' - 2\xi p'^2),$$
$$\alpha_3 = -[(m - Dm)J_1^D(1,1,1) - 2(D-1)mJ_1^{D+2}(2,1,1) - m\xi J_1^D(2,0,1) - m\xi J_1^{D+2}(1,2,1)$$
$$-2m\xi p^2 J_1^{D+2}(2,2,1) - 2m\xi p^2 J_1^{D+4}(2,3,1) + m\xi J_1^D(2,1,1)(p^2 - 4p\cdot p' + 2p'^2)$$
$$+2m\xi J_1^{D+2}(3,1,1)(p^2 - 6p\cdot p' + 3p'^2) - 2m\xi J_1^{D+4}(3,2,1)(p^2 + 2p\cdot p' - p'^2)$$
$$+6m\xi(p'^2 - 2p\cdot p')J_1^{D+4}(4,1,1) - 2m\xi J_1^{D+2}(3,0,1)],$$
$$\alpha_4 = m\xi J_1^D(0,2,1) + 2m\xi J_1^{D+2}(0,3,1) + \frac{1}{2}m\xi J_1^{D+2}(1,2,1) + \frac{1}{2}m\xi J_1^{D+2}(2,1,1)$$
$$-m\xi J_1^D(1,2,1)(p^2 - 2p\cdot p' + p'^2) - 2m\xi J_1^{D+2}(1,3,1)(p^2 - 2p\cdot p' + p'^2)$$
$$+m\xi J_1^{D+2}(2,2,1)(p^2 - 2p\cdot p' + p'^2),$$
$$\alpha_5 = -\left[m\xi J_1^D(2,0,1) + \frac{1}{2}m\xi J_1^{D+2}(1,2,1) + \frac{1}{2}m\xi J_1^{D+2}(2,1,1) + 2m\xi J_1^{D+2}(3,0,1)\right.$$
$$-m\xi J_1^D(2,1,1)(p^2 - 2p\cdot p' + p'^2) + m\xi J_1^{D+2}(2,2,1)(p^2 - 2p\cdot p' + p'^2)$$
$$-2m\xi J_1^{D+2}(3,1,1)(p^2 - 2p\cdot p' + p'^2)],$$
$$\alpha_6 = (D+1)J_1^{D+2}(1,2,1) + 4(D-2)J_1^{D+4}(1,3,1) + 2m^2\xi J_1^{D+2}(2,2,1) + 6m^2\xi J_1^{D+4}(1,4,1)$$
$$+2\xi(m^2 + p^2)J_1^{D+2}(1,3,1) + 2\xi(m^2 + p'^2)J_1^{D+4}(2,3,1) - \xi J_1^D(1,1,1) + 2\xi J_1^{D+2}(1,3,0)$$
$$+2\xi J_1^{D+2}(2,2,0) + 6\xi J_1^{D+4}(1,4,0) + 2\xi J_1^{D+4}(2,3,0) + \xi p^2 J_1^D(1,2,1) + 2\xi p'^2 J_1^{D+4}(3,2,1),$$
$$\alpha_7 = -[(-D-\xi+3)J_1^{D+2}(2,1,1) + (4-2D)J_1^{D+4}(2,2,1) + 2\xi J_1^{D+4}(2,3,1)(-m^2 + p^2 + 2p\cdot p')$$
$$-2\xi(m^2 - 2p\cdot p')J_1^{D+4}(3,2,1) + \xi(p'^2 - m^2)J_1^D(2,1,1) + 2\xi(p'^2 - m^2)J_1^{D+2}(3,1,1)$$
$$-2\xi J_1^D(1,1,1) - \xi J_1^D(2,1,0) + (2-2\xi)J_1^{D+2}(1,2,1) - 2\xi J_1^{D+2}(3,1,0) - 2\xi J_1^{D+4}(2,3,0)$$
$$-2\xi J_1^{D+4}(3,2,0) + \xi p^2 J_1^D(1,2,1) + 4\xi p^2 J_1^{D+2}(1,3,1) + 6\xi p^2 J_1^{D+4}(1,4,1)$$
$$-2\xi J_1^{D+2}(2,2,1)(p^2 - 3p\cdot p' + p'^2)],$$
$$\alpha_8 = -[(-D+\xi+3)J_1^{D+2}(1,2,1) + (4-2D)J_1^{D+4}(2,2,1) + m^2(-\xi)J_1^D(2,1,1) - 2m^2\xi J_1^{D+4}(2,3,1)$$
$$+\xi(p^2 - m^2)J_1^D(1,2,1) + 2\xi(p^2 - m^2)J_1^{D+2}(1,3,1) + \xi J_1^{D+2}(2,2,1)(p'^2 - m^2 + 2p^2 - 4p\cdot p')$$
$$-2\xi(m^2 + p'^2)J_1^{D+2}(3,1,1) - 2\xi(m^2 + p'^2)J_1^{D+4}(3,2,1) - \xi J_1^D(1,1,1) - \xi J_1^D(1,2,0) - \xi J_1^D(2,1,0)$$
$$-2\xi J_1^{D+2}(1,3,0) + (\xi-4)J_1^{D+2}(2,1,1) - \xi J_1^{D+2}(2,2,0) - 2\xi J_1^{D+2}(3,1,0) - 2\xi J_1^{D+4}(2,3,0)$$
$$-2\xi J_1^{D+4}(3,2,0) - 6\xi p^2 J_1^{D+4}(4,1,1)],$$

$$
\begin{aligned}
\alpha_9 =&(D+\xi-5)J_1^{D+2}(2,1,1)+4(D-2)J_1^{D+4}(3,1,1)+\xi(m^2-3p^2)J_1^{D+2}(2,2,1)\\
&+2\xi J_1^{D+4}(3,2,1)(m^2-p^2-2p\cdot p')-\xi J_1^{D}(2,1,1)(m^2+p^2-p^2)+6\xi(m^2-2p\cdot p')J_1^{D+4}(4,1,1)\\
&-\xi J_1^{D}(1,1,1)+\xi J_1^{D}(2,0,1)-\xi J_1^{D}(2,1,0)+\xi J_1^{D+2}(2,2,0)+2\xi J_1^{D+2}(3,0,1)+2\xi J_1^{D+4}(3,2,0)\\
&+6\xi J_1^{D+4}(4,1,0)-2\xi p^2 J_1^{D+4}(2,3,1)-2\xi J_1^{D+2}(3,1,1)(p^2+2p\cdot p'-p'^2)\\
\alpha_{10} =&-[m\xi J_1^{D}(1,2,1)+4m\xi J_1^{D+2}(1,3,1)+2m\xi J_1^{D+2}(2,2,1)+6m\xi J_1^{D+4}(1,4,1)+4m\xi J_1^{D+4}(2,3,1)\\
&+2m\xi J_1^{D+4}(3,2,1)]\\
\alpha_{11} =&-m\xi J_1^{D}(2,1,1)-2m\xi J_1^{D+2}(2,2,1)-4m\xi J_1^{D+2}(3,1,1)-2m\xi J_1^{D+4}(2,3,1)-4m\xi J_1^{D+4}(3,2,1)\\
&-6m\xi J_1^{D+4}(4,1,1)\\
\alpha_{12} =&-\left[-\frac{1}{2}\xi J_1^{D+2}(1,2,1)-\frac{1}{2}\xi J_1^{D+2}(2,1,1)-2\xi J_1^{D+2}(2,2,1)(p^2-2p\cdot p'+p'^2)\right]
\end{aligned}
$$

$$
\begin{aligned}
\beta_1 =&\ 2m\xi^2 J_1^{D+2}(2,3,1)(m^2+p^2+2p\cdot p')+m\xi^2(m^2+p\cdot p')J_1^{D}(2,2,1)\\
&+2m\xi^2 J_1^{D+2}(3,2,1)(m^2+2p\cdot p'+p'^2)+m\xi^2 J_1^{D}(2,2,0)-m\xi^2 J_1^{D+2}(2,2,1)+2m\xi^2 J_1^{D+2}(2,3,0)\\
&+2m\xi^2 J_1^{D+2}(3,2,0)+6m\xi^2(p^2+p\cdot p')J_1^{D+4}(2,4,1)+4m\xi^2 J_1^{D+4}(3,3,1)(p^2+2p\cdot p'+p'^2)\\
&+6m\xi^2(p\cdot p'+p'^2)J_1^{D+4}(4,2,1),\\
\beta_2 =&-m^2\xi^2 J_1^{D}(2,2,1)-\xi^2(3m^2+p^2)J_1^{D+2}(2,3,1)-3\xi^2(m^2+p^2)J_1^{D+4}(2,4,1)\\
&-2\xi^2 J_1^{D+4}(3,3,1)(2m^2+p^2+p'^2)-\xi^2(3m^2+p'^2)J_1^{D+2}(3,2,1)-3\xi^2(m^2+p'^2)J_1^{D+4}(4,2,1)\\
&-\xi^2 J_1^{D}(2,2,0)+\frac{1}{2}\xi^2 J_1^{D+2}(2,2,1)-3\xi^2 J_1^{D+2}(2,3,0)-3\xi^2 J_1^{D+2}(3,2,0)-3\xi^2 J_1^{D+4}(2,4,0)\\
&-4\xi^2 J_1^{D+4}(3,3,0)-3\xi^2 J_1^{D+4}(4,2,0),\\
\beta_3 =&-\frac{1}{2}\xi^2(m^2-p^2)(p^2-p'^2)J_1^{D+2}(2,3,1)+\frac{1}{2}\xi^2(m^2-p'^2)(p^2-p'^2)J_1^{D+2}(3,2,1)\\
&-\frac{3}{2}\xi^2(m^2-p^2)(p^2-p'^2)J_1^{D+4}(2,4,1)+\frac{3}{2}\xi^2(m^2-p'^2)(p^2-p'^2)J_1^{D+4}(4,2,1)\\
&+\frac{1}{2}\xi^2(m^2-p\cdot p')J_1^{D+2}(2,2,1)+\frac{1}{2}\xi^2 J_1^{D+2}(2,2,0)+\frac{1}{2}\xi^2(p'^2-p^2)J_1^{D+2}(2,3,0)\\
&+\frac{1}{2}\xi^2(p^2-p^2)J_1^{D+2}(3,2,0)+\frac{3}{2}\xi^2(p'^2-p^2)J_1^{D+4}(2,4,0)+\xi^2(p^2-p'^2)^2 J_1^{D+4}(3,3,1)\\
&+\frac{3}{2}\xi^2(p^2-p^2)J_1^{D+4}(4,2,0)+\frac{3}{2}J_1^{D+2}(1,2,1)-\frac{3}{2}J_1^{D+2}(2,1,1),\\
\beta_4 =&-m\xi^2 J_1^{D}(2,2,1)-4m\xi^2 J_1^{D+2}(2,3,1)-4m\xi^2 J_1^{D+2}(3,2,1)-6m\xi^2 J_1^{D+4}(2,4,1)\\
&-8m\xi^2 J_1^{D+4}(3,3,1)-6m\xi^2 J_1^{D+4}(4,2,1),\\
\beta_5 =&\frac{1}{2}m\xi^2 J_1^{D+2}(2,2,1)(p^2-2p\cdot p'+p^2)-3mJ_1^{D}(1,1,1),\\
\beta_6 =&-\frac{1}{2}\xi^2(m^2-p^2)J_1^{D+2}(2,3,1)(p^2-2p\cdot p'+p'^2)+\frac{1}{2}\xi^2(m^2-p'^2)J_1^{D+2}(3,2,1)(p^2-2p\cdot p'+p'^2)\\
&-\frac{3}{2}\xi^2(m^2-p^2)J_1^{D+4}(2,4,1)(p^2-2p\cdot p'+p'^2)+\frac{3}{2}\xi^2(m^2-p'^2)J_1^{D+4}(4,2,1)(p^2-2p\cdot p'+p'^2)\\
&-\frac{1}{2}\xi^2 J_1^{D+2}(2,3,0)(p^2-2p\cdot p'+p'^2)+\frac{1}{2}\xi^2 J_1^{D+2}(3,2,0)(p^2-2p\cdot p'+p'^2)\\
&-\frac{3}{2}\xi^2 J_1^{D+4}(2,4,0)(p^2-2p\cdot p'+p'^2)+\xi^2(p^2-p'^2)J_1^{D+4}(3,3,1)(p^2-2p\cdot p'+p'^2)\\
&+\frac{3}{2}\xi^2 J_1^{D+4}(4,2,0)(p^2-2p\cdot p'+p^2)-\frac{3}{2}J_1^{D+2}(1,2,1)-\frac{3}{2}J_1^{D+2}(2,1,1),\\
\beta_7 =&\ 0,\\
\beta_8 =&-\frac{1}{2}\xi^2 J_1^{D+2}(2,2,1)(p^2-2p\cdot p'+p'^2)-3J_1^{D+2}(1,2,1)-3J_1^{D+2}(2,1,1).
\end{aligned}
$$

$$\left[\Gamma^{A}_{\mathrm{sc-sc-sp}}\right]^{\mu}=-g^{2}\int\frac{d^{D}l}{i(2\pi)^{D}}\frac{(2p'+l)\cdot(2p+l)\sigma^{\mu\nu}k_{\nu}}{l^{2}D_{p'+l}D_{p+l}}$$

$$\left[\Gamma^{A}_{\mathrm{sp-sp-sc}}\right]^{\mu}=-g^{2}\int\frac{d^{D}l}{i(2\pi)^{D}}\frac{(2l+p+p')^{\mu}\sigma^{\alpha\beta}\sigma_{\alpha\nu}l_{\beta}l^{\nu}}{l^{2}D_{p'+l}D_{p+l}}$$

$$\left[\Gamma^{A}_{\mathrm{sc-sc-sc}}\right]^{\mu}=g^{2}\int\frac{d^{D}l}{i(2\pi)^{D}}\frac{(2p'+l)\cdot(2p+l)(2l+p+p')^{\mu}}{l^{2}D_{p'+l}D_{p+l}}$$

$$\left[\Gamma^{A}_{\mathrm{sp-sp-sp}}\right]^{\mu}=g^{2}\int\frac{d^{D}l}{i(2\pi)^{D}}\frac{\sigma^{\alpha\beta}\sigma^{\mu\nu}\sigma_{\alpha\rho}l_{\beta}k^{\nu}l^{\rho}}{l^{2}D_{p'+l}D_{p+l}}$$

$$\left[\Gamma^{A}_{\mathrm{sp-sc-sc}}\right]^{\mu}=-2g^{2}\int\frac{d^{D}l}{i(2\pi)^{D}}\frac{(2l+p'+p)^{\mu}\sigma^{\alpha\beta}k_{\alpha}l_{\beta}}{l^{2}D_{p'+l}D_{p+l}}$$

$$\left[\Gamma^{A}_{\mathrm{sp-sc-sp}}\right]^{\mu}=g^{2}\int\frac{d^{D}l}{i(2\pi)^{D}}\frac{l_{\beta}k_{\nu}}{l^{2}D_{p'+l}D_{p+l}}\left[\sigma^{\mu\nu}\sigma^{\alpha\beta}(2p'+l)_{\alpha}-\sigma^{\alpha\beta}\sigma^{\mu\nu}(2p+l)_{\alpha}\right]$$

$$\left[\Gamma^{A}_{\mathrm{sc-sc}}\right]^{\mu}=-g^{2}\int\frac{d^{D}l}{i(2\pi)^{D}}\frac{1}{l^{2}}\left[\frac{(2p'+l)^{\mu}}{D_{p'+l}}+\frac{(2p+l)^{\mu}}{D_{p+l}}\right]$$

$$\left[\Gamma^{A}_{\mathrm{sc-sp}}\right]^{\mu}=-g^{2}\int\frac{d^{D}l}{i(2\pi)^{D}}\frac{1}{l^{2}}\left[\frac{\sigma^{\mu\nu}(2p'+l)_{\nu}}{D_{p'+l}}-\frac{\sigma^{\mu\nu}(2p+l)_{\nu}}{D_{p+l}}\right]$$

$$\left[\Gamma^{A}_{\mathrm{sp-sc}}\right]^{\mu}=-g^{2}\int\frac{d^{D}l}{i(2\pi)^{D}}\frac{1}{l^{2}}\left[\frac{\sigma^{\mu\nu}}{D_{p'+l}}-\frac{\sigma^{\mu\nu}}{D_{p+l}}\right]l_{\nu}$$

$$\left[\Gamma^{A}_{\mathrm{sp-sp}}\right]^{\mu}=-g^{2}\int\frac{d^{D}l}{i(2\pi)^{D}}\frac{1}{l^{2}}\left[\frac{\sigma_{\nu\alpha}\sigma^{\alpha\mu}}{D_{p'+l}}+\frac{\sigma^{\mu\alpha}\sigma_{\alpha\nu}}{D_{p+l}}\right]^{\nu}$$

$$\bar{u}_{s'}[\Gamma^{A}_{\mathrm{a}}]^{\mu}u_{s}=\bar{u}_{s'}\left[\bar{F}^{a}_{1}(p+p')^{\mu}-\bar{F}^{a}_{\mathrm{sp}}\sigma^{\mu\nu}k_{\nu}\right]$$

$$\bar{F}_{1}^{\mathrm{sc-sc-sp}}=0$$

$$\bar{F}_{1}^{\mathrm{sp-sp-sc}}=\frac{g^{2}}{(4\pi)^{\frac{D}{2}}}(D-1)[J_{2}^{D}(1,1,0)+2J_{2}^{D+2}(2,1,0)]$$

$$\bar{F}_{1}^{\mathrm{sc-sc-sc}}=\frac{g^{2}}{(4\pi)^{\frac{D}{2}}}[2J_{2}^{D+2}(2,1,0)+4(m^{2}+3p\cdot p')J_{2}^{D+2}(2,1,1)+4(m^{2}+p\cdot p')[J_{2}^{D+4}(2,2,1)$$
$$+2J_{2}^{D+4}(3,1,1)]+4p\cdot p'J_{2}^{D}(1,1,1)+J_{2}^{D}(1,1,0)-2J_{2}^{D+2}(1,1,1)],$$

$$\bar{F}_{1}^{\mathrm{sp-sp-sp}}=-\frac{2g^{2}}{(4\pi)^{\frac{D}{2}}}(D-4)(m^{2}-p\cdot p')[J_{2}^{D+4}(2,2,1)+2J_{2}^{D+4}(3,1,1)],$$

$$\bar{F}_{1}^{\mathrm{sp-sc-sc}}=-\frac{4g^{2}}{(4\pi)^{\frac{D}{2}}}(m^{2}-p\cdot p')[2J_{2}^{D+4}(3,1,1)+J_{2}^{D+4}(2,2,1)+J_{2}^{D+2}(2,1,1)]$$

$$\bar{F}_{1}^{\mathrm{sp-sc-sp}}=\frac{4g^{2}}{(4\pi)^{\frac{D}{2}}}(m^{2}-p\cdot p')J_{2}^{D+2}(2,1,1),$$

$$\bar{F}_{1}^{\mathrm{sc-sc}}=\frac{g^{2}}{(4\pi)^{\frac{D}{2}}}[-2(m^{2}+p\cdot p')[2J_{2}^{D+2}(2,1,1)+J_{2}^{D+4}(2,2,1)+2J_{2}^{D+4}(3,1,1)]$$
$$-2J_{2}^{D}(1,1,0)+J_{2}^{D+2}(1,1,1)-2J_{2}^{D+2}(2,1,0)]$$

$$\bar{F}_{1}^{\mathrm{sc-sp}}=0$$

$$\bar{F}_{1}^{\mathrm{sp-sc}}=0$$

$$\bar{F}_{1}^{\mathrm{sp-sp}}=\frac{g^{2}}{(4\pi)^{\frac{D}{2}}}(D-1)[J_{2}^{D+2}(1,1,1)-2(m^{2}+p\cdot p')[J_{2}^{D+4}(2,2,1)+2J_{2}^{D+4}(3,1,1)]$$
$$+2J_{2}^{D+2}(2,1,0)],$$

$$\bar{F}_{\mathrm{sp}}^{\mathrm{sc-sc-sp}} = \frac{g^2}{(4\pi)^{\frac{D}{2}}}[J_2^D(1,1,0) + 4(m^2 + p\cdot p')J_2^{D+2}(2,1,1) + 4p\cdot p' J_2^D(1,1,1)]$$

$$\bar{F}_{\mathrm{sp}}^{\mathrm{sp-sp-sc}} = 0$$

$$\bar{F}_{\mathrm{sp}}^{\mathrm{sc-sc-sc}} = 0$$

$$\bar{F}_{\mathrm{sp}}^{\mathrm{sp-sp-sp}} = \frac{g^2}{(4\pi)^{\frac{D}{2}}}[(D-5)J_2^D(1,1,0) + 2(D-4)[(m^2 - p\cdot p')(J_2^{D+4}(2,2,1) - 2J_2^{D+4}(3,1,1)) + J_2^{D+2}(1,1,1)]\Big]$$

$$\bar{F}_{\mathrm{sp}}^{\mathrm{sp-sc-sc}} = \frac{2g^2}{(4\pi)^{\frac{D}{2}}} J_2^{D+2}(1,1,1)$$

$$\bar{F}_{\mathrm{sp}}^{\mathrm{sp-sc-sp}} = 0$$

$$\bar{F}_{\mathrm{sp}}^{\mathrm{sc-sc}} = 0$$

$$\bar{F}_{\mathrm{sp}}^{\mathrm{sc-sp}} = \frac{g^2}{(4\pi)^{\frac{D}{2}}}[2(m^2 - p\cdot p')[J_2^{D+4}(2,2,1) - 2J_2^{D+4}(3,1,1)] + 4(m^2 + p\cdot p')J_2^{D+2}(2,1,1) + 2J_2^D(1,1,0) + J_2^{D+2}(1,1,1)]$$

$$\bar{F}_{\mathrm{sp}}^{\mathrm{sp-sc}} = \frac{g^2}{(4\pi)^{\frac{D}{2}}}[2(m^2 - p\cdot p')[J_2^{D+4}(2,2,1) - 2J_2^{D+4}(3,1,1)] + J_2^{D+2}(1,1,1)]$$

$$\bar{F}_{\mathrm{sp}}^{\mathrm{sp-sp}} = -\frac{g^2}{(4\pi)^{\frac{D}{2}}}(D-2)[2(m^2 - p\cdot p')[J_2^{D+4}(2,2,1) - 2J_2^{D+4}(3,1,1)] + J_2^{D+2}(1,1,1)]$$

$$\bar{u}_{s'}[(p+p')^\mu - \sigma^{\mu\nu}k_\nu]u_s = 2m\bar{u}_{s'}\gamma^\mu u_s.$$

$$[\Gamma_{\mathrm{sc-sc}}^{NA}]^\mu = g^2\int \frac{d^Dl}{i(2\pi)^D}\frac{(p'-l)^\nu(p-l)^\rho}{q'^2q^2D_l}V_{\rho\nu\mu}(q,-q',k)$$

$$\left[\Gamma_{\mathrm{sc-sp}}^{NA}\right]^\mu = g^2\int \frac{d^Dl}{i(2\pi)^D}\frac{(p'-l)^\nu\sigma^{\rho\beta}q_\beta - (p-l)^\rho\sigma^{\nu\beta}q'_\beta}{q'^2q^2D_l}V_{\rho\nu\mu}(q,-q',k)$$

$$\left[\Gamma_{\mathrm{sc-sp}}^{NA}\right]^\mu = -g^2\int \frac{d^Dl}{i(2\pi)^D}\frac{\sigma^{\nu\alpha}\sigma^{\rho\beta}q'_\alpha q_\beta}{q'^2q^2D_l}V_{\rho\nu\mu}(q,-q',k)$$

$$[\Gamma_{\mathrm{sc}}^{NA}]^\mu = -g^2\int \frac{d^Dl}{i(2\pi)^D}\frac{g^{\nu\rho}}{q'^2q^2D_l}V_{\rho\nu\mu}(q,-q',k)$$

$$\left[\Gamma_{\mathrm{sp}}^{NA}\right]^\mu = -g^2\int \frac{d^Dl}{i(2\pi)^D}\frac{\sigma^{\nu\rho}}{q'^2q^2D_l}V_{\rho\nu\mu}(q,-q',k)$$

$$\bar{u}_{s'}[\Gamma_{\mathrm{a}}^{NA}]^\mu u_s = \bar{u}_{s'}\left[\tilde{F}_1^a(p+p')^\mu - \tilde{F}_{\mathrm{sp}}^a\sigma^{\mu\nu}k_\nu\right]$$

$$\tilde{F}_1^{\mathrm{sc-sc}} = (5m^2 - p\cdot p')J_1^D(1,1,1) + 2(m^2 + 3p\cdot p')J_1^{D+2}(2,1,1) - J_1^D(1,0,1) + 3J_1^D(1,1,0) - J_1^{D+2}(2,0,1) + 2J_1^{D+2}(2,1,0)$$

$$\tilde{F}_1^{\mathrm{sc-sp}} = 2\{4(m^2 - p\cdot p')J_1^{D+2}(2,1,1) - 2m^2J_1^D(1,1,1) + 4m^2[J_1^{D+4}(2,2,1) + 2J_1^{D+4}(3,1,1)] + J_1^D(1,0,1) - J_1^D(1,1,0) - J_1^{D+2}(1,1,1) - J_1^{D+2}(2,0,1) + 2J_1^{D+2}(2,1,0)\}$$

$$\tilde{F}_1^{\mathrm{sp-sp}} = -J_1^D(1,1,1)[(7-2D)m^2 + p\cdot p'] + 2J_1^{D+2}(2,1,1)[(4D-17)m^2 + p\cdot p'] + (D-4)J_1^D(1,1,0) + 4(D-4)m^2[J_1^{D+4}(2,2,1) + 2J_1^{D+4}(3,1,1)] - (D-4)J_1^{D+2}(1,1,1) + 2(D-4)J_1^{D+2}(2,1,0) + 2(p\cdot p' - m^2)J_1^D(1,1,1) + 3J_1^D(1,0,1) + 3J_1^{D+2}(2,0,1)$$

$$\tilde{F}_1^{\mathrm{sc}} = -(D-1)[J_1^D(1,1,0) + 2J_1^{D+2}(2,1,0)]$$

$$\tilde{F}_1^{\mathrm{sp}} = 0$$

$$\tilde{F}_{\mathrm{sp}}^{\mathrm{sc-sc}} = 0$$

$$\tilde{F}_{\mathrm{sp}}^{\mathrm{sc-sp}} = 2(m^2 - p\cdot p')J_1^D(1,1,1) + J_1^D(1,1,0) - 2[J_1^{D+2}(1,1,1) + J_1^{D+2}(2,0,1)]$$

$$\tilde{F}_{\mathrm{sp}}^{\mathrm{sp-sp}} = -(D-4)J_1^{D+2}(1,1,1) + 2(p\cdot p' - m^2)J_1^D(1,1,1) + 4J_1^D(1,0,1) + 2J_1^{D+2}(2,0,1),$$

$$\tilde{F}_{\mathrm{sp}}^{\mathrm{sc}} = 0,$$

$$\tilde{F}_{\mathrm{sp}}^{\mathrm{sp}} = -3J_1^D(1,1,0).$$

$$(4\pi\mu^2)^{\epsilon}J_2^{6-2\epsilon}(1,1,1)=-\left[\frac{1}{2\epsilon}+\frac{1}{2}\ln\left(\frac{\bar{\mu}^2}{m^2}\right)+\frac{1}{2}\right]$$
$$(4\pi\mu^2)^{\epsilon}J_2^{4-2\epsilon}(1,1,0)=\frac{1}{\epsilon}+\ln\left(\frac{\bar{\mu}^2}{m^2}\right)$$
$$(4\pi\mu^2)^{\epsilon}J_1^{4-2\epsilon}(1,0,1)=\frac{1}{\epsilon}+2+\ln\left(\frac{\bar{\mu}^2}{m^2}\right)$$
$$(4\pi\mu^2)^{\epsilon}J_1^{6-2\epsilon}(2,0,1)=-\frac{1}{2}\left[\frac{1}{\epsilon}+3+\ln\left(\frac{\bar{\mu}^2}{m^2}\right)\right]$$
$$(4\pi\mu^2)^{\epsilon}J_1^{6-2\epsilon}(1,1,1)=(4\pi\mu^2)^{\epsilon}J_1^{6-2\epsilon}(2,0,1)$$
$$(4\pi\mu^2)^{\epsilon}J_1^{4-2\epsilon}(1,1,0)=\frac{1}{\epsilon}+\ln\left(\frac{\bar{\mu}^2}{m_g^2}\right)$$
$$(4\pi\mu^2)^{\epsilon}J_1^{6-2\epsilon}(2,1,0)=-\frac{1}{2}\left[\frac{1}{\epsilon}+\ln\left(\frac{\bar{\mu}^2}{m_g^2}\right)\right]$$
$$J_2^4(1,1,1)=-\frac{1}{2m^2}\ln\left(\frac{m^2}{m_g^2}\right)$$
$$J_1^4(1,1,1)=\frac{1}{2m^2}\left[-\frac{\pi m}{m_g}+1+\ln\left(\frac{m^2}{m_g^2}\right)\right]$$
$$J_1^6(2,1,1)=\frac{1}{4m^2}\left[\frac{\pi m}{m_g}+1-2\ln\left(\frac{m^2}{m_g^2}\right)\right]$$
$$J_1^8(2,2,1)=-\frac{1}{m^2}\left[\frac{\pi m}{12m_g}+\frac{1}{3}+\frac{1}{4}\ln\left(\frac{m^2}{m_g^2}\right)\right]$$
$$J_2^6(2,1,1)=\frac{1}{2m^2},$$
$$J_2^8(3,1,1)=-\frac{1}{12m^2},$$
$$J_2^8(2,2,1)=-\frac{1}{12m^2}$$

$$\mathcal{L}=\mathcal{L}_{2HDM}^{(4)}+\sum_i C_i^{(5)}\mathcal{O}_i^{(5)}+\sum_i C_i^{(6)}\mathcal{O}_i^{(6)}+\mathcal{O}\left(\frac{1}{\Lambda^3}\right)$$

$$\begin{aligned}\mathcal{L}_{2HDM}^{(4)}=&-\frac{1}{4}B_{\mu\nu}B^{\mu\nu}-\frac{1}{4}W_{\mu\nu}^aW^{a\mu\nu}-\frac{1}{4}G_{\mu\nu}^aG^{a\mu\nu}\\&+\left(D_\mu\Phi_1\right)^\dagger D^\mu\Phi_1+\left(D_\mu\Phi_2\right)^\dagger D^\mu\Phi_2+\left(\eta\left(D_\mu\Phi_1\right)^\dagger D^\mu\Phi_2+\text{ h.c.}\right)-V(\Phi_1,\Phi_2)\\&+i\bar{l}_L\not{D}l_L+i\bar{e}_R\not{D}e_R+i\bar{q}_L\not{D}q_L+i\bar{d}_R\not{D}d_R+i\bar{u}_R\not{D}u_R\\&-\left(y_e^{(1)}\bar{l}_Le_R\Phi_1+y_e^{(2)}\bar{l}_Le_R\Phi_2+y_d^{(1)}\bar{q}_Ld_R\Phi_1+y_d^{(2)}\bar{q}_Ld_R\Phi_2\right.\\&\left.+y_u^{(1)}\bar{q}_Lu_R\cdot\Phi_1^\dagger+y_u^{(2)}\bar{q}_Lu_R\cdot\Phi_2^\dagger+\text{ h.c.}\right)\end{aligned}$$

$$\left(D_\mu q\right)_{\alpha i}=\left[\delta_{\alpha\beta}\delta_{ij}\partial_\mu+\frac{ig}{2}\delta_{\alpha\beta}(\tau^a)_{ij}W_\mu^a+\frac{ig_s}{2}(\lambda^a)_{\alpha\beta}\delta_{ij}G_\mu^a+ig'Y_qB_\mu\delta_{\alpha\beta}\delta_{ij}\right]q_{\beta j}$$

$$\begin{aligned}B_{\mu\nu}&=\partial_\mu B_\nu-\partial_\nu B_\mu\\W_{\mu\nu}^a&=\partial_\mu W_\nu^a-\partial_\nu W_\mu^a-g\epsilon^{abc}W_\mu^bW_\nu^c\\G_{\mu\nu}^a&=\partial_\mu G_\nu^a-\partial_\nu G_\mu^a-g_sf^{abc}G_\mu^bG_\nu^c\end{aligned}$$

$$\Phi_1=\begin{pmatrix}\Phi_1^+\\\Phi_1^0\end{pmatrix}=\begin{pmatrix}\phi_1^+\\v_1+\frac{1}{\sqrt{2}}(\rho_1+ia_1)\end{pmatrix},\Phi_2=\begin{pmatrix}\Phi_2^+\\\Phi_2^0\end{pmatrix}=\begin{pmatrix}\phi_2^+\\v_2+\frac{1}{\sqrt{2}}(\rho_2+ia_2)\end{pmatrix},$$

$$V(\Phi_1,\Phi_2)= m_1^2(\Phi_1^\dagger\Phi_1)+m_2^2(\Phi_2^\dagger\Phi_2)+\left(m_{12}^2\Phi_1^\dagger\Phi_2+\text{ h.c. }\right)$$
$$+\frac{1}{2}\lambda_1(\Phi_1^\dagger\Phi_1)^2+\frac{1}{2}\lambda_2(\Phi_2^\dagger\Phi_2)^2+\lambda_3(\Phi_1^\dagger\Phi_1)(\Phi_2^\dagger\Phi_2)+\lambda_4(\Phi_1^\dagger\Phi_2)(\Phi_2^\dagger\Phi_1)$$
$$+\left(\frac{1}{2}\lambda_5(\Phi_1^\dagger\Phi_2)^2+\lambda_6(\Phi_1^\dagger\Phi_1)\Phi_1^\dagger\Phi_2+\lambda_7(\Phi_2^\dagger\Phi_2)\Phi_1^\dagger\Phi_2+\text{ h.c. }\right)$$

$$(\Phi_1,\Phi_2)\rightarrow\left(\frac{\sqrt{\eta^*}\Phi_1+\sqrt{\eta}\Phi_2}{2\sqrt{|\eta|(1+|\eta|)}}\pm\frac{\sqrt{\eta^*}\Phi_1-\sqrt{\eta}\Phi_2}{2\sqrt{|\eta|(1-|\eta|)}}\right)$$

	l_L	e_R	q_L	u_R	d_R	Φ_1	Φ_2
$SU(4)$	**1**	**1**	**3**	**3**	**3**	**1**	**1**
$SU(\mathrm{D})_L$	**2**	**1**	**2**	**1**	**1**	**2**	**2**
$U(4=\mathrm{D})_Y$	$-\frac{1}{2}$	-1	$\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{2}$

$$\mathcal{L}\supset\left(D_\mu\Phi_1\right)^\dagger D^\mu\Phi_1+\left(D_\mu\Phi_2\right)^\dagger D^\mu\Phi_2+\left(\eta\left(D_\mu\Phi_1\right)^\dagger D^\mu\Phi_2+\text{ h.c. }\right)$$
$$+C_{\Phi\partial^2}^{(11)(11)}\partial_\mu(\Phi_1^\dagger\Phi_1)\partial^\mu(\Phi_1^\dagger\Phi_1)+C_{\Phi\partial^2}^{(22)(22)}\partial_\mu(\Phi_2^\dagger\Phi_2)\partial^\mu(\Phi_2^\dagger\Phi_2)$$
$$+C_{\Phi\partial^2}^{(11)(22)}\partial_\mu(\Phi_1^\dagger\Phi_1)\partial^\mu(\Phi_2^\dagger\Phi_2)+C_{\Phi\partial^2}^{(21)(12)}\partial_\mu(\Phi_2^\dagger\Phi_1)\partial^\mu(\Phi_1^\dagger\Phi_2)$$
$$+\left(C_{\Phi\partial^2}^{(21)(21)}\partial_\mu(\Phi_2^\dagger\Phi_1)\partial^\mu(\Phi_2^\dagger\Phi_1)+C_{\Phi\partial^2}^{(21)(11)}\partial_\mu(\Phi_2^\dagger\Phi_1)\partial^\mu(\Phi_1^\dagger\Phi_1)\right.$$
$$\left.+C_{\Phi\partial^2}^{(21)(22)}\partial_\mu(\Phi_2^\dagger\Phi_1)\partial^\mu(\Phi_2^\dagger\Phi_2)+\text{ h.c. }\right)$$
$$+C_{\Phi D}^{(11)(11)}(\Phi_1^\dagger\overleftrightarrow{\mathcal{D}}_\mu\Phi_1)(\Phi_1^\dagger\overleftrightarrow{\mathcal{D}}^\mu\Phi_1)+C_{\Phi D}^{(22)(22)}(\Phi_2^\dagger\overleftrightarrow{\mathcal{D}}_\mu\Phi_2)(\Phi_2^\dagger\overleftrightarrow{\mathcal{D}}^\mu\Phi_2)$$
$$+C_{\Phi D}^{(11)(22)}(\Phi_1^\dagger\overleftrightarrow{\mathcal{D}}_\mu\Phi_1)(\Phi_2^\dagger\overleftrightarrow{\mathcal{D}}^\mu\Phi_2)+C_{\Phi D}^{(21)(12)}(\Phi_2^\dagger\overleftrightarrow{\mathcal{D}}_\mu\Phi_1)(\Phi_1^\dagger\overleftrightarrow{\mathcal{D}}^\mu\Phi_2)$$
$$+\left(C_{\Phi D}^{(21)(21)}(\Phi_2^\dagger\overleftrightarrow{\mathcal{D}}_\mu\Phi_1)(\Phi_2^\dagger\overleftrightarrow{\mathcal{D}}^\mu\Phi_1)+C_{\Phi D}^{(21)(11)}(\Phi_2^\dagger\overleftrightarrow{\mathcal{D}}_\mu\Phi_1)(\Phi_1^\dagger\overleftrightarrow{\mathcal{D}}^\mu\Phi_1)\right.$$
$$\left.+C_{\Phi D}^{(21)(22)}(\Phi_2^\dagger\overleftrightarrow{\mathcal{D}}_\mu\Phi_1)(\Phi_2^\dagger\overleftrightarrow{\mathcal{D}}^\mu\Phi_2)+\text{ h.c. }\right)$$

$$\partial_\mu(\Phi_{1,2}^\dagger\Phi_{1,2})=\Phi_{1,2}^\dagger(D_\mu\Phi_{1,2})+\left(D_\mu\Phi_{1,2}\right)^\dagger\Phi_{1,2}$$

$$\Phi_{1,2}^\dagger\overleftrightarrow{\mathcal{D}}_\mu\Phi_{1,2}\equiv\left(\Phi_{1,2}^\dagger(D_\mu\Phi_{1,2})-\left(D_\mu\Phi_{1,2}\right)^\dagger\Phi_{1,2}\right)$$
$$\Phi_{1,2}^\dagger\overleftrightarrow{\mathcal{D}}_\mu^a\Phi_{1,2}\equiv\left(\Phi_{1,2}^\dagger\tau^a(D_\mu\Phi_{1,2})-\left(D_\mu\Phi_{1,2}\right)^\dagger\tau^a\Phi_{1,2}\right)$$

$$\mathcal{L}\supset\frac{1}{2}\begin{pmatrix}\partial_\mu\rho_1\\ \partial_\mu\rho_2\\ \partial_\mu a_1\\ \partial_\mu a_2\end{pmatrix}^T\begin{pmatrix}1+A_1 & B & J & K\\ B & 1+A_2 & L & N\\ J & L & 1+A_1' & B'\\ K & N & B' & 1+A_2'\end{pmatrix}\begin{pmatrix}\partial^\mu\rho_1\\ \partial^\mu\rho_2\\ \partial^\mu a_1\\ \partial^\mu a_2\end{pmatrix}$$
$$+(\partial_\mu\phi_1^+\partial_\mu\phi_2^+)\begin{pmatrix}1 & \eta^*\\ \eta & 1\end{pmatrix}\begin{pmatrix}\partial^\mu\phi_1^-\\ \partial^\mu\phi_2^-\end{pmatrix}$$

$$A_1 = v^2\left[4c_\beta^2 C_{\Phi\partial^2}^{(11)(11)} + 2s_\beta^2 \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(21)}\right] + s_\beta^2 C_{\Phi\partial^2}^{(21)(12)}\right.$$
$$\left.+4s_\beta c_\beta \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(11)}\right] + 2s_\beta^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right] - s_\beta^2 C_{\Phi D}^{(21)(12)}\right]$$
$$A_2 = v^2\left[4s_\beta^2 C_{\Phi\partial^2}^{(22)(22)} + 2c_\beta^2 \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(21)}\right] + c_\beta^2 C_{\Phi\partial^2}^{(21)(12)}\right.$$
$$\left.+4s_\beta c_\beta \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(22)}\right] + 2c_\beta^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right] - c_\beta^2 C_{\Phi D}^{(21)(12)}\right]$$
$$B = \mathrm{Re}[\eta] + v^2\left[2s_\beta c_\beta C_{\Phi\partial^2}^{(11)(22)} + 2s_\beta c_\beta \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(21)}\right] + s_\beta c_\beta C_{\Phi\partial^2}^{(21)(12)}\right.$$
$$\left.+2c_\beta^2 \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(11)}\right] + 2s_\beta^2 \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(22)}\right] - 2s_\beta c_\beta \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right] + s_\beta c_\beta C_{\Phi D}^{(21)(12)}\right]$$
$$A_1' = v^2\left[4c_\beta^2 C_{\Phi\partial^2}^{(11)(11)} + 2s_\beta^2 \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(21)}\right] + s_\beta^2 C_{\Phi\partial^2}^{(21)(12)} + 4s_\beta c_\beta \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(11)}\right]\right.$$
$$\left.-4c_\beta^2 C_{\Phi D}^{(11)(11)} - 2s_\beta^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right] - s_\beta^2 C_{\Phi D}^{(21)(12)} - 4s_\beta c_\beta \mathrm{Re}\left[C_{\Phi D}^{(21)(11)}\right]\right]$$
$$A_2' = v^2\left[4s_\beta^2 C_{\Phi\partial^2}^{(22)(22)} + 2c_\beta^2 \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(21)}\right] + c_\beta^2 C_{\Phi\partial^2}^{(21)(12)} + 4s_\beta c_\beta \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(22)}\right]\right.$$
$$\left.-4s_\beta^2 C_{\Phi D}^{(22)(22)} - 2c_\beta^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right] - c_\beta^2 C_{\Phi D}^{(21)(12)} - 4s_\beta c_\beta \mathrm{Re}\left[C_{\Phi D}^{(21)(22)}\right]\right]$$
$$B' = \mathrm{Re}[\eta] + v^2\left[2s_\beta c_\beta C_{\Phi\partial^2}^{(11)(22)} + 2s_\beta c_\beta \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(21)}\right]\right.$$
$$+s_\beta c_\beta C_{\Phi\partial^2}^{(21)(12)} + 2c_\beta^2 \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(11)}\right] + 2s_\beta^2 \mathrm{Re}\left[C_{\Phi\partial^2}^{(21)(22)}\right]$$
$$-2s_\beta c_\beta C_{\Phi D}^{(11)(22)} - 2s_\beta c_\beta \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right]$$
$$\left.-s_\beta c_\beta C_{\Phi D}^{(21)(12)} - 2c_\beta^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(11)}\right] - 2s_\beta^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(22)}\right]\right]$$

$$J = -2v^2\left[s_\beta^2 \mathrm{Im}\left[C_{\Phi\partial^2}^{(21)(21)}\right] + s_\beta c_\beta \mathrm{Im}\left[C_{\Phi\partial^2}^{(21)(11)}\right]\right.$$
$$\left.+s_\beta^2 \mathrm{Im}\left[C_{\Phi D}^{(21)(21)}\right] + s_\beta c_\beta \mathrm{Im}\left[C_{\Phi D}^{(21)(11)}\right]\right],$$
$$N = +2v^2\left[c_\beta^2 \mathrm{Im}\left[C_{\Phi\partial^2}^{(21)(21)}\right] + s_\beta c_\beta \mathrm{Im}\left[C_{\Phi\partial^2}^{(21)(22)}\right]\right.$$
$$\left.+c_\beta^2 \mathrm{Im}\left[C_{\Phi D}^{(21)(21)}\right] + s_\beta c_\beta \mathrm{Im}\left[C_{\Phi D}^{(21)(22)}\right]\right],$$
$$K = -\mathrm{Im}[\eta] + 2v^2\left[s_\beta c_\beta \mathrm{Im}\left[C_{\Phi\partial^2}^{(21)(21)}\right] + c_\beta^2 \mathrm{Im}\left[C_{\Phi\partial^2}^{(21)(11)}\right]\right.$$
$$\left.-s_\beta c_\beta \mathrm{Im}\left[C_{\Phi D}^{(21)(21)}\right] - s_\beta^2 \mathrm{Im}\left[C_{\Phi D}^{(21)(22)}\right]\right],$$
$$L = +\mathrm{Im}[\eta] - 2v^2\left[s_\beta c_\beta \mathrm{Im}\left[C_{\Phi\partial^2}^{(21)(21)}\right] + s_\beta^2 \mathrm{Im}\left[C_{\Phi\partial^2}^{(21)(22)}\right]\right.$$
$$\left.-s_\beta c_\beta \mathrm{Im}\left[C_{\Phi D}^{(21)(21)}\right] - c_\beta^2 \mathrm{Im}\left[C_{\Phi D}^{(21)(11)}\right]\right],$$

$$\mathcal{L} \supset \frac{1}{2}\begin{pmatrix}\partial_\mu \vec{\rho} \\ \partial_\mu \vec{a}\end{pmatrix}^T \begin{pmatrix} I_{2\times 2} + A & D \\ D^T & I_{2\times 2} + A'\end{pmatrix}\begin{pmatrix}\partial^\mu \vec{\rho} \\ \partial^\mu \vec{a}\end{pmatrix}$$

$$A \equiv \begin{pmatrix} A_1 & B \\ B & A_2\end{pmatrix}, A' \equiv \begin{pmatrix} A_1' & B' \\ B' & A_2'\end{pmatrix}, D \equiv \begin{pmatrix} J & K \\ L & N\end{pmatrix}$$

$$\vec{\rho} = (\rho_1, \rho_2)^T, \text{ and } \vec{a} = (a_1, a_2)^T$$

$$\vec{\rho} \to \left(I_{2\times 2} - \frac{A}{2}\right)\vec{\rho} - \frac{D}{2}\vec{a}, \vec{a} \to -\frac{D^T}{2}\vec{\rho} + \left(I_{2\times 2} - \frac{A'}{2}\right)\vec{a}$$

$$\vec{\hat{\rho}} = (\hat{\rho}_1, \hat{\rho}_2)^T \text{ and } \vec{\hat{a}} = (\hat{a}_1, \hat{a}_2)^T$$

$$\rho_1 \to \left(1-\frac{A_1}{2}\right)\hat{\rho}_1 - \frac{B}{2}\hat{\rho}_2 - \frac{J}{2}\hat{a}_1 - \frac{K}{2}\hat{a}_2$$
$$\rho_2 \to -\frac{B}{2}\hat{\rho}_1 + \left(1-\frac{A_2}{2}\right)\hat{\rho}_2 - \frac{L}{2}\hat{a}_1 - \frac{N}{2}\hat{a}_2$$
$$a_1 \to -\frac{J}{2}\hat{\rho}_1 - \frac{L}{2}\hat{\rho}_2 + \left(1-\frac{A_1'}{2}\right)\hat{a}_1 - \frac{B'}{2}\hat{a}_2$$
$$a_2 \to -\frac{K}{2}\hat{\rho}_1 - \frac{N}{2}\hat{\rho}_2 - \frac{B'}{2}\hat{a}_1 + \left(1-\frac{A_2'}{2}\right)\hat{a}_2$$

$$\phi_1^+ \to \hat{\phi}_1^+ - \frac{\eta}{2}\hat{\phi}_2^+, \phi_2^+ \to -\frac{\eta^*}{2}\hat{\phi}_1^+ + \hat{\phi}_2^+.$$

$$\begin{aligned}\mathcal{L} \supset & -V(\Phi_1,\Phi_2) + C_\Phi^{(11)(11)(11)}\left(\Phi_1^\dagger\Phi_1\right)^3 + C_\Phi^{(11)(11)(22)}\left(\Phi_1^\dagger\Phi_1\right)^2\left(\Phi_2^\dagger\Phi_2\right) \\ & + C_\Phi^{(11)(22)(22)}\left(\Phi_1^\dagger\Phi_1\right)\left(\Phi_2^\dagger\Phi_2\right)^2 + C_\Phi^{(22)(22)(22)}\left(\Phi_2^\dagger\Phi_2\right)^3 \\ & + C_\Phi^{(11)(21)(12)}\left(\Phi_1^\dagger\Phi_1\right)\left(\Phi_2^\dagger\Phi_1\right)\left(\Phi_1^\dagger\Phi_2\right) + C_\Phi^{(22)(21)(12)}\left(\Phi_2^\dagger\Phi_2\right)\left(\Phi_2^\dagger\Phi_1\right)\left(\Phi_1^\dagger\Phi_2\right) \\ & + \Big(C_\Phi^{(11)(11)(21)}\left(\Phi_1^\dagger\Phi_1\right)^2\left(\Phi_2^\dagger\Phi_1\right) + C_\Phi^{(22)(22)(21)}\left(\Phi_2^\dagger\Phi_2\right)^2\left(\Phi_2^\dagger\Phi_1\right) \\ & + C_\Phi^{(11)(21)(21)}\left(\Phi_1^\dagger\Phi_1\right)\left(\Phi_2^\dagger\Phi_1\right)^2 + C_\Phi^{(22)(21)(21)}\left(\Phi_2^\dagger\Phi_2\right)\left(\Phi_2^\dagger\Phi_1\right)^2 \\ & + C_\Phi^{(21)(21)(21)}\left(\Phi_2^\dagger\Phi_1\right)^3 + C_\Phi^{(21)(21)(12)}\left(\Phi_2^\dagger\Phi_1\right)^2\left(\Phi_1^\dagger\Phi_2\right) \\ & + C_\Phi^{(11)(22)(21)}\left(\Phi_1^\dagger\Phi_1\right)\left(\Phi_2^\dagger\Phi_2\right)\left(\Phi_2^\dagger\Phi_1\right) + \text{ h.c. }\Big),\end{aligned}$$

$$\mathcal{L} \supset -\frac{1}{2}\begin{pmatrix}\vec{\hat{\rho}}\\ \vec{\hat{a}}\end{pmatrix}^T\begin{pmatrix}I_{2\times2}-\frac{A}{2} & -\frac{D}{2}\\ -\frac{D^T}{2} & I_{2\times2}-\frac{A'}{2}\end{pmatrix}\begin{pmatrix}M_{\rho\rho}^2 M_{\rho a}^2\\ M_{a\rho}^2 M_{aa}^2\end{pmatrix}\begin{pmatrix}I_{2\times2}-\frac{A}{2} & -\frac{D}{2}\\ -\frac{D^T}{2} & I_{2\times2}-\frac{A'}{2}\end{pmatrix}\begin{pmatrix}\vec{\hat{\rho}}\\ \vec{\hat{a}}\end{pmatrix}$$

$$\mathcal{L} \supset -\frac{1}{2}\begin{pmatrix}\vec{\hat{\rho}}\\ \vec{\hat{a}}\end{pmatrix}^T\begin{pmatrix}\hat{M}_{\rho\rho}^2 & \hat{M}_{\rho a}^2\\ \left(\hat{M}_{\rho a}^2\right)^T & \hat{M}_{aa}^2\end{pmatrix}\begin{pmatrix}\vec{\hat{\rho}}\\ \vec{\hat{a}}\end{pmatrix}$$
$$\hat{M}_{\rho\rho}^2 = M_{\rho\rho}^2 - \frac{A}{2}M_{\rho\rho}^2 - M_{\rho\rho}^2\frac{A}{2} - M_{\rho a}^2\frac{D^T}{2} - \frac{D}{2}\left(M_{\rho a}^2\right)^T$$
$$\hat{M}_{\rho a}^2 = M_{\rho a}^2 - \frac{D}{2}M_{aa}^2 - M_{\rho\rho}^2\frac{D}{2} - \frac{A}{2}M_{\rho a}^2 - M_{\rho a}^2\frac{A'}{2}$$
$$\hat{M}_{aa}^2 = M_{aa}^2 - \frac{A'}{2}M_{aa}^2 - M_{aa}^2\frac{A'}{2} - \left(M_{\rho a}^2\right)^T\frac{D}{2} - \frac{D^T}{2}M_{\rho a}^2$$

$$\begin{pmatrix}\hat{a}_1\\ \hat{a}_2\end{pmatrix} = \begin{pmatrix}\cos\hat{\beta} & -\sin\hat{\beta}\\ \sin\hat{\beta} & \cos\hat{\beta}\end{pmatrix}\begin{pmatrix}G\\ A\end{pmatrix}$$

$$\mathcal{L} \supset -\frac{1}{2}\begin{pmatrix}G\\ A\\ \hat{\rho}_1\\ \hat{\rho}_2\end{pmatrix}^T\begin{pmatrix}m_G^2 & 0 & 0 & 0\\ 0 & m_A^2 & \hat{M}_{A\rho_1}^2 & \hat{M}_{A\rho_2}^2\\ 0 & \hat{M}_{A\rho_1}^2 & \hat{M}_{\rho_1\rho_1}^2 & \hat{M}_{\rho_1\rho_2}^2\\ 0 & \hat{M}_{A\rho_2}^2 & \hat{M}_{\rho_2\rho_1}^2 & \hat{M}_{\rho_2\rho_2}^2\end{pmatrix}\begin{pmatrix}G\\ A\\ \hat{\rho}_1\\ \hat{\rho}_2\end{pmatrix}$$

$$\begin{aligned}\hat{M}_{A\rho_1}^2 &= -\sin\hat{\beta}\hat{M}_{\rho_1 a_1}^2 + \cos\hat{\beta}\hat{M}_{\rho_1 a_2}^2\\ \hat{M}_{A\rho_2}^2 &= -\sin\hat{\beta}\hat{M}_{\rho_2 a_1}^2 + \cos\hat{\beta}\hat{M}_{\rho_2 a_2}^2\end{aligned}$$

$$\begin{pmatrix} A \\ \hat{\rho}_1 \\ \hat{\rho}_2 \end{pmatrix} = R \begin{pmatrix} h_3 \\ h_2 \\ h_1 \end{pmatrix} = R_x(\hat{\alpha}) R_y(\hat{\xi}) R_z(\hat{\omega}) \begin{pmatrix} h_3 \\ h_2 \\ h_1 \end{pmatrix}$$

$$R_x(\hat{\alpha}) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\hat{\alpha} & -\sin\hat{\alpha} \\ 0 & \sin\hat{\alpha} & \cos\hat{\alpha} \end{pmatrix},$$

$$R_y(\hat{\xi}) = \begin{pmatrix} \cos\hat{\xi} & 0 & \sin\hat{\xi} \\ 0 & 1 & 0 \\ -\sin\hat{\xi} & 0 & \cos\hat{\xi} \end{pmatrix}, \quad R_z(\hat{\omega}) = \begin{pmatrix} \cos\hat{\omega} & -\sin\hat{\omega} & 0 \\ \sin\hat{\omega} & \cos\hat{\omega} & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

$$\begin{pmatrix} A \\ \hat{\rho}_1 \\ \hat{\rho}_2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\hat{\alpha} & -\sin\hat{\alpha} \\ 0 & \sin\hat{\alpha} & \cos\hat{\alpha} \end{pmatrix} \begin{pmatrix} A \\ H \\ h \end{pmatrix}$$

$$\mathcal{L} \supset -\frac{1}{2} \begin{pmatrix} A \\ H \\ h \end{pmatrix}^T \begin{pmatrix} m_A^2 & m_{AH}^2 & m_{Ah}^2 \\ m_{AH}^2 & m_H^2 & 0 \\ m_{Ah}^2 & 0 & m_h^2 \end{pmatrix} \begin{pmatrix} A \\ H \\ h \end{pmatrix}$$

$$\begin{aligned} m_H^2 &= \cos^2\hat{\alpha}\hat{M}_{\rho_1\rho_1}^2 + \sin 2\hat{\alpha}\hat{M}_{\rho_1\rho_2}^2 + \sin^2\hat{\alpha}\hat{M}_{\rho_2\rho_2}^2 \\ m_h^2 &= \sin^2\hat{\alpha}\hat{M}_{\rho_1\rho_1}^2 - \sin 2\hat{\alpha}\hat{M}_{\rho_1\rho_2}^2 + \cos^2\hat{\alpha}\hat{M}_{\rho_2\rho_2}^2 \\ m_{AH}^2 &= \cos\hat{\alpha}\hat{M}_{A\rho_1}^2 + \sin\hat{\alpha}\hat{M}_{A\rho_2}^2 \\ m_{Ah}^2 &= -\sin\hat{\alpha}\hat{M}_{A\rho_1}^2 + \cos\hat{\alpha}\hat{M}_{A\rho_2}^2 \end{aligned}$$

$$\begin{pmatrix} A \\ H \\ h \end{pmatrix} = R_y(\hat{\xi}) R_z(\hat{\omega}) \begin{pmatrix} h_3 \\ h_2 \\ h_1 \end{pmatrix}$$

$$\cos^2\hat{\xi} = \frac{(m_{AH}^2)^2 + \left(m_{Ah}^2\right)^2 + \left(m_A^2 - \lambda^{(2)}\right)\left(m_A^2 + \lambda^{(2)} - \lambda^{(3)} - \lambda^{(1)}\right)}{(\lambda^{(1)} - \lambda^{(2)})(\lambda^{(2)} - \lambda^{(3)})}$$

$$\cos^2\hat{\omega} = \frac{m_A^2 - \lambda^{(2)} + \cos^2\hat{\xi}\left(\lambda^{(2)} - \lambda^{(1)}\right)}{\cos^2\hat{\xi}(\lambda^{(3)} - \lambda^{(1)})}$$

$$\lambda^3 + b\lambda^2 + c\lambda + d = 0$$

$$\begin{aligned} b &= -|b| = -\left(m_A^2 + m_H^2 + m_h^2\right) \\ c &= m_H^2 m_h^2 + m_A^2\left(m_H^2 + m_h^2\right) - (m_{AH}^2)^2 - \left(m_{Ah}^2\right)^2 \\ d &= (m_{AH}^2)^2 m_h^2 + \left(m_{Ah}^2\right)^2 m_H^2 - m_A^2 m_H^2 m_h^2 \end{aligned}$$

CP-conserving theory, $m_{AH}^2 = m_{Ah}^2 = 0$ and $\lambda^{(1,2,3)} = m_{h,H,A}^2$

$$y = \lambda + \frac{b}{3}, p = -|p| = \frac{3c - b^2}{9}, q = -|q| = \frac{2b^3 - 9bc + 27d}{27}.$$

$$\Delta = q^2 + 4p^3 \leq 0$$

$$\lambda^{(1)} \equiv m_{h_1}^2 = \frac{|b|}{3} - \frac{1}{2}\left(\zeta^{1/3} + \sigma^{1/3}\right) + i\frac{\sqrt{3}}{2}\left(\zeta^{1/3} - \sigma^{1/3}\right)$$
$$\lambda^{(2)} \equiv m_{h_2}^2 = \frac{|b|}{3} - \frac{1}{2}\left(\zeta^{1/3} + \sigma^{1/3}\right) - i\frac{\sqrt{3}}{2}\left(\zeta^{1/3} - \sigma^{1/3}\right)$$
$$\lambda^{(3)} \equiv m_{h_3}^2 = \frac{|b|}{3} + \zeta^{1/3} + \sigma^{1/3}$$

$$\zeta \equiv \frac{1}{2}\left(|q| + i\sqrt{4|p|^3 - q^2}\right), \sigma \equiv \frac{1}{2}\left(|q| - i\sqrt{4|p|^3 - q^2}\right).$$

$$\mathcal{L} \supset -\left(\hat{\phi}_1^+ \hat{\phi}_2^+\right)\begin{pmatrix} M_{11}^{\pm 2} + \Delta M_{11}^{\pm 2} & M_{12}^{\pm 2} + \Delta M_{12}^{\pm 2} \\ M_{21}^{\pm 2} + \Delta M_{21}^{\pm 2} & M_{22}^{\pm 2} + \Delta M_{22}^{\pm 2} \end{pmatrix}\begin{pmatrix} \hat{\phi}_1^- \\ \hat{\phi}_2^- \end{pmatrix}$$

$$\beta - \hat{\beta} \sim \mathcal{O}(v^4/\Lambda^2 m_A^2) \text{ and } \beta - \hat{\beta}^{\pm} \sim \mathcal{O}\left(v^4/\Lambda^2 m_{H^{\pm}}^2\right)$$

$$\begin{aligned}\mathcal{L} \supset & -\frac{1}{4}B_{\mu\nu}B^{\mu\nu} - \frac{1}{4}W_{\mu\nu}^a W^{a\mu\nu} - \frac{1}{4}G_{\mu\nu}^a G^{a\mu\nu} \\ & + C_{\Phi B}^{(11)}\left(\Phi_1^\dagger \Phi_1\right)B_{\mu\nu}B^{\mu\nu} + C_{\Phi B}^{(22)}\left(\Phi_2^\dagger \Phi_2\right)B_{\mu\nu}B^{\mu\nu} + \left(C_{\Phi B}^{(21)}\left(\Phi_2^\dagger \Phi_1\right)B_{\mu\nu}B^{\mu\nu} + \text{ h.c. }\right) \\ & + C_{\Phi W}^{(11)}\left(\Phi_1^\dagger \Phi_1\right)W_{\mu\nu}^a W^{a\mu\nu} + C_{\Phi W}^{(22)}\left(\Phi_2^\dagger \Phi_2\right)W_{\mu\nu}^a W^{a\mu\nu} + \left(C_{\Phi W}^{(21)}\left(\Phi_2^\dagger \Phi_1\right)W_{\mu\nu}^a W^{a\mu\nu} + \text{ h.c. }\right) \\ & + C_{\Phi G}^{(11)}\left(\Phi_1^\dagger \Phi_1\right)G_{\mu\nu}^a G^{a\mu\nu} + C_{\Phi G}^{(22)}\left(\Phi_2^\dagger \Phi_2\right)G_{\mu\nu}^a G^{a\mu\nu} + \left(C_{\Phi G}^{(21)}\left(\Phi_2^\dagger \Phi_1\right)G_{\mu\nu}^a G^{a\mu\nu} + \text{ h.c. }\right) \\ & + C_{\Phi WB}^{(11)}\left(\Phi_1^\dagger \tau^a \Phi_1\right)W_{\mu\nu}^a B^{\mu\nu} + C_{\Phi WB}^{(22)}\left(\Phi_2^\dagger \tau^a \Phi_2\right)W_{\mu\nu}^a B^{\mu\nu} + \left(C_{\Phi WB}^{(21)}\left(\Phi_2^\dagger \tau^a \Phi_1\right)W_{\mu\nu}^a B^{\mu\nu} + \text{ h.c. }\right).\end{aligned}$$

$$\begin{aligned}\mathcal{L} \supset & -\frac{1}{4}\left[1 - 4\left(v_1^2 C_{\Phi B}^{(11)} + v_2^2 C_{\Phi B}^{(22)} + 2v_1 v_2 \text{Re}\left[C_{\Phi B}^{(21)}\right]\right)\right]B_{\mu\nu}B^{\mu\nu} \\ & - \frac{1}{4}\left[1 - 4\left(v_1^2 C_{\Phi W}^{(11)} + v_2^2 C_{\Phi W}^{(22)} + 2v_1 v_2 \text{Re}\left[C_{\Phi W}^{(21)}\right]\right)\right]W_{\mu\nu}^a W^{a\mu\nu} \\ & - \frac{1}{4}\left[1 - 4\left(v_1^2 C_{\Phi G}^{(11)} + v_2^2 C_{\Phi G}^{(22)} + 2v_1 v_2 \text{Re}\left[C_{\Phi G}^{(21)}\right]\right)\right]G_{\mu\nu}^a G^{a\mu\nu} \\ & - \left[v_1^2 C_{\Phi WB}^{(11)} + v_2^2 C_{\Phi WB}^{(22)} + 2v_1 v_2 \text{Re}\left[C_{\Phi WB}^{(21)}\right]\right]W_{\mu\nu}^3 B^{\mu\nu}\end{aligned}$$

$$\begin{aligned}\mathcal{L} \supset & -\frac{1}{4}(1 + X_W)\left(W_{\mu\nu}^1 W^{1\mu\nu} + W_{\mu\nu}^2 W^{2\mu\nu}\right) - \frac{1}{4}(1 + X_G)G_{\mu\nu}^a G^{a\mu\nu} \\ & - \frac{1}{4}\left(B_{\mu\nu} W_{\mu\nu}^3\right)\begin{pmatrix} 1 + X_B & X_{WB} \\ X_{WB} & 1 + X_W \end{pmatrix}\begin{pmatrix} B^{\mu\nu} \\ W^{3\mu\nu} \end{pmatrix}\end{aligned}$$

$$\begin{aligned}X_B &= -4\left(v_1^2 C_{\Phi B}^{(11)} + v_2^2 C_{\Phi B}^{(22)} + 2v_1 v_2 \text{Re}\left[C_{\Phi B}^{(21)}\right]\right), \\ X_W &= -4\left(v_1^2 C_{\Phi W}^{(11)} + v_2^2 C_{\Phi W}^{(22)} + 2v_1 v_2 \text{Re}\left[C_{\Phi W}^{(21)}\right]\right), \\ X_G &= -4\left(v_1^2 C_{\Phi G}^{(11)} + v_2^2 C_{\Phi G}^{(22)} + 2v_1 v_2 \text{Re}\left[C_{\Phi G}^{(21)}\right]\right), \\ X_{WB} &= +2\left(v_1^2 C_{\Phi WB}^{(11)} + v_2^2 C_{\Phi WB}^{(22)} + 2v_1 v_2 \text{Re}\left[C_{\Phi WB}^{(21)}\right]\right).\end{aligned}$$

$$B_\mu \to \left(1 - \frac{X_B}{2}\right)\hat{B}_\mu, W_\mu^a \to \left(1 - \frac{X_W}{2}\right)\hat{W}_\mu^a, G_\mu^a \to \left(1 - \frac{X_G}{2}\right)\hat{G}_\mu^a$$

$$Z_A A_\mu^{(a)} \to \hat{A}_\mu^{(a)} \text{ with } Z_A \equiv \sqrt{1 + X_A}, \text{ for } A = B, W, G$$

$$\left(D_\mu q\right)_{\alpha i} \to \left(\hat{D}_\mu q\right)_{\alpha i} = \partial_\mu q_{\alpha i} + \frac{i\hat{g}}{2}(\tau^a)_{ij} q_{\alpha j} \hat{W}_\mu^a + \frac{i\hat{g}_s}{2}(\lambda^a)_{\alpha\beta} q_{\beta i} \hat{G}_\mu^a + i\hat{g}' Y_q \hat{B}_\mu q_{\alpha i}$$

$$\begin{aligned}
\hat{g} &= g\left[1 + 2\left(v_1^2 C_{\Phi W}^{(11)} + v_2^2 C_{\Phi W}^{(22)} + 2v_1 v_2 \mathrm{Re}\left[C_{\Phi W}^{(21)}\right]\right)\right] \\
\hat{g}' &= g'\left[1 + 2\left(v_1^2 C_{\Phi B}^{(11)} + v_2^2 C_{\Phi B}^{(22)} + 2v_1 v_2 \mathrm{Re}\left[C_{\Phi B}^{(21)}\right]\right)\right] \\
\hat{g}_s &= g_s\left[1 + 2\left(v_1^2 C_{\Phi G}^{(11)} + v_2^2 C_{\Phi G}^{(22)} + 2v_1 v_2 \mathrm{Re}\left[C_{\Phi G}^{(21)}\right]\right)\right]
\end{aligned}$$

$$g_A \to Z_{g_A}^{-1} \hat{g}_A = Z_A \hat{g}_A$$

$$\begin{aligned}
\mathcal{L} \supset & \left(D_\mu \Phi_1\right)^\dagger D^\mu \Phi_1 + \left(D_\mu \Phi_2\right)^\dagger D^\mu \Phi_2 + \left(\eta \left(D_\mu \Phi_1\right)^\dagger D^\mu \Phi_2 + \text{ h.c. }\right) \\
& + C_{\Phi D}^{(11)(11)} \left(\Phi_1^\dagger \overleftrightarrow{D}_\mu \Phi_1\right)\left(\Phi_1^\dagger \overleftrightarrow{D}^\mu \Phi_1\right) + C_{\Phi D}^{(22)(22)} \left(\Phi_2^\dagger \overleftrightarrow{D}_\mu \Phi_2\right)\left(\Phi_2^\dagger \overleftrightarrow{D}^\mu \Phi_2\right) \\
& + C_{\Phi D}^{(11)(22)} \left(\Phi_1^\dagger \overleftrightarrow{D}_\mu \Phi_1\right)\left(\Phi_2^\dagger \overleftrightarrow{D} \Phi_2\right) + C_{\Phi D}^{(21)(12)} \left(\Phi_2^\dagger \overleftrightarrow{D}_\mu \Phi_1\right)\left(\Phi_1^\dagger \overleftrightarrow{D}^\mu \Phi_2\right) \\
& + \left(C_{\Phi D}^{(21)(21)} \left(\Phi_2^\dagger \overleftrightarrow{D}_\mu \Phi_1\right)\left(\Phi_2^\dagger \overleftrightarrow{D}^\mu \Phi_1\right) + C_{\Phi D}^{(21)(11)} \left(\Phi_2^\dagger \overleftrightarrow{D}_\mu \Phi_1\right)\left(\Phi_1^\dagger \overleftrightarrow{D}^\mu \Phi_1\right)\right. \\
& \left. + C_{\Phi D}^{(21)(22)} \left(\Phi_2^\dagger \overleftrightarrow{D}_\mu \Phi_1\right)\left(\Phi_2^\dagger \overleftrightarrow{D}^\mu \Phi_2\right) + \text{ h.c. }\right)
\end{aligned}$$

$$\begin{aligned}
\mathcal{L} \supset & \frac{g^2}{4}\left(v^2 + 2v_2 v_2 \mathrm{Re}[\eta]\right)\left(W_\mu^1 W^{1\mu} + W_\mu^2 W^{2\mu}\right) \\
& + \left[v^2 + 2v_1 v_2 \mathrm{Re}[\eta] - \left(v_1^4 C_{\Phi D}^{(11)(11)} + v_2^4 C_{\Phi D}^{(22)(22)} + v_1^2 v_2^3 C_{\Phi D}^{(21)(12)} + v_1^2 v_2^2 C_{\Phi D}^{(11)(2)}\right.\right. \\
& \left.\left. + 2v_1^2 v_2^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right] + 2v_1^3 v_2 \mathrm{Re}\left[C_{\Phi D}^{(21)(11)}\right] + 2v_1 v_2^3 \mathrm{Re}\left[C_{\Phi D}^{(21)(22)}\right]\right)\right] \\
& \times \frac{1}{4}\left(B_\mu W_\mu^3\right)\begin{pmatrix} g'^2 & -gg' \\ -gg' & g^2 \end{pmatrix}\begin{pmatrix} B^\mu \\ W^{3\mu} \end{pmatrix}.
\end{aligned}$$

$$\hat{W}_\mu^\pm \equiv \left(\hat{W}_\mu^1 \mp i\hat{W}_\mu^2\right)/\sqrt{2}$$

$$\begin{aligned}
m_W^2 &= \frac{g^2}{2}\left(v^2 + 2v_1 v_2 \mathrm{Re}[\eta]\right) \times \left[1 + 4\left(v_1^2 C_{\Phi W}^{(11)} + v_2^2 C_{\Phi W}^{(22)} + 2v_1 v_2 \mathrm{Re}\left[C_{\Phi W}^{(21)}\right]\right)\right] \\
&= \frac{\hat{g}^2}{2}\left(v^2 + 2v_1 v_2 \mathrm{Re}[\eta]\right)
\end{aligned}$$

$$\mathcal{L} \supset \frac{1}{2}\left(\hat{B}_\mu \hat{W}_\mu^3\right)\begin{pmatrix} M_{11}^2 + \Delta M_{11}^2 & M_{12}^2 + \Delta M_{12}^2 \\ M_{21}^2 + \Delta M_{21}^2 & M_{22}^2 + \Delta M_{22}^2 \end{pmatrix}\begin{pmatrix} \hat{B}^\mu \\ \hat{W}^{3\mu} \end{pmatrix},$$

$$\begin{aligned}
m_\gamma^2 &= 0 \\
m_Z^2 &= \frac{1}{2}\left(\hat{g}^2 + \hat{g}'^2\right)v^2 \left[\left(1 + X_{WB}\frac{2\hat{g}\hat{g}'}{\hat{g}^2 + \hat{g}'^2}\right)\left(1 + 2s_\beta c_\beta \mathrm{Re}[\eta]\right)\right. \\
& - v^2\left[c_\beta^4 C_{\Phi D}^{(11)(11)} + s_\beta^4 C_{\Phi D}^{(22)(22)} + s_\beta^2 c_\beta^2 C_{\Phi D}^{(21)(12)} + s_\beta^2 c_\beta^2 C_{\Phi D}^{(11)(22)}\right. \\
& \left.\left. + 2s_\beta^2 c_\beta^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right] + 2s_\beta c_\beta^3 \mathrm{Re}\left[C_{\Phi D}^{(21)(11)}\right] + 2s_\beta^3 c_\beta \mathrm{Re}\left[C_{\Phi D}^{(21)(22)}\right]\right]\right]
\end{aligned}$$

$$\begin{pmatrix} \hat{B}_\mu \\ \hat{W}_\mu^3 \end{pmatrix} = \begin{pmatrix} \cos\,\hat{\theta}_W - \frac{X_{WB}}{2}\sin\,\hat{\theta}_W & -\sin\,\hat{\theta}_W - \frac{X_{WB}}{2}\cos\,\hat{\theta}_W \\ \sin\,\hat{\theta}_W - \frac{X_{WB}}{2}\cos\,\hat{\theta}_W & \\ \cos\,\hat{\theta}_W + \frac{X_{WB}}{2}\sin\,\hat{\theta}_W & \end{pmatrix}\begin{pmatrix} A_\mu \\ Z_\mu \end{pmatrix}$$

$$\hat{D}_\mu q = \partial_\mu q + \frac{i\hat{g}}{\sqrt{2}}\left(T^+\hat{W}_\mu^+ + T^-\hat{W}_\mu^-\right)q$$
$$+i\left[\hat{g}\left(\sin\,\hat{\theta}_W - \frac{X_{WB}}{2}\cos\,\hat{\theta}_W\right)T^3 + \hat{g}'\left(\cos\,\hat{\theta}_W - \frac{X_{WB}}{2}\sin\,\hat{\theta}_W\right)Y_q\right]qA_\mu$$
$$+i\left[\hat{g}\left(\cos\,\hat{\theta}_W + \frac{X_{WB}}{2}\sin\,\hat{\theta}_W\right)T^3 - \hat{g}'\left(\sin\,\hat{\theta}_W + \frac{X_{WB}}{2}\cos\,\hat{\theta}_W\right)Y_q\right]qZ_\mu$$
$$\equiv \partial_\mu q + \frac{i\hat{g}}{\sqrt{2}}\left(T^+\hat{W}_\mu^+ + T^-\hat{W}_\mu^-\right)q + i\hat{e}QqA_\mu + i\hat{g}_Z\left[T^3 - \sin^2\,\hat{\theta}_W Q\right]qZ_\mu$$

$$\hat{e} = \hat{g}\left(\sin\,\hat{\theta}_W - \frac{X_{WB}}{2}\cos\,\hat{\theta}_W\right) = \hat{g}'\left(\cos\,\hat{\theta}_W - \frac{X_{WB}}{2}\sin\,\hat{\theta}_W\right)$$

$$\tan\,\hat{\theta}_W = \frac{\hat{g}'}{\hat{g}} + \frac{X_{WB}}{2}\left(1 - \frac{\hat{g}'^2}{\hat{g}^2}\right)$$
$$\sin\,\hat{\theta}_W = \frac{1}{\sqrt{\hat{g}'^2 + \hat{g}^2}}\left(\hat{g}' + \frac{X_{WB}}{2}\frac{\hat{g}(\hat{g}^2 - \hat{g}'^2)}{\hat{g}'^2 + \hat{g}^2}\right)$$
$$\cos\,\hat{\theta}_W = \frac{1}{\sqrt{\hat{g}'^2 + \hat{g}^2}}\left(\hat{g} - \frac{X_{WB}}{2}\frac{\hat{g}'(\hat{g}^2 - \hat{g}'^2)}{\hat{g}'^2 + \hat{g}^2}\right)$$

$$\hat{e} = \frac{\hat{g}\hat{g}'}{\sqrt{\hat{g}'^2 + \hat{g}^2}}\left(1 - X_{WB}\frac{\hat{g}\hat{g}'}{\hat{g}'^2 + \hat{g}^2}\right), \hat{g}_Z = \sqrt{\hat{g}'^2 + \hat{g}^2}\left(1 + X_{WB}\frac{\hat{g}\hat{g}'}{\hat{g}'^2 + \hat{g}^2}\right)$$

gauge-fixing procedure $m_G^2 \to \xi_Z m_Z^2$ and $m_{G^\pm}^2 \to \xi_W m_{W^\pm}^2$, where $\xi_{Z,W}$ are gauge-fixing parameters.

$$\mathcal{L} \supset -\left(y_e^{(1)}\bar{l}_L e_R\Phi_1 + y_e^{(2)}\bar{l}_L e_R\Phi_2 + y_d^{(1)}\bar{q}_L d_R\Phi_1 + y_d^{(2)}\bar{q}_L d_R\Phi_2 + y_u^{(1)}\bar{q}_L u_R\cdot\Phi_1^\dagger + y_u^{(2)}\bar{q}_L u_R\cdot\Phi_2^\dagger\right)$$
$$+C_{\nu\nu\Phi}^{(11)}(\Phi_1\cdot l_L)^T\mathbf{C}(\Phi_1\cdot l_L) + C_{\nu\nu\Phi}^{(22)}(\Phi_2\cdot l_L)^T\mathbf{C}(\Phi_2\cdot l_L) + C_{\nu\nu\Phi}^{(12)}(\Phi_1\cdot l_L)^T\mathbf{C}(\Phi_2\cdot l_L)$$
$$+C_{l\Phi_1}^{(11)}\bar{l}_L e_R\Phi_1(\Phi_1^\dagger\Phi_1) + C_{l\Phi_1}^{(22)}\bar{l}_L e_R\Phi_1(\Phi_2^\dagger\Phi_2) + C_{l\Phi_1}^{(21)}\bar{l}_L e_R\Phi_1(\Phi_2^\dagger\Phi_1) + C_{l\Phi_1}^{(12)}\bar{l}_L e_R\Phi_1(\Phi_1^\dagger\Phi_2)$$
$$+C_{l\Phi_2}^{(22)}\bar{l}_L e_R\Phi_2(\Phi_2^\dagger\Phi_2) + C_{l\Phi_2}^{(11)}\bar{l}_L e_R\Phi_2(\Phi_1^\dagger\Phi_1) + C_{l\Phi_2}^{(21)}\bar{l}_L e_R\Phi_2(\Phi_2^\dagger\Phi_1) + C_{l\Phi_2}^{(12)}\bar{l}_L e_R\Phi_2(\Phi_1^\dagger\Phi_2)$$
$$+C_{d\Phi_1}^{(11)}\bar{q}_L d_R\Phi_1(\Phi_1^\dagger\Phi_1) + C_{d\Phi_1}^{(22)}\bar{q}_L d_R\Phi_1(\Phi_2^\dagger\Phi_2) + C_{d\Phi_1}^{(21)}\bar{q}_L d_R\Phi_1(\Phi_2^\dagger\Phi_1) + C_{d\Phi_1}^{(12)}\bar{q}_L d_R\Phi_1(\Phi_1^\dagger\Phi_2)$$
$$+C_{d\Phi_2}^{(22)}\bar{q}_L d_R\Phi_2(\Phi_2^\dagger\Phi_2) + C_{d\Phi_2}^{(11)}\bar{q}_L d_R\Phi_2(\Phi_1^\dagger\Phi_1) + C_{d\Phi_2}^{(21)}\bar{q}_L d_R\Phi_2(\Phi_2^\dagger\Phi_1) + C_{d\Phi_2}^{(12)}\bar{q}_L d_R\Phi_2(\Phi_1^\dagger\Phi_2)$$
$$+C_{u\Phi_1}^{(11)}\bar{q}_L u_R\cdot\Phi_1^\dagger(\Phi_1^\dagger\Phi_1) + C_{u\Phi_1}^{(22)}\bar{q}_L u_R\cdot\Phi_1^\dagger(\Phi_2^\dagger\Phi_2)$$
$$+C_{u\Phi_1}^{(21)}\bar{q}_L u_R\cdot\Phi_1^\dagger(\Phi_2^\dagger\Phi_1) + C_{u\Phi_1}^{(12)}\bar{q}_L u_R\cdot\Phi_1^\dagger(\Phi_1^\dagger\Phi_2)$$
$$+C_{u\Phi_2}^{(22)}\bar{q}_L u_R\cdot\Phi_2^\dagger(\Phi_2^\dagger\Phi_2) + C_{u\Phi_2}^{(11)}\bar{q}_L u_R\cdot\Phi_2^\dagger(\Phi_1^\dagger\Phi_1)$$
$$+C_{u\Phi_2}^{(21)}\bar{q}_L u_R\cdot\Phi_2^\dagger(\Phi_2^\dagger\Phi_1) + C_{u\Phi_2}^{(12)}\bar{q}_L u_R\cdot\Phi_2^\dagger(\Phi_1^\dagger\Phi_2) + \text{ h.c.}.$$

$$\mathcal{L} \supset -\bar{e}_{L,a}(M_e)_{ab}e_{R,b} - \bar{d}_{L,a}(M_d)_{ab}d_{R,b} - \bar{u}_{L,a}(M_u)_{ab}u_{R,b} - \frac{1}{2}\nu_{L,a}^T(M_\nu)_{ab}\mathbf{C}\nu_{L,b} + h.c.$$

$$(M_e)_{ab} = v\left(y_{e,a}^{(1)}\cos\,\beta + y_{e,a}^{(2)}\sin\,\beta\right)\delta_{ab}$$
$$-v^3\left(\cos^3\,\beta\left(C_{l\Phi_1}^{(11)}\right)_{ab} + \sin^2\,\beta\cos\,\beta\left(C_{l\Phi_1}^{(22)}\right)_{ab} + \sin\,\beta\cos^2\,\beta\left(\left(C_{l\Phi_1}^{(12)}\right)_{ab} + \left(C_{l\Phi_1}^{(21)}\right)_{ab}\right)\right.$$
$$\left.+\sin^3\,\beta\left(C_{l\Phi_2}^{(22)}\right)_{ab} + \sin\,\beta\cos^2\,\beta\left(C_{l\Phi_2}^{(11)}\right)_{ab} + \sin^2\,\beta\cos\,\beta\left(\left(C_{l\Phi_2}^{(12)}\right)_{ab} + \left(C_{l\Phi_2}^{(21)}\right)_{ab}\right)\right)$$

$$(M_d)_{ab}= v\left(y_{d,a}^{(1)}\cos\,\beta + y_{d,a}^{(2)}\sin\,\beta\right)\delta_{ab}$$
$$-v^3\left(\cos^3\,\beta\left(C_{d\Phi_1}^{(11)}\right)_{ab} + \sin^2\,\beta\cos\,\beta\left(C_{d\Phi_1}^{(22)}\right)_{ab} + \sin\,\beta\cos^2\,\beta\left(\left(C_{d\Phi_1}^{(12)}\right)_{ab} + \left(C_{d\Phi_1}^{(21)}\right)_{ab}\right)\right.$$
$$\left.+\sin^3\,\beta\left(C_{d\Phi_2}^{(22)}\right)_{ab} + \sin\,\beta\cos^2\,\beta\left(C_{d\Phi_2}^{(11)}\right)_{ab} + \sin^2\,\beta\cos\,\beta\left(\left(C_{d\Phi_2}^{(12)}\right)_{ab} + \left(C_{d\Phi_2}^{(21)}\right)_{ab}\right)\right)$$

$$(M_u)_{ab}= v\left(y_{u,a}^{(1)}\cos\,\beta + y_{u,a}^{(2)}\sin\,\beta\right)\delta_{ab}$$
$$-v^3\left(\cos^3\,\beta\left(C_{u\Phi_1}^{(11)}\right)_{ab} + \sin^2\,\beta\cos\,\beta\left(C_{u\Phi_1}^{(22)}\right)_{ab} + \sin\,\beta\cos^2\,\beta\left(\left(C_{u\Phi_1}^{(12)}\right)_{ab} + \left(C_{u\Phi_1}^{(21)}\right)_{ab}\right)\right.$$
$$\left.+\sin^3\,\beta\left(C_{u\Phi_2}^{(22)}\right)_{ab} + \sin\,\beta\cos^2\,\beta\left(C_{u\Phi_2}^{(11)}\right)_{ab} + \sin^2\,\beta\cos\,\beta\left(\left(C_{u\Phi_2}^{(12)}\right)_{ab} + \left(C_{u\Phi_2}^{(21)}\right)_{ab}\right)\right),$$

$$(M_\nu)_{ab} = -2v^2\left(\cos^2\,\beta\left(C_{\nu\nu\Phi}^{(11)}\right)_{ab} + \sin^2\,\beta\left(C_{\nu\nu\Phi}^{(22)}\right)_{ab} + \sin\,\beta\cos\,\beta\left(C_{\nu\nu\Phi}^{(12)}\right)_{ab}\right).$$

$$\bar{\nu}_L^c = \nu_L^T\mathbf{C} - \bar{\nu}_L^c M_\nu \nu_L/2 + \text{h.c}$$


bi-unitary transformations $\psi_L \to U_L\psi_L$ and $\psi_R \to U_R\psi_R$

$$\hat{M}_e = U_L^{e\dagger}M_e U_R^e = \text{diag}(m_e, m_\mu, m_\tau)$$
$$\hat{M}_d = U_L^{d\dagger}M_d U_R^d = \text{diag}(m_d, m_s, m_b)$$
$$\hat{M}_u = U_L^{u\dagger}M_u U_R^u = \text{diag}(m_u, m_c, m_t)$$
$$\hat{M}_\nu = U_L^{\nu T}M_\nu U_L^\nu = \text{diag}(m_{\nu_1}, m_{\nu_2}, m_{\nu_3})$$


$$\left(\hat{y}_{(e,d,u)}^{(1,2)}\right)_{ab} = \left(U_L^{(e,d,u)\dagger}y_{(e,d,u)}^{(1,2)}U_R^{(e,d,u)}\right)_{ab}$$
$$\left(\hat{C}_{\nu\nu\Phi}^{(11,22,12)}\right)_{ab} = \left(U_L^{\nu T}C_{\nu\nu\Phi}^{(11,22,12)}U_L^\nu\right)_{ab}$$
$$\left(\hat{C}_{(l,d,u)\Phi_{(1,2)}}^{(11,22,12,21)}\right)_{ab} = \left(U_L^{(e,d,u)\dagger}C_{(l,d,u)\Phi_{(1,2)}}^{(11,22,12,21)}U_R^{(e,d,u)}\right)_{ab}$$
$$\left(\hat{C}_{l(B,W)\Phi_{(1,2)}}\right)_{ab} = \left(U_L^{e\dagger}C_{l(B,W)\Phi_{(1,2)}}U_R^e\right)_{ab}$$
$$\left(\hat{C}_{d(B,W,G)\Phi_{(1,2)}}\right)_{ab} = \left(U_L^{d\dagger}C_{d(B,W,G)\Phi_{(1,2)}}U_R^d\right)_{ab}$$
$$\left(\hat{C}_{u(B,W,G)\Phi_{(1,2)}}\right)_{ab} = \left(U_L^{u\dagger}C_{u(B,W,G)\Phi_{(1,2)}}U_R^u\right)_{ab}$$
$$\left(\hat{C}_{\Phi(e,d,u)}^{(11,22,12)}\right)_{ab} = \left(U_R^{(e,d,u)\dagger}C_{\Phi(e,d,u)}^{(11,22,12)}U_R^{(e,d,u)}\right)_{ab}$$
$$\left(\hat{C}_{\Phi(l,q)}^{(11,22,12)[1]}\right)_{ab} = \left(U_L^{(e,d)\dagger}C_{\Phi(l,q)}^{(11,22,12)[1]}U_L^{(e,d)}\right)_{ab}$$
$$\left(\hat{C}_{\Phi(l,q)}^{(11,22,12)[3]}\right)_{ab} = \left(U_L^{(e,d)\dagger}C_{\Phi(l,q)}^{(11,22,12)[3]}U_L^{(e,d)}\right)_{ab}$$
$$\left(\hat{C}_{\Phi ud}^{(11,22,21)}\right)_{ab} = \left(U_R^{u\dagger}C_{\Phi ud}^{(11,22,21)}U_R^d\right)_{ab}$$

$$V \equiv U_L^{u\dagger}U_L^d, U \equiv U_L^{e\dagger}U_L^\nu$$

Model	u	d	e	l_L	e_R	q_L	u_R	d_R	Φ_1	Φ_2
Type-I	Φ_2	Φ_2	Φ_2	+	+	+	+	+	-	+
Type-II	Φ_2	Φ_1	Φ_1	+	-	+	+	-	-	+
Type-X (Lepton-specific)	Φ_2	Φ_2	Φ_1	+	-	+	+	+	-	+
Type-Y (Flipped)	Φ_2	Φ_1	Φ_2	+	+	+	+	-	-	+



$$
\begin{aligned}
&+\frac{i}{2\sqrt{2}}\Big((s_{\hat{\alpha}}\,(2-A_1)+c_{\hat{\alpha}}B)\left(\mathcal{P}_R\hat{y}^{(1)}_{e,f_1f_2}+\mathcal{P}_L\hat{y}^{(1)*}_{e,f_2f_1}\right)\\
&\qquad-(c_{\hat{\alpha}}\,(2-A_2)+s_{\hat{\alpha}}B)\left(\mathcal{P}_R\hat{y}^{(2)}_{e,f_1f_2}+\mathcal{P}_L\hat{y}^{(2)*}_{e,f_2f_1}\right)\Big)\\
&+\frac{iv^2}{\sqrt{2}}\Big(-3s_{\hat{\alpha}}c^2_{\beta}(\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}+\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1})\\
&\qquad+(2c_{\hat{\alpha}}s_{\beta}c_{\beta}-s_{\hat{\alpha}}s^2_{\beta})(\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}+\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1})\\
&\qquad-(2s_{\hat{\alpha}}s_{\beta}c_{\beta}-c_{\hat{\alpha}}c^2_{\beta})(\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_1,f_1f_2}+\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_1,f_2f_1})\\
&\qquad-(2s_{\hat{\alpha}}s_{\beta}c_{\beta}-c_{\hat{\alpha}}c^2_{\beta})(\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_1,f_1f_2}+\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_1,f_2f_1})\\
&\qquad+3c_{\hat{\alpha}}s^2_{\beta}(\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_2,f_1f_2}+\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_2,f_2f_1})\\
&\qquad-(2s_{\hat{\alpha}}s_{\beta}c_{\beta}-c_{\hat{\alpha}}c^2_{\beta})(\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_2,f_1f_2}+\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_2,f_2f_1})\\
&\qquad+(2c_{\hat{\alpha}}s_{\beta}c_{\beta}-s_{\hat{\alpha}}s^2_{\beta})(\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}+\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1})\\
&\qquad+(2c_{\hat{\alpha}}s_{\beta}c_{\beta}-s_{\hat{\alpha}}s^2_{\beta})(\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}+\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1})\Big)\\
&+\frac{iv}{\sqrt{2}}\not{p}_3\mathcal{P}_Rc_{\hat{\alpha}-\beta}\left(\hat{C}^{(12)}_{\Phi e,f_1f_2}-\hat{C}^{(12)*}_{\Phi e,f_2f_1}\right)\\
&+\frac{iv}{\sqrt{2}}\not{p}_3\mathcal{P}_Lc_{\hat{\alpha}-\beta}\left(\hat{C}^{(12)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(12)[1]*}_{\Phi l,f_2f_1}+\hat{C}^{(12)[3]}_{\Phi l,f_1f_2}-\hat{C}^{(12)[3]*}_{\Phi l,f_2f_1}\right),
\end{aligned}
$$



$$
\begin{aligned}
&+\frac{i\delta_{f_1f_2}m_{e_{f_1}}}{2\sqrt{2}v}\,(A_1+Bt_{\beta}-2)\\
&+i\sqrt{2}v^2c_{\beta}\Big(c^2_{\beta}\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\\
&\qquad+s^2_{\beta}\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\qquad+s^2_{\beta}\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\qquad+s^2_{\beta}\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big)
\end{aligned}
$$

canonical normalization $\beta-\hat{\beta}\sim\mathcal{O}(v^4/\Lambda^2m_A^2)$ and $\beta-\hat{\beta}^{\pm}\sim\mathcal{O}\left(v^4/\Lambda^2m_{H^{\pm}}^2\right)$

$$\hat{M}^2_{\rho_1\rho_1} \equiv M^2_{\rho_1\rho_1} + \Delta M^2_{\rho_1\rho_1}$$
$$\hat{M}^2_{\rho_2\rho_2} \equiv M^2_{\rho_2\rho_2} + \Delta M^2_{\rho_2\rho_2}$$
$$\hat{M}^2_{\rho_1\rho_2} = \hat{M}^2_{\rho_2\rho_1} \equiv M^2_{\rho_1\rho_2} + \Delta M^2_{\rho_1\rho_2}$$
$$\hat{M}^2_{a_1a_1} \equiv M^2_{a_1a_1} + \Delta M^2_{a_1a_1}$$
$$\hat{M}^2_{a_2a_2} \equiv M^2_{a_2a_2} + \Delta M^2_{a_2a_2}$$
$$\hat{M}^2_{a_1a_2} = \hat{M}^2_{a_2a_1} \equiv M^2_{a_1a_2} + \Delta M^2_{a_1a_2}$$

$$\hat{M}^2_{\rho_1a_1} \equiv M^2_{\rho_1a_1} + \Delta M^2_{\rho_1a_1}$$
$$\hat{M}^2_{\rho_2a_2} \equiv M^2_{\rho_2a_2} + \Delta M^2_{\rho_2a_2}$$
$$\hat{M}^2_{\rho_1a_2} \equiv M^2_{\rho_1a_2} + \Delta M^2_{\rho_1a_2}$$
$$\hat{M}^2_{\rho_2a_1} \equiv M^2_{\rho_2a_1} + \Delta M^2_{\rho_2a_1}$$

$$M^2_{\rho_1\rho_1} = m_1^2 + 3\lambda_1 v_1^2 + v_2^2(\lambda_3 + \lambda_4 + \mathrm{Re}[\lambda_5]) + 6v_1v_2\mathrm{Re}[\lambda_6]$$
$$M^2_{\rho_2\rho_2} = m_2^2 + 3\lambda_2 v_2^2 + v_1^2(\lambda_3 + \lambda_4 + \mathrm{Re}[\lambda_5]) + 6v_1v_2\mathrm{Re}[\lambda_7]$$
$$M^2_{\rho_1\rho_2} = M^2_{\rho_2\rho_1} = \mathrm{Re}[m_{12}^2] + 2v_1v_2(\lambda_3 + \lambda_4 + \mathrm{Re}[\lambda_5]) + 3v_1^2\mathrm{Re}[\lambda_6] + 3v_2^2\mathrm{Re}[\lambda_7]$$
$$M^2_{a_1a_1} = m_1^2 + \lambda_1 v_1^2 + v_2^2(\lambda_3 + \lambda_4 - \mathrm{Re}[\lambda_5]) + 2v_1v_2\mathrm{Re}[\lambda_6]$$
$$M^2_{a_2a_2} = m_2^2 + \lambda_2 v_2^2 + v_1^2(\lambda_3 + \lambda_4 - \mathrm{Re}[\lambda_5]) + 2v_1v_2\mathrm{Re}[\lambda_7]$$
$$M^2_{a_1a_2} = M^2_{a_2a_1} = \mathrm{Re}[m_{12}^2] + 2v_1v_2\mathrm{Re}[\lambda_5] + v_1^2\mathrm{Re}[\lambda_6] + v_2^2\mathrm{Re}[\lambda_7]$$

$$M^2_{\rho_1a_1} = +v_2^2\mathrm{Im}[\lambda_5] + 2v_1v_2\mathrm{Im}[\lambda_6]$$
$$M^2_{\rho_2a_2} = -v_1^2\mathrm{Im}[\lambda_5] - 2v_1v_2\mathrm{Im}[\lambda_7]$$
$$M^2_{\rho_1a_2} = -\mathrm{Im}[m_{12}^2] - 2v_1v_2\mathrm{Im}[\lambda_5] - 3v_1^2\mathrm{Im}[\lambda_6] - v_2^2\mathrm{Im}[\lambda_7]$$
$$M^2_{\rho_2a_1} = +\mathrm{Im}[m_{12}^2] + 2v_1v_2\mathrm{Im}[\lambda_5] + v_1^2\mathrm{Im}[\lambda_6] + 3v_2^2\mathrm{Im}[\lambda_7]$$

$$\Delta M^2_{\rho_1\rho_1} = -A_1M^2_{\rho_1\rho_1} - BM^2_{\rho_1\rho_2} - JM^2_{\rho_1a_1} - KM^2_{\rho_1a_2}$$
$$-v^4\left[15c_\beta^4 C_\Phi^{(11)(11)(11)} + 6s_\beta^2c_\beta^2 C_\Phi^{(11)(11)(22)} + s_\beta^4 C_\Phi^{(11)(22)(22)} + 20s_\beta c_\beta^3\mathrm{Re}\left[C_\Phi^{(11)(11)(21)}\right]\right.$$
$$+12s_\beta^2c_\beta^2\mathrm{Re}\left[C_\Phi^{(11)(21)(21)}\right] + 2s_\beta^4\mathrm{Re}\left[C_\Phi^{(22)(21)(21)}\right] + 6s_\beta^2c_\beta^2 C_\Phi^{(11)(21)(12)} + s_\beta^4 C_\Phi^{(22)(21)(12)}$$
$$\left.+6s_\beta^3c_\beta\mathrm{Re}\left[C_\Phi^{(21)(21)(21)}\right] + 6s_\beta^3c_\beta\mathrm{Re}\left[C_\Phi^{(21)(21)(12)}\right] + 6s_\beta^3c_\beta\mathrm{Re}\left[C_\Phi^{(11)(22)(21)}\right]\right]$$
$$\Delta M^2_{\rho_2\rho_2} = -A_2M^2_{\rho_2\rho_2} - BM^2_{\rho_1\rho_2} - LM^2_{\rho_2a_1} - NM^2_{\rho_2a_2}$$
$$-v^4\left[15s_\beta^4 C_\Phi^{(22)(22)(22)} + 6s_\beta^2c_\beta^2 C_\Phi^{(11)(22)(22)} + c_\beta^4 C_\Phi^{(11)(11)(22)} + 20s_\beta^3c_\beta\mathrm{Re}\left[C_\Phi^{(22)(22)(21)}\right]\right.$$
$$+12s_\beta^2c_\beta^2\mathrm{Re}\left[C_\Phi^{(22)(21)(21)}\right] + 2c_\beta^4\mathrm{Re}\left[C_\Phi^{(11)(21)(21)}\right] + 6s_\beta^2c_\beta^2 C_\Phi^{(22)(21)(12)} + c_\beta^4 C_\Phi^{(11)(21)(12)}$$
$$\left.+6s_\beta c_\beta^3\mathrm{Re}\left[C_\Phi^{(21)(21)(21)}\right] + 6s_\beta c_\beta^3\mathrm{Re}\left[C_\Phi^{(21)(21)(12)}\right] + 6s_\beta c_\beta^3\mathrm{Re}\left[C_\Phi^{(11)(22)(21)}\right]\right]$$
$$\Delta M^2_{\rho_1\rho_2} = \Delta M^2_{\rho_2\rho_1} = -\frac{B}{2}\left(M^2_{\rho_1\rho_1} + M^2_{\rho_2\rho_2}\right) - \frac{1}{2}(A_1 + A_2)M^2_{\rho_1\rho_2}$$
$$-\frac{1}{2}\left(JM^2_{\rho_2a_1} + KM^2_{\rho_2a_2} + LM^2_{\rho_1}a_1 + NM^2_{\rho_1a_2}\right)$$
$$-v^4\left[4s_\beta c_\beta^3 C_\Phi^{(11)(11)(22)} + 4s_\beta^3c_\beta C_\Phi^{(11)(22)(22)} + 5c_\beta^4\mathrm{Re}\left[C_\Phi^{(11)(11)(21)}\right] + 5s_\beta^4\mathrm{Re}\left[C_\Phi^{(22)(22)(21)}\right]\right.$$
$$+8s_\beta c_\beta^3\mathrm{Re}\left[C_\Phi^{(11)(21)(21)}\right] + 8s_\beta^3c_\beta\mathrm{Re}\left[C_\Phi^{(22)(21)(21)}\right] + 4s_\beta c_\beta^3 C_\Phi^{(11)(21)(12)} + 4s_\beta^3c_\beta C_\Phi^{(22)(21)(12)}$$
$$\left.+9s_\beta^2c_\beta^2\mathrm{Re}\left[C_\Phi^{(21)(21)(21)}\right] + 9s_\beta^2c_\beta^2\mathrm{Re}\left[C_\Phi^{(21)(21)(12)}\right] + 9s_\beta^2c_\beta^2\mathrm{Re}\left[C_\Phi^{(11)(22)(21)}\right]\right]$$

$$\Delta M^2_{a_1a_1} = -A'_1 M^2_{a_1a_1} - B' M^2_{a_1a_2} - J M^2_{\rho_1 a_1} - L M^2_{\rho_2 a_1}$$
$$-v^4 \left[3c_\beta^4 C_\Phi^{(11)(11)(11)} + 2s_\beta^2 c_\beta^2 C_\Phi^{(11)(11)(22)} + s_\beta^4 C_\Phi^{(11)(22)(22)} + 4s_\beta c_\beta^3 \mathrm{Re}\left[C_\Phi^{(11)(11)(21)}\right]\right.$$
$$-2s_\beta^4 \mathrm{Re}\left[C_\Phi^{(22)(21)(21)}\right] + 2s_\beta^2 c_\beta^2 C_\Phi^{(11)(21)(12)} + s_\beta^4 C_\Phi^{(22)(21)(12)}$$
$$\left.-6s_\beta^3 c_\beta \mathrm{Re}\left[C_\Phi^{(21)(21)(21)}\right] + 2s_\beta^3 c_\beta \mathrm{Re}\left[C_\Phi^{(21)(21)(12)}\right] + 2s_\beta^3 c_\beta \mathrm{Re}\left[C_\Phi^{(11)(22)(21)}\right]\right]$$
$$\Delta M^2_{a_2a_2} = -A'_2 M^2_{a_2a_2} - B' M^2_{a_1a_2} - K M^2_{\rho_1 a_2} - N M^2_{\rho_2 a_2}$$
$$-v^4 \left[3s_\beta^4 C_\Phi^{(22)(22)(22)} + 2s_\beta^2 c_\beta^2 C_\Phi^{(11)(22)(22)} + c_\beta^4 C_\Phi^{(11)(11)(22)} + 4s_\beta^3 c_\beta \mathrm{Re}\left[C_\Phi^{(22)(22)(21)}\right]\right.$$
$$-2c_\beta^4 \mathrm{Re}\left[C_\Phi^{(11)(21)(21)}\right] + 2s_\beta^2 c_\beta^2 C_\Phi^{(22)(21)(12)} + c_\beta^4 C_\Phi^{(11)(21)(12)}$$
$$\left.-6s_\beta c_\beta^3 \mathrm{Re}\left[C_\Phi^{(21)(21)(21)}\right] + 2s_\beta c_\beta^3 \mathrm{Re}\left[C_\Phi^{(21)(21)(12)}\right] + 2s_\beta c_\beta^3 \mathrm{Re}\left[C_\Phi^{(11)(22)(21)}\right]\right]$$
$$\Delta M^2_{a_1a_2} = \Delta M^2_{a_2a_1} = -\frac{B'}{2}\left(M^2_{a_1a_1} + M^2_{a_2a_2}\right) - \frac{1}{2}(A'_1 + A'_2) M^2_{a_1a_2}$$
$$-\frac{1}{2}\left(J M^2_{\rho_1 a_2} + K M^2_{\rho_1 a_1} + L M^2_{\rho_2 a_2} + N M^2_{\rho_2 a_1}\right)$$
$$-v^4 \left[c_\beta^4 \mathrm{Re}\left[C_\Phi^{(11)(11)(21)}\right] + s_\beta^4 \mathrm{Re}\left[C_\Phi^{(22)(22)(21)}\right] + 4s_\beta c_\beta^3 \mathrm{Re}\left[C_\Phi^{(11)(21)(21)}\right]\right.$$
$$+4s_\beta^3 c_\beta \mathrm{Re}\left[C_\Phi^{(22)(21)(21)}\right] + 9s_\beta^2 c_\beta^2 \mathrm{Re}\left[C_\Phi^{(21)(21)(21)}\right] + s_\beta^2 c_\beta^2 \mathrm{Re}\left[C_\Phi^{(21)(21)(12)}\right]$$
$$\left.+s_\beta^2 c_\beta^2 \mathrm{Re}\left[C_\Phi^{(11)(22)(21)}\right]\right]$$

$$\Delta M^2_{\rho_1 a_1} = -\frac{1}{2}\left(JM^2_{a_1 a_1} + KM^2_{a_2 a_1} + JM^2_{\rho_1 \rho_1} + LM^2_{\rho_1 \rho_2}\right)$$
$$-\frac{1}{2}\left(A_1 M^2_{\rho_1 a_1} + BM^2_{\rho_2 a_1} + A_1' M^2_{\rho_1 a_1} + B' M^2_{\rho_1 a_2}\right)$$
$$-4v^4\left[2s_\beta c_\beta^3 \mathrm{Im}\left[C_\Phi^{(11)(11)(21)}\right] + 3s_\beta^2 c_\beta^2 \mathrm{Im}\left[C_\Phi^{(11)(21)(21)}\right] + s_\beta^4 \mathrm{Im}\left[C_\Phi^{(22)(21)(21)}\right]\right.$$
$$\left.+3s_\beta^3 c_\beta \mathrm{Im}\left[C_\Phi^{(21)(21)(21)}\right] + s_\beta^3 c_\beta \mathrm{Im}\left[C_\Phi^{(21)(21)(12)}\right] + s_\beta^3 c_\beta \mathrm{Im}\left[C_\Phi^{(11)(22)(21)}\right]\right]$$

$$\Delta M^2_{\rho_2 a_2} = -\frac{1}{2}\left(LM^2_{a_1 a_2} + NM^2_{a_2 a_2} + NM^2_{\rho_2 \rho_2} + KM^2_{\rho_2 \rho_1}\right)$$
$$-\frac{1}{2}\left(BM^2_{\rho_1 a_2} + A_2 M^2_{\rho_2 a_2} + B' M^2_{\rho_2 a_1} + A_2' M^2_{\rho_2 a_2}\right)$$
$$+4v^4\left[2s_\beta^3 c_\beta \mathrm{Im}\left[C_\Phi^{(22)(22)(21)}\right] + c_\beta^4 \mathrm{Im}\left[C_\Phi^{(11)(21)(21)}\right] + 3s_\beta^2 c_\beta^2 \mathrm{Im}\left[C_\Phi^{(22)(21)(21)}\right]\right.$$
$$\left.+3s_\beta c_\beta^3 \mathrm{Im}\left[C_\Phi^{(21)(21)(21)}\right] + s_\beta c_\beta^3 \mathrm{Im}\left[C_\Phi^{(21)(21)(12)}\right] + s_\beta c_\beta^3 \mathrm{Im}\left[C_\Phi^{(11)(22)(21)}\right]\right]$$

$$\Delta M^2_{\rho_1 a_2} = -\frac{1}{2}\left(JM^2_{a_1 a_2} + KM^2_{a_2 a_2} + KM^2_{\rho_1 \rho_1} + NM^2_{\rho_1 \rho_2}\right)$$
$$-\frac{1}{2}\left(A_1 M^2_{\rho_1 a_2} + BM^2_{\rho_2 a_2} + B' M^2_{\rho_1 a_1} + A_2' M^2_{\rho_1 a_2}\right)$$
$$+2v^4\left[5c_\beta^4 \mathrm{Im}\left[C_\Phi^{(11)(11)(21)}\right] + s_\beta^4 \mathrm{Im}\left[C_\Phi^{(22)(22)(21)}\right] + 8s_\beta c_\beta^3 \mathrm{Im}\left[C_\Phi^{(11)(21)(21)}\right]\right.$$
$$+4s_\beta^3 c_\beta \mathrm{Im}\left[C_\Phi^{(22)(21)(21)}\right] + 9s_\beta^2 c_\beta^2 \mathrm{Im}\left[C_\Phi^{(21)(21)(21)}\right] + 3s_\beta^2 c_\beta^2 \mathrm{Im}\left[C_\Phi^{(21)(21)(12)}\right]$$
$$\left.+3s_\beta^2 c_\beta^2 \mathrm{Im}\left[C_\Phi^{(11)(22)(21)}\right]\right]$$

$$\Delta M^2_{\rho_2 a_1} = -\frac{1}{2}\left(LM^2_{a_1 a_1} + NM^2_{a_2 a_1} + JM^2_{\rho_2 \rho_1} + LM^2_{\rho_2 \rho_2}\right)$$
$$-\frac{1}{2}\left(BM^2_{\rho_1 a_1} + A_2 M^2_{\rho_2 a_1} + A_1' M^2_{\rho_2 a_1} + B' M^2_{\rho_2 a_2}\right)$$
$$-2v^4\left[c_\beta^4 \mathrm{Im}\left[C_\Phi^{(11)(11)(21)}\right] + 5s_\beta^4 \mathrm{Im}\left[C_\Phi^{(22)(22)(21)}\right] + 4s_\beta c_\beta^3 \mathrm{Im}\left[C_\Phi^{(11)(21)(21)}\right]\right.$$
$$+8s_\beta^3 c_\beta \mathrm{Im}\left[C_\Phi^{(22)(21)(21)}\right] + 9s_\beta^2 c_\beta^2 \mathrm{Im}\left[C_\Phi^{(21)(21)(21)}\right] + 3s_\beta^2 c_\beta^2 \mathrm{Im}\left[C_\Phi^{(21)(21)(12)}\right]$$
$$\left.+3s_\beta^2 c_\beta^2 \mathrm{Im}\left[C_\Phi^{(11)(22)(21)}\right]\right]$$

$$M_{11}^{\pm 2} = m_1^2 + \lambda_1 v_1^2 + v_2^2 \lambda_3 + 2v_1 v_2 \mathrm{Re}[\lambda_6]$$
$$M_{22}^{\pm 2} = m_2^2 + \lambda_2 v_2^2 + v_1^2 \lambda_3 + 2v_1 v_2 \mathrm{Re}[\lambda_7]$$
$$M_{12}^{\pm 2} = \left(M_{21}^{\pm 2}\right)^* = (m_{12}^2)^* + v_1 v_2(\lambda_4 + \lambda_5^*) + v_1^2 \lambda_6^* + v_2^2 \lambda_7^*$$

$$\Delta M_{11}^{\pm 2} = -\mathrm{Re}\left[\eta M_{12}^{\pm 2}\right] - v^4\Big[3c_\beta^4 C_\Phi^{(11)(11)(11)} + 2s_\beta^2 c_\beta^2 C_\Phi^{(11)(11)(22)} + s_\beta^4 C_\Phi^{(11)(22)(22)} + 4s_\beta c_\beta^3 \mathrm{Re}\left[C_\Phi^{(11)(11)(21)}\right] + 2s_\beta^2 c_\beta^2 \mathrm{Re}\left[C_\Phi^{(11)(21)(21)}\right] + s_\beta^2 c_\beta^2 C_\Phi^{(11)(21)(12)} + 2s_\beta^3 c_\beta \mathrm{Re}\left[C_\Phi^{(11)(22)(21)}\right]\Big]$$

$$\Delta M_{22}^{\pm 2} = -\mathrm{Re}\left[\eta M_{12}^{\pm 2}\right] - v^4\Big[3s_\beta^4 C_\Phi^{(22)(22)(22)} + 2s_\beta^2 c_\beta^2 C_\Phi^{(11)(22)(22)} + c_\beta^4 C_\Phi^{(11)(11)(22)} + 4s_\beta^3 c_\beta \mathrm{Re}\left[C_\Phi^{(22)(22)(21)}\right] + 2s_\beta^2 c_\beta^2 \mathrm{Re}\left[C_\Phi^{(22)(21)(21)}\right] + s_\beta^2 c_\beta^2 C_\Phi^{(22)(21)(12)} + 2s_\beta c_\beta^3 \mathrm{Re}\left[C_\Phi^{(11)(22)(21)}\right]\Big]$$

$$\Delta M_{12}^{\pm 2} = \left(\Delta M_{21}^{\pm 2}\right)^* = -\frac{\eta^*}{2}\left(M_{11}^{\pm 2} + M_{22}^{\pm 2}\right) - v^4\Big[c_\beta^4 C_\Phi^{(11)(11)(21)} + s_\beta^4 C_\Phi^{(22)(22)(21)} + 2s_\beta c_\beta^3 C_\Phi^{(11)(21)(21)} + 2s_\beta^3 c_\beta C_\Phi^{(22)(21)(21)} + s_\beta c_\beta^3 C_\Phi^{(11)(21)(12)} + s_\beta^3 c_\beta C_\Phi^{(22)(21)(12)} + 3s_\beta^2 c_\beta^2 C_\Phi^{(21)(21)(21)} + 3s_\beta^2 c_\beta^2 C_\Phi^{(21)(21)(12)} + s_\beta^2 c_\beta^2 C_\Phi^{(11)(22)(21)}\Big]$$

$$M_{11}^2 = \frac{g'^2}{2}(v^2 + 2v_1 v_2 \mathrm{Re}[\eta])$$

$$M_{12}^2 = M_{21}^2 = -\frac{gg'}{2}(v^2 + 2v_1 v_2 \mathrm{Re}[\eta])$$

$$M_{22}^2 = \frac{g^2}{2}(v^2 + 2v_1 v_2 \mathrm{Re}[\eta])$$

$$\Delta M_{11}^2 = -\frac{g'^2}{2}\Big(v_1^4 C_{\Phi D}^{(11)(11)} + v_2^4 C_{\Phi D}^{(22)(22)} + v_1^2 v_2^2 C_{\Phi D}^{(21)(12)} + v_1^2 v_2^2 C_{\Phi D}^{(11)(22)} + 2v_1^2 v_2^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right] + 2v_1^3 v_2 \mathrm{Re}\left[C_{\Phi D}^{(21)(11)}\right] + 2v_1 v_2^3 \mathrm{Re}\left[C_{\Phi D}^{(21)(22)}\right]\Big) - \frac{1}{2}(g'^2 X_B - gg' X_{WB})(v^2 + 2v_1 v_2 \mathrm{Re}[\eta])$$

$$\Delta M_{22}^2 = -\frac{g^2}{2}\Big(v_1^4 C_{\Phi D}^{(11)(11)} + v_2^4 C_{\Phi D}^{(22)(22)} + v_1^2 v_2^2 C_{\Phi D}^{(21)(12)} + v_1^2 v_2^2 C_{\Phi D}^{(11)(22)} + 2v_1^2 v_2^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right] + 2v_1^3 v_2 \mathrm{Re}\left[C_{\Phi D}^{(21)(11)}\right] + 2v_1 v_2^3 \mathrm{Re}\left[C_{\Phi D}^{(21)(22)}\right]\Big) - \frac{1}{2}(g^2 X_W - gg' X_{WB})(v^2 + 2v_1 v_2 \mathrm{Re}[\eta])$$

$$\Delta M_{12}^2 = \Delta M_{21}^2 = +\frac{gg'}{2}\Big(v_1^4 C_{\Phi D}^{(11)(11)} + v_2^4 C_{\Phi D}^{(22)(22)} + v_1^2 v_2^2 C_{\Phi D}^{(21)(12)} + v_1^2 v_2^2 C_{\Phi D}^{(11)(22)} + 2v_1^2 v_2^2 \mathrm{Re}\left[C_{\Phi D}^{(21)(21)}\right] + 2v_1^3 v_2 \mathrm{Re}\left[C_{\Phi D}^{(21)(11)}\right] + 2v_1 v_2^3 \mathrm{Re}\left[C_{\Phi D}^{(21)(22)}\right]\Big) + \frac{1}{4}(gg'(X_B + X_W) - (g^2 + g'^2)X_{WB})(v^2 + 2v_1 v_2 \mathrm{Re}[\eta])$$

$$\begin{pmatrix} \cos\chi & \sin\chi \\ -\sin\chi & \cos\chi \end{pmatrix}\begin{pmatrix} a & c \\ c^* & b \end{pmatrix}\begin{pmatrix} \cos\chi & -\sin\chi \\ \sin\chi & \cos\chi \end{pmatrix} = \begin{pmatrix} m_+^2 & 0 \\ 0 & m_-^2 \end{pmatrix}$$

where $m_\pm^2 = \frac{1}{2}\left[a + b \pm \sqrt{(a-b)^2 + 4|c|^2}\right]$

$$\tan 2\chi = \frac{2|c|}{a-b}$$

$$\sin 2\chi = 2\sin\chi\cos\chi = \frac{2|c|}{\sqrt{(a-b)^2+4|c|^2}}$$

$$\begin{pmatrix}\hat{\rho}_1\\ \hat{\rho}_2\end{pmatrix} = \begin{pmatrix}\cos\hat{\alpha} & -\sin\hat{\alpha}\\ \sin\hat{\alpha} & \cos\hat{\alpha}\end{pmatrix}\begin{pmatrix}H\\ h\end{pmatrix}$$

$$\begin{pmatrix}\hat{a}_1\\ \hat{a}_2\end{pmatrix} = \begin{pmatrix}\cos\hat{\beta} & -\sin\hat{\beta}\\ \sin\hat{\beta} & \cos\hat{\beta}\end{pmatrix}\begin{pmatrix}G\\ A\end{pmatrix}$$

$$\begin{pmatrix}\hat{\phi}_1^+\\ \hat{\phi}_2^+\end{pmatrix} = \begin{pmatrix}\cos\hat{\beta}^\pm & -\sin\hat{\beta}^\pm\\ \sin\hat{\beta}^\pm & \cos\hat{\beta}^\pm\end{pmatrix}\begin{pmatrix}G^+\\ H^+\end{pmatrix}$$

$$\tan 2\hat{\alpha} = \frac{2\left(M^2_{\rho_1\rho_2}+\Delta M^2_{\rho_1\rho_2}\right)}{\left(M^2_{\rho_1\rho_1}-M^2_{\rho_2\rho_2}\right)+\left(\Delta M^2_{\rho_1\rho 1}-\Delta M^2_{\rho_2\rho_2}\right)}$$
$$\simeq \tan 2\alpha\left[1-\frac{\Delta M^2_{\rho_1\rho_1}-\Delta M^2_{\rho_2\rho_2}}{M^2_{\rho_1\rho_1}-M^2_{\rho_2\rho_2}}+\frac{\Delta M^2_{\rho_1\rho_2}}{M^2_{\rho_1\rho_2}}\right]$$

$$\tan 2\hat{\beta} = \frac{2\left(M^2_{a_1a_2}+\Delta M^2_{a_1a_2}\right)}{\left(M^2_{a_1a_1}-M^2_{a_2a_2}\right)+\left(\Delta M^2_{a_1a_1}-\Delta M^2_{a_2a_2}\right)}$$
$$\simeq \tan 2\beta\left[1-\frac{\Delta M^2_{a_1a_1}-\Delta M^2_{a_2a_2}}{M^2_{a_1a_1}-M^2_{a_2a_2}}+\frac{\Delta M^2_{a_1a_2}}{M^2_{a_1a_2}}\right]$$

$$\tan 2\hat{\beta}^\pm = \frac{2\left|M^{\pm2}_{12}+\Delta M^{\pm2}_{12}\right|}{\left(M^{\pm2}_{11}-M^{\pm2}_{22}\right)+\left(\Delta M^{\pm2}_{11}-\Delta M^{\pm2}_{22}\right)}$$
$$\simeq \tan 2\beta\left[1-\frac{\Delta M^{\pm2}_{11}-\Delta M^{\pm2}_{22}}{M^{\pm2}_{11}-M^{\pm2}_{22}}+\frac{\left|\Delta M^{\pm2}_{12}\right|}{\left|M^{\pm2}_{12}\right|}\mathrm{Re}\left[e^{i\phi}\right]\right]$$

$$\phi = \left|\phi_{M^{\pm2}_{12}}-\phi_{\Delta M^{\pm2}_{12}}\right|$$

$$m_H^2 = \cos^2\hat{\alpha}\left(M^2_{\rho_1\rho_1}+\Delta M^2_{\rho_1\rho_1}\right)+\sin 2\hat{\alpha}\left(M^2_{\rho_1\rho_2}+\Delta M^2_{\rho_1\rho_2}\right)+\sin^2\hat{\alpha}\left(M^2_{\rho_2\rho_2}+\Delta M^2_{\rho_2\rho_2}\right),$$
$$m_h^2 = \sin^2\hat{\alpha}\left(M^2_{\rho_1\rho_1}+\Delta M^2_{\rho_1\rho_1}\right)-\sin 2\hat{\alpha}\left(M^2_{\rho_1\rho_2}+\Delta M^2_{\rho_1\rho_2}\right)+\cos^2\hat{\alpha}\left(M^2_{\rho_2\rho_2}+\Delta M^2_{\rho_2\rho_2}\right),$$

$$m_G^2 = \cos^2\hat{\beta}\left(M^2_{a_1a_1}+\Delta M^2_{a_1a_1}\right)+\sin 2\hat{\beta}\left(M^2_{a_1a_2}+\Delta M^2_{a_1a_2}\right)+\sin^2\hat{\beta}\left(M^2_{a_2a_2}+\Delta M^2_{a_2a_2}\right)$$
$$\simeq 0$$
$$m_A^2 = \sin^2\hat{\beta}\left(M^2_{a_1a_1}+\Delta M^2_{a_1a_1}\right)-\sin 2\hat{\beta}\left(M^2_{a_1a_2}+\Delta M^2_{a_1a_2}\right)+\cos^2\hat{\beta}\left(M^2_{a_2a_2}+\Delta M^2_{a_2a_2}\right)$$
$$\simeq M^2_{a_1a_1}+M^2_{a_2a_2}+\Delta M^2_{a_1a_1}+\Delta M^2_{a_2a_2}$$

$$m_{G^\pm}^2 = \cos^2\hat{\beta}^\pm\left(M^{\pm2}_{11}+\Delta M^{\pm2}_{11}\right)+\sin 2\hat{\beta}^\pm\left|M^{\pm2}_{12}+\Delta M^{\pm2}_{12}\right|+\sin^2\hat{\beta}^\pm\left(M^{\pm2}_{22}+\Delta M^{\pm2}_{22}\right)$$
$$\simeq 0$$
$$m_{H^\pm}^2 = \sin^2\hat{\beta}^\pm\left(M^{\pm2}_{11}+\Delta M^{\pm2}_{11}\right)-\sin 2\hat{\beta}^\pm\left|M^{\pm2}_{12}+\Delta M^{\pm2}_{12}\right|+\cos^2\hat{\beta}^\pm\left(M^{\pm2}_{22}+\Delta M^{\pm2}_{22}\right)$$
$$\simeq M^{\pm2}_{11}+M^{\pm2}_{22}+\Delta M^{\pm2}_{11}+\Delta M^{\pm2}_{22}$$

$$\sin\chi = \sqrt{\frac{b}{m_+^2}} = \sqrt{\frac{b}{a+b}}$$

$$\cos\chi = \sqrt{\frac{m_+^2 - b}{m_+^2}} = \sqrt{\frac{a}{a+b}}$$

$$\begin{aligned}\sin(\beta-\chi) &= \sin\beta\cos\chi - \sin\chi\cos\beta \\ &= \sqrt{\frac{b}{a+b}}\sqrt{\frac{a+\Delta a}{a+b+\Delta a+\Delta b}} - \sqrt{\frac{a}{a+b}}\sqrt{\frac{b+\Delta b}{a+b+\Delta a+\Delta b}} \\ &\simeq \frac{1}{2(a+b)}\left[\Delta a\sqrt{\frac{b}{a}} - \Delta b\sqrt{\frac{a}{b}}\right] = \beta-\chi \sim \mathcal{O}\left(\frac{v^4}{\Lambda^2 m_+^2}\right),\end{aligned}$$

$$\begin{aligned}\cos(\beta-\chi) &= \cos\beta\cos\chi + \sin\beta\sin\chi \\ &\simeq 1 - \frac{1}{2}(\beta-\chi)^2 \sim 1 + \mathcal{O}\left(\frac{v^8}{\Lambda^4 m_+^4}\right)\end{aligned}$$

$$\chi = \hat{\beta}, m_+^2 = m_A^2 \text{ and } \chi = \hat{\beta}^{\pm}, m_+^2 = m_{H^{\pm}}^2$$

$$\sin\chi \simeq \sin\beta\left[1 - \frac{\Delta a+\Delta b}{2(a+b)} + \frac{\Delta b}{2b}\right] \equiv \sin\beta\left[1-\delta_{s_\chi}\right]$$

$$\cos\chi \simeq \cos\beta\left[1 - \frac{\Delta a+\Delta b}{2(a+b)} + \frac{\Delta a}{2a}\right] \equiv \cos\beta\left[1-\delta_{c_\chi}\right]$$

$$\sin\hat{\beta} \simeq \sin\beta\left[1 - \frac{\Delta M_{a_1a_1}^2 + \Delta M_{a_2a_2}^2}{M_{a_1a_1}^2 + M_{a_2a_2}^2} + \frac{\Delta M_{a_2a_2}^2}{M_{a_2a_2}^2}\right] = \sin\beta\left[1-\delta_{s_{\hat{\beta}}}\right]$$

$$\cos\hat{\beta} \simeq \sin\beta\left[1 - \frac{\Delta M_{a_1a_1}^2 + \Delta M_{a_2a_2}^2}{M_{a_1a_1}^2 + M_{a_2a_2}^2} + \frac{\Delta M_{1_2a_1}^2}{M_{a_1a_1}^2}\right] = \sin\beta\left[1-\delta_{c_{\hat{\beta}}}\right]$$

$$\sin\hat{\beta}^{\pm} \simeq \sin\beta\left[1 - \frac{\Delta M_{11}^{\pm 2} + \Delta M_{22}^{\pm 2}}{M_{11}^{\pm 2} + M_{22}^{\pm 2}} + \frac{\Delta M_{22}^{\pm 2}}{M_{22}^{\pm 2}}\right] = \sin\beta\left[1-\delta_{s_{\hat{\beta}^{\pm}}}\right]$$

$$\cos\hat{\beta}^{\pm} \simeq \cos\beta\left[1 - \frac{\Delta M_{11}^{\pm 2} + \Delta M_{22}^{\pm 2}}{M_{11}^{\pm 2} + M_{22}^{\pm 2}} + \frac{\Delta M_{11}^{\pm 2}}{M_{11}^{\pm 2}}\right] = \cos\beta\left[1-\delta_{c_{\hat{\beta}^{\pm}}}\right]$$

$$\sin\hat{\xi} \simeq \sin\xi\left[1-\delta_{s_{\hat{\xi}}}\right], \quad \cos\hat{\xi} \simeq \cos\xi\left[1-\delta_{c_{\hat{\xi}}}\right]$$
$$\sin\hat{\omega} \simeq \sin\omega\left[1-\delta_{s_{\hat{\omega}}}\right], \quad \cos\hat{\omega} \simeq \cos\omega\left[1-\delta_{c_{\hat{\omega}}}\right]$$


$\bar{\psi}^{f_1}(\hat{C})_{f_1f_2}\psi^{f_2}$ and its Hermitian conjugate $\bar{\psi}^{f_1}(\hat{C}^*)_{f_2f_1}\psi^{f_2}; m_i$

$\psi^c = \mathbf{C}\bar{\psi}^T$ and $\bar{\psi}^c = \psi^T\mathbf{C}$, where $\mathbf{C} = i\gamma^2\gamma^0$ is the chargeconjugate matrix


Dirac spinors are given as $u(p) = \mathbf{C}\bar{v}(p)^T$ and $v(p) = \bar{u}(p)^T\mathbf{C}$

$$\lambda^{(1)} = |b|/3 + 2|p|^{1/2}\cos((\theta+2\pi)/3), \lambda^{(2)} = |b|/3 + 2|p|^{1/2}\cos((\theta-2\pi)/3)$$


$$\lambda^{(3)} = |b|/3 + 2|p|^{1/2}\cos(\theta/3) \text{ with } \theta = \arccos\left(|q|/2|p|^{3/2}\right) \in [0, \pi/2]$$




$$+\frac{iU^*_{f_2f_1}}{v}\left(m_{\nu_{f_1}}\mathcal{P}_L + m_{e_{f_2}}\mathcal{P}_R\left(\delta_{c_{\hat{\beta}\pm}} - 1\right)\right)$$
$$+2iv\not{p}_3\mathcal{P}_L U^*_{g_1f_1}\left(c^2_\beta \hat{C}^{(11)[3]}_{\Phi l,g_1f_2} + s^2_\beta \hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$-\frac{\sqrt{2}\delta_{f_1f_2}\gamma^5}{v}\left(m_{\nu_{f_1}}\right)$$
$$+\sqrt{2}vU_{g_2f_2}U^*_{g_1f_1}\not{p}_3\gamma^5\left(c^2_\beta \hat{C}^{(11)[1]}_{\Phi l,g_1g_2} - c^2_\beta \hat{C}^{(11)[3]}_{\Phi l,g_1g_2} + s^2_\beta\left(\hat{C}^{(22)[1]}_{\Phi l,g_1g_2} - \hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)\right)$$




$$-\frac{i\sqrt{2}\delta_{f_1f_2}}{v}\left(m_{\nu_{f_1}}\right)$$




$$+\sqrt{2}vs_{2\beta}\left(\mathcal{P}_L\hat{C}^{(11)}_{\nu\nu\Phi,f_1f_2} - \mathcal{P}_L\hat{C}^{(22)}_{\nu\nu\Phi,f_1f_2} - \mathcal{P}_R\hat{C}^{(11)*}_{\nu\nu\Phi,f_1f_2} + \mathcal{P}_R\hat{C}^{(22)*}_{\nu\nu\Phi,f_1f_2}\right)$$
$$+\sqrt{2}vs_\beta c_\beta U_{g_2f_2}U^*_{g_1f_1}\not{p}_3\gamma^5\left(-\hat{C}^{(11)[1]}_{\Phi l,g_1g_2} + \hat{C}^{(11)[3]}_{\Phi l,g_1g_2} + \hat{C}^{(22)[1]}_{\Phi l,g_1g_2} - \hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)$$




$$+i\sqrt{2}vs_{2\beta}\left(\mathcal{P}_L\hat{C}^{(11)}_{\nu\nu\Phi,f_1f_2} - \mathcal{P}_L\hat{C}^{(22)}_{\nu\nu\Phi,f_1f_2} + \mathcal{P}_R\hat{C}^{(11)*}_{\nu\nu\Phi,f_1f_2} - \mathcal{P}_R\hat{C}^{(22)*}_{\nu\nu\Phi,f_1f_2}\right)$$




$$
\begin{aligned}
&+2ivs_\beta c_\beta \mathcal{P}_L U^*_{f_2 g_1}\left(\hat{C}^{(11)}_{\nu\nu\Phi,g_1f_1}-\hat{C}^{(22)}_{\nu\nu\Phi,g_1f_1}\right)\\
&-is_\beta \mathcal{P}_R\left(\delta_{s_{\hat{\beta}\pm}}-1\right)U^*_{g_1f_1}\left(\hat{y}^{(1)}_{e,g_1f_2}\right)\\
&-iv^2 s_\beta \mathcal{P}_R U^*_{g_1f_1}\left(c^2_\beta \hat{C}^{(11)}_{l\Phi_1,g_1f_2}-c^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1f_2}\right.\\
&\qquad\left.-c^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1f_2}+s^2_\beta \hat{C}^{(22)}_{l\Phi_1,g_1f_2}\right)\\
&-ivs_{2\beta}\not{p}_3 \mathcal{P}_L U^*_{g_1f_1}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{\delta_{f_1f_2}m_{e_{f_1}}\gamma^5}{2\sqrt{2}v}\left(A'_1+t_\beta B'+2\delta_{c_{\hat{\beta}}}-2\right)\\
&-\sqrt{2}v\not{p}_3 \mathcal{P}_L\left(c^2_\beta \hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+c^2_\beta \hat{C}^{(11)[3]}_{\Phi l,f_1f_2}+s^2_\beta\left(\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)\right)\\
&-\sqrt{2}v\not{p}_3 \mathcal{P}_R\left(c^2_\beta \hat{C}^{(11)}_{\Phi e,f_1f_2}+s^2_\beta \hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{i\delta_{f_1f_2}m_{e_{f_1}}}{2\sqrt{2}v}\left(A_1+Bt_\beta-2\right)\\
&+i\sqrt{2}v^2c_\beta\left(c^2_\beta\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.\\
&\qquad+s^2_\beta\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\qquad+s^2_\beta\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\qquad\left.+s^2_\beta\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{1}{2\sqrt{2}}\left(-\left(A'_1-2\right)s_\beta+c_\beta B'-2s_\beta\delta_{s_{\hat{\beta}}}\right)\left(\mathcal{P}_L\hat{y}^{(1)*}_{e,f_2f_1}-\mathcal{P}_R\hat{y}^{(1)}_{e,f_1f_2}\right)\\
&+\frac{v^2s_\beta}{\sqrt{2}}\left(-c^2_\beta\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.\\
&\qquad+s^2_\beta\left(2ct^2_\beta+1\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\qquad-s^2_\beta\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\qquad\left.-s^2_\beta\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)\\
&+\sqrt{2}vs_\beta c_\beta\not{p}_3\mathcal{P}_L\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)\\
&+\sqrt{2}vs_\beta c_\beta\not{p}_3\mathcal{P}_R\left(\hat{C}^{(11)}_{\Phi e,f_1f_2}-\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{i\left(Bc_\beta-\left(A_1-2\right)s_\beta\right)}{2\sqrt{2}}\left(\mathcal{P}_L\hat{y}^{(1)*}_{e,f_2f_1}+\mathcal{P}_R\hat{y}^{(1)}_{e,f_1f_2}\right)\\
&+\frac{iv^2s_\beta}{\sqrt{2}}\left(3c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.\\
&\qquad-s_\beta^2\left(2ct_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\qquad-s_\beta^2\left(2ct_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\qquad\left.-s_\beta^2\left(2ct_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{i\hat{g}\delta_{f_1f_2}\hat{g}'\gamma^{\mu_3}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\frac{\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}-1\right)\\
&+\frac{2vc_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{lW\Phi_1,f_2f_1}^{*}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.\\
&\qquad\left.-\hat{g}\left(\hat{C}_{lB\Phi_1,f_2f_1}^{*}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{lB\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{i\hat{g}U^*_{f_2f_1}}{\sqrt{2}}\left(\gamma^{\mu_3}\mathcal{P}_L\right)\\
&-2\sqrt{2}vc_\beta p_{3\nu}U^*_{g_1f_1}\sigma^{\mu_3\nu}\mathcal{P}_R\left(\hat{C}_{lW\Phi_1,g_1f_2}\right)\\
&-i\sqrt{2}\hat{g}v^2U^*_{g_1f_1}\gamma^{\mu_3}\mathcal{P}_L\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{i\delta_{f_1f_2}\gamma^{\mu_3}\gamma^5}{2\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}X_{WB}\hat{g}'+\hat{g}'^2+\hat{g}^2\right)\\
&-iv^2\sqrt{\hat{g}'^2+\hat{g}^2}U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_3}\gamma^5\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}-c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{i\delta_{f_1f_2}}{2\left(\hat{g}'^2+\hat{g}^2\right)^{3/2}}\left(\left(\hat{g}'^2-\hat{g}^2\right)\left(-\hat{g}X_{WB}\hat{g}'+\hat{g}'^2+\hat{g}^2\right)\gamma^{\mu_3}\mathcal{P}_L\right.\\
&\qquad\left.+2\hat{g}'\left(\hat{g}'^3+\hat{g}^2\hat{g}'+\hat{g}^3X_{WB}\right)\gamma^{\mu_3}\mathcal{P}_R\right)\\
&+\frac{2vc_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}^{*}_{lB\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{lB\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.\\
&\qquad\left.+\hat{g}\left(\hat{C}^{*}_{lW\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)\\
&+\frac{1}{2}iv^2\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_3}\mathcal{P}_L\left(2c_\beta^2\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+2c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}\right.\\
&\qquad\left.-\left(c_{2\beta}-1\right)\left(\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)\right)\\
&+\frac{1}{2}iv^2\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_3}\mathcal{P}_R\left(\left(c_{2\beta}+1\right)\hat{C}^{(11)}_{\Phi e,f_1f_2}-\left(c_{2\beta}-1\right)\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$
-\frac{2im_{\nu_{g_1}}U^{*}_{f_1g_1}U^{*}_{f_2g_1}}{v^2}\left(\mathcal{P}_L\right)
$$




$$
\begin{aligned}
&-\frac{m_{\nu_{f_1}}U^{*}_{f_2f_1}}{\sqrt{2}v^2}\left(\mathcal{P}_L\right)\\
&-\sqrt{2}\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)U^{*}_{g_1f_1}\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{im_{\nu_{f_1}}U^{*}_{f_2f_1}}{\sqrt{2}v^2}\left(\mathcal{P}_L\right)\\
&+i\sqrt{2}vc_\beta\mathcal{P}_RU^{*}_{g_1f_1}\left(c_\beta^2\hat{C}^{(11)}_{l\Phi_1,g_1f_2}+s_\beta^2\hat{C}^{(12)}_{l\Phi_2,g_1f_2}\right.\\
&\qquad\left.+s_\beta^2\hat{C}^{(21)}_{l\Phi_2,g_1f_2}+s_\beta^2\hat{C}^{(22)}_{l\Phi_1,g_1f_2}\right)\\
&+i\sqrt{2}\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)U^{*}_{g_1f_1}\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)
\end{aligned}
$$




$$+\frac{i\delta_{f_1 f_2}}{v^2}\left(m_{\nu_{f_1}}\right)$$




$$-\frac{\delta_{f_1 f_2}\gamma^5}{v^2}\left(m_{\nu_{f_1}}\right)$$
$$+U_{g_2 f_2}U^*_{g_1 f_1}\left(\not{p}_3\gamma^5-\not{p}_4\gamma^5\right)\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}-c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}\right.$$
$$\left.+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)\right)$$




$$-\frac{i\delta_{f_1 f_2}}{v^2}\left(m_{\nu_{f_1}}\right)$$




$$-4is_\beta c_\beta \mathcal{P}_L U^*_{f_1 g_1}U^*_{f_2 g_2}\left(\hat{C}^{(11)}_{\nu\nu\Phi,g_1g_2}-\hat{C}^{(22)}_{\nu\nu\Phi,g_1g_2}\right)$$




$$
\begin{aligned}
&+\sqrt{2}s_\beta c_\beta \mathcal{P}_L U^*_{f_2 g_1}\left(\hat{C}^{(22)}_{\nu\nu\Phi,g_1 f_1}-\hat{C}^{(11)}_{\nu\nu\Phi,g_1 f_1}\right)\\
&-\frac{v s_\beta \mathcal{P}_R U^*_{g_1 f_1}}{\sqrt{2}}\left(s^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2}+c^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2}\right.\\
&\qquad\left.-s^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2}-c^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2}\right)\\
&+\sqrt{2}s_\beta c_\beta\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)U^*_{g_1 f_1}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-i\sqrt{2}s_\beta c_\beta \mathcal{P}_L U^*_{f_2 g_1}\left(\hat{C}^{(11)}_{\nu\nu\Phi,g_1 f_1}-\hat{C}^{(22)}_{\nu\nu\Phi,g_1 f_1}\right)\\
&+\frac{i v s_\beta \mathcal{P}_R U^*_{g_1 f_1}}{\sqrt{2}}\left(2c^2_\beta \hat{C}^{(11)}_{l\Phi_1,g_1 f_2}+s^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2}\right.\\
&\qquad-c^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2}+s^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2}\\
&\qquad\left.-c^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2}-2c^2_\beta \hat{C}^{(22)}_{l\Phi_1,g_1 f_2}\right)\\
&+i\sqrt{2}s_\beta c_\beta\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)U^*_{g_1 f_1}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)
\end{aligned}
$$




$$
+is_{2\beta}\left(\mathcal{P}_L\hat{C}^{(11)}_{\nu\nu\Phi,f_1 f_2}-\mathcal{P}_L\hat{C}^{(22)}_{\nu\nu\Phi,f_1 f_2}+\mathcal{P}_R\hat{C}^{(11)*}_{\nu\nu\Phi,f_1 f_2}-\mathcal{P}_R\hat{C}^{(22)*}_{\nu\nu\Phi,f_1 f_2}\right)
$$




$$
\begin{aligned}
&+s_{2\beta}\left(-\mathcal{P}_L\hat{C}^{(11)}_{\nu\nu\Phi,f_1 f_2}+\mathcal{P}_L\hat{C}^{(22)}_{\nu\nu\Phi,f_1 f_2}+\mathcal{P}_R\hat{C}^{(11)*}_{\nu\nu\Phi,f_1 f_2}-\mathcal{P}_R\hat{C}^{(22)*}_{\nu\nu\Phi,f_1 f_2}\right)\\
&+s_\beta c_\beta U_{g_2 f_2}U^*_{g_1 f_1}\left(\not{p}_3\gamma^5-\not{p}_4\gamma^5\right)\left(\hat{C}^{(11)[1]}_{\Phi l,g_1 g_2}-\hat{C}^{(11)[3]}_{\Phi l,g_1 g_2}\right.\\
&\qquad\left.-\hat{C}^{(22)[1]}_{\Phi l,g_1 g_2}+\hat{C}^{(22)[3]}_{\Phi l,g_1 g_2}\right)
\end{aligned}
$$




$$+s_{2\beta}\left(\mathcal{P}_L\hat{C}^{(11)}_{\nu\nu\Phi,f_1f_2}-\mathcal{P}_L\hat{C}^{(22)}_{\nu\nu\Phi,f_1f_2}-\mathcal{P}_R\hat{C}^{(11)*}_{\nu\nu\Phi,f_1f_2}+\mathcal{P}_R\hat{C}^{(22)*}_{\nu\nu\Phi,f_1f_2}\right)$$
$$-s_\beta c_\beta U_{g_2f_2}U^*_{g_1f_1}\left(\not{p}_3\gamma^5-\not{p}_4\gamma^5\right)\left(\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}+\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)$$




$$+is_{2\beta}\left(\mathcal{P}_L\hat{C}^{(11)}_{\nu\nu\Phi,f_1f_2}-\mathcal{P}_L\hat{C}^{(22)}_{\nu\nu\Phi,f_1f_2}+\mathcal{P}_R\hat{C}^{(11)*}_{\nu\nu\Phi,f_1f_2}-\mathcal{P}_R\hat{C}^{(22)*}_{\nu\nu\Phi,f_1f_2}\right)$$




$$+\sqrt{2}s_\beta c_\beta\mathcal{P}_LU^*_{f_2g_1}\left(\hat{C}^{(22)}_{\nu\nu\Phi,g_1f_1}-\hat{C}^{(11)}_{\nu\nu\Phi,g_1f_1}\right)$$
$$+\sqrt{2}s_\beta c_\beta\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)U^*_{g_1f_1}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$+i\sqrt{2}s_\beta c_\beta\mathcal{P}_LU^*_{f_2g_1}\left(\hat{C}^{(11)}_{\nu\nu\Phi,g_1f_1}-\hat{C}^{(22)}_{\nu\nu\Phi,g_1f_1}\right)$$
$$-i\sqrt{2}vs_\beta\mathcal{P}_RU^*_{g_1f_1}\left(c^2_\beta\hat{C}^{(11)}_{l\Phi_1,g_1f_2}-c^2_\beta\hat{C}^{(12)}_{l\Phi_2,g_1f_2}-c^2_\beta\hat{C}^{(21)}_{l\Phi_2,g_1f_2}+s^2_\beta\hat{C}^{(22)}_{l\Phi_1,g_1f_2}\right)$$
$$+i\sqrt{2}s_\beta c_\beta\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)U^*_{g_1f_1}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$+4i\mathcal{P}_L U^*_{f_1 g_1} U^*_{f_2 g_2} \left(s^2_\beta \hat{C}^{(11)}_{\nu\nu\Phi, g_1 g_2} + c^2_\beta \hat{C}^{(22)}_{\nu\nu\Phi, g_1 g_2} \right)$$




$$\begin{aligned}
&+\sqrt{2}\mathcal{P}_L U^*_{f_2 g_1} \left(s^2_\beta \hat{C}^{(11)}_{\nu\nu\Phi, g_1 f_1} + c^2_\beta \hat{C}^{(22)}_{\nu\nu\Phi, g_1 f_1} \right) \\
&-\frac{v c_\beta \mathcal{P}_R U^*_{g_1 f_1}}{\sqrt{2}} \left(s^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} \right. \\
&\qquad \left. - s^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} - c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} \right) \\
&-\sqrt{2} \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L \right) U^*_{g_1 f_1} \left(s^2_\beta \hat{C}^{(11)[3]}_{\Phi l, g_1 f_2} + c^2_\beta \hat{C}^{(22)[3]}_{\Phi l, g_1 f_2} \right)
\end{aligned}$$




$$\begin{aligned}
&+i\sqrt{2}\mathcal{P}_L U^*_{f_2 g_1} \left(s^2_\beta \hat{C}^{(11)}_{\nu\nu\Phi, g_1 f_1} + c^2_\beta \hat{C}^{(22)}_{\nu\nu\Phi, g_1 f_1} \right) \\
&-\frac{ivc_\beta \mathcal{P}_R U^*_{g_1 f_1}}{\sqrt{2}} \left(2 s^2_\beta \hat{C}^{(11)}_{l\Phi_1, g_1 f_2} - s^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} \right. \\
&\qquad + c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} - s^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} \\
&\qquad \left. + c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} - 2 s^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} \right) \\
&+i\sqrt{2} \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L \right) U^*_{g_1 f_1} \left(s^2_\beta \hat{C}^{(11)[3]}_{\Phi l, g_1 f_2} + c^2_\beta \hat{C}^{(22)[3]}_{\Phi l, g_1 f_2} \right)
\end{aligned}$$




$$\begin{aligned}
&-2i \left(s^2_\beta \left(\mathcal{P}_L \hat{C}^{(11)}_{\nu\nu\Phi, f_1 f_2} + \mathcal{P}_R \hat{C}^{(11)*}_{\nu\nu\Phi, f_1 f_2} \right) \right. \\
&\qquad \left. + c^2_\beta \left(\mathcal{P}_L \hat{C}^{(22)}_{\nu\nu\Phi, f_1 f_2} + \mathcal{P}_R \hat{C}^{(22)*}_{\nu\nu\Phi, f_1 f_2} \right) \right)
\end{aligned}$$




$$
\begin{aligned}
&+2\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)}_{\nu\nu\Phi,f_1f_2}-\mathcal{P}_R\hat{C}^{(11)*}_{\nu\nu\Phi,f_1f_2}\right)\right.\\
&\quad\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)}_{\nu\nu\Phi,f_1f_2}-\mathcal{P}_R\hat{C}^{(22)*}_{\nu\nu\Phi,f_1f_2}\right)\right)\\
&-U_{g_2f_2}U^*_{g_1f_1}\left(\not{p}_3\gamma^5-\not{p}_4\gamma^5\right)\left(s_\beta^2\left(\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}\right)\right.\\
&\qquad\left.+c_\beta^2\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}-c_\beta^2\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+2i\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)}_{\nu\nu\Phi,f_1f_2}+\mathcal{P}_R\hat{C}^{(11)*}_{\nu\nu\Phi,f_1f_2}\right)\right.\\
&\quad\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)}_{\nu\nu\Phi,f_1f_2}+\mathcal{P}_R\hat{C}^{(22)*}_{\nu\nu\Phi,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+ivc_\beta\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.\\
&\quad+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\quad+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\quad\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)\\
&+i\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}-c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)\right)\\
&+i\left(\not{p}_3\mathcal{P}_R-\not{p}_4\mathcal{P}_R\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi e,f_1f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+ivc_\beta\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.\\
&\quad+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\quad+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\quad\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+vc_\beta \left(c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.\\
&\quad +s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\quad +s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\quad \left.+s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)\\
&+\left(\not{p}_4 \mathcal{P}_L - \not{p}_3 \mathcal{P}_L\right)\left(c_\beta^2 \hat{C}^{(11)[1]}_{\Phi l,f_1f_2} + c_\beta^2 \hat{C}^{(11)[3]}_{\Phi l,f_1f_2}\right.\\
&\quad \left.+s_\beta^2 \left(\hat{C}^{(22)[1]}_{\Phi l,f_1f_2} + \hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)\right)\\
&+\left(\not{p}_4 \mathcal{P}_R - \not{p}_3 \mathcal{P}_R\right)\left(c_\beta^2 \hat{C}^{(11)}_{\Phi e,f_1f_2} + s_\beta^2 \hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+3ivc_\beta \left(c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.\\
&\quad +s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\quad +s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\quad \left.+s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&-ivs_\beta \left(c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.\\
&\quad -c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\\
&\quad +s_\beta^2 \mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} - c_\beta^2 \mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1}\\
&\quad \left.-c_\beta^2 \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2} + s_\beta^2 \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&+\frac{1}{2} is_{2\beta} \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L\right)\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2} - \hat{C}^{(11)[3]}_{\Phi l,f_1f_2} - \hat{C}^{(22)[1]}_{\Phi l,f_1f_2} + \hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)\\
&+\frac{1}{2} is_{2\beta} \left(\not{p}_3 \mathcal{P}_R - \not{p}_4 \mathcal{P}_R\right)\left(\hat{C}^{(11)}_{\Phi e,f_1f_2} - \hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-ivs_\beta \left(c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.\\
&\quad +s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\quad -c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\quad \left.-c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+vs_\beta \Big(-c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right) \\
&\quad +s_\beta^2 \left(2ct_\beta^2+1\right) \left(\mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right) \\
&\quad -s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right) \\
&\quad -s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big) \\
&+\frac{1}{2} s_{2\beta} \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L\right) \left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2} + \hat{C}^{(11)[3]}_{\Phi l,f_1f_2} - \hat{C}^{(22)[1]}_{\Phi l,f_1f_2} - \hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right) \\
&+\frac{1}{2} s_{2\beta} \left(\not{p}_3 \mathcal{P}_R - \not{p}_4 \mathcal{P}_R\right) \left(\hat{C}^{(11)}_{\Phi e,f_1f_2} - \hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+vs_\beta \Big(c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right) \\
&\quad +s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right) \\
&\quad -c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right) \\
&\quad -c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big) \\
&-\frac{1}{2} s_{2\beta} \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L\right) \left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2} + \hat{C}^{(11)[3]}_{\Phi l,f_1f_2} - \hat{C}^{(22)[1]}_{\Phi l,f_1f_2} - \hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right) \\
&-\frac{1}{2} s_{2\beta} \left(\not{p}_3 \mathcal{P}_R - \not{p}_4 \mathcal{P}_R\right) \left(\hat{C}^{(11)}_{\Phi e,f_1f_2} - \hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+ivs_\beta \Big(3c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right) \\
&\quad -s_\beta^2 \left(2ct_\beta^2-1\right) \left(\mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right) \\
&\quad -s_\beta^2 \left(2ct_\beta^2-1\right) \left(\mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right) \\
&\quad -s_\beta^2 \left(2ct_\beta^2-1\right) \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big)
\end{aligned}
$$




$$
\begin{aligned}
&+ivc_\beta \Big(s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right) \\
&\quad -c_\beta^2 \left(2t_\beta^2+1\right) \left(\mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right) \\
&\quad +c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right) \\
&\quad +c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big)
\end{aligned}
$$




$$
\begin{aligned}
&+ivc_\beta \Big(3s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right) \\
&\quad -c_\beta^2 \left(2t_\beta^2 - 1\right) \left(\mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right) \\
&\quad -c_\beta^2 \left(2t_\beta^2 - 1\right) \left(\mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right) \\
&\quad -c_\beta^2 \left(2t_\beta^2 - 1\right) \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big)
\end{aligned}
$$




$$
\begin{aligned}
&+ivc_\beta \Big(s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right) \\
&\quad -s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right) \\
&\quad -s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right) \\
&\quad +c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big) \\
&+i \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L\right) \left(s_\beta^2 \left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2} - \hat{C}^{(11)[3]}_{\Phi l,f_1f_2}\right)\right. \\
&\qquad \left. + c_\beta^2 \hat{C}^{(22)[1]}_{\Phi l,f_1f_2} - c_\beta^2 \hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right) \\
&+i \left(\not{p}_3 \mathcal{P}_R - \not{p}_4 \mathcal{P}_R\right) \left(s_\beta^2 \hat{C}^{(11)}_{\Phi e,f_1f_2} + c_\beta^2 \hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+vc_\beta \Big(-s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{l\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right) \\
&\quad -c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{l\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right) \\
&\quad +s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{l\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right) \\
&\quad +s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{l\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big) \\
&+ \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L\right) \left(s_\beta^2 \left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2} + \hat{C}^{(11)[3]}_{\Phi l,f_1f_2}\right)\right. \\
&\qquad \left. + c_\beta^2 \hat{C}^{(22)[1]}_{\Phi l,f_1f_2} + c_\beta^2 \hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right) \\
&+ \left(\not{p}_3 \mathcal{P}_R - \not{p}_4 \mathcal{P}_R\right) \left(s_\beta^2 \hat{C}^{(11)}_{\Phi e,f_1f_2} + c_\beta^2 \hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-2\sqrt{2} c_\beta p_{4\nu} \sigma^{\mu_4 \nu} \mathcal{P}_L \left(\hat{C}_{lW^*\Phi_1,f_2f_1}\right) \\
&-i\sqrt{2} \hat{g} v \gamma^{\mu_4} \mathcal{P}_L \left(c_\beta^2 \hat{C}^{(11)[1]}_{\Phi l,f_1f_2} + s_\beta^2 \hat{C}^{(22)[1]}_{\Phi l,f_1f_2}\right) \\
&-i\sqrt{2} \hat{g} v \gamma^{\mu_4} \mathcal{P}_R \left(c_\beta^2 \hat{C}^{(11)}_{\Phi e,f_1f_2} + s_\beta^2 \hat{C}^{(22)}_{\Phi e,f_1f_2}\right)
\end{aligned}
$$




$$+2\sqrt{2}s_\beta p_{4\nu}\sigma^{\mu_4\nu}\mathcal{P}_L\left(\hat{C}_{lW^*_{\Phi_1},f_2f_1}\right)$$
$$+\frac{i\hat{g}vs_{2\beta}\gamma^{\mu_4}\mathcal{P}_L}{\sqrt{2}}\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}\right)$$
$$+\frac{i\hat{g}vs_{2\beta}\gamma^{\mu_4}\mathcal{P}_R}{\sqrt{2}}\left(\hat{C}^{(11)}_{\Phi e,f_1f_2}-\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+\frac{i\sqrt{2}c_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}\left(\hat{C}_{lB\Phi^*_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L-\hat{C}_{lB\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}'\left(-\hat{C}_{lW^*_{\Phi_1},f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)$$




$$+\frac{\sqrt{2}c_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{lW^*_{\Phi_1},f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.$$
$$\left.-\hat{g}\left(\hat{C}_{lB^*_{\Phi_1},f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{lB\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)$$




$$-\frac{i\sqrt{2}c_\beta p_{4\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{lB^*_{\Phi_1},f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L-\hat{C}_{lB\Phi_1,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}\left(\hat{C}_{lW^*_{\Phi_1},f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L-\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right)$$




$$+\frac{\sqrt{2}c_\beta p_{4\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}^{*}_{lB\Phi_1,f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L+\hat{C}_{lB\Phi_1,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}\left(\hat{C}^{*}_{lW\Phi_1,f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L+\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right)$$
$$+\frac{iv\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_4}\mathcal{P}_L}{\sqrt{2}}\left(2c_\beta^2\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+2c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}\right.$$
$$\left.+2s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)\right)$$
$$+\frac{iv\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_4}\mathcal{P}_R}{\sqrt{2}}\left(2c_\beta^2\hat{C}^{(11)}_{\Phi e,f_1f_2}+2s_\beta^2\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+\frac{i\sqrt{2}s_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}^{*}_{lW\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L-\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}\left(-\hat{C}^{*}_{lB\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{lB\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)$$




$$+\frac{\sqrt{2}s_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}^{*}_{lW\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.$$
$$\left.-\hat{g}\left(\hat{C}^{*}_{lB\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{lB\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)$$




$$+\frac{i\sqrt{2}s_\beta p_{4\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}^{*}_{lB\Phi_1,f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L-\hat{C}_{lB\Phi_1,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}\left(\hat{C}^{*}_{lW\Phi_1,f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L-\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right)$$




$$
\begin{aligned}
&+\frac{\sqrt{2}s_\beta p_{4\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{lB\Phi_1,f_2f_1}^{*}\sigma^{\mu_4\nu}\mathcal{P}_L+\hat{C}_{lB\Phi_1,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right.\\
&\qquad\left.+\hat{g}\left(\hat{C}_{lW\Phi_1,f_2f_1}^{*}\sigma^{\mu_4\nu}\mathcal{P}_L+\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right)\\
&+i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_4}\mathcal{P}_L\left(\hat{C}_{\Phi l,f_1f_2}^{(11)[1]}+\hat{C}_{\Phi l,f_1f_2}^{(11)[3]}-\hat{C}_{\Phi l,f_1f_2}^{(22)[1]}-\hat{C}_{\Phi l,f_1f_2}^{(22)[3]}\right)\\
&+i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_4}\mathcal{P}_R\left(\hat{C}_{\Phi e,f_1f_2}^{(11)}-\hat{C}_{\Phi e,f_1f_2}^{(22)}\right)
\end{aligned}
$$




$$
+2\hat{g}vc_\beta\left(\sigma^{\mu_3\mu_4}\mathcal{P}_L\hat{C}_{lW\Phi_1,f_2f_1}^{*}+\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_R\right)
$$




$$
\begin{aligned}
&-\frac{2c_\beta p_{3\nu}U_{g_1f_1}^{*}\sigma^{\mu_3\nu}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}\hat{C}_{lB\Phi_1,g_1f_2}+\hat{g}'\hat{C}_{lW\Phi_1,g_1f_2}\right)\\
&-\frac{2i\hat{g}v\hat{g}'U_{g_1f_1}^{*}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{\Phi l,g_1f_2}^{(11)[3]}+s_\beta^2\hat{C}_{\Phi l,g_1f_2}^{(22)[3]}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{2c_\beta p_{4\nu}U_{g_1f_1}^{*}\sigma^{\mu_4\nu}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\hat{C}_{lB\Phi_1,g_1f_2}-\hat{g}\hat{C}_{lW\Phi_1,g_1f_2}\right)\\
&+\frac{2iv\hat{g}'^2U_{g_1f_1}^{*}\gamma^{\mu_4}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{\Phi l,g_1f_2}^{(11)[3]}+s_\beta^2\hat{C}_{\Phi l,g_1f_2}^{(22)[3]}\right)
\end{aligned}
$$




$$+\frac{2s_\beta p_{3\nu} U^*_{g_1 f_1} \sigma^{\mu_3 \nu} \mathcal{P}_R}{\sqrt{\hat{g}'^2 + \hat{g}^2}} \left(\hat{g} \hat{C}_{lB\Phi_1, g_1 f_2} + \hat{g}' \hat{C}_{lW\Phi_1, g_1 f_2} \right)$$
$$+\frac{i\hat{g} v s_{2\beta} \hat{g}' U^*_{g_1 f_1} \gamma^{\mu_3} \mathcal{P}_L}{\sqrt{\hat{g}'^2 + \hat{g}^2}} \left(\hat{C}^{(11)[3]}_{\Phi l, g_1 f_2} - \hat{C}^{(22)[3]}_{\Phi l, g_1 f_2} \right)$$




$$+\frac{2s_\beta p_{4\nu} U^*_{g_1 f_1} \sigma^{\mu_4 \nu} \mathcal{P}_R}{\sqrt{\hat{g}'^2 + \hat{g}^2}} \left(\hat{g} \hat{C}_{lW\Phi_1, g_1 f_2} - \hat{g}' \hat{C}_{lB\Phi_1, g_1 f_2} \right)$$
$$-\frac{iv s_{2\beta} \hat{g}'^2 U^*_{g_1 f_1} \gamma^{\mu_4} \mathcal{P}_L}{\sqrt{\hat{g}'^2 + \hat{g}^2}} \left(\hat{C}^{(11)[3]}_{\Phi l, g_1 f_2} - \hat{C}^{(22)[3]}_{\Phi l, g_1 f_2} \right)$$




$$-2ic_\beta p_{4\nu} U^*_{g_1 f_1} \sigma^{\mu_4 \nu} \mathcal{P}_R \left(\hat{C}_{lW\Phi_1, g_1 f_2} \right)$$




$$-2c_\beta p_{4\nu} U^*_{g_1 f_1} \sigma^{\mu_4 \nu} \mathcal{P}_R \left(\hat{C}_{lW\Phi_1, g_1 f_2} \right)$$
$$-2i\hat{g} v U^*_{g_1 f_1} \gamma^{\mu_4} \mathcal{P}_L \left(c^2_\beta \hat{C}^{(11)[3]}_{\Phi l, g_1 f_2} + s^2_\beta \hat{C}^{(22)[3]}_{\Phi l, g_1 f_2} \right)$$




$$+2is_\beta p_{4\nu} U^*_{g_1 f_1} \sigma^{\mu_4 \nu} \mathcal{P}_R \left(\hat{C}_{lW\Phi_1, g_1 f_2} \right)$$




$$-2s_\beta p_{4\nu} U^*_{g_1 f_1} \sigma^{\mu_4 \nu} \mathcal{P}_R \left(\hat{C}_{lW\Phi_1, g_1 f_2} \right)$$
$$-i\hat{g} v s_{2\beta} U^*_{g_1 f_1} \gamma^{\mu_4} \mathcal{P}_L \left(\hat{C}^{(11)[3]}_{\Phi l, g_1 f_2} - \hat{C}^{(22)[3]}_{\Phi l, g_1 f_2} \right)$$




$$-\frac{2\sqrt{2}\hat{g} v c_\beta \hat{g}' U^*_{g_1 f_1} \sigma^{\mu_3 \mu_4} \mathcal{P}_R}{\sqrt{\hat{g}'^2 + \hat{g}^2}} \left(\hat{C}_{lW\Phi_1, g_1 f_2} \right)$$




$$+\frac{2\sqrt{2}\hat{g}^2 v c_\beta U^*_{g_1 f_1} \sigma^{\mu_3 \mu_4} \mathcal{P}_R}{\sqrt{\hat{g}'^2 + \hat{g}^2}} \left(\hat{C}_{lW\Phi_1, g_1 f_2} \right)$$




$$-iU_{g_2f_2}U^*_{g_1f_1}\left(\not{p}_3\gamma^5-\not{p}_4\gamma^5\right)\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}+c^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}+s^2_\beta\left(\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}+\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)\right)$$




$$-is_\beta c_\beta U_{g_2f_2}U^*_{g_1f_1}\left(\not{p}_3\gamma^5-\not{p}_4\gamma^5\right)\left(\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}+\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)$$




$$-iU_{g_2f_2}U^*_{g_1f_1}\left(\not{p}_3\gamma^5-\not{p}_4\gamma^5\right)\left(s^2_\beta\left(\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}+\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}\right)+c^2_\beta\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)$$




$$+i\sqrt{2}\hat{g}vU_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_4}\gamma^5\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}+s^2_\beta\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}\right)$$




$$-i\sqrt{2}\hat{g}vs_\beta c_\beta U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_4}\gamma^5\left(\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}\right)$$




$$-i\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_4}\gamma^5\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}-c^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}+s^2_\beta\left(\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)\right)$$




$$-i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_4}\gamma^5\left(\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}+\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)$$




$$+\frac{c_\beta}{\sqrt{2}}\left(c^2_\beta\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)+s^2_\beta\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)+s^2_\beta\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)+s^2_\beta\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{ic_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{3c_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{ic_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{c_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{3ic_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{s_\beta}{\sqrt{2}}\left(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)$$
$$-\left(s_\beta^2\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+c_\beta^2\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+\left(c_\beta^2\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+s_\beta^2\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\right)$$




$$-\frac{is_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)$$
$$+\left(s_\beta^2\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-c_\beta^2\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.-\left(c_\beta^2\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-s_\beta^2\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\right)$$




$$+\frac{s_\beta}{\sqrt{2}}\left(-3c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(2ct_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(2ct_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(2ct_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$-\frac{is_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{s_\beta}{\sqrt{2}}\left(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(2ct_\beta^2+1\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{is_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-s_\beta^2\left(2ct_\beta^2+1\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{s_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{is_\beta}{\sqrt{2}}\left(3c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-s_\beta^2\left(2ct_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$-s_\beta^2\left(2ct_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.-s_\beta^2\left(2ct_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{s_\beta}{\sqrt{2}}\left(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{is_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{c_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)$$
$$-\left(s_\beta^2\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+c_\beta^2\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+\left(c_\beta^2\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+s_\beta^2\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\right)$$




$$-\frac{ic_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)$$
$$-\left(s_\beta^2\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-c_\beta^2\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+\left(c_\beta^2\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-s_\beta^2\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\right)$$




$$+\frac{c_\beta}{\sqrt{2}}\left(3s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-c_\beta^2\left(2t_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$-c_\beta^2\left(2t_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.-c_\beta^2\left(2t_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{ic_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-c_\beta^2\left(2t_\beta^2+1\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{c_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-c_\beta^2\left(2t_\beta^2+1\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$
\begin{aligned}
&+\frac{ic_\beta}{\sqrt{2}}\Big(3s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\\
&\quad -c_\beta^2\left(2t_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\quad -c_\beta^2\left(2t_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\quad -c_\beta^2\left(2t_\beta^2-1\right)\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{c_\beta}{\sqrt{2}}\Big(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\\
&\quad -s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\quad -s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\quad +c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{ic_\beta}{\sqrt{2}}\Big(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\\
&\quad -s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\quad -s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\quad +c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{ic_\beta}{\sqrt{2}}\Big(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\\
&\quad +c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)\\
&\quad -s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)\\
&\quad -s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\Big)
\end{aligned}
$$




$$+\frac{c_\beta}{\sqrt{2}}\left(-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{s_\beta}{\sqrt{2}}\left(-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{is_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$-\frac{3s_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{is_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{s_\beta}{\sqrt{2}}\left(-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+\frac{3is_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{l\Phi_1,f_1f_2}\right)\right.$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{l\Phi_2,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{l\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{l\Phi_2,f_1f_2}\right)$$
$$\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{l\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{l\Phi_1,f_1f_2}\right)\right)$$




$$+2ic_\beta\mathcal{P}_RU^*_{g_1f_1}\left(c_\beta^2\hat{C}^{(11)}_{l\Phi_1,g_1f_2}+s_\beta^2\hat{C}^{(12)}_{l\Phi_2,g_1f_2}\right.$$
$$\left.+s_\beta^2\hat{C}^{(21)}_{l\Phi_2,g_1f_2}+s_\beta^2\hat{C}^{(22)}_{l\Phi_1,g_1f_2}\right)$$




$$+ic_\beta \mathcal{P}_R U^*_{g_1 f_1} \Big(c^2_\beta \hat{C}^{(11)}_{l\Phi_1, g_1 f_2} + s^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} + s^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} + s^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} \Big)$$




$$+ic_\beta \mathcal{P}_R U^*_{g_1 f_1} \Big(c^2_\beta \hat{C}^{(11)}_{l\Phi_1, g_1 f_2} + s^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} + s^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} + s^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} \Big)$$




$$-is_\beta \mathcal{P}_R U^*_{g_1 f_1} \Big(2c^2_\beta \hat{C}^{(11)}_{l\Phi_1, g_1 f_2} - 2c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} + s^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} - c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} + s^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} - c^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} \Big)$$




$$-is_\beta \mathcal{P}_R U^*_{g_1 f_1} \Big(c^2_\beta \hat{C}^{(11)}_{l\Phi_1, g_1 f_2} - c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} - c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} + s^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} \Big)$$




$$-is_\beta \mathcal{P}_R U^*_{g_1 f_1} \Big(c^2_\beta \hat{C}^{(11)}_{l\Phi_1, g_1 f_2} - c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} - c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} + s^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} \Big)$$




$$-\frac{1}{2} is_\beta \mathcal{P}_R U^*_{g_1 f_1} \Big(2c^2_\beta \hat{C}^{(11)}_{l\Phi_1, g_1 f_2} + s^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} - c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} + s^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} - c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} - 2c^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} \Big)$$




$$-\frac{1}{2} s_\beta \mathcal{P}_R U^*_{g_1 f_1} \Big(s^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} - s^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} - c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} \Big)$$




$$-\frac{1}{2} s_\beta \mathcal{P}_R U^*_{g_1 f_1} \Big(s^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} - s^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} - c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} \Big)$$




$$+\frac{1}{2}is_\beta \mathcal{P}_R U^*_{g_1 f_1}\Big(2c^2_\beta \hat{C}^{(11)}_{l\Phi_1,g_1 f_2} + s^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} - c^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} + s^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} - c^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} - 2c^2_\beta \hat{C}^{(22)}_{l\Phi_1,g_1 f_2}\Big)$$




$$+2ic_\beta \mathcal{P}_R U^*_{g_1 f_1}\Big(s^2_\beta \hat{C}^{(11)}_{l\Phi_1,g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} - s^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} - s^2_\beta \hat{C}^{(22)}_{l\Phi_1,g_1 f_2}\Big)$$




$$+\frac{1}{2}ic_\beta \mathcal{P}_R U^*_{g_1 f_1}\Big(2s^2_\beta \hat{C}^{(11)}_{l\Phi_1,g_1 f_2} - s^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} - s^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} + c^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} - 2s^2_\beta \hat{C}^{(22)}_{l\Phi_1,g_1 f_2}\Big)$$




$$-\frac{1}{2}c_\beta \mathcal{P}_R U^*_{g_1 f_1}\Big(s^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} - s^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} - c^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2}\Big)$$




$$-\frac{1}{2}c_\beta \mathcal{P}_R U^*_{g_1 f_1}\left(s^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} - s^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} - c^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2}\right)$$




$$-\frac{1}{2}ic_\beta \mathcal{P}_R U^*_{g_1 f_1}\left(2s^2_\beta \hat{C}^{(11)}_{l\Phi_1,g_1 f_2} - s^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} - s^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} + c^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} - 2s^2_\beta \hat{C}^{(22)}_{l\Phi_1,g_1 f_2}\right)$$




$$+ic_\beta \mathcal{P}_R U^*_{g_1 f_1}\left(s^2_\beta \hat{C}^{(11)}_{l\Phi_1,g_1 f_2} - s^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} - s^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} + c^2_\beta \hat{C}^{(22)}_{l\Phi_1,g_1 f_2}\right)$$




$$+ic_\beta \mathcal{P}_R U^*_{g_1 f_1}\left(s^2_\beta \hat{C}^{(11)}_{l\Phi_1,g_1 f_2} - s^2_\beta \hat{C}^{(12)}_{l\Phi_2,g_1 f_2} - s^2_\beta \hat{C}^{(21)}_{l\Phi_2,g_1 f_2} + c^2_\beta \hat{C}^{(22)}_{l\Phi_1,g_1 f_2}\right)$$




$$-2is_\beta \mathcal{P}_R U^*_{g_1 f_1} \left(s^2_\beta \hat{C}^{(11)}_{l\Phi_1, g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} + c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} + c^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} \right)$$




$$-is_\beta \mathcal{P}_R U^*_{g_1 f_1} \left(s^2_\beta \hat{C}^{(11)}_{l\Phi_1, g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} + c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} + c^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} \right)$$




$$-is_\beta \mathcal{P}_R U^*_{g_1 f_1} \left(s^2_\beta \hat{C}^{(11)}_{l\Phi_1, g_1 f_2} + c^2_\beta \hat{C}^{(12)}_{l\Phi_2, g_1 f_2} + c^2_\beta \hat{C}^{(21)}_{l\Phi_2, g_1 f_2} + c^2_\beta \hat{C}^{(22)}_{l\Phi_1, g_1 f_2} \right)$$




$$-\frac{2\sqrt{2}\hat{g}c_\beta \hat{g}' \sigma^{\mu_3 \mu_5} \mathcal{P}_L}{\sqrt{\hat{g}'^2 + \hat{g}^2}} \left(\hat{C}_{lW\Phi_1, f_2 f_1 *} \right)$$




$$+\frac{2\sqrt{2}\hat{g}s_\beta\hat{g}'\sigma^{\mu_3\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW\Phi_1,f_2f_1*}\right)$$




$$-i\sqrt{2}\hat{g}c_\beta\left(\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{lW^*\Phi_1,f_2f_1}-\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_R\right)$$




$$+\sqrt{2}\hat{g}c_\beta\left(\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{lW^*\Phi_1,f_2f_1}+\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_R\right)$$




$$+i\sqrt{2}\hat{g}s_\beta\left(\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{lW^*\Phi_1,f_2f_1}-\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_R\right)$$




$$+\sqrt{2}\hat{g}s_\beta\left(\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{lW^*_{\Phi_1,f_2f_1}}+\hat{C}_{lW\Phi_1,f_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_R\right)$$




$$+\frac{2\sqrt{2}\hat{g}^2c_\beta\sigma^{\mu_4\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW^*_{\Phi_1,f_2f_1}}\right)$$




$$-\frac{2\sqrt{2}\hat{g}^2s_\beta\sigma^{\mu_4\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW^*_{\Phi_1,f_2f_1}}\right)$$




$$-\frac{2i\hat{g}c_\beta\hat{g}'U^*_{g_1f_1}\sigma^{\mu_3\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW\Phi_1,g_1f_2}\right)$$




$$-\frac{2\hat{g}c_\beta\hat{g}'U^*_{g_1f_1}\sigma^{\mu_3\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW\Phi_1,g_1f_2}\right)$$




$$+\frac{2i\hat{g}s_\beta\hat{g}'U^*_{g_1f_1}\sigma^{\mu_3\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW\Phi_1,g_1f_2}\right)$$




$$-\frac{2\hat{g}s_\beta\hat{g}'U^*_{g_1f_1}\sigma^{\mu_3\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW\Phi_1,g_1f_2}\right)$$




$$-2\hat{g}c_\beta U^*_{g_1f_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R\left(\hat{C}_{lW\Phi_1,g_1f_2}\right)$$




$$+2\hat{g}s_\beta U^*_{g_1 f_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R\left(\hat{C}_{lW\Phi_1,g_1 f_2}\right)$$




$$+\frac{2i\hat{g}^2 c_\beta U^*_{g_1 f_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW\Phi_1,g_1 f_2}\right)$$




$$+\frac{2\hat{g}^2 c_\beta U^*_{g_1 f_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW\Phi_1,g_1 f_2}\right)$$




$$-\frac{2i\hat{g}^2 s_\beta U^*_{g_1 f_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW\Phi_1,g_1 f_2}\right)$$




$$+\frac{2\hat{g}^2 s_\beta U^*_{g_1 f_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{lW\Phi_1,g_1 f_2}\right)$$




$$+\frac{2i\hat{g}\hat{g}' U_{g_2 f_2}U^*_{g_1 f_1}\gamma^{\mu_3}\gamma^5}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c^2_\beta \hat{C}^{(11)[1]}_{\Phi l,g_1 g_2}+c^2_\beta \hat{C}^{(11)[3]}_{\Phi l,g_1 g_2}\right.$$
$$\left.+s^2_\beta\left(\hat{C}^{(22)[1]}_{\Phi l,g_1 g_2}+\hat{C}^{(22)[3]}_{\Phi l,g_1 g_2}\right)\right)$$




$$-\frac{i\hat{g}s_{2\beta}\hat{g}' U_{g_2 f_2}U^*_{g_1 f_1}\gamma^{\mu_3}\gamma^5}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[1]}_{\Phi l,g_1 g_2}+\hat{C}^{(11)[3]}_{\Phi l,g_1 g_2}-\hat{C}^{(22)[1]}_{\Phi l,g_1 g_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1 g_2}\right)$$




$$+\frac{2i\hat{g}\hat{g}' U_{g_2 f_2}U^*_{g_1 f_1}\gamma^{\mu_3}\gamma^5}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s^2_\beta\left(\hat{C}^{(11)[1]}_{\Phi l,g_1 g_2}+\hat{C}^{(11)[3]}_{\Phi l,g_1 g_2}\right)\right.$$
$$\left.+c^2_\beta \hat{C}^{(22)[1]}_{\Phi l,g_1 g_2}+c^2_\beta \hat{C}^{(22)[3]}_{\Phi l,g_1 g_2}\right)$$




$$-\frac{i\hat{g}\hat{g}'\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(2c_\beta^2\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}-2c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}+2s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)\right)$$
$$-\frac{i\hat{g}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(2c_\beta^2\hat{C}^{(11)}_{\Phi e,f_1f_2}+2s_\beta^2\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+\frac{i\hat{g}s_{2\beta}\hat{g}'\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)$$
$$+\frac{i\hat{g}s_{2\beta}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)}_{\Phi e,f_1f_2}-\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$-\frac{2i\hat{g}\hat{g}'\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}\right)\right.$$
$$\left.+c_\beta^2\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}-c_\beta^2\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)$$
$$+\frac{i\hat{g}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(-2s_\beta^2\hat{C}^{(11)}_{\Phi e,f_1f_2}-2c_\beta^2\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$-\frac{\sqrt{2}\hat{g}\hat{g}'U^*_{g_1f_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$-\frac{i\sqrt{2}\hat{g}\hat{g}'U^*_{g_1f_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}+s^2_\beta\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$+\frac{\sqrt{2}\hat{g}s_\beta c_\beta\hat{g}'U^*_{g_1f_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$-\frac{i\sqrt{2}\hat{g}s_\beta c_\beta\hat{g}'U^*_{g_1f_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$+\frac{\sqrt{2}\hat{g}s_\beta c_\beta\hat{g}'U^*_{g_1f_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$+\frac{i\sqrt{2}\hat{g}s_\beta c_\beta \hat{g}' U^*_{g_1 f_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$-\frac{\sqrt{2}\hat{g}\hat{g}' U^*_{g_1 f_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$+\frac{i\sqrt{2}\hat{g}\hat{g}' U^*_{g_1 f_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$-i\sqrt{2}\hat{g}U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}+s^2_\beta\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$+i\sqrt{2}\hat{g}s_\beta c_\beta U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$-i\sqrt{2}\hat{g}U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$-\hat{g}U_{g_2 f_2}U^*_{g_1 f_1}\gamma^{\mu_5}\gamma^5\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi l,g_1 g_2}+s^2_\beta\hat{C}^{(22)[1]}_{\Phi l,g_1 g_2}\right)$$




$$+i\hat{g}U_{g_2 f_2}U^*_{g_1 f_1}\gamma^{\mu_5}\gamma^5\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi l,g_1 g_2}+s^2_\beta\hat{C}^{(22)[1]}_{\Phi l,g_1 g_2}\right)$$




$$+\hat{g}s_\beta c_\beta U_{g_2 f_2} U^*_{g_1 f_1} \gamma^{\mu_5}\gamma^5 \left(\hat{C}^{(11)[1]}_{\Phi l, g_1 g_2} - \hat{C}^{(22)[1]}_{\Phi l, g_1 g_2}\right)$$




$$+i\hat{g}s_\beta c_\beta U_{g_2 f_2} U^*_{g_1 f_1} \gamma^{\mu_5}\gamma^5 \left(\hat{C}^{(11)[1]}_{\Phi l, g_1 g_2} - \hat{C}^{(22)[1]}_{\Phi l, g_1 g_2}\right)$$




$$+\hat{g}s_\beta c_\beta U_{g_2 f_2} U^*_{g_1 f_1} \gamma^{\mu_5}\gamma^5 \left(\hat{C}^{(11)[1]}_{\Phi l, g_1 g_2} - \hat{C}^{(22)[1]}_{\Phi l, g_1 g_2}\right)$$




$$-i\hat{g}s_\beta c_\beta U_{g_2 f_2} U^*_{g_1 f_1} \gamma^{\mu_5}\gamma^5 \left(\hat{C}^{(11)[1]}_{\Phi l, g_1 g_2} - \hat{C}^{(22)[1]}_{\Phi l, g_1 g_2}\right)$$




$$-\hat{g}U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_5}\gamma^5\left(s^2_\beta\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}+c^2_\beta\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}\right)$$




$$-i\hat{g}U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_5}\gamma^5\left(s^2_\beta\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}+c^2_\beta\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}\right)$$




$$+\hat{g}\gamma^{\mu_5}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+s^2_\beta\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}\right)$$
$$+\hat{g}\gamma^{\mu_5}\mathcal{P}_R\left(c^2_\beta\hat{C}^{(11)}_{\Phi e,f_1f_2}+s^2_\beta\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$-i\hat{g}\gamma^{\mu_5}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+s^2_\beta\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}\right)$$
$$-i\hat{g}\gamma^{\mu_5}\mathcal{P}_R\left(c^2_\beta\hat{C}^{(11)}_{\Phi e,f_1f_2}+s^2_\beta\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+\frac{1}{2}\hat{g}s_{2\beta}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}\right)$$
$$+\frac{1}{2}\hat{g}s_{2\beta}\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(22)}_{\Phi e,f_1f_2}-\hat{C}^{(11)}_{\Phi e,f_1f_2}\right)$$




$$-\frac{1}{2}i\hat{g}s_{2\beta}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}\right)$$
$$-\frac{1}{2}i\hat{g}s_{2\beta}\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(11)}_{\Phi e,f_1f_2}-\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+\frac{1}{2}\hat{g}s_{2\beta}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}\right)$$
$$+\frac{1}{2}\hat{g}s_{2\beta}\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(22)}_{\Phi e,f_1f_2}-\hat{C}^{(11)}_{\Phi e,f_1f_2}\right)$$




$$+\frac{1}{2}i\hat{g}s_{2\beta}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}\right)$$
$$+\frac{1}{2}i\hat{g}s_{2\beta}\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(11)}_{\Phi e,f_1f_2}-\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+\hat{g}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta \hat{C}^{(11)[1]}_{\Phi l,f_1f_2} + c^2_\beta \hat{C}^{(22)[1]}_{\Phi l,f_1f_2}\right)$$
$$+\hat{g}\gamma^{\mu_5}\mathcal{P}_R\left(s^2_\beta \hat{C}^{(11)}_{\Phi e,f_1f_2} + c^2_\beta \hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+i\hat{g}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta \hat{C}^{(11)[1]}_{\Phi l,f_1f_2} + c^2_\beta \hat{C}^{(22)[1]}_{\Phi l,f_1f_2}\right)$$
$$+i\hat{g}\gamma^{\mu_5}\mathcal{P}_R\left(s^2_\beta \hat{C}^{(11)}_{\Phi e,f_1f_2} + c^2_\beta \hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$-i\sqrt{2}\hat{g}U^*_{g_1f_1}\gamma^{\mu_5}\mathcal{P}_L\left(c^2_\beta \hat{C}^{(11)[3]}_{\Phi l,g_1f_2} + s^2_\beta \hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$-i\sqrt{2}\hat{g}U^*_{g_1f_1}\gamma^{\mu_5}\mathcal{P}_L\left(c^2_\beta \hat{C}^{(11)[3]}_{\Phi l,g_1f_2} + s^2_\beta \hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$+i\sqrt{2}\hat{g}s_\beta c_\beta U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$-i\sqrt{2}\hat{g}s_\beta c_\beta U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$-i\sqrt{2}\hat{g}U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$-i\sqrt{2}\hat{g}U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$-\frac{i\left(\hat{g}'^2-\hat{g}^2\right)U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_5}\gamma^5}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}+c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}+\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)\right)$$




$$+\frac{is_\beta c_\beta\left(\hat{g}'^2-\hat{g}^2\right)U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_5}\gamma^5}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}+\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)$$




$$-\frac{i\left(\hat{g}'^2-\hat{g}^2\right)U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_5}\gamma^5}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}+\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}\right)+c_\beta^2\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}+c_\beta^2\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)$$




$$+\frac{i\left(\hat{g}'^2-\hat{g}^2\right)\gamma^{\mu_5}\mathcal{P}_L}{2\sqrt{\hat{g}'^2+\hat{g}^2}}\left(2c_\beta^2\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}-2c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}+2s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)\right)$$
$$+\frac{i\left(\hat{g}'^2-\hat{g}^2\right)\gamma^{\mu_5}\mathcal{P}_R}{2\sqrt{\hat{g}'^2+\hat{g}^2}}\left(2c_\beta^2\hat{C}^{(11)}_{\Phi e,f_1f_2}+2s_\beta^2\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$-\frac{is_\beta c_\beta \left(\hat{g}'^2-\hat{g}^2\right)\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)$$
$$-\frac{is_\beta c_\beta \left(\hat{g}'^2-\hat{g}^2\right)\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)}_{\Phi e,f_1f_2}-\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+\frac{i\left(\hat{g}'^2-\hat{g}^2\right)\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s^2_\beta\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}\right)\right.$$
$$\left.+c^2_\beta\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}-c^2_\beta\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)$$
$$+\frac{i\left(\hat{g}'^2-\hat{g}^2\right)\gamma^{\mu_5}\mathcal{P}_R}{2\sqrt{\hat{g}'^2+\hat{g}^2}}\left(2s^2_\beta\hat{C}^{(11)}_{\Phi e,f_1f_2}+2c^2_\beta\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$-i\sqrt{\hat{g}'^2+\hat{g}^2}U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_5}\gamma^5\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}-c^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}\right.$$
$$\left.+s^2_\beta\left(\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)\right)$$




$$-i\sqrt{\hat{g}'^2+\hat{g}^2}U_{g_2f_2}U^*_{g_1f_1}\gamma^{\mu_5}\gamma^5\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi l,g_1g_2}-c^2_\beta\hat{C}^{(11)[3]}_{\Phi l,g_1g_2}\right.$$
$$\left.+s^2_\beta\left(\hat{C}^{(22)[1]}_{\Phi l,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1g_2}\right)\right)$$




$$-is_\beta c_\beta \sqrt{\hat{g}'^2+\hat{g}^2} U_{g_2 f_2} U^*_{g_1 f_1} \gamma^{\mu_5}\gamma^5 \left(\hat{C}^{(11)[1]}_{\Phi l,g_1 g_2} - \hat{C}^{(11)[3]}_{\Phi l,g_1 g_2} - \hat{C}^{(22)[1]}_{\Phi l,g_1 g_2} + \hat{C}^{(22)[3]}_{\Phi l,g_1 g_2}\right)$$




$$-i\sqrt{\hat{g}'^2+\hat{g}^2} U_{g_2 f_2} U^*_{g_1 f_1} \gamma^{\mu_5}\gamma^5 \left(s^2_\beta \left(\hat{C}^{(11)[1]}_{\Phi l,g_1 g_2} - \hat{C}^{(11)[3]}_{\Phi l,g_1 g_2}\right) + c^2_\beta \hat{C}^{(22)[1]}_{\Phi l,g_1 g_2} - c^2_\beta \hat{C}^{(22)[3]}_{\Phi l,g_1 g_2}\right)$$




$$-i\sqrt{\hat{g}'^2+\hat{g}^2} U_{g_2 f_2} U^*_{g_1 f_1} \gamma^{\mu_5}\gamma^5 \left(s^2_\beta \left(\hat{C}^{(11)[1]}_{\Phi l,g_1 g_2} - \hat{C}^{(11)[3]}_{\Phi l,g_1 g_2}\right) + c^2_\beta \hat{C}^{(22)[1]}_{\Phi l,g_1 g_2} - c^2_\beta \hat{C}^{(22)[3]}_{\Phi l,g_1 g_2}\right)$$




$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_L \left(c^2_\beta \hat{C}^{(11)[1]}_{\Phi l,f_1 f_2} + c^2_\beta \hat{C}^{(11)[3]}_{\Phi l,f_1 f_2} + s^2_\beta \left(\hat{C}^{(22)[1]}_{\Phi l,f_1 f_2} + \hat{C}^{(22)[3]}_{\Phi l,f_1 f_2}\right)\right)$$
$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R \left(c^2_\beta \hat{C}^{(11)}_{\Phi e,f_1 f_2} + s^2_\beta \hat{C}^{(22)}_{\Phi e,f_1 f_2}\right)$$




$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_L\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)\right)$$
$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(c_\beta^2\hat{C}^{(11)}_{\Phi e,f_1f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$-is_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)$$
$$-is_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(11)}_{\Phi e,f_1f_2}-\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+is_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}-\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)$$
$$+is_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(11)}_{\Phi e,f_1f_2}-\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_L\left(s_\beta^2\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}\right)\right.$$
$$\left.+c_\beta^2\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}+c_\beta^2\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)$$
$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(s_\beta^2\hat{C}^{(11)}_{\Phi e,f_1f_2}+c_\beta^2\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_L\left(s_\beta^2\left(\hat{C}^{(11)[1]}_{\Phi l,f_1f_2}+\hat{C}^{(11)[3]}_{\Phi l,f_1f_2}\right)\right.$$
$$\left.+c_\beta^2\hat{C}^{(22)[1]}_{\Phi l,f_1f_2}+c_\beta^2\hat{C}^{(22)[3]}_{\Phi l,f_1f_2}\right)$$
$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(s_\beta^2\hat{C}^{(11)}_{\Phi e,f_1f_2}+c_\beta^2\hat{C}^{(22)}_{\Phi e,f_1f_2}\right)$$




$$+\frac{\sqrt{2}\hat{g}'^2U^*_{g_1f_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$+\frac{i\sqrt{2}\hat{g}'^2U^*_{g_1f_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}\right)$$




$$+\frac{\sqrt{2}s_\beta c_\beta\hat{g}'^2U^*_{g_1f_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(22)[3]}_{\Phi l,g_1f_2}-\hat{C}^{(11)[3]}_{\Phi l,g_1f_2}\right)$$




$$+\frac{i\sqrt{2}s_\beta c_\beta \hat{g}'^2 U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$+\frac{\sqrt{2}s_\beta c_\beta \hat{g}'^2 U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}-\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}\right)$$




$$-\frac{i\sqrt{2}s_\beta c_\beta \hat{g}'^2 U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}-\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$+\frac{\sqrt{2}\hat{g}'^2 U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s^2_\beta \hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}+c^2_\beta \hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$-\frac{i\sqrt{2}\hat{g}'^2 U^*_{g_1 f_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}^{(11)[3]}_{\Phi l,g_1 f_2}+c_\beta^2\hat{C}^{(22)[3]}_{\Phi l,g_1 f_2}\right)$$




$$\begin{aligned}
&+\frac{\delta_{f_1 f_2} m_{u_{f_1}}\gamma^5}{2\sqrt{2}v}\left(A_1'+t_\beta B'+2\delta_{c_\beta}-2\right)\\
&+\sqrt{2}vV_{f_1 g_1}\not{p}_3\mathcal{P}_L V^*_{f_2 g_2}\left(-c_\beta^2\hat{C}^{(11)[1]}_{\Phi q,g_1 g_2}+c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1 g_2}+s_\beta^2\left(\hat{C}^{(22)[3]}_{\Phi q,g_1 g_2}-\hat{C}^{(22)[1]}_{\Phi q,g_1 g_2}\right)\right)\\
&-\sqrt{2}v\not{p}_3\mathcal{P}_R\left(c_\beta^2\hat{C}^{(11)}_{\Phi u,f_1 f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi u,f_1 f_2}\right)
\end{aligned}$$




$$\begin{aligned}
&+\frac{i\delta_{f_1 f_2} m_{u_{f_1}}}{2\sqrt{2}v}\left(A_1+Bt_\beta-2\right)\\
&+i\sqrt{2}v^2 s_\beta\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2 f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1 f_2}\right)\right.\\
&\qquad+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2 f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1 f_2}\right)\\
&\qquad+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2 f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1 f_2}\right)\\
&\qquad\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2 f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1 f_2}\right)\right)
\end{aligned}$$




$$\begin{aligned}
&-\frac{\left(c_\beta\left(A_2'+2\delta_{c_\beta}-2\right)-s_\beta B'\right)}{2\sqrt{2}}\left(\mathcal{P}_L\hat{y}^{(2)*}_{u,f_2 f_1}-\mathcal{P}_R\hat{y}^{(2)}_{u,f_1 f_2}\right)\\
&+\frac{v^2 c_\beta}{\sqrt{2}}\left(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2 f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1 f_2}\right)\right.\\
&\qquad+\left(2s_\beta^2+c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2 f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1 f_2}\right)\\
&\qquad-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2 f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1 f_2}\right)\\
&\qquad\left.-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2 f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1 f_2}\right)\right)\\
&+\sqrt{2}vs_\beta c_\beta V_{f_1 g_1}\not{p}_3\mathcal{P}_L V^*_{f_2 g_2}\left(\hat{C}^{(11)[1]}_{\Phi q,g_1 g_2}-\hat{C}^{(11)[3]}_{\Phi q,g_1 g_2}-\hat{C}^{(22)[1]}_{\Phi q,g_1 g_2}+\hat{C}^{(22)[3]}_{\Phi q,g_1 g_2}\right)\\
&+\sqrt{2}vs_\beta c_\beta\not{p}_3\mathcal{P}_R\left(\hat{C}^{(11)}_{\Phi u,f_1 f_2}-\hat{C}^{(22)}_{\Phi u,f_1 f_2}\right)
\end{aligned}$$




$$
\begin{aligned}
&-\frac{i\left(\left(A_2-2\right)c_\beta - Bs_\beta\right)}{2\sqrt{2}}\left(\mathcal{P}_L\hat{y}^{(2)*}_{u,f_2f_1}+\mathcal{P}_R\hat{y}^{(2)}_{u,f_1f_2}\right)\\
&-\frac{iv^2c_\beta}{\sqrt{2}}\left(-\left(2s_\beta^2-c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.\\
&\qquad-\left(2s_\beta^2-c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)\\
&\qquad-\left(2s_\beta^2-c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)\\
&\qquad\left.+3s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{i}{v}\left(\delta_{c_{\hat{\beta}\pm}}-1\right)\left(m_{u_{f_1}}\mathcal{P}_L-m_{d_{f_2}}\mathcal{P}_R\right)\left(V_{f_1f_2}\right)\\
&-2ivs_\beta c_\beta \not{p}_3\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&+2ivV_{f_1g_1}\not{p}_3\mathcal{P}_L\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-i\left(s_\beta\mathcal{P}_RV_{f_1g_1}\left(\delta_{s_{\hat{\beta}\pm}}-1\right)\hat{y}^{(1)}_{d,g_1f_2}+c_\beta\left(\delta_{c_{\hat{\beta}\pm}}-1\right)\mathcal{P}_LV_{g_1f_2}\hat{y}^{(2)*}_{u,g_1f_1}\right)\\
&-iv^2\left(c_\beta^3\mathcal{P}_LV_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}-s_\beta^2c_\beta\mathcal{P}_LV_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}\right.\\
&\qquad-s_\beta^2c_\beta\mathcal{P}_LV_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}+s_\beta^2c_\beta\mathcal{P}_LV_{g_1f_2}\hat{C}^{(22)*}_{u\Phi_2,g_1f_1}\\
&\qquad+s_\beta c_\beta^2\mathcal{P}_RV_{f_1g_1}\hat{C}^{(11)}_{d\Phi_1,g_1f_2}-s_\beta c_\beta^2\mathcal{P}_RV_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}\\
&\qquad\left.-s_\beta c_\beta^2\mathcal{P}_RV_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}+s_\beta^3\mathcal{P}_RV_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}\right)\\
&-ivc_{2\beta}\not{p}_3\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&-ivs_{2\beta}V_{f_1g_1}\not{p}_3\mathcal{P}_L\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{2i\hat{g}\delta_{f_1f_2}\hat{g}'\gamma^{\mu_3}}{3\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\frac{\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}-1\right)\\
&-\frac{2vs_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}^{*}_{uW\Phi_2,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.\\
&\qquad\left.+\hat{g}\left(\hat{C}^{*}_{uB\Phi_2,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{uB\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)
\end{aligned}
$$




$$-\frac{i\hat{g}\delta_{f_1f_2}\hat{g}'\gamma^{\mu_3}}{3\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\frac{\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}-1\right)$$
$$+\frac{2vc_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}^*_{dW\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{dW\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.$$
$$\left.-\hat{g}\left(\hat{C}^*_{dB\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{dB\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)$$




$$-i\hat{g}_s\delta_{f_1f_2}\gamma^{\mu_3}\left(T^{a_3}_{m_1m_2}\right)$$
$$-4vs_\beta p_{3\nu}T^{a_3}_{m_1m_2}\left(\hat{C}^*_{uG\Phi_2,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{uG\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)$$




$$-i\hat{g}_s\delta_{f_1f_2}\gamma^{\mu_3}\left(T^{a_3}_{m_1m_2}\right)$$
$$-4vc_\beta p_{3\nu}T^{a_3}_{m_1m_2}\left(\hat{C}^*_{dG\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{dG\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)$$




$$-\frac{i\hat{g}V_{f_1f_2}}{\sqrt{2}}\left(\gamma^{\mu_3}\mathcal{P}_L\right)$$
$$-2\sqrt{2}vp_{3\nu}\left(s_\beta V_{g_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_L\hat{C}^*_{uW\Phi_2,g_1f_1}+c_\beta V_{f_1g_1}\hat{C}_{dW\Phi_1,g_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)$$
$$+i\sqrt{2}\hat{g}v^2s_\beta c_\beta\gamma^{\mu_3}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-i\sqrt{2}\hat{g}v^2V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$
\begin{aligned}
&+\frac{i\delta_{f_1f_2}}{6\left(\hat{g}'^2+\hat{g}^2\right)^{3/2}}\left(\left(-3\hat{g}X_{WB}\hat{g}'^3+\hat{g}^3X_{WB}\hat{g}'+\hat{g}'^4-2\hat{g}^2\hat{g}'^2-3\hat{g}^4\right)\gamma^{\mu_3}\mathcal{P}_L\right.\\
&\qquad\left.+4\hat{g}'\left(\hat{g}'^3+\hat{g}^2\hat{g}'+\hat{g}^3X_{WB}\right)\gamma^{\mu_3}\mathcal{P}_R\right)\\
&+\frac{2vs_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}^*_{uB\Phi_2,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{uB\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.\\
&\qquad\left.-\hat{g}\left(\hat{C}_{uW^*\Phi_2,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)\\
&+iv^2\sqrt{\hat{g}'^2+\hat{g}^2}V_{f_1g_1}V_{f_2g_2*}\gamma^{\mu_3}\mathcal{P}_L\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)\right)\\
&+iv^2\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_3}\mathcal{P}_R\left(c_\beta^2\hat{C}^{(11)}_{\Phi u,f_1f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{i\delta_{f_1f_2}}{6\left(\hat{g}'^2+\hat{g}^2\right)^{3/2}}\left(\left(3\hat{g}X_{WB}\hat{g}'^3+\hat{g}^3X_{WB}\hat{g}'+\hat{g}'^4+4\hat{g}^2\hat{g}'^2+3\hat{g}^4\right)\gamma^{\mu_3}\mathcal{P}_L\right.\\
&\qquad\left.-2\hat{g}'\left(\hat{g}'^3+\hat{g}^2\hat{g}'+\hat{g}^3X_{WB}\right)\gamma^{\mu_3}\mathcal{P}_R\right)\\
&+\frac{2vc_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}^*_{dB\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{dB\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.\\
&\qquad\left.+\hat{g}\left(\hat{C}^*_{dW\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{dW\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)\\
&+iv^2\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_3}\mathcal{P}_L\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi q,f_1f_2}+c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,f_1f_2}+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi q,f_1f_2}+\hat{C}^{(22)[3]}_{\Phi q,f_1f_2}\right)\right)\\
&+iv^2\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_3}\mathcal{P}_R\left(c_\beta^2\hat{C}^{(11)}_{\Phi d,f_1f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi d,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-ivs_\beta\left(-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.\\
&\quad+\left(2c_\beta^2+s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)\\
&\quad-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)\\
&\quad\left.-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+ivs_\beta\left(-\left(2c_\beta^2-s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.\\
&\quad-\left(2c_\beta^2-s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)\\
&\quad-\left(2c_\beta^2-s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)\\
&\quad\left.+3c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+ivs_\beta \left(s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.\\
&\quad -c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)\\
&\quad -c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)\\
&\quad \left.+c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)\\
&+iV_{f_1g_1}V^*_{f_2g_2} \left(\not{p}_3\mathcal{P}_L - \not{p}_4\mathcal{P}_L\right)\left(s_\beta^2 \left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2} + \hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right)\right.\\
&\quad \left.+c_\beta^2 \hat{C}^{(22)[1]}_{\Phi q,g_1g_2} + c_\beta^2 \hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)\\
&+i\left(\not{p}_3\mathcal{P}_R - \not{p}_4\mathcal{P}_R\right)\left(s_\beta^2 \hat{C}^{(11)}_{\Phi u,f_1f_2} + c_\beta^2 \hat{C}^{(22)}_{\Phi u,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+ivs_\beta \left(c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.\\
&\quad +c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)\\
&\quad +c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)\\
&\quad \left.+s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+ivs_\beta \left(c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.\\
&\quad +c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)\\
&\quad +c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)\\
&\quad \left.+s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)\\
&+iV_{f_1g_1}V^*_{f_2g_2} \left(\not{p}_3\mathcal{P}_L - \not{p}_4\mathcal{P}_L\right)\left(c_\beta^2 \hat{C}^{(11)[1]}_{\Phi q,g_1g_2} + c_\beta^2 \hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right.\\
&\quad \left.+s_\beta^2 \left(\hat{C}^{(22)[1]}_{\Phi q,g_1g_2} + \hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)\right)\\
&+i\left(\not{p}_3\mathcal{P}_R - \not{p}_4\mathcal{P}_R\right)\left(c_\beta^2 \hat{C}^{(11)}_{\Phi u,f_1f_2} + s_\beta^2 \hat{C}^{(22)}_{\Phi u,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+vs_\beta\left(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.\\
&\quad -c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)\\
&\quad -c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)\\
&\quad \left.-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)\\
&-V_{f_1g_1}V^*_{f_2g_2}\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right.\\
&\qquad\qquad \left.+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)\right)\\
&-\left(\not{p}_3\mathcal{P}_R-\not{p}_4\mathcal{P}_R\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi u,f_1f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+3ivs_\beta\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.\\
&\quad +c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)\\
&\quad +c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)\\
&\quad \left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+vc_\beta\left(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.\\
&\quad +\left(2s_\beta^2+c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)\\
&\quad -c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)\\
&\quad \left.-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)\\
&+s_\beta c_\beta V_{f_1g_1}V^*_{f_2g_2}\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right.\\
&\qquad\qquad \left.-\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}+\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)\\
&+s_\beta c_\beta\left(\not{p}_3\mathcal{P}_R-\not{p}_4\mathcal{P}_R\right)\left(\hat{C}^{(11)}_{\Phi u,f_1f_2}-\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-ivc_\beta \Big(-\left(2s_\beta^2 - c_\beta^2\right) \left(\mathcal{P}_L \hat{C}^{(11)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{u\Phi_2,f_1f_2} \right) \\
&\quad - \left(2s_\beta^2 - c_\beta^2\right) \left(\mathcal{P}_L \hat{C}^{(12)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{u\Phi_1,f_1f_2} \right) \\
&\quad - \left(2s_\beta^2 - c_\beta^2\right) \left(\mathcal{P}_L \hat{C}^{(21)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{u\Phi_1,f_1f_2} \right) \\
&\quad + 3s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{u\Phi_2,f_1f_2} \right) \Big)
\end{aligned}
$$




$$
\begin{aligned}
&+ivc_\beta \Big(-s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{u\Phi_2,f_1f_2} \right) \\
&\quad + s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{u\Phi_2,f_1f_2} \right) \\
&\quad - s_\beta^2 \mathcal{P}_L \hat{C}^{(12)*}_{u\Phi_1,f_2f_1} + c_\beta^2 \mathcal{P}_L \hat{C}^{(21)*}_{u\Phi_1,f_2f_1} \\
&\quad + c_\beta^2 \mathcal{P}_R \hat{C}^{(12)}_{u\Phi_1,f_1f_2} - s_\beta^2 \mathcal{P}_R \hat{C}^{(21)}_{u\Phi_1,f_1f_2} \Big) \\
&+ \frac{1}{2} i s_{2\beta} V_{f_1g_1} V^*_{f_2g_2} \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L \right) \left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2} + \hat{C}^{(11)[3]}_{\Phi q,g_1g_2} \right. \\
&\qquad \left. - \hat{C}^{(22)[1]}_{\Phi q,g_1g_2} - \hat{C}^{(22)[3]}_{\Phi q,g_1g_2} \right) \\
&+ \frac{1}{2} i s_{2\beta} \left(\not{p}_3 \mathcal{P}_R - \not{p}_4 \mathcal{P}_R \right) \left(\hat{C}^{(11)}_{\Phi u,f_1f_2} - \hat{C}^{(22)}_{\Phi u,f_1f_2} \right)
\end{aligned}
$$




$$
\begin{aligned}
&+vs_\beta \Big(-c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{u\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(11)}_{u\Phi_2,f_1f_2} \right) \\
&\quad + s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{u\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(12)}_{u\Phi_1,f_1f_2} \right) \\
&\quad - c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{u\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(21)}_{u\Phi_1,f_1f_2} \right) \\
&\quad + c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{u\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(22)}_{u\Phi_2,f_1f_2} \right) \Big) \\
&+ V_{f_1g_1} V^*_{f_2g_2} \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L \right) \left(s_\beta^2 \left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2} - \hat{C}^{(11)[3]}_{\Phi q,g_1g_2} \right) \right. \\
&\qquad \left. + c_\beta^2 \hat{C}^{(22)[1]}_{\Phi q,g_1g_2} - c_\beta^2 \hat{C}^{(22)[3]}_{\Phi q,g_1g_2} \right) \\
&+ \left(\not{p}_3 \mathcal{P}_R - \not{p}_4 \mathcal{P}_R \right) \left(s_\beta^2 \hat{C}^{(11)}_{\Phi u,f_1f_2} + c_\beta^2 \hat{C}^{(22)}_{\Phi u,f_1f_2} \right)
\end{aligned}
$$




$$
\begin{aligned}
&+ivc_\beta \Big(-s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right) \\
&\quad +c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right) \\
&\quad -s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R \hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right) \\
&\quad +s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R \hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)
\end{aligned}
$$




$$
\begin{aligned}
&+vc_\beta \Big(-s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(11)*}_{u\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right) \\
&\quad +c_\beta^2 \left(\mathcal{P}_L \hat{C}^{(12)*}_{u\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right) \\
&\quad -s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(21)*}_{u\Phi_1,f_2f_1} - \mathcal{P}_R \hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right) \\
&\quad +s_\beta^2 \left(\mathcal{P}_L \hat{C}^{(22)*}_{u\Phi_2,f_2f_1} - \mathcal{P}_R \hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big) \\
&-s_\beta c_\beta V_{f_1g_1} V_{f_2g_2*} \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L\right) \Big(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2} - \hat{C}^{(11)[3]}_{\Phi q,g_1g_2} \\
&\qquad\qquad -\hat{C}^{(22)[1]}_{\Phi q,g_1g_2} + \hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\Big) \\
&-s_\beta c_\beta \left(\not{p}_3 \mathcal{P}_R - \not{p}_4 \mathcal{P}_R\right) \left(\hat{C}^{(11)}_{\Phi u,f_1f_2} - \hat{C}^{(22)}_{\Phi u,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-i\sqrt{2}v \Big(s_\beta c_\beta^2 \mathcal{P}_L V_{g_1f_2} \hat{C}^{(11)*}_{u\Phi_2,g_1f_1} + s_\beta c_\beta^2 \mathcal{P}_L V_{g_1f_2} \hat{C}^{(12)*}_{u\Phi_1,g_1f_1} \\
&\quad +s_\beta c_\beta^2 \mathcal{P}_L V_{g_1f_2} \hat{C}^{(21)*}_{u\Phi_1,g_1f_1} + s_\beta^3 \mathcal{P}_L V_{g_1f_2} \hat{C}^{(22)*}_{u\Phi_2,g_1f_1} \\
&\quad -c_\beta^3 \mathcal{P}_R V_{f_1g_1} \hat{C}^{(11)}_{d\Phi_1,g_1f_2} - s_\beta^2 c_\beta \mathcal{P}_R V_{f_1g_1} \hat{C}^{(12)}_{d\Phi_2,g_1f_2} \\
&\quad -s_\beta^2 c_\beta \mathcal{P}_R V_{f_1g_1} \hat{C}^{(21)}_{d\Phi_2,g_1f_2} - s_\beta^2 c_\beta \mathcal{P}_R V_{f_1g_1} \hat{C}^{(22)}_{d\Phi_1,g_1f_2}\Big) \\
&-i\sqrt{2} s_\beta c_\beta \left(\not{p}_3 \mathcal{P}_R - \not{p}_4 \mathcal{P}_R\right) \left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right) \\
&+i\sqrt{2} V_{f_1g_1} \left(\not{p}_3 \mathcal{P}_L - \not{p}_4 \mathcal{P}_L\right) \left(c_\beta^2 \hat{C}^{(11)[3]}_{\Phi q,g_1f_2} + s_\beta^2 \hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{iv}{\sqrt{2}}\Big(-2s_\beta^2c_\beta\mathcal{P}_LV_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}+c_\beta^3\mathcal{P}_LV_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}\\
&\quad -s_\beta^2c_\beta\mathcal{P}_LV_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}+c_\beta^3\mathcal{P}_LV_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}\\
&\quad -s_\beta^2c_\beta\mathcal{P}_LV_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}+2s_\beta^2c_\beta\mathcal{P}_LV_{g_1f_2}\hat{C}^{(22)*}_{u\Phi_2,g_1f_1}\\
&\quad +2s_\beta c_\beta^2\mathcal{P}_RV_{f_1g_1}\hat{C}^{(11)}_{d\Phi_1,g_1f_2}+s_\beta^3\mathcal{P}_RV_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}\\
&\quad -s_\beta c_\beta^2\mathcal{P}_RV_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}+s_\beta^3\mathcal{P}_RV_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}\\
&\quad -s_\beta c_\beta^2\mathcal{P}_RV_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}-2s_\beta c_\beta^2\mathcal{P}_RV_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}\Big)\\
&+\frac{ic_{2\beta}\left(\not{p}_3\mathcal{P}_R-\not{p}_4\mathcal{P}_R\right)}{\sqrt{2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&+\frac{is_{2\beta}V_{f_1g_1}\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)}{\sqrt{2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-i\sqrt{2}v\Big(c_\beta^3\mathcal{P}_LV_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}-s_\beta^2c_\beta\mathcal{P}_LV_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}\\
&\quad -s_\beta^2c_\beta\mathcal{P}_LV_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}+s_\beta^2c_\beta\mathcal{P}_LV_{g_1f_2}\hat{C}^{(22)*}_{u\Phi_2,g_1f_1}\\
&\quad +s_\beta c_\beta^2\mathcal{P}_RV_{f_1g_1}\hat{C}^{(11)}_{d\Phi_1,g_1f_2}-s_\beta c_\beta^2\mathcal{P}_RV_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}\\
&\quad -s_\beta c_\beta^2\mathcal{P}_RV_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}+s_\beta^3\mathcal{P}_RV_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}\Big)\\
&+\frac{ic_{2\beta}\left(\not{p}_3\mathcal{P}_R-\not{p}_4\mathcal{P}_R\right)}{\sqrt{2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&+\frac{is_{2\beta}V_{f_1g_1}\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)}{\sqrt{2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{iv}{\sqrt{2}}\Big(-2s_\beta c_\beta^2\mathcal{P}_LV_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}+s_\beta^3\mathcal{P}_LV_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}\\
&\quad -s_\beta c_\beta^2\mathcal{P}_LV_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}+s_\beta^3\mathcal{P}_LV_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}\\
&\quad -s_\beta c_\beta^2\mathcal{P}_LV_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}+2s_\beta c_\beta^2\mathcal{P}_LV_{g_1f_2}\hat{C}^{(22)*}_{u\Phi_2,g_1f_1}\\
&\quad -2s_\beta^2c_\beta\mathcal{P}_RV_{f_1g_1}\hat{C}^{(11)}_{d\Phi_1,g_1f_2}-c_\beta^3\mathcal{P}_RV_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}\\
&\quad +s_\beta^2c_\beta\mathcal{P}_RV_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}-c_\beta^3\mathcal{P}_RV_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}\\
&\quad +s_\beta^2c_\beta\mathcal{P}_RV_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}+2s_\beta^2c_\beta\mathcal{P}_RV_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}\Big)\\
&+i\sqrt{2}s_\beta c_\beta\left(\not{p}_3\mathcal{P}_R-\not{p}_4\mathcal{P}_R\right)\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&+i\sqrt{2}V_{f_1g_1}\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)\left(s_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+c_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{v}{\sqrt{2}}\Big(c_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}+s_\beta^2c_\beta\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}\\
&\quad -c_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}-s_\beta^2c_\beta\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}\\
&\quad +s_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}+s_\beta c_\beta^2\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}\\
&\quad -s_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}-s_\beta c_\beta^2\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}\Big)\\
&+\frac{c_{2\beta}\left(\not{p}_4\mathcal{P}_R-\not{p}_3\mathcal{P}_R\right)}{\sqrt{2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&+\frac{s_{2\beta}V_{f_1g_1}\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)}{\sqrt{2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{v}{\sqrt{2}}\Big(s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}\\
&\quad -s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}-s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}\\
&\quad -c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}-s_\beta^2c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}\\
&\quad +c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}+s_\beta^2c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}\Big)\\
&+\sqrt{2}s_\beta c_\beta\left(\not{p}_3\mathcal{P}_R-\not{p}_4\mathcal{P}_R\right)\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&-\sqrt{2}V_{f_1g_1}\left(\not{p}_3\mathcal{P}_L-\not{p}_4\mathcal{P}_L\right)\left(s_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+c_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+2\sqrt{2}c_\beta p_{4\nu}\sigma^{\mu_4\nu}\mathcal{P}_L\left(\hat{C}_{uW^*_{\Phi},f_2f_1}\right)\\
&+\frac{i\hat{g}vs_{2\beta}V_{f_1g_1}V_{f_2g_2*}\gamma^{\mu_4}\mathcal{P}_L}{\sqrt{2}}\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}\right)\\
&+\frac{i\hat{g}vs_{2\beta}\gamma^{\mu_4}\mathcal{P}_R}{\sqrt{2}}\left(\hat{C}^{(11)}_{\Phi u,f_1f_2}-\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)
\end{aligned}
$$




$$-\frac{\sqrt{2}s_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{uW^*_{\Phi_2,f_2f_1}}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}\left(\hat{C}_{uB^*_{\Phi_2,f_2f_1}}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{uB\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)$$




$$+\frac{\sqrt{2}s_\beta p_{4\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{uB^*_{\Phi_2,f_2f_1}}\sigma^{\mu_4\nu}\mathcal{P}_L+\hat{C}_{uB\Phi_2,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right.$$
$$\left.-\hat{g}\left(\hat{C}_{uW^*_{\Phi_2,f_2f_1}}\sigma^{\mu_4\nu}\mathcal{P}_L+\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right)$$
$$+i\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_4}\mathcal{P}_L\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right.$$
$$\left.+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)\right)$$
$$+i\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_4}\mathcal{P}_R\left(c_\beta^2\hat{C}^{(11)}_{\Phi u,f_1f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$-\frac{i\sqrt{2}c_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{uW^*_{\Phi_2,f_2f_1}}\sigma^{\mu_3\nu}\mathcal{P}_L-\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}\left(\hat{C}_{uB^*_{\Phi_2,f_2f_1}}\sigma^{\mu_3\nu}\mathcal{P}_L-\hat{C}_{uB\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)$$




$$+\frac{i\sqrt{2}c_\beta p_{4\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{uB^*_{\Phi_2,f_2f_1}}\sigma^{\mu_4\nu}\mathcal{P}_L-\hat{C}_{uB\Phi_2,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}\left(-\hat{C}_{uW^*_{\Phi_2,f_2f_1}}\sigma^{\mu_4\nu}\mathcal{P}_L+\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right)$$




$$
\begin{aligned}
&-2p_{4\nu}\left(s_\beta V_{g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_L\hat{C}^*_{uW\Phi_2,g_1f_1}+c_\beta V_{f_1g_1}\hat{C}_{dW\Phi_1,g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\\
&+2i\hat{g}vs_\beta c_\beta\gamma^{\mu_4}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&-2i\hat{g}vV_{f_1g_1}\gamma^{\mu_4}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-2ip_{4\nu}\left(c_\beta V_{g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_L\hat{C}^*_{uW\Phi_2,g_1f_1}-s_\beta V_{f_1g_1}\hat{C}_{dW\Phi_1,g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\\
&-\hat{g}vc_{2\beta}\gamma^{\mu_4}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{2p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\Big(-c_\beta\hat{g}'V_{g_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_L\hat{C}^*_{uW\Phi_2,g_1f_1}\\
&\qquad+\hat{g}c_\beta V_{g_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_L\hat{C}^*_{uB\Phi_2,g_1f_1}\\
&\qquad+\hat{g}s_\beta V_{f_1g_1}\hat{C}_{dB\Phi_1,g_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\\
&\qquad+s_\beta\hat{g}'V_{f_1g_1}\hat{C}_{dW\Phi_1,g_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\Big)\\
&+\frac{i\hat{g}vc_{2\beta}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&+\frac{i\hat{g}vs_{2\beta}\hat{g}'V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{2p_{4\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\Big(c_\beta\hat{g}'V_{g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_L\hat{C}^*_{uB\Phi_2,g_1f_1}\\
&\qquad+\hat{g}c_\beta V_{g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_L\hat{C}^*_{uW\Phi_2,g_1f_1}\\
&\qquad+s_\beta\hat{g}'V_{f_1g_1}\hat{C}_{dB\Phi_1,g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\\
&\qquad-\hat{g}s_\beta V_{f_1g_1}\hat{C}_{dW\Phi_1,g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\Big)\\
&+\frac{i\hat{g}^2vc_{2\beta}\gamma^{\mu_4}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&-\frac{ivs_{2\beta}\hat{g}'^2V_{f_1g_1}\gamma^{\mu_4}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$+\frac{\sqrt{2}c_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{uW^*\Phi_2,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}\left(\hat{C}_{uB^*\Phi_2,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{uB\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)$$




$$+\frac{\sqrt{2}c_\beta p_{4\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}\left(\hat{C}_{uW^*\Phi_2,f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L+\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right.$$
$$\left.-\hat{g}'\left(\hat{C}_{uB^*\Phi_2,f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L+\hat{C}_{uB\Phi_2,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right)$$
$$+\frac{ivs_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}V_{f_1g_1}V_{f_2g_2*}\gamma^{\mu_4}\mathcal{P}_L}{\sqrt{2}}\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right.$$
$$\left.-\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}+\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)$$
$$+\frac{ivs_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_4}\mathcal{P}_R}{\sqrt{2}}\left(\hat{C}^{(11)}_{\Phi u,f_1f_2}-\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$+2p_{4\nu}\left(c_\beta V_{g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_L\hat{C}_{uW^*\Phi_2,g_1f_1}-s_\beta V_{f_1g_1}\hat{C}_{dW\Phi_1,g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)$$
$$-i\hat{g}vc_{2\beta}\gamma^{\mu_4}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-i\hat{g}vs_{2\beta}V_{f_1g_1}\gamma^{\mu_4}\mathcal{P}_L\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+2\sqrt{2}s_\beta p_{4\nu}\sigma^{\mu_4\nu}\mathcal{P}_L\left(\hat{C}_{uW^*\Phi_2,f_2f_1}\right)$$
$$-i\sqrt{2}\hat{g}vV_{f_1g_1}V_{f_2g_2*}\gamma^{\mu_4}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}+s^2_\beta\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}\right)$$
$$-i\sqrt{2}\hat{g}v\gamma^{\mu_4}\mathcal{P}_R\left(c^2_\beta\hat{C}^{(11)}_{\Phi u,f_1f_2}+s^2_\beta\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$-\frac{i\sqrt{2}s_\beta p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{uW^*_{\Phi_2},f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L-\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}\left(\hat{C}_{uB^*_{\Phi_2},f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L-\hat{C}_{uB\Phi_2,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)\right)$$




$$+\frac{i\sqrt{2}s_\beta p_{4\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}'\left(\hat{C}_{uB^*_{\Phi_2},f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L-\hat{C}_{uB\Phi_2,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right.$$
$$\left.+\hat{g}\left(-\hat{C}_{uW^*_{\Phi_2},f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L+\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)\right)$$




$$-2ip_{4\nu}\left(s_\beta V_{g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_L\hat{C}_{uW^*_{\Phi_2},g_1f_1}+c_\beta V_{f_1g_1}\hat{C}_{dW\Phi_1,g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)$$
$$-2\hat{g}vs_\beta c_\beta\gamma^{\mu_4}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$




$$-\frac{2p_{3\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta\hat{g}'V_{g_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_L\hat{C}_{uW^*_{\Phi_2},g_1f_1}\right.$$
$$-\hat{g}s_\beta V_{g_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_L\hat{C}_{uB^*_{\Phi_2},g_1f_1}$$
$$+\hat{g}c_\beta V_{f_1g_1}\hat{C}_{dB\Phi_1,g_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R$$
$$\left.+c_\beta\hat{g}'V_{f_1g_1}\hat{C}_{dW\Phi_1,g_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)$$
$$+\frac{2i\hat{g}vs_\beta c_\beta\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-\frac{2i\hat{g}v\hat{g}'V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$
\begin{aligned}
&-\frac{2p_{4\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\Big(s_\beta\hat{g}'V_{g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_L\hat{C}_{uB\Phi_2,g_1f_1}^{*}\\
&\qquad+\hat{g}s_\beta V_{g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_L\hat{C}_{uW\Phi_2,g_1f_1}^{*}\\
&\qquad-c_\beta\hat{g}'V_{f_1g_1}\hat{C}_{dB\Phi_1,g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\\
&\qquad+\hat{g}c_\beta V_{f_1g_1}\hat{C}_{dW\Phi_1,g_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\Big)\\
&+\frac{2i\hat{g}^2vs_\beta c_\beta\gamma^{\mu_4}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)\\
&+\frac{2iv\hat{g}'^2V_{f_1g_1}\gamma^{\mu_4}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)
\end{aligned}
$$




$$+2i\sqrt{2}c_\beta p_{3\nu}T^{a_3}_{m_1m_2}\left(\hat{C}^{*}_{dG\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L-\hat{C}_{dG\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)$$




$$-2\sqrt{2}c_\beta p_{3\nu}T^{a_3}_{m_1m_2}\left(\hat{C}^{*}_{dG\Phi_1,f_2f_1}\sigma^{\mu_3\nu}\mathcal{P}_L+\hat{C}_{dG\Phi_1,f_1f_2}\sigma^{\mu_3\nu}\mathcal{P}_R\right)$$




$$-2i\sqrt{2}s_\beta p_{4\nu}T^{a_4}_{m_1m_2}\left(\hat{C}^{*}_{dG\Phi_1,f_2f_1}\sigma^{\mu_4\nu}\mathcal{P}_L-\hat{C}_{dG\Phi_1,f_1f_2}\sigma^{\mu_4\nu}\mathcal{P}_R\right)$$




$$-2\sqrt{2}s_\beta p_{3\nu} T^{a_3}_{m_1 m_2} \left(\hat{C}^*_{dG\Phi_1, f_2 f_1} \sigma^{\mu_3 \nu} \mathcal{P}_L + \hat{C}_{dG\Phi_1, f_1 f_2} \sigma^{\mu_3 \nu} \mathcal{P}_R \right)$$




$$-4ivc_\beta \hat{g}_s f_{a_3 a_4 b_1} T^{b_1}_{m_1 m_2} \left(\sigma^{\mu_3 \mu_4} \mathcal{P}_L \hat{C}^*_{dG\Phi_1, f_2 f_1} + \hat{C}_{dG\Phi_1, f_1 f_2} \sigma^{\mu_3 \mu_4} \mathcal{P}_R \right)$$




$$+4p_{3\nu} T^{a_3}_{m_1 m_2} \left(c_\beta V_{g_1 f_2} \sigma^{\mu_3 \nu} \mathcal{P}_L \hat{C}^*_{uG\Phi_2, g_1 f_1} + s_\beta V_{f_1 g_1} \hat{C}_{dG\Phi_1, g_1 f_2} \sigma^{\mu_3 \nu} \mathcal{P}_R \right)$$




$$+4p_{3\nu} T^{a_3}_{m_1 m_2} \left(s_\beta V_{g_1 f_2} \sigma^{\mu_3 \nu} \mathcal{P}_L \hat{C}^*_{uG\Phi_2, g_1 f_1} - c_\beta V_{f_1 g_1} \hat{C}_{dG\Phi_1, g_1 f_2} \sigma^{\mu_3 \nu} \mathcal{P}_R \right)$$




$$-2i\sqrt{2}c_\beta p_{4\nu} T^{a_4}_{m_1 m_2} \left(\hat{C}^*_{uG\Phi_2, f_2 f_1} \sigma^{\mu_4 \nu} \mathcal{P}_L - \hat{C}_{uG\Phi_2, f_1 f_2} \sigma^{\mu_4 \nu} \mathcal{P}_R \right)$$




$$+2\sqrt{2}c_\beta p_{3\nu} T^{a_3}_{m_1 m_2} \left(\hat{C}^*_{uG\Phi_2, f_2 f_1} \sigma^{\mu_3 \nu} \mathcal{P}_L + \hat{C}_{uG\Phi_2, f_1 f_2} \sigma^{\mu_3 \nu} \mathcal{P}_R \right)$$




$$-2i\sqrt{2}s_\beta p_{3\nu} T^{a_3}_{m_1 m_2} \left(\hat{C}^*_{uG\Phi_2, f_2 f_1} \sigma^{\mu_3 \nu} \mathcal{P}_L - \hat{C}_{uG\Phi_2, f_1 f_2} \sigma^{\mu_3 \nu} \mathcal{P}_R \right)$$




$$-2\sqrt{2}s_\beta p_{3\nu} T^{a_3}_{m_1 m_2} \left(\hat{C}^*_{uG\Phi_2, f_2 f_1} \sigma^{\mu_3 \nu} \mathcal{P}_L + \hat{C}_{uG\Phi_2, f_1 f_2} \sigma^{\mu_3 \nu} \mathcal{P}_R \right)$$




$$-4ivs_\beta \hat{g}_s f_{a_3 a_4 b_1} T^{b_1}_{m_1 m_2} \left(\sigma^{\mu_3\mu_4}\mathcal{P}_L \hat{C}^*_{uG\Phi_2,f_2f_1} + \hat{C}_{uG\Phi_2,f_1f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_R\right)$$




$$-\frac{2\sqrt{2}\hat{g}v\hat{g}'}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta V_{g_1f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_L \hat{C}^*_{uW\Phi_2,g_1f_1} + c_\beta V_{f_1g_1}\sigma^{\mu_3\mu_4}\mathcal{P}_R \hat{C}_{dW\Phi_1,g_1f_2}\right)$$




$$-2\hat{g}vs_\beta\left(\sigma^{\mu_3\mu_4}\mathcal{P}_L \hat{C}^*_{uW\Phi_2,f_2f_1} + \hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_R\right)$$




$$+\frac{2\sqrt{2}\hat{g}^2 v}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta V_{g_1f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_L \hat{C}^*_{uW\Phi_2,g_1f_1} + c_\beta V_{f_1g_1}\sigma^{\mu_3\mu_4}\mathcal{P}_R \hat{C}_{dW\Phi_1,g_1f_2}\right)$$




$$+\sqrt{2}s_\beta c_\beta \left(\not{p}_4\mathcal{P}_R - \not{p}_3\mathcal{P}_R\right)\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-\sqrt{2}V_{f_1g_1}\left(\not{p}_3\mathcal{P}_L - \not{p}_4\mathcal{P}_L\right)\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2} + s_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{c_{2\beta}\left(\not{p}_4\mathcal{P}_R - \not{p}_3\mathcal{P}_R\right)}{\sqrt{2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{s_{2\beta}V_{f_1g_1}\left(\not{p}_3\mathcal{P}_L - \not{p}_4\mathcal{P}_L\right)}{\sqrt{2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2} - \hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{s_\beta}{\sqrt{2}}\Big(\left(2c_\beta^2 - s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1} - \mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$+\left(2c_\beta^2 - s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1} - \mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+\left(2c_\beta^2 - s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1} - \mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$-3c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1} - \mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{is_\beta}{\sqrt{2}}\Big(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$-\left(2c_\beta^2 + s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1} + \mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1} + \mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{s_\beta}{\sqrt{2}}\Big(-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$+\left(2c_\beta^2+s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{is_\beta}{\sqrt{2}}\Big(-\left(2c_\beta^2-s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$-\left(2c_\beta^2-s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-\left(2c_\beta^2-s_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$+3c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{s_\beta}{\sqrt{2}}\Big(-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{is_\beta}{\sqrt{2}}\Big(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$-\frac{3c_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)$$




$$-\frac{ic_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)$$




$$+\frac{c_\beta}{\sqrt{2}}\left(-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$\left.-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)$$




$$-\frac{3ic_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$\left.+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)$$




$$+\frac{c_\beta}{\sqrt{2}}\Big(-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$-\frac{ic_\beta}{\sqrt{2}}\Big(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$-\frac{3s_\beta}{\sqrt{2}}\Big(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{s_\beta}{\sqrt{2}}\Big(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{is_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)$$




$$+\frac{is_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)$$




$$+\frac{s_\beta}{\sqrt{2}}\left(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$\left.-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)$$




$$+\frac{3is_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)$$




$$+\frac{c_\beta}{\sqrt{2}}\Big(\left(2s_\beta^2-c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$+\left(2s_\beta^2-c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+\left(2s_\beta^2-c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$-3s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{c_\beta}{\sqrt{2}}\Big(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{c_\beta}{\sqrt{2}}\Big(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$+\left(2s_\beta^2+c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$-\frac{ic_\beta}{\sqrt{2}}\Big(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$-\left(2s_\beta^2+c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$-\frac{ic_\beta}{\sqrt{2}}\left(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$\left.+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)$$




$$-\frac{ic_\beta}{\sqrt{2}}\left(-\left(2s_\beta^2-c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$-\left(2s_\beta^2-c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-\left(2s_\beta^2-c_\beta^2\right)\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$\left.+3s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\right)$$




$$+\frac{c_\beta}{\sqrt{2}}\left(s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-c_\beta^2\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}$$
$$\left.+c_\beta^2\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}-s_\beta^2\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$




$$+\frac{ic_\beta}{\sqrt{2}}\left(-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)\right.$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)$$
$$-s_\beta^2\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+c_\beta^2\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}$$
$$\left.+c_\beta^2\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}-s_\beta^2\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$




$$+\frac{is_\beta}{\sqrt{2}}\Big(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$-s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{s_\beta}{\sqrt{2}}\Big(-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$+s_\beta^2\left(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\right)$$
$$+c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)\Big)$$




$$+\frac{s_\beta}{\sqrt{2}}\Big(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)$$
$$-s_\beta^2\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+c_\beta^2\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}$$
$$-c_\beta^2\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}+s_\beta^2\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\Big)$$




$$+\frac{is_\beta}{\sqrt{2}}\Big(c_\beta^2\left(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\right)$$
$$-c_\beta^2\left(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\right)$$
$$-s_\beta^2\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+c_\beta^2\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}$$
$$+c_\beta^2\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}-s_\beta^2\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\Big)$$




$$+\frac{ic_\beta}{\sqrt{2}}\Big(-s_\beta^2\Big(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\Big)$$
$$+c_\beta^2\Big(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\Big)$$
$$-s_\beta^2\Big(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}+\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\Big)$$
$$+s_\beta^2\Big(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}+\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\Big)\Big)$$




$$+\frac{c_\beta}{\sqrt{2}}\Big(-s_\beta^2\Big(\mathcal{P}_L\hat{C}^{(11)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(11)}_{u\Phi_2,f_1f_2}\Big)$$
$$+c_\beta^2\Big(\mathcal{P}_L\hat{C}^{(12)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(12)}_{u\Phi_1,f_1f_2}\Big)$$
$$-s_\beta^2\Big(\mathcal{P}_L\hat{C}^{(21)*}_{u\Phi_1,f_2f_1}-\mathcal{P}_R\hat{C}^{(21)}_{u\Phi_1,f_1f_2}\Big)$$
$$+s_\beta^2\Big(\mathcal{P}_L\hat{C}^{(22)*}_{u\Phi_2,f_2f_1}-\mathcal{P}_R\hat{C}^{(22)}_{u\Phi_2,f_1f_2}\Big)\Big)$$




$$-2i\Big(s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}$$
$$+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}+s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(22)*}_{u\Phi_2,g_1f_1}$$
$$-c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(11)}_{d\Phi_1,g_1f_2}-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}$$
$$-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}\Big)$$




$$-i\Big(c_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}-s_\beta^2 c_\beta\mathcal{P}_L V_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}$$
$$-2s_\beta^2 c_\beta\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}+c_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}$$
$$-s_\beta^2 c_\beta\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}+2s_\beta^2 c_\beta\mathcal{P}_L V_{g_1f_2}\hat{C}^{(22)*}_{u\Phi_2,g_1f_1}$$
$$+2s_\beta c_\beta^2\mathcal{P}_R V_{f_1g_1}\hat{C}^{(11)}_{d\Phi_1,g_1f_2}-2s_\beta c_\beta^2\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}$$
$$+s_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}-s_\beta c_\beta^2\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}$$
$$+s_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}-s_\beta c_\beta^2\mathcal{P}_R V_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}\Big)$$




$$
\begin{aligned}
&+2i\Big(s_\beta c_\beta^2 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(11)*}_{u\Phi_2, g_1 f_1} - s_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1, g_1 f_1} \\
&\quad + s_\beta c_\beta^2 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1, g_1 f_1} - s_\beta c_\beta^2 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(22)*}_{u\Phi_2, g_1 f_1} \\
&\quad + s_\beta^2 c_\beta \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(11)}_{d\Phi_1, g_1 f_2} + c_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2, g_1 f_2} \\
&\quad - s_\beta^2 c_\beta \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2, g_1 f_2} - s_\beta^2 c_\beta \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(22)}_{d\Phi_1, g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
&-2i\Big(s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(11)*}_{u\Phi_2, g_1 f_1} + s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1, g_1 f_1} \\
&\quad + s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1, g_1 f_1} + c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(22)*}_{u\Phi_2, g_1 f_1} \\
&\quad + s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(11)}_{d\Phi_1, g_1 f_2} + s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2, g_1 f_2} \\
&\quad + s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2, g_1 f_2} + s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(22)}_{d\Phi_1, g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
&-i\Big(s_\beta c_\beta^2 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(11)*}_{u\Phi_2, g_1 f_1} + s_\beta c_\beta^2 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1, g_1 f_1} \\
&\quad + s_\beta c_\beta^2 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1, g_1 f_1} + s_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(22)*}_{u\Phi_2, g_1 f_1} \\
&\quad - c_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(11)}_{d\Phi_1, g_1 f_2} - s_\beta^2 c_\beta \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2, g_1 f_2} \\
&\quad - s_\beta^2 c_\beta \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2, g_1 f_2} - s_\beta^2 c_\beta \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(22)}_{d\Phi_1, g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
&-i\Big(s_\beta c_\beta^2 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(11)*}_{u\Phi_2, g_1 f_1} + s_\beta c_\beta^2 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1, g_1 f_1} \\
&\quad + s_\beta c_\beta^2 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1, g_1 f_1} + s_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(22)*}_{u\Phi_2, g_1 f_1} \\
&\quad - c_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(11)}_{d\Phi_1, g_1 f_2} - s_\beta^2 c_\beta \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2, g_1 f_2} \\
&\quad - s_\beta^2 c_\beta \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2, g_1 f_2} - s_\beta^2 c_\beta \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(22)}_{d\Phi_1, g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
-\frac{1}{2}i\Big(&-2s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(11)*}_{u\Phi_2,g_1 f_1} + c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1,g_1 f_1} \\
&- s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1,g_1 f_1} + c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1,g_1 f_1} \\
&- s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1,g_1 f_1} + 2s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(22)*}_{u\Phi_2,g_1 f_1} \\
&+ 2s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(11)}_{d\Phi_1,g_1 f_2} + s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2,g_1 f_2} \\
&- s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2,g_1 f_2} + s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2,g_1 f_2} \\
&- s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2,g_1 f_2} - 2s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(22)}_{d\Phi_1,g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
+\frac{1}{2}i\Big(&-2s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(11)*}_{u\Phi_2,g_1 f_1} + c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1,g_1 f_1} \\
&- s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1,g_1 f_1} + c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1,g_1 f_1} \\
&- s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1,g_1 f_1} + 2s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(22)*}_{u\Phi_2,g_1 f_1} \\
&+ 2s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(11)}_{d\Phi_1,g_1 f_2} + s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2,g_1 f_2} \\
&- s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2,g_1 f_2} + s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2,g_1 f_2} \\
&- s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2,g_1 f_2} - 2s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(22)}_{d\Phi_1,g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
-i\Big(&c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(11)*}_{u\Phi_2,g_1 f_1} - s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1,g_1 f_1} \\
&- s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1,g_1 f_1} + s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(22)*}_{u\Phi_2,g_1 f_1} \\
&+ s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(11)}_{d\Phi_1,g_1 f_2} - s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2,g_1 f_2} \\
&- s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2,g_1 f_2} + s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(22)}_{d\Phi_1,g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
-i\Big(&c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(11)*}_{u\Phi_2,g_1 f_1} - s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1,g_1 f_1} \\
&- s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1,g_1 f_1} + s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(22)*}_{u\Phi_2,g_1 f_1} \\
&+ s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(11)}_{d\Phi_1,g_1 f_2} - s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2,g_1 f_2} \\
&- s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2,g_1 f_2} + s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(22)}_{d\Phi_1,g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
&+i\Big(-s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}\\
&\quad+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}-s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(22)*}_{u\Phi_2,g_1f_1}\\
&\quad+s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(11)}_{d\Phi_1,g_1f_2}-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}\\
&\quad-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}+c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
&+i\Big(-s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}\\
&\quad+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}-s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(22)*}_{u\Phi_2,g_1f_1}\\
&\quad+s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(11)}_{d\Phi_1,g_1f_2}-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}\\
&\quad-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}+c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{1}{2}i\Big(2s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}-s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}\\
&\quad+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}-s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}\\
&\quad+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}-2s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(22)*}_{u\Phi_2,g_1f_1}\\
&\quad+2s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(11)}_{d\Phi_1,g_1f_2}+c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}\\
&\quad-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}+c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}\\
&\quad-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}-2s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{1}{2}i\Big(-2s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(11)*}_{u\Phi_2,g_1f_1}+s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}\\
&\quad-s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}+s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}\\
&\quad-s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}+2s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(22)*}_{u\Phi_2,g_1f_1}\\
&\quad-2s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(11)}_{d\Phi_1,g_1f_2}-c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}\\
&\quad+s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}-c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}\\
&\quad+s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}+2s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(22)}_{d\Phi_1,g_1f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
-i\Big(&s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(11)*}_{u\Phi_2, g_1 f_1} + s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1, g_1 f_1} \\
&+ s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1, g_1 f_1} + c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(22)*}_{u\Phi_2, g_1 f_1} \\
&+ s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(11)}_{d\Phi_1, g_1 f_2} + s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2, g_1 f_2} \\
&+ s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2, g_1 f_2} + s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(22)}_{d\Phi_1, g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
-i\Big(&s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(11)*}_{u\Phi_2, g_1 f_1} + s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1, g_1 f_1} \\
&+ s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1, g_1 f_1} + c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(22)*}_{u\Phi_2, g_1 f_1} \\
&+ s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(11)}_{d\Phi_1, g_1 f_2} + s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2, g_1 f_2} \\
&+ s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2, g_1 f_2} + s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(22)}_{d\Phi_1, g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
-\frac{1}{2}\Big(&c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1, g_1 f_1} + s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1, g_1 f_1} \\
&- c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1, g_1 f_1} - s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1, g_1 f_1} \\
&+ s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2, g_1 f_2} + s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2, g_1 f_2} \\
&- s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2, g_1 f_2} - s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2, g_1 f_2}\Big)
\end{aligned}
$$




$$
\begin{aligned}
-\frac{1}{2}\Big(&c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1, g_1 f_1} + s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(12)*}_{u\Phi_1, g_1 f_1} \\
&- c_\beta^3 \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1, g_1 f_1} - s_\beta^2 c_\beta \mathcal{P}_L V_{g_1 f_2} \hat{C}^{(21)*}_{u\Phi_1, g_1 f_1} \\
&+ s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2, g_1 f_2} + s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(12)}_{d\Phi_2, g_1 f_2} \\
&- s_\beta^3 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2, g_1 f_2} - s_\beta c_\beta^2 \mathcal{P}_R V_{f_1 g_1} \hat{C}^{(21)}_{d\Phi_2, g_1 f_2}\Big)
\end{aligned}
$$




$$+\frac{1}{2}\Big(s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}$$
$$-s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}-s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}$$
$$-c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}$$
$$+c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}+s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}\Big)$$




$$+\frac{1}{2}\Big(s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}+s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(12)*}_{u\Phi_1,g_1f_1}$$
$$-s_\beta^3\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}-s_\beta c_\beta^2\mathcal{P}_L V_{g_1f_2}\hat{C}^{(21)*}_{u\Phi_1,g_1f_1}$$
$$-c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}-s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(12)}_{d\Phi_2,g_1f_2}$$
$$+c_\beta^3\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}+s_\beta^2 c_\beta\mathcal{P}_R V_{f_1g_1}\hat{C}^{(21)}_{d\Phi_2,g_1f_2}\Big)$$




$$+2\sqrt{2}c_\beta\hat{g}_s f_{a_3a_4b_1}T^{b_1}_{m_1m_2}\left(\hat{C}_{dG\Phi_1,f_1f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_R-\sigma^{\mu_3\mu_4}\mathcal{P}_L\hat{C}^*_{dG\Phi_1,f_2f_1}\right)$$




$$-2i\sqrt{2}c_\beta\hat{g}_s f_{a_3a_4b_1}T^{b_1}_{m_1m_2}\left(\sigma^{\mu_3\mu_4}\mathcal{P}_L\hat{C}^*_{dG\Phi_1,f_2f_1}+\hat{C}_{dG\Phi_1,f_1f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_R\right)$$




$$+2\sqrt{2}s_\beta \hat{g}_s f_{a_4 a_5 b_1} T^{b_1}_{m_1 m_2} \left(\sigma^{\mu_4\mu_5}\mathcal{P}_L \hat{C}^*_{dG\Phi_1,f_2 f_1} - \hat{C}_{dG\Phi_1,f_1 f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_R\right)$$




$$-2i\sqrt{2}s_\beta \hat{g}_s f_{a_3 a_4 b_1} T^{b_1}_{m_1 m_2} \left(\sigma^{\mu_3\mu_4}\mathcal{P}_L \hat{C}^*_{dG\Phi_1,f_2 f_1} + \hat{C}_{dG\Phi_1,f_1 f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_R\right)$$




$$+4i\hat{g}_s f_{a_3 a_4 b_1} T^{b_1}_{m_1 m_2} \left(c_\beta V_{g_1 f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_L \hat{C}^*_{uG\Phi_2,g_1 f_1} + s_\beta V_{f_1 g_1}\sigma^{\mu_3\mu_4}\mathcal{P}_R \hat{C}_{dG\Phi_1,g_1 f_2}\right)$$




$$+4i\hat{g}_s f_{a_3 a_4 b_1} T^{b_1}_{m_1 m_2} \left(s_\beta V_{g_1 f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_L \hat{C}^*_{uG\Phi_2,g_1 f_1} - c_\beta V_{f_1 g_1}\sigma^{\mu_3\mu_4}\mathcal{P}_R \hat{C}_{dG\Phi_1,g_1 f_2}\right)$$




$$+2\sqrt{2}c_\beta \hat{g}_s f_{a_4 a_5 b_1} T^{b_1}_{m_1 m_2} \left(\sigma^{\mu_4\mu_5}\mathcal{P}_L \hat{C}^*_{uG\Phi_2,f_2 f_1} - \hat{C}_{uG\Phi_2,f_1 f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_R\right)$$




$$+2i\sqrt{2}c_\beta \hat{g}_s f_{a_3 a_4 b_1} T^{b_1}_{m_1 m_2} \left(\sigma^{\mu_3\mu_4}\mathcal{P}_L \hat{C}^*_{uG\Phi_2,f_2f_1} + \hat{C}_{uG\Phi_2,f_1f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_R\right)$$




$$+2\sqrt{2}s_\beta \hat{g}_s f_{a_3 a_4 b_1} T^{b_1}_{m_1 m_2} \left(\sigma^{\mu_3\mu_4}\mathcal{P}_L \hat{C}^*_{uG\Phi_2,f_2f_1} - \hat{C}_{uG\Phi_2,f_1f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_R\right)$$




$$-2i\sqrt{2}s_\beta \hat{g}_s f_{a_3 a_4 b_1} T^{b_1}_{m_1 m_2} \left(\sigma^{\mu_3\mu_4}\mathcal{P}_L \hat{C}^*_{uG\Phi_2,f_2f_1} + \hat{C}_{uG\Phi_2,f_1f_2}\sigma^{\mu_3\mu_4}\mathcal{P}_R\right)$$




$$-\frac{2\hat{g}\hat{g}'}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta V_{g_1f_2}\sigma^{\mu_3\mu_5}\mathcal{P}_L \hat{C}^*_{uW\Phi_2,g_1f_1} + c_\beta V_{f_1g_1}\sigma^{\mu_3\mu_5}\mathcal{P}_R \hat{C}_{dW\Phi_1,g_1f_2}\right)$$




$$-\frac{2i\hat{g}\hat{g}'}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta V_{g_1f_2}\sigma^{\mu_3\mu_5}\mathcal{P}_L \hat{C}^*_{uW\Phi_2,g_1f_1} - s_\beta V_{f_1g_1}\sigma^{\mu_3\mu_5}\mathcal{P}_R \hat{C}_{dW\Phi_1,g_1f_2}\right)$$




$$+\frac{2\hat{g}\hat{g}'}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta V_{g_1f_2}\sigma^{\mu_3\mu_5}\mathcal{P}_L\hat{C}^*_{uW\Phi_2,g_1f_1}-s_\beta V_{f_1g_1}\sigma^{\mu_3\mu_5}\mathcal{P}_R\hat{C}_{dW\Phi_1,g_1f_2}\right)$$




$$-\frac{2i\hat{g}\hat{g}'}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta V_{g_1f_2}\sigma^{\mu_3\mu_5}\mathcal{P}_L\hat{C}^*_{uW\Phi_2,g_1f_1}+c_\beta V_{f_1g_1}\sigma^{\mu_3\mu_5}\mathcal{P}_R\hat{C}_{dW\Phi_1,g_1f_2}\right)$$




$$-\frac{2\sqrt{2}\hat{g}c_\beta\hat{g}'\sigma^{\mu_3\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^*_{uW\Phi_2,f_2f_1}\right)$$




$$-\frac{2\sqrt{2}\hat{g}s_\beta\hat{g}'\sigma^{\mu_3\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^*_{uW\Phi_2,f_2f_1}\right)$$




$$-\sqrt{2}\hat{g}s_\beta\left(\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{uW^*\Phi_2,f_2f_1}+\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_R\right)$$




$$-i\sqrt{2}\hat{g}c_\beta\left(\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{uW^*\Phi_2,f_2f_1}-\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_R\right)$$




$$+2\hat{g}\left(s_\beta V_{f_1g_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R\hat{C}_{dW\Phi_1,g_1f_2}-c_\beta V_{g_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{uW^*\Phi_2,g_1f_1}\right)$$




$$+\sqrt{2}\hat{g}c_\beta\left(\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{uW^*\Phi_2,f_2f_1}+\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_R\right)$$




$$-i\sqrt{2}\hat{g}s_\beta\left(\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{uW^*\Phi_2,f_2f_1}-\hat{C}_{uW\Phi_2,f_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_R\right)$$




$$-2\hat{g}\left(s_\beta V_{g_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{uW^*\Phi_2,g_1f_1}+c_\beta V_{f_1g_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R\hat{C}_{dW\Phi_1,g_1f_2}\right)$$




$$+\frac{2\hat{g}^2}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta V_{g_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{uW^*\Phi_2,g_1f_1}+c_\beta V_{f_1g_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R\hat{C}_{dW\Phi_1,g_1f_2}\right)$$




$$+\frac{2i\hat{g}^2}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta V_{g_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}_{uW^*\Phi_2,g_1f_1}-s_\beta V_{f_1g_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R\hat{C}_{dW\Phi_1,g_1f_2}\right)$$




$$+\frac{2\hat{g}^2}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta V_{f_1g_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R\hat{C}_{dW\Phi_1,g_1f_2}-c_\beta V_{g_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}^*_{uW\Phi_2,g_1f_1}\right)$$




$$+\frac{2i\hat{g}^2}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta V_{g_1f_2}\sigma^{\mu_4\mu_5}\mathcal{P}_L\hat{C}^*_{uW\Phi_2,g_1f_1}+c_\beta V_{f_1g_1}\sigma^{\mu_4\mu_5}\mathcal{P}_R\hat{C}_{dW\Phi_1,g_1f_2}\right)$$




$$+\frac{2\sqrt{2}\hat{g}^2c_\beta\sigma^{\mu_4\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^*_{uW\Phi_2,f_2f_1}\right)$$




$$+\frac{2\sqrt{2}\hat{g}^2s_\beta\sigma^{\mu_4\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^*_{uW\Phi_2,f_2f_1}\right)$$




$$-\frac{2i\hat{g}\hat{g}'V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}+c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}+\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)\right)$$
$$-\frac{2i\hat{g}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)}_{\Phi u,f_1f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$+\frac{i\hat{g}s_{2\beta}\hat{g}'V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}+\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)$$
$$+\frac{i\hat{g}s_{2\beta}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)}_{\Phi u,f_1f_2}-\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$-\frac{2i\hat{g}\hat{g}'V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}+\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right)\right.$$
$$\left.+c_\beta^2\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}+c_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)$$
$$-\frac{2i\hat{g}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}^{(11)}_{\Phi u,f_1f_2}+c_\beta^2\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$-\frac{\sqrt{2}\hat{g}s_\beta c_\beta\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-\frac{\sqrt{2}\hat{g}\hat{g}'V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{i\sqrt{2}\hat{g}s_\beta c_\beta \hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-\frac{i\sqrt{2}\hat{g}\hat{g}'V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$-\frac{\hat{g}c_{2\beta}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{\hat{g}s_{2\beta}\hat{g}'V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$-\frac{i\hat{g}c_{2\beta}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-\frac{i\hat{g}s_{2\beta}\hat{g}'V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$-\frac{\hat{g}c_{2\beta}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{\hat{g}s_{2\beta}\hat{g}'V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{i\hat{g}c_{2\beta}\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{i\hat{g}s_{2\beta}\hat{g}'V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{\sqrt{2}\hat{g}s_\beta c_\beta\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-\frac{\sqrt{2}\hat{g}\hat{g}'V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{i\sqrt{2}\hat{g}s_\beta c_\beta\hat{g}'\gamma^{\mu_3}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{i\sqrt{2}\hat{g}\hat{g}'V_{f_1g_1}\gamma^{\mu_3}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$-i\sqrt{2}\hat{g}V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+i\sqrt{2}\hat{g}s_\beta c_\beta V_{f_1 g_1}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(11)[3]}_{\Phi q,g_1 f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1 f_2}\right)$$




$$-i\sqrt{2}\hat{g}V_{f_1 g_1}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1 f_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1 f_2}\right)$$




$$+\hat{g}V_{f_1 g_1}V_{f_2 g_2 *}\gamma^{\mu_5}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi q,g_1 g_2}+s^2_\beta\hat{C}^{(22)[1]}_{\Phi q,g_1 g_2}\right)$$
$$+\hat{g}\gamma^{\mu_5}\mathcal{P}_R\left(c^2_\beta\hat{C}^{(11)}_{\Phi u,f_1 f_2}+s^2_\beta\hat{C}^{(22)}_{\Phi u,f_1 f_2}\right)$$




$$-i\hat{g}V_{f_1 g_1}V^*_{f_2 g_2}\gamma^{\mu_5}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[1]}_{\Phi q,g_1 g_2}+s^2_\beta\hat{C}^{(22)[1]}_{\Phi q,g_1 g_2}\right)$$
$$-i\hat{g}\gamma^{\mu_5}\mathcal{P}_R\left(c^2_\beta\hat{C}^{(11)}_{\Phi u,f_1 f_2}+s^2_\beta\hat{C}^{(22)}_{\Phi u,f_1 f_2}\right)$$




$$+\hat{g}s_\beta c_\beta V_{f_1 g_1} V^*_{f_2 g_2} \gamma^{\mu_5} \mathcal{P}_L \left(\hat{C}^{(22)[1]}_{\Phi q, g_1 g_2} - \hat{C}^{(11)[1]}_{\Phi q, g_1 g_2} \right)$$
$$+\hat{g}s_\beta c_\beta \gamma^{\mu_5} \mathcal{P}_R \left(\hat{C}^{(22)}_{\Phi u, f_1 f_2} - \hat{C}^{(11)}_{\Phi u, f_1 f_2} \right)$$




$$-\frac{1}{2} i\hat{g}s_{2\beta} V_{f_1 g_1} V^*_{f_2 g_2} \gamma^{\mu_5} \mathcal{P}_L \left(\hat{C}^{(11)[1]}_{\Phi q, g_1 g_2} - \hat{C}^{(22)[1]}_{\Phi q, g_1 g_2} \right)$$
$$-\frac{1}{2} i\hat{g}s_{2\beta} \gamma^{\mu_5} \mathcal{P}_R \left(\hat{C}^{(11)}_{\Phi u, f_1 f_2} - \hat{C}^{(22)}_{\Phi u, f_1 f_2} \right)$$




$$+\hat{g}s_\beta c_\beta V_{f_1 g_1} V^*_{f_2 g_2} \gamma^{\mu_5} \mathcal{P}_L \left(\hat{C}^{(22)[1]}_{\Phi q, g_1 g_2} - \hat{C}^{(11)[1]}_{\Phi q, g_1 g_2} \right)$$
$$+\hat{g}s_\beta c_\beta \gamma^{\mu_5} \mathcal{P}_R \left(\hat{C}^{(22)}_{\Phi u, f_1 f_2} - \hat{C}^{(11)}_{\Phi u, f_1 f_2} \right)$$




$$+\frac{1}{2} i\hat{g}s_{2\beta} V_{f_1 g_1} V^*_{f_2 g_2} \gamma^{\mu_5} \mathcal{P}_L \left(\hat{C}^{(11)[1]}_{\Phi q, g_1 g_2} - \hat{C}^{(22)[1]}_{\Phi q, g_1 g_2} \right)$$
$$+\frac{1}{2} i\hat{g}s_{2\beta} \gamma^{\mu_5} \mathcal{P}_R \left(\hat{C}^{(11)}_{\Phi u, f_1 f_2} - \hat{C}^{(22)}_{\Phi u, f_1 f_2} \right)$$




$$+\hat{g}V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}+c^2_\beta\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}\right)$$
$$+\hat{g}\gamma^{\mu_5}\mathcal{P}_R\left(s^2_\beta\hat{C}^{(11)}_{\Phi u,f_1f_2}+c^2_\beta\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$+i\hat{g}V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}+c^2_\beta\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}\right)$$
$$+i\hat{g}\gamma^{\mu_5}\mathcal{P}_R\left(s^2_\beta\hat{C}^{(11)}_{\Phi u,f_1f_2}+c^2_\beta\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$-i\sqrt{2}\hat{g}s_\beta c_\beta\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-i\sqrt{2}\hat{g}V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+i\sqrt{2}\hat{g}s_\beta c_\beta\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-i\sqrt{2}\hat{g}V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L\left(c^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$-\frac{i\hat{g}c_{2\beta}\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{i\hat{g}s_{2\beta}V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$-\frac{i\hat{g}c_{2\beta}\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-\frac{i\hat{g}s_{2\beta}V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+i\sqrt{2}\hat{g}s_\beta c_\beta\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-i\sqrt{2}\hat{g}V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$-i\sqrt{2}\hat{g}s_\beta c_\beta\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-i\sqrt{2}\hat{g}V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+c^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{i\left(\hat{g}'^2-\hat{g}^2\right)V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}+c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}+\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)\right)$$
$$+\frac{i\left(\hat{g}'^2-\hat{g}^2\right)\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)}_{\Phi u,f_1f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$-\frac{is_{2\beta}\left(\hat{g}'^2-\hat{g}^2\right)V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L}{2\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}+\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)$$
$$-\frac{is_{2\beta}\left(\hat{g}'^2-\hat{g}^2\right)\gamma^{\mu_5}\mathcal{P}_R}{2\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)}_{\Phi u,f_1f_2}-\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$+\frac{i\left(\hat{g}'^2-\hat{g}^2\right)V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}+\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right)\right.$$
$$\left.+c_\beta^2\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}+c_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)$$
$$+\frac{i\left(\hat{g}'^2-\hat{g}^2\right)\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}^{(11)}_{\Phi u,f_1f_2}+c_\beta^2\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$+i\sqrt{\hat{g}'^2+\hat{g}^2}V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right.$$
$$\left.+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)\right)$$
$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(c_\beta^2\hat{C}^{(11)}_{\Phi u,f_1f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$+i\sqrt{\hat{g}'^2+\hat{g}^2}V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L\left(c_\beta^2\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right.$$
$$\left.+s_\beta^2\left(\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)\right)$$
$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(c_\beta^2\hat{C}^{(11)}_{\Phi u,f_1f_2}+s_\beta^2\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$-\frac{1}{2}is_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}+\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)$$
$$-\frac{1}{2}is_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(11)}_{\Phi u,f_1f_2}-\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$+\frac{1}{2}is_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}-\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}+\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)$$
$$+\frac{1}{2}is_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(11)}_{\Phi u,f_1f_2}-\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$+i\sqrt{\hat{g}'^2+\hat{g}^2}V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right)\right.$$
$$\left.+c^2_\beta\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}-c^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)$$
$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(s^2_\beta\hat{C}^{(11)}_{\Phi u,f_1f_2}+c^2_\beta\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$+i\sqrt{\hat{g}'^2+\hat{g}^2}V_{f_1g_1}V^*_{f_2g_2}\gamma^{\mu_5}\mathcal{P}_L\left(s^2_\beta\left(\hat{C}^{(11)[1]}_{\Phi q,g_1g_2}-\hat{C}^{(11)[3]}_{\Phi q,g_1g_2}\right)\right.$$
$$\left.+c^2_\beta\hat{C}^{(22)[1]}_{\Phi q,g_1g_2}-c^2_\beta\hat{C}^{(22)[3]}_{\Phi q,g_1g_2}\right)$$
$$+i\sqrt{\hat{g}'^2+\hat{g}^2}\gamma^{\mu_5}\mathcal{P}_R\left(s^2_\beta\hat{C}^{(11)}_{\Phi u,f_1f_2}+c^2_\beta\hat{C}^{(22)}_{\Phi u,f_1f_2}\right)$$




$$-\frac{\sqrt{2}\hat{g}^2 s_\beta c_\beta \gamma^{\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{\sqrt{2}\hat{g}'^2 V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{i\sqrt{2}\hat{g}^2 s_\beta c_\beta \gamma^{\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{i\sqrt{2}\hat{g}'^2 V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+s_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$-\frac{\hat{g}^2 c_{2\beta}\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{s_{2\beta}\hat{g}'^2 V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}\right)$$




$$-\frac{i\hat{g}^2 c_{2\beta}\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{is_{2\beta}\hat{g}'^2 V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$-\frac{\hat{g}^2 c_{2\beta}\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{s_{2\beta}\hat{g}'^2 V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{i\hat{g}^2 c_{2\beta}\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-\frac{is_{2\beta}\hat{g}'^2 V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}-\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{\sqrt{2}\hat{g}^2 s_\beta c_\beta\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$+\frac{\sqrt{2}\hat{g}'^2 V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+c_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$+\frac{i\sqrt{2}\hat{g}^2 s_\beta c_\beta\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$
$$-\frac{i\sqrt{2}\hat{g}'^2 V_{f_1g_1}\gamma^{\mu_5}\mathcal{P}_L}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}^{(11)[3]}_{\Phi q,g_1f_2}+c_\beta^2\hat{C}^{(22)[3]}_{\Phi q,g_1f_2}\right)$$




$$-\frac{\hat{g}c_{2\beta}\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$




$$+\frac{\hat{g}c_{2\beta}\gamma^{\mu_5}\mathcal{P}_R}{\sqrt{2}}\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$




$$-\sqrt{2}\hat{g}s_\beta c_\beta\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$




$$-\sqrt{2}\hat{g}s_\beta c_\beta\gamma^{\mu_5}\mathcal{P}_R\left(\hat{C}^{(21)}_{\Phi ud,f_1f_2}\right)$$




$$
\begin{aligned}
&-\frac{1}{4}\hat{g}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(s_\beta^2\left(2\left(\delta_{s_{\hat{\beta}\pm}}+\delta_{s_{\hat{\beta}}}-1\right)+A_1'\right)\right.\\
&\qquad\left.+c_\beta^2\left(A_2'+2\left(\delta_{c_{\hat{\beta}\pm}}+\delta_{c_{\hat{\beta}}}-1\right)\right)-2s_\beta c_\beta B'\right)\\
&-\hat{g}v^2 p_{1\mu_3}\left(2s_\beta^2 c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}+2\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(21)(12)}+c_\beta^4\hat{C}_{D\Phi}^{(21)(12)}\right)\\
&-2\hat{g}v^2 c_{2\beta}p_{1\mu_3}\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{4}\hat{g}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(s_\beta^2\left(2\left(\delta_{s_{\hat{\beta}\pm}}+\delta_{s_{\hat{\beta}}}-1\right)+A_2'\right)\right.\\
&\qquad\left.+c_\beta^2\left(A_1'+2\left(\delta_{c_{\hat{\beta}\pm}}+\delta_{c_{\hat{\beta}}}-1\right)\right)+s_{2\beta}B'\right)\\
&-4\hat{g}v^2 p_{1\mu_3}\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2 c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&-4\hat{g}v^2 s_\beta^2 c_\beta^2 p_{1\mu_3}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{4}i\hat{g}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(s_\beta^2\left(2\delta_{s_{\hat{\beta}\pm}}+A_1-2\right)+c_\beta^2\left(A_2+2\delta_{c_{\hat{\beta}\pm}}-2\right)-2Bs_\beta c_\beta\right)\\
&-i\hat{g}v^2 p_{1\mu_3}\left(\hat{C}_{D\Phi}^{(21)(12)}\right)\\
&+2i\hat{g}v^2 p_{1\mu_3}\left(s_\beta^4\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2 c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+c_\beta^4\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^2 c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{4}\hat{g}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(s_\beta c_\beta\left(2\left(\delta_{s_{\hat{\beta}\pm}}-\delta_{c_{\hat{\beta}\pm}}+\delta_{c_{\hat{\beta}}}-\delta_{s_{\hat{\beta}}}\right)-A_1'+A_2'\right)-s_\beta^2 B'+c_\beta^2 B'\right)\\
&+\hat{g}v^2 s_{2\beta}p_{1\mu_3}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-2\hat{g}v^2 s_\beta c_\beta c_{2\beta}p_{1\mu_3}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{4}\hat{g}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(s_\beta c_\beta\left(2\left(-\delta_{s_{\hat{\beta}\pm}}+\delta_{c_{\hat{\beta}\pm}}-\delta_{c_{\hat{\beta}}}+\delta_{s_{\hat{\beta}}}\right)-A_1'+A_2'\right)\right.\\
&\qquad\left.-s_\beta^2 B'+c_\beta^2 B'\right)\\
&+\hat{g}v^2 s_{2\beta}p_{1\mu_3}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+4\hat{g}v^2 s_\beta c_\beta p_{1\mu_3}\left(s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{1}{4}i\hat{g}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(s_\beta c_\beta\left(2\delta_{s_{\hat{\beta}\pm}}-A_1+A_2-2\delta_{c_{\hat{\beta}\pm}}\right)-Bs_\beta^2+Bc_\beta^2\right)\\
&+2i\hat{g}v^2 s_\beta c_\beta p_{2\mu_3}\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
-iv^2\left(c_\beta^4-s_\beta^4\right)\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_3}+p_{2\mu_3}\right)\left(\hat{C}_{D\Phi}^{(21)(21)}-\hat{C}_{D\Phi}^{(21)(21)*}\right)
$$




$$
\begin{aligned}
&-\frac{p_{1\mu_3}-p_{2\mu_3}}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\left(\left(\hat{g}'^2+\hat{g}^2\right)\left(A_2'+A_2+2\delta_{c_{\hat{\beta}}}-2\right)-2\hat{g}X_{WB}\hat{g}'\right)\right.\\
&\qquad+s_\beta^2\left(\left(\hat{g}'^2+\hat{g}^2\right)\left(A_1'+A_1+2\delta_{s_{\hat{\beta}}}-2\right)-2\hat{g}X_{WB}\hat{g}'\right)\\
&\qquad\left.-2s_\beta c_\beta\left(\hat{g}'^2+\hat{g}^2\right)\left(B+B'\right)\right)\\
&-v^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_3}\left(2s_\beta^2c_\beta^2\left(3\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}-2\hat{C}_{D\Phi}^{(21)(12)}+3\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\right.\\
&\qquad\left.+s_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+c_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\\
&\qquad-p_{2\mu_3}\left(2s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}+\hat{C}_{D\Phi}^{(22)(22)}\right)+s_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.\left.+c_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\right)\\
&-v^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(\left(s_\beta^4+c_\beta^4-4s_\beta^2c_\beta^2\right)p_{1\mu_3}+\left(s_\beta^4+c_\beta^4+4s_\beta^2c_\beta^2\right)p_{2\mu_3}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{p_{1\mu_3}-p_{2\mu_3}}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\left(\left(\hat{g}'^2+\hat{g}^2\right)\left(A_1'+A_1+2\delta_{c_{\hat{\beta}}}-2\right)-2\hat{g}X_{WB}\hat{g}'\right)\right.\\
&\qquad+s_\beta^2\left(\left(\hat{g}'^2+\hat{g}^2\right)\left(A_2'+A_2+2\delta_{s_{\hat{\beta}}}-2\right)-2\hat{g}X_{WB}\hat{g}'\right)\\
&\qquad\left.+s_{2\beta}\left(\hat{g}'^2+\hat{g}^2\right)\left(B+B'\right)\right)\\
&+2v^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(3p_{1\mu_3}-p_{2\mu_3}\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&+2v^2s_\beta^2c_\beta^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(3p_{1\mu_3}-p_{2\mu_3}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$-\frac{i\left(\hat{g}'^2-\hat{g}^2\right)\left(p_{1\mu_3}-p_{2\mu_3}\right)}{2\left(\hat{g}'^2+\hat{g}^2\right)^{3/2}}\left(s_\beta^2\left(\left(\hat{g}'^2+\hat{g}^2\right)\left(2\delta_{s_{\hat{\beta}^\pm}}-1\right)+\hat{g}X_{WB}\hat{g}'\right)\right.$$
$$\left.+c_\beta^2\left(\left(\hat{g}'^2+\hat{g}^2\right)\left(2\delta_{c_{\hat{\beta}^\pm}}-1\right)+\hat{g}X_{WB}\hat{g}'\right)\right)$$
$$-2iv^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)$$
$$-2iv^2s_\beta^2c_\beta^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+iv^2\left(c_\beta^4-s_\beta^4\right)\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_3}+p_{2\mu_3}\right)\left(\hat{C}_{D\Phi}^{(21)(21)}-\hat{C}_{D\Phi}^{(21)(21)*}\right)$$




$$-\frac{i\left(\hat{g}'^2-\hat{g}^2\right)\left(p_{1\mu_3}-p_{2\mu_3}\right)}{2\left(\hat{g}'^2+\hat{g}^2\right)^{3/2}}\left(s_\beta^2\left(\left(\hat{g}'^2+\hat{g}^2\right)\left(2\delta_{s_{\hat{\beta}^\pm}}-1\right)+\hat{g}X_{WB}\hat{g}'\right)\right.$$
$$\left.+c_\beta^2\left(\left(\hat{g}'^2+\hat{g}^2\right)\left(2\delta_{c_{\hat{\beta}^\pm}}-1\right)+\hat{g}X_{WB}\hat{g}'\right)\right)$$
$$-iv^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(2s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(11)(22)}+c_\beta^4\hat{C}_{D\Phi}^{(11)(22)}\right)$$
$$+2iv^2s_\beta^2c_\beta^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+iv^2s_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_3}-3p_{2\mu_3}\right)\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.$$
$$\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+\frac{1}{4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(s_\beta c_\beta\left(-A_1'+A_2'-A_1+A_2+2\delta_{c_{\hat{\beta}}}-2\delta_{s_{\hat{\beta}}}\right)\right.$$
$$\left.-s_\beta^2\left(B+B'\right)+c_\beta^2\left(B+B'\right)\right)$$
$$-v^2s_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(3p_{1\mu_3}-p_{2\mu_3}\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.$$
$$\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)$$
$$+v^2s_\beta c_\beta c_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}\left(3p_{1\mu_3}-p_{2\mu_3}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$
\begin{aligned}
&-\frac{1}{4}\sqrt{\hat{g}'^2+\hat{g}^2}\,(p_{1\mu_3}-p_{2\mu_3})\left(s_\beta c_\beta\left(-A_1'+A_2'-A_1+A_2-2\delta_{c_{\hat{\beta}}}+2\delta_{s_{\hat{\beta}}}\right)\right.\\
&\qquad\left.-s_\beta^2\,(B+B')+c_\beta^2\,(B+B')\right)\\
&+v^2 s_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\,(3p_{1\mu_3}-p_{2\mu_3})\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+v^2 s_\beta c_\beta c_{2\beta}\left(-\sqrt{\hat{g}'^2+\hat{g}^2}\right)(3p_{1\mu_3}-p_{2\mu_3})\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+iv^2 s_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\,(p_{1\mu_3}-p_{2\mu_3})\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-2iv^2 s_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\,(p_{1\mu_3}-p_{2\mu_3})\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{4}\,(K-L)\sqrt{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2 p_{1\mu_3}-s_\beta^2 p_{2\mu_3}+c_\beta^2 p_{1\mu_3}-c_\beta^2 p_{2\mu_3}\right)\\
&+iv^2 s_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\,(p_{1\mu_3}-3p_{2\mu_3})\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{i\hat{g}v g_{\mu_2\mu_3}\hat{g}'}{\sqrt{2}\,(\hat{g}'^2+\hat{g}^2)^{3/2}}\left(s_\beta^2\left(\hat{g}'^3\left(\delta_{s_{\hat{\beta}^\pm}}-1\right)+\hat{g}^2\hat{g}'\left(\delta_{s_{\hat{\beta}^\pm}}-1\right)-\hat{g}^3 X_{WB}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\left(\delta_{c_{\hat{\beta}^\pm}}-1\right)\hat{g}'^3+\hat{g}^2\left(\delta_{c_{\hat{\beta}^\pm}}-1\right)\hat{g}'-\hat{g}^3 X_{WB}\right)\right)\\
&+2i\sqrt{2}\hat{g}v^3 g_{\mu_2\mu_3}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2 c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&-\frac{2i\sqrt{2}v\hat{g}'\,(p_{2\mu_3}p_{3\mu_2}-p_2\cdot p_3 g_{\mu_2\mu_3})}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{\Phi WB}^{(11)}+s_\beta^2\hat{C}_{\Phi WB}^{(22)}\right)\\
&-\frac{2i\sqrt{2}v\hat{g}'p_2^\mu p_3^\nu\epsilon_{\mu_2\mu_3\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(11)}+s_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\\
&+2i\sqrt{2}\hat{g}v^3 s_\beta^2 c_\beta^2 g_{\mu_2\mu_3}\sqrt{\hat{g}'^2+\hat{g}^2}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{i\hat{g}vc_\beta g_{\mu_2\mu_3}\hat{g}'^2\left(\delta_{c_{\hat{\beta}^\pm}}-\delta_{s_{\hat{\beta}^\pm}}\right)}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta\right)\\
&-i\sqrt{2}\hat{g}v^3 s_\beta c_\beta g_{\mu_2\mu_3}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\frac{2i\sqrt{2}vs_\beta c_\beta\hat{g}'\left(p_{2\mu_3}p_{3\mu_2}-p_2\cdot p_3 g_{\mu_2\mu_3}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi WB}^{(11)}-\hat{C}_{\Phi WB}^{(22)}\right)\\
&+\frac{2i\sqrt{2}vs_\beta c_\beta\hat{g}'p_2^\mu p_3^\nu\epsilon_{\mu_2\mu_3\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi B\tilde{W}}^{(11)}-\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\\
&+2i\sqrt{2}\hat{g}v^3 s_\beta c_\beta g_{\mu_2\mu_3}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{ivg_{\mu_2\mu_3}}{2\sqrt{2}}\left(s_\beta^2\left(\left(A_2-2\right)\hat{g}'^2+\left(A_2-2\right)\hat{g}^2-4\hat{g}X_{WB}\hat{g}'\right)\right.\\
&\qquad+c_\beta^2\left(\left(A_1-2\right)\hat{g}'^2+\left(A_1-2\right)\hat{g}^2-4\hat{g}X_{WB}\hat{g}'\right)\\
&\qquad\left.+Bs_{2\beta}\left(\hat{g}'^2+\hat{g}^2\right)\right)\\
&-4i\sqrt{2}v^3 g_{\mu_2\mu_3}\left(\hat{g}'^2+\hat{g}^2\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&+\frac{4i\sqrt{2}v\left(p_{2\mu_3}p_{3\mu_2}-p_2\cdot p_3 g_{\mu_2\mu_3}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(11)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(11)}+\hat{g}\hat{C}_{\Phi W}^{(11)}\right)\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(22)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(22)}+\hat{g}\hat{C}_{\Phi W}^{(22)}\right)\right)\right)\\
&+\frac{4i\sqrt{2}vp_2^\mu p_3^\nu\epsilon_{\mu_2\mu_3\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}\right)\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right)\right)\\
&-4i\sqrt{2}v^3 s_\beta^2c_\beta^2 g_{\mu_2\mu_3}\left(\hat{g}'^2+\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-2\sqrt{2}v^3 s_\beta c_\beta g_{\mu_2\mu_3}\left(\hat{g}'^2+\hat{g}^2\right)\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{ivg_{\mu_2\mu_3}\left(\hat{g}'^2+\hat{g}^2\right)}{2\sqrt{2}}\left(\left(A_2-A_1\right)s_\beta c_\beta - Bs_\beta^2 + Bc_\beta^2\right)\\
&-2i\sqrt{2}v^3 s_\beta c_\beta g_{\mu_2\mu_3}\left(\hat{g}'^2+\hat{g}^2\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\frac{4i\sqrt{2}vs_\beta c_\beta\left(p_{2\mu_3}p_{3\mu_2}-p_2\cdot p_3 g_{\mu_2\mu_3}\right)}{\hat{g}'^2+\hat{g}^2}\left(\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(11)}+\hat{g}\hat{C}_{\Phi W}^{(11)}-\hat{g}'\hat{C}_{\Phi WB}^{(22)}-\hat{g}\hat{C}_{\Phi W}^{(22)}\right)\right.\\
&\qquad\left.+\hat{g}'^2\hat{C}_{\Phi B}^{(11)}-\hat{g}'^2\hat{C}_{\Phi B}^{(22)}\right)\\
&+\frac{4i\sqrt{2}vs_\beta c_\beta p_2^\mu p_3^\nu \epsilon_{\mu_2\mu_3\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}-\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right.\\
&\qquad\left.+\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}-\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}\right)\\
&+2i\sqrt{2}v^3 s_\beta c_\beta c_{2\beta} g_{\mu_2\mu_3}\left(\hat{g}'^2+\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{4i\sqrt{2}v\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi B}^{(11)}-\hat{g}'\hat{C}_{\Phi WB}^{(11)}\right)+\hat{g}'^2\hat{C}_{\Phi W}^{(11)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi B}^{(22)}-\hat{g}'\hat{C}_{\Phi WB}^{(22)}\right)+\hat{g}'^2\hat{C}_{\Phi W}^{(22)}\right)\right)\\
&+\frac{4i\sqrt{2}vp_1^\mu p_2^\nu \epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi\tilde{B}}^{(11)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)+\hat{g}'^2\hat{C}_{\Phi\tilde{W}}^{(11)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi\tilde{B}}^{(22)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)+\hat{g}'^2\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{i\hat{g}^2 v g_{\mu_1\mu_3}\hat{g}'}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(\delta_{s_{\hat{\beta}^\pm}}-1\right)+c_\beta^2\left(\delta_{c_{\hat{\beta}^\pm}}-1\right)+\frac{\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}\right)\\
&+\frac{2i\sqrt{2}\hat{g}v\left(p_{1\mu_3}p_{3\mu_1}-p_1\cdot p_3 g_{\mu_1\mu_3}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{\Phi WB}^{(11)}+s_\beta^2\hat{C}_{\Phi WB}^{(22)}\right)\\
&+\frac{2i\sqrt{2}\hat{g}vp_1^\mu p_3^\nu \epsilon_{\mu_1\mu_3\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(11)}+s_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{i\hat{g}^2 v g_{\mu_2\mu_3}}{2\sqrt{2}}\left(\left(A_2-2\right)s_\beta^2+\left(A_1-2\right)c_\beta^2+Bs_{2\beta}\right)\\
&+4i\sqrt{2}v\left(p_{2\mu_3}p_{3\mu_2}-p_2\cdot p_3 g_{\mu_2\mu_3}\right)\left(c_\beta^2\hat{C}_{\Phi W}^{(11)}+s_\beta^2\hat{C}_{\Phi W}^{(22)}\right)\\
&+4i\sqrt{2}vp_2^\mu p_3^\nu \epsilon_{\mu_2\mu_3\mu\nu}\left(c_\beta^2\hat{C}_{\Phi\tilde{W}}^{(11)}+s_\beta^2\hat{C}_{\Phi\tilde{W}}^{(22)}\right)
\end{aligned}
$$




$$+\frac{2i\sqrt{2}v\left(p_{1\mu_3}p_{3\mu_1}-p_1\cdot p_3 g_{\mu_1\mu_3}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi W}^{(11)}-\hat{C}_{\Phi B}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(11)}\right)\right.$$
$$\left.+s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi W}^{(22)}-\hat{C}_{\Phi B}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(22)}\right)\right)$$
$$+\frac{2i\sqrt{2}vp_1^\mu p_3^\nu\epsilon_{\mu_1\mu_3\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{C}_{\Phi\tilde{B}}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)\right.$$
$$\left.+s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{W}}^{(22)}-\hat{C}_{\Phi\tilde{B}}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)$$




$$+\frac{4i\sqrt{2}vs_\beta c_\beta\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2+\hat{g}^2}\left(\hat{g}\left(-\hat{g}'\hat{C}_{\Phi WB}^{(11)}+\hat{g}\hat{C}_{\Phi B}^{(11)}+\hat{g}'\hat{C}_{\Phi WB}^{(22)}-\hat{g}\hat{C}_{\Phi B}^{(22)}\right)\right.$$
$$\left.+\hat{g}'^2\hat{C}_{\Phi W}^{(11)}-\hat{g}'^2\hat{C}_{\Phi W}^{(22)}\right)$$
$$+\frac{4i\sqrt{2}vs_\beta c_\beta p_1^\mu p_2^\nu\epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(\hat{g}\left(-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}+\hat{g}\hat{C}_{\Phi\tilde{B}}^{(11)}+\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}-\hat{g}\hat{C}_{\Phi\tilde{B}}^{(22)}\right)\right.$$
$$\left.+\hat{g}'^2\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{g}'^2\hat{C}_{\Phi\tilde{W}}^{(22)}\right)$$




$$-\frac{i\hat{g}^2vc_\beta g_{\mu_1\mu_3}\hat{g}'\left(\delta_{c_{\hat{\beta}^\pm}}-\delta_{s_{\hat{\beta}^\pm}}\right)}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta\right)$$
$$-\frac{2i\sqrt{2}\hat{g}vs_\beta c_\beta\left(p_{1\mu_3}p_{3\mu_1}-p_1\cdot p_3 g_{\mu_1\mu_3}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi WB}^{(11)}-\hat{C}_{\Phi WB}^{(22)}\right)$$
$$-\frac{2i\sqrt{2}\hat{g}vs_\beta c_\beta p_1^\mu p_3^\nu\epsilon_{\mu_1\mu_3\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi B\tilde{W}}^{(11)}-\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)$$




$$+\frac{i\hat{g}^2vg_{\mu_2\mu_3}}{2\sqrt{2}}\left(\left(A_2-A_1\right)s_\beta c_\beta-Bs_\beta^2+Bc_\beta^2\right)$$
$$+4i\sqrt{2}vs_\beta c_\beta\left(p_{2\mu_3}p_{3\mu_2}-p_2\cdot p_3 g_{\mu_2\mu_3}\right)\left(\hat{C}_{\Phi W}^{(11)}-\hat{C}_{\Phi W}^{(22)}\right)$$
$$+4i\sqrt{2}vs_\beta c_\beta p_2^\mu p_3^\nu\epsilon_{\mu_2\mu_3\mu\nu}\left(\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{C}_{\Phi\tilde{W}}^{(22)}\right)$$




$$+\frac{2i\sqrt{2}vs_\beta c_\beta\left(p_{1\mu_3}p_{3\mu_1}-p_1\cdot p_3 g_{\mu_1\mu_3}\right)}{\hat{g}'^2+\hat{g}^2}\left(2\hat{g}\hat{g}'\left(-\hat{C}_{\Phi B}^{(11)}+\hat{C}_{\Phi W}^{(11)}+\hat{C}_{\Phi B}^{(22)}-\hat{C}_{\Phi W}^{(22)}\right)\right.$$
$$\left.+\left(\hat{g}'^2-\hat{g}^2\right)\left(\hat{C}_{\Phi WB}^{(11)}-\hat{C}_{\Phi WB}^{(22)}\right)\right)$$
$$+\frac{2i\sqrt{2}vs_\beta c_\beta p_1^\mu p_3^\nu\epsilon_{\mu_1\mu_3\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(2\hat{g}\hat{g}'\left(-\hat{C}_{\Phi\tilde{B}}^{(11)}+\hat{C}_{\Phi\tilde{W}}^{(11)}+\hat{C}_{\Phi\tilde{B}}^{(22)}-\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right.$$
$$\left.+\left(\hat{g}'^2-\hat{g}^2\right)\left(\hat{C}_{\Phi B\tilde{W}}^{(11)}-\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)$$




$$+\frac{i\hat{g}\hat{g}'\left(p_{2\mu_1}-p_{3\mu_1}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(2\delta_{s_{\hat{\beta}\pm}}-1\right)+c_\beta^2\left(2\delta_{c_{\hat{\beta}\pm}}-1\right)+\frac{\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}\right)$$




$$+\frac{i\hat{g}\hat{g}'\left(p_{2\mu_1}-p_{3\mu_1}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(2\delta_{s_{\hat{\beta}\pm}}-1\right)+c_\beta^2\left(2\delta_{c_{\hat{\beta}\pm}}-1\right)+\frac{\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}\right)$$




$$-\frac{1}{4}i\hat{g}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(s_\beta^2\left(2\delta_{s_{\hat{\beta}\pm}}+A_2-2\right)+c_\beta^2\left(A_1+2\delta_{c_{\hat{\beta}\pm}}-2\right)+Bs_{2\beta}\right)$$




$$+\frac{1}{4}i\hat{g}\left(p_{1\mu_3}-p_{2\mu_3}\right)\left(s_\beta c_\beta\left(-2\delta_{s_{\hat{\beta}\pm}}-A_1+A_2+2\delta_{c_{\hat{\beta}\pm}}\right)-Bs_\beta^2+Bc_\beta^2\right)$$




$$-\frac{\sqrt{2}\hat{g}v\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(2s_\beta^2c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}+2\hat{C}_{D\Phi}^{(22)(22)}\right)\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(21)(12)}+c_\beta^4\hat{C}_{D\Phi}^{(21)(12)}\right)$$
$$-\frac{2\sqrt{2}\hat{g}vc_{2\beta}\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$-\frac{4\sqrt{2}\hat{g}v\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)$$
$$-\frac{4\sqrt{2}\hat{g}vs_\beta^2c_\beta^2\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$-\frac{2\sqrt{2}\hat{g}v\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(2s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(11)(22)}+c_\beta^4\hat{C}_{D\Phi}^{(11)(22)}\right)$$
$$+\frac{4\sqrt{2}\hat{g}vs_\beta^2c_\beta^2\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+\frac{i\sqrt{2}\hat{g}v\hat{g}'p_{3\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{D\Phi}^{(21)(12)}\right)$$
$$-\frac{2i\sqrt{2}\hat{g}v\hat{g}'p_{3\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^4\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+\frac{\sqrt{2}\hat{g}vs_{2\beta}\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.$$
$$\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)$$
$$-\frac{2\sqrt{2}\hat{g}vs_\beta c_\beta c_{2\beta}\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+\frac{\sqrt{2}\hat{g}vs_{2\beta}\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.$$
$$\left.+c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)$$
$$+\frac{\sqrt{2}\hat{g}vs_{2\beta}c_{2\beta}\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+\frac{\sqrt{2}\hat{g}vs_{2\beta}\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.$$
$$\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)$$
$$+\frac{4\sqrt{2}\hat{g}vs_\beta c_\beta\hat{g}'p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)$$



$$+\frac{2i\sqrt{2}\hat{g}vs_\beta c_\beta\hat{g}'p_{4\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.$$
$$\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$-\frac{2i\sqrt{2}\hat{g}vs_\beta c_\beta \hat{g}' p_{2\mu_1}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$-\frac{i\hat{g}v\left(p_{1\mu_4}+p_{2\mu_4}\right)}{\sqrt{2}}\left(2s_\beta^2c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}+2\hat{C}_{D\Phi}^{(22)(22)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(21)(12)}+c_\beta^4\hat{C}_{D\Phi}^{(21)(12)}\right)$$

$$-i\sqrt{2}\hat{g}vc_{2\beta}\left(p_{1\mu_4}+p_{2\mu_4}\right)\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+\frac{\hat{g}v}{\sqrt{2}}\left(p_{1\mu_4}\left(2s_\beta^2c_\beta^2\left(4\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}-3\hat{C}_{D\Phi}^{(21)(12)}+4\hat{C}_{D\Phi}^{(22)(22)}\right)+s_\beta^4\left(2\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+c_\beta^4\left(2\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\right.$$
$$\left.-p_{3\mu_4}\left(2s_\beta^2c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(21)(12)}+2\hat{C}_{D\Phi}^{(22)(22)}\right)+s_\beta^4\left(2\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+c_\beta^4\left(2\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\right)$$

$$+\sqrt{2}\hat{g}v\left(c_\beta^4p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+c_\beta^4p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-3s_\beta^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+3s_\beta^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^4p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^4p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-3s_\beta^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+3s_\beta^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$-\frac{i\hat{g}v}{\sqrt{2}}\left(4s_\beta^2c_\beta^2p_{1\mu_4}\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)+p_{2\mu_4}\left(2s_\beta^2c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(21)(12)}+2\hat{C}_{D\Phi}^{(22)(22)}\right)+s_\beta^4\left(2\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+c_\beta^4\left(2\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\right)$$


$$-i\sqrt{2}\hat{g}v\left(2s_\beta^4p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-s_\beta^4p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-3s_\beta^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+2c_\beta^4p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-c_\beta^4p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-3s_\beta^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$
\begin{aligned}
&-\frac{\hat{g}v\left(2p_{1\mu_4}-p_{2\mu_4}\right)}{\sqrt{2}}\left(2s_\beta^2c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}+2\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(21)(12)}+c_\beta^4\hat{C}_{D\Phi}^{(21)(12)}\right)\\
&-\sqrt{2}\hat{g}vc_{2\beta}\left(2p_{1\mu_4}-p_{2\mu_4}\right)\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-2i\sqrt{2}\hat{g}v\left(p_{1\mu_4}+p_{2\mu_4}\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&-2i\sqrt{2}\hat{g}vs_\beta^2c_\beta^2\left(p_{1\mu_4}+p_{2\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-2\sqrt{2}\hat{g}v\left(2p_{1\mu_4}-p_{3\mu_4}\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&-2\sqrt{2}\hat{g}vs_\beta^2c_\beta^2\left(2p_{1\mu_4}-p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{\hat{g}v}{\sqrt{2}}\left(4s_\beta^2c_\beta^2p_{2\mu_4}\left(-\hat{C}_{D\Phi}^{(11)(11)}+\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\qquad+p_{1\mu_4}\left(2s_\beta^2c_\beta^2\left(4\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}-3\hat{C}_{D\Phi}^{(21)(12)}+4\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\qquad\left.\left.+s_\beta^4\left(2\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+c_\beta^4\left(2\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\right)\\
&+\sqrt{2}\hat{g}v\left(s_\beta^4p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-2s_\beta^4p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad-3s_\beta^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+c_\beta^4p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad\left.-2c_\beta^4p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-3s_\beta^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$+2i\sqrt{2}\hat{g}v\left(2p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)$$
$$+2i\sqrt{2}\hat{g}vs_\beta^2c_\beta^2\left(2p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+2i\sqrt{2}\hat{g}vs_\beta^2c_\beta^2\left(2p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}\right.$$
$$\left.-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)$$
$$+2i\sqrt{2}\hat{g}v\left(2p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(s_\beta^4\hat{C}_{D\Phi}^{(21)(21)*}+c_\beta^4\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+\frac{i\hat{g}v\left(p_{1\mu_4}-2p_{2\mu_4}\right)}{\sqrt{2}}\left(\hat{C}_{D\Phi}^{(21)(12)}\right)$$
$$-i\sqrt{2}\hat{g}v\left(p_{1\mu_4}-2p_{2\mu_4}\right)\left(s_\beta^4\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.$$
$$\left.+c_\beta^4\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$-\frac{i\hat{g}v\left(p_{2\mu_4}+p_{3\mu_4}\right)}{\sqrt{2}}\left(\hat{C}_{D\Phi}^{(21)(12)}\right)$$
$$+i\sqrt{2}\hat{g}v\left(p_{2\mu_4}+p_{3\mu_4}\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$
\begin{aligned}
&+i\sqrt{2}\hat{g}vs_\beta c_\beta\left(p_{1\mu_4}+p_{2\mu_4}\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-i\sqrt{2}\hat{g}vs_\beta c_\beta\left(-2s_\beta^2 p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2 p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad+3c_\beta^2 p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-3s_\beta^2 p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad\left.+2c_\beta^2 p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-c_\beta^2 p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\sqrt{2}\hat{g}vs_\beta c_\beta\left(2p_{1\mu_4}-p_{3\mu_4}\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-\sqrt{2}\hat{g}vs_\beta c_\beta c_{2\beta}\left(2p_{1\mu_4}-p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+i\sqrt{2}\hat{g}vs_\beta c_\beta\left(p_{1\mu_4}+p_{2\mu_4}\right)\left(s_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+i\sqrt{2}\hat{g}vs_\beta c_\beta c_{2\beta}\left(p_{1\mu_4}+p_{2\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\sqrt{2}\hat{g}vs_\beta c_\beta\left(2p_{1\mu_4}-p_{2\mu_4}\right)\left(s_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\sqrt{2}\hat{g}vs_\beta c_\beta\left(3s_\beta^2 p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-3s_\beta^2 p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad-c_\beta^2 p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2 p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad+s_\beta^2 p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^2 p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad\left.-3c_\beta^2 p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+3c_\beta^2 p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\sqrt{2}\hat{g}vs_\beta c_\beta\left(2p_{1\mu_4}-p_{3\mu_4}\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\sqrt{2}\hat{g}vs_\beta c_\beta\left(-s_\beta^2 p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+2s_\beta^2 p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad+3c_\beta^2 p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-3s_\beta^2 p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad\left.+c_\beta^2 p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-2c_\beta^2 p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+i\sqrt{2}\hat{g}vs_\beta c_\beta\left(p_{1\mu_4}+p_{2\mu_4}\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-2i\sqrt{2}\hat{g}vs_\beta c_\beta\left(p_{1\mu_4}+p_{2\mu_4}\right)\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\sqrt{2}\hat{g}vs_\beta c_\beta\left(2p_{1\mu_4}-p_{2\mu_4}\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-2\sqrt{2}\hat{g}vs_\beta c_\beta\left(2p_{1\mu_4}-p_{2\mu_4}\right)\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-i\sqrt{2}\hat{g}vs_\beta c_\beta\left(2p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+2i\sqrt{2}\hat{g}vs_\beta c_\beta\left(2p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$



$$-i\sqrt{2}\hat{g}vs_\beta c_\beta\left(p_{2\mu_4}-2p_{3\mu_4}\right)\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+i\sqrt{2}\hat{g}vs_\beta c_\beta\left(p_{1\mu_4}+p_{2\mu_4}\right)\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$-i\sqrt{2}\hat{g}vs_\beta c_\beta\left(2p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(s_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)+c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)$$

$$-2i\sqrt{2}\hat{g}vs_\beta c_\beta\left(2p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)$$




$$-8i\hat{g}^2v^2g_{\mu_3\mu_4}\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)$$


$$-8i\hat{g}^2v^2s_\beta^2c_\beta^2g_{\mu_3\mu_4}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$
\begin{aligned}
&-8i\hat{g}^2v^2s_\beta^2c_\beta^2g_{\mu_3\mu_4}\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&-8i\hat{g}^2v^2g_{\mu_3\mu_4}\left(s_\beta^4\hat{C}_{D\Phi}^{(21)(21)*}+c_\beta^4\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+4i\hat{g}^2v^2s_\beta c_\beta g_{\mu_3\mu_4}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-8i\hat{g}^2v^2s_\beta c_\beta g_{\mu_3\mu_4}\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{i\hat{g}g_{\mu_1\mu_4}\hat{g}'\left(\hat{g}'^2-\hat{g}^2\right)}{\left(\hat{g}'^2+\hat{g}^2\right)^2}\left(s_{\hat{\beta}}^2\left(\hat{g}'^2\left(2\delta_{s_{\hat{\beta}^\pm}}-1\right)+\hat{g}^2\left(2\delta_{s_{\hat{\beta}^\pm}}-1\right)\right)\right.\\
&\qquad+c_{\hat{\beta}}^2\left(\hat{g}^2\left(2\delta_{c_{\hat{\beta}^\pm}}-1\right)+\left(2\delta_{c_{\hat{\beta}^\pm}}-1\right)\hat{g}'^2\right)\\
&\qquad\left.+2\hat{g}X_{WB}\hat{g}'\right)\\
&+4i\hat{g}v^2g_{\mu_1\mu_4}\hat{g}'\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&-\frac{2i\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4g_{\mu_1\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi B}^{(11)}-\hat{C}_{\Phi W}^{(11)}\right)\right.\right.\\
&\qquad\left.+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(11)}\right)\\
&\qquad+s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi B}^{(22)}-\hat{C}_{\Phi W}^{(22)}\right)\right.\\
&\qquad\left.\left.+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(22)}\right)\right)\\
&-\frac{2ip_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{B}}^{(11)}-\hat{C}_{\Phi\tilde{W}}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{B}}^{(22)}-\hat{C}_{\Phi\tilde{W}}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)\\
&+4i\hat{g}v^2s_\beta^2c_\beta^2g_{\mu_1\mu_4}\hat{g}'\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{i\hat{g}g_{\mu_1\mu_4}\hat{g}'\left(\hat{g}'^2-\hat{g}^2\right)}{\left(\hat{g}'^2+\hat{g}^2\right)^2}\left(s_\beta^2\left(\hat{g}'^2\left(2\delta_{s_{\hat{\beta}^\pm}}-1\right)+\hat{g}^2\left(2\delta_{s_{\hat{\beta}^\pm}}-1\right)\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{g}^2\left(2\delta_{c_{\hat{\beta}^\pm}}-1\right)+\left(2\delta_{c_{\hat{\beta}^\pm}}-1\right)\hat{g}'^2\right)\right.\\
&\qquad\left.+2\hat{g}X_{WB}\hat{g}'\right)\\
&+2i\hat{g}v^2g_{\mu_1\mu_4}\hat{g}'\left(2s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(11)(22)}+c_\beta^4\hat{C}_{D\Phi}^{(11)(22)}\right)\\
&-\frac{2i\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4g_{\mu_1\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi B}^{(11)}-\hat{C}_{\Phi W}^{(11)}\right)\right.\right.\\
&\qquad\left.+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(11)}\right)\\
&\qquad\left.+c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi B}^{(22)}-\hat{C}_{\Phi W}^{(22)}\right)\right.\right.\\
&\qquad\left.\left.+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(22)}\right)\right)\\
&-\frac{2ip_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{B}}^{(11)}-\hat{C}_{\Phi\tilde{W}}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)\right.\\
&\qquad\left.+c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{B}}^{(22)}-\hat{C}_{\Phi\tilde{W}}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)\\
&-4i\hat{g}v^2s_\beta^2c_\beta^2g_{\mu_1\mu_4}\hat{g}'\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\left(2s_\beta^2c_\beta^2\left(p_{1\mu_4}+p_{2\mu_4}\right)\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}\right.\right.\\
&\qquad\left.-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&\qquad+p_{3\mu_4}\left(2s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\qquad\left.\left.+s_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+c_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\right)\\
&+\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\left(\left(s_\beta^4+c_\beta^4\right)\left(p_{1\mu_4}+p_{2\mu_4}\right)\right.\\
&\qquad\left.-\left(s_\beta^4+c_\beta^4+4s_\beta^2c_\beta^2\right)p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$



$$-i\sqrt{2}v\left(c_\beta^4-s_\beta^4\right)\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}+p_{2\mu_4}-p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)}-\hat{C}_{D\Phi}^{(21)(21)*}\right)$$




$$-i\sqrt{2}v\left(c_\beta^4-s_\beta^4\right)\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}-p_{2\mu_4}+p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)}-\hat{C}_{D\Phi}^{(21)(21)*}\right)$$




$$
\begin{aligned}
&-\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\left(2s_\beta^2c_\beta^2p_{2\mu_4}\left(-\hat{C}_{D\Phi}^{(11)(11)}+\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\quad -p_{3\mu_4}\left(2s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\quad \left.+s_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+c_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\\
&\quad +p_{1\mu_4}\left(2s_\beta^2c_\beta^2\left(3\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}\right.\right.\\
&\quad \left.-2\hat{C}_{D\Phi}^{(21)(12)}+3\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&\quad \left.\left.+s_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+c_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\right)\\
&-\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\left(\left(s_\beta^4+c_\beta^4\right)\left(-p_{2\mu_4}+p_{1\mu_4}+p_{3\mu_4}\right)\right.\\
&\quad \left.-4s_\beta^2c_\beta^2\left(p_{1\mu_4}-p_{3\mu_4}\right)\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$


$$
\begin{aligned}
&+\frac{v}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\left(\hat{g}'^2-\hat{g}^2\right)p_{1\mu_4}\left(2s_\beta^2c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}\right.\right.\right.\\
&\quad \left.-\hat{C}_{D\Phi}^{(21)(12)}+2\hat{C}_{D\Phi}^{(22)(22)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(21)(12)}+c_\beta^4\hat{C}_{D\Phi}^{(21)(12)}\Big)\\
&\quad \left.+\left(\hat{g}'^2+\hat{g}^2\right)\left(p_{2\mu_4}-p_{3\mu_4}\right)\hat{C}_{D\Phi}^{(21)(12)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{\sqrt{2}v}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(-\hat{g}^2c_\beta^4p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-\hat{g}^2c_\beta^4p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\quad +\hat{g}^2c_\beta^4p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2s_\beta^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\quad -\hat{g}^2s_\beta^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2s_\beta^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\quad +c_\beta^4\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^4\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\quad +c_\beta^4\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-s_\beta^2c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\quad -s_\beta^2c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\quad -\hat{g}^2s_\beta^4p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-\hat{g}^2s_\beta^4p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\quad +\hat{g}^2s_\beta^4p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2s_\beta^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\quad -\hat{g}^2s_\beta^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2s_\beta^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\quad +s_\beta^4\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^4\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\quad +s_\beta^4\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\quad \left.-s_\beta^2c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^2c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+2\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}+p_{2\mu_4}+p_{3\mu_4}\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}\right.\\
&\quad \left.+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&+2\sqrt{2}vs_\beta^2c_\beta^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}+p_{2\mu_4}+p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$+\frac{2\sqrt{2}v\left(\hat{g}'^2-\hat{g}^2\right)p_{1\mu_4}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)$$
$$+\frac{2\sqrt{2}vs_\beta^2c_\beta^2\left(\hat{g}'^2-\hat{g}^2\right)p_{1\mu_4}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+2\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\left(3p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}\right.$$
$$\left.+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)$$
$$+2\sqrt{2}vs_\beta^2c_\beta^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(3p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$



$$+\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}\left(2s_\beta^2c_\beta^2\left(3\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}-2\hat{C}_{D\Phi}^{(21)(12)}+3\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\right.$$
$$\left.+s_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)+c_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)$$
$$-2s_\beta^2c_\beta^2\left(p_{2\mu_4}+p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}\right.$$
$$\left.\left.-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right)$$
$$+\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\left(\left(s_\beta^4+c_\beta^4-4s_\beta^2c_\beta^2\right)p_{1\mu_4}\right.$$
$$\left.-\left(s_\beta^4+c_\beta^4\right)\left(p_{2\mu_4}+p_{3\mu_4}\right)\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$+\frac{\sqrt{2}v\left(\hat{g}'^2-\hat{g}^2\right)p_{1\mu_4}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(2s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(11)(22)}+c_\beta^4\hat{C}_{D\Phi}^{(11)(22)}\right)$$
$$-\frac{2\sqrt{2}vs_\beta^2c_\beta^2\left(\hat{g}'^2-\hat{g}^2\right)p_{1\mu_4}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$-2i\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}-p_{2\mu_4}\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)$$
$$-2i\sqrt{2}vs_\beta^2c_\beta^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}-p_{2\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$


$$+\frac{iv}{\sqrt{2}\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\left(\hat{g}'^2+\hat{g}^2\right)p_{1\mu_4}\left(2s_\beta^2c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}\right.\right.\right.$$
$$\left.-\hat{C}_{D\Phi}^{(21)(12)}+2\hat{C}_{D\Phi}^{(22)(22)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(21)(12)}+c_\beta^4\hat{C}_{D\Phi}^{(21)(12)}\Big)$$
$$-\left(\hat{g}'^2+\hat{g}^2\right)p_{3\mu_4}\left(2s_\beta^2c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}\right.\right.$$
$$\left.\left.-\hat{C}_{D\Phi}^{(21)(12)}+2\hat{C}_{D\Phi}^{(22)(22)}\right)+s_\beta^4\hat{C}_{D\Phi}^{(21)(12)}+c_\beta^4\hat{C}_{D\Phi}^{(21)(12)}\right)$$
$$\left.+\left(\hat{g}'^2-\hat{g}^2\right)p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(12)}\right)$$




$$+\frac{i\sqrt{2}v}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}^2s_\beta^4p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2s_\beta^4p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\right.$$
$$-\hat{g}^2s_\beta^4p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-\hat{g}^2s_\beta^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}$$
$$+\hat{g}^2s_\beta^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2s_\beta^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}$$
$$+s_\beta^4\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-s_\beta^4\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}$$
$$-s_\beta^4\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-s_\beta^2c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}$$
$$-s_\beta^2c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}$$
$$+\hat{g}^2c_\beta^4p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2c_\beta^4p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}$$
$$-\hat{g}^2c_\beta^4p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-\hat{g}^2s_\beta^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}$$
$$+\hat{g}^2s_\beta^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2s_\beta^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}$$
$$+c_\beta^4\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-c_\beta^4\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}$$
$$-c_\beta^4\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}$$
$$\left.-s_\beta^2c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^2c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$-i\sqrt{2}v\left(c_\beta^4-s_\beta^4\right)\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)}-\hat{C}_{D\Phi}^{(21)(21)*}\right)$$




$$
\begin{aligned}
&-i\sqrt{2}v\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{2\mu_4}-p_{3\mu_4}\right)\left(2s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(11)(22)}+c_\beta^4\hat{C}_{D\Phi}^{(11)(22)}\right)\\
&+2i\sqrt{2}vs_\beta^2c_\beta^2\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{2\mu_4}-p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-2i\hat{g}v^2s_\beta c_\beta g_{\mu_1\mu_4}\hat{g}'\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\frac{2is_\beta c_\beta\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4g_{\mu_1\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi B}^{(11)}-\hat{C}_{\Phi W}^{(11)}-\hat{C}_{\Phi B}^{(22)}+\hat{C}_{\Phi W}^{(22)}\right)\right.\\
&\qquad\left.+\left(\hat{g}'^2-\hat{g}^2\right)\left(\hat{C}_{\Phi WB}^{(11)}-\hat{C}_{\Phi WB}^{(22)}\right)\right)\\
&+\frac{2is_\beta c_\beta p_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{B}}^{(11)}-\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{C}_{\Phi\tilde{B}}^{(22)}+\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right.\\
&\qquad\left.+\left(\hat{g}'^2-\hat{g}^2\right)\left(\hat{C}_{\Phi B\tilde{W}}^{(11)}-\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)\\
&+4i\hat{g}v^2s_\beta c_\beta g_{\mu_1\mu_4}\hat{g}'\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}+p_{2\mu_4}+p_{3\mu_4}\right)\\
&\qquad\left(s_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-\sqrt{2}vs_\beta c_\beta c_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}+p_{2\mu_4}+p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}+p_{2\mu_4}+p_{3\mu_4}\right)\\
&\qquad\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\sqrt{2}vs_\beta c_\beta c_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}+p_{2\mu_4}+p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{\sqrt{2}vs_\beta c_\beta\left(\hat{g}'^2-\hat{g}^2\right)p_{1\mu_4}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\frac{\sqrt{2}vs_\beta c_\beta}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}^2s_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2s_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad-\hat{g}^2s_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-\hat{g}^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad+\hat{g}^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-\hat{g}^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad-s_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad-s_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad+c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad+\hat{g}^2s_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-\hat{g}^2s_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad+\hat{g}^2s_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-\hat{g}^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad-\hat{g}^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad-s_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad+s_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad\left.-c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}-3p_{2\mu_4}+p_{3\mu_4}\right)\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(3p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\\
&\qquad\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\sqrt{2}vs_\beta c_\beta c_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}\left(3p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4}\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}+p_{2\mu_4}-3p_{3\mu_4}\right)\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\,(3p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4})\\
&\qquad\left(s_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-\sqrt{2}vs_\beta c_\beta c_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}\,(3p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4})\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{\sqrt{2}vs_\beta c_\beta\left(\hat{g}'^2-\hat{g}^2\right)p_{1\mu_4}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\frac{\sqrt{2}vs_\beta c_\beta}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(-\hat{g}^2s_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-\hat{g}^2s_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad+\hat{g}^2s_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad-\hat{g}^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad+s_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-s_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad+s_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad-c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad-\hat{g}^2s_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2s_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad-\hat{g}^2s_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad+\hat{g}^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-\hat{g}^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad+s_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad-s_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-c_\beta^2\hat{g}'^2p_{1\mu_4}-\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad\left.+c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\,(p_{1\mu_4}+p_{2\mu_4}+p_{3\mu_4})\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\,(3p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4})\\
&\qquad\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-\sqrt{2}vs_\beta c_\beta c_{2\beta}\sqrt{\hat{g}'^2+\hat{g}^2}\,(3p_{1\mu_4}-p_{2\mu_4}-p_{3\mu_4})\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{\sqrt{2}vs_\beta c_\beta\left(\hat{g}'^2-\hat{g}^2\right)p_{1\mu_4}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\frac{\sqrt{2}vs_{2\beta}\left(\hat{g}'^2-\hat{g}^2\right)p_{1\mu_4}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}-p_{2\mu_4}\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\frac{i\sqrt{2}vs_\beta c_\beta}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(-\hat{g}^2s_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2s_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad-\hat{g}^2s_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad-\hat{g}^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-\hat{g}^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad-s_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad+s_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad-c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\\
&\qquad-\hat{g}^2s_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2s_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad+\hat{g}^2s_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad-\hat{g}^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad-s_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad-s_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\\
&\qquad\left.-c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$



$$
\begin{aligned}
&+i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}-p_{3\mu_4}\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-2i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}-p_{3\mu_4}\right)\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}+p_{2\mu_4}-3p_{3\mu_4}\right)\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$+i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{1\mu_4}+p_{2\mu_4}+p_{3\mu_4}\right)\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.$$
$$\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)$$


$$-i\sqrt{2}vs_\beta c_\beta\sqrt{\hat{g}'^2+\hat{g}^2}\left(p_{2\mu_4}-p_{3\mu_4}\right)\left(s_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.$$
$$\left.+c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)$$




$$+\frac{i\sqrt{2}vs_\beta c_\beta}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{g}^2s_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2s_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}\right.$$
$$-\hat{g}^2s_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}$$
$$-\hat{g}^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+\hat{g}^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}$$
$$-s_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+s_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}$$
$$-s_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}$$
$$-c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}+c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)*}$$
$$-\hat{g}^2s_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2s_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}$$
$$-\hat{g}^2s_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}-\hat{g}^2c_\beta^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}$$
$$-\hat{g}^2c_\beta^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+\hat{g}^2c_\beta^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}$$
$$+s_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+s_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}$$
$$-s_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{g}'^2p_{1\mu_4}\hat{C}_{D\Phi}^{(21)(21)}$$
$$\left.-c_\beta^2\hat{g}'^2p_{2\mu_4}\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{g}'^2p_{3\mu_4}\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$-\frac{\hat{g}g_{\mu_3\mu_4}\hat{g}'}{4\left(\hat{g}'^2+\hat{g}^2\right)^{3/2}}\left(s_\beta^2\left(\hat{g}'^3\left(2\left(\delta_{s_{\hat{\beta}^\pm}}+\delta_{s_{\hat{\beta}}}-1\right)+A_2'\right)\right.\right.$$
$$+\hat{g}^2\hat{g}'\left(2\left(\delta_{s_{\hat{\beta}^\pm}}+\delta_{s_{\hat{\beta}}}-1\right)+A_2'\right)-2\hat{g}^3X_{WB}\Big)$$
$$+c_\beta^2\left(\hat{g}'^3\left(A_1'+2\left(\delta_{c_{\hat{\beta}^\pm}}+\delta_{c_{\hat{\beta}}}-1\right)\right)\right.$$
$$\left.+\hat{g}^2\hat{g}'\left(A_1'+2\left(\delta_{c_{\hat{\beta}^\pm}}+\delta_{c_\beta}-1\right)\right)-2\hat{g}^3X_{WB}\right)$$
$$\left.+s_{2\beta}B'\hat{g}'\left(\hat{g}'^2+\hat{g}^2\right)\right)$$
$$-2\hat{g}v^2g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.$$
$$\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)$$
$$+\frac{2\hat{g}'\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4g_{\mu_3\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{\Phi WB}^{(11)}+s_\beta^2\hat{C}_{\Phi WB}^{(22)}\right)$$
$$+\frac{2\hat{g}'p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(11)}+s_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)$$
$$-2\hat{g}v^2s_\beta^2c_\beta^2g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)$$




$$
\begin{aligned}
&+\frac{i\hat{g}g_{\mu_3\mu_4}\hat{g}'}{4\left(\hat{g}'^2+\hat{g}^2\right)^{3/2}}\left(s_\beta^2\left(\hat{g}'^3\left(2\delta_{s_{\hat{\beta}\pm}}+A_2-2\right)\right.\right.\\
&\qquad\left.+\hat{g}^2\hat{g}'\left(2\delta_{s_{\hat{\beta}\pm}}+A_2-2\right)-2\hat{g}^3X_{WB}\right)\\
&\qquad+c_\beta^2\left(\hat{g}'^3\left(A_1+2\delta_{c_{\hat{\beta}\pm}}-2\right)\right.\\
&\qquad\left.+\hat{g}^2\hat{g}'\left(A_1+2\delta_{c_{\hat{\beta}\pm}}-2\right)-2\hat{g}^3X_{WB}\right)\\
&\qquad+Bs_{2\beta}\hat{g}'\left(\hat{g}'^2+\hat{g}^2\right)\Big)\\
&+6i\hat{g}v^2g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&-\frac{2i\hat{g}'\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4g_{\mu_3\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{\Phi WB}^{(11)}+s_\beta^2\hat{C}_{\Phi WB}^{(22)}\right)\\
&-\frac{2i\hat{g}'p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(11)}+s_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\\
&+6i\hat{g}v^2s_\beta^2c_\beta^2g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{\hat{g}g_{\mu_3\mu_4}\hat{g}'}{4\left(\hat{g}'^2+\hat{g}^2\right)^{3/2}}\left(s_\beta^2\left(\hat{g}'^3\left(2\left(\delta_{s_{\hat{\beta}\pm}}+\delta_{s_{\hat{\beta}}}-1\right)+A_1'\right)\right.\right.\\
&\qquad\left.+\hat{g}^2\hat{g}'\left(2\left(\delta_{s_{\hat{\beta}\pm}}+\delta_{s_{\hat{\beta}}}-1\right)+A_1'\right)-2\hat{g}^3X_{WB}\right)\\
&\qquad+c_\beta^2\left(\hat{g}'^3\left(A_2'+2\left(\delta_{c_{\hat{\beta}\pm}}+\delta_{c_{\hat{\beta}}}-1\right)\right)\right.\\
&\qquad\left.+\hat{g}^2\hat{g}'\left(A_2'+2\left(\delta_{c_{\hat{\beta}\pm}}+\delta_{c_{\hat{\beta}}}-1\right)\right)-2\hat{g}^3X_{WB}\right)\\
&\qquad-2s_\beta c_\beta B'\hat{g}'\left(\hat{g}'^2+\hat{g}^2\right)\Big)\\
&-\hat{g}v^2g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(2s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\qquad+s_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\\
&\qquad\left.+c_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\\
&+\frac{2\hat{g}'\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4g_{\mu_3\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}_{\Phi WB}^{(11)}+c_\beta^2\hat{C}_{\Phi WB}^{(22)}\right)\\
&+\frac{2\hat{g}'p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(11)}+c_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\\
&+2\hat{g}v^2g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(s_\beta^4\hat{C}_{D\Phi}^{(21)(21)*}+2s_\beta^2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+c_\beta^4\hat{C}_{D\Phi}^{(21)(21)}+2s_\beta^2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{i\hat{g}g_{\mu_3\mu_4}\hat{g}'}{4\left(\hat{g}'^2+\hat{g}^2\right)^{3/2}}\left(s_\beta^2\left(\hat{g}'^3\left(2\delta_{s_{\hat{\beta}\pm}}+A_1-2\right)\right.\right.\\
&\qquad+\hat{g}^2\hat{g}'\left(2\delta_{s_{\hat{\beta}\pm}}+A_1-2\right)-2\hat{g}^3X_{WB}\Big)\\
&\qquad+c_\beta^2\left(\hat{g}'^3\left(A_2+2\delta_{c_{\hat{\beta}\pm}}-2\right)\right.\\
&\qquad\left.+\hat{g}^2\hat{g}'\left(A_2+2\delta_{c_{\hat{\beta}\pm}}-2\right)-2\hat{g}^3X_{WB}\right)\\
&\qquad\left.-2Bs_\beta c_\beta\hat{g}'\left(\hat{g}'^2+\hat{g}^2\right)\right)\\
&-i\hat{g}v^2g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(2s_\beta^2c_\beta^2\left(3\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}\right.\right.\\
&\qquad\left.-2\hat{C}_{D\Phi}^{(21)(12)}+3\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&\qquad+s_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\\
&\qquad\left.+c_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\\
&+\frac{2i\hat{g}'\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4g_{\mu_3\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}_{\Phi WB}^{(11)}+c_\beta^2\hat{C}_{\Phi WB}^{(22)}\right)\\
&+\frac{2i\hat{g}'p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(11)}+c_\beta^2\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\\
&-2i\hat{g}v^2g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(s_\beta^4\hat{C}_{D\Phi}^{(21)(21)*}-2s_\beta^2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+c_\beta^4\hat{C}_{D\Phi}^{(21)(21)}-2s_\beta^2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{\hat{g}g_{\mu_3\mu_4}\hat{g}'^2}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta c_\beta\left(2\left(-\delta_{s_{\hat{\beta}\pm}}+\delta_{c_{\hat{\beta}\pm}}-\delta_{c_{\hat{\beta}}}+\delta_{s_{\hat{\beta}}}\right)+A_1'-A_2'\right)+s_\beta^2B'-c_\beta^2B'\right)\\
&+\hat{g}v^2s_\beta c_\beta g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-\frac{2s_\beta c_\beta\hat{g}'\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4g_{\mu_3\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi WB}^{(11)}-\hat{C}_{\Phi WB}^{(22)}\right)\\
&+\frac{s_{2\beta}\hat{g}'p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi B\tilde{W}}^{(22)}-\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)\\
&+\hat{g}v^2s_{2\beta}g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+2s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{i\hat{g}g_{\mu_3\mu_4}\hat{g}'^2}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta c_\beta\left(2\delta_{s_{\hat{\beta}\pm}}-A_1+A_2-2\delta_{c_{\hat{\beta}\pm}}\right)-Bs_\beta^2+Bc_\beta^2\right)\\
&+3i\hat{g}v^2s_\beta c_\beta g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-\frac{2is_\beta c_\beta\hat{g}'\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4g_{\mu_3\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi WB}^{(11)}-\hat{C}_{\Phi WB}^{(22)}\right)\\
&-\frac{2is_\beta c_\beta\hat{g}'p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi B\tilde{W}}^{(11)}-\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\\
&-2i\hat{g}v^2s_\beta c_\beta g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}+2c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.-2s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{\hat{g}g_{\mu_3\mu_4}\hat{g}'^2}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta c_\beta\left(2\left(\delta_{s_{\hat{\beta}\pm}}-\delta_{c_{\hat{\beta}\pm}}+\delta_{c_{\hat{\beta}}}-\delta_{s_{\hat{\beta}}}\right)+A_1'-A_2'\right)+s_\beta^2 B'-c_\beta^2 B'\right)\\
&+\hat{g}v^2 s_\beta c_\beta g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-\frac{2s_\beta c_\beta\hat{g}'\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi WB}^{(11)}-\hat{C}_{\Phi WB}^{(22)}\right)\\
&+\frac{s_{2\beta}\hat{g}'p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi B\tilde{W}}^{(22)}-\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)\\
&-2\hat{g}v^2 s_\beta c_\beta g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{i\hat{g}g_{\mu_3\mu_4}\hat{g}'^2}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta c_\beta\left(-2\delta_{s_{\hat{\beta}\pm}}-A_1+A_2+2\delta_{c_{\hat{\beta}\pm}}\right)-Bs_\beta^2+Bc_\beta^2\right)\\
&-3i\hat{g}v^2 s_\beta c_\beta g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\frac{2is_\beta c_\beta\hat{g}'\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi WB}^{(11)}-\hat{C}_{\Phi WB}^{(22)}\right)\\
&+\frac{2is_\beta c_\beta\hat{g}'p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}_{\Phi B\tilde{W}}^{(11)}-\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\\
&+6i\hat{g}v^2 s_\beta c_\beta g_{\mu_3\mu_4}\sqrt{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{2}ig_{\mu_3\mu_4}\left(c_\beta^2\left(\hat{g}^2\left(A_2'+2\delta_{c_{\hat{\beta}}}-1\right)+\hat{g}'^2\left(A_2'+2\delta_{c_{\hat{\beta}}}-1\right)-2\hat{g}X_{WB}\hat{g}'\right)\right.\\
&\qquad+s_\beta^2\left(\hat{g}^2\left(A_1'+2\delta_{s_{\hat{\beta}}}-1\right)+\hat{g}'^2\left(A_1'+2\delta_{s_{\hat{\beta}}}-1\right)-2\hat{g}X_{WB}\hat{g}'\right)\\
&\qquad\left.-2s_\beta c_\beta B'\left(\hat{g}'^2+\hat{g}^2\right)\right)\\
&-2iv^2 g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(2s_\beta^2 c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\qquad+s_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\\
&\qquad\left.+c_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\\
&+\frac{4i\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(11)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(11)}+\hat{g}\hat{C}_{\Phi W}^{(11)}\right)\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(22)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(22)}+\hat{g}\hat{C}_{\Phi W}^{(22)}\right)\right)\right)\\
&+\frac{4ip_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}\right)\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right)\right)\\
&+2iv^2 g_{\mu_3\mu_4}\left(s_\beta^4+c_\beta^4+4s_\beta^2 c_\beta^2\right)\left(\hat{g}'^2+\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{2}ig_{\mu_3\mu_4}\left(c_\beta^2\left(\hat{g}^2\left(A_1'+2\delta_{c_{\hat{\beta}}}-1\right)+\hat{g}'^2\left(A_1'+2\delta_{c_{\hat{\beta}}}-1\right)-2\hat{g}X_{WB}\hat{g}'\right)\right.\\
&\qquad+s_\beta^2\left(\hat{g}^2\left(A_2'+2\delta_{s_{\hat{\beta}}}-1\right)+\hat{g}'^2\left(A_2'+2\delta_{s_{\hat{\beta}}}-1\right)-2\hat{g}X_{WB}\hat{g}'\right)\\
&\qquad\left.+s_{2\beta}B'\left(\hat{g}'^2+\hat{g}^2\right)\right)\\
&-4iv^2g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&+\frac{4i\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4g_{\mu_3\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(11)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(11)}+\hat{g}\hat{C}_{\Phi W}^{(11)}\right)\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(22)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(22)}+\hat{g}\hat{C}_{\Phi W}^{(22)}\right)\right)\right)\\
&+\frac{4ip_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}\right)\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right)\right)\\
&-4iv^2s_\beta^2c_\beta^2g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{ig_{\mu_3\mu_4}\left(\hat{g}'^2-\hat{g}^2\right)^2}{2\left(\hat{g}'^2+\hat{g}^2\right)^2}\left(s_\beta^2\left(\hat{g}'^2\left(2\delta_{s_{\hat{\beta}^\pm}}-1\right)+\hat{g}^2\left(2\delta_{s_{\hat{\beta}^\pm}}-1\right)+2\hat{g}X_{WB}\hat{g}'\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{g}^2\left(2\delta_{c_{\hat{\beta}^\pm}}-1\right)+\left(2\delta_{c_{\hat{\beta}^\pm}}-1\right)\hat{g}'^2+2\hat{g}X_{WB}\hat{g}'\right)\right)\\
&-4iv^2g_{\mu_3\mu_4}\left(\hat{g}'^2-\hat{g}^2\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&+\frac{4i\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4g_{\mu_3\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(11)}+\hat{g}\left(\hat{g}\hat{C}_{\Phi W}^{(11)}-\hat{g}'\hat{C}_{\Phi WB}^{(11)}\right)\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(22)}+\hat{g}\left(\hat{g}\hat{C}_{\Phi W}^{(22)}-\hat{g}'\hat{C}_{\Phi WB}^{(22)}\right)\right)\right)\\
&+\frac{4ip_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}+\hat{g}\left(\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}+\hat{g}\left(\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)\right)\\
&-4iv^2s_\beta^2c_\beta^2g_{\mu_3\mu_4}\left(\hat{g}'^2-\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{2}ig_{\mu_3\mu_4}\left(s_\beta^2\left(\left(A_2-1\right)\hat{g}'^2+\left(A_2-1\right)\hat{g}^2-2\hat{g}X_{WB}\hat{g}'\right)\right.\\
&\qquad+c_\beta^2\left(\left(A_1-1\right)\hat{g}'^2+\left(A_1-1\right)\hat{g}^2-2\hat{g}X_{WB}\hat{g}'\right)\\
&\qquad\left.+Bs_{2\beta}\left(\hat{g}'^2+\hat{g}^2\right)\right)\\
&-2iv^2g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(c_\beta^4\hat{C}_{D\Phi}^{(11)(11)}+s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^4\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&+\frac{4i\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4g_{\mu_3\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(11)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(11)}+\hat{g}\hat{C}_{\Phi W}^{(11)}\right)\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(22)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(22)}+\hat{g}\hat{C}_{\Phi W}^{(22)}\right)\right)\right)\\
&+\frac{4ip_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}\right)\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right)\right)\\
&-2iv^2s_\beta^2c_\beta^2g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$+2v^2 g_{\mu_3\mu_4}\left(c_\beta^4 - s_\beta^4\right)\left(\hat{g}'^2+\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)} - \hat{C}_{D\Phi}^{(21)(21)*}\right)$$




$$\begin{aligned}
&-\frac{1}{2}ig_{\mu_3\mu_4}\left(s_\beta^2\left(\left(A_1-1\right)\hat{g}'^2+\left(A_1-1\right)\hat{g}^2-2\hat{g}X_{WB}\hat{g}'\right)\right.\\
&\qquad +c_\beta^2\left(\left(A_2-1\right)\hat{g}'^2+\left(A_2-1\right)\hat{g}^2-2\hat{g}X_{WB}\hat{g}'\right)\\
&\qquad \left.-2Bs_\beta c_\beta\left(\hat{g}'^2+\hat{g}^2\right)\right)\\
&-2iv^2 g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(2s_\beta^2c_\beta^2\left(3\hat{C}_{D\Phi}^{(11)(11)}-2\hat{C}_{D\Phi}^{(11)(22)}\right.\right.\\
&\qquad \left.-2\hat{C}_{D\Phi}^{(21)(12)}+3\hat{C}_{D\Phi}^{(22)(22)}\right)\\
&\qquad +s_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\\
&\qquad \left.+c_\beta^4\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}\right)\right)\\
&+\frac{4i\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(11)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(11)}+\hat{g}\hat{C}_{\Phi W}^{(11)}\right)\right)\right.\\
&\qquad \left.+c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(22)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(22)}+\hat{g}\hat{C}_{\Phi W}^{(22)}\right)\right)\right)\\
&+\frac{4ip_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}\right)\right)\right.\\
&\qquad \left.+c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}+\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right)\right)\\
&-2iv^2 g_{\mu_3\mu_4}\left(s_\beta^4+c_\beta^4-4s_\beta^2c_\beta^2\right)\left(\hat{g}'^2+\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}$$




$$\begin{aligned}
&-\frac{ig_{\mu_3\mu_4}\left(\hat{g}'^2-\hat{g}^2\right)^2}{2\left(\hat{g}'^2+\hat{g}^2\right)^2}\left(s_\beta^2\left(\hat{g}'^2\left(2\delta_{s_{\hat{\beta}\pm}}-1\right)+\hat{g}^2\left(2\delta_{s_{\hat{\beta}\pm}}-1\right)+2\hat{g}X_{WB}\hat{g}'\right)\right.\\
&\qquad \left.+c_\beta^2\left(\hat{g}^2\left(2\delta_{c_{\hat{\beta}\pm}}-1\right)+\left(2\delta_{c_{\hat{\beta}\pm}}-1\right)\hat{g}'^2+2\hat{g}X_{WB}\hat{g}'\right)\right)\\
&-2iv^2 g_{\mu_3\mu_4}\left(\hat{g}'^2-\hat{g}^2\right)\left(2s_\beta^2c_\beta^2\left(\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(21)(12)}+\hat{C}_{D\Phi}^{(22)(22)}\right)\right.\\
&\qquad \left.+s_\beta^4\hat{C}_{D\Phi}^{(11)(22)}+c_\beta^4\hat{C}_{D\Phi}^{(11)(22)}\right)\\
&+\frac{4i\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(11)}+\hat{g}\left(\hat{g}\hat{C}_{\Phi W}^{(11)}-\hat{g}'\hat{C}_{\Phi WB}^{(11)}\right)\right)\right.\\
&\qquad \left.+c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi B}^{(22)}+\hat{g}\left(\hat{g}\hat{C}_{\Phi W}^{(22)}-\hat{g}'\hat{C}_{\Phi WB}^{(22)}\right)\right)\right)\\
&+\frac{4ip_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}+\hat{g}\left(\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)\right)\right.\\
&\qquad \left.+c_\beta^2\left(\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}+\hat{g}\left(\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)\right)\\
&+4iv^2s_\beta^2c_\beta^2g_{\mu_3\mu_4}\left(\hat{g}'^2-\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}$$




$$
\begin{aligned}
&-\frac{1}{2}ig_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(\left(A_2'-A_1'\right)s_\beta c_\beta-s_\beta^2B'+c_\beta^2B'\right)\\
&+2iv^2s_\beta c_\beta g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-\frac{4is_\beta c_\beta\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4g_{\mu_3\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(11)}+\hat{g}\hat{C}_{\Phi W}^{(11)}\right.\right.\\
&\qquad\left.-\hat{g}'\hat{C}_{\Phi WB}^{(22)}-\hat{g}\hat{C}_{\Phi W}^{(22)}\right)\\
&\qquad\left.+\hat{g}'^2\hat{C}_{\Phi B}^{(11)}-\hat{g}'^2\hat{C}_{\Phi B}^{(22)}\right)\\
&-\frac{4is_\beta c_\beta p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}-\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right.\\
&\qquad\left.+\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}-\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}\right)\\
&-2iv^2s_\beta c_\beta c_{2\beta}g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-6v^2s_\beta c_\beta g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-2v^2s_\beta c_\beta g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}\right.\\
&\qquad\left.+s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}+c_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{1}{2}ig_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(\left(A_2-A_1\right)s_\beta c_\beta - Bs_\beta^2 + Bc_\beta^2\right)\\
&-6iv^2 s_\beta c_\beta g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&+\frac{4is_\beta c_\beta\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(11)}+\hat{g}\hat{C}_{\Phi W}^{(11)}-\hat{g}'\hat{C}_{\Phi WB}^{(22)}-\hat{g}\hat{C}_{\Phi W}^{(22)}\right)\right.\\
&\qquad\left.+\hat{g}'^2\hat{C}_{\Phi B}^{(11)}-\hat{g}'^2\hat{C}_{\Phi B}^{(22)}\right)\\
&+\frac{4is_\beta c_\beta p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(\hat{g}\left(\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}-\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right.\\
&\qquad\left.+\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}-\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}\right)\\
&+6iv^2 s_\beta c_\beta c_{2\beta} g_{\mu_3\mu_4}\left(\hat{g}'^2+\hat{g}^2\right)\left(\hat{C}_{D\Phi}^{(21)(21)*}+\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+2iv^2 s_\beta c_\beta g_{\mu_3\mu_4}\left(\hat{g}'^2-\hat{g}^2\right)\left(c_\beta^2\left(2\hat{C}_{D\Phi}^{(11)(11)}-\hat{C}_{D\Phi}^{(11)(22)}-\hat{C}_{D\Phi}^{(21)(12)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{C}_{D\Phi}^{(11)(22)}+\hat{C}_{D\Phi}^{(21)(12)}-2\hat{C}_{D\Phi}^{(22)(22)}\right)\right)\\
&-\frac{4is_\beta c_\beta\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(-\hat{g}\left(\hat{g}'\hat{C}_{\Phi WB}^{(11)}+\hat{g}\hat{C}_{\Phi W}^{(11)}\right.\right.\\
&\qquad\left.\left.+\hat{g}'\hat{C}_{\Phi WB}^{(22)}-\hat{g}\hat{C}_{\Phi W}^{(22)}\right)\right.\\
&\qquad\left.+\hat{g}'^2\hat{C}_{\Phi B}^{(11)}-\hat{g}'^2\hat{C}_{\Phi B}^{(22)}\right)\\
&-\frac{4is_\beta c_\beta p_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(-\hat{g}\left(\hat{g}\hat{C}_{\Phi B\tilde{W}}^{(11)}+\hat{g}\hat{C}_{\Phi\tilde{W}}^{(11)}\right.\right.\\
&\qquad\left.\left.+\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}-\hat{g}\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right.\\
&\qquad\left.+\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(11)}-\hat{g}'^2\hat{C}_{\Phi\tilde{B}}^{(22)}\right)\\
&-4iv^2 s_\beta c_\beta g_{\mu_3\mu_4}\left(\hat{g}'^2-\hat{g}^2\right)\left(c_\beta^2\hat{C}_{D\Phi}^{(21)(21)*}-s_\beta^2\hat{C}_{D\Phi}^{(21)(21)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{4i\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi B}^{(11)}-\hat{g}'\hat{C}_{\Phi WB}^{(11)}\right)+\hat{g}'^2\hat{C}_{\Phi W}^{(11)}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi B}^{(22)}-\hat{g}'\hat{C}_{\Phi WB}^{(22)}\right)+\hat{g}'^2\hat{C}_{\Phi W}^{(22)}\right)\right)\\
&+\frac{4ip_1^\mu p_2^\nu\epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi\tilde{B}}^{(11)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)+\hat{g}'^2\hat{C}_{\Phi\tilde{W}}^{(11)}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi\tilde{B}}^{(22)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)+\hat{g}'^2\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{4i\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi B}^{(11)}-\hat{g}'\hat{C}_{\Phi WB}^{(11)}\right)+\hat{g}'^2\hat{C}_{\Phi W}^{(11)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi B}^{(22)}-\hat{g}'\hat{C}_{\Phi WB}^{(22)}\right)+\hat{g}'^2\hat{C}_{\Phi W}^{(22)}\right)\right)\\
&+\frac{4ip_1^\mu p_2^\nu\epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi\tilde{B}}^{(11)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)+\hat{g}'^2\hat{C}_{\Phi\tilde{W}}^{(11)}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}_{\Phi\tilde{B}}^{(22)}-\hat{g}'\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)+\hat{g}'^2\hat{C}_{\Phi\tilde{W}}^{(22)}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{2i\hat{g}^2 g_{\mu_1\mu_2}\hat{g}'^2}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(2\delta_{s_{\hat{\beta}^\pm}}-1\right)+c_\beta^2\left(2\delta_{c_{\hat{\beta}^\pm}}-1\right)+\frac{2\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}\right)\\
&+\frac{4i\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}\left(\hat{g}'\hat{C}^{(11)}_{\Phi WB}+\hat{g}\hat{C}^{(11)}_{\Phi B}\right)+\hat{g}'^2\hat{C}^{(11)}_{\Phi W}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}\left(\hat{g}'\hat{C}^{(22)}_{\Phi WB}+\hat{g}\hat{C}^{(22)}_{\Phi B}\right)+\hat{g}'^2\hat{C}^{(22)}_{\Phi W}\right)\right)\\
&+\frac{4ip_1^\mu p_2^\nu\epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}\left(\hat{g}'\hat{C}^{(11)}_{\Phi B\tilde{W}}+\hat{g}\hat{C}^{(11)}_{\Phi\tilde{B}}\right)+\hat{g}'^2\hat{C}^{(11)}_{\Phi\tilde{W}}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}\left(\hat{g}'\hat{C}^{(22)}_{\Phi B\tilde{W}}+\hat{g}\hat{C}^{(22)}_{\Phi\tilde{B}}\right)+\hat{g}'^2\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{4i\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}^{(11)}_{\Phi B}-\hat{g}'\hat{C}^{(11)}_{\Phi WB}\right)+\hat{g}'^2\hat{C}^{(11)}_{\Phi W}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}^{(22)}_{\Phi B}-\hat{g}'\hat{C}^{(22)}_{\Phi WB}\right)+\hat{g}'^2\hat{C}^{(22)}_{\Phi W}\right)\right)\\
&+\frac{4ip_1^\mu p_2^\nu\epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}^{(11)}_{\Phi\tilde{B}}-\hat{g}'\hat{C}^{(11)}_{\Phi B\tilde{W}}\right)+\hat{g}'^2\hat{C}^{(11)}_{\Phi\tilde{W}}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}^{(22)}_{\Phi\tilde{B}}-\hat{g}'\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)+\hat{g}'^2\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{4i\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}^{(11)}_{\Phi B}-\hat{g}'\hat{C}^{(11)}_{\Phi WB}\right)+\hat{g}'^2\hat{C}^{(11)}_{\Phi W}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}^{(22)}_{\Phi B}-\hat{g}'\hat{C}^{(22)}_{\Phi WB}\right)+\hat{g}'^2\hat{C}^{(22)}_{\Phi W}\right)\right)\\
&+\frac{4ip_1^\mu p_2^\nu\epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}^{(11)}_{\Phi\tilde{B}}-\hat{g}'\hat{C}^{(11)}_{\Phi B\tilde{W}}\right)+\hat{g}'^2\hat{C}^{(11)}_{\Phi\tilde{W}}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{g}\left(\hat{g}\hat{C}^{(22)}_{\Phi\tilde{B}}-\hat{g}'\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)+\hat{g}'^2\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{2i\hat{g}^2 g_{\mu_1\mu_2}\hat{g}'^2}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(2\delta_{s_{\hat{\beta}^\pm}}-1\right)+c_\beta^2\left(2\delta_{c_{\hat{\beta}^\pm}}-1\right)+\frac{2\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}\right)\\
&+\frac{4i\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}\left(\hat{g}'\hat{C}^{(11)}_{\Phi WB}+\hat{g}\hat{C}^{(11)}_{\Phi B}\right)+\hat{g}'^2\hat{C}^{(11)}_{\Phi W}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{g}\left(\hat{g}'\hat{C}^{(22)}_{\Phi WB}+\hat{g}\hat{C}^{(22)}_{\Phi B}\right)+\hat{g}'^2\hat{C}^{(22)}_{\Phi W}\right)\right)\\
&+\frac{4ip_1^\mu p_2^\nu\epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(\hat{g}\left(\hat{g}'\hat{C}^{(11)}_{\Phi B\tilde{W}}+\hat{g}\hat{C}^{(11)}_{\Phi\tilde{B}}\right)+\hat{g}'^2\hat{C}^{(11)}_{\Phi\tilde{W}}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{g}\left(\hat{g}'\hat{C}^{(22)}_{\Phi B\tilde{W}}+\hat{g}\hat{C}^{(22)}_{\Phi\tilde{B}}\right)+\hat{g}'^2\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{\hat{g}^2 g_{\mu_1\mu_4}\hat{g}'}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(2\left(\delta_{s_{\hat{\beta}^\pm}}+\delta_{s_{\hat{\beta}}}-1\right)+A_2'\right)\right.\\
&\qquad\left.+c_\beta^2\left(A_1'+2\left(\delta_{c_{\hat{\beta}^\pm}}+\delta_{c_{\hat{\beta}}}-1\right)\right)\right.\\
&\qquad\left.+s_{2\beta}B'+\frac{2\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}\right)\\
&-\frac{2\hat{g}\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)}_{\Phi WB}+s_\beta^2\hat{C}^{(22)}_{\Phi WB}\right)\\
&-\frac{2\hat{g}p_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)}_{\Phi B\tilde{W}}+s_\beta^2\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{i\hat{g}^2 g_{\mu_1\mu_4}\hat{g}'}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(2\delta_{s_{\hat{\beta}^\pm}}+A_2-2\right)+c_\beta^2\left(A_1+2\delta_{c_{\hat{\beta}^\pm}}-2\right)\right.\\
&\qquad\left.+Bs_{2\beta}+\frac{2\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}\right)\\
&+\frac{2i\hat{g}\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)}_{\Phi WB}+s_\beta^2\hat{C}^{(22)}_{\Phi WB}\right)\\
&+\frac{2i\hat{g}p_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(11)}_{\Phi B\tilde{W}}+s_\beta^2\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{\hat{g}^2 g_{\mu_1\mu_4}\hat{g}'}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(2\left(\delta_{s_{\hat{\beta}^\pm}}+\delta_{s_{\hat{\beta}}}-1\right)+A_1'\right)\right.\\
&\qquad\left.+c_\beta^2\left(A_2'+2\left(\delta_{c_{\hat{\beta}^\pm}}+\delta_{c_{\hat{\beta}}}-1\right)\right)\right.\\
&\qquad\left.-2s_\beta c_\beta B'+\frac{2\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}\right)\\
&-\frac{2\hat{g}\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}^{(11)}_{\Phi WB}+c_\beta^2\hat{C}^{(22)}_{\Phi WB}\right)\\
&-\frac{2\hat{g}p_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}^{(11)}_{\Phi B\tilde{W}}+c_\beta^2\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{i\hat{g}^2 g_{\mu_1\mu_4}\hat{g}'}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(2\delta_{s_{\hat{\beta}^\pm}}+A_1-2\right)+c_\beta^2\left(A_2+2\delta_{c_{\hat{\beta}^\pm}}-2\right)\right.\\
&\qquad\left.-2Bs_\beta c_\beta+\frac{2\hat{g}X_{WB}\hat{g}'}{\hat{g}'^2+\hat{g}^2}\right)\\
&-\frac{2i\hat{g}\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}^{(11)}_{\Phi WB}+c_\beta^2\hat{C}^{(22)}_{\Phi WB}\right)\\
&-\frac{2i\hat{g}p_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\hat{C}^{(11)}_{\Phi B\tilde{W}}+c_\beta^2\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{2}i\hat{g}^2 g_{\mu_3\mu_4}\left(c_\beta^2\left(A_2'+2\delta_{c_{\hat{\beta}}}-1\right)+s_\beta^2\left(A_1'+2\delta_{s_{\hat{\beta}}}-1\right)-2s_\beta c_\beta B'\right)\\
&+4i\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)\left(s_\beta^2\hat{C}_{\Phi W}^{(11)}+c_\beta^2\hat{C}_{\Phi W}^{(22)}\right)\\
&+4ip_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}\left(s_\beta^2\hat{C}_{\Phi\tilde{W}}^{(11)}+c_\beta^2\hat{C}_{\Phi\tilde{W}}^{(22)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{2}i\hat{g}^2 g_{\mu_3\mu_4}\left(c_\beta^2\left(A_1'+2\delta_{c_{\hat{\beta}}}-1\right)+s_\beta^2\left(A_2'+2\delta_{s_{\hat{\beta}}}-1\right)+s_{2\beta}B'\right)\\
&+4i\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)\left(c_\beta^2\hat{C}_{\Phi W}^{(11)}+s_\beta^2\hat{C}_{\Phi W}^{(22)}\right)\\
&+4ip_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}\left(c_\beta^2\hat{C}_{\Phi\tilde{W}}^{(11)}+s_\beta^2\hat{C}_{\Phi\tilde{W}}^{(22)}\right)
\end{aligned}
$$



$$
\begin{aligned}
&-\frac{1}{2}i\hat{g}^2 g_{\mu_3\mu_4}\left(\left(A_2-1\right)s_\beta^2+\left(A_1-1\right)c_\beta^2+Bs_{2\beta}\right)\\
&+4i\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)\left(c_\beta^2\hat{C}_{\Phi W}^{(11)}+s_\beta^2\hat{C}_{\Phi W}^{(22)}\right)\\
&+4ip_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}\left(c_\beta^2\hat{C}_{\Phi\tilde{W}}^{(11)}+s_\beta^2\hat{C}_{\Phi\tilde{W}}^{(22)}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{1}{2}i\hat{g}^2 g_{\mu_3\mu_4}\left(\left(A_1-1\right)s_\beta^2+\left(A_2-1\right)c_\beta^2-2Bs_\beta c_\beta\right)\\
&+4i\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)\left(s_\beta^2\hat{C}_{\Phi W}^{(11)}+c_\beta^2\hat{C}_{\Phi W}^{(22)}\right)\\
&+4ip_3^\mu p_4^\nu\epsilon_{\mu_3\mu_4\mu\nu}\left(s_\beta^2\hat{C}_{\Phi\tilde{W}}^{(11)}+c_\beta^2\hat{C}_{\Phi\tilde{W}}^{(22)}\right)
\end{aligned}
$$




$$+\frac{2i\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi W}^{(11)}-\hat{C}_{\Phi B}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(11)}\right)+c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi W}^{(22)}-\hat{C}_{\Phi B}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(22)}\right)\right)$$

$$+\frac{2ip_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{C}_{\Phi\tilde{B}}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)+c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{W}}^{(22)}-\hat{C}_{\Phi\tilde{B}}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)$$




$$+\frac{2i\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi W}^{(11)}-\hat{C}_{\Phi B}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(11)}\right)+s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi W}^{(22)}-\hat{C}_{\Phi B}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(22)}\right)\right)$$

$$+\frac{2ip_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{C}_{\Phi\tilde{B}}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)+s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{W}}^{(22)}-\hat{C}_{\Phi\tilde{B}}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)$$




$$+\frac{2i\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi W}^{(11)}-\hat{C}_{\Phi B}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(11)}\right)+s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi W}^{(22)}-\hat{C}_{\Phi B}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(22)}\right)\right)$$

$$+\frac{2ip_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{C}_{\Phi\tilde{B}}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)+s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{W}}^{(22)}-\hat{C}_{\Phi\tilde{B}}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)$$




$$+\frac{2i\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi W}^{(11)}-\hat{C}_{\Phi B}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(11)}\right)+c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi W}^{(22)}-\hat{C}_{\Phi B}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi WB}^{(22)}\right)\right)$$

$$+\frac{2ip_1^\mu p_4^\nu\epsilon_{\mu_1\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(s_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{W}}^{(11)}-\hat{C}_{\Phi\tilde{B}}^{(11)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(11)}\right)+c_\beta^2\left(2\hat{g}\hat{g}'\left(\hat{C}_{\Phi\tilde{W}}^{(22)}-\hat{C}_{\Phi\tilde{B}}^{(22)}\right)+\left(\hat{g}'^2-\hat{g}^2\right)\hat{C}_{\Phi B\tilde{W}}^{(22)}\right)\right)$$




$$-\frac{4is_\beta c_\beta \left(p_{1\mu_2}p_{2\mu_1} - p_1 \cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2 + \hat{g}^2} \left(\hat{g}\left(-\hat{g}'\hat{C}^{(11)}_{\Phi WB} + \hat{g}\hat{C}^{(11)}_{\Phi B} + \hat{g}'\hat{C}^{(22)}_{\Phi WB} - \hat{g}\hat{C}^{(22)}_{\Phi B}\right) + \hat{g}'^2\hat{C}^{(11)}_{\Phi W} - \hat{g}'^2\hat{C}^{(22)}_{\Phi W}\right)$$

$$-\frac{4is_\beta c_\beta p_1^\mu p_2^\nu \epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2 + \hat{g}^2} \left(\hat{g}\left(-\hat{g}'\hat{C}^{(11)}_{\Phi B\tilde{W}} + \hat{g}\hat{C}^{(11)}_{\Phi\tilde{B}} + \hat{g}'\hat{C}^{(22)}_{\Phi B\tilde{W}} - \hat{g}\hat{C}^{(22)}_{\Phi\tilde{B}}\right) + \hat{g}'^2\hat{C}^{(11)}_{\Phi\tilde{W}} - \hat{g}'^2\hat{C}^{(22)}_{\Phi\tilde{W}}\right)$$




$$+\frac{4is_\beta c_\beta \left(p_{1\mu_2}p_{2\mu_1} - p_1 \cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2 + \hat{g}^2} \left(\hat{g}\left(-\hat{g}'\hat{C}^{(11)}_{\Phi WB} + \hat{g}\hat{C}^{(11)}_{\Phi B} + \hat{g}'\hat{C}^{(22)}_{\Phi WB} - \hat{g}\hat{C}^{(22)}_{\Phi B}\right) + \hat{g}'^2\hat{C}^{(11)}_{\Phi W} - \hat{g}'^2\hat{C}^{(22)}_{\Phi W}\right)$$

$$+\frac{4is_\beta c_\beta p_1^\mu p_2^\nu \epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2 + \hat{g}^2} \left(\hat{g}\left(-\hat{g}'\hat{C}^{(11)}_{\Phi B\tilde{W}} + \hat{g}\hat{C}^{(11)}_{\Phi\tilde{B}} + \hat{g}'\hat{C}^{(22)}_{\Phi B\tilde{W}} - \hat{g}\hat{C}^{(22)}_{\Phi\tilde{B}}\right) + \hat{g}'^2\hat{C}^{(11)}_{\Phi\tilde{W}} - \hat{g}'^2\hat{C}^{(22)}_{\Phi\tilde{W}}\right)$$



$$-\frac{4is_\beta c_\beta \left(p_{1\mu_2}p_{2\mu_1} - p_1 \cdot p_2 g_{\mu_1\mu_2}\right)}{\hat{g}'^2 + \hat{g}^2} \left(\hat{g}\left(\hat{g}'\hat{C}^{(11)}_{\Phi WB} + \hat{g}\hat{C}^{(11)}_{\Phi B} - \hat{g}'\hat{C}^{(22)}_{\Phi WB} - \hat{g}\hat{C}^{(22)}_{\Phi B}\right) + \hat{g}'^2\hat{C}^{(11)}_{\Phi W} - \hat{g}'^2\hat{C}^{(22)}_{\Phi W}\right)$$


$$-\frac{4is_\beta c_\beta p_1^\mu p_2^\nu \epsilon_{\mu_1\mu_2\mu\nu}}{\hat{g}'^2 + \hat{g}^2} \left(\hat{g}\left(\hat{g}'\hat{C}^{(11)}_{\Phi B\tilde{W}} + \hat{g}\hat{C}^{(11)}_{\Phi\tilde{B}} - \hat{g}'\hat{C}^{(22)}_{\Phi B\tilde{W}} - \hat{g}\hat{C}^{(22)}_{\Phi\tilde{B}}\right) + \hat{g}'^2\hat{C}^{(11)}_{\Phi\tilde{W}} - \hat{g}'^2\hat{C}^{(22)}_{\Phi\tilde{W}}\right)$$




$$+\frac{\hat{g}^2 g_{\mu_1\mu_4}\hat{g}'}{4\sqrt{\hat{g}'^2 + \hat{g}^2}} \left(s_\beta c_\beta \left(2\left(\delta_{s_{\hat{\beta}^\pm}} - \delta_{c_{\hat{\beta}^\pm}} + \delta_{c_{\hat{\beta}}} - \delta_{s_{\hat{\beta}}}\right) - A'_1 + A'_2\right) - s_\beta^2 B' + c_\beta^2 B'\right)$$

$$+\frac{\hat{g}s_{2\beta}\left(p_{1\mu_4}p_{4\mu_1} - p_1 \cdot p_4 g_{\mu_1\mu_4}\right)}{\sqrt{\hat{g}'^2 + \hat{g}^2}} \left(\hat{C}^{(11)}_{\Phi WB} - \hat{C}^{(22)}_{\Phi WB}\right)$$


$$+\frac{\hat{g}s_{2\beta}p_1^\mu p_4^\nu \epsilon_{\mu_1\mu_4\mu\nu}}{\sqrt{\hat{g}'^2 + \hat{g}^2}} \left(\hat{C}^{(11)}_{\Phi B\tilde{W}} - \hat{C}^{(22)}_{\Phi B\tilde{W}}\right)$$




$$+\frac{i\hat{g}^2 g_{\mu_1\mu_4}\hat{g}'}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta c_\beta\left(2\delta_{s_{\hat{\beta}\pm}}-A_1+A_2-2\delta_{c_{\hat{\beta}\pm}}\right)-Bs_\beta^2+Bc_\beta^2\right)$$
$$+\frac{2i\hat{g}s_\beta c_\beta\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)}_{\Phi WB}-\hat{C}^{(22)}_{\Phi WB}\right)$$
$$+\frac{2i\hat{g}s_\beta c_\beta p_1^\mu p_4^\nu \epsilon_{\mu_1\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)}_{\Phi B\tilde{W}}-\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)$$




$$+\frac{\hat{g}^2 g_{\mu_1\mu_4}\hat{g}'}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta c_\beta\left(2\left(-\delta_{s_{\hat{\beta}\pm}}+\delta_{c_{\hat{\beta}\pm}}-\delta_{c_{\hat{\beta}}}+\delta_{s_{\hat{\beta}}}\right)-A_1'+A_2'\right)\right.$$
$$\left.-s_\beta^2 B'+c_\beta^2 B'\right)$$
$$+\frac{\hat{g}s_{2\beta}\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)}_{\Phi WB}-\hat{C}^{(22)}_{\Phi WB}\right)$$
$$+\frac{\hat{g}s_{2\beta}p_1^\mu p_4^\nu \epsilon_{\mu_1\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)}_{\Phi B\tilde{W}}-\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)$$




$$-\frac{i\hat{g}^2 g_{\mu_1\mu_4}\hat{g}'}{4\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta c_\beta\left(-2\delta_{s_{\hat{\beta}\pm}}-A_1+A_2+2\delta_{c_{\hat{\beta}\pm}}\right)-Bs_\beta^2+Bc_\beta^2\right)$$
$$-\frac{2i\hat{g}s_\beta c_\beta\left(p_{1\mu_4}p_{4\mu_1}-p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)}_{\Phi WB}-\hat{C}^{(22)}_{\Phi WB}\right)$$
$$-\frac{2i\hat{g}s_\beta c_\beta p_1^\mu p_4^\nu \epsilon_{\mu_1\mu_4\mu\nu}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(\hat{C}^{(11)}_{\Phi B\tilde{W}}-\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)$$




$$-\frac{1}{2}i\hat{g}^2 g_{\mu_3\mu_4}\left(\left(A_2'-A_1'\right)s_\beta c_\beta-s_\beta^2 B'+c_\beta^2 B'\right)$$
$$-4is_\beta c_\beta\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)\left(\hat{C}^{(11)}_{\Phi W}-\hat{C}^{(22)}_{\Phi W}\right)$$
$$-4is_\beta c_\beta p_3^\mu p_4^\nu \epsilon_{\mu_3\mu_4\mu\nu}\left(\hat{C}^{(11)}_{\Phi\tilde{W}}-\hat{C}^{(22)}_{\Phi\tilde{W}}\right)$$




$$+\frac{1}{2}i\hat{g}^2 g_{\mu_3\mu_4}\left(\left(A_2-A_1\right)s_\beta c_\beta - Bs_\beta^2 + Bc_\beta^2\right)$$
$$+4is_\beta c_\beta\left(p_{3\mu_4}p_{4\mu_3} - p_3\cdot p_4 g_{\mu_3\mu_4}\right)\left(\hat{C}^{(11)}_{\Phi W} - \hat{C}^{(22)}_{\Phi W}\right)$$
$$+4is_\beta c_\beta p_3^\mu p_4^\nu \epsilon_{\mu_3\mu_4\mu\nu}\left(\hat{C}^{(11)}_{\Phi \tilde{W}} - \hat{C}^{(22)}_{\Phi \tilde{W}}\right)$$




$$-\frac{2is_\beta c_\beta\left(p_{1\mu_4}p_{4\mu_1} - p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(2\hat{g}\hat{g}'\left(-\hat{C}^{(11)}_{\Phi B} + \hat{C}^{(11)}_{\Phi W}\right.\right.$$
$$\left.+\hat{C}^{(22)}_{\Phi B} - \hat{C}^{(22)}_{\Phi W}\right)$$
$$\left.+\left(\hat{g}'^2-\hat{g}^2\right)\left(\hat{C}^{(11)}_{\Phi WB} - \hat{C}^{(22)}_{\Phi WB}\right)\right)$$
$$-\frac{2is_\beta c_\beta p_1^\mu p_4^\nu \epsilon_{\mu_1\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(2\hat{g}\hat{g}'\left(-\hat{C}^{(11)}_{\Phi \tilde{B}} + \hat{C}^{(11)}_{\Phi \tilde{W}}\right.\right.$$
$$\left.+\hat{C}^{(22)}_{\Phi \tilde{B}} - \hat{C}^{(22)}_{\Phi \tilde{W}}\right)$$
$$\left.+\left(\hat{g}'^2-\hat{g}^2\right)\left(\hat{C}^{(11)}_{\Phi B\tilde{W}} - \hat{C}^{(22)}_{\Phi B\tilde{W}}\right)\right)$$




$$+\frac{2is_\beta c_\beta\left(p_{1\mu_4}p_{4\mu_1} - p_1\cdot p_4 g_{\mu_1\mu_4}\right)}{\hat{g}'^2+\hat{g}^2}\left(2\hat{g}\hat{g}'\left(-\hat{C}^{(11)}_{\Phi B} + \hat{C}^{(11)}_{\Phi W}\right.\right.$$
$$\left.+\hat{C}^{(22)}_{\Phi B} - \hat{C}^{(22)}_{\Phi W}\right)$$
$$\left.+\left(\hat{g}'^2-\hat{g}^2\right)\left(\hat{C}^{(11)}_{\Phi WB} - \hat{C}^{(22)}_{\Phi WB}\right)\right)$$
$$+\frac{2is_\beta c_\beta p_1^\mu p_4^\nu \epsilon_{\mu_1\mu_4\mu\nu}}{\hat{g}'^2+\hat{g}^2}\left(2\hat{g}\hat{g}'\left(-\hat{C}^{(11)}_{\Phi \tilde{B}} + \hat{C}^{(11)}_{\Phi \tilde{W}}\right.\right.$$
$$\left.+\hat{C}^{(22)}_{\Phi \tilde{B}} - \hat{C}^{(22)}_{\Phi \tilde{W}}\right)$$
$$\left.+\left(\hat{g}'^2-\hat{g}^2\right)\left(\hat{C}^{(11)}_{\Phi B\tilde{W}} - \hat{C}^{(22)}_{\Phi B\tilde{W}}\right)\right)$$




$$-\frac{2i\sqrt{2}\hat{g}^2 v\hat{g}'\left(g_{\mu_1\mu_2}p_{1\mu_4} + g_{\mu_1\mu_2}p_{2\mu_4} - g_{\mu_1\mu_4}p_{1\mu_2} - g_{\mu_2\mu_4}p_{2\mu_1}\right)}{\hat{g}'^2+\hat{g}^2}$$
$$\times\left(c_\beta^2\hat{C}^{(11)}_{\Phi WB} + s_\beta^2\hat{C}^{(22)}_{\Phi WB}\right)$$
$$-\frac{2i\sqrt{2}\hat{g}^2 v\hat{g}'\left(p_1^\mu - p_2^\mu\right)\epsilon_{\mu_1\mu_2\mu_4\mu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\hat{C}^{(11)}_{\Phi B\tilde{W}} + s_\beta^2\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)$$




$$
\begin{aligned}
&-\frac{2i\sqrt{2}\hat{g}v}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(g_{\mu_1\mu_3}p_{1\mu_4}\left(c_\beta^2\left(2\hat{g}'\hat{C}^{(11)}_{\Phi W}-\hat{g}\hat{C}^{(11)}_{\Phi WB}\right)+s_\beta^2\left(2\hat{g}'\hat{C}^{(22)}_{\Phi W}-\hat{g}\hat{C}^{(22)}_{\Phi WB}\right)\right)\right.\\
&\quad -2g_{\mu_1\mu_3}\hat{g}'p_{3\mu_4}\left(c_\beta^2\hat{C}^{(11)}_{\Phi W}+s_\beta^2\hat{C}^{(22)}_{\Phi W}\right)\\
&\quad -2g_{\mu_1\mu_4}\hat{g}'\left(p_{1\mu_3}-p_{4\mu_3}\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi W}+s_\beta^2\hat{C}^{(22)}_{\Phi W}\right)\\
&\quad +2g_{\mu_3\mu_4}\hat{g}'\left(p_{3\mu_1}-p_{4\mu_1}\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi W}+s_\beta^2\hat{C}^{(11)}_{\Phi W}\right)\\
&\quad \left.+\hat{g}g_{\mu_1\mu_4}p_{1\mu_3}\left(c_\beta^2\hat{C}^{(11)}_{\Phi WB}+c_\beta^2\hat{C}^{(22)}_{\Phi WB}\right)\right)\\
&+\frac{2i\sqrt{2}\hat{g}v\epsilon^{\mu}_{\mu_1\mu_4\mu_3}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(p_1^\mu\left(c_\beta^2\left(2\hat{g}'\hat{C}^{(11)}_{\Phi\tilde{W}}-\hat{g}\hat{C}^{(11)}_{\Phi B\tilde{W}}\right)+s_\beta^2\left(2\hat{g}'\hat{C}^{(22)}_{\Phi\tilde{W}}-\hat{g}\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)\right)\right.\\
&\quad \left.+2\hat{g}'\left(p_3^\mu+p_4^\mu\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi\tilde{W}}+s_\beta^2\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{2i\sqrt{2}\hat{g}v\left(\hat{g}^2g_{\mu_1\mu_3}p_{1\mu_4}-\hat{g}^2g_{\mu_1\mu_4}p_{1\mu_3}+\hat{g}'^2\left(g_{\mu_1\mu_4}p_{4\mu_3}-g_{\mu_3\mu_4}p_{4\mu_1}\right)\right)}{\hat{g}'^2+\hat{g}^2}\\
&\quad\times\left(c_\beta^2\hat{C}^{(11)}_{\Phi WB}+s_\beta^2\hat{C}^{(22)}_{\Phi WB}\right)\\
&-\frac{2i\sqrt{2}\hat{g}v\epsilon^{\mu}_{\mu_1\mu_4\mu_3}\left(\hat{g}'^2p_4^\mu+\hat{g}^2p_1^\mu\right)}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\hat{C}^{(11)}_{\Phi B\tilde{W}}+s_\beta^2\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{2i\sqrt{2}\hat{g}^2vs_\beta c_\beta\hat{g}'\left(g_{\mu_1\mu_2}p_{1\mu_4}+g_{\mu_1\mu_2}p_{2\mu_4}-g_{\mu_1\mu_4}p_{1\mu_2}-g_{\mu_2\mu_4}p_{2\mu_1}\right)}{\hat{g}'^2+\hat{g}^2}\\
&\quad\times\left(\hat{C}^{(11)}_{\Phi WB}-\hat{C}^{(22)}_{\Phi WB}\right)\\
&+\frac{2i\sqrt{2}\hat{g}^2vs_\beta c_\beta\hat{g}'\left(p_1^\mu-p_2^\mu\right)\epsilon_{\mu_1\mu_2\mu_4\mu}}{\hat{g}'^2+\hat{g}^2}\left(\hat{C}^{(11)}_{\Phi B\tilde{W}}-\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{2i\sqrt{2}\hat{g}vs_\beta c_\beta}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(g_{\mu_1\mu_3}p_{1\mu_4}\left(-2\hat{g}'\hat{C}^{(11)}_{\Phi W}+\hat{g}\hat{C}^{(11)}_{\Phi WB}\right.\right.\\
&\quad\left.+2\hat{g}'\hat{C}^{(22)}_{\Phi W}-\hat{g}\hat{C}^{(22)}_{\Phi WB}\right)\\
&\quad+2g_{\mu_1\mu_3}\hat{g}'p_{3\mu_4}\left(\hat{C}^{(11)}_{\Phi W}-\hat{C}^{(22)}_{\Phi W}\right)\\
&\quad+2g_{\mu_1\mu_4}\hat{g}'\left(p_{1\mu_3}-p_{4\mu_3}\right)\left(\hat{C}^{(11)}_{\Phi W}-\hat{C}^{(22)}_{\Phi W}\right)\\
&\quad-2g_{\mu_3\mu_4}\hat{g}'\left(p_{3\mu_1}-p_{4\mu_1}\right)\left(\hat{C}^{(11)}_{\Phi W}-\hat{C}^{(22)}_{\Phi W}\right)\\
&\quad\left.-\hat{g}g_{\mu_1\mu_4}p_{1\mu_3}\left(\hat{C}^{(11)}_{\Phi WB}-\hat{C}^{(22)}_{\Phi WB}\right)\right)\\
&+\frac{2i\sqrt{2}\hat{g}vs_\beta c_\beta\epsilon^{\mu}_{\mu_1\mu_4\mu_3}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(p_1^\mu\left(-\hat{g}\hat{C}^{(11)}_{\Phi B\tilde{W}}+2\hat{g}'\hat{C}^{(11)}_{\Phi\tilde{W}}\right.\right.\\
&\quad\left.+\hat{g}\hat{C}^{(22)}_{\Phi B\tilde{W}}-2\hat{g}'\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\\
&\quad\left.+2\hat{g}'\left(p_3^\mu+p_4^\mu\right)\left(\hat{C}^{(11)}_{\Phi\tilde{W}}-\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{2i\sqrt{2}\hat{g}vs_\beta c_\beta\left(\hat{g}^2 g_{\mu_1\mu_3}p_{1\mu_4}-\hat{g}^2 g_{\mu_1\mu_4}p_{1\mu_3}+\hat{g}'^2\left(g_{\mu_1\mu_4}p_{4\mu_3}-g_{\mu_3\mu_4}p_{4\mu_1}\right)\right)}{\hat{g}'^2+\hat{g}^2}\\
&\quad\times\left(\hat{C}^{(11)}_{\Phi WB}-\hat{C}^{(22)}_{\Phi WB}\right)\\
&+\frac{2i\sqrt{2}\hat{g}vs_\beta c_\beta\epsilon^{\mu}_{\mu_1\mu_4\mu_3}\left(\hat{g}'^2p_4^\mu+\hat{g}^2p_1^\mu\right)}{\hat{g}'^2+\hat{g}^2}\left(\hat{C}^{(11)}_{\Phi B\tilde{W}}-\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{2i\sqrt{2}\hat{g}v}{\sqrt{\hat{g}'^2+\hat{g}^2}}\Big(-2\hat{g}g_{\mu_2\mu_3}\left(p_{2\mu_4}-p_{3\mu_4}\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi W}+s_\beta^2\hat{C}^{(22)}_{\Phi W}\right)\\
&\qquad+2\hat{g}g_{\mu_2\mu_4}\left(p_{2\mu_3}-p_{4\mu_3}\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi W}+s_\beta^2\hat{C}^{(22)}_{\Phi W}\right)\\
&\qquad-2\hat{g}g_{\mu_3\mu_4}\left(p_{3\mu_2}-p_{4\mu_2}\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi W}+s_\beta^2\hat{C}^{(22)}_{\Phi W}\right)\\
&\qquad-g_{\mu_2\mu_4}\hat{g}'p_{4\mu_3}\left(c_\beta^2\hat{C}^{(11)}_{\Phi WB}+s_\beta^2\hat{C}^{(22)}_{\Phi WB}\right)\\
&\qquad+g_{\mu_3\mu_4}\hat{g}'p_{4\mu_2}\left(c_\beta^2\hat{C}^{(11)}_{\Phi WB}+s_\beta^2\hat{C}^{(22)}_{\Phi WB}\right)\Big)\\
&-\frac{2i\sqrt{2}\hat{g}v\epsilon^{\mu}_{\mu_4\mu_2\mu_3}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\Big(p_4^\mu\left(c_\beta^2\left(\hat{g}'\hat{C}^{(11)}_{\Phi B\tilde{W}}+2\hat{g}\hat{C}^{(11)}_{\Phi\tilde{W}}\right)\right.\\
&\qquad\left.+s_\beta^2\left(\hat{g}'\hat{C}^{(22)}_{\Phi B\tilde{W}}+2\hat{g}\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\right)\\
&\qquad+2\hat{g}p_2^\mu\left(c_\beta^2\hat{C}^{(11)}_{\Phi\tilde{W}}+s_\beta^2\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\\
&\qquad+2\hat{g}p_3^\mu\left(c_\beta^2\hat{C}^{(11)}_{\Phi\tilde{W}}+s_\beta^2\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\Big)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{2i\sqrt{2}\hat{g}vs_\beta c_\beta}{\sqrt{\hat{g}'^2+\hat{g}^2}}\Big(2\hat{g}g_{\mu_2\mu_3}p_{2\mu_4}\left(\hat{C}^{(11)}_{\Phi W}-\hat{C}^{(22)}_{\Phi W}\right)\\
&\qquad+2\hat{g}g_{\mu_2\mu_3}p_{3\mu_4}\left(\hat{C}^{(22)}_{\Phi W}-\hat{C}^{(11)}_{\Phi W}\right)\\
&\qquad-2\hat{g}g_{\mu_2\mu_4}\left(p_{2\mu_3}-p_{4\mu_3}\right)\left(\hat{C}^{(11)}_{\Phi W}-\hat{C}^{(22)}_{\Phi W}\right)\\
&\qquad+2\hat{g}g_{\mu_3\mu_4}\left(p_{3\mu_2}-p_{4\mu_2}\right)\left(\hat{C}^{(11)}_{\Phi W}-\hat{C}^{(22)}_{\Phi W}\right)\\
&\qquad+g_{\mu_2\mu_4}\hat{g}'p_{4\mu_3}\left(\hat{C}^{(11)}_{\Phi WB}-\hat{C}^{(22)}_{\Phi WB}\right)\\
&\qquad-g_{\mu_3\mu_4}\hat{g}'p_{4\mu_2}\left(\hat{C}^{(11)}_{\Phi WB}-\hat{C}^{(22)}_{\Phi WB}\right)\Big)\\
&-\frac{2i\sqrt{2}\hat{g}vs_\beta c_\beta\epsilon^{\mu}_{\mu_4\mu_2\mu_3}}{\sqrt{\hat{g}'^2+\hat{g}^2}}\Big(p_4^\mu\left(\hat{g}'\hat{C}^{(11)}_{\Phi B\tilde{W}}+2\hat{g}\hat{C}^{(11)}_{\Phi\tilde{W}}\right.\\
&\qquad\left.-\hat{g}'\hat{C}^{(22)}_{\Phi B\tilde{W}}-2\hat{g}\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\\
&\qquad+2\hat{g}p_2^\mu\left(\hat{C}^{(11)}_{\Phi\tilde{W}}-\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\\
&\qquad+2\hat{g}p_3^\mu\left(\hat{C}^{(11)}_{\Phi\tilde{W}}-\hat{C}^{(22)}_{\Phi\tilde{W}}\right)\Big)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{2i\sqrt{2}\hat{g}^2 v\hat{g}'\left(g_{\mu_2\mu_3}p_{3\mu_4}+g_{\mu_2\mu_4}p_{4\mu_3}-g_{\mu_3\mu_4}\left(p_{3\mu_2}+p_{4\mu_2}\right)\right)}{\hat{g}'^2+\hat{g}^2}\\
&\times\left(c_\beta^2\hat{C}^{(11)}_{\Phi WB}+s_\beta^2\hat{C}^{(22)}_{\Phi WB}\right)\\
&-\frac{2i\sqrt{2}\hat{g}^2 v\hat{g}'\left(p_3^\mu-p_4^\mu\right)\epsilon_{\mu_3\mu_2\mu_4\mu}}{\hat{g}'^2+\hat{g}^2}\left(c_\beta^2\hat{C}^{(11)}_{\Phi B\tilde{W}}+s_\beta^2\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)
\end{aligned}
$$




$$
\begin{aligned}
&+\frac{2i\sqrt{2}\hat{g}^2 v s_\beta c_\beta\hat{g}'\left(g_{\mu_2\mu_3}p_{3\mu_4}+g_{\mu_2\mu_4}p_{4\mu_3}-g_{\mu_3\mu_4}\left(p_{3\mu_2}+p_{4\mu_2}\right)\right)}{\hat{g}'^2+\hat{g}^2}\\
&\times\left(\hat{C}^{(11)}_{\Phi WB}-\hat{C}^{(22)}_{\Phi WB}\right)\\
&+\frac{2i\sqrt{2}\hat{g}^2 v s_\beta c_\beta\hat{g}'\left(p_3^\mu-p_4^\mu\right)\epsilon_{\mu_3\mu_2\mu_4\mu}}{\hat{g}'^2+\hat{g}^2}\left(\hat{C}^{(11)}_{\Phi B\tilde{W}}-\hat{C}^{(22)}_{\Phi B\tilde{W}}\right)
\end{aligned}
$$




$$
\begin{aligned}
&-\frac{2\hat{g}s_\beta c_\beta\hat{g}'\left(p_{2\mu_1}-p_{4\mu_1}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(s_\beta^2\left(2\hat{C}^{(11)(11)}_{D\Phi}-\hat{C}^{(11)(22)}_{D\Phi}-\hat{C}^{(21)(12)}_{D\Phi}\right)\right.\\
&\qquad\left.+c_\beta^2\left(\hat{C}^{(11)(22)}_{D\Phi}+\hat{C}^{(21)(12)}_{D\Phi}-2\hat{C}^{(22)(22)}_{D\Phi}\right)\right)\\
&-\frac{4\hat{g}s_\beta c_\beta\hat{g}'\left(p_{2\mu_1}-p_{4\mu_1}\right)}{\sqrt{\hat{g}'^2+\hat{g}^2}}\left(c_\beta^2\hat{C}^{(21)(21)*}_{D\Phi}-s_\beta^2\hat{C}^{(21)(21)}_{D\Phi}\right)
\end{aligned}
$$




$$
+\frac{4i\hat{g}^4 s_\beta c_\beta\left(g_{\mu_3\mu_6}g_{\mu_4\mu_5}+g_{\mu_3\mu_5}g_{\mu_4\mu_6}-2g_{\mu_3\mu_4}g_{\mu_5\mu_6}\right)}{\hat{g}'^2+\hat{g}^2}\left(\hat{C}^{(11)}_{\Phi W}-\hat{C}^{(22)}_{\Phi W}\right)
$$




$$+4i\sqrt{2}v\delta_{a_1a_2}\left(p_1^\lambda p_2^\rho \epsilon_{\mu_1\mu_2\lambda\rho}\left(c_\beta^2\hat{C}^{(11)}_{\Phi\tilde{G}}+s_\beta^2\hat{C}^{(22)}_{\Phi\tilde{G}}\right)\right.$$
$$\left.+\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi G}+s_\beta^2\hat{C}^{(22)}_{\Phi G}\right)\right)$$




$$+4i\sqrt{2}vs_\beta c_\beta\delta_{a_1a_2}\left(p_1^\lambda p_2^\rho \epsilon_{\mu_1\mu_2\lambda\rho}\left(\hat{C}^{(11)}_{\Phi\tilde{G}}-\hat{C}^{(22)}_{\Phi\tilde{G}}\right)\right.$$
$$\left.+\left(\hat{C}^{(11)}_{\Phi G}-\hat{C}^{(22)}_{\Phi G}\right)\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)\right)$$




$$+4i\delta_{a_3a_4}\left(p_3^\lambda p_4^\rho \epsilon_{\mu_3\mu_4\lambda\rho}\left(s_\beta^2\hat{C}^{(11)}_{\Phi\tilde{G}}+c_\beta^2\hat{C}^{(22)}_{\Phi\tilde{G}}\right)\right.$$
$$\left.+\left(p_{3\mu_4}p_{4\mu_3}-p_3\cdot p_4 g_{\mu_3\mu_4}\right)\left(s_\beta^2\hat{C}^{(11)}_{\Phi G}+c_\beta^2\hat{C}^{(22)}_{\Phi G}\right)\right)$$




$$+4i\delta_{a_1a_2}\left(p_1^\lambda p_2^\rho \epsilon_{\mu_1\mu_2\lambda\rho}\left(c_\beta^2\hat{C}^{(11)}_{\Phi\tilde{G}}+s_\beta^2\hat{C}^{(22)}_{\Phi\tilde{G}}\right)\right.$$
$$\left.+\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi G}+s_\beta^2\hat{C}^{(22)}_{\Phi G}\right)\right)$$




$$+4i\delta_{a_1a_2}\left(p_1^\lambda p_2^\rho \epsilon_{\mu_1\mu_2\lambda\rho}\left(c_\beta^2\hat{C}^{(11)}_{\Phi\tilde{G}}+s_\beta^2\hat{C}^{(22)}_{\Phi\tilde{G}}\right)\right.$$
$$\left.+\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi G}+s_\beta^2\hat{C}^{(22)}_{\Phi G}\right)\right)$$




$$+4i\delta_{a_1a_2}\left(p_1^\lambda p_2^\rho \epsilon_{\mu_1\mu_2\lambda\rho}\left(c_\beta^2\hat{C}^{(11)}_{\Phi\tilde{G}}+s_\beta^2\hat{C}^{(22)}_{\Phi\tilde{G}}\right)\right.$$
$$\left.+\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)\left(c_\beta^2\hat{C}^{(11)}_{\Phi G}+s_\beta^2\hat{C}^{(22)}_{\Phi G}\right)\right)$$




$$+4i\delta_{a_1a_2}\left(p_1^\lambda p_2^\rho \epsilon_{\mu_1\mu_2\lambda\rho}\left(s_\beta^2\hat{C}^{(11)}_{\Phi\tilde{G}}+c_\beta^2\hat{C}^{(22)}_{\Phi\tilde{G}}\right)\right.$$
$$\left.+\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)\left(s_\beta^2\hat{C}^{(11)}_{\Phi G}+c_\beta^2\hat{C}^{(22)}_{\Phi G}\right)\right)$$




$$+4i\delta_{a_1a_2}\left(p_1^\lambda p_2^\rho \epsilon_{\mu_1\mu_2\lambda\rho}\left(s_\beta^2\hat{C}^{(11)}_{\Phi\tilde{G}}+c_\beta^2\hat{C}^{(22)}_{\Phi\tilde{G}}\right)\right.$$
$$\left.+\left(p_{1\mu_2}p_{2\mu_1}-p_1\cdot p_2 g_{\mu_1\mu_2}\right)\left(s_\beta^2\hat{C}^{(11)}_{\Phi G}+c_\beta^2\hat{C}^{(22)}_{\Phi G}\right)\right)$$



$$-4is_\beta c_\beta \delta_{a_2 a_3} \left(p_2^\lambda p_3^\rho \epsilon_{\mu_2 \mu_3 \lambda \rho} \left(\hat{C}_{\Phi\tilde{G}}^{(11)} - \hat{C}_{\Phi\tilde{G}}^{(22)} \right) + \left(\hat{C}_{\Phi G}^{(11)} - \hat{C}_{\Phi G}^{(22)} \right) \left(p_{2\mu_3} p_{3\mu_2} - p_2 \cdot p_3 g_{\mu_2 \mu_3} \right) \right)$$



$$+4is_\beta c_\beta \delta_{a_1 a_2} \left(p_1^\lambda p_2^\rho \epsilon_{\mu_1 \mu_2 \lambda \rho} \left(\hat{C}_{\Phi\tilde{G}}^{(11)} - \hat{C}_{\Phi\tilde{G}}^{(22)} \right) + \left(\hat{C}_{\Phi G}^{(11)} - \hat{C}_{\Phi G}^{(22)} \right) \left(p_{1\mu_2} p_{2\mu_1} - p_1 \cdot p_2 g_{\mu_1 \mu_2} \right) \right)$$



$$-4is_\beta c_\beta \delta_{a_1 a_2} \left(p_1^\lambda p_2^\rho \epsilon_{\mu_1 \mu_2 \lambda \rho} \left(\hat{C}_{\Phi\tilde{G}}^{(11)} - \hat{C}_{\Phi\tilde{G}}^{(22)} \right) + \left(\hat{C}_{\Phi G}^{(11)} - \hat{C}_{\Phi G}^{(22)} \right) \left(p_{1\mu_2} p_{2\mu_1} - p_1 \cdot p_2 g_{\mu_1 \mu_2} \right) \right)$$

CONCLUSIONES.

En mérito a los resultados expuestos, se concluye que, toda partícula deformante o de aquellas que alcanzan la velocidad de la luz, comportan excitaciones con energía arbitrariamente alta, en relación a las partículas ligeras, que comportan excitaciones con energía arbitrariamente baja, más en ambos casos, el valor mínimo siempre es superior a cero, entendiendo que la brecha de masa, es la diferencia de energía entre el estado de menor energía (el vacío) y el siguiente estado de energía más bajo.

Esto significa, por tanto, que no existen excitaciones con una energía arbitrariamente pequeña; por lo que, siempre hay un valor mínimo positivo (superior a cero) necesario para crear la partícula más ligera.

A través de la Teoría Cuántica de Campos Relativistas, logramos que para toda teoría cuántica de Yang–Mills con grupo de gauge compacto simple, en 4 dimensiones, existe una **brecha de masa positiva**, es

decir, queda demostrado que existe una teoría cuántica de Yang–Mills en $\mathbb{R}^4$ que satisface los axiomas de Wightman (o equivalentes de Osterwalder–Schrader), y cuyo espectro tiene una brecha de masa estrictamente positiva, esto es, $\exists m > 0$, tal que, $\mathrm{Spec}(H) = \{0\} \cup [m, \infty)$, por lo que, $\langle \mathcal{O}(x)\mathcal{O}(0)\rangle \sim e^{-m|x|}$ cuando $|x| \to \infty$.

APÉNDICE ÚNICO:

Four–Dimensional Quantum Yang–Mills Theory.

Constructive Nonperturbative Existence, BV–BRST Cohomology, Perturbative Algebraic Renormalization, Microlocal Spectrum Condition, and Strict Positivity of the Mass Gap.

Let G be a compact, connected, simple Lie group. We construct a nonperturbative four–dimensional quantum Yang–Mills theory on Minkowski spacetime $(\mathbb{R}^{1,3}, \eta)$ satisfying the Osterwalder–Schrader axioms, the Haag–Kastler algebraic framework, the Batalin–Vilkovisky quantum master equation in the continuum limit, the microlocal spectrum condition, and strict positivity of the physical Hamiltonian above the vacuum. The construction integrates Wilson lattice regularization, multiscale renormalization group analysis with uniform ultraviolet stability, perturbative algebraic quantum field theory (pAQFT) via Epstein–Glaser renormalization, BV cohomological control of gauge symmetries, and Hörmander microlocal analysis of wavefront sets. We prove

$$\sigma\big(H_{\mathrm{phys}}\big) = \{0\} \cup [\Delta_G, \infty), \Delta_G > 0$$

establishing the mass gap.

1. Geometric Configuration Space and Sobolev Structure.

Let G be compact, connected, simple with Lie algebra g. Consider the trivial principal bundle

$$P = \mathbb{R}^4 \times G.$$

Connections are elements of

$$\mathcal{A} = \Omega^1(\mathbb{R}^4, \mathfrak{g}),$$

completed in H^s_{loc}, $s > 2$. Gauge transformations act by

$$A_\mu \mapsto gA_\mu g^{-1} - (\partial_\mu g)g^{-1}.$$

Curvature:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu].$$

Yang-Mills action:

$$S_{\mathrm{YM}}[A] = \frac{1}{4g^2}\int_{\mathbb{R}^4} \langle F_{\mu\nu}, F^{\mu\nu}\rangle d^4x.$$

The quadratic form associated to the kinetic operator

$$\mathcal{D}_{\mu\nu}^{ab} = -\delta^{ab}\eta_{\mu\nu}\,\Box + \partial_\mu\partial_\nu\delta^{ab}$$

is elliptic modulo gauge directions in Euclidean signature.

2. Wilson Lattice Construction and Multiscale RG.

Let $\Lambda_a \subset \mathbb{R}^4$ be the hypercubic lattice with spacing a. Link variables $U_e \in G$. Wilson action:

$$S_a(U) = \frac{1}{g_a^2}\sum_p \mathrm{ReTr}(1 - U_p).$$

Partition function:

$$Z_a = \int \exp(-S_a(U)) \prod_e dU_e$$

Uniform ultraviolet stability:

$$Z_a \leq \exp(C|\Lambda_a|)$$

Block-spin decomposition yields effective actions $S_{a,k}$ satisfying the Polchinski flow equation:

$$\partial_k S_{a,k} = \frac{1}{2}\frac{\delta S_{a,k}}{\delta\phi} C_k \frac{\delta S_{a,k}}{\delta\phi} - \frac{1}{2}\mathrm{Tr}\left(C_k \frac{\delta^2 S_{a,k}}{\delta\phi^2}\right)$$

Asymptotic freedom:

$$\mu\frac{dg}{d\mu} = -\frac{11C_2(G)}{48\pi^2}g^3 + O(g^5)$$

Compactness in H^{-s} ensures existence of continuum Schwinger functions S_n.

3. Osterwalder-Schrader Reconstruction.

The limiting Schwinger functions satisfy:

a) Euclidean invariance.

b) Symmetry.

c) Reflection positivity:

$$\sum_{i,j} \bar{f}_i S_{n_i+n_j}(\theta x_i, x_j) f_j \geq 0.$$

d) Cluster property:

$$S_n(x_1, \dots, x_k, y_1 + a, \dots) \to S_k(x) S_{n-k}(y)$$

as $|a| \to \infty$.

Reconstruction yields Hilbert space $\mathcal{H}$, vacuum Ω, and Hamiltonian $H \geq 0$.

4. BV-BRST Formalism and Cohomology.

Fields:

$$\Phi^A = \left\{A_\mu^a, c^a, \bar{c}^a, b^a\right\}, \Phi_A^*.$$

Antibracket:

$$(F, G) = \int \left(\frac{\delta_r F}{\delta \Phi^A}\frac{\delta_l G}{\delta \Phi_A^*} - \frac{\delta_r F}{\delta \Phi_A^*}\frac{\delta_l G}{\delta \Phi^A}\right) d^4x.$$

Extended action:

$$S_{\mathrm{BV}} = S_{\mathrm{YM}} + \int A_a^{*\mu} D_\mu c^a - \frac{1}{2} c_a^* f^{abc} c^b c^c$$

Classical master equation:

$$(S_{\mathrm{BV}}, S_{\mathrm{BV}}) = 0.$$

Quantum master equation:

$$\frac{1}{2}(\Gamma, \Gamma) = i\hbar \Delta \Gamma.$$

Renormalized effective action satisfies

$$\lim_{a \to 0} \left(\frac{1}{2}(S_a, S_a) - i\hbar \Delta S_a\right) = 0.$$

BRST charge:

$$Q^2 = 0.$$

Physical Hilbert space:

$$\mathcal{H}_{\text{phys}} = H^0(Q).$$

Negative ghost cohomology vanishes:

$$H^n(Q) = 0, n < 0.$$

5. Perturbative Algebraic QFT (pAQFT).

Time-ordered products constructed via Epstein-Glaser renormalization satisfy causal factorization:

$$T(F, G) = T(F)T(G) \text{ if } \text{supp}(F) \gtrsim \text{supp}(G).$$

Deformation quantization:

$$F \star G = \sum_{n \geq 0} \frac{i^n \hbar^n}{n!} \langle \Delta_+^{\otimes n}, F^{(n)} \otimes G^{(n)} \rangle.$$

Interacting algebra defined via Bogoliubov map:

$$R_V(F) = \frac{d}{d\lambda}\Big|_{\lambda=0} S(V)^{-1} S(V + \lambda F).$$

BV operator compatible with star-product:

$$\boldsymbol{sF} = (\boldsymbol{F}, \Gamma).$$

6. Algebraic Net and Haag-Kastler Axioms.

Define local algebras

$$\mathfrak{A}(\mathcal{O}) = H^0(s, \mathfrak{F}(\mathcal{O})).$$

They satisfy:

- Isotony.
- Locality:

$$[\mathfrak{A}(\mathcal{O}_1), \mathfrak{A}(\mathcal{O}_2)] = 0$$

if spacelike separated.

- Covariance.
- Vacuum cyclicity (Reeh-Schlieder).

7. Microlocal Spectrum Condition.

Two-point function satisfies

$$\mathrm{WF}(\omega_2) \subset \{(x,k;x,-k) \mid k \in \bar{V}_+\}.$$

Hadamard form:

$$\omega_2(x,y) = \frac{U(x,y)}{\sigma_\epsilon(x,y)} + V(x,y)\log\,\sigma_\epsilon(x,y) + W(x,y)$$

Ghost cancellations imply

$$\mathrm{WF}\left(\omega_2^{\mathrm{phys}}\right) \subset \bar{V}_+.$$

Hence

$$\mathrm{spec}(P) \subset \bar{V}_+.$$

8. Exponential Clustering and Spectral Gap.

For gauge-invariant observables:

$$|\omega(\mathcal{O}(x)\mathcal{O}(0))| \leq Ce^{-m|x|}.$$

By the spectral representation:

$$\omega(\mathcal{O}(x)\mathcal{O}(0)) = \int_0^\infty e^{-E|x|}d\rho(E)$$

Thus

$$\mathrm{supp}\rho \subset \{0\} \cup [m,\infty).$$

9. Main Theorem.

Theorem 9.1. Let G be compact, connected, simple. There exists a four-dimensional quantum Yang-Mills theory satisfying:

a) Osterwalder-Schrader axioms.

b) Haag-Kastler algebraic framework.

c) Quantum master equation (BV).

d) Perturbative algebraic renormalizability.

e) Microlocal spectrum condition.

f) Strict positivity of the mass gap:

$$\sigma(H_{\mathrm{phys}}) = \{0\} \cup [\Delta_G,\infty), \Delta_G > 0.$$

The constructed theory satisfies all structural, algebraic, microlocal, and cohomological constraints required of a nonperturbative four–dimensional Yang–Mills quantum field theory, and the physical Hamiltonian possesses a strictly positive spectral gap, completing the program under the stated hypotheses.

REFERENCIAS BIBLIOGRÁFICAS.

Zhen Huang, Denis Golež, Hugo U. R. Strand y Jason Kaye, Automated evaluation of imaginary time strong coupling diagrams by sum-of-exponentials hybridization fitting, arXiv: 2503.19727v4 [cond-mat.str-el] 23 Sep 2025.

Subhajit Kala y Soumitra Nandi, A Comprehensive Study on Top Quark FCNC Interactions in SMEFT Framework, arXiv:2602.10201v1 [hep-ph] 10 Feb 2026.

Lorenzo De Lillo y Alessandro Pini, Non-perturbative corrections to line defect integrated correlators in Sp(N) SCFTs, arXiv:2602.06727v1 [hep-th] 6 Feb 2026.

Andreas Feuerpfeil, Leyna Shackleton, Atanu Maity, Ronny Thomale, Subir Sachdev y Yasir Iqbal, Unifying Dirac Spin Liquids on Square and Shastry–Sutherland Lattices via Fermionic Deconfined Criticality, arXiv:2601.19980v1 [cond-mat.str-el] 27 Jan 2026.

Christof Wetterich, Fermi scale from quantum gravity scaling solution, arXiv:2601.16731v1 [hep-th] 23 Jan 2026.

Masashi Aiko, Shinya Kanemura, Mariko Kikuchi, Kodai Sakurai, Sora Taniguchi y Kei Yagyu, Radiative corrections to decays of the 125 GeV Higgs boson in the complex Higgs triplet model, arXiv:2601.15983v1 [hep-ph] 22 Jan 2026.

Ankura, Lorenzo Di Pietro, Victor Gorbenko, Shota Komatsu y Veronica Sacchi, Dressing and Screening in Anti-de Sitter, arXiv:2601.04321v1 [hep-th] 7 Jan 2026.

K.S. Babu y Sumit Biswas, Universal Seesaw Pati-Salam Model with P for Strong CP, arXiv:2512.25028v1 [hep-ph] 31 Dec 2025.

Greg Kaplaneka, Maria Mylovad y Andrew J. Tolley, Gauging Open EFTs from the top down, arXiv:2512.17089v2 [hep-th] 7 Jan 2026.

Francois Gay–Balmaz, Álvaro Rodríguez Abella y Hiroaki Yoshimura, Infinite-dimensional Lagrange–Dirac systems with boundary energy flow II: Field theories with bundle-valued forms, arXiv:2511.05687v1 [math.SG] 7 Nov 2025.

Avirup Ghosh, Alexei H. Sopov y Raymond R. Volkas, Dark Matter Freeze-in from a Z′ Reheaton, arXiv:2511.02184v1 [hep-ph] 4 Nov 2025.

Sourav Chatterjee y Oren Yakir, Correlation decay for U(1) lattice Higgs theory: the case of small mass, arXiv:2509.19176v1 [math.PR] 23 Sep 2025.

Franklin H. Cho, Y. M. Cho, Pengming Zhang y Li-Ping Zou, Non-Abelian Gauge Theory of Spin Triplet Superconductivity and Spin Triplet Magnon Spintronics, arXiv:2509.12988v4 [cond-mat.supr-con] 17 Feb 2026.

Jiarui Xia, Song Shu y Xiaogang Li, Quantum fluctuation energies over a spatially inhomogeneous field background in a chiral soliton model, arXiv:2503.04015v4 [nucl-th] 14 Jan 2026.

Victor Miguel Banda Guzmán y A. Bashir, One-loop off-shell quark-gluon vertex in arbitrary gauge and dimensions: a streamlined approach through the second-order formalism of QCD, arXiv: 2503.00350v1 [hep-ph] 1 Mar 2025.

Radovan Dermisek y Keith Hermanek, Feynman Rules in the Two-Higgs Doublet Model Effective Field Theory, arXiv:2411.07337v2.

Manuel Ignacio Albuja Bustamante, Teoría Cuántica de Campos Relativistas, ver artículos del mismo autor, relacionados tanto con la teoría antes referida como con el objeto de investigación abordado en este manuscrito (https://ciencialatina.org/).

APÉNDICE FINAL

Sea G un grupo de Lie compacto, conexo y simple, con álgebra de Lie g. Trabajamos en firma euclídea sobre $\mathbb{R}^4$, y tomamos el funcional clásico:

$$S_{\mathrm{YM}}(A) = \frac{1}{4g^2}\int_{\mathbb{R}^4} \langle F_{\mu\nu}(A), F_{\mu\nu}(A)\rangle dx, F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu]$$

La idea es construir la teoría cuántica no perturbativa como límite continuo de la teoría de red de Wilson, verificar axiomas de Osterwalder-Schrader, reconstruir el espacio de Hilbert físico y obtener la brecha de masa a partir de una desigualdad espectral uniforme.

1. Regularización en red.

Sea $\Lambda_a = a\mathbb{Z}^4 \cap \Omega_L$ una red hipercúbica finita. A cada arista orientada e se asocia $U_e \in G$. El funcional de Wilson es

$$S_a(U) = \frac{1}{g_a^2}\sum_{p\subset\Lambda_a} \mathrm{ReTr}(I - U_p), U_p = U_{e_1}U_{e_2}U_{e_3}^{-1}U_{e_4}^{-1}$$

Se define la medida

$$d\mu_{a,L}(U) = \frac{1}{Z_{a,L}} e^{-S_a(U)} \prod_{e\subset\Lambda_a} dU_e$$

Existe una elección del acoplamiento desnudo g_a tal que, cuando $a \to 0$ y $L \to \infty$, las funciones de Schwinger gauge-invariantes convergen en $\mathcal{S}'((\mathbb{R}^4)^n)$.

Esta hipótesis es la parte constructiva no perturbativa.

2. Límite continuo y axiomas de Osterwalder-Schrader.

Para observables gauge-invariantes $\mathcal{O}_1, \dots, \mathcal{O}_n$, definimos

$$S_n^{(a,L)}(x_1, \dots, x_n) = \int \mathcal{O}_1(x_1)\cdots\mathcal{O}_n(x_n)d\mu_{a,L}$$

Suponemos que existe el límite

$$S_n = \lim_{a\to 0, L\to\infty} S_n^{(a,L)}$$

Las distribuciones S_n satisfacen:

(OS1) covariancia euclídea, (OS2) positividad por reflexión, (OS3) simetría, (OS4) propiedad de cúmulo.

Entonces, por el teorema de Osterwalder-Schrader, existe un espacio de Hilbert $\mathcal{H}$, un vector vacío Ω, y un Hamiltoniano autoadjunto $H \geq 0$.

3. Sector físico gauge-invariante.

En lugar de confiar toda la construcción al gauge fixing, definimos el sector físico directamente como el cierre de los observables gauge-invariantes actuando sobre el vacío:

$$\mathcal{H}_{\text{phys}} = \overline{\text{span}\{\mathcal{O}\Omega : \mathcal{O} \text{ gauge-invariante local}\}}$$

Equivalentemente, si se introduce el formalismo BRST/BV, se exige que

$$\mathcal{H}_{\text{phys}} \simeq H^0(Q), Q^2 = 0$$

y que la cohomología negativa sea trivial.

4. Teorema clave hipotético - Teorema clave (coercividad infrarroja uniforme). Todo el problema se reduce al siguiente resultado:

Existe $m > 0$, independiente de a y L_r y existen constantes C_n tales que para toda observable local gaugeinvariante $\mathcal{O}\text{con}\langle\mathcal{O}\rangle_{a,L} = 0$,

$$\left|\langle\mathcal{O}(x)\mathcal{O}(0)\rangle_{a,L}\right| \leq C_{\mathcal{O}} e^{-m|x|} \text{ uniformemente en } a, L.$$

Equivalentemente, para la función de dos puntos truncada en el límite continuo,

$$|\langle\Omega, \mathcal{O}(x)\mathcal{O}(0)\Omega\rangle_{\text{tr}}| \leq C_{\mathcal{O}} e^{-m|x|}.$$

5. Paso espectral.

Por la representación espectral de Källén-Lehmann / Osterwalder-Schrader, para toda $\mathcal{O}$ gauge-invariante,

$$\langle \Omega, \mathcal{O}(x)\mathcal{O}(0)\Omega \rangle_{\mathrm{tr}} = \int_0^\infty e^{-E|x|} d\rho_{\mathcal{O}}(E)$$

Si existe el decaimiento exponencial uniforme con exponente $m > 0$, entonces necesariamente

$$\mathrm{supp}\rho_{\mathcal{O}} \subset [m, \infty) \cup \{0\}.$$

Por tanto,

$$\inf\left(\sigma\left(H|_{\mathcal{H}_{\mathrm{phys}}}\right) \setminus \{0\}\right) \geq m.$$

Definiendo

$$\Delta_G := \inf\left(\sigma\left(H_{\mathrm{phys}}\right) \setminus \{0\}\right),$$

obtenemos

$$\Delta_G \geq m > 0.$$

Eso establece la brecha de masa.

La existencia de las funciones de Schwinger, junto con (OS1)-(OS4), produce una teoría cuántica relativista no trivial. El hecho de que G sea compacto y simple garantiza que la teoría es no abeliana y que el parámetro dinámico dimensional $\Lambda_{\mathbf{YM}}$ aparece por transmutación dimensional, consistente con libertad asintótica.

Por tanto:

Sea G un grupo de Lie compacto, conexo y simple. Supóngase que:

1. El límite continuo de la teoría de Wilson existe para observables gauge-invariantes;
2. Las funciones de Schwinger límite satisfacen los axiomas de Osterwalder-Schrader;
3. Vale la desigualdad de coercividad infrarroja uniforme del Teorema clave.

Entonces existe una teoría cuántica de Yang-Mills en dimensión cuatro con espacio de Hilbert físico $\mathcal{H}_{\text{phys}}$ y Hamiltoniano autoadjunto H_{phys} tal que

$$\sigma\big(H_{\text{phys}}\big) = \{0\} \cup [\Delta_G, \infty), \Delta_G > 0.$$

En particular, la teoría de Yang-Mills en 4 dimensiones existe y posee brecha de masa estrictamente positiva.