Ciencia Latina Revista Científica Multidisciplinar, Ciudad de México, México.
ISSN 2707-2207 / ISSN 2707-2215 (en línea), enero-febrero 2026,
Volumen 10, Número 1.

https://doi.org/10.37811/cl_rcm.v10i1

TEORÍA CUÁNTICA DE CAMPOS RELATIVISTAS: UNA ALTERNATIVA DE SOLUCIÓN AL PROBLEMA DEL MILENIO DE YANG – MILLS. UN INTENTO POR UNIFICAR LA RELATIVIDAD GENERAL Y LA MECÁNICA CUÁNTICA. VOLUMEN IV.

RELATIVISTIC QUANTUM FIELD THEORY: AN ALTERNATIVE SOLUTION TO THE YANG–MILLS MILLENNIUM PROBLEM. AN ATTEMPT TO UNIFY GENERAL RELATIVITY AND QUANTUM MECHANICS. VOLUME IV.

Manuel Ignacio Albuja Bustamante
Investigador Independiente

DOI: https://doi.org/10.37811/cl_rcm.v10i1.22959

TEORÍA CUÁNTICA DE CAMPOS RELATIVISTAS: UNA ALTERNATIVA DE SOLUCIÓN AL PROBLEMA DEL MILENIO DE YANG – MILLS. UN INTENTO POR UNIFICAR LA RELATIVIDAD GENERAL Y LA MECÁNICA CUÁNTICA. VOLUMEN IV.

Manuel Ignacio Albuja Bustamante[1]
ignaciomanuelalbujabustamante@gmail.com
https://orcid.org/0009-0005-0115-767X
Investigador Independiente

RESUMEN.
En este trabajo, compuesto por diez volúmenes, abordaremos aspectos esenciales de la Teoría Cuántica de Campos Relativistas (TCCR), con propósitos de optimización de los cálculos expuestos en trabajos anteriores pero sobre todo, posicionar la referida teoría, como una alternativa de solución al problema del milenio de Yang – Mills y la brecha de masa. La idea esencial es la misma, todo espacio – tiempo cuántico, es decir, todo campo cuántico, es curvo y esa deformación ocurre por la gravedad y supergravedad cuánticas, según sea el caso, que provocan las partículas oscuras o estrella, al momento de interactuar con un campo gravitónico o supergravitónico, según corresponda, o en relación a la criticidad de su centro de masa y/o energía, lo que afecta su espín, velocidad y momento angular y por ende, sus trayectorias orbitales. Por tanto, la TCCR, no es un intento por cuantizar la gravedad, sino por introducir la gravedad, como principio de mínima acción de un sistema cuántico y de sus estados fundamentales.
Las métricas siguen siendo las mismas, es decir, que para un campo cuántico curvo o geométricamente deformado, la densidad lagrangiana/hamiltoniana equivale a: $\mathcal{LH}_{curvature} = \langle \int \hat{e}^{i\hbar t} \sqrt{\bar{\bar{g}}^{\mu\nu}} \otimes \overleftrightarrow{m\bar{\psi}\bar{\psi}} - \partial^2 \triangle' \rangle' \langle \otimes_{\Re}^{\mathfrak{G}} | d^4x/\partial\mathcal{R} \rangle' \int \left\| \overline{\frac{\partial \phi'_{\sigma\rho}}{\partial \phi^{\dagger}_{\sigma\rho}}} \right\| -$

$\left\langle \frac{\partial \phi^{\star}_{\sigma\rho}}{\partial^{\mathfrak{H}}_{\mu\nu}} \middle| \partial \uparrow / \partial t \setminus \partial \downarrow / \partial t \partial^2 \square \middle| \begin{matrix} \blacksquare^{\circlearrowleft} \\ \blacksquare_{\circlearrowright} \end{matrix} \partial^2 \varphi / \partial \psi \blacksquare \right\rangle \wedge_{\nu}^{\mu} \sum_{\substack{0 \leq \iota \leq m \\ 0<J<n}} P(\iota, J) \prod_{k=1}^{n} A_k \cup_{n=1}^{m} (X_n \cap Y_n) \cup_{n=1}^{m} (X_n \cap Y_n) \circledcirc \Lambda_{\nu}^{\mu} \odot \Gamma_{\nu}^{\mu},$

respecto de una partícula pesada ρ, sea oscura o blanca (partícula estrella), según corresponda, a propósito de la criticidad de su masa y/o energía $\langle 0 | \sum_{\delta} \partial \mathfrak{m} / \partial \mathfrak{e} \rangle$ o de su interacción con un gravitón o un gravitino, según corresponda, en coordenadas $\langle \rho^{\mu} \rho^{\nu} \rho^{\sigma} \rho^{\varrho} \rangle$, esto último, lo que ocurre por permeabilización del campo gravitónico o supergravitónico en $\blacksquare = \int \langle \partial \mathfrak{G} / \partial \subseteq \mathfrak{G} \rangle$, lo que corresponde al espacio – cuántico deformado en $\mathfrak{C}_{\mathfrak{H}\Re} = \langle \sum_{\blacksquare}^{\sigma\rho} \mathcal{R}_{\nu}^{\mu\dagger} | \otimes \mathcal{H}_{\mu}^{\nu\star} \rangle$ lo que en dimensiones $\mathbb{R}^{\eta}$, representa, gravedad o supergravedad cuánticas por curvatura o supercurvatura del espacio - tiempo cuántico multidimensional.

Palabras Clave: Supergravedad cuántica, gravedad cuántica, partícula oscura, partícula estrella, hiperpartículas, suprapartículas, teoría cuántica de campos relativistas, problema del milenio de Yang – Mills y la brecha de masa, partículas ligeras, curvatura, supercurvatura, multidimensiones, agujeros cuánticos.

[1] Autor principal
Correspondencia: ignaciomanuelalbujabustamante@gmail.com

RELATIVISTIC QUANTUM FIELD THEORY: AN ALTERNATIVE SOLUTION TO THE YANG–MILLS MILLENNIUM PROBLEM. AN ATTEMPT TO UNIFY GENERAL RELATIVITY AND QUANTUM MECHANICS. VOLUME IV.

ABSTRACT.

In this work, composed of ten volumes, we will address essential aspects of the Quantum Theory of Relativistic Fields (TCCR), with the purpose of optimizing the calculations exposed in previous works but above all, positioning the aforementioned theory as an alternative solution to the Yang-Mills millennium problem and the mass gap. The essential idea is the same, all quantum space-time, that is, every quantum field, is curved and that deformation occurs due to quantum gravity and supergravity, as the case may be, caused by dark particles or stars, when interacting with a gravitonic or supergravitonic field, as appropriate, or in relation to the criticality of its center of mass and/or energy which affects their spin, velocity and angular momentum and therefore, their orbital trajectories. Therefore, the TCCR is not an attempt to quantize gravity, but to introduce gravity, as the principle of least action of a quantum system and its fundamental states.

The metrics remain the same, i.e., for a curved or geometrically warped quantum field, the Lagrangian/Hamiltonian density is equal to: $\mathcal{LH}_{curvature} = \langle \int \hat{e}^{i\hbar t}\sqrt{\bar{g}^{\mu\nu}} \otimes \overleftrightarrow{m\psi}\bar{\psi} - \partial^2 \triangle' \rangle' \langle \otimes_{\Re}^{\mathfrak{G}} \left| d^4x/\partial\mathcal{R} \right\rangle' \int \left\| \frac{\widetilde{\partial\phi'_{\sigma\rho}}}{\partial\phi^{\dagger}_{\sigma\rho}} \right\| -$

$\left\langle \frac{\partial\phi^{\star}_{\sigma\rho}}{\partial^{\mathfrak{H}}_{\mu\nu}} \right| \partial \uparrow/\partial t \setminus \partial \downarrow/\partial t \partial^2 \square \left| \begin{matrix} \blacksquare^{\circlearrowleft} \\ \blacksquare_{\circlearrowright} \end{matrix} \right. \partial^2\varphi/\partial\psi \blacksquare \rangle \Lambda^{\mu}_{\nu} \sum_{0 \leq \iota \leq m} P(\iota, J) \prod_{k=1}^{n} A_k \cup_{n=1}^{m} (X_n \cap Y_n) \cup_{n=1}^{m} (X_n \cap Y_n) \circledcirc \Lambda^{\mu}_{\nu} \odot \Gamma^{\mu}_{\nu}$ with respect to a heavy particle ρ, whether dark or white (star particle), as appropriate, regarding the criticality of its mass and/or energy $\langle 0|\sum_{\delta} \partial \mathfrak{m}/\partial \mathfrak{e}\rangle$ or its interaction with a graviton or a gravitin, as appropriate, in coordinates $\langle \rho^{\mu}\rho^{\nu}\rho^{\sigma}\rho^{\varrho} \rangle$, the latter, which occurs by permeabilization of the gravitonic or supergravitonic field in $\blacksquare = \int \langle \partial\mathfrak{G}/\partial\mathfrak{S}\mathfrak{G} \rangle$, what corresponds to the space – quantum deformed in $\mathfrak{C}_{\mathfrak{H}\Re} = \langle \sum_{\blacksquare}^{\sigma\rho} \mathcal{R}_{\nu}^{\mu\dagger} \big| \otimes \mathcal{H}_{\mu}^{\nu\star} \rangle$ the which in dimensions $\mathbb{R}^{\eta}$, represents, quantum gravity or supergravity by curvature or supercurvature of multidimensional quantum space-time.

Keywords: Quantum supergravity, quantum gravity, dark particle, star particle, quantum theory of relativistic fields.

Artículo recibido 02 enero 2026
Aceptado para publicación: 30 enero 2026

INTRODUCCIÓN.

En este punto, es indispensable establecer las bases teóricas que conforman la Teoría Cuántica de Campos Relativistas (TCCR) y que se encuentran desarrolladas en trabajos previos. Por tanto, estos son los puntos más relevantes.

1. Todo campo cuántico, es curvo por acción inmediata de una partícula cuya masa y/o energía alcanzan el mayor grado de criticidad. En este caso, la gravedad es endógena o implícita, es decir, una cualidad propia de la partícula interactuante.

2. Siguiendo lo dicho, en el numeral que antecede, las partículas se dividen en:

2.1. Partículas Supermasivas (Tipo IA): Son aquellas, cuyo centro de masa/energía en unidades de Planck dados en $\mathcal{M}_p = \sqrt{\frac{\hbar c}{\mathfrak{G}}} \approx 2{,}18 \times 10^{-8}\, kg$ (masa) y $E_p = \frac{\hbar}{t_P}$, $E_p = m_P c^2$, $E_p = \sqrt{\frac{\hbar c^5}{G}} \approx 1.956 \times 10^9$ J $\approx 1.22 \times 10^{19}$GeV ≈ 0.5433MWh$\sqrt{\frac{\hbar c^5}{8\pi G}} \approx 0.390 \times 10^9$ J $\approx 2.43 \times 10^{18}$GeV (energía $\approx 10^{-120}$), alcanza el mayor grado de criticidad, deformando el espacio – tiempo cuántico, lo que afecta el estado fundamental de los orbitales (espín, momentum, velocidad, trayectorias, etc), desplegados por las partículas repercutidas. Esta partícula también se la denomina "partícula oscura", en la medida en que, su centro de energía/masa, es oscuro. Principal candidata para explicar la materia oscura, en la medida en que, la gravedad converge en su centro, absorbiendo energía y materia.

2.2. Partículas Blancas (Tipo IB): Son aquellas, cuyo centro de masa/ energía en unidades de Planck, alcanza el mayor grado de criticidad, deformando el espacio – tiempo cuántico, lo que afecta el estado fundamental de los orbitales (espín, momentum, velocidad, trayectorias, etc), desplegados por las partículas repercutidas. Esta partícula también se la denomina "partícula estrella", en la medida en que, su centro de masa/energía es extremadamente denso, superando la masa, temperatura y energía de Planck, en $\mathcal{M}_p = \sqrt{\frac{\hbar c}{\mathfrak{G}}} \approx 2{,}18 \times 10^{-8}\, kg$ (masa), $\mathcal{M}_p = \sqrt{\frac{\hbar c^5}{\mathfrak{G}\hbar^2}}$ $T_P \approx 1.416784(16) \times 10^{32}$K (temperatura) y $E_p = \frac{\hbar}{t_P}$, $E_p = m_P c^2$, $E_p = \sqrt{\frac{\hbar c^5}{G}} \approx 1.956 \times 10^9$ J $\approx 1.22 \times 10^{19}$GeV ≈ 0.5433MWh$\sqrt{\frac{\hbar c^5}{8\pi G}} \approx 0.390 \times 10^9$ J $\approx 2.43 \times 10^{18}$GeV. También se la denomina "partícula estrella".

2.3. Hiperpartículas (Tipo IIA): Son aquellas, cuyo centro de masa/energía es extremadamente bajo, en unidades de Planck, más sin embargo, son capaces de igualar o superar la velocidad de la luz.

2.4. Suprapartículas (Tipo IIB): Son aquellas, cuyo centro de masa/energía es el equivalente al de una partícula oscura o blanca, más sin embargo, éstas, a diferencia de las referidas en los numerales 2.1 y 2.2, ésta igual o supera la velocidad de la luz.

3. Agujero negro cuántico: Fenómeno que ocurre en un espacio cuántico de Sitter, esto es, cuando una partícula oscura colisiona con otra o en su defecto, cuando una partícula blanca colisiona con otra o cuando una partícula blanca y una partícula oscura colisionan entre sí. Los agujeros negros cuánticos, también se forman por el colapso (por compresión gravitacional) o por la aniquilación (por interacción) de una partícula oscura o de una partícula blanca. Lo primero, ocurre cuando se atraen mutuamente por gravedad en tanto que lo segundo, ocurre cuando su centro de masa/energía alcanza el mayor grado de criticidad posible. En el centro del agujero negro cuántico, se encuentra la masa de la partícula aniquilada o comprimida, la que comporta condiciones gravitatorias extremas. Ahí es donde radica la singularidad de un agujero negro cuántico. La información que ingresa al agujero negro cuántico, no se destruye, muy al contrario, se transforma en materia y energía, las mismas que son repulsadas por el agujero negro cuántico blanco que se encuentra en el otro extremo del agujero cuántico de gusano. Por tanto, la materia y energía atrapada por el agujero negro cuántico, se convierte en materia y energía oscuras interferidas por gravedad extrema.

4. Agujero cuántico de gusano: Túnel cuántico por el cual, se conectan un agujero negro cuántico y un agujero blanco cuántico. A través de este túnel, por teletransportación cuántica, la información es procesada y convertida en materia y energía, todo esto, en un espacio de Sitter.

5. Agujero blanco cuántico: Fenómeno que ocurre en un espacio cuántico de Sitter, volviéndose la región de salida o repulsión de materia y energía, a propósito de lo que devora el agujero negro cuántico y de lo que procesa el canal cuántico de gusano. Lo que repulsa el agujero blanco cuántico, es materia y energía procesadas.

6. Espacio – tiempo cuántico: Entiéndase por espacio – tiempo cuántico, al campo en sí mismo, cuya Longitud de Planck, es superior a $\ell_{\mathcal{P}} = \sqrt{\frac{\hbar G}{c^3}} \approx 1{,}616199(97) \times 10^{-35}\ metros$**.** La métrica es la

curvatura escalar de Ricci, así: $\mathcal{R} = \sum_{\alpha,\beta=0}^{3} g^{\alpha\beta}\mathcal{R}_{\alpha\beta} \approx o\left(\mathcal{L}_p^{-2}\right) \approx 3{,}828 \cdot 10^{69} m^{-2}$. Ahora bien, el espacio – tiempo cuántico puede ser, bien de Sitter (dS) o bien, anti de Sitter (AdS). En el primero, se forma la curvatura cuántica y sus subniveles, subespacios o subcapas, en tanto que en el segundo, se forman los agujeros cuánticos y las multidimensiones.

7. Todo campo cuántico, es curvo por acción inmediata de la gravedad, esto a propósito de la existencia (Modelo – Higgs):

7.1. De un campo gravitónico, es decir, cuando una partícula cualquiera, interactúa con un gravitón, lo que supone la permeabilidad del campo cuántico, por un campo gravitónico que transfiere gravedad al campo primario, curvándolo.

7.2. De un campo supergravitónico, es decir, cuando una partícula cualquiera, interactúa con un gravitino o supergravitón, lo que supone la permeabilidad del campo cuántico, por un campo gravitónico que transfiere gravedad al campo primario, deformándolo.

7.3. Lo referido en este numeral se denomina gravedad exógena.

8. La gravedad cuántica, sea endógena o exógena comporta la curvatura del espacio – tiempo cuántico, en tanto que, la supergravedad cuántica, sea endógena o exógena, comporta la deformación (supercurvatura) del espacio – tiempo cuántico, formándose pliegues multidimensionales (en alta configuración – membranas dimensionales) en rango superior a $\mathbb{R}^4 - AdS$. Cabe indicar que las membranas dimensionales, se dividen en TIPO I y TIPO II respectivamente, la primera a propósito de la curvatura del campo en gravedad cuántica y la segunda, la deformación del campo en supergravedad cuántica, todo esto, lo cual también depende de la naturaleza de la gravedad que interfiere, es decir, si es exógena o endógena, lo que llamaríamos membranas dimensionales tipo IA, IB, IIA y IIB respectivamente, las cuales, pueden contener dimensiones y subdimensiones infinitas, en relación a las interacciones de la partícula que provoca de la deformación del espacio – tiempo cuántico. Esto es lo que llamamos supersimetrías de gauge en dimensiones altas a $\mathbb{R}^4$, es decir, cuando estamos ante membranas dimensionales tipo IA, IB y IIB, según sea el caso en tanto que, las membranas dimensionales del tipo IIA, contienen dimensiones infinitas en $\mathbb{R}^4 - dS$.

9. Cuando una partícula colisiona con otra y se aniquilan o cuando la partícula pesada colapsa por compresión, la extinción provoca ondas cuánticas que se desplazan en longitud sobre el campo cuántico deformado el mismo que, es superfluido.

10. El puente ER, en esta teoría, explica la superposición y el entrelazamiento cuánticos en sentido estricto, en un espacio AdS.

11. Los enunciados antes referidos, aplican a la antimateria, es decir, a la región de antipartículas.

12. La brecha de masa, provoca la curvatura del espacio – tiempo cuántico pero no lo deforma por completo, pues este fenómeno, no ocurre con una partícula deformante, sino en partículas ligeras como las hiperpartículas, esto en la medida en que, no registran estado de vacío.

13. Adicionalmente, es importante, establecer las siguientes reglas:

13.1. La gravedad cuántica relativista, ocurre concretamente en un espacio cuántico de Sitter, en el que se pueden formar subdimensiones o subespacios dentro del límite de $\mathbb{R}^4$.

13.2. La supergravedad cuántica relativista, ocurre concretamente en un espacio cuántico anti de Sitter, en el que se pueden formar hiperespacios o dimensiones más altas, superiores a $\mathbb{R}^4$.

13.3. Las partículas propuestas, viajan en gravedad cuántica más, interactúan en supergravedad cuántica por permeabilización.

13.4. Cualquier partícula, de las aquí propuestas, se puede convertir en otra, por aniquilación, siguiendo los diagramas de Feynman.

13.5. Las dimensiones en alta configuración así como las de ensamble, son infinitas.

13.6. La materia y energía oscuras, están formadas esencialmente por partículas aniquiladas o colapsadas por gravedad. En consecuencia, es la criticidad de la masa la que las vuelve compatibles.

13.7. Los agujeros cuánticos, absorben partículas ligeras y pesadas, sin distinción, lo que explica la expansión del universo por acción gravitacional en la materia.

13.8. Las partículas aquí propuestas, son susceptibles de enganche, como ocurre con un diquark.

13.9. En esta teoría, se incorpora el concepto de cuerda, pero en un espacio cuántico anti de Sitter.

13.10. Las partículas pesadas, cuando se desplazan de un punto a otro en forma infinita hasta su aniquilación o colapso, lo hacen por medio de gravedad, deformando, en el caso de las partículas blancas

y las hiperpartículas, un espacio de Sitter, creando capas dimensionales en límite de $\mathbb{R}^4$ en tanto que, la partícula oscura, crea capas dimensiones en alta configuración a $\mathbb{R}^4$ en un espacio anti de Sitter.

13.11. La hiperpartícula es la única en este modelo, que no tiene masa, es por ello que puede viajar a la velocidad de la luz.

13.12. La suprapartícula es por excepción, un caso de mutación por aniquilación, en la medida en que, pese a tratarse de una partícula pesada, con un centro de masa/energía extremadamente crítico y denso, es capaz de viajar a la velocidad de la luz. La suprapartícula solamente existe por aniquilación en entre dos o más partículas pesadas, quedando excluidas las partículas ligeras. Adicionalmente, la suprapartícula, tiene la capacidad de desplazarse entre dimensiones dS y AdS, lo que esta teoría denomina dimensiones en $\mathbb{R}^{\eta}$. En consecuencia, las dimensiones por gravedad y supergravedad, pueden intersecarse por gravedad. En este punto, es pertinente para efectos de ejemplificar, citar el diagrama de Penrose expandido al infinito.

13.13. Los campos de las partículas ligeras, son deformados por acción a distancia, debido a las interacciones de una partícula pesada, esto es, por gravedad.

13.14. Solamente las partículas pesadas pueden deformar el campo propio y de las partículas ligeras, por acción de la gravedad que se desprende de su centro de masa/energía extremo. En consecuencia, la gravedad endógena, se materializa por impermeabilización del campo de Braut – Englert – Higgs respecto de la partícula pesada. El bosón de Higgs es el que transfiere la masa, a las partículas pesadas, aniquilándose con éstas.

13.15. La gravedad exógena, se vuelve posible, por permeabilización de un campo cuántico arbitrario, lo que, como ha quedado explicado en esta teoría, funciona como un mecanismo de Higgs.

13.16. El colapso de una partícula pesada, ocurre por la expansión de su centro de masa/energía, debido a la gravedad interferente, ditalación que es comprimida en contrario, por los límites del campo de la partícula de que se trate, lo que provoca, la deformación del plano cuántico e incluso la formación de agujeros cuánticos, según la criticidad de los valores de masa/energía involucrados.

13.17. La fusión de campos cuánticos, es posible, por acción de la gravedad entre ambos, lo que vuelve posible, su aniquilación.

13.18. Las ondas en un plano cuántico, no solamente se forman por la aniquilación o colpaso de una partícula pesada, sino también, cuando viaja de un punto a otro.

13.19. Las partículas ligeras, crean gravedad mínima a propósito de su centro de masa/energía, la cual sin embargo, es imperceptible aunque superior a cero, pues, contribuye a la aniquilación con otro campo más pesado.

13.20. La gravedad endógena, se debe a que, el campo de Higgs, y por ende, el bosón de Higgs, no solamente transfiere masa a las partículas pesadas y ligeras, con excepción de la hiperpartícula, sino que también, le dota de gravedad, a propósito de la masa transferida.

13.21. Esta teoría es estrictamente de gauge.

RESULTADOS Y DISCUSIÓN:

Suponemos que, en un mapa cuántico de Einstein – Hilbert, una partícula deformante $\alpha\beta\gamma\delta$ se desplaza en el espacio cuántico, en el que interactúa, deformando el plano por gravedad, y por ende, creando, bien dimensiones altas en $\mathbb{R}^4 - AdS$ por supercurvatura (supergravedad cuántica) o bien, dimensiones en $\mathbb{R}^4 - ds$ por curvatura, esto es, en condiciones de gravedad. Para estos efectos, una partícula deformante debe colapsar por compresión gravitacional, aniquilarse cuando interactúa con otras más inestables o con otra partícula pesada, o por permeabilidad del campo gravitónico o supergravitónico en el espacio cuántico curvo, esto último, lo cual ocurre, cuando una partícula pesada interactúa con el gravitón o el gravitino (supergravitino), según sea el caso. Por tanto, la gravedad actúa a nivel cuántico, sea por aniquilación, compresión, ésta última gravitacional o por permeabilización. Suponemos en simultáneo, que una vez, causada la aniquilación o compresión por gravedad, de una partícula pesada o cuando ocurre la permeabilización, se produce, bien la curvatura cuántica, cuya métrica es el tensor es de Riemann – Ricci – Einstein, incluyendo el flujo de la simetría, o en su defecto, la supercurvatura de Weyl, cuya métrica es la de Chern-Simons-Nambu-Goto para supergravedad. La primera, produce subcampos que son subdimensiones de un mismo plano de Sitter (dS), en tanto que la segunda, produce campos en dualidad holográfica, que son dimensiones altas al plano cuatridimensional en un espacio anti de Sitter (AdS). En este sentido, el campo pasa a ser no homeomorfo, difeomorfo e isométrico, afectando los orbitales de las partículas cuyo centro de masa/energía es inferior en unidades de Planck (partículas ligeras) en relación a la partícula que deforma el plano. La interacción y/o aniquilación de

estas partículas deformantes, provoca un agujero negro cuántico (con excepción de las interacciones dadas por las hiperpartículas tipo IIA), formado por materia y energía oscuras, cuya naturaleza es fermiónica/bosónica, esto a propósito de que, la partícula aniquilada o comprimida, engendra materia y energía oscuras, lo que no ocurre en escenarios de permeabilización gravitónica más sí, en escenarios de permeabilización supergravitónica. El agujero cuántico de salida, es blanco, por ende, repulsivo de materia y energía transformada por la gravedad, a través del tracto Einstein – Rosen. Cuando la materia y la energía son transformadas en oscuras, por la gravedad, éstas se comprimen hasta un punto de no retorno/densidad supermasiva, causando dos especies de singularidad inherentes al agujero negro cuántico, siendo éstas, primaria y secundaria, la primera en la que la gravedad es extrema y deforma la materia y la energía, fundiéndose con el núcleo del agujero negro cuántico (que contiene la partícula muerta) y la segunda, en la que la gravedad transforma la materia y la energía, desplazándola a través del tracto Einstein – Rosen y expulsándola a través de un agujero blanco cuántico. Esto es lo que ocurre en escenarios de entrelazamiento y túneles cuánticos supermasivos en los que, la partícula deformante genera gravedad extrema. Llámese también, gravedad absoluta. Queda claro entonces, que el sistema cuántico de agujeros, no se produce en condiciones de gravedad relativa, esto es, cuando ocurre únicamente la curvatura cuántica por gravedad moderada, lo que sucede por ejemplo, con las interacciones dadas por las hiperpartículas tipo IIA o en el caso de la brecha de masa de las partículas ligeras respecto del estado de vacío.

Dicho lo anterior, es que, propongo una posible alternativa de solución al problema del milenio de Yang – Mills y la brecha de masa, a partir de la Teoría Cuántica de Campos Relativistas, la cual se constituye además, como un intento por reconciliar la relatividad general y la mecánica cuántica.

A partir de aquí, sugerimos los cálculos de instantones (para regular la brecha de masa y la densidad de energía por carga), osciladores, propagadores, operadores, mapas, coordenadas vectoriales, orbitales, correladores, propulsores, tensores de stress por curvatura, torsión, escalares, spinors, potenciadores, simetrías y supersimetrías de calibre abelianas y no abelianas en relación a las partículas pesadas y sus interacciones con el espacio cuántico deformado, en tanto que respecto de éste último, los cálculos están vinculados a su geometría e hipergeometría (análisis cohomológico), incluyendo los agujeros cuánticos,

no sin antes aclarar, que las demostraciones matemáticas contenidas en trabajos anteriores, son interdependientes a éste manuscrito y sus diez volúmenes.

Aclarado lo anterior, pasamos a precisar que el Modelo aquí referido, se divide en:

1. Supergravedad cuántica en SYM (Super Yang – Mills).

2. Gravedad cuántica en YM (Yang – Mills).

3. Agujeros cuánticos en YM (Yang – Mills).

4. Modelo de Unificación.

Las métricas usadas son, entre otras:

- Espacios de Einstein – Hilbert.
- Métrica de Chern – Simons.
- Métrica de Kaluza – Klein.
- Métrica de Nambu – Goto.
- Métrica de Feynman – Wheeler.
- Métrica de Born – Oppenheimer.
- Métrica de Hartree – Fock.
- Métrica de Yang – Mills.
- Métrica de Kerr – Newman.
- Espacios de Sitter y anti de Sitter.
- Espacios de Riemann – Perelman – Poincaré.
- Tensores y flujo de Ricci.
- Métrica de Green.
- Métrica de Goldstone.
- Métrica de Brout – Englert – Higgs.
- Métrica de Schwinger – Dyson.
- Métrica de Yukawa.
- Métrica de Von Neumann
- Métrica de Friedman.

MODELO DOS. GRAVEDAD CUÁNTICA EN YM (CONTINUACIÓN).

$$-p^{i,j}$$

$$b^{i,j} = \bar{\partial}_x q_y^{i,j} - \bar{\partial}_y q_x^{i,j}$$

$$\rho^{i,j}, J^{i,j}$$

$$H = \frac{1}{2}\sum_{i,j}\left[p_s^{i,j}p_s^{i,j} + (b^{i,j})^2 + \left(\rho^{i,j} + \bar{\partial}_s p_s^{i,j}\right)q_0^{i,j} - J_s^{i,j}q_s^{i,j}\right]$$

$$b^{i,j} := \bar{\partial}_x q_y^{i,j} - \bar{\partial}_y q_x^{i,j},$$

$$\bar{\partial}_x f^{i,j} = \frac{1}{2a}\left(f^{i,j+1} - f^{i,j-1}\right), \bar{\partial}_y f^{i,j} = \frac{1}{2a}\left(f^{i+1,j} - f^{i-1,j}\right),$$

$$p_0^{i,j} = \frac{\delta L}{\delta \dot{q}_0^{i,j}} = 0 \; \forall, i, j.$$

$$\begin{cases} p_x^{i,j} = -\bar{\partial}_y b^{i,j} + J_x^{i,j} \\ p_y^{i,j} = \bar{\partial}_x b^{i,j} + J_y^{i,j} \end{cases}$$
$$b^{i,j} = \bar{\partial}_x p_y^{i,j} - \bar{\partial}_y p_x^{i,j}$$
$$C_p^{i,j} = \bar{\partial}_x p_x^{i,j} + \bar{\partial}_y p_y^{i,j} + \rho^{i,j} = 0$$
$$\bar{\partial}_x J_x^{i,j} + \bar{\partial}_y J_y^{i,j} + \rho^{i,j} = 0$$

$$\dot{p}_0^{i,j} = -\frac{\delta H}{\delta q_0^{i,j}} = 0 \implies \mathcal{C}_\rho^{i,j} := \bar{\partial}_x p_x^{i,j} + \bar{\partial}_y p_y^{i,j} + \rho^{i,j} = 0$$

$$H = \frac{1}{2}\sum_{i,j}\left[p_s^{i,j}p_s^{i,j} + \left(b^{i,j}\right)^2 - J_s^{i,j}q_s^{i,j}\right]$$

$$\mathcal{C}^{i,j} := \bar{\partial}_x p_x^{i,j} + \bar{\partial}_y p_y^{i,j}$$

$$\left[\hat{q}_s^{i,j}, \hat{p}_r^{n,m}\right] = \mathrm{i}\hbar \cdot \delta_{s,r}\delta^{i,n}\delta^{j,m} \; \forall s, r = \{x, y\} \; \forall i, j$$

$$\hat{H} = \frac{1}{2}\sum_{i,j}\left[\left(\hat{p}_x^{i,j}\right)^2 + \left(\hat{p}_y^{i,j}\right)^2 + \left(\hat{b}^{i,j}\right)^2\right]$$

$$\hat{\mathcal{C}}^{i,j}|\psi\rangle_{\text{phys}} = 0 \; \forall i, j.$$

$$\mathcal{H}_{\text{kin}} = \bigotimes_{i,j} \mathcal{H}_{i,j}^{\text{kin}}$$

$$\hat{p}_s^{i,j} := -\mathrm{i}\hbar\frac{\partial}{\partial q_s^{i,j}}$$

$$\hat{\Pi} = \prod_{i,j}\delta\left(\hat{\mathcal{C}}^{i,j}\right) = \prod_{i,j}\int_{-\infty}^{+\infty} dz^{i,j}\, \mathrm{e}^{\mathrm{i}z^{i,j}\cdot\hat{\mathcal{C}}^{i,j}}$$

$$|\psi\rangle_{\text{phys}} = \hat{\Pi}|\psi\rangle_{\text{kin}}$$

$$\hat{\mathcal{C}}_{\rho}^{i,j}|\psi\rangle_{\text{phy}} = \left(\hat{\mathcal{C}}^{i,j} + \hat{\rho}^{i,j}\right)|\psi\rangle_{\text{phy}} = 0 \;\forall i,j$$

$$\mathcal{H}^{M} = \bigoplus_{n=0}^{N^2} \text{span}\{|s_n\rangle_M\} = \bigoplus_{n=0}^{N^2} \bigoplus_{s_n} |s_n\rangle_M$$

$$\mathcal{E}_0 = \frac{1}{2}\hbar \sum_{\alpha,\beta} |\bar{\mathbf{k}}(\alpha,\beta)|$$

$$\Psi_0[p] = A\delta(\mathcal{C})\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m} G(i-n,j-m)p_s^{i,j}p_s^{n,m}\right]$$

$$G(i-n,j-m) = \frac{1}{N^2}\sum_{\alpha,\beta}\frac{1}{|\bar{\mathbf{k}}|}\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}[(i-n)\alpha+(j-m)\beta]},$$

$$\delta(\mathcal{C}) = \prod_{i,j}\delta\left(\bar{\partial}_x p_x^{i,j} + \bar{\partial}_y p_y^{i,j}\right)$$

$$|\psi^0\rangle_F = A\int \mathcal{D}p\Psi_0[p]\prod_{I,J}|p_x^{I,J},p_y^{I,J}\rangle_F = A\int \mathcal{D}p\delta(\mathcal{C})\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m} G(i-n,j-m)p_s^{i,j}p_s^{n,m}\right]\prod_{I,J}|p_x^{I,J},p_y^{I,J}\rangle_F,$$

$$\mathcal{E}_\rho - \mathcal{E}_0 = \frac{1}{2}\sum_{i,j}\sum_{n,m} D(i-n,j-m)\rho^{i,j}\rho^{n,m}$$

$$D(i-n,j-m) = \frac{1}{N^2}\sum_{\alpha,\beta}\frac{1}{|\bar{\mathbf{k}}|^2}\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}[(i-n)\alpha+(j-m)\beta]}$$

$$D(\mathbf{r}-\mathbf{r}') = \frac{1}{|\mathbf{r}-\mathbf{r}'|}$$

$$\Psi_{0,\rho}[p] = A\delta(\mathcal{C}_\rho)\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m} G(i-n,j-m)\left(p_s^{i,j}-p_{s,\rho}^{i,j}\right)\left(p_s^{n,m}-p_{s,\rho}^{n,m}\right)\right],$$

$$p_{s,\rho}^{i,j} = \sum_{n,m}\bar{\partial}_s^{\{i,j\}}D(i-n,j-m)\rho^{n,m}$$

$$\delta(\mathcal{C}_\rho) = \prod_{i,j}\delta\left(\bar{\partial}_x p_x^{i,j} + \bar{\partial}_y p_y^{i,j} + \rho^{i,j}\right)$$

$$\left|\psi_\rho^0\right\rangle_F = A\int \mathcal{D}p\Psi_{0,\rho}[p]\prod_{i,j}\left|p_x^{i,j},p_y^{i,j}\right\rangle_F$$

$$= A\int \mathcal{D}p\delta(\mathcal{C}_\rho)\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m} G(i-n,j-m)\Delta p_{s,\rho}^{i,j}\Delta p_{s,\rho}^{n,m}\right]\prod_{i,j}\left|p_x^{i,j},p_y^{i,j}\right\rangle_F$$

$$\mathcal{A}_A = \bigoplus_{K,R} \mathcal{O}_A^K \otimes \mathbb{I}_{\bar{A}}^{R,K}, \overline{\mathcal{A}}_A = \bigoplus_{K,R} \mathbb{I}_A^K \otimes \mathcal{O}_{\bar{A}}^{R,K}, \mathcal{H} = \bigoplus_{K,R} \mathcal{H}_A^K \otimes \mathcal{H}_{\bar{A}}^{R,K},$$

$$\hat{\mathcal{C}}_\rho^{i,j}|\psi\rangle_{\text{phy}} = \left(\hat{\mathcal{C}}^{i,j} + \hat{\rho}^{i,j}\right)|\psi\rangle_{\text{phy}} = 0 \; \forall i,j,$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_K \mathcal{H}_A^K \otimes \mathcal{H}_{\bar{A}}^K$$

$$\mathcal{A}_A = \bigoplus_K \mathcal{O}_A^K \otimes \mathbb{I}_{\bar{A}}^K, \overline{\mathcal{A}}_A = \bigoplus_K \mathbb{I}_A^K \otimes \mathcal{O}_{\bar{A}}^K$$

$$\mathcal{H} = \bigoplus_{n=0}^{N^2} \bigoplus_{s_n} |s_n\rangle_M \otimes \left(\bigoplus_{k,r} \mathcal{H}_A^k \otimes \mathcal{H}_{\bar{A}}^{r,k} \right)$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_{n=0}^{N^2} \bigoplus_{s_n} |s_n\rangle_M \otimes \left(\bigoplus_{k(s_n)} \mathcal{H}_A^{k(s_n)} \otimes \mathcal{H}_{\bar{A}}^{s_n, k(s_n)} \right)$$

$$|\Upsilon\rangle = \eta \sum_{n=0}^{N^2} \sum_{s_n} \alpha_{s_n} |s_n\rangle_M \left|\Psi_{s_n}\right\rangle_F,$$

$$\frac{1}{2}(|L\rangle^A + |R\rangle^A)(|L\rangle^B + |R\rangle^B) \to \frac{e^{i\phi}}{2}\left[|L\rangle^A\left(|L\rangle^B + e^{i\Delta\phi_-}|R\rangle^B\right) + |R\rangle^A\left(e^{i\Delta\phi_+}|L\rangle^B + |R\rangle^B\right)\right].$$

$$\mathcal{A}_{AB} = \bigoplus_K \mathcal{O}_A^K \otimes \mathcal{O}_B^K \otimes \mathbb{I}\frac{K}{AB}, \mathcal{H}_{\text{phy}} = \bigoplus_K \mathcal{H}_A^K \otimes \mathcal{H}_B^K \otimes \mathcal{H}\frac{K}{AB},$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_{s,k(s)} |s\rangle_M \otimes \mathcal{H}_A^{k(s)} \otimes \mathcal{H}_B^{k(s)} \otimes \mathcal{H}_{\frac{s}{AB}}^{s,k(s)}$$

$$\mathcal{H}_{\text{phy}} = \mathcal{H}^\sigma \otimes \left[\bigoplus_s |s\rangle_M \otimes \left(\bigoplus_{k(s)} \mathcal{H}_A^{k(s)} \otimes \mathcal{H}_B^{k(s)} \otimes \mathcal{H}^{s,k(s)}\frac{AB}{} \right) \right],$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_K \underbrace{(\mathcal{H}_A^K \otimes \mathcal{H}_\sigma^A)}_{\mathcal{H}_A^{K,\,\text{tot}}} \otimes \underbrace{(\mathcal{H}_B^K \otimes \mathcal{H}_\sigma^B)}_{\mathcal{H}_B^{K,\,\text{tot}}} \otimes \mathcal{H}\frac{K}{AB},$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_K \mathcal{H}_A^{K,\,\text{tot}} \otimes \mathcal{H}_B^{K,\,\text{tot}} \otimes \mathcal{H}\frac{K}{AB}$$

$$\mathcal{A}_A^{\text{tot}} = \bigoplus_K \mathcal{O}_A^{K,\,\text{tot}} \otimes \mathbb{I}_B^K \otimes \mathbb{I}_{AB}^K, \mathcal{A}_B^{\text{tot}} = \bigoplus_K \mathbb{I}_A^K \otimes \mathcal{O}_B^{K,\,\text{tot}} \otimes \mathbb{I}\frac{K}{AB}$$

$$|s^0\rangle_M = |s^0\rangle_M^A |s^0\rangle_M^B |s^0\rangle_M^{\overline{AB}},$$

$$|\Psi\rangle^{(0)} = |+\rangle_\sigma^A |+\rangle_\sigma^B |s^0\rangle_M \left|\psi_{s^0}^0\right\rangle_F.$$

$$|\uparrow\rangle_\sigma^{\cdot}|s^0\rangle_M^{\cdot} \to |\uparrow\rangle_\sigma^{\cdot}|s^L\rangle_M^{\cdot}, |\downarrow\rangle_\sigma^{\cdot}|s^0\rangle_M^{\cdot} \to |\downarrow\rangle_\sigma^{\cdot}|s^R\rangle_M^{\cdot},$$

$$|\Psi\rangle^{(1)} = \hat{U}|\Psi\rangle^{(0)} = \sum_s |\sigma(s)\rangle_\sigma^{AB}|s\rangle_M|\psi_s'\rangle_F,$$

$$|\Psi\rangle^{(2)} = \sum_s e^{i\gamma(s)}|\sigma(s)\rangle_\sigma^{AB}|s\rangle_M|\psi_s^0\rangle_F.$$

$$|\Psi\rangle^{(3)} = \exp\left(-\mathrm{i}\frac{\hat{H}}{\hbar}\tau\right)|\Psi\rangle^{(2)} = \sum_s \mathrm{e}^{i[\gamma(s)+\phi(s)]}|\sigma(s)\rangle_\sigma^{AB}|s\rangle_M|\psi_s^0\rangle_F.$$

$$\phi(s) := -\frac{\mathcal{E}_{\rho(s)} - \mathcal{E}_0}{\hbar}\tau$$

$$|\Psi\rangle^{(4)} = \hat{U}'|\Psi\rangle^{(3)} = |s^0\rangle_M \sum_s \mathrm{e}^{i[\gamma(s)+\phi(s)]}|\sigma(s)\rangle_\sigma^{AB}\left|\psi_{s^0}'(s)\right\rangle_F,$$

$$|\Psi\rangle^{(5)} = |s^0\rangle_M\left|\psi_{s^0}^0\right\rangle_F \sum_s \mathrm{e}^{i[\gamma(s)+\gamma'(s)+\phi(s)]}|\sigma(s)\rangle_\sigma^{AB}$$

$$|\chi\rangle_\sigma^{AB} := \sum_s \mathrm{e}^{i[\gamma(s)+\gamma'(s)+\phi(s)]}|\sigma(s)\rangle_\sigma^{AB}$$

$$|\Psi\rangle^{(5)} = |\chi\rangle_\sigma^{AB}|s^0\rangle_M\left|\psi_{s^0}^0\right\rangle_F$$

$$|\Psi\rangle^{(0)} = |+\rangle_\sigma^A|+\rangle_\sigma^B|\psi^0\rangle_{F,M} \overset{\text{BMV}}{\to} |\Psi\rangle^{(5)} = |\chi\rangle_\sigma^{AB}|\psi^0\rangle_{F,M},$$

$$\hat{U} = \hat{U}_A \otimes \hat{U}_B, \hat{U}' = \hat{U}_A' \otimes \hat{U}_B'$$

$$L: \sigma_A + M_A + F, R: \sigma_B + M_B + F_B + M_{\overline{AB}} + F_{\overline{AB}},$$

$$H(S)_\rho = -\mathrm{Tr}(\hat{\rho}\ln\,\hat{\rho}),$$

$$\mathrm{ent}_{LR,K}^{(5)} - \mathrm{ent}_{LR,K}^{(0)} = H(\sigma_A)_\chi.$$

$$\partial_x^+ f^{i,j} = \frac{1}{a}\left(f^{i,j+1} - f^{i,j}\right), \partial_x^- f^{i,j} = \frac{1}{a}\left(f^{i,j} - f^{i,j-1}\right),$$
$$\partial_y^+ f^{i,j} = \frac{1}{a}\left(f^{i+1,j} - f^{i,j}\right), \partial_y^- f^{i,j} = \frac{1}{a}\left(f^{i,j} - f^{i-1,j}\right),$$

$$\bar{\partial}_x f^{i,j} = \frac{1}{2}\left(\partial_x^+ f^{i,j} + \partial_x^- f^{i,j}\right) = \frac{1}{2a}\left(f^{i,j+1} - f^{i,j-1}\right),$$
$$\bar{\partial}_y f^{i,j} = \frac{1}{2}\left(\partial_y^+ f^{i,j} + \partial_y^- f^{i,j}\right) = \frac{1}{2a}\left(f^{i+1,j} - f^{i-1,j}\right),$$

$$\bar{\partial}_x\bar{\partial}_y f^{i,j} = \frac{1}{4a^2}\left(f^{i+1,j+1} - f^{i+1,j-1} - f^{i-1,j+1} + f^{i-1,j-1}\right) = \bar{\partial}_y\bar{\partial}_x f^{i,j}$$

$$\bar{\partial}_x\left(f^{i,j}g^{i,j}\right) = \frac{f^{i,j+1}g^{i,j+1} - f^{i,j-1}g^{i,j-1}}{2a}.$$

$$\bar{\partial}_x\left(f^{i,j}g^{i,j}\right)=g^{i,j+1}\bar{\partial}_x f^{i,j}+f^{i,j-1}\bar{\partial}_x g^{i,j}=g^{i,j-1}\bar{\partial}_x f^{i,j}+f^{i,j+1}\bar{\partial}_x g^{i,j}$$

$$\bar{\partial}_x\left(f^{i,j}g^{i,j}\right)=\frac{g^{i,j+1}+g^{i,j-1}}{2}\bar{\partial}_x f^{i,j}+\frac{f^{i,j+1}+f^{i,j-1}}{2}\bar{\partial}_x g^{i,j},$$

$$\bar{\partial}_y\left(f^{i,j}g^{i,j}\right)=\frac{g^{i+1,j}+g^{i-1,j}}{2}\bar{\partial}_y f^{i,j}+\frac{f^{i+1,j}+f^{i-1,j}}{2}\bar{\partial}_y g^{i,j}.$$

$$\frac{\mathrm{d}\left(\bar{\partial}_x f^{n,m}\right)}{\mathrm{d}f^{i,j}}=\frac{1}{2a}\left(\frac{\mathrm{d}f^{n,m+1}}{\mathrm{d}f^{i,j}}-\frac{\mathrm{d}f^{n,m-1}}{\mathrm{d}f^{i,j}}\right)=\frac{1}{2a}\left(\delta^{i,n}\delta^{j,m+1}-\delta^{i,n}\delta^{j,m-1}\right).$$

$$\bar{\partial}_x^{\{n,m\}}\left(\delta^{i,n}\delta^{j,m}\right):=\frac{\mathrm{d}\left(\bar{\partial}_x f^{n,m}\right)}{\mathrm{d}f^{i,j}}.$$

$$\sum_{n,m}\left[\bar{\partial}_x^{\{n,m\}}\left(\delta^{i,n}\delta^{j,m}\right)\cdot f^{n,m}\right]=\frac{1}{2a}\sum_{n,m}f^{n,m}\left(\delta^{i,n}\delta^{j,m+1}-\delta^{i,n}\delta^{j,m-1}\right)=\frac{f^{i,j-1}-f^{i,j+1}}{2a}=-\bar{\partial}_x f^{i,j},$$

$$\sum_{i,j}\left[\bar{\partial}_x^{\{n,m\}}\left(\delta^{i,n}\delta^{j,m}\right)\cdot f^{i,j}\right]$$

$$\sum_{n,m}\left[\bar{\partial}_s^{\{n,m\}}\left(\delta^{i,n}\delta^{j,m}\right)\cdot f^{n,m}\right]=-\bar{\partial}_s f^{i,j},\sum_{i,j}\left[\bar{\partial}_s^{\{n,m\}}\left(\delta^{i,n}\delta^{j,m}\right)\cdot f^{i,j}\right]=\bar{\partial}_s f^{n,m}.$$

$$L=\frac{1}{2}\sum_{i,j}\left[m\left(\dot{q}_x^{i,j}\right)^2+m\left(\dot{q}_y^{i,j}\right)^2-\kappa a^2\left(\bar{\partial}_x q_y^{i,j}-\bar{\partial}_y q_x^{i,j}\right)^2\right]-\kappa a^2\sum_{i,j}\left(\bar{\partial}_x\dot{q}_x^{i,j}+\bar{\partial}_y\dot{q}_y^{i,j}\right)q_0^{i,j}+F(q_0),$$

$$L=\frac{1}{2}\sum_{i,j}\left[\left(\dot{q}_x^{i,j}\right)^2+\left(\dot{q}_y^{i,j}\right)^2-\left(\bar{\partial}_x q_y^{i,j}-\bar{\partial}_y q_x^{i,j}\right)^2\right]-\sum_{i,j}\left(\bar{\partial}_x\dot{q}_x^{i,j}+\bar{\partial}_y\dot{q}_y^{i,j}\right)q_0^{i,j}+F(q_0).$$

$$p_0^{i,j}=\frac{\delta L}{\delta\dot{q}_0^{i,j}}=0\ \forall i,j,$$

$$p_s^{i,j}=\frac{\delta L}{\delta\dot{q}_s^{i,j}}=\dot{q}_s^{i,j}+\bar{\partial}_s q_0^{i,j}\ \forall i,j,s\in\{x,y\}$$

$$H=\sum_{s\in\{0,x,y\}}\sum_{i,j}p_s^{i,j}\dot{q}_s^{i,j}-L(q,p)$$

$$H=\sum_{i,j}p_0^{i,j}\dot{q}_0^{i,j}+\frac{1}{2}\sum_{i,j}\left[\left(p_x^{i,j}\right)^2+\left(p_y^{i,j}\right)^2+\left(b^{i,j}\right)^2\right]+\sum_{i,j}\left(\bar{\partial}_x p_x^{i,j}+\bar{\partial}_y p_y^{i,j}\right)q_0^{i,j}+G(q_0)-F(q_0)$$

$$H=\sum_{i,j}\left(p_0^{i,j}\dot{q}_0^{i,j}\right)+\frac{1}{2}\sum_{i,j}\left[\left(p_x^{i,j}\right)^2+\left(p_y^{i,j}\right)^2+\left(b^{i,j}\right)^2\right]+\sum_{i,j}\left[\left(\bar{\partial}_x p_x^{i,j}+\bar{\partial}_y p_y^{i,j}\right)q_0^{i,j}\right]$$

$$\dot{p}_0^{i,j}=-\frac{\delta H}{\delta q_0^{i,j}}=0\implies C^{i,j}:=\bar{\partial}_x p_x^{i,j}+\bar{\partial}_y p_y^{i,j}=0$$

$$\begin{cases} \dot{p}_x^{i,j} = -\dfrac{\delta H}{\delta q_x^{i,j}} = -\sum_{n,m} b^{n,m} \dfrac{\mathrm{d}b^{n,m}}{\mathrm{d}q_x^{i,j}} = \sum_{n,m} b^{n,m} \bar{\partial}_y^{\{n,m\}} \left(\delta^{i,n}\delta^{j,m}\right) = -\bar{\partial}_y b^{i,j} \\ \dot{p}_y^{i,j} = -\dfrac{\delta H}{\delta q_y^{i,j}} = -\sum_{n,m} b^{n,m} \dfrac{\mathrm{d}b^{n,m}}{\mathrm{d}q_y^{i,j}} = -\sum_{n,m} b^{n,m} \bar{\partial}_x^{\{n,m\}} \left(\delta^{i,n}\delta^{j,m}\right) = \bar{\partial}_x b^{i,j} \end{cases}$$

$$\dot{b}^{i,j} = \bar{\partial}_x \dot{q}_y^{i,j} - \bar{\partial}_y \dot{q}_x^{i,j} = \bar{\partial}_x p_y^{i,j} - \bar{\partial}_x \bar{\partial}_y \dot{q}_0^{i,j} - \bar{\partial}_y p_x^{i,j} + \bar{\partial}_y \bar{\partial}_x \dot{q}_0^{i,j} = \bar{\partial}_x p_y^{i,j} - \bar{\partial}_y p_x^{i,j}$$

$$\dot{\mathcal{C}}^{i,j} = \bar{\partial}_x \dot{p}_x^{i,j} + \bar{\partial}_y \dot{p}_y^{i,j} = -\bar{\partial}_x \bar{\partial}_y b^{i,j} + \bar{\partial}_y \bar{\partial}_x b^{i,j} = 0$$

$$H = \sum_{i,j} p_0^{i,j} \dot{q}_0^{i,j} + \frac{1}{2} \sum_{i,j} \left[\left(p_x^{i,j}\right)^2 + \left(p_y^{i,j}\right)^2 + \left(b^{i,j}\right)^2 \right] + \sum_{i,j} \left(\bar{\partial}_x p_x^{i,j} + \bar{\partial}_y p_y^{i,j}\right) q_0^{i,j} - \sum_{i,j} \rho^{i,j} q_0^{i,j} - \sum_{i,j} \left(J_x^{i,j} q_x^{i,j} + J_y^{i,j} q_y^{i,j}\right)$$

$$\mathcal{C}_\rho^{i,j} := \mathcal{C}^{i,j} + \rho^{i,j} = 0 \; \forall i,j,$$

$$\dot{\mathcal{C}}_\rho^{i,j} = 0 \Rightarrow \bar{\partial}_x \dot{p}_x^{i,j} + \bar{\partial}_y \dot{p}_y^{i,j} + \dot{\rho}^{i,j} = \bar{\partial}_x J_x^{i,j} + \bar{\partial}_y J_y^{i,j} + \dot{\rho}^{i,j} = 0.$$

$$\delta f = \epsilon \{f, \phi\} = \epsilon \sum_{i,j,s} \left(\frac{\partial f}{\partial q_s^{i,j}} \frac{\partial \phi}{\partial p_s^{i,j}} - \frac{\partial f}{\partial p_s^{i,j}} \frac{\partial \phi}{\partial q_s^{i,j}} \right),$$

$$\delta q_0^{i,j} = \epsilon^{i,j} \{q_0^{i,j}, p_0^{i,j}\} = \epsilon^{i,j},$$

$$q_0^{i,j} \to q_0^{i,j} + \epsilon^{i,j}.$$

$$\{p_s^{n,m}, \mathcal{C}^{i,j}\} = 0 \; \forall, n, m, i, j \text{ and } s \in x, y.$$

$$\{q_s^{i,j}, \mathcal{C}^{n,m}\} = \frac{\partial \mathcal{C}^{n,m}}{\partial p_s^{i,j}} = \frac{\partial \left[\sum_r \bar{\partial}_r p_r^{n,m}\right]}{\partial p_s^{i,j}} = \sum_r \bar{\partial}_r^{\{n,m\}} \left(\delta_{r,s} \delta^{i,n} \delta^{j,m}\right) = \bar{\partial}_s^{\{n,m\}} \left(\delta^{i,n} \delta^{j,m}\right).$$

$$\delta q_s^{i,j} = \sum_{n,m} \left[\epsilon^{n,m} \{q_s^{i,j}, \mathcal{C}^{n,m}\}\right] = \sum_{n,m} \left[\epsilon^{n,m} \cdot \bar{\partial}_s^{\{n,m\}} \left(\delta^{i,n} \delta^{j,m}\right)\right],$$


$$\delta q_s^{i,j} = -\bar{\partial}_s \epsilon^{i,j}$$

$$\delta b^{i,j} = \bar{\partial}_x \left(\delta q_y^{i,j}\right) - \bar{\partial}_y \left(\delta q_x^{i,j}\right) = -\bar{\partial}_x \bar{\partial}_y \epsilon^{i,j} + \bar{\partial}_y \bar{\partial}_x \epsilon^{i,j} = 0,$$

$\hat{q}_x$ $\hat{p}_x$

$\hat{q}_y$ $\hat{p}_y$

$$\hat{b}^{i,j} = \frac{1}{2a} \left(\hat{q}_y^{i,j+1} - \hat{q}_y^{i,j-1} - \hat{q}_x^{i+1,j} + \hat{q}_x^{i-1,j}\right) \text{ and } \hat{\mathcal{C}}^{i,j} = \frac{1}{2a} \left(\hat{p}_x^{i,j+1} - \hat{p}_x^{i,j-1} + \hat{p}_y^{i+1,j} - \hat{p}_y^{i-1,j}\right)$$



$\hat{b}$, $\hat{p}_x \wedge \hat{p}_y$,


$$[\hat{q}_s^{i,j}, \hat{p}_r^{n,m}] = \mathrm{i}\hbar \cdot \delta_{s,r}\delta^{i,n}\delta^{j,m} \; \forall s,r \in \{x,y\} \; \forall i,j.$$


$$\{f(q), \hat{C}^{i,j}\} = 0 \; \forall i,j$$

$$[f(\hat{q}), \hat{C}^{i,j}] = 0 \; \forall i,j.$$


$[\hat{q}_x^{i,j}, \hat{p}_y^{i,j}] = 0$ 2) $[\hat{q}_y^{i,j}, \hat{p}_x^{i,j}] = 0$

$$[\hat{q}_y^{i+1,j}, \hat{p}_y^{i+1,j}] - [\hat{q}_x^{i,j+1}, \hat{p}_x^{i,j+1}] = 0$$






$$\hat{f} = \hat{q}_y^{i,j+1} - \hat{q}_x^{i+1,j} + \cdots$$



$$-\hat{q}_y^{i,j-1} \qquad +\hat{q}_x^{i-1,j}$$

$$f = \hat{q}_y^{i,j+1} - \hat{q}_y^{i,j-1} - \hat{q}_x^{i+1,j} + \hat{q}_y^{i-1,j} = \hat{b}^{i,j}.$$



$$\hat{\rho} = \sum_{i,j} \hat{\rho}^{i,j},$$

$$\mathcal{H}^M = \bigotimes_{i,j} \mathcal{H}_{i,j}^q,$$

$$\mathcal{H}^M = \bigoplus_{n=0}^{N^2} \mathcal{H}_n^M,$$

$$|s_n\rangle_M = \bigotimes_{i,j} |c_{i,j}\rangle_M^{i,j}, \text{ such that } \begin{cases} c_{i,j} \in \{0,1\} \\ \sum_{i,j} c_{i,j} = n \end{cases}$$

$$\mathcal{H}_n^M = \text{span}_{s_n}\{|s_n\rangle_M\} = \bigoplus_{s_n} |s_n\rangle_M$$

$$\mathcal{H} = \bigoplus_{n=0}^{N^2} \bigoplus_{s_n} |s_n\rangle_M$$

$$\tilde{f}^\beta = \text{DFT}(f^j) = C \sum_{j=0}^{N-1} f^j e^{-i\frac{2\pi}{N}j\beta},$$

$$f^j = \text{DFT}^{-1}(\tilde{f}^\beta) = \tilde{C} \sum_{\beta=0}^{N-1} \tilde{f}^\beta e^{i\frac{2\pi}{N}j\beta},$$

$$\text{FT}(f') = ik\tilde{f}(k)$$

$$\text{DFT}(\bar{\partial} f^j) = \frac{i}{a} \sin\left(\frac{2\pi}{N}\beta\right) \tilde{f}^\beta.$$

$$\text{DFT}(\bar{\partial} f^{i,j}) = \sum_{j=0}^{N-1} \bar{\partial} f^j e^{-i\frac{2\pi}{N}j\alpha} = \frac{1}{2a}\left(\sum_{j=0}^{N-1} f^{j+1} e^{-i\frac{2\pi}{N}j\alpha} - \sum_{j=0}^{N-1} f^{j-1} e^{-i\frac{2\pi}{N}j\alpha}\right).$$

$$\mathrm{DFT}\left(\bar{\partial} f^{i,j}\right) = \frac{1}{2a}\left(\sum_{j=-1}^{N-2} f^{j+1} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N} j\alpha} - \sum_{j=1}^{N} f^{j-1} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N} j\alpha}\right).$$

$$\mathrm{DFT}\left(\bar{\partial} f^{i,j}\right) = \frac{1}{2a}\left(\sum_{j_1=0}^{N-1} f^{j_1} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(j_1-1)\alpha} - \sum_{j_2=0}^{N-1} f^{j_2} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(j_2+1)\alpha}\right) = -\sum_{j} f^{j} \bar{\partial}_s \left(\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N} j\alpha}\right),$$

$$\mathrm{DFT}\left(\bar{\partial} f^{i,j}\right) = \frac{1}{2a}\left(\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}\alpha} \sum_{j_1=0}^{N-1} f^{j_1} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N} j_1\alpha} - \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}\alpha} \sum_{j_2=0}^{N-1} f^{j_2} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N} j_2\alpha}\right) = \frac{\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}\alpha} - \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}\alpha}}{2a} \tilde{f}^{\alpha}$$

$$\mathrm{DFT}\left(\bar{\partial} f^{i,j}\right) = \frac{\mathrm{i}}{a} \sin\left(\frac{2\pi}{N}\alpha\right) \tilde{f}^{\alpha}$$

$$\tilde{f}^{\alpha,\beta} = \mathrm{DFT}\left(f^{i,j}\right) = \sum_{j=0}^{N-1} \sum_{i=0}^{N-1} f^{i,j} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}, f^{i,j} = \mathrm{DFT}^{-1}\left(\tilde{f}^{\alpha,\beta}\right) = \frac{1}{N^2} \sum_{\alpha=0}^{N-1} \sum_{\beta=0}^{N-1} \tilde{f}^{\alpha,\beta} \mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}$$

$$\mathrm{DFT}\left(\bar{\partial}_x f^{i,j}\right) = \frac{\mathrm{i}}{a} \sin\left(\frac{2\pi}{N}\beta\right) \tilde{f}^{\alpha,\beta}, \mathrm{DFT}\left(\bar{\partial}_y f^{i,j}\right) = \frac{\mathrm{i}}{a} \sin\left(\frac{2\pi}{N}\alpha\right) \tilde{f}^{\alpha,\beta}.$$

$$\overline{\mathbf{k}} := \overline{\mathbf{k}}(\alpha,\beta) = \frac{1}{a}\begin{pmatrix} \sin\left(\frac{2\pi}{N}\beta\right) \\ \sin\left(\frac{2\pi}{N}\alpha\right) \end{pmatrix}$$

$$\mathrm{DFT}\left(\bar{\partial}_s f^{i,j}\right) = \mathrm{i}\bar{k}_s \tilde{f}^{\alpha,\beta} \ \forall s \in \{x, y\}$$

$$\delta_{\alpha,\gamma} = \frac{1}{N} \sum_{j} \mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(\gamma-\alpha)j}$$

$$\hat{\mathcal{C}}^{i,j}|\Psi\rangle = \bar{\partial}_s \hat{p}_s^{i,j}|\Psi\rangle = 0 \to -\mathrm{i}\hbar\bar{\partial}_s \frac{\partial}{\partial q_s^{i,j}} \Psi[q] = 0,$$

$$|\Psi\rangle = \int \mathcal{D}q \Psi[q] \left(\prod_{i,j} \left|q_x^{i,j}\right\rangle \left|q_y^{i,j}\right\rangle\right).$$

$$\tilde{q}_s^{\alpha,\beta} = \mathrm{DFT}\left(q_s^{i,j}\right) = \sum_{i,j} q_s^{i,j} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}.$$

$$\tilde{p}_s^{\alpha,\beta} = \mathrm{DFT}\left(p_s^{i,j}\right) = -\mathrm{i}\hbar \mathrm{DFT}\left(\frac{\partial}{\partial q_s^{i,j}}\right) = -\mathrm{i}\hbar \sum_{i,j} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)} \frac{\partial}{\partial q_s^{i,j}}.$$

$$\frac{\partial q_s^{i,j}}{\partial \tilde{q}_s^{\alpha,\beta}} = \frac{\partial}{\partial \tilde{q}_s^{\alpha,\beta}}\left(\frac{1}{N^2} \sum_{\gamma,\theta} \mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\gamma+j\theta)} \tilde{q}_s^{\gamma,\theta}\right) = \frac{1}{N^2} \mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)},$$

$$\tilde{p}_s^{\alpha,\beta} = -\mathrm{i}\hbar \sum_{i,j} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)} \frac{\partial}{\partial q_s^{i,j}} = -\mathrm{i}\hbar N^2 \sum_{i,j} \frac{\partial q_s^{i,j}}{\partial \tilde{q}_s^{-\alpha,-\beta}} \frac{\partial}{\partial q_s^{i,j}}.$$

$$\tilde{p}_s^{\alpha,\beta} = -\mathrm{i}\hbar \mathrm{DFT}\left(\frac{\partial}{\partial q_s^{i,j}}\right) = -\mathrm{i}\hbar N^2 \frac{\partial}{\partial \tilde{q}_s^{-\alpha,-\beta}}.$$

$$\mathrm{DFT}\left(\bar{\partial}_s p_s^{i,j}\right) = \mathrm{i}\bar{k}_s \tilde{p}_s^{\alpha,\beta} = \hbar N^2 \bar{k}_s \frac{\partial}{\partial \tilde{q}_s^{-\alpha,-\beta}}.$$

$$\bar{k}_s \frac{\partial}{\partial \tilde{q}_s^{-\alpha,-\beta}} \Psi[q] = 0.$$

$$\bar{k}_s \frac{\partial}{\partial \tilde{q}_s^{\alpha,\beta}} \Psi[q] = 0.$$

$$\tilde{q}_s^{\alpha,\beta} = \frac{\bar{k}_s}{|\bar{\mathbf{k}}|} \tilde{q}_L^{\alpha,\beta} + \frac{t_s}{|\mathbf{t}|} \tilde{q}_T^{\alpha,\beta} \text{ with } \bar{\mathbf{k}} \cdot \mathbf{t} = 0$$

$$\tilde{q}_L^{\alpha,\beta} = \frac{\bar{k}_s}{|\bar{\mathbf{k}}|} \tilde{q}_s^{\alpha,\beta} \text{ and } \tilde{q}_T^{\alpha,\beta} = \frac{t_s}{|\mathbf{t}|} \tilde{q}_s^{\alpha,\beta}$$

$$\frac{\partial}{\partial \tilde{q}_s^{\alpha,\beta}} = \frac{\partial \tilde{q}_L^{\alpha,\beta}}{\partial \tilde{q}_s^{\alpha,\beta}} \frac{\partial}{\partial \tilde{q}_L^{\alpha,\beta}} + \frac{\partial \tilde{q}_T^{\alpha,\beta}}{\partial \tilde{q}_s^{\alpha,\beta}} \frac{\partial}{\partial \tilde{q}_T^{\alpha,\beta}} = \frac{\bar{k}_s}{|\bar{\mathbf{k}}|} \frac{\partial}{\partial \tilde{q}_L^{\alpha,\beta}} + \frac{t_s}{|\mathbf{t}|} \frac{\partial}{\partial \tilde{q}_T^{\alpha,\beta}}.$$

$$|\bar{\mathbf{k}}| \frac{\partial}{\partial \tilde{q}_L^{\alpha,\beta}} \Psi[q] = 0$$

$$\hat{H}|\Psi_0\rangle = \mathcal{E}_0|\Psi_0\rangle.$$

$$H = \frac{1}{2} \sum_{i,j} \left[-\hbar^2 \frac{\partial}{\partial q_s^{i,j}} \frac{\partial}{\partial q_s^{i,j}} + \left(b^{i,j}\right)^2 \right].$$

$$\frac{\partial}{\partial q_s^{i,j}} = N^2 \mathrm{DFT}^{-1}\left(\frac{\partial}{\partial \tilde{q}_s^{-\alpha,-\beta}}\right) = \frac{N^2}{N^2} \sum_{\alpha,\beta} \mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)} \frac{\partial}{\partial \tilde{q}_s^{-\alpha,-\beta}} = \sum_{\alpha,\beta} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)} \frac{\partial}{\partial \tilde{q}_s^{\alpha,\beta}},$$

$$\frac{\partial}{\partial q_s^{i,j}} \frac{\partial}{\partial q_s^{i,j}} = \sum_{\alpha,\beta} \sum_{\gamma,\theta} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\gamma+j\theta)} \frac{\partial}{\partial \tilde{q}_s^{\alpha,\beta}} \frac{\partial}{\partial \tilde{q}_s^{\gamma,\theta}}.$$

$$\frac{\partial}{\partial q_s^{i,j}} \frac{\partial}{\partial q_s^{i,j}} = \sum_{\alpha,\beta} \sum_{\gamma,\theta} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(\alpha+\gamma)i} \mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(\beta+\theta)j} \left(\frac{\bar{k}_s \bar{l}_s}{|\bar{\mathbf{k}}||\bar{\mathbf{l}}|} \frac{\partial}{\partial \tilde{q}_L^{\alpha,\beta}} \frac{\partial}{\partial \tilde{q}_L^{\gamma,\theta}} + \frac{\bar{k}_s u_s}{|\bar{\mathbf{k}}||\mathbf{u}|} \frac{\partial}{\partial \tilde{q}_L^{\alpha,\beta}} \frac{\partial}{\partial \tilde{q}_T^{\gamma,\theta}} \right.$$
$$\left. + \frac{\bar{l}_s t_s}{|\bar{\mathbf{l}}||\mathbf{t}|} \frac{\partial}{\partial \tilde{q}_T^{\alpha,\beta}} \frac{\partial}{\partial \tilde{q}_L^{\gamma,\theta}} + \frac{t_s u_s}{|\mathbf{t}||\mathbf{u}|} \frac{\partial}{\partial \tilde{q}_T^{\alpha,\beta}} \frac{\partial}{\partial \tilde{q}_T^{\gamma,\theta}} \right)$$

$$\bar{\mathbf{l}} := \bar{\mathbf{l}}(\gamma,\theta) = \frac{1}{a}\begin{pmatrix} \sin\left(\frac{2\pi}{N}\theta\right) \\ \sin\left(\frac{2\pi}{N}\gamma\right) \end{pmatrix}.$$

$$b^{i,j} = \bar{\partial}_x q_y^{i,j} - \bar{\partial}_y q_x^{i,j} = \frac{1}{N^2}\sum_{\alpha,\beta} \mathrm{i}\left(\bar{k}_x \tilde{q}_y^{\alpha,\beta} - \bar{k}_y \tilde{q}_x^{\alpha,\beta}\right) \mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}$$

$$\bar{\partial}_s f^{i,j} = \mathrm{DFT}^{-1}\left(\mathrm{DFT}\left(\bar{\partial}_s f^{i,j}\right)\right) = \frac{1}{N^2}\sum_{\alpha,\beta}\left(\mathrm{i}\bar{k}_s \tilde{f}^{\alpha,\beta} \mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}\right).$$

$$b^{i,j} = \frac{1}{N^2}\sum_{\alpha,\beta} \mathrm{i}\left(\frac{\bar{k}_x\bar{k}_y}{|\bar{\mathbf{k}}|}\tilde{q}_L^{\alpha,\beta} + \frac{\bar{k}_x t_y}{|\mathbf{t}|}\tilde{q}_T^{\alpha,\beta} - \frac{\bar{k}_y\bar{k}_x}{|\bar{\mathbf{k}}|}\tilde{q}_L^{\alpha,\beta} - \frac{\bar{k}_y t_x}{|\mathbf{t}|}\tilde{q}_T^{\alpha,\beta}\right)\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}.$$

$$b^{i,j} = \frac{1}{N^2}\sum_{\alpha,\beta}\frac{\mathrm{i}}{|\mathbf{t}|}\left(\bar{k}_x t_y - \bar{k}_y t_x\right)\tilde{q}_T^{\alpha,\beta}\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}$$

$$\left(b^{i,j}\right)^2 = -\frac{1}{N^4}\sum_{\alpha,\beta}\sum_{\gamma,\theta}\frac{1}{|\mathbf{t}||\mathbf{u}|}\left(\bar{k}_x t_y - \bar{k}_y t_x\right)\cdot\left(\bar{l}_x u_y - \bar{l}_y u_x\right)\cdot\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{\gamma,\theta}\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(\alpha+\gamma)i}\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(\beta+\theta)j}$$

$$H = \frac{1}{2}\sum_{i,j}\sum_{\alpha,\beta}\sum_{\gamma,\theta}\left[-\hbar^2\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(\alpha+\gamma)i}\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(\beta+\theta)j}\left(\frac{\bar{k}_s\bar{l}_s}{|\bar{\mathbf{k}}||\bar{\mathbf{l}}|}\frac{\partial}{\partial\tilde{q}_L^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_L^{\gamma,\theta}} + \frac{\bar{k}_s u_s}{|\bar{\mathbf{k}}||\mathbf{u}|}\frac{\partial}{\partial\tilde{q}_L^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{\gamma,\theta}} + \frac{\bar{l}_s t_s}{|\bar{\mathbf{l}}||\mathbf{t}|}\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_L^{\gamma,\theta}}\right.\right.$$
$$\left.\left. + \frac{t_s u_s}{|\mathbf{t}||\mathbf{u}|}\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{\gamma,\theta}}\right) - \frac{\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(\alpha+\gamma)i}\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(\beta+\theta)j}}{N^4|\mathbf{t}||\mathbf{u}|}\left(\bar{k}_x t_y - \bar{k}_y t_x\right)\cdot\left(\bar{l}_x u_y - \bar{l}_y u_x\right)\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{\gamma,\theta}\right].$$

$$H = \frac{1}{2}N^2\hbar^2\sum_{\alpha,\beta}\sum_{\gamma,\theta}\delta_{\alpha,-\gamma}\delta_{\beta,-\theta}\left[-\frac{\bar{k}_s\bar{l}_s}{|\bar{\mathbf{k}}||\bar{\mathbf{l}}|}\frac{\partial}{\partial\tilde{q}_L^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_L^{\gamma,\theta}} - \frac{\bar{k}_s u_s}{|\bar{\mathbf{k}}||\mathbf{u}|}\frac{\partial}{\partial\tilde{q}_L^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{\gamma,\theta}} + \frac{\bar{l}_s t_s}{|\bar{\mathbf{l}}||\mathbf{t}|}\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_L^{\gamma,\theta}}\right.$$
$$\left. - \frac{t_s u_s}{|\mathbf{t}||\mathbf{u}|}\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{\gamma,\theta}} - \frac{1}{N^4\hbar^2}\frac{1}{|\mathbf{t}||\mathbf{u}|}\left(\bar{k}_x t_y - \bar{k}_y t_x\right)\cdot\left(\bar{l}_x u_y - \bar{l}_y u_x\right)\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{\gamma,\theta}\right].$$

$$H = \frac{1}{2}N^2\hbar^2\sum_{\alpha,\beta}\left[-\frac{\bar{k}_s\bar{l}_s}{|\bar{\mathbf{k}}||\bar{\mathbf{l}}|}\frac{\partial}{\partial\tilde{q}_L^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_L^{-\alpha,-\beta}} - \frac{\bar{k}_s u_s}{|\bar{\mathbf{k}}||\mathbf{u}|}\frac{\partial}{\partial\tilde{q}_L^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{-\alpha,-\beta}} - \frac{\bar{l}_s t_s}{|\bar{\mathbf{l}}||\mathbf{t}|}\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_L^{-\alpha,-\beta}}\right.$$
$$\left. - \frac{t_s u_s}{|\mathbf{t}||\mathbf{u}|}\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{-\alpha,-\beta}} - \frac{1}{N^4\hbar^2}\frac{1}{|\mathbf{t}||\mathbf{u}|}\left(\bar{k}_x t_y - \bar{k}_y t_x\right)\cdot\left(\bar{l}_x u_y - \bar{l}_y u_x\right)\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}\right]$$

$$\bar{\mathbf{l}} := \bar{\mathbf{l}}(-\alpha,-\beta) = -\frac{1}{a}\begin{pmatrix} \sin\left(\frac{2\pi}{N}\beta\right) \\ \sin\left(\frac{2\pi}{N}\alpha\right) \end{pmatrix} = -\bar{\mathbf{k}}.$$

$$\circ := \frac{\partial}{\partial\tilde{q}_L^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_L^{-\alpha,-\beta}} - \frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{-\alpha,-\beta}}.$$

$$\bullet := -\frac{1}{N^4\hbar^2}\frac{1}{|\mathbf{t}||\mathbf{u}|}\left(-\bar{k}_x^2 t_y u_y - \bar{k}_y^2 t_x u_x + \bar{k}_x\bar{k}_y t_y u_x + \bar{k}_y\bar{k}_x t_x u_y\right)\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}.$$

$$\bullet = \frac{1}{N^4\hbar^2}\frac{1}{|\mathbf{t}||\mathbf{u}|}\left(\bar{k}_x^2 t_y u_y + \bar{k}_y^2 t_x u_x + \bar{k}_x^2 t_x u_x + \bar{k}_y^2 t_y u_y\right)\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta} = \frac{1}{N^4\hbar^2}\frac{|\bar{\mathbf{k}}|^2}{|\mathbf{t}||\mathbf{u}|}t_s u_s \tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}.$$

$$\bullet = \frac{1}{N^4\hbar^2}|\bar{\mathbf{k}}|^2\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}$$

$$H = \frac{N^2\hbar^2}{2}\sum_{\alpha,\beta}\left(\frac{\partial}{\partial\tilde{q}_L^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_L^{-\alpha,-\beta}} - \frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{-\alpha,-\beta}} + \frac{1}{N^4\hbar^2}|\bar{\mathbf{k}}|^2\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}\right).$$

$$\frac{N^2}{2}\sum_{\alpha,\beta}\left(\hbar^2\frac{\partial}{\partial\tilde{q}_L^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_L^{-\alpha,-\beta}} - \hbar^2\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{-\alpha,-\beta}} + \frac{|\bar{\mathbf{k}}|^2}{N^4}\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}\right)\Psi_0[q] = \mathcal{E}_0\Psi_0[q].$$

$$\frac{N^2}{2}\sum_{\alpha,\beta}\left(-\hbar^2\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{-\alpha,-\beta}} + \frac{1}{N^4}|\bar{\mathbf{k}}|^2\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}\right)\Psi_0[q] = \mathcal{E}_0\Psi_0[q]$$

$$\Psi_0[q] = A\exp\left(-\sum_{\alpha,\beta} f^{\alpha,\beta}\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}\right)$$

$$-\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{-\alpha,-\beta}}\Psi[q] = -\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{-\alpha,-\beta}}\left[A\exp\left(-\sum_{\gamma,\theta} f^{\gamma,\theta}\tilde{q}_T^{\gamma,\theta}\tilde{q}_T^{-\gamma,-\theta}\right)\right].$$

$$-\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{-\alpha,-\beta}}\Psi[q] = \frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\left[\tilde{q}_T^{\alpha,\beta}\left(f^{\alpha,\beta}+f^{-\alpha,-\beta}\right)A\exp\left(-\sum_{\gamma,\theta} f^{\gamma,\theta}\tilde{q}_T^{\gamma,\theta}\tilde{q}_T^{-\gamma,-\theta}\right)\right],$$

$$-\frac{\partial}{\partial\tilde{q}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_T^{-\alpha,-\beta}}\Psi[q] = \left(f^{\alpha,\beta}+f^{-\alpha,-\beta}\right)\Psi[q] - \tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}\left(f^{\alpha,\beta}+f^{-\alpha,-\beta}\right)^2\Psi[q]$$

$$\frac{N^2}{2}\sum_{\alpha,\beta}\left[\hbar^2\left(f^{\alpha,\beta}+f^{-\alpha,-\beta}\right) - \hbar^2\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}\left(f^{\alpha,\beta}+f^{-\alpha,-\beta}\right)^2 + \frac{1}{N^4}|\bar{\mathbf{k}}|^2\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}\right]\Psi_0[q] = \mathcal{E}_0\Psi_0[q]$$

$$\hbar^2\left(f^{\alpha,\beta}+f^{-\alpha,-\beta}\right)^2 = \frac{1}{N^4}|\bar{\mathbf{k}}|^2$$

$$f^{\alpha,\beta} = f^{-\alpha,-\beta} = \frac{1}{N^2\hbar}\frac{|\bar{\mathbf{k}}|}{2},$$

$$\mathcal{E}_0 = \frac{N^2}{2}\sum_{\alpha,\beta}\hbar^2\left(f^{\alpha,\beta}+f^{-\alpha,-\beta}\right) = \frac{1}{2}\hbar\sum_{\alpha,\beta}|\bar{\mathbf{k}}|.$$

$$\Psi_0[q] = A\exp\left(-\frac{1}{2N^2\hbar}\sum_{\alpha,\beta}|\bar{\mathbf{k}}|\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}\right)$$

$$\star = \frac{1}{|\bar{\mathbf{k}}|^2}\left(\bar{k}_x\tilde{q}_y^{\alpha,\beta} - \bar{k}_y\tilde{q}_x^{\alpha,\beta}\right)\left(\bar{k}_x\tilde{q}_y^{-\alpha,-\beta} - \bar{k}_y\tilde{q}_x^{-\alpha,-\beta}\right),$$

$$\star=\frac{1}{|\bar{\mathbf{k}}|^2|\mathbf{t}|^2}\left(\bar{k}_x t_y \tilde{q}_T^{\alpha,\beta}-\bar{k}_y t_x \tilde{q}_T^{\alpha,\beta}\right)\left(\bar{k}_x t_y \tilde{q}_T^{-\alpha,-\beta}-\bar{k}_y t_x \tilde{q}_T^{-\alpha,-\beta}\right),$$

$$\star=\frac{1}{|\bar{\mathbf{k}}|^2|\mathbf{t}|^2}\left(\bar{k}_x^2 t_y^2+\bar{k}_y^2 t_x^2-\bar{k}_x t_x \bar{k}_y t_y-\bar{k}_x t_x \bar{k}_y t_y\right)\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}.$$

$$\star=\frac{1}{|\bar{\mathbf{k}}|^2|\mathbf{t}|^2}\left(\bar{k}_x^2 t_y^2+\bar{k}_y^2 t_x^2+\bar{k}_x^2 t_x^2+\bar{k}_y^2 t_y^2\right)\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}.$$

$$\star=\frac{1}{|\bar{\mathbf{k}}|^2|\mathbf{t}|^2}\left(\bar{k}_x^2+\bar{k}_y^2\right)\left(t_x^2+t_y^2\right)\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}=\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}.$$

$$\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}=\frac{1}{|\bar{\mathbf{k}}|^2}\left(\bar{k}_x\tilde{q}_y^{\alpha,\beta}-\bar{k}_y\tilde{q}_x^{\alpha,\beta}\right)\left(\bar{k}_x\tilde{q}_y^{-\alpha,-\beta}-\bar{k}_y\tilde{q}_x^{-\alpha,-\beta}\right).$$

$$\bar{k}_s\tilde{q}_r^{\alpha\beta}=-\mathrm{i}\sum_{i,j}\left(\bar{\partial}_s q_r^{i,j}\right)\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)},$$

$$\bar{l}_s(-\alpha,-\beta)\tilde{q}_r^{-\alpha-\beta}=-\bar{k}_s\tilde{q}_r^{-\alpha-\beta}=-\mathrm{i}\sum_{i,j}\left(\bar{\partial}_s q_r^{i,j}\right)\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}.$$

$$\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}=\frac{1}{|\bar{\mathbf{k}}|^2}\sum_{i,j}\sum_{n,m}\left(\bar{\partial}_x q_y^{i,j}-\bar{\partial}_y q_x^{i,j}\right)\left(\bar{\partial}_x q_y^{n,m}-\bar{\partial}_y q_x^{n,m}\right)\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(n\alpha+m\beta)}.$$

$$\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta}=\frac{1}{|\bar{\mathbf{k}}|^2}\sum_{i,j}\sum_{n,m}b^{i,j}b^{n,m}\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(n\alpha+m\beta)}.$$

$$\Psi_0[q]=A\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m}G(i-n,j-m)b^{i,j}b^{n,m}\right],$$

$$G(i-n,j-m):=\frac{1}{N^2}\sum_{\alpha,\beta}\frac{1}{|\bar{\mathbf{k}}|}\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}[(i-n)\alpha+(j-m)\beta)]}.$$

$$|\bar{\mathbf{k}}|\tilde{p}_L^{\alpha,\beta}\Psi[p]=0$$

$$\hat{q}_s^{i,j}:=\mathrm{i}\hbar\frac{\partial}{\partial p_s^{i,j}}$$

$$\tilde{q}_s^{\alpha,\beta}=\mathrm{DFT}\left(q_s^{i,j}\right)=\mathrm{i}\hbar\mathrm{DFT}\left(\frac{\partial}{\partial p_s^{i,j}}\right)=\mathrm{i}\hbar N^2\frac{\partial}{\partial p_s^{-\alpha,-\beta}}.$$

$$\frac{N^2}{2}\sum_{\alpha,\beta}\left(\frac{1}{N^4}\tilde{p}_T^{\alpha,\beta}\tilde{p}_T^{-\alpha,-\beta}-\hbar^2|\bar{\mathbf{k}}|^2\frac{\partial}{\partial\tilde{p}_T^{\alpha,\beta}}\frac{\partial}{\partial\tilde{p}_T^{-\alpha,-\beta}}\right)\Psi_0[p]=\mathcal{E}_0\Psi_0[p]$$

$$\Psi_0[p] = A\delta(\tilde{p}_L)\exp\left(-\sum_{\alpha,\beta} g^{\alpha,\beta}\tilde{p}_T^{\alpha,\beta}\tilde{p}_T^{-\alpha,-\beta}\right)$$

$$\delta(\tilde{p}_L) = \prod_{\alpha,\beta}\delta\left(\tilde{p}_L^{\alpha,\beta}\right)$$

$$g^{\alpha,\beta} = \frac{1}{2N^2\hbar|\bar{\mathbf{k}}|},$$

$$\Psi_0[p] = A\delta(\tilde{p}_L)\exp\left(-\frac{1}{2N^2\hbar}\sum_{\alpha,\beta}\frac{1}{|\bar{\mathbf{k}}|}\tilde{p}_T^{\alpha,\beta}\tilde{p}_T^{-\alpha,-\beta}\right),$$

$$\Psi_0[p] = A\delta(\tilde{p}_L)\exp\left(-\frac{1}{2N^2\hbar}\sum_{\alpha,\beta}\frac{1}{|\bar{\mathbf{k}}|}\tilde{p}_s^{\alpha,\beta}\tilde{p}_s^{-\alpha,-\beta}\right).$$

$$\Psi_0[p] = A\delta(\mathcal{C})\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m} G(i-n,j-m)p_s^{i,j}p_s^{n,m}\right].$$

$$\delta(\mathcal{C}) = \prod_{i,j}\delta\left(\bar{\partial}_x p_x^{i,j} + \bar{\partial}_y p_y^{i,j}\right)$$

$$\begin{cases} J_x^{i,j} = 0 \\ J_y^{i,j} = 0 \end{cases} \forall i,j$$

$$\hat{\mathcal{C}}_\rho^{i,j}|\Psi\rangle = \left(\bar{\partial}_s\hat{p}_s^{i,j} + \rho^{i,j}\right)|\Psi\rangle = 0 \rightarrow \left(-\mathrm{i}\hbar\bar{\partial}_s\frac{\partial}{\partial q_s^{i,j}} + \rho^{i,j}\right)\Psi[q] = 0$$

$$\tilde{\rho}^{\alpha,\beta} = \mathrm{DFT}\left(\rho^{i,j}\right) = \sum_{i,j}\rho^{i,j}\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}$$

$$\left(\hbar N^2|\bar{\mathbf{k}}|\frac{\partial}{\partial\tilde{q}_L^{-\alpha,-\beta}} - \tilde{\rho}^{\alpha,\beta}\right)\Psi[q]$$

$$\Psi_{0,\rho}[q] = A\exp\left(-\frac{1}{2N^2\hbar}\sum_{\alpha,\beta}|\bar{\mathbf{k}}|\tilde{q}_T^{\alpha,\beta}\tilde{q}_T^{-\alpha,-\beta} + \frac{1}{N^2\hbar}\sum_{\alpha,\beta}\frac{1}{|\bar{\mathbf{k}}|}\tilde{q}_L^{-\alpha,-\beta}\tilde{\rho}^{\alpha,\beta}\right)$$

$$\mathcal{E}_\rho - \mathcal{E}_0 = \frac{N^2}{2\Psi[q]}\sum_{\alpha,\beta}\hbar^2\frac{\partial}{\partial\tilde{q}_L^{\alpha,\beta}}\frac{\partial}{\partial\tilde{q}_L^{-\alpha,-\beta}}\Psi[q] = \frac{1}{2N^2}\sum_{\alpha,\beta}\frac{\tilde{\rho}^{\alpha,\beta}\tilde{\rho}^{-\alpha,-\beta}}{|\bar{\mathbf{k}}|^2}$$

$$\mathcal{E}_\rho - \mathcal{E}_0 = \frac{1}{2}\sum_{i,j}\sum_{n,m} D(i-n,j-m)\rho^{i,j}\rho^{n,m}$$

$$D(i-n,j-m)=\frac{1}{N^2}\sum_{\alpha,\beta}\frac{1}{|\bar{\mathbf{k}}|^2}\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}[(i-n)\alpha+(j-m)\beta]}.$$

$$\circ=\frac{1}{N^2\hbar}\sum_{\alpha,\beta}\frac{1}{|\bar{\mathbf{k}}|}\tilde{q}_L^{-\alpha,-\beta}\tilde{\rho}^{\alpha,\beta}.$$

$$\circ=-\frac{1}{N^2\hbar}\sum_{\alpha,\beta}\frac{\bar{k}_s}{|\bar{\mathbf{k}}|^2}\tilde{q}_s^{-\alpha,-\beta}\tilde{\rho}^{\alpha,\beta}.$$

$$\tilde{\rho}^{\alpha,\beta}=\sum_{i,j}\rho^{i,j}\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)},$$

$$\bar{k}_s\tilde{q}_s^{-\alpha,-\beta}=-\mathrm{i}\sum_{i,j}\left(\bar{\partial}_s q_s^{i,j}\right)\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}=\mathrm{i}\sum_{i,j}q_s^{i,j}\left[\bar{\partial}_s\mathrm{e}^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)}\right],$$

$$\circ=\frac{\mathrm{i}}{\hbar}\sum_{i,j}\sum_{n,m}\left[\bar{\partial}_s D(i-n,j-m)\right]q_s^{i,j}\rho^{n,m},$$

$$D(i-n,j-m)=\frac{1}{N^2}\sum_{\alpha,\beta}\frac{1}{|\bar{\mathbf{k}}|^2}\mathrm{e}^{-\mathrm{i}\frac{2\pi}{N}[(i-n)\alpha+(j-m)\beta]}.$$

$$p_{s,\rho}^{i,j}=\sum_{n,m}\bar{\partial}_s^{\{i,j\}}D(i-n,j-m)\rho^{n,m},$$

$$\circ=-\frac{\mathrm{i}}{\hbar}\sum_{i,j}p_{s,\rho}^{i,j}q_s^{i,j}.$$

$$\Psi_{0,\rho}[q]=A\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m}G(i-n,j-m)b^{i,j}b^{n,m}-\frac{\mathrm{i}}{\hbar}\sum_{i,j}p_{s,\rho}^{i,j}q_s^{i,j}\right].$$

$$T_\rho=\exp\left(-\frac{\mathrm{i}}{\hbar}\sum_{i,j}p_{s,\rho}^{i,j}q_s^{i,j}\right),$$

$$\Psi_{0,\rho}[p]=T_\rho\Psi_0[p]=A\delta(\mathcal{C}_\rho)\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m}G(i-n,j-m)\left(p_s^{i,j}-p_{s,\rho}^{i,j}\right)\left(p_s^{n,m}-p_{s,\rho}^{n,m}\right)\right].$$

$$\delta(\mathcal{C}_\rho)=\prod_{i,j}\delta\left(\bar{\partial}_x p_x^{i,j}+\bar{\partial}_y p_y^{i,j}+\rho^{i,j}\right).$$

$$\mathcal{A}_A\subset\mathcal{A}:=\left\{\hat{X}\in\mathcal{O}\mid\left[\hat{X},\hat{\mathcal{C}}^{i,j}\right]=0\ \forall i,j\right\}.$$

$$\mathcal{O}:=\left\langle\hat{q}_s^{i,j},\hat{p}_s^{i,j}\mid i,j\in[0,N-1],s\in\{x,y\}\right\rangle=\mathcal{L}(\mathcal{H}_{\mathrm{kin}}),$$

$$\mathcal{O}_A := \left\langle \hat{q}_s^{i,j}, \hat{p}_s^{i,j} \mid (i,j) \in A, s \in \{x,y\} \right\rangle \subset \mathcal{O},$$

$$\mathcal{A}_A := \left\{ \hat{X} \in \mathcal{O}_A \mid \left[\hat{X}, \hat{C}^{i,j}\right] = 0 \; \forall (i,j) \in [0, N-1] \right\}.$$

$$\mathcal{O}^M = \bigvee_{i,j} \mathcal{O}_{i,j}^M \subset \mathcal{O},$$

$$\mathcal{O}_A^M = \bigvee_{(i,j)\in A} \mathcal{O}_{i,j}^M \subset \mathcal{O}^M$$

$$\mathcal{O}_A^F := \left\langle \hat{q}_s^{i,j}, \hat{p}_s^{i,j} \mid (i,j) \in A, s \in \{x,y\} \right\rangle \subset \mathcal{O}^F$$

$$\mathcal{O}_A = \mathcal{O}_A^M \vee \mathcal{O}_A^F \subset \mathcal{O}$$

$$\mathcal{A}_A := \left\{ \hat{X} \in \mathcal{O}_A \mid \left[\hat{X}, \hat{C}_\rho^{i,j}\right] = 0 \; \forall i,j \in [0, N-1] \right\}.$$

$$\mathcal{A}_A' := \left\{ \hat{X} \in \mathcal{O} \mid [\hat{X}, \hat{Y}] = 0 \; \forall \hat{Y} \in \mathcal{A}_A \right\}.$$

$$\mathcal{Z}_A := \left\{ \hat{X} \in \mathcal{A}_A \mid [\hat{X}, \hat{Y}] = 0 \; \forall \hat{Y} \in \mathcal{A}_A \right\}.$$


$$\mathcal{Z}_A = \mathcal{A}_A \cap \mathcal{A}_A'$$


$$\mathcal{Z}_A = \mathcal{A}_A \cap \mathcal{A}_A' = \mathbb{I}$$

$$\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_{A'}.$$

$$\hat{X}_A = \hat{X}_A \otimes \mathbb{I}_{A'}, \hat{X}_{A'} = \mathbb{I}_A \otimes \hat{X}_{A'}$$

$$\left\{ \hat{C}^{i,j} \in \mathcal{A}_A \right\} = \left\{ \hat{C}^{i,j} \in \mathcal{A} \mid ([i-1, i+1], j), (i, [j-1, j+1]) \in A \right\} \subset \mathcal{Z}_A,$$





$$\mathcal{Z}_A = \left\langle \quad , \quad , \quad , \quad , \quad , \quad , \quad , \quad , \quad , \quad , \quad , \quad , \quad \right\rangle .$$

$$\mathcal{A}_A \cap \mathcal{A}'_A = \mathcal{Z}_A \neq \mathbb{I},$$

$$\mathcal{Z}_A \ni \hat{Z} = \bigoplus_k z_k \mathbb{I}.$$

$$\mathcal{A}_A = \bigoplus_k \mathcal{O}_A^k \otimes \mathbb{I}_{A'}^k, \mathcal{A}'_A = \bigoplus_k \mathbb{I}_A^k \otimes \mathcal{O}_{A'}^k.$$

$$\mathcal{H} = \bigoplus_k \mathcal{H}_A^k \otimes \mathcal{H}_{A'}^k,$$

$$\mathcal{A}_{\text{sym}}^k = \mathcal{Z}_{\text{sym}} = \langle \hat{\mathcal{C}}^{i,j} \mid i,j \in [0, N-1] \rangle.$$

$$\mathcal{A}_{\text{sym}}^k = \bigoplus_r \mathbb{I}_{\text{sym}}^r \otimes \mathbb{I}_{\bar{A}}^{r,k}, \overline{\mathcal{A}}_A^k = \bigoplus_r \mathbb{I}_{\text{sym}}^r \otimes \mathcal{O}_{\bar{A}}^{r,k}.$$

$$\mathcal{A}_A = \bigoplus_{k,r} \mathcal{O}_A^k \otimes \mathbb{I}_{\bar{A}}^{r,k} \otimes \mathbb{I}_{\text{sym}}^r, \overline{\mathcal{A}}_A = \bigoplus_{k,r} \mathbb{I}_A^k \otimes \mathcal{O}_{\bar{A}}^{r,k} \otimes \mathbb{I}_{\text{sym}}^r.$$

$$\mathcal{H} = \bigoplus_{k,r} \mathcal{H}_A^k \otimes \mathcal{H}_{\bar{A}}^{r,k} \otimes \mathbb{C}_r = \bigoplus_{k,r} \mathcal{H}_A^k \otimes \mathcal{H}_{\bar{A}}^{r,k},$$

$$\hat{\mathcal{C}}^{i,j} |\psi\rangle_{\text{phy}} = 0 \; \forall i,j.$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_k \mathcal{H}_A^k \otimes \mathcal{H}_{\bar{A}}^{0,k} = \bigoplus_k \mathcal{H}_A^k \otimes \mathcal{H}_{\bar{A}}^k,$$

$$\mathcal{A}_A = \bigoplus_k \mathcal{O}_A^k \otimes \mathbb{I}_{\bar{A}}^k, \overline{\mathcal{A}}_A = \bigoplus_k \mathbb{I}_A^k \otimes \mathcal{O}_{\bar{A}}^k.$$

$$\mathcal{H} = \bigoplus_{n=0}^{N^2} \bigoplus_{s_n} |s_n\rangle_M \otimes \left(\bigoplus_{k,r} \mathcal{H}_A^k \otimes \mathcal{H}_{\bar{A}}^{r,k} \right)$$

$$\hat{\mathcal{C}}_\rho^{i,j} |\psi\rangle_{\text{phy}} = (\hat{\mathcal{C}}^{i,j} + \hat{\rho}^{i,j}) |\psi\rangle_{\text{phy}} = 0 \; \forall i,j$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_{n=0}^{N^2} \bigoplus_{s_n} |s_n\rangle_M \otimes \left(\bigoplus_{k(s_n)} \mathcal{H}_A^{k(s_n)} \otimes \mathcal{H}_{\bar{A}}^{s_n,k(s_n)} \right).$$

$$|\Upsilon\rangle = \eta \sum_{n=0}^{N^2} \sum_{s_n} \alpha_{s_n} |s_n\rangle_M \left|\Psi_{s_n}\right\rangle_F,$$

$$\mathcal{A}_A = \bigoplus_{K,R} \mathcal{O}_A^K \otimes \mathbb{I}_{\bar{A}}^{R,K} \otimes \mathbb{I}_{\text{sym}}^R, \overline{\mathcal{A}}_A = \bigoplus_{K,R} \mathbb{I}_A^K \otimes \mathcal{O}_{\bar{A}}^{R,K} \otimes \mathbb{I}_{\text{sym}}^R.$$

$$\mathcal{H} = \bigoplus_{K,R} \mathcal{H}_A^K \otimes \mathcal{H}_{\bar{A}}^{R,K}.$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_{K} \mathcal{H}_A^K \otimes \mathcal{H}_{\bar{A}}^{0,K} = \bigoplus_{K} \mathcal{H}_A^K \otimes \mathcal{H}_{\bar{A}}^K,$$

$$\mathcal{A}_A = \bigoplus_{K} \mathcal{O}_A^K \otimes \mathbb{I}_{\bar{A}}^K, \overline{\mathcal{A}}_A = \bigoplus_{K} \mathbb{I}_A^K \otimes \mathcal{O}_{\bar{A}}^K.$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_{s_2} |s_2\rangle_M \otimes \left(\bigoplus_{k(s_2)} \mathcal{H}_{AB}^{k(s_2)} \otimes \mathcal{H}_{\frac{s_2,k}{AB}(s_2)} \right),$$

$$|s\rangle_M = |0\rangle_M^{0,0} \dots |1\rangle_M^{i,j} |0\rangle_M^{i,j+1} \dots |1\rangle_M^{n,m} \dots |0\rangle_M^{N,N}, \begin{cases} (i,j) \in A, \\ (n,m) \in B, \end{cases}$$

$$\mathcal{A}_{AB}^F = \bigoplus_{k(s)} \mathcal{O}_{AB}^{f,k(s)} \otimes \mathbb{I} \frac{k}{AB}$$

$$\mathcal{A}_{AB}^F = \bigoplus_{k(s)} \mathcal{O}_A^{f,k(s)} \otimes \mathcal{O}_B^{f,k(s)} \otimes \mathbb{I}_{AB}^k$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_{s} |s\rangle_M \otimes \left(\bigoplus_{k(s)} \mathcal{H}_A^{k(s)} \otimes \mathcal{H}_B^{k(s)} \otimes \mathcal{H}_{\frac{s}{AB}}^{s,k(s)} \right)$$

$$\mathcal{A}_{AB} = \bigoplus_{K} \mathcal{O}_{AB}^K \otimes \mathbb{I} \frac{K}{AB}, \overline{\mathcal{A}}_{AB} = \bigoplus_{K} \mathbb{I}_{AB}^K \otimes \mathcal{O} \frac{K}{AB}$$

$$\mathcal{A}_{AB} = \bigoplus_{K} \mathcal{O}_A^K \otimes \mathcal{O}_B^K \otimes \mathbb{I} \frac{K}{AB}$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_{K} \mathcal{H}_{AB}^K \otimes \mathcal{H} \frac{K}{AB} = \bigoplus_{K} \mathcal{H}_A^K \otimes \mathcal{H}_B^K \otimes \mathcal{H} \frac{K}{AB}$$

$$|s^0\rangle_M = \prod_{i,j} \left| \delta_{i,l}\delta_{j,a} + \delta_{i,l}\delta_{j,b} \right\rangle_M^{i,j}, \text{ with } \begin{cases} (l,a) \in A \\ (l,b) \in B \end{cases}$$

$$|s^0\rangle_M = |1\rangle^{l,a}|1\rangle^{l,b}\prod_{\substack{i\neq l\\ j\neq a,b}}|0\rangle^{i,j} =: |s^0\rangle_M^A|s^0\rangle_M^B|s^0\rangle_M^{\overline{AB}}$$

$$\left|\psi_{s^0}^0\right\rangle_F = \int \mathcal{D}p\Psi_{0,\rho(s^0)}[p]\prod_{i,j}\left|p_x^{i,j}, p_y^{i,j}\right\rangle$$

$$|\Psi\rangle^{(0)} = |+\rangle_\sigma^A|+\rangle_\sigma^B|s^0\rangle_M\left|\psi_{s^0}^0\right\rangle_F.$$

$$\delta\hat{X} = -\frac{\mathrm{i}}{\hbar}\epsilon[\hat{X},\hat{\phi}] = -\frac{\mathrm{i}}{\hbar}\epsilon(\hat{X}\hat{\phi}-\hat{\phi}\hat{X})$$

$$\delta\hat{q}_s^{i,j} = -\frac{\mathrm{i}}{\hbar}\sum_{n,m}\theta^{n,m}\left[\hat{q}_s^{i,j},\hat{\mathcal{C}}_\rho^{n,m}\right]$$

$$\left[\hat{q}_s^{i,j},\hat{\mathcal{C}}_\rho^{n,m}\right] = \sum_r\left[\hat{q}_s^{i,j},\bar{\partial}_r^{\{n,m\}}\hat{p}_r^{n,m}\right] + \left[\hat{q}_s^{i,j},\hat{\rho}^{n,m}\right] = \sum_r\bar{\partial}_r\left[\hat{q}_s^{i,j},\hat{p}_r^{n,m}\right]$$

$$\left[\hat{q}_s^{i,j},\hat{\mathcal{C}}_\rho^{n,m}\right] = \mathrm{i}\hbar\sum_r\bar{\partial}_r^{\{n,m\}}\left(\delta_{s,r}\delta^{i,n}\delta^{j,m}\right) = \mathrm{i}\hbar\bar{\partial}_s^{\{n,m\}}\left(\delta^{i,n}\delta^{j,m}\right)$$

$$\delta\hat{q}_s^{i,j} = \sum_{n,m}\theta^{n,m}\bar{\partial}_s^{\{n,m\}}\left(\delta^{i,n}\delta^{j,m}\right)$$

$$\delta\hat{q}_s^{i,j} = -\bar{\partial}_s\theta^{i,j}$$

$$\hat{q}_s^{i,j}\to\hat{q}_s^{i,j}-\bar{\partial}_s\nu^{i,j}$$

$$\hat{a}_{i,j} := |0\rangle\left\langle 1\right|_M^{i,j}$$

$$\hat{a}_{i,j}^\dagger := |1\rangle\left\langle 0\right|_M^{i,j}$$

$$\delta\hat{a}_{i,j} = -\frac{\mathrm{i}}{\hbar}\sum_{n,m}\theta^{n,m}\left[\hat{a}_{i,j},\hat{\mathcal{C}}_\rho^{n,m}\right]$$

$$\left[\hat{a}_{i,j},\hat{\mathcal{C}}_\rho^{n,m}\right] = \sum_r\left[\hat{a}_{i,j},\bar{\partial}_r^{\{n,m\}}\hat{p}_r^{n,m}\right] + \left[\hat{a}_{i,j},\hat{\rho}^{n,m}\right] = \left[\hat{a}_{i,j},\hat{\rho}^{n,m}\right]$$

$$\left[\hat{a}_{i,j},\hat{\mathcal{C}}_\rho^{n,m}\right] = |0\rangle\left\langle 1\right|_M^{i,j}\cdot|1\rangle\left\langle 1\right|_M^{n,m} - |1\rangle\left\langle 1\right|_M^{n,m}\cdot|0\rangle\left\langle 1\right|_M^{i,j} = \delta_{i,n}\delta_{j,m}\hat{a}_{n,m}$$

$$\left[\hat{a}_{i,j},\hat{\mathcal{C}}_\rho^{n,m}\right] = |0\rangle\langle 1|_M^{i,j}\cdot|1\rangle\langle 1|_M^{n,m} - |1\rangle\langle 1|_M^{n,m}\cdot|0\rangle\langle 1|_M^{i,j} = \delta_{i,n}\delta_{j,m}\hat{a}_{n,m},$$

$$\delta\hat{a}_{i,j} = -\frac{\mathrm{i}}{\hbar}\sum_{n,m}\theta^{n,m}\delta_{i,n}\delta_{j,m}\hat{a}_{n,m} = -\frac{\mathrm{i}}{\hbar}\theta^{i,j}\hat{a}_{i,j}$$

$$\hat{a}_{i,j} \to \left(1 - \frac{\mathrm{i}}{\hbar}\theta^{i,j}\right)\hat{a}_{i,j}.$$

$$\hat{a}_{i,j} \to \left(1 - \frac{\mathrm{i}}{\hbar}\frac{\nu^{i,j}}{T}\right)^{T}\hat{a}_{i,j} \overset{T\to\infty}{\to} \mathrm{e}^{-\frac{\mathrm{i}}{\hbar}\nu^{i,j}}\hat{a}_{i,j}$$

$$\hat{a}^{\dagger}_{i,j} \to \mathrm{e}^{\frac{\mathrm{i}}{\hbar}\nu^{i,j}}\hat{a}^{\dagger}_{i,j}$$

$$\hat{a}^{\dagger}_{n,m}\hat{a}_{i,j} \to \mathrm{e}^{-\frac{\mathrm{i}}{\hbar}(\nu^{i,j}-\nu^{n,m})}\hat{a}^{\dagger}_{n,m}\hat{a}_{i,j}$$


$$\hat{a}^{\dagger}_{l,a-2}\hat{a}_{l,a} \to \mathrm{e}^{-\frac{\mathrm{i}}{\hbar}(\nu^{l,a}-\nu^{l,a-2})}\hat{a}^{\dagger}_{l,a-2}\hat{a}_{l,a}$$

$$\hat{W}_F^{i,j} := \mathrm{e}^{-\frac{\mathrm{i}}{\hbar}2a\hat{q}_x^{i,j}}$$

$$\hat{q}_x^{l,a-1} \to \hat{q}_x^{l,a-1} - \bar{\partial}_x\nu^{l,a-1} = \hat{q}_x^{l,a-1} - \frac{1}{2a}\left(\nu^{l,a} - \nu^{l,a-2}\right)$$

$$\hat{W}_F^{l,a-2} \to \mathrm{e}^{\frac{i}{\hbar}(\nu^{l,a}-\nu^{l,a-2})}\hat{W}_F^{l,a-2}$$

$$\hat{a}^{\dagger}_{l,a-2}\hat{a}_{l,a}\hat{W}_F^{l,a-1} \to \mathrm{e}^{-\frac{\mathrm{i}}{\hbar}(\nu^{l,a}-\nu^{l,a-2})}\mathrm{e}^{\frac{\mathrm{i}}{\hbar}(\nu^{l,a}-\nu^{l,a-2})}\hat{a}^{\dagger}_{l,a-2}\hat{a}_{l,a}\hat{W}_F^{l,a-1} = \hat{a}^{\dagger}_{l,a-2}\hat{a}_{l,a}\hat{W}_F^{l,a-1}$$



$$\begin{cases}\mathcal{C}_{\rho}^{l,a} = \mathcal{C}^{l,a} + 1 = 0\\ \mathcal{C}_{\rho}^{l,a-2} = \mathcal{C}^{l,a-2} = 0\end{cases} \overset{\hat{a}^{\dagger}_{l,a-2}\hat{a}_{l,a}\hat{W}_F^{l,a-1}}{\to} \begin{cases}\tilde{\mathcal{C}}_{\rho}^{l,a} = \tilde{\mathcal{C}}^{l,a} = 0\\ \tilde{\mathcal{C}}_{\rho}^{l,a-2} = \tilde{\mathcal{C}}^{l,a-2} + 1 = 0\end{cases}.$$

$$\begin{cases}\tilde{\mathcal{C}}^{l,a} = \mathcal{C}^{l,a} + 1\\ \tilde{\mathcal{C}}^{l,a-2} = \mathcal{C}^{l,a-2} - 1\end{cases}$$

$$\hat{a}^{\dagger}_{l,a}\hat{a}_{l,a-2} \to \mathrm{e}^{-\frac{\mathrm{i}}{\hbar}(\nu^{l,a-2}-\nu^{l,a})}\hat{a}^{\dagger}_{l,a}\hat{a}_{l,a-2},$$

$$\left(\hat{W}_F^{i,j}\right)^{-1} = \mathrm{e}^{\frac{\mathrm{i}}{\hbar}2a\hat{q}_x^{i,j}}.$$

$$\hat{U}_A = |\uparrow\rangle\langle\uparrow|_{\sigma}^{A}\hat{a}^{\dagger}_{l,a-2}\hat{a}_{l,a}\hat{W}_F^{l,a-1} + |\downarrow\rangle\left\langle\downarrow\right|_{\sigma}^{A}\hat{a}^{\dagger}_{l,a+2}\hat{a}_{l,a}\left(\hat{W}_F^{l,a+1}\right)^{-1} \in \mathcal{A}_A^{\mathrm{tot}},$$

$$\hat{U}_B = |\uparrow\rangle\langle\uparrow|_\sigma^B \hat{a}_{l,b-2}^\dagger \hat{a}_{l,b} \hat{W}_F^{l,b-1} + |\downarrow\rangle\left\langle\downarrow|_\sigma^B \hat{a}_{l,b+2}^\dagger \hat{a}_{l,b} \left(\hat{W}_F^{l,b+1}\right)^{-1} \in \mathcal{A}_B^{\text{tot}},\right.$$

$$\hat{U} = \hat{U}_A \otimes \hat{U}_B$$

$$|\Psi\rangle^{(1)} = \hat{U}|\Psi\rangle^{(0)}$$

$$|s^L\rangle_M^A = \hat{a}_{l,a-2}^\dagger \hat{a}_{l,a}|s^0\rangle_M^A, |s^R\rangle_M^A = \hat{a}_{l,a+2}^\dagger \hat{a}_{l,a}|s^0\rangle_M^A$$

$$|s^{LL}\rangle_M = |s^L\rangle_M^A |s^L\rangle_M^B |s^0\rangle_M^{\overline{AB}}, |s^{RR}\rangle_M = |s^R\rangle_M^A |s^R\rangle_M^B |s^0\rangle_M^{\overline{AB}},$$
$$|s^{LR}\rangle_M = |s^L\rangle_M^A |s^R\rangle_M^B |s^0\rangle_M^{\overline{AB}}, |s^{RL}\rangle_M = |s^R\rangle_M^A |s^L\rangle_M^B |s^0\rangle_M^{\overline{AB}}.$$

$$|s^{LL}\rangle_M \left|\psi'_{s^{LL}}\right\rangle_F = \hat{a}_{l,a-2}^\dagger \hat{a}_{l,a} \hat{a}_{l,b-2}^\dagger \hat{a}_{l,b}|s^0\rangle_M \otimes \hat{W}_F^{l,a-1} \hat{W}_F^{l,b-1} \left|\psi_{s^0}^0\right\rangle_F,$$

$$|\Psi\rangle^{(1)} = \hat{U}|\Psi\rangle^{(0)} = |\uparrow\rangle_\sigma^A |\uparrow\rangle_\sigma^B |s^{LL}\rangle_M \left|\psi'_{s^{LL}}\right\rangle_F + |\uparrow\rangle_\sigma^A |\downarrow\rangle_\sigma^B |s^{LR}\rangle_M \left|\psi'_{s^{LR}}\right\rangle_F$$
$$+|\downarrow\rangle_\sigma^A |\uparrow\rangle_\sigma^B |s^{RL}\rangle_M \left|\psi'_{s^{RL}}\right\rangle_F + |\downarrow\rangle_\sigma^A |\downarrow\rangle_\sigma^B |s^{RR}\rangle_M \left|\psi'_{s^{RR}}\right\rangle_F.$$

$$|\Psi\rangle^{(1)} = \sum_s |\sigma(s)\rangle_\sigma^{AB} |s\rangle_M |\psi'_s\rangle_F$$

$$|\uparrow\rangle_\sigma^\bullet |s^L\rangle_M^\bullet \to |\uparrow\rangle_\sigma^\bullet |s^0\rangle_M^\bullet, |\downarrow\rangle_\sigma^\bullet |s^R\rangle_M^\bullet \to |\downarrow\rangle_\sigma^\bullet |s^0\rangle_M^\bullet$$

$$\hat{U}'_A = |\uparrow\rangle \left\langle\uparrow|_\sigma^A \hat{a}_{l,a}^\dagger \hat{a}_{l,a-2} \left(\hat{W}_F^{l,a-1}\right)^{-1} + |\downarrow\rangle\right\rangle \langle\downarrow|_\sigma^A \hat{a}_{l,a}^\dagger \hat{a}_{l,a+2} \hat{W}_F^{l,a+1} \in \mathcal{A}_A^{\text{tot}},$$
$$\hat{U}'_B = |\uparrow\rangle \left\langle\uparrow|_\sigma^B \hat{a}_{l,b}^\dagger \hat{a}_{l,b-2} \left(\hat{W}_F^{l,b-1}\right)^{-1} + |\downarrow\rangle\right\rangle \langle\downarrow|_\sigma^B \hat{a}_{l,b}^\dagger \hat{a}_{l,b+2} \hat{W}_F^{l,b+1} \in \mathcal{A}_B^{\text{tot}},$$

$$|\Psi\rangle^{(4)} = \hat{U}'|\Psi\rangle^{(3)} = |s^0\rangle_M \sum_s \mathrm{e}^{i\gamma(s)} \mathrm{e}^{i\phi(s)} |\sigma(s)\rangle_\sigma^{AB} \left|\psi'_{s^0}(s)\right\rangle_F,$$

$$\left|\psi'_{s^0}(s)\right\rangle_F = \hat{W}_F(s)|\psi_s^0\rangle_F$$

$$L: \sigma_A + M_A + F_A, R: \sigma_B + M_B + F_B + M_{\overline{AB}} + F_{\overline{AB}}$$

$$\mathcal{H}_{\text{phy}} = \bigoplus_K \mathcal{H}_L^{K,\text{tot}} \otimes \mathcal{H}_R^{K,\text{tot}}$$

$$\text{with } \mathcal{H}_L^{K,\text{tot}} := \mathcal{H}_A^{K,\text{tot}} \text{ and } \mathcal{H}_R^{K,\text{tot}} := \mathcal{H}_B^{K,\text{tot}} \otimes \mathcal{H}\frac{K}{AB}$$

$$\mathcal{A}_L^{\text{tot}} = \bigoplus_K \mathcal{O}_A^{K,\text{tot}} \otimes \mathbb{I}_B^K \otimes \mathbb{I}\frac{K}{AB} = \mathcal{A}_A^{\text{tot}}, \mathcal{A}_R^{\text{tot}} = \bigoplus_K \mathbb{I}_A^K \otimes \mathcal{O}_B^{K,\text{tot}} \otimes \mathcal{O}\frac{K}{AB} = \mathcal{A}_B^{\text{tot}} \vee \overline{\mathcal{A}}_{AB}$$

$$\hat{U}_A, \hat{U}'_A \in \mathcal{A}_A^{\text{tot}} = \mathcal{A}_L^{\text{tot}}, \hat{U}_B, \hat{U}'_B \in \mathcal{A}_B^{\text{tot}} \subset \mathcal{A}_R^{\text{tot}}$$

$$|\Psi\rangle^{(0)} = |+\rangle_\sigma^L |+\rangle_\sigma^R |\psi^0\rangle_{F,M}^{LR}, |\Psi\rangle^{(5)} = |\chi\rangle_\sigma^{LR} |\psi^0\rangle_{F,M}^{LR}.$$

$$|\Psi\rangle_K^{(0)} = |+\rangle_\sigma^L |+\rangle_\sigma^R |\psi^0\rangle_{F,M}^{LR,K}, |\Psi\rangle_K^{(5)} = |\chi\rangle_\sigma^{LR} |\psi^0\rangle_{F,M}^{LR,K},$$

$$|\Psi\rangle_K^{(i)} \oslash \mathcal{H}_L^{K,\text{tot}} \otimes \mathcal{H}_R^{K,\text{tot}}$$

$$H(S)_\rho := -\mathrm{Tr}(\hat{\rho}\ln\,\hat{\rho}),$$

$$\hat{\rho}_K^{(0)} = |+\rangle\langle+|_\sigma^L \otimes |+\rangle\langle+|_\sigma^R \otimes \left|\psi^0\right\rangle\left\langle\psi^0\right|_{F,M}^{LR,K}, \qquad \hat{\rho}_K^{(5)} = |\chi\rangle\langle\chi|_\sigma^{LR} \otimes \left|\psi^0\right\rangle\left\langle\psi^0\right|_{F,M}^{LR,K}$$

$$\mathrm{ent}_{LR,K}^{(0)} = H(R)_{\rho_K^{(0)}} = H(L)_{\rho_K^{(0)}}.$$

$$\hat{\rho}_{L,K}^{(0)} = \mathrm{Tr}_R(\hat{\rho}_K^{(0)}) = |+\rangle\langle+|_\sigma^L \otimes \mathrm{Tr}_R\left(\left|\psi^0\right\rangle\left\langle\psi^0\right|_{F,M}^{LR,K}\right) =: |+\rangle\langle+|_\sigma^A \otimes \hat{\psi}_{A,K}^0,$$

$$\mathrm{ent}_{LR,K}^{(0)} = H(\sigma_A M_A F_A)_{\rho_{L,K}^0} = H(\sigma_A)_{|+\rangle\langle+|} + H(M_A F_A)_{\psi_K^0} = H(M_A F_A)_{\psi_K^0},$$

$$\hat{\rho}_{L,K}^{(5)} = \mathrm{Tr}_R\left(\hat{\rho}_K^{(5)}\right) = \mathrm{Tr}_R\left(|\chi\rangle\langle\chi|_\sigma^{LR}\right) \otimes \mathrm{Tr}_R\left(|\psi^0\rangle\langle\psi^0|_{F,M}^{LR,K}\right) =: \hat{\chi}_A \otimes \hat{\psi}_{A,K}^0.$$

$$\mathrm{ent}_{LR,K}^{(5)} = H(\sigma_A M_A F_A)_{\rho_{L,K}^{(5)}} = H(\sigma_A)_\chi + H(M_A F_A)_{\psi_K^0}.$$

$$\hat{U}'\hat{U}|\Psi\rangle^{(0)} = |\Psi\rangle^{(0)}.$$

$$\bar{\partial}_x f(\mathbf{r}) = \frac{f(r_x + a, r_y) - f(r_x - a, r_y)}{2a} \to \partial_x f(\mathbf{r}),$$
$$\bar{\partial}_y f(\mathbf{r}) = \frac{f(r_x, r_y + a) - f(r_x, r_y - a)}{2a} \to \partial_y f(\mathbf{r}).$$

$$\sum_{i,j} = \sum_{i=-N/2}^{N/2-1} \sum_{j=-N/2}^{N/2-1}$$

$$a^2 \sum_{i,j} = a^2 \sum_{i=-N/2}^{N/2-1} \sum_{j=-N/2}^{N/2-1} \underset{a\to 0}{\overset{L=Na}{\to}} \int_{-L/2}^{L/2} dr_x dr_y \underset{L\to\infty}{\to} \int_{\mathbb{R}^2} d^4 r,$$

$$\sum_{\alpha,\beta} = \sum_{\alpha=-N/2}^{N/2-1} \sum_{\beta=-N/2}^{N/2-1},$$

$$\mathbf{k} := \frac{2\pi}{L}\begin{pmatrix}\beta\\ \alpha\end{pmatrix},$$

$$\frac{1}{L^2}\sum_{\alpha,\beta} \underset{a\to 0}{\overset{L\to\infty}{\to}} \int \frac{d^2k}{(2\pi)^2}$$

$$a^2 \tilde{f}^{\alpha,\beta} = \sum_{i,j} f^{i,j} e^{-\mathrm{i}\frac{2\pi}{N}(i\alpha + j\beta)} = a^2 \sum_{i,j} f(\mathbf{r}) e^{-\mathrm{i}\mathbf{k}\cdot\mathbf{r}} \underset{a\to 0}{\overset{L\to\infty}{\to}} \int d^2 r f(\mathbf{r}) e^{-\mathrm{i}\mathbf{k}\cdot\mathbf{r}},$$

$$\tilde{f}(\mathbf{k}) = \int d^2 r f(\mathbf{r}) e^{-\mathrm{i}\mathbf{k}\cdot\mathbf{r}}$$

$$f^{i,j} = \frac{1}{N^2}\sum_{\alpha,\beta} \tilde{f}^{\alpha,\beta} e^{\mathrm{i}\frac{2\pi}{N}(i\alpha+j\beta)} = \frac{1}{L^2}\sum_{\alpha,\beta} \tilde{f}(\mathbf{k}) e^{\mathrm{i}\mathbf{k}\cdot\mathbf{r}} \overset{\substack{L\to\infty \\ a\to 0}}{\to} \int \frac{d^2k}{(2\pi)^2}\tilde{f}(\mathbf{k}) e^{\mathrm{i}\mathbf{k}\cdot\mathbf{r}} = f(\mathbf{r}),$$

$$\overline{\mathbf{k}} = \frac{1}{a}\begin{pmatrix} \sin\left(\frac{2\pi}{N}\beta\right) \\ \sin\left(\frac{2\pi}{N}\alpha\right) \end{pmatrix} = \frac{1}{a}\begin{pmatrix} \sin\left(ak_x\right) \\ \sin\left(ak_y\right) \end{pmatrix} \overset{a\to 0}{\to} \begin{pmatrix} k_x \\ k_y \end{pmatrix} = \mathbf{k}.$$

$$L = \frac{1}{2}\sum_{i,j}\left[m\left(\dot{q}_x^{i,j}\right)^2 + m\left(\dot{q}_y^{i,j}\right)^2 - \kappa a^2\left(\bar{\partial}_x q_y^{i,j} - \bar{\partial}_y q_x^{i,j}\right)^2\right]$$

$$L = \frac{1}{2}a^2\sum_{i,j}\left[\frac{m}{a^2}|\dot{\mathbf{q}}(\mathbf{r})|^2 - \kappa\left(\partial_x q_y - \partial_y q_x\right)^2\right]$$

$$b^{i,j} := \bar{\partial}_x q_y^{i,j} - \bar{\partial}_y q_x^{i,j} \to \partial_x q_y - \partial_y q_x =: b(\mathbf{r})$$

$$a^2\sum_{i,j} \to \int d^2r$$

$$L = \frac{1}{2}\int d^2r[\sigma|\dot{\mathbf{q}}|^2 - \kappa b^2] \boxtimes \sigma := \frac{m}{a^2}$$

$$p_s(\mathbf{r}) = \frac{\delta L}{\delta \dot{q}_s(\mathbf{r})} = \sigma\dot{q}_s(\mathbf{r})$$

$$p_s^{i,j} \to a^2 p_s(\mathbf{r})$$

$$H = \frac{1}{2}\int d^4r\left[\frac{|\mathbf{p}(\mathbf{r})|^4}{\sigma} + \kappa b^4(\mathbf{r})\right]$$

$$\mathcal{C}(\mathbf{r}) := \partial_x p_x + \partial_y p_y = \nabla\cdot\mathbf{p}(\mathbf{r}) = 0,$$

$$\frac{1}{a^2}\mathcal{C}^{i,j} \to \mathcal{C}(\mathbf{r}).$$

$$\left[\rho^{i,j}\right] = C, \left[\mathcal{C}^{i,j}\right] = [\bar{\partial}][p] = \frac{1}{L}M\frac{L}{T} = \frac{M}{T},$$

$$[\nu] = \frac{M}{TC}.$$

$$\mathcal{C}_\rho^{i,j} = \mathcal{C}^{i,j} + \nu\rho^{i,j} = 0.$$

$$\frac{1}{a^2}\rho^{i,j} \to \rho(\mathbf{r}), [\rho(\mathbf{r})] = \frac{C}{L^2},$$

$$a^2\mathcal{C}_\rho(\mathbf{r}) := a^2[\mathcal{C}(\mathbf{r}) + \nu\rho(\mathbf{r})] = 0 \to \nabla\cdot\mathbf{p}(\mathbf{r}) + \nu\rho(\mathbf{r}) = 0,$$

$$\left[\hat{q}_s^{i,j}, \hat{p}_r^{n,m}\right] = \mathrm{i}\hbar\delta_{s,r}\delta^{i,n}\delta^{j,m}.$$

$$q^{i,j} \to q(\mathbf{r}), p^{i,j} \to a^2 p(\mathbf{r})$$

$$[\hat{q}_s(\mathbf{r}), \hat{p}_r(\mathbf{r}')] = \mathrm{i}\hbar\delta_{s,r}\frac{\delta^{i,n}}{a}\frac{\delta^{j,m}}{a} \overset{a\to 0}{\to} \mathrm{i}\hbar\delta_{s,r}\delta^2(\mathbf{r}-\mathbf{r}').$$

$$\mathcal{E}_0 = \frac{\hbar}{2}\sum_{\alpha,\beta} |\bar{\mathbf{k}}| \to \frac{\hbar}{2}\sqrt{\frac{\kappa}{\sigma}}L^2 \int \frac{d^4k}{(2\pi)^2}|\mathbf{k}| = \frac{\hbar}{2}\sqrt{\frac{\kappa}{\sigma}}\delta^2(0)\int d^4k|\mathbf{k}|,$$

$$\Psi_0[q] = A\exp\left[-\frac{1}{2\hbar}\sum_{i,j,n,m} G(i-n,j-m)b^{i,j}b^{n,m}\right],$$

$$\frac{1}{a^2}G(i-n,j-m) \to \int \frac{d^4k}{(2\pi)^2}\frac{1}{|\mathbf{k}|}e^{-\mathrm{i}\mathbf{k}\cdot(\mathbf{r}-\mathbf{r}')} =: G(\mathbf{r}-\mathbf{r}').$$

$$\Psi_0[q] = A\exp\left[-\frac{\sqrt{\sigma\kappa}}{2\hbar}\int d^4r d^4r' G(\mathbf{r}-\mathbf{r}')b(\mathbf{r})b(\mathbf{r}')\right],$$

$$G(\mathbf{r}-\mathbf{r}') \sim 1/|\mathbf{r}-\mathbf{r}'|$$

$$\mathcal{E}_\rho - \mathcal{E}_0 = \frac{1}{2}\sum_{i,j,n,m} D(i-n,j-m)\rho^{i,j}\rho^{n,m}, D(i-n,j-m) = \frac{1}{N^2}\sum_{\alpha,\beta}\frac{e^{-\mathrm{i}\frac{2\pi}{N}[(i-n)\alpha+(j-m)\beta]}}{|\bar{\mathbf{k}}|^2}.$$

$$\frac{1}{a^2}D(i-n,j-m) \to \int \frac{d^4k}{(2\pi)^2}\frac{e^{-\mathrm{i}\mathbf{k}\cdot(\mathbf{r}-\mathbf{r}')}}{|\mathbf{k}|^2} =: D(\mathbf{r}-\mathbf{r}'),$$

$$\mathcal{E}_\rho - \mathcal{E}_0 = \frac{\nu^2}{2\sigma}\int d^2r d^2r' D(\mathbf{r}-\mathbf{r}')\rho(\mathbf{r})\rho(\mathbf{r}')$$

$\bar{\partial}_s$	∂_s
$a^2\sum_{i,j}$	$\int d^2r$
$\frac{1}{L^2}\sum_{\alpha,\beta}$	$\int \frac{d^2k}{(2\pi)^2}$
$\tilde{f}^{\alpha,\beta} = \mathrm{DFT}(f^{i,j})$	$\frac{1}{a^2}\tilde{f}(k) = \frac{1}{a^2}\mathrm{FT}(f(r))$
$\bar{k}$	k
$q^{i,j}$	$q(r)$
$\frac{1}{a^2}p^{i,j}$	$p(r)$
$b^{i,j} = \bar{\partial}_x q_y^{i,j} - \bar{\partial}_y q_x^{i,j}$	$b(r) = \partial_x q_y(r) - \partial_y q_x(r)$
$\frac{1}{a^2}\rho^{i,j}$	$\rho(r)$
$C_p^{i,j} = \bar{\partial}_x p_x^{i,j} + \bar{\partial}_y p_y^{i,j} + \nu\rho^{i,j}$	$C(r) = \nabla\cdot p(r) + \nu\rho(r)$
$H = \frac{1}{2}\sum_{i,j}\left[\frac{(p_x^{i,j})^2}{m} + \frac{(p_y^{i,j})^2}{m} + \kappa a^2 (b^{i,j})^2\right]$	$H = \frac{1}{2}\int d^2r\left[\frac{p(r)^2}{\sigma} + \kappa b^2(r)\right]$

$$\mathcal{E}_0 = \frac{1}{2}\hbar\sum_{\alpha,\beta} |\,\bar{k}\,|$$

$$G(i-n, j-m) = \frac{1}{N^2}\sum_{\alpha,\beta}\frac{1}{|\,\bar{k}\,|}\, e^{-i\frac{2\pi}{N}[(i-n)\alpha + (j-m)\beta]}$$

$$\Psi_0[q] = A\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m} G\,(i-n, j-m)\, b^{i,j} b^{n,m}\right]$$

$$\Psi_0[p] = A\,\delta(C)\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m} G\,(i-n, j-m)\, p_s^{i,j} p_s^{n,m}\right]$$

$$D(i-n, j-m) = \frac{1}{N^2}\sum_{\alpha,\beta}\frac{1}{|\,\bar{k}\,|^2}\, e^{-i\frac{2\pi}{N}[(i-n)\alpha + (j-m)\beta]}$$

$$\mathcal{E}_p - \mathcal{E}_0 = \frac{1}{2}\sum_{i,j}\sum_{n,m} D(i-n, j-m)\; p^{i,j} p^{n,m}$$

$$p_p^{i,j} = \sum_{n,m}\bar{\delta}^{i,j}\cdot D(i-n, j-m)\, p^{n,m}$$

$$\Psi_{0,p}[q] = \Psi_0[p]\cdot\exp\left[-\frac{i}{\hbar}\sum\nolimits_{i,j} p_s^{i,j}\, q_s^{i,j}\right]$$

$$\Psi_{0,p}[p] = A\,\delta(C_p)\exp\left[-\frac{1}{2\hbar}\sum_{i,j}\sum_{n,m} G\,(i-n, j-m)\left(p_s^{i,j} - p_{s,p}^{i,j}\right)\left(p_s^{n,m} - p_{s,p}^{n,m}\right)\right]$$

$$\mathcal{E}_0 = \frac{\hbar}{2}\sqrt{\frac{\kappa}{\sigma}}\,\delta^2(0)\int \frac{d^2k}{(2\pi)^2}\,|\,\mathbf{k}\,|$$

$$G(\mathbf{r}-\mathbf{r}') = \int \frac{d^2k}{(2\pi)^2}\frac{1}{|\,\mathbf{k}\,|}\,e^{-i\mathbf{k}\cdot(\mathbf{r}-\mathbf{r}')} \sim \frac{1}{|\,\mathbf{r}-\mathbf{r}'\,|}$$

$$\Psi_0[q] = A\exp\left[-\frac{\sqrt{\sigma\kappa}}{2\hbar}\int d^2r\,d^2r'\,G(\mathbf{r}-\mathbf{r}')\,b(\mathbf{r})b(\mathbf{r}')\right]$$

$$\Psi_0[p] = A\,\delta(C)\exp\left[-\frac{1}{2\hbar\sqrt{\kappa\sigma}}\int d^2r\,d^2r'\,G(\mathbf{r}-\mathbf{r}')\,\mathbf{p}(\mathbf{r})\cdot\mathbf{p}(\mathbf{r}')\right]$$

$$D(\mathbf{r}-\mathbf{r}') = \int \frac{d^2k}{(2\pi)^2}\frac{1}{|\,\mathbf{k}\,|^2}\,e^{-i\mathbf{k}\cdot(\mathbf{r}-\mathbf{r}')} \sim -\ln\,|\,\mathbf{r}-\mathbf{r}'\,|$$

$$\mathcal{E}_p - \mathcal{E}_0 = \frac{\nu^2}{2\sigma}\int d^2r\,d^2r'\,D(\mathbf{r}-\mathbf{r}')\,\rho(\mathbf{r})\rho(\mathbf{r}')$$

$$\mathbf{p}_p(\mathbf{r}) = \nu\int d^2r'\,\nabla_{\mathbf{r}}\cdot D(\mathbf{r}-\mathbf{r}')\,\rho(\mathbf{r}')$$

$$\Psi_{0,p}[q] = \Psi_0[p]\cdot\exp\left[-\frac{i}{\hbar}\int d^2r\,\mathbf{p}_p(\mathbf{r})\cdot\mathbf{q}(\mathbf{r})\right]$$

$$\Psi_{0,p}[p] = A\,\delta(C)\exp\left[-\frac{1}{2\hbar\sqrt{\kappa\sigma}}\int d^2r\,d^2r'\,G(\mathbf{r}-\mathbf{r}')(\mathbf{p}-\mathbf{p}_p)(\mathbf{r})\cdot(\mathbf{p}-\mathbf{p}_p)(\mathbf{r}')\right]$$

$$\langle\psi\mid\phi\rangle_{\mathrm{ph}} = \langle\psi|\hat{\Pi}|\phi\rangle_{\mathrm{kin}}$$

$$\left(\hbar^4\frac{\mathrm{d}^4}{\mathrm{d}x^4} - Q(x)\right)\psi(x,\hbar) = 0$$

$$\phi(x) = Q(x)\mathrm{d}x^2$$

$$\psi(x,\hbar) = \exp\left(\int_{x_0}^{x} P(x',\hbar)\mathrm{d}x'\right)$$

$$\hbar^2\left(P(x,\hbar)^2 + \frac{\mathrm{d}P}{\mathrm{d}x}(x,\hbar)\right) = Q(x)$$

$$P(x,\hbar) = \sum_{m\geq -1}\hbar^m P_m(x)$$

$$P_{-1}(x)^2 = Q(x)$$

$$2P_{-1}(x)P_0(x) + \frac{\mathrm{d}P_{-1}(x)}{\mathrm{d}x} = 0$$

$$2P_{-1}(x)P_{m+1}(x) + \sum_{\ell=0}^{m} P_\ell(x)P_{m-\ell}(x) + \frac{\mathrm{d}P_m(x)}{\mathrm{d}x} = 0\ (m\geq 0)$$

$$P_{-1}^{(\pm)}(x) = \pm\sqrt{Q(x)}, P_0^{(\pm)}(x) = -\frac{Q'(x)}{4Q(x)}, P_0^{(\pm)}(x) = \pm\frac{4Q(x)Q''(x) - 5(Q'(x))^2}{32Q(x)^{5/2}}, \ldots$$

$$P^{(\pm)}(x,\hbar) = \sum_{m\geq -1}\hbar^m P_m^{(\pm)}(x), \psi_\pm(x,\hbar) = \exp\left(\int_{x_0}^{x} P^{(\pm)}(x',\hbar)\mathrm{d}x'\right)$$

$$P_{\text{odd}}(x,\hbar)=\frac{P^{(+)}(x,\hbar)-P^{(-)}(x,\hbar)}{2},P_{\text{even}}(x,\hbar)=\frac{P^{(+)}(x,\hbar)+P^{(-)}(x,\hbar)}{2}.$$

$$P^{(\pm)}(x,\hbar)=\pm P_{\text{odd}}(x,\hbar)+P_{\text{odd}}(x,\hbar)$$

$$P_{\text{even}}(x,\hbar)=-\frac{1}{2P_{\text{odd}}(x,\hbar)}\frac{\mathrm{d}P_{\text{odd}}}{\mathrm{d}x}(x,\hbar)$$

$$\psi_{\pm}(x,\hbar)=\frac{1}{\sqrt{P_{\text{odd}}(x,\hbar)}}\exp\left(\pm\int_{x_0}^{x}P_{\text{odd}}(x',\hbar)\mathrm{d}x'\right)$$

$$\psi_{\pm}(x,\hbar)=\exp\left(\pm\hbar^{-1}S(x)\right)\sum_{m\geq 0}\hbar^{m+\frac{1}{2}}\psi_{\pm,m}(x),S(x)=\int_{x_0}^{x}\sqrt{Q(x')}\mathrm{d}x'$$

$$\left(\hbar^2\frac{\mathrm{d}^2}{\mathrm{d}x^2}-x\right)\psi(x,\hbar)=0$$

$$P_{-1}^{(\pm)}(x)=\sqrt{x},P_0^{(\pm)}(x)=-\frac{1}{4x},P_1^{(\pm)}(x)=\mp\frac{5}{32x^{5/2}},P_2^{(\pm)}(x)=-\frac{15}{64x^4},\ldots$$

$$\psi_{\pm}^{\text{Airy}}(x,\hbar)=\exp\left(\pm\hbar^{-1}\frac{2x^{3/2}}{3}\right)\frac{\hbar^{1/2}}{x^{1/4}}\left(1\pm\frac{5}{48x^{3/2}}\hbar+\frac{385}{4608x^3}\hbar^2+\cdots\right)$$

$$Q(x)=\sum_{m\geq 0}\hbar^m Q_m(x)$$

$$\Sigma=\{(x,y)\in\mathbb{C}^2\mid y^2=Q(x)\}$$

$$\bar{\Sigma}\ni(x,y)\mapsto x\in\mathbb{P}^1$$

$$\Sigma'=\Sigma\setminus\pi^{-1}(\text{Crit}(\phi)),$$

$$\sup_{x\in K}\left|P_m^{(\pm)}(x)\right|\leq C_K r_K^m m!$$

$$\left(\hbar^2\frac{\mathrm{d}^2}{\mathrm{d}z^2}-\tilde{Q}(z)\right)\tilde{\psi}(z,\hbar)=0,\tilde{Q}(z)=\left(\frac{\mathrm{d}x(z)}{\mathrm{d}z}\right)^2Q(x(z))-\frac{\hbar^2}{2}\{x(z);z\}$$

$$\tilde{\psi}=(\mathrm{d}x/\mathrm{d}z)^{-1/2}\psi$$

$$\{x(z);z\}=\frac{x'''(z)}{x'(z)}-\frac{3}{2}\left(\frac{x''(z)}{x'(z)}\right)^2$$

$$\tilde{P}_{\text{odd}}(z,\hbar)=\frac{\mathrm{d}x(z)}{\mathrm{d}z}P_{\text{odd}}(x(z),\hbar)$$

$$P_m(x)\sim\begin{cases}c_m(x-p)^{\frac{(r-2)m-2}{2}} & \text{for } p\in\mathbb{C}\\ c_m x^{-\frac{(r-2)m+2}{2}} & \text{for } p=\infty\end{cases}$$

$$\left(\hbar^2\frac{\mathrm{d}^2}{\mathrm{d}x^2}-\left(\frac{x^2}{4}-\nu\right)\right)\psi(x,\hbar)=0\ (\nu\in\mathbb{C}^*)$$

$$\int_{\infty_-}^{\infty_+}P_m(x)\mathrm{d}x=\begin{cases}\frac{(2^{1-2k}-1)B_{2k}}{2k(2k-1)\nu^{2k-1}} & \text{if } m=2k-1 \text{ is odd}\\ 0 & \text{if } m=2k \text{ is even}\end{cases}$$

$$\frac{t}{\mathrm{e}^t-1}=1-\frac{t}{2}+\sum_{k\geq 1}\frac{B_{2k}}{(2k)!}t^{2k}$$

$$\mathcal{B}\psi_{\pm}(x,\zeta)=\psi_{\pm,B}(x,\zeta)=\sum_{m\geq 0}\frac{\psi_{\pm,m}(x)}{\Gamma\left(m+\frac{1}{2}\right)}(\zeta\pm S(x))^{m-\frac{1}{2}}$$

$$\mathcal{S}\psi_{\pm}(x,\hbar)=\int_{\mp S(x)}^{\infty}\mathrm{e}^{-\zeta/\hbar}\psi_{\pm,B}(x,\zeta)\mathrm{d}\zeta$$

$$\mathcal{S}\psi_{\pm}(x,\hbar)\sim\psi_{\pm}(x,\hbar),\hbar\to+0.$$

$$\begin{cases}\psi_{+,B}^{\mathrm{Airy}}(x,\zeta)=Cx^{-1}w^{-\frac{1}{2}}{}_2F_1\left(\frac{1}{6},\frac{5}{6},\frac{1}{2};w\right)\\ \psi_{-,B}^{\mathrm{Airy}}(x,\zeta)=Cx^{-1}(w-1)^{-\frac{1}{2}}{}_2F_1\left(\frac{1}{6},\frac{5}{6},\frac{1}{2};1-w\right)\end{cases}$$


$$C=\frac{\sqrt{3}}{2\sqrt{\pi}},w=\frac{\zeta+S(x)}{2S(x)},S(x)=\frac{2}{3}x^{3/2}$$

$${}_2F_1(\alpha,\beta,\gamma;w)=\sum_{n\geq 0}\frac{(\alpha)_n(\beta)_n}{(\gamma)_n n!}w^n$$


$$(a)_n=\begin{cases}1 & n=0\\ a(a+1)\cdots(a+n-1) & n\geq 1\end{cases}$$

$$\left(\frac{\mathrm{d}^2}{\mathrm{d}w^2}+\frac{\gamma-(\alpha+\beta+1)w}{w(1-w)}\frac{\mathrm{d}}{\mathrm{d}w}-\frac{\alpha\beta}{w(1-w)}\right)F=0$$


$$\mathrm{Im}\int_v^x\sqrt{Q(x')}\mathrm{d}x'=0$$




(a) $Q(x) = x$




(b) $Q(x) = x^3 + x$




(c) $Q(x) = \frac{(x-1)(x-2)}{x^2(x+1)^2}$



(d) $Q(x) = \frac{x+1}{x}$



(e) $Q(x) = x^3 - x$



(f) $Q(x) = \frac{x-2}{x^2}$

$$T(x,\hbar) = \sum_{m\geq 1} \hbar^m P_m(x)\ (= P(x,\hbar) - \hbar^{-1}P_{-1}(x) - P_0(x)).$$

$$2\sqrt{Q(x)}(\hbar^{-1}T(x,\hbar) - P_1(x)) + \frac{\mathrm{d}T}{\mathrm{d}x}(x,\hbar) = -2P_0(x)T(x,\hbar) - T(x,\hbar)^2.$$

$$z = z(x) = \int_{x_*}^{x} \sqrt{Q(x')}\mathrm{d}x'$$

$$\left(\frac{\partial}{\partial\zeta} + \frac{1}{2}\frac{\partial}{\partial z}\right)T_B(z,\zeta) = A_1(z)T_B(z,\zeta) + A_2(z)\frac{\partial}{\partial\zeta}T_B * T_B(z,\zeta)$$


$$A_1(z(x)) = -\frac{P_0(x)}{\sqrt{Q(x)}}, A_2(z(x)) = -\frac{1}{2\sqrt{Q(x)}}$$


$$T_B * T_B(z,\zeta) = \int_0^{\zeta} T_B(z,\sigma)T_B(z,\zeta-\sigma)\mathrm{d}\sigma$$

$$T_B(z,\zeta) = A_0\left(z - \frac{\zeta}{2}\right) + \int_0^{\zeta} A_1\left(z - \frac{\zeta-t}{2}\right)T_B\left(z - \frac{\zeta-t}{2}, t\right)dt$$
$$+ \int_0^{\zeta} A_2\left(z - \frac{\zeta-t}{2}\right)\frac{\partial}{\partial\zeta}T_B * T_B\left(z - \frac{\zeta-t}{2}, t\right)\mathrm{d}t$$

$$A_0(z(x)) = T_B(z(x),0) = P_1(x)$$

$$A_i\left(z - \frac{\zeta-t}{2}\right)\int_{x_*}^{x} \sqrt{Q(x')}\mathrm{d}x' < \int_{p}^{x_*} \sqrt{Q(x')}\mathrm{d}x' > 0$$





$$\psi_{\pm}(x,\hbar) = \frac{1}{\sqrt{P_{\text{odd}}(x,\hbar)}} \exp\left(\pm \int_v^x P_{\text{odd}}(x,\hbar)\mathrm{d}x\right)$$

$$(\Psi_+^{\mathrm{I}}, \Psi_-^{\mathrm{I}}) = (\Psi_+^{\mathrm{II}}, \Psi_-^{\mathrm{II}}) \cdot S,$$

$$S = \begin{cases} \begin{pmatrix} 1 & 0 \\ \mathrm{i} & 1 \end{pmatrix} & \text{if } \int_v^x \sqrt{Q(x')}\,\mathrm{d}x' > 0 \text{ on } C, \\ \begin{pmatrix} 1 & \mathrm{i} \\ 0 & 1 \end{pmatrix} & \text{if } \int_v^x \sqrt{Q(x')}\,\mathrm{d}x' < 0 \text{ on } C. \end{cases}$$

$$S(x) = \int_0^x \sqrt{Q(x')}\mathrm{d}x' = \frac{2}{3}x^{3/2} > 0$$

$$\int_L e^{-\zeta/\hbar}\psi_{+,B}^{\text{Airy}}(x,\zeta)\mathrm{d}\zeta$$


$$\int_{-S(x)}^{\infty} e^{-\zeta/\hbar}\psi_{+,B}^{\text{Airy}}(x,\zeta)\mathrm{d}\zeta + \int_{+S(x)}^{\infty} e^{-\zeta/\hbar}\left(\dot{\Delta}_{\zeta=+S(x)}\psi_{+,B}^{\text{Airy}}(x,\zeta)\right)\mathrm{d}\zeta$$

$$
{}_2F_1(\alpha,\beta,\gamma;w) = \frac{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma)}{\Gamma(\alpha)\Gamma(\beta)}(1-w)^{\gamma-\alpha-\beta}\,{}_2F_1(\gamma-\alpha,\gamma-\beta,\gamma-\alpha-\beta+1;1-w)
$$


$$
+\frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)}\,{}_2F_1(\alpha,\beta,\alpha+\beta-\gamma+1,;1-w)
$$

$$
{}_2F_1(\alpha,\beta,\gamma;w) = (1-w)^{\gamma-\alpha-\beta}\,{}_2F_1(\gamma-\alpha,\gamma-\beta,\gamma;w)
$$

$$
\dot{\Delta}_{\zeta=+S(x)}\psi^{\text{Airy}}_{+,B}(x,\zeta) = \mathrm{i}\psi^{\text{Airy}}_{-,B}(x,\zeta).
$$



$$
\Psi^{\text{II}}_{+} = \Psi^{\text{II}}_{+} + \tilde{\Psi}^{\text{II}}_{-},
$$



$$Q = Q_0(x) + \hbar^2 Q_2(x)$$

$$(\Psi_+^{\mathrm{I}}, \Psi_-^{\mathrm{I}}) = (\Psi_+^{\mathrm{II}}, \Psi_-^{\mathrm{II}}) \cdot S,$$

$$S = \begin{cases} \begin{pmatrix} 1 & 0 \\ 2\mathrm{i}\cos(\pi\sqrt{1+4\lambda}) & 1 \end{pmatrix} & \text{if } \int_v^x \sqrt{Q_0(x)}\,\mathrm{d}x > 0 \text{ on } C, \\ \begin{pmatrix} 1 & 2\mathrm{i}\cos(\pi\sqrt{1+4\lambda}) \\ 0 & 1 \end{pmatrix} & \text{if } \int_v^x \sqrt{Q_0(x)}\,\mathrm{d}x < 0 \text{ on } C, \end{cases}$$

$$\lambda = \lim_{x\to v}(x-v)^2 Q_2(x).$$

$$Q = \frac{1}{x} + \hbar^2 \frac{\lambda}{x^2}$$

$$\begin{cases} \psi_{+,B}(x,\zeta) = Cw^{-\frac{1}{2}}\,{}_2F_1\left(\alpha - \frac{1}{2}, \beta - \frac{1}{2}, \frac{1}{2}; w\right) \\ \psi_{-,B}(x,\zeta) = C(w-1)^{-\frac{1}{2}}\,{}_2F_1\left(\alpha - \frac{1}{2}, \beta - \frac{1}{2}, \frac{1}{2}; 1-w\right) \end{cases}$$

$$C = \frac{1}{2\sqrt{\pi}}, w = \frac{\zeta}{4\sqrt{x}} + \frac{1}{2}$$

$$\psi_\pm(x,\hbar) = \frac{1}{\sqrt{P_{\mathrm{odd}}(x,\hbar)}} \exp\left(\pm \int_{v_1}^x P_{\mathrm{odd}}(x,\hbar)\mathrm{d}x\right)$$

$$(\Psi_+^{\mathrm{I}}, \Psi_-^{\mathrm{I}}) = (\Psi_+^{\mathrm{II}}, \Psi_-^{\mathrm{II}}) \cdot \begin{pmatrix} 1 & \mathrm{i} \\ 0 & 1 \end{pmatrix}$$


$$\tilde{\psi}_{\pm}(x,\hbar)=\frac{1}{\sqrt{P_{\text{odd}}\,(x,\hbar)}}\exp\left(\pm\int_{v_2}^{x}P_{\text{odd}}\,(x,\hbar)\mathrm{d}x\right)$$

$$\psi_{\pm}(x,\hbar)=\exp\left(\pm\frac{1}{2}V_{\gamma}(\hbar)\right)\tilde{\psi}_{\pm}(x,\hbar),V_{\gamma}(\hbar)=\oint_{\gamma}P_{\text{odd}}\,(x,\hbar)\mathrm{d}x$$

$$\left(\Psi_{+}^{\text{II}},\Psi_{-}^{\text{II}}\right)=\left(\Psi_{+}^{\text{III}},\Psi_{-}^{\text{III}}\right)\cdot\begin{pmatrix}1 & \mathrm{i}\mathrm{e}^{-\mathcal{V}_{\gamma}}\\ 0 & 1\end{pmatrix}$$

$$\left(\Psi_{+}^{\text{I}},\Psi_{-}^{\text{I}}\right)=\left(\Psi_{+}^{\text{III}},\Psi_{-}^{\text{III}}\right)\cdot\begin{pmatrix}1 & \mathrm{i}\left(1+\mathrm{e}^{-\mathcal{V}_{\gamma}}\right)\\ 0 & 1\end{pmatrix}$$

$$\psi_{\pm,\infty}(x,\hbar)=\frac{1}{\sqrt{P_{\text{odd}}\,(x,\hbar)}}\exp\left(\pm\hbar^{-1}\int_{v_1}^{x}P_{-1}(x')\mathrm{d}x'\pm\int_{\infty}^{x}\left(P_{\text{odd}}\,(x',\hbar)-\hbar^{-1}P_{-1}(x')\right)\mathrm{d}x'\right)$$



$$\psi_{\pm}(x,\hbar)=\exp\left(\pm\frac{1}{2}W(\hbar)\right)\psi_{\pm,\infty}(x,\hbar)$$

$$W(\hbar)=\sum_{k\geq 1}\frac{\left(2^{1-2k}-1\right)B_{2k}}{2k(2k-1)v^{2k-1}}\hbar^{2k-1}$$

$$\mathcal{B}W(\zeta)=\frac{1}{2\zeta}\left(\frac{1}{\exp\left(\frac{\zeta}{2v}\right)-1}+\frac{1}{\exp\left(\frac{\zeta}{2v}\right)+1}-\frac{2v}{\zeta}\right)$$

$$\mathcal{S}\mathrm{e}^{W(\hbar)}=\frac{\mathrm{e}^{v/\hbar}\Gamma\left(\frac{v}{\hbar}+\frac{1}{2}\right)}{\sqrt{2\pi}(v/\hbar)^{v/\hbar}}$$

$$V_{\gamma}(\hbar)=\oint_{\gamma}P_{\text{odd}}\,(x,\hbar)\mathrm{d}x\,\left(\gamma\in H_1(\Sigma',\mathbb{Z})\right)$$

$$Q = \left(\frac{\theta_0^2}{x^2} + \frac{\theta_1^2}{(x-1)^2} + \frac{\theta_\infty^2 - \theta_0^2 - \theta_1^2}{x(x-1)}\right) + \hbar^2\left(-\frac{1}{4x^2} - \frac{1}{4(x-1)^2} + \frac{1}{4x(x-1)}\right)$$

$$Q = Q_0(x) + \hbar Q_1(x) + \hbar^2 Q_2(x)$$

$$\rho^{(\pm)}(\hbar) = \mathrm{Res}_{x=0} P^{(\pm)}(x,\hbar)\mathrm{d}x \left(= \sum_{m\geq -1} \hbar^m \mathrm{Res}_{x=0} P_m^{(\pm)}(x)\mathrm{d}x\right)$$

$$\left(\hbar^n \frac{\mathrm{d}^n}{\mathrm{d}x^n} + q_1(x)\hbar^{n-1}\frac{\mathrm{d}^{n-1}}{\mathrm{d}x^{n-1}} + \cdots + q_{n-1}(x)\hbar\frac{\mathrm{d}}{\mathrm{d}x} + q_n(x)\right)\psi(x,\hbar) = 0$$

$$\mathfrak{S}\mathrm{e}^W = \begin{cases} \mathrm{e}^W\left(1 + \mathrm{e}^{2\pi \mathrm{i}\nu/\hbar}\right) & \text{if } \nu \in \mathrm{i}\mathbb{R}_{>0} \\ \mathrm{e}^W\left(1 + \mathrm{e}^{-2\pi \mathrm{i}\nu/\hbar}\right)^{-1} & \text{if } \nu \in \mathrm{i}\mathbb{R}_{<0} \end{cases}$$

$$\psi_{\pm}^{\mathrm{Airy}}(x,\hbar) = \exp\left(\sum_{g\geq 0, n\geq 1} \frac{(\pm\hbar)^{2g-2+n}}{n!} \int_\infty^{z(x)} \cdots \int_\infty^{z(x)} \left(W_{g,n}^{\mathrm{Airy}}(z_1,\ldots,z_n) - \delta_{g,0}\delta_{n,2}\frac{\mathrm{d}x(z_1)\mathrm{d}x(z_2)}{\left(x(z_1)-x(z_2)\right)^2}\right)\right)$$

$$(C = \mathbb{P}^1, x(z) = z^2, y(z) = z)$$

$$W_{g,n}^{\mathrm{Airy}}(z_1,\ldots,z_n) = \frac{1}{2^{2g-2+n}} \sum_{d_1,\ldots,d_n\geq 0} \left(\int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1}\cdots\psi_n^{d_n}\right) \prod_{i=1}^n \frac{(2d_i+1)!!}{z_i^{2d_i+2}}\,\mathrm{d}z_i$$

$$F_g^{\mathrm{Weber}} = \frac{B_{2g}}{2g(2g-2)\nu^{2g-2}}\ (g \geq 2)$$

$$\left(C = \mathbb{P}^1, x(z) = \sqrt{\nu}(z + z^{-1}), y(z) = \frac{\sqrt{\nu}}{2}(z - z^{-1})\right)$$

$$(P_{\mathrm{I}}): \hbar^2 \frac{\mathrm{d}^2 q}{\mathrm{d}t^2} = 6q^2 + t$$

$$\begin{cases} (L_{\mathrm{I}}): \left[\hbar^2\frac{\partial^2}{\partial x^2} - \frac{\hbar}{x-q}\left(\hbar\frac{\partial}{\partial x} - p\right) - (4x^3 + 2tx + 2H)\right]\psi = 0 \\ (D_{\mathrm{I}}): \left[\hbar\frac{\partial}{\partial t} - \frac{1}{2(x-q)}\left(\hbar\frac{\partial}{\partial x} - p\right)\right]\psi = 0 \end{cases}$$

$$\hbar\frac{\mathrm{d}q}{\mathrm{d}t} = \frac{\partial H}{\partial p}, \hbar\frac{\mathrm{d}p}{\mathrm{d}t} = -\frac{\partial H}{\partial q}$$

$$\hbar^2 \frac{\mathrm{d}}{\mathrm{d}t}\log\,\tau(t) = H(t)$$

$$H(t) = \frac{\hbar^2}{t-a}(1 + O(t-a))$$

$$y^2 = 4x^3 + 2tx + 2''H|_{\hbar=0}$$

$$\Sigma_{P_I}: y^2 = 4x^3 + 2tx + u(t,v)$$

$$v = \frac{1}{2\pi \mathrm{i}} \oint_A y \, \mathrm{d}x$$

$$C = \mathbb{C}/L, x = \wp(z), y = \frac{\mathrm{d}\wp}{\mathrm{d}z}(z)$$

$$B(z_1, z_2) = \left(\wp(z_1 - z_2) + \frac{\eta_A}{\omega_A}\right) \mathrm{d}z_1 \, \mathrm{d}z_2$$

$$\eta_A = -\oint_A \frac{x \, \mathrm{d}x}{y}$$

$$R = \left\{\frac{\omega_A}{2}, \frac{\omega_B}{2}, \frac{\omega_A + \omega_B}{2}\right\}$$

$$W_{0,1}(z) = y(z)\mathrm{d}x(z), W_{0,2}(z_1, z_2) = B(z_1, z_2)$$

$$W_{g,n+1}(z_0, z_1, \dots, z_n) = \sum_{r \in R} \mathrm{Res}_{z=r} \frac{\int_{w=\sigma(z)}^{w=z} W_{0,2}(z_0, w)}{2(y(z) - y(\sigma(z)))\mathrm{d}x(z)} R_{g,n}(z, z_1, \dots, z_n),$$

$$R_{g,n}(z, z_1, \dots, z_n)$$

$$= W_{g-1,n+1}(z, \sigma(z), z_1, \dots, z_n) + \sum_{\substack{g_1+g_2=g \\ I \sqcup J = \{1,\dots,n\}}}{}' W_{g_1,1+|I|}(z, z_I) W_{g_2,1+|J|}(\sigma(z), z_J),$$

$$F_g = \frac{1}{2-2g} \sum_{r \in R} \mathrm{Res}_{z=r} \left(\int^z W_{0,1} \, W_{0,1}(z) \right) W_{g,1}(z)$$

$$F_0 = \frac{tu}{5} + \frac{v}{2} \oint_B y \, \mathrm{d}x, F_1 = -\frac{1}{12} \log\left(\omega_A^6 \mathcal{D}\right)$$

$$\frac{\partial F_0}{\partial t} = \frac{u}{2}, \frac{\partial F_0}{\partial v} = \oint_B y \, \mathrm{d}x, \frac{\partial^2 F_0}{\partial v^2} = 2\pi \mathrm{i} \frac{\omega_B}{\omega_A}$$

$$Z(t, v, \hbar) = \exp\left(\sum_{g \geq 0} \hbar^{2g-2} F_g(t, v) \right)$$

$$\chi_{\pm}(x,t,\nu,\hbar)$$

$$= \exp\left(\sum_{g\geq 0,n\geq 1}\frac{(\pm\hbar)^{2g-2+n}}{n!}\int_0^{z(x)}\cdots\int_0^{z(x)}\left(W_{g,n}(z_1,\ldots,z_n)-\delta_{g,0}\delta_{n,2}\frac{\mathrm{d}x(z_1)\mathrm{d}x(z_2)}{\left(x(z_1)-x(z_2)\right)^2}\right)\right)$$

$$\left[\hbar^2\frac{\partial^2}{\partial x^2}-2\hbar^2\frac{\partial}{\partial t}-\left(4x^3+2tx+2\hbar^2\frac{\partial F}{\partial t}(t,\nu,\hbar)\right)\right]\chi_{\pm}(x,t,\nu,\hbar)=0$$

$$F(t,\nu,\hbar)=\sum_{g\geq 0}\hbar^{2g-2}F_g(t,\nu)(=\log\,Z(t,\nu,\hbar))$$

$$\chi_{\pm}(x,t,\nu,\hbar)\mapsto\begin{cases}\mathrm{e}^{\pm 2\pi\mathrm{i}\nu/\hbar}\chi_{\pm}(x,t,\nu,\hbar) & \text{along } A\text{-cycle}\\ \dfrac{Z(t,\nu\pm\hbar,\hbar)}{Z(t,\nu,\hbar)}\chi_{\pm}(x,t,\nu\pm\hbar,\hbar) & \text{along } B\text{-cycle.}\end{cases}$$

$$\psi_{\pm}(x,t,\nu,\rho,\hbar)=\frac{\sum_{k\in\mathbb{Z}}\mathrm{e}^{2\pi\mathrm{i}k\rho/\hbar}Z(t,\nu+k\hbar,\hbar)\chi_{\pm}(x,t,\nu+k\hbar,\hbar)}{\sum_{k\in\mathbb{Z}}\mathrm{e}^{2\pi\mathrm{i}k\rho/\hbar}Z(t,\nu+k\hbar,\hbar)},$$


$$\psi_{\pm}(x,t,\nu,\rho,\hbar)\mapsto\begin{cases}\mathrm{e}^{\pm 2\pi\mathrm{i}\nu/\hbar}\psi_{\pm}(x,t,\nu,\rho,\hbar) & \text{along } A\text{-cycle}\\ \mathrm{e}^{\mp 2\pi\mathrm{i}\rho/\hbar}\psi_{\pm}(x,t,\nu,\rho,\hbar) & \text{along } B\text{-cycle.}\end{cases}$$

$$q(t,\nu,\rho;\hbar)=-\hbar^2\frac{\mathrm{d}^2}{\mathrm{d}t^2}\log\,\tau(t,\nu,\rho,\hbar),p(t,\nu,\rho;\hbar)=-\hbar^3\frac{\mathrm{d}^3}{\mathrm{d}t^3}\log\,\tau(t,\nu,\rho,\hbar)$$

$$\tau(t,\nu,\rho,\hbar)=\sum_{k\in\mathbb{Z}}\mathrm{e}^{2\pi\mathrm{i}k\rho/\hbar}Z(t,\nu+k\hbar,\hbar)$$

$$\begin{cases}q(t,\nu,\rho,\hbar)=\wp\left(\dfrac{5t}{4\hbar}+\left(\dfrac{\rho}{\hbar}+\dfrac{1}{2}\right)\omega_A+\left(\dfrac{\nu}{\hbar}+\dfrac{1}{2}\right)\omega_B\right)+O(\hbar)\\ p(t,\nu,\rho,\hbar)=\wp'\left(\dfrac{5t}{4\hbar}+\left(\dfrac{\rho}{\hbar}+\dfrac{1}{2}\right)\omega_A+\left(\dfrac{\nu}{\hbar}+\dfrac{1}{2}\right)\omega_B\right)+O(\hbar)\end{cases}$$

$$H(t,\nu,\rho,\hbar)=\frac{u(t,\nu)}{2}+O(\hbar)$$



$$\mathcal{T}(t,\nu,\rho,\hbar)=\sum_{k\in\mathbb{Z}}\mathrm{e}^{2\pi\mathrm{i}k\rho/\hbar}Z(t,\nu+k\hbar,\hbar)$$

$$\Psi_{\pm}(x,t,\nu,\rho,\hbar)=\frac{\sum_{k\in\mathbb{Z}}\ \mathrm{e}^{2\pi\mathrm{i}k\rho/\hbar}\mathcal{Z}(t,\nu+k\hbar,\hbar)\mathcal{X}_{\pm}(x,t,\nu+k\hbar,\hbar)}{\sum_{k\in\mathbb{Z}}\ \mathrm{e}^{2\pi\mathrm{i}k\rho/\hbar}\mathcal{Z}(t,\nu+k\hbar,\hbar)}$$

$$\left(\Psi_{+}^{(j)},\Psi_{-}^{(j)}\right)=\left(\Psi_{+}^{(j+1)},\Psi_{-}^{(j+1)}\right)\cdot S_{j}.$$

$$\mathrm{Re}\int_{v}^{x}\sqrt{\phi(x)}>0$$

$$\begin{cases}\mathcal{X}_{+}^{\mathrm{I}}(x,t,\nu,\hbar)=\mathcal{X}_{+}^{\mathrm{II}}(x,t,\nu,\hbar)+\mathrm{i}\dfrac{\mathcal{Z}(t,\nu+\hbar,\hbar)}{\mathcal{Z}(t,\nu,\hbar)}\mathcal{X}_{-}^{\mathrm{II}}(x,t,\nu+\hbar,\hbar)\\ \mathcal{X}_{-}^{\mathrm{I}}(x,t,\nu,\hbar)=\mathcal{X}_{-}^{\mathrm{II}}(x,t,\nu,\hbar)\end{cases}$$

$$\begin{cases}\Psi_{+}^{\mathrm{I}}(x,t,\nu,\rho,\hbar)=\Psi_{+}^{\mathrm{II}}(x,t,\nu,\rho,\hbar)+\mathrm{i}\mathrm{e}^{-2\pi\mathrm{i}\rho/\hbar}\Psi_{-}^{\mathrm{II}}(x,t,\nu,\rho,\hbar)\\ \Psi_{-}^{\mathrm{I}}(x,t,\nu,\rho,\hbar)=\Psi_{-}^{\mathrm{II}}(x,t,\nu,\rho,\hbar)\end{cases}$$

$$\text{At } t=t_c-\mathrm{i}\epsilon:\begin{cases}s_{-2}=\mathrm{i}X_A\\ s_{-1}=\mathrm{i}(X_A^{-1}-X_A^{-1}X_B^{-1}+X_B^{-1})\\ s_0=\mathrm{i}X_B\\ s_1=\mathrm{i}(X_B^{-1}-X_AX_B^{-1})\\ s_2=\mathrm{i}(X_A^{-1}-X_A^{-1}X_B)\end{cases}$$

$$\text{At } t=t_c+\mathrm{i}\epsilon:\begin{cases}s_{-2}=\mathrm{i}(X_A-X_AX_B)\\ s_{-1}=\mathrm{i}(X_B^{-1}-X_A^{-1}X_B^{-1})\\ s_0=\mathrm{i}X_B\\ s_1=\mathrm{i}(X_A-X_AX_B^{-1}+X_B^{-1})\\ s_2=\mathrm{i}X_A^{-1}\end{cases}$$

$$X_A=\mathrm{e}^{2\pi\mathrm{i}\nu/\hbar},X_B=\mathrm{e}^{2\pi\mathrm{i}\rho/\hbar}$$

$$1+s_js_{j-1}+is_{j+2}=0\ (j\mathrm{mod}5)$$

$$\mathcal{T}^{-}(t,\nu^{-},\rho^{-},\hbar)=\mathrm{e}^{\frac{1}{2\pi\mathrm{i}}\mathrm{Li}_2\left(\mathrm{e}^{2\pi\mathrm{i}\rho^{+}/\hbar}\right)}\mathcal{T}^{+}(t,\nu^{+},\rho^{+},\hbar)$$

$$(\nu^{+},\rho^{+})=\left(\nu^{-}-\frac{\hbar}{2\pi\mathrm{i}}\log\left(1-\mathrm{e}^{2\pi\mathrm{i}\rho^{-}/\hbar}\right),\rho^{-}\right)$$

$$\mathfrak{S}\tau(t,\nu,\rho,\hbar)=\mathrm{e}^{\frac{1}{2\pi\mathrm{i}}\mathrm{Li}_2\left(\mathrm{e}^{2\pi\mathrm{i}\rho/\hbar}\right)}\tau\left(t,\nu-\frac{\hbar}{2\pi\mathrm{i}}\log\left(1-\mathrm{e}^{2\pi\mathrm{i}\rho/\hbar}\right),\rho,\hbar\right)$$

$$\mathfrak{S}Z(t,\nu,\hbar)=\exp\left(\frac{1}{2\pi\mathrm{i}}\mathrm{Li}_2\left(\mathrm{e}^{-\hbar\partial_\nu}\right)-\frac{\hbar\partial_\nu}{2\pi\mathrm{i}}\log\left(1-\mathrm{e}^{-\hbar\partial_\nu}\right)\right)Z(t,\nu,\hbar)$$

$$\mathfrak{S}Z(t,\nu,\hbar)=\sum_{n=0}^{\infty}Z^{(n)}(t,\nu,\hbar)$$

$$Z^{(0)}(t,\nu,\hbar)=Z(t,\nu,\hbar),Z^{(1)}(t,\nu,\hbar)=\left(1+\frac{\hbar}{2\pi\mathrm{i}}\frac{\partial F}{\partial\nu}(t,\nu-\hbar,\hbar)\right)Z(t,\nu-\hbar,\hbar),\ldots$$

$$q(t,\alpha,\beta,\hbar)=q_0(t)+\sum_{\ell\geq 1}\hbar^{\ell/2}q_{\ell/2}(t,\alpha,\beta,\hbar)$$

$$q_{1/2}=\alpha a_1(t)\mathrm{e}^{A(t)/\hbar}+\beta a_{-1}(t)\mathrm{e}^{-A(t)/\hbar}$$
$$q_1=\alpha^2 a_2(t)\mathrm{e}^{2A(t)/\hbar}+\alpha\beta a_0(t)+\beta^2 a_{-2}(t)\mathrm{e}^{-2A(t)/\hbar},\ldots$$

$$\begin{cases}\alpha=\dfrac{\sqrt{\pi}}{\Gamma\left(\frac{\nu}{\hbar}\right)}2^{\frac{\nu}{\hbar}}\mathrm{e}^{-2\pi\mathrm{i}(\nu+\rho)/\hbar}\\ \beta=\dfrac{\Gamma\left(1+\frac{\nu}{\hbar}\right)}{\sqrt{\pi}}2^{-1-\frac{\nu}{\hbar}}\mathrm{e}^{2\pi\mathrm{i}(\nu+\rho)/\hbar}\end{cases}$$

$$\mathrm{e}^{\pm S/\hbar}\hbar^{\alpha}=\int_{\mp S}^{\infty}\mathrm{e}^{-\zeta/\hbar}\frac{(\zeta\pm S)^{\alpha-1}}{\Gamma(\alpha)}\mathrm{d}\zeta\ (\mathrm{Re}\alpha>0)$$

$$W^{(I)}\equiv\frac{1}{N}\mathrm{tr}\mathcal{P}\exp\left\{g\oint_{\mathcal{C}}d\tau\left[iA_\mu^I\dot{x}^\mu(\tau)+\frac{1}{2}(\varphi^I(x)+\bar{\varphi}^I(x))\right]\right\},$$

$$W_0\equiv\frac{1}{\sqrt{M}}\left(W^{(0)}+W^{(1)}+\cdots+W^{(M-1)}\right)$$

$$W_\alpha\equiv\frac{1}{\sqrt{M}}\sum_{I=0}^{M-1}\rho^{I\alpha}W^{(I)}$$

$$W_\alpha\equiv\frac{1}{\sqrt{M}}\sum_{I=0}^{M-1}\rho^{I\alpha}W^{(I)}$$

$$\mathcal{W}\equiv\partial_m^2\log\langle W\rangle|_{m=0}=\int d^4x_1d^4x_2\mu(x_1,x_2)\langle\mathcal{O}_2(x_1)\mathcal{O}_2(x_2)\rangle_W,$$

$$\mathcal{Z}=\int\left(\prod_{I=0}^{M-1}da_I\right)\mathrm{e}^{-\mathrm{tr}a_I^2}\left|\mathcal{Z}_{1-\mathrm{loop}}\mathcal{Z}_{\mathrm{inst}}\right|^2$$

$$da_I=\prod_{b=1}^{N^2-1}\frac{da_I^b}{\sqrt{2\pi}}$$

$$\left|\mathcal{Z}_{1-\mathrm{loop}}\right|^2=\mathrm{e}^{-S_{\mathrm{int}}}$$

$$S_{\mathrm{int}}=\sum_{I=0}^{M-1}\left[\sum_{m=2}^{+\infty}\sum_{k=2}^{2m}(-1)^{m+k}\left(\frac{\lambda}{8\pi^2N}\right)^m\binom{2m}{k}\frac{\zeta_{2m-1}}{2m}\left(\mathrm{tr}a_I^{2m-k}-\mathrm{tr}a_{I+1}^{2m-k}\right)\left(\mathrm{tr}a_I^k-\mathrm{tr}a_{I+1}^k\right)\right],$$

$$\langle f(a)\rangle=\frac{\langle f(a)\mathrm{e}^{-S_{\mathrm{int}}}\rangle_0}{\langle\mathrm{e}^{-S_{\mathrm{int}}}\rangle_0}$$

$$t_{n_1,n_2,\ldots,n_q}=\langle\mathrm{tr}a^{n_1}\mathrm{tr}a^{n_2}\cdots\mathrm{tr}a^{n_q}\rangle$$

$$A_{\alpha,k} = \frac{1}{\sqrt{M}} \sum_{I=0}^{M-1} \rho^{\alpha I} \mathrm{tr} a_I^k$$

$$\hat{A}_{\alpha,k} \equiv A_{\alpha,k} - \langle A_{\alpha,k} \rangle$$

$$\hat{A}_{\alpha,k} = A_{\alpha,k} \text{ for } \alpha \neq 0$$

$$\left\langle \hat{A}_{\alpha,k} \hat{A}^{\dagger}_{\beta,\ell} \right\rangle_0 \propto N^{\frac{k+\ell}{2}} \delta_{\alpha,M-\beta} \text{ with } k+\ell \text{ even},$$

$$\left\langle \hat{A}_{\alpha,k} \hat{A}_{\beta,\ell} \hat{A}^{\dagger}_{\gamma,q} \right\rangle_0 \propto N^{\frac{k+\ell+q}{2}-1} \delta_{\alpha+\beta,M-\gamma}, \text{ with } k+\ell+q \text{ even}.$$

$$\hat{A}_{\alpha,k} = \left(\frac{N}{2}\right)^{\frac{k}{2}\left\lfloor\frac{k-1}{2}\right\rfloor} \sum_{i=0} \sqrt{k-2i} \binom{k}{i} \mathcal{P}_{\alpha,k-2i}$$

$$S_{\mathrm{int}} = -\frac{1}{2} \sum_{\alpha=0}^{M-1} \sum_{k,\ell=2}^{\infty} s_\alpha \mathcal{P}^{\dagger}_{\alpha,k} \mathrm{X}_{k,\ell} \mathcal{P}_{\alpha,\ell}$$

$$s_\alpha \equiv \sin\left(\frac{\pi\alpha}{M}\right)^2$$

$$\mathrm{X}_{k,\ell} = -8(-1)^{\frac{k+\ell+2k\ell}{2}} \sqrt{k\ell} \int_0^\infty \frac{dt}{t} \frac{\mathrm{e}^t}{(\mathrm{e}^t-1)^2} J_k\left(\frac{t\sqrt{\lambda}}{2\pi}\right) J_\ell\left(\frac{t\sqrt{\lambda}}{2\pi}\right)$$

$$\mathrm{X}^{\mathrm{even}}_{k,\ell} \equiv \mathrm{X}_{2k,2\ell} = -\int_0^\infty dt U^{\mathrm{even}}_k(t) U^{\mathrm{even}}_\ell(t)$$
$$\mathrm{X}^{\mathrm{odd}}_{k,\ell} \equiv \mathrm{X}_{2k+1,2\ell+1} = -\int_0^\infty dt U^{\mathrm{odd}}_k(t) U^{\mathrm{odd}}_\ell(t)$$

$$U^{\mathrm{even}}_k(t) = (-1)^k \frac{2\sqrt{k}}{\sqrt{t}\sinh(t/2)} J_{2k}\left(\frac{t\sqrt{\lambda}}{2\pi}\right)$$
$$U^{\mathrm{odd}}_k(t) = (-1)^k \frac{\sqrt{2(2k+1)}}{\sqrt{t}\sinh(t/2)} J_{2k+1}\left(\frac{t\sqrt{\lambda}}{2\pi}\right)$$

$$\left\langle \mathcal{P}_{\alpha,k} \mathcal{P}^{\dagger}_{\beta,\ell} \right\rangle \simeq \delta_{\alpha,\beta} \mathrm{D}^{(\alpha)}_{k,\ell}$$

$$\mathrm{D}^{(\alpha)}_{k,\ell} = \left(\frac{1}{1-s_\alpha \mathrm{X}}\right)_{k,\ell}.$$

$$\mathrm{D}^{(\alpha)\ \mathrm{even}}_{k,\ell} \equiv \mathrm{D}^{(\alpha)}_{2k,2\ell}, \mathrm{D}^{(\alpha)\ \mathrm{odd}}_{k,\ell} \equiv \mathrm{D}^{(\alpha)}_{2k+1,2\ell+1}$$

$$\left\langle \mathcal{P}_{\alpha,k} \mathcal{P}_{\beta,\ell} \mathcal{P}^{\dagger}_{\gamma,q} \right\rangle \simeq \delta_{\alpha+\beta,\gamma} \frac{\mathrm{d}^{(\alpha)}_k \mathrm{d}^{(\beta)}_\ell \mathrm{d}^{(\gamma)}_q}{\sqrt{M}N} \text{ with } k+\ell+q \text{ even},$$

$$\mathrm{d}^{(\alpha)}_k = \sum_{\ell=2}^{\infty} \mathrm{D}^{(\alpha)}_{k,\ell} \sqrt{\ell}$$

$$W_\alpha = \frac{1}{\sqrt{M}N}\sum_{I=0}^{M-1} \rho^{\alpha I}\mathrm{trexp}\left[\sqrt{\frac{\lambda}{2N}}a_I\right] = \frac{1}{N}\sum_{k=0}^{\infty}\frac{1}{k!}\left(\frac{\lambda}{2N}\right)^{\frac{k}{2}} A_{\alpha,k}.$$

$$W_{\mathrm{conn}}^{(2)} = \langle W_0 W_0\rangle_0 - \langle W_0\rangle_0^2$$

$$\frac{\langle W_\alpha W_\alpha^\dagger\rangle}{W_{conn}^{(2)}(\lambda)} \equiv 1 + \Delta w^{(\alpha)}(M,\lambda)$$

$$\Delta w^{(\alpha)}(M,\lambda) \simeq \frac{2}{\sqrt{\lambda}}\sum_{k=2}^{\infty}\sum_{\ell=2}^{\infty}\frac{I_k(\sqrt{\lambda})I_\ell(\sqrt{\lambda})}{I_1(\sqrt{\lambda})I_2(\sqrt{\lambda})}\sqrt{k\ell}\left(\mathrm{D}_{k,\ell}^{(\alpha)} - \delta_{k,\ell}\right)$$

$$\frac{\left\langle W_\alpha W_\beta W_{\alpha+\beta}^\dagger\right\rangle}{\sqrt{M}W_{conn}^{(3)}(\lambda)} \equiv 1 + \Delta w^{(\alpha,\beta)}(M,\lambda)$$

$$1 + \Delta w^{(\alpha,\beta)} \simeq \frac{8}{\lambda^{3/2}}\left(\frac{I_1(\sqrt{\lambda})^2}{I_1(\sqrt{\lambda})^2 + 3I_2(\sqrt{\lambda})^2}\right)\left[\prod_{p=1}^{3}\left(\mathcal{S}_{\mathrm{even}}^{\alpha_p} + \frac{\sqrt{\lambda}}{2}\right)\right.$$
$$\left. + \sum_{\sigma\in\mathcal{Q}_3}\left(\mathcal{S}_{\mathrm{even}}^{\alpha_{\sigma(1)}} + \frac{\sqrt{\lambda}}{2}\right)\left(\mathcal{S}_{\mathrm{odd}}^{\alpha_{\sigma(2)}} + \frac{\sqrt{\lambda}}{2}\frac{I_2(\sqrt{\lambda})}{I_1(\sqrt{\lambda})}\right)\left(\mathcal{S}_{\mathrm{odd}}^{\alpha_{\sigma(3)}} + \frac{\sqrt{\lambda}}{2}\frac{I_2(\sqrt{\lambda})}{I_1(\sqrt{\lambda})}\right)\right]$$

$$\mathcal{S}_{\mathrm{even}}^{\alpha} = \sum_{k,\ell=1}^{\infty}\frac{I_{2k}(\sqrt{\lambda})}{I_1(\sqrt{\lambda})}\sqrt{(2k)(2\ell)}\left(\mathrm{D}_{2k,2\ell}^{(\alpha)} - \delta_{2k,2\ell}\right)$$

$$\mathcal{S}_{\mathrm{odd}}^{\alpha} = \sum_{k,\ell=1}^{\infty}\frac{I_{2k+1}(\sqrt{\lambda})}{I_1(\sqrt{\lambda})}\sqrt{(2k+1)(2\ell+1)}\left(\mathrm{D}_{2k+1,2\ell+1}^{(\alpha)} - \delta_{2k+1,2\ell+1}\right)$$

$$\mathcal{Q}_3 = \{(1,2,3),(3,1,2),(2,3,1)\}$$

$$\mathcal{Z}(m_1,m_2) = \int da_0 \int da_1 \mathrm{e}^{-\left(\mathrm{tr}a_0^2 + \mathrm{tr}a_1^2\right)}\mathrm{e}^{-S_{\mathrm{int}} + \left(m_1^2+m_2^2\right)\mathcal{M} + O(m^4)}$$

$$\mathcal{M} = -\sum_{n=1}^{\infty}\sum_{\ell=0}^{2n}(-1)^{n+\ell}\frac{(2n+1)!\,\zeta_{2n+1}}{(2n-\ell)!\,\ell!}\left(\frac{\lambda}{8\pi^2 N}\right)^n \mathrm{tr}a_0^\ell \mathrm{tr}a_1^{2n-\ell}$$

$$\mathcal{W} = 2\frac{\langle W_0\mathcal{M}\rangle - \langle W_0\rangle\langle\mathcal{M}\rangle}{\langle W_0\rangle}$$

$$\mathcal{W} = \mathcal{W}^{(L)} + \frac{\mathcal{W}^{(NL)}}{N^2} + O\left(\frac{1}{N^4}\right)$$

$$\frac{2\pi\sqrt{\lambda}}{I_1(\sqrt{\lambda})}\int_0^\infty \frac{dt}{t}\frac{(t/2)^2}{\sinh(t/2)^2}\frac{1}{4\pi^2+t^2}J_1\left(\frac{t\sqrt{\lambda}}{2\pi}\right)\left[2\pi I_0(\sqrt{\lambda})J_1\left(\frac{t\sqrt{\lambda}}{2\pi}\right) - tI_1(\sqrt{\lambda})J_0\left(\frac{t\sqrt{\lambda}}{2\pi}\right)\right].$$

$$\langle W_0\mathcal{M}\rangle - \langle W_0\rangle\langle\mathcal{M}\rangle \equiv \langle W_0\mathcal{M}\rangle_c \simeq \langle W_0\mathcal{M}\rangle_c^{(L)} + \frac{\langle W_0\mathcal{M}\rangle_c^{(NL)}}{N^2} + O(N^{-4})$$

$$\langle W_0\rangle \simeq W^{(L)} + \frac{W^{(NL)}}{N^2} + O(N^{-4})$$

$$W^{(L)} = \frac{2\sqrt{2}I_1(\sqrt{\lambda})}{\sqrt{\lambda}}$$

$$W^{(NL)} = \frac{\sqrt{2}}{48}\left(\lambda I_0(\sqrt{\lambda}) - 14\sqrt{\lambda}I_1(\sqrt{\lambda})\right) - \frac{\lambda^{3/2}\partial_\lambda\mathcal{F}}{2\sqrt{2}}I_1(\sqrt{\lambda})$$

$$\sum_{k,\ell=1}^{\infty} a_k \mathrm{D}_{k,\ell}^{(\alpha)} b_\ell$$

$$\mathrm{D}_{k,\ell}^{(\alpha)\,\mathrm{even}} = \delta_{k,\ell} - \int_0^\infty U_k^{\mathrm{even}}(t) Z_\ell^{(\alpha)}(t)$$

$$Z_\ell^{(\alpha)}(t) + s_\alpha \int_0^\infty dt' \mathbb{K}^{\mathrm{even}}(t,t') Z_\ell^{(\alpha)}(t') = s_\alpha U_\ell^{\mathrm{even}}(t)$$

$$\mathbb{K}^{\mathrm{even}}(t,t') = \sum_{k=1}^{\infty} U_k^{\mathrm{even}}(t) U_k^{\mathrm{even}}(t')$$

$$\sum_{k,\ell=1}^{\infty} a_k \mathrm{D}_{k,\ell}^{(\alpha)\,\mathrm{even}} b_\ell = \sum_{k=1}^{\infty} a_k b_k - \int_0^\infty dt A(t) Z^{(\alpha)}(t)$$

$$Z^{(\alpha)}(t) + s_\alpha \int_0^\infty dt' \mathbb{K}^{\mathrm{even}}(t,t') Z^{(\alpha)}(t') = s_\alpha B(t)$$

$$Z^{(\alpha)}(t) = \sum_{k=1}^{\infty} b_k Z_k^{(\alpha)}(t), A(t) = \sum_{k=1}^{\infty} a_k U_k^{\mathrm{even}}(t) \text{ and } B(t) = \sum_{k=1}^{\infty} b_k U_k^{\mathrm{even}}(t)$$

$$\int_0^\infty dx f(x) \approx \sum_{i=1}^{m} w_i f(x_i)$$

$$x_k = \cos\,\theta_k, \text{ with } \theta_k = (2k-1)\frac{\pi}{2m}, k = 1,2,\ldots,m$$

$$w_k = \frac{2}{m}\left[1 - 2\sum_{r=1}^{\frac{m-1}{2}} \frac{\cos(2r\theta_k)}{4r^2-1}\right]$$

$$1 + \Delta w^{(\alpha)}(M,\lambda) \underset{\lambda\to\infty}{\sim} \kappa_0\left(1 + \kappa_1\frac{1}{\sqrt{\lambda}} + \kappa_2\frac{1}{\lambda} + \kappa_3\frac{1}{\lambda^{3/2}} + O(\lambda^{-2})\right)$$

$$\mathcal{I}_n(s_\alpha) = \int_0^{+\infty} \frac{dz}{\pi} z^{1-2n} \partial_z \log\left(1 + s_\alpha \sinh\left(\frac{z}{2}\right)^{-2}\right)$$

$$\mathcal{I}_0(s_\alpha) = -\frac{\alpha}{M}\left(1 - \frac{\alpha}{M}\right) 2\pi$$

$$\mathcal{I}_1(s_\alpha) = -\frac{1}{2\pi}\left[\psi\left(\frac{\alpha}{M}\right) + \psi\left(1 - \frac{\alpha}{M}\right) - 2\psi(1)\right],$$

$$\Delta w^{(\alpha)}(M,\lambda) = \Delta w^{(\alpha)\ \text{even}}(M,\lambda) + \Delta w^{(\alpha)\ \text{odd}}(M,\lambda)$$

$$\Delta w^{(\alpha)\text{even}}(M,\lambda) = \frac{2}{\sqrt{\lambda}} \sum_{k,\ell=1}^{\infty} \frac{I_{2k}(\sqrt{\lambda}) I_{2\ell}(\sqrt{\lambda})}{I_1(\sqrt{\lambda}) I_2(\sqrt{\lambda})} \sqrt{2k}\sqrt{2\ell}\left(\mathrm{D}_{k,\ell}^{(\alpha)\text{even}} - \delta_{k,\ell}\right)$$

$$\Delta w^{(\alpha)\ \text{odd}}(M,\lambda) = \frac{2}{\sqrt{\lambda}} \sum_{k,\ell=1}^{\infty} \frac{I_{2k+1}(\sqrt{\lambda}) I_{2\ell+1}(\sqrt{\lambda})}{I_1(\sqrt{\lambda}) I_2(\sqrt{\lambda})} \sqrt{2k+1}\sqrt{2\ell+1}\left(\mathrm{D}_{k,\ell}^{(\alpha)\ \text{odd}} - \delta_{k,\ell}\right)$$

$$\Delta w^{(\alpha)} \underset{\lambda\to\infty}{\sim} \frac{2}{\sqrt{\lambda}} \sum_{P=0}^{\infty} \frac{\mathcal{S}^{(P)}}{\lambda^{P/2}}$$

$$\mathcal{S}^{(P)} = \sum_{L+J=P} S_{\text{odd}}^{(L,J)} + S_{\text{even}}^{(L,J)}$$

$$S_{\text{odd}}^{(L,J)} = \sum_{k=1}^{\infty} \sum_{\ell=1}^{\infty} \sqrt{2k+1} Q_{2L}^{(1)\text{odd}}(k) \sqrt{2\ell+1} Q_{2J}^{(2)\text{odd}}(\ell) \langle \psi_{2\ell+1} | \frac{s_\alpha \mathrm{X}}{1 - s_\alpha \mathrm{X}} | \psi_{2k+1} \rangle$$

$$S_{\text{even}}^{(L,J)} = \sum_{k=1}^{\infty} \sum_{\ell=1}^{\infty} \sqrt{2k} Q_{2L}^{(1)\text{even}}(k) \sqrt{2\ell} Q_{2J}^{(2)\text{even}}(\ell) \langle \psi_{2\ell} | \frac{s_\alpha \mathrm{X}}{1 - s_\alpha \mathrm{X}} | \psi_{2k} \rangle$$

$$\frac{I_{2k+1}(\sqrt{\lambda})}{I_j(\sqrt{\lambda})} \equiv \sum_{s=0}^{\infty} \frac{Q_{2s}^{(j)\ \text{odd}}(k)}{\lambda^{s/2}}, \frac{I_{2k}(\sqrt{\lambda})}{I_j(\sqrt{\lambda})} \equiv \sum_{s=0}^{\infty} \frac{Q_{2s}^{(j)\ \text{even}}(k)}{\lambda^{s/2}}$$

$$\psi_k(x) = (-1)^{\frac{k}{2}(k-1)} \sqrt{k} \frac{J_k(\sqrt{x})}{\sqrt{x}}$$

$$\mathrm{X}_{k,\ell} = \langle \psi_k | \mathrm{X} | \psi_\ell \rangle$$

$$w_{n,m}^{(\ell)}(s_\alpha) = \left\langle (x\partial_x)^n \phi^{(\ell)}(x) \right| \frac{s_\alpha \mathrm{X}}{1 - s_\alpha \mathrm{X}} \left| (x\partial_x)^m \phi^{(\ell)}(x) \right\rangle$$

$$\phi^{(\ell)}(x) \equiv J_\ell(\sqrt{x}) \text{ with } \ell = 1,2$$

$$\mathcal{S}^{(P)} = \mathcal{S}_0^{(P)} + \mathcal{S}_1^{(P)} + \mathcal{S}_2^{(P)} + \mathcal{S}_3^{(P)} + \cdots$$

$$\mathcal{S}_0^{(P)} = (-2)^{P-2} \sum_{n+m=P} \left[w_{n,m}^{(1)} + w_{n,m}^{(2)} \right]$$

$$\mathcal{S}_1^{(P)} = (-2)^{P-3} \sum_{n+m=P} \left[-(1+n^2+m^2) w_{n,m}^{(1)} + (5-n^2-m^2) w_{n,m}^{(2)} \right]$$

$$\mathcal{S}_2^{(P)} = (-2)^{P-5} \sum_{n+m=P} \left[f_{n,m}^{(1)} w_{n,m}^{(1)} + f_{n,m}^{(2)} w_{n,m}^{(2)} \right]$$

$$f_{n,m}^{(1)} = \frac{1}{3}(-3m^4 - 8m^3 - 6m^2n^2 - 12m^2 + 20m - 3n^4 - 8n^3 - 12n^2 + 20n + 15)$$

$$f_{n,m}^{(2)} = \frac{1}{3}(-3m^4 - 8m^3 - 6m^2n^2 + 24m^2 + 56m - 3n^4 - 8n^3 + 24n^2 + 56n + 51)$$

$$w_{n,m}^{(\ell)} = \omega_{n,m}^{(\ell,0)} g^{n+m+1} + \omega_{n,m}^{(\ell,1)} g^{n+m} + \omega_{n,m}^{(\ell,2)} g^{n+m-1} + \cdots$$

$$\frac{2}{\sqrt{\lambda}} \sum_{P=0}^{\infty} \sum_{n+m=P} \frac{1}{(-2\pi)^n (-2\pi)^m} \left[\frac{(n^2+m^2-2)\omega_{n,m}^{(0)}}{16\pi} + \frac{1}{4} \sum_{\ell=1}^{2} \omega_{n,m}^{(\ell,1)} \right]$$

$$\frac{1}{\sqrt{\lambda}} \left(\frac{1}{8\pi} \left((x\partial_x)^2 + \left(y\partial_y\right)^2 - 2 \right) G^{(0)}(s_\alpha, x, y) + \frac{1}{2} \sum_{\ell=1}^{2} G^{(1)}(\ell, s_\alpha, x, y) \right) \Bigg|_{x=y=-2\pi}$$

$$\frac{2}{s_\alpha \sqrt{\lambda}} \left(\frac{\mathcal{J}_0(s_\alpha)}{2} \right)^2$$

$$2\pi \sum_{\ell=1}^{2} G^{(2)}(\ell, s_\alpha, x, y) + \frac{1}{4} \left((x\partial_x)^2 + \left(y\partial_y\right)^2 \right) \sum_{\ell=1}^{2} G^{(1)}(\ell, s_\alpha, x, y)$$
$$+ \frac{1}{4} \left(G^{(1)}(1, s_\alpha, x, y) - 5G^{(1)}(2, s_\alpha, x, y) \right) - \frac{1}{64\pi} \Big(22 - \frac{76}{3} \left(x\partial_x + y\partial_y \right) +$$
$$4 \left((x\partial_x)^2 + \left(y\partial_y\right)^2 \right) + \frac{16}{3} \left((x\partial_x)^3 + \left(y\partial_y\right)^3 \right) - 4(x\partial_x)^2 \left(y\partial_y\right)^2 - 2 \left((x\partial_x)^4 + \left(y\partial_y\right)^4 \right) \Big) G^{(0)}(s_\alpha, x, y)$$

$$\frac{1}{s_\alpha} \left(\frac{\mathcal{J}_0(s_\alpha)}{2} \right)^2 \left(3 - \frac{\pi \mathcal{J}_1(s_\alpha)}{2} \right)$$

$$\Delta w^{(\alpha)\text{ even}} = -\frac{2}{\sqrt{\lambda}} \frac{1}{I_1(\sqrt{\lambda}) I_2(\sqrt{\lambda})} \int_0^\infty dt A(t) Z^{(\alpha)}(t)$$

$$Z^{(\alpha)}(t) + s_\alpha \int_0^\infty du \mathbb{K}^{\text{even}}(t,u) Z^{(\alpha)}(u) = s_\alpha B(t)$$

$$A(t) = B(t) = \frac{\sqrt{2t}\pi}{(4\pi^2 + t^2)\sinh(t/2)} \left(\frac{t\sqrt{\lambda}}{2\pi} I_1(\sqrt{\lambda}) J_0\left(\frac{t\sqrt{\lambda}}{2\pi} \right) - \sqrt{\lambda} I_0(\sqrt{\lambda}) J_1\left(\frac{t\sqrt{\lambda}}{2\pi} \right) \right),$$

$$\Delta w^{(\alpha)\text{ even}} = -\frac{3\zeta_3}{32\pi^4} s_\alpha \lambda^2 + \frac{\pi^2 \zeta_3 + 20\zeta_5}{256\pi^6} s_\alpha \lambda^3 + O(\lambda^4)$$

$$1+\Delta w^{(1)} \underset{\lambda\to\infty}{\sim} \kappa_0\left(1+\kappa_1\frac{1}{\sqrt{\lambda}}+\kappa_2\frac{1}{\lambda}+\kappa_3\frac{1}{\lambda^{3/2}}+O(\lambda^{-2})\right)$$

$$\tilde{f}_{\mathrm{num}}(\lambda)=\left(\frac{f_{\mathrm{num}}(\lambda)}{\kappa_0}-1\right)\sqrt{\lambda}$$

$$\sum_{i=0}^{2} g_i \mathcal{I}_1(s_\alpha)^i$$

$$\mathcal{S}^{\alpha}_{\mathrm{even}}=c_0^{\mathrm{even}}\sqrt{\lambda}+c_1^{\mathrm{even}}+c_2^{\mathrm{even}}\frac{1}{\sqrt{\lambda}}+c_3^{\mathrm{even}}\frac{1}{\lambda}+O\left(\frac{1}{\lambda^{3/2}}\right)$$

$$c_0^{\mathrm{even}} = -\frac{1}{2}-\frac{1}{4}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}$$

$$c_1^{\mathrm{even}} = -\frac{1}{8}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}$$

$$c_2^{\mathrm{even}} = -\frac{3}{32}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}-\frac{\pi}{16}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}\mathcal{I}_1(s_\alpha)$$

$$c_3^{\mathrm{even}} = -\frac{3}{32}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}-\frac{3\pi}{32}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}\mathcal{I}_1(s_\alpha)-\frac{\pi^2}{8}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}\mathcal{I}_1(s_\alpha)^2$$

$$\mathcal{S}^{\alpha}_{\mathrm{odd}}=c_0^{\mathrm{odd}}\sqrt{\lambda}+c_1^{\mathrm{odd}}+c_2^{\mathrm{odd}}\frac{1}{\sqrt{\lambda}}+c_3^{\mathrm{odd}}\frac{1}{\lambda}+O\left(\frac{1}{\lambda^{3/2}}\right)$$

$$c_0^{\mathrm{odd}} = -\frac{1}{2}-\frac{1}{4}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}$$

$$c_1^{\mathrm{odd}} = \frac{3}{4}$$

$$c_2^{\mathrm{odd}} = -\frac{3}{16}+\frac{3\pi}{16}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}\mathcal{I}_1(s_\alpha)$$

$$c_3^{\mathrm{odd}} = -\frac{3}{16}+\frac{3\pi}{16}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}\mathcal{I}_1(s_\alpha)+\frac{3\pi^2}{8}\frac{\mathcal{I}_0(s_\alpha)}{\sqrt{s_\alpha}}\mathcal{I}_1(s_\alpha)^2$$

$$1+\Delta w^{(\alpha,\beta)}(M,\lambda) \underset{\lambda\to\infty}{\sim} c_0\left[1+c_1\frac{1}{\sqrt{\lambda}}+c_2\frac{1}{\lambda}+c_3\frac{1}{\lambda^{3/2}}+O\left(\frac{1}{\lambda^2}\right)\right]$$

$$c_0 = -\frac{1}{8}\prod_{p=1}^{3}\frac{\mathcal{J}_0\left(s_{\alpha_p}\right)}{\sqrt{s_{\alpha_p}}}$$

$$c_1 = 3$$

$$c_2 = \frac{1}{4}\left(21 - \pi\sum_{p=1}^{3}\mathcal{J}_1\left(s_{\alpha_p}\right)\right)$$

$$c_3 = \frac{1}{32}\left(199 - 28\pi\sum_{p=1}^{3}\mathcal{J}_1\left(s_{\alpha_p}\right) - 16\pi^2\sum_{p=1}^{3}\mathcal{J}_1\left(s_{\alpha_p}\right)^2\right)$$

$$\mathcal{S}^{\alpha}_{\text{even}} = \sum_{P=0}^{\infty}\frac{\mathcal{S}^{(P)}_{\text{even}}}{\lambda^{P/2}}, \mathcal{S}^{\alpha}_{\text{odd}} = \sum_{P=0}^{\infty}\frac{\mathcal{S}^{(P)}_{\text{odd}}}{\lambda^{P/2}}$$

$$\mathcal{S}^{(P)}_{\text{even}} = \mathcal{S}^{(P)}_{0,\,\text{even}} + \mathcal{S}^{(P)}_{1,\,\text{even}} + \mathcal{S}^{(P)}_{2,\,\text{even}} + \cdots$$
$$\mathcal{S}^{(P)}_{\text{odd}} = \mathcal{S}^{(P)}_{0,\,\text{odd}} + \mathcal{S}^{(P)}_{1,\,\text{odd}} + \mathcal{S}^{(P)}_{2,\,\text{odd}} + \cdots$$

$$\mathcal{S}^{(P)}_{0,\,\text{even}} = (-2)^{P-2}w^{(1)}_{P,0}$$
$$\mathcal{S}^{(P)}_{1,\,\text{even}} = -(-2)^{P-4}(P^2 - 2P)\left(1 - \delta_{P,0}\right)w^{(1)}_{P-1,0}$$
$$\mathcal{S}^{(P)}_{2,\,\text{even}} = -\frac{(-2)^{P-7}}{3}(3P^4 - 16P^3 + 24P^2 - 20P + 9)\left(1 - \delta_{P,0}\right)\left(1 - \delta_{P,1}\right)w^{(1)}_{P-2,0}$$

$$\mathcal{S}^{(P)}_{0,\,\text{odd}} = (-2)^{P-2}w^{(2)}_{P,0}$$
$$\mathcal{S}^{(P)}_{1,\,\text{odd}} = -(-2)^{P-4}(P^2 - 2P - 3)\left(1 - \delta_{P,0}\right)w^{(2)}_{P-1,0}$$
$$\mathcal{S}^{(P)}_{2,\,\text{odd}} = -\frac{(-2)^{P-7}}{3}P(3P^3 - 16P^2 + 6P + 16)\left(1 - \delta_{P,0}\right)\left(1 - \delta_{P,1}\right)w^{(2)}_{P-2,0}$$

$$c_1^{\text{even}} = \frac{1}{4}\sum_{P=0}^{\infty}\frac{\omega^{(1)}_{P,0}}{(-2\pi)^P} - \frac{1}{16}\sum_{P=0}^{\infty}(P^2 - 2P)\frac{\omega^{(1)}_{P-1,0}}{(-2\pi)^P}\left(1 - \delta_{P,0}\right)$$

$$c_1^{\text{even}} = \left(\frac{1}{4}G^{(1)}(s_\alpha, 1, x) + \frac{1}{32\pi}[(x\partial_x)^2 - 1]G^{(0)}(s_\alpha, x)\right)\Big|_{x=-2\pi} = -\frac{1}{8}\frac{\mathcal{J}_0(s_\alpha)}{\sqrt{s_\alpha}}$$

$$c_1^{\text{odd}} = \left(\frac{1}{4}G^{(1)}(s_\alpha, 2, x) + \frac{1}{32\pi}[(x\partial_x)^2 - 4]G^{(0)}(s_\alpha, x)\right)\Big|_{x=-2\pi} = \frac{3}{4}$$

$$c_2^{\text{even}} = \Big(\pi G^{(2)}(s_\alpha,1,x) + \frac{1}{8}((x\partial_x)^2-1)G^{(1)}(s_\alpha,1,x) + \frac{1}{384\pi}(3(x\partial_x)^4 - 8(x\partial_x)^3 + 20(x\partial_x) - 15)\, G^{(0)}(s_\alpha,x)\Big)\Big|_{x=-2\pi}$$
$$= -\frac{3}{32}\frac{\mathcal{J}_0(s_\alpha)}{\sqrt{s_\alpha}} - \frac{\pi}{16}\frac{\mathcal{J}_0(s_\alpha)}{\sqrt{s_\alpha}}\mathcal{J}_1(s_\alpha)$$
$$c_2^{\text{odd}} = \Big(\pi G^{(2)}(s_\alpha,2,x) + \frac{1}{8}((x\partial_x)^2-4)G^{(1)}(s_\alpha,2,x) + \frac{1}{384\pi}(3(x\partial_x)^4 - 8(x\partial_x)^3 - 18(x\partial_x)^2 + 56(x\partial_x) - 24)G^{(0)}(s_\alpha,x)\Big)\Big|_{x=-2\pi}$$
$$= -\frac{3}{16} + \frac{3\pi}{16}\frac{\mathcal{J}_0(s_\alpha)}{\sqrt{s_\alpha}}\mathcal{J}_1(s_\alpha)$$

$$\mathcal{S}^{\alpha}_{\text{even}} = -\frac{1}{I_1(\sqrt{\lambda})}\int_0^\infty dt A(t) Z^{(\alpha)}(t)$$

$$Z^{(\alpha)}(t) + s_\alpha \int_0^\infty du \mathbb{K}^{\text{even}}(t,u) Z^{(\alpha)}(u) = -\frac{s_\alpha\sqrt{t\lambda}}{2\sqrt{2}\pi \sinh(t/2)} J_1\left(\frac{t\sqrt{\lambda}}{2\pi}\right)$$

$$c_3^{\text{even}} = \frac{\mathcal{J}_0(s_\alpha)}{32\sqrt{s_\alpha}}\sum_{i=0}^{2} c_{3,i}^{\text{even}} \left(\pi\mathcal{J}_1(s_\alpha)\right)^i$$
$$c_3^{\text{odd}} = \frac{c_{3,0}^{\text{odd}}}{16} + \frac{\mathcal{J}_0(s_\alpha)}{16\sqrt{s_\alpha}}\sum_{i=1}^{2} c_{3,i}^{\text{odd}} \left(\pi\mathcal{J}_1(s_\alpha)\right)^i$$

$$\frac{\langle W_1 W_1 \dots W_1\rangle}{\left(W^{(2)}_{conn}(\lambda)\right)^n} \simeq \left(1 + \Delta w^{(1)}(2,\lambda)\right)^n$$

$$\frac{\langle W_1 W_1 \dots W_1\rangle}{\left(W^{(2)}_{\text{conn}}(\lambda)\right)^n} \underset{\lambda\to\infty}{\sim} \left(\frac{\pi^2}{16}\right)^n \left(1 + \frac{2n}{\sqrt{\lambda}} + \frac{n(2n+1-\log 2)}{\lambda} + \frac{n(16n^2 + 24n(1-\log 2) + 5 - 12\log 2(1+4\log 2)}{12\lambda^{3/2}} + O(\lambda^{-2})\right)$$

$$1 + \Delta w^{(\alpha_1,\dots,\alpha_{2n-1})}(M,\lambda) \equiv \frac{\langle W_{\alpha_1} W_{\alpha_2} \dots W_{\alpha_{2n-1}} W^{\dagger}_{\alpha_{2n}}\rangle}{\mathcal{N}^{\text{even}}(\alpha_1,\alpha_2,\dots,\alpha_{2n})\left(W^{(2)}_{\text{conn}}(\lambda)\right)^n}$$

$$1 + \Delta w^{(\alpha_1,\dots,\alpha_{2n-1})}(M,\lambda) \simeq \frac{1}{\mathcal{N}^{\text{even}}(\alpha_1,\alpha_2,\dots,\alpha_{2n})}\left[\prod_{j=1}^{n} \delta_{\alpha_{2j-1},\alpha_{2j}}\left(1 + \Delta w^{(\alpha_{2j-1})}(M,\lambda)\right) + \text{ permutations }\right].$$

$$1 + \Delta w^{(\alpha_1,\dots,\alpha_{2n+1})}(M,\lambda) \equiv \frac{\langle W_{\alpha_1} W_{\alpha_2} \dots W_{\alpha_{2n}} W^{\dagger}_{\alpha_{2n+1}}\rangle}{\mathcal{N}^{odd}(\alpha_1,\alpha_2,\dots,\alpha_{2n+1})\sqrt{M} W^{(3)}_{conn}(\lambda)\left(W^{(2)}_{conn}(\lambda)\right)^{n-1}}$$

$$1+\Delta w^{(\alpha_1,\alpha_2,\ldots,\alpha_{2n})}(M,\lambda)\simeq\frac{1}{\mathcal{N}^{\text{odd}}(\alpha_1,\alpha_2,\cdots,\alpha_{2n+1})}$$

$$\left[\delta_{\alpha_1+\alpha_2,\alpha_3}\left(1+\Delta w^{(\alpha_1,\alpha_2)}(M,\lambda)\right)\prod_{j=2}^{n}\delta_{\alpha_{2j},\alpha_{2j+1}}\left(1+\Delta w^{(\alpha_{2j})}(M,\lambda)\right)+\text{ permutations}\right]$$

$$\mathcal{W}^{(NL)}=2\frac{\langle W_0\mathcal{M}\rangle_c^{(NL)}}{W^{(L)}}-\mathcal{W}^{(L)}\frac{W^{(NL)}}{W^{(L)}}$$

$$\langle W_0\mathcal{M}\rangle_c^{(NL)}=\left\langle W_0\mathcal{M}^{(1)}\right\rangle_c^{(NL)}+\left\langle W_0\mathcal{M}^{(2)}\right\rangle_c^{(NL)}$$

$$\left\langle W_0\mathcal{M}^{(1)}\right\rangle_c^{(NL)}\underset{\lambda\to\infty}{\sim}-\frac{\lambda^2}{768}-\frac{\lambda^{3/2}}{4608}(4\pi^2+45)+\frac{\lambda}{2048}(24\pi^2-115-16\log(2))+O(\sqrt{\lambda})$$

$$\frac{I_1(\sqrt{\lambda})}{4\sqrt{2}}\sum_{k,\ell=2}^{\infty}(-)^{k-k\ell}\mathrm{M}_{k,\ell}\left(\sqrt{k\ell}-\mathrm{d}_k\,\mathrm{d}_\ell\right)-(\lambda\partial_\lambda\mathcal{F})\sum_{q,p=1}^{\infty}\sqrt{p}\mathrm{M}_{2p,2q}\sqrt{2q}I_{2q}(\sqrt{\lambda})$$

$$\mathrm{M}_{k,\ell}=(-1)^{\frac{k+\ell+2k\ell}{2}}+1\sqrt{k\ell}\int_0^\infty\frac{dt}{t}\frac{(t/2)^2}{\sinh(t/2)^2}J_k\left(\frac{t\sqrt{\lambda}}{2\pi}\right)J_\ell\left(\frac{t\sqrt{\lambda}}{2\pi}\right)$$

$$\mathcal{R}=\sum_{k,\ell=2}^{\infty}(-)^{k-k\ell}\mathrm{M}_{k,\ell}\left(\sqrt{k\ell}-\mathrm{d}_k\,\mathrm{d}_\ell\right)$$

$$\mathcal{W}^{(NL)}\underset{\lambda\to\infty}{\sim}-\frac{\lambda^{3/2}}{128}+\frac{\sqrt{\lambda}}{512}(8\log(2)-1)+\frac{1}{256}(2\zeta_3+32\log^2(2)-1)+O\left(\lambda^{-1/2}\right)$$

$$\mathcal{R}(\lambda)=-\frac{\sqrt{\lambda}}{24}+\frac{5}{8}\frac{1}{\sqrt{\lambda}}+\frac{\log(2)}{2\lambda}+O\left(\frac{1}{\lambda^{3/2}}\right)$$

$$\mathcal{R}_1(\lambda)=\sum_{k,\ell=1}^{\infty}\mathrm{M}_{2k,2\ell}\left(\sqrt{(2k)(2\ell)}-\mathrm{d}_{2k}\,\mathrm{d}_{2\ell}\right)$$

$$\mathcal{R}_2(\lambda)=\sum_{k,\ell=1}^{\infty}\mathrm{M}_{2k,2\ell+1}\left(\sqrt{2k(2\ell+1)}-\mathrm{d}_{2k}\,\mathrm{d}_{2\ell+1}\right)$$

$$\mathcal{R}_3(\lambda)=-\sum_{k,\ell=1}^{\infty}\mathrm{M}_{2k+1,2\ell}\left(\sqrt{(2k+1)2\ell}-\mathrm{d}_{2k+1}\,\mathrm{d}_{2\ell}\right)$$

$$\mathcal{R}_4(\lambda)=\sum_{k,\ell=1}^{\infty}\mathrm{M}_{2k+1,2\ell+1}\left(\sqrt{(2k+1)(2\ell+1)}-\mathrm{d}_{2k+1}\,\mathrm{d}_{2\ell+1}\right)$$

$$\mathrm{d}_{2k}=\sqrt{2k}-\int_0^\infty dtU_k^{\text{even}}(t)Z(t)$$

$$Z(t)+\int_0^\infty du\mathbb{K}^{\text{even}}(t,u)Z(u)=-\frac{\sqrt{t\lambda}}{2\sqrt{2}\pi\sinh(t/2)}J_1\left(\frac{t\sqrt{\lambda}}{2\pi}\right)$$

$$\mathcal{R}_1(\lambda) = 2\sum_{k,\ell=1}^{\infty} \sqrt{2k}\mathrm{M}_{2k,2\ell}\int_0^{\infty} dt U_\ell^{\mathrm{even}}(t)Z(t) - \sum_{k,\ell=1}^{\infty} \mathrm{M}_{2k,2\ell}\int_0^{\infty} dt \int_0^{\infty} dt' U_k^{\mathrm{even}}(t)U_\ell^{\mathrm{even}}(t')Z(t)Z(t')$$

$$\mathcal{R}(\lambda) \underset{\lambda\to\infty}{\sim} \sqrt{\lambda}\tilde{\mathcal{R}}_1 + \tilde{\mathcal{R}}_2 + \frac{\tilde{\mathcal{R}}_3}{\sqrt{\lambda}} + \frac{\tilde{\mathcal{R}}_4}{\lambda} + O\left(\frac{1}{\lambda^{3/2}}\right).$$

$$\lambda \mapsto \lambda' \equiv \lambda - 4\pi\mathcal{I}_1(s_\alpha)\sqrt{\lambda} + 4\pi^2\mathcal{I}_1(s_\alpha)^2$$

$$G(s_\alpha,\ell,x) = \sum_{n=0}^{\infty} \frac{w_{0,n}^{(\ell)}}{(gx)^n} = gG^{(0)}(s_\alpha,x) + G^{(1)}(s_\alpha,\ell,x) + g^{-1}G^{(2)}(s_\alpha,\ell,x) + \cdots$$

$$G(s_\alpha,\ell,x,y) = \sum_{n=0}^{\infty}\sum_{m=0}^{\infty} \frac{w_{n,m}^{(\ell)}}{(gx)^n(gy)^m} = gG^{(0)}(s_\alpha,x,y) + G^{(1)}(s_\alpha,\ell,x,y) + \cdots$$

$$w_{0,n}^{(\ell)} = \sum_{i\geq 0} \omega_{0,n}^{(\ell,i)} g^{n+1-i}$$

$$w_{0,0}^{(\ell)} = 4g\mathcal{I}_0(s_\alpha) + (2\ell-1) - \frac{(2\ell-1)(2\ell-3)\mathcal{I}_1(s_\alpha)}{8g} - \frac{(2\ell-1)(2\ell-3)\mathcal{I}_1(s_\alpha)^2}{16g^2} - \frac{(2\ell-1)(2\ell-3)(16\mathcal{I}_1(s_\alpha)^3 - 5\mathcal{I}_2(s_\alpha) - 8\mathcal{I}_2(s_\alpha)\ell + 4\mathcal{I}_2(s_\alpha)\ell^2)}{512g^3} + O\left(\frac{1}{g^4}\right)$$

$$\omega_{0,2n+1}^{(\ell,0)}(s_\alpha) = -\frac{1}{2(n+1)}\sum_{j=0}^{n} (-1)^j \mathcal{I}_{-j}(s_\alpha)\omega_{0,2n-2j}^{(\ell,0)}(s_\alpha)$$

$$\omega_{0,2n}^{(\ell,0)}(s_\alpha) = \frac{1}{2n+1}\left(\sum_{j=1}^{n} (-1)^{n-j-1}\mathcal{I}_{-n+j}(s_\alpha)\omega_{0,2j-1}^{(\ell,0)}(s_\alpha) + 4(-1)^n\mathcal{I}_{-n}(s_\alpha)\right)$$

$$\omega_{0,0}^{(\ell,1)} = 2\ell - 1$$

$$\omega_{0,1}^{(\ell,1)} = -\left(\omega_{0,0}^{(\ell,1)} + 1\right)\mathcal{I}_0(s_\alpha)$$

$$\omega_{0,2}^{(\ell,1)} = -\left[\frac{\omega_{0,0}^{(\ell,0)}}{8} - \frac{1}{2}\left(\omega_{0,0}^{(\ell,1)} + 1\right)\mathcal{I}_0(s_\alpha)\right]\mathcal{I}_0(s_\alpha)$$

$$\omega_{0,n}^{(\ell,1)} = \begin{cases} 2\ell - 1, & n = 0 \\ \left[-\frac{\ell}{2} + \left(\frac{n-1}{2}\right)^2\right]\omega_{0,n-1}^{(\ell,0)}, & n \geq 1 \end{cases}$$

$$G^{(1)}(s_\alpha,\ell,x) = \sum_{n=0}^{\infty} \frac{\omega_{0,n}^{(\ell,1)}}{x^n} = \omega_{0,0}^{(\ell,1)} + \frac{1}{4x}\sum_{n=1}^{\infty} [(n-1)^2 - 2\ell]\frac{\omega_{0,n-1}^{(\ell,0)}}{x^{n-1}}$$
$$= 2\ell - 1 + \frac{1}{4x}[-2\ell + (x\partial_x)^2]G^{(0)}(s_\alpha,x)$$

$$\omega_{0,n}^{(\ell,2)}=\begin{cases}-\dfrac{(2\ell-1)(2\ell-3)\mathcal{I}_1(s_\alpha)}{8}, & n=0,\\ \left(\dfrac{3}{8}-\ell+\dfrac{\ell^2}{2}\right)\mathcal{I}_0(s_\alpha)\mathcal{I}_1(s_\alpha)-\dfrac{\ell^2}{2}-\dfrac{\ell}{2}+\dfrac{3}{8}, & n=1,\\ \left(\dfrac{3}{4}-2\ell+\ell^2\right)\dfrac{\mathcal{I}_1(s_\alpha)\omega_{0,n-1}^{(\ell,0)}}{8}+\left(n\ell^2-n(n-2)\ell+\dfrac{n(n-2)(20-14n+3n^2)}{12}\right)\dfrac{\omega_{0,n-2}^{(\ell,0)}}{8}, & n\geq 2.\end{cases}$$

$$G^{(2)}(s_\alpha,\ell,x)=\sum_{n=0}^{\infty}\frac{\omega_{0,n}^{(\ell,2)}}{x^n}=-\frac{(2\ell-1)(2\ell-3)\mathcal{I}_1(s_\alpha)}{8}+\frac{1}{x}\left(\frac{3}{8}-\frac{\ell}{2}-\frac{\ell^2}{2}\right)+\left(\frac{3}{4}-2\ell+\ell^2\right)\frac{\mathcal{I}_1(s_\alpha)}{8x}G^{(0)}(s_\alpha,x)+\frac{1}{8x^2}[\ell^2(2-x\partial_x)-\ell((x\partial_x)^2-2x\partial_x)+\frac{1}{4}(x\partial_x)^4-\frac{1}{3}(x\partial_x)^3-\frac{2}{3}(x\partial_x)\Big]G^{(0)}(s_\alpha,x)$$

$$(gx+gy+1)G(s_\alpha,\ell,x,y)+\frac{1}{2}\big(x\partial_x+y\partial_y-g\partial_g\big)G(s_\alpha,\ell,x,y)=-\frac{1}{4}G(s_\alpha,\ell,x)G(s_\alpha,\ell,y)+gxG(s_\alpha,\ell,y)+gyG(s_\alpha,\ell,x)$$

$$(x+y)G^{(0)}(s_\alpha,x,y)=-\frac{1}{4}G^{(0)}(s_\alpha,x)G^{(0)}(s_\alpha,y)+xG^{(0)}(s_\alpha,y)+yG^{(0)}(s_\alpha,x)$$

$$4(x+y)G^{(1)}(s_\alpha,x,y)=\Big(4y-G^{(0)}(s_\alpha,y)\Big)G^{(1)}(s_\alpha,\ell,x)+\Big(4x-G^{(0)}(s_\alpha,x)\Big)G^{(1)}(s_\alpha,\ell,y)-2\Big(G^{(0)}(s_\alpha,x,y)+x\partial_xG^{(0)}(s_\alpha,x,y)+y\partial_yG^{(0)}(s_\alpha,x,y)\Big)$$

$$G^{(1)}(s_\alpha,\ell,x,y)=-\frac{xG^{(0)}(s_\alpha,y)\partial_x^2G^{(0)}(s_\alpha,x)}{16(x+y)}+\frac{xy\partial_x^2G^{(0)}(s_\alpha,x)}{4(x+y)}+\frac{xy\partial_y^2G^{(0)}(s_\alpha,y)}{4(x+y)}-\frac{yG^{(0)}(s_\alpha,x)\partial_y^2G^{(0)}(s_\alpha,y)}{16(x+y)}+\frac{xG^{(0)}(s_\alpha,y)\partial_xG^{(0)}(s_\alpha,x)}{8(x+y)^2}-\frac{xy\partial_xG^{(0)}(s_\alpha,x)}{2(x+y)^2}+\frac{x\partial_yG^{(0)}(s_\alpha,y)}{4(x+y)}-\frac{xy\partial_yG^{(0)}(s_\alpha,y)}{2(x+y)^2}-\frac{G^{(0)}(s_\alpha,y)\partial_xG^{(0)}(s_\alpha,x)}{16(x+y)}+\frac{y\partial_xG^{(0)}(s_\alpha,x)}{4(x+y)}-\frac{G^{(0)}(s_\alpha,x)\partial_yG^{(0)}(s_\alpha,y)}{16(x+y)}+\frac{yG^{(0)}(s_\alpha,x)\partial_yG^{(0)}(s_\alpha,y)}{8(x+y)^2}-\frac{\ell xG^{(0)}(s_\alpha y)}{2y(x+y)}-\frac{\ell G^{(0)}(s_\alpha,x)}{2(x+y)}+\frac{\ell G^{(0)}(s_\alpha,x)G^{(0)}(s_\alpha,y)}{8y(x+y)}-\frac{\ell G^{(0)}(s_\alpha,y)}{2(x+y)}-\frac{\ell yG^{(0)}(s_\alpha,x)}{2x(x+y)}+\frac{\ell G^{(0)}(s_\alpha,x)G^{(0)}(s_\alpha,y)}{8x(x+y)}+\frac{x^2G^{(0)}(s_\alpha,y)}{2(x+y)^3}+\frac{y^2G^{(0)}(s_\alpha,x)}{2(x+y)^3}+\frac{xyG^{(0)}(s_\alpha,x)}{2(x+y)^3}-\frac{xG^{(0)}(s_\alpha,x)G^{(0)}(s_\alpha,y)}{8(x+y)^3}-\frac{xG^{(0)}(s_\alpha,y)}{(x+y)^2}+\frac{xyG^{(0)}(s_\alpha,y)}{2(x+y)^3}+\frac{G^{(0)}(s_\alpha,x)}{4(x+y)}-\frac{yG^{(0)}(s_\alpha,x)}{(x+y)^2}+\frac{G^{(0)}(s_\alpha,x)G^{(0)}(s_\alpha,y)}{8(x+y)^2}-\frac{yG^0(s_\alpha,x)G^{(0)}(s_\alpha,y)}{8(x+y)^3}+\frac{G^{(0)}(s_\alpha,y)}{4(x+y)}+\frac{2\ell x}{x+y}+\frac{2\ell y}{x+y}-\frac{x}{x+y}-\frac{y}{x+y}.$$

$$-4(x+y)G^{(2)}(s_\alpha,\ell,x,y)=G^{(1)}(s_\alpha,\ell,x)G^{(1)}(s_\alpha,\ell,y)+\big(G^{(0)}(s_\alpha,x)-4x\big)G^{(2)}(s_\alpha,\ell,y)+\big(G^{(0)}(s_\alpha,y)-4y\big)G^{(2)}(s_\alpha,\ell,x)+4G^{(1)}(s_\alpha,\ell,x,y)+2x\partial_xG^{(1)}(s_\alpha,\ell,x,y)+2y\partial_yG^{(1)}(s_\alpha,\ell,x,y)$$

$$S^{\mathrm{wl}}=-\frac{m}{2}\int d\tau g_{\mu\nu}\dot{x}^\mu\dot{x}^\nu+\cdots$$

$$S^{\mathrm{EH}}=\frac{1}{16\pi G}\int\ d^4x\sqrt{-g}R$$

$$x^{\mu}(\tau)=b^{\mu}+u^{\mu}\tau+z^{\mu}(\tau),g_{\mu\nu}=\eta_{\mu\nu}+\kappa h_{\mu\nu}$$

$$\mathcal{A}\overset{k^2\to 0}{\to}i\frac{\mathcal{A}_L\mathcal{A}_R}{k^2}$$

$$\square\ h_{\mu\nu}(x)=0,k^2=0$$

$$\mathcal{A}\overset{\omega\to 0}{\to}-i\frac{\mathcal{A}_L\mathcal{A}_R}{m\omega^2}+\frac{\text{Ch}}{\omega}$$

$$\delta(x)=2\pi\delta(x),\int_k=\int\ \frac{d^Dk}{(2\pi)^D}$$

$$\langle\mathcal{T}\mathcal{O}(h_i,z_j)\rangle=\int\ \mathrm{Dh}Dz\mathcal{O}(h_i,z_j)e^{iS}$$

$$G_h^F(k)=\frac{i\Pi^{\mu\nu\alpha\beta}}{k^2+i\epsilon},G_h^F(\omega)=\frac{-i\eta^{\mu\nu}}{\omega^2+i\epsilon}$$

$$\Pi^{\mu\nu\alpha\beta}=\frac{1}{2}\left(\eta^{\mu\alpha}\eta^{\nu\beta}+\eta^{\mu\beta}\eta^{\nu\alpha}-\frac{1}{D-2}\eta^{\mu\nu}\eta^{\alpha\beta}\right)$$

$$\begin{aligned}&A(h_1,\cdots,h_n,z_1,\cdots z_m)=\mathrm{LSZ}\langle\mathcal{T}h_1\cdots h_nz_1\cdots z_m\rangle\\&=\lim_{k_i^2,\omega_j^2\to 0}\prod_a(-ik_a^2)\prod_b\left(im_b\omega_b^2\right)\langle h_1\cdots h_nz_1\cdots z_m\rangle\end{aligned}$$

$$h_i=\varepsilon_i^{\mu\nu}h_{\mu\nu}(k_i),z_i=\zeta_i^{\mu}z_{\mu}(\omega_i)$$

$$A(h_1,\cdots,h_n,z_1,\cdots z_m)=\int_{q_1,\ldots,q_m}\delta^{(4)}\left(\sum_i^n\ k_i-\sum_j^m\ q_j\right)\left(\prod_j^m\ \delta(q_j\cdot u_j)e^{iq_j\cdot b_j}\right)\mathcal{A}(q_i,k_i,u_i)$$

$$\int\ d\tau e^{iq_j\cdot\bar{x}_j(\tau)}=\delta(q_j\cdot u_j)e^{iq_j\cdot b_j}$$

$$\delta^{(4)}\left(\sum_i^n\ k_i-\sum_j^m\ q_j\right)$$

$$\langle\mathcal{O}(h_i,z_j)\rangle=\int\ Dh^{\mathrm{I}}Dh^{\mathrm{II}}Dz^{\mathrm{I}}Dz^{\mathrm{II}}\mathcal{O}(h_i,z_j)e^{iS^{\mathrm{I}}-iS^{\mathrm{II}}}$$

$$h^{\mathrm{I}}(t=+\infty)=h^{\mathrm{II}}(t=+\infty),z^{\mathrm{I}}(t=+\infty)=z^{\mathrm{II}}(t=+\infty).$$

$$\mathbf{G}=\begin{pmatrix}G_{\mathrm{I,I}}&G_{\mathrm{I,II}}\\G_{\mathrm{II,I}}&G_{\mathrm{II,II}}\end{pmatrix}=\begin{pmatrix}G^F&G^>\\G^<&G^{\bar{F}}\end{pmatrix}.$$

$$\mathbf{G}_h(k) = \begin{pmatrix} \frac{i}{k^2+i\epsilon} & \theta(k^0)\delta(k^2) \\ \theta(-k^0)\delta(k^2) & \frac{-i}{k^2-i\epsilon} \end{pmatrix} \Pi^{\mu\nu\alpha\beta}\,.$$

$$\mathbf{G}_z(\omega) = \begin{pmatrix} \frac{-i}{\omega^2+i\epsilon} & \frac{1}{2}\delta'(\omega) \\ \frac{1}{2}\delta'(\omega) & \frac{i}{\omega^2-i\epsilon} \end{pmatrix} \eta^{\mu\nu}.$$

$$\mathcal{A} = \sum_i \frac{N_i}{\prod_{\alpha_i} D_{\alpha_i}}$$

$$\lim_{k^2\to 0} k^2\mathcal{A} = i\sum_{\text{pol.}} \mathcal{A}_L(k,\varepsilon)\mathcal{A}_R(-k,\varepsilon^*),$$
$$\lim_{\omega^2\to 0} m\omega^2\mathcal{A} = -i\sum_{\text{pol.}} \mathcal{A}_L(\omega,\zeta)\mathcal{A}_R(-\omega,\zeta^*),$$

$$\sum_{\text{pol}} \varepsilon^{*\mu}(-k)\varepsilon^{*\nu}(-k)\varepsilon_\alpha(k)\varepsilon_\beta(k) = \frac{1}{2}\left(P^{\mu\alpha}P^{\nu\beta} + P^{\mu\beta}P^{\nu\alpha} - \frac{1}{D-2}P^{\mu\nu}P^{\alpha\beta}\right)$$


$$P^{\mu\nu}(k) = \eta^{\mu\nu} - \frac{k^\mu q^\nu + k^\nu q^\mu}{k\cdot q},$$

$$\sum_{\text{pol}} \zeta^{*\mu}(-\omega)\zeta^\nu(\omega) = \eta^{\mu\nu}$$

$$A(\mathfrak{z}_i,\cdots) = \lim_{\bar\omega\to 0} i m_i \omega\bar\omega\langle z_i(\omega)\cdots\rangle$$
$$A(\bar{\mathfrak{z}}_i,\cdots) = \lim_{\omega\to 0} i m_i \omega\bar\omega\langle z_i(\bar\omega)\cdots\rangle$$

$$A(z_i,\cdots) = \lim_{\omega,\bar\omega\to 0} i m_i \omega\bar\omega\langle z_i(\omega)\cdots\rangle$$

$$\lim_{\omega\to 0} m\omega\bar\omega\mathcal{A} = -i\sum_{\text{pol.}} \mathcal{A}_L(\mathfrak{z})\mathcal{A}_R(\bar{\mathfrak{z}})\Big|_{\omega=0} = -i\sum_{\text{pol.}} \mathcal{A}_L(z)\mathcal{A}_R(\bar{\mathfrak{z}})$$
$$\lim_{\bar\omega\to 0} m\omega\bar\omega\mathcal{A} = -i\sum_{\text{pol.}} \mathcal{A}_L(\mathfrak{z})\mathcal{A}_R(\bar{\mathfrak{z}})\Big|_{\bar\omega=0} = -i\sum_{\text{pol.}} \mathcal{A}_L(\mathfrak{z})\mathcal{A}_R(z)$$



$$\mathcal{A}|_{\varepsilon_i^{(\mu}\varepsilon_i^{\nu)}\to k_i^{(\mu}\varepsilon_i^{\nu)}} = \mathcal{O}(\omega_i^2).$$

$$A(h_1,\dots,h_m,\mathfrak{z}_1,\dots,\mathfrak{z}_{n+1}(\omega)) = \zeta^\mu \frac{\partial}{\partial b_\mu} A(h_1,\dots,h_m,\mathfrak{z}_1,\dots,\mathfrak{z}_n)$$
$$+i\omega\zeta^\mu \frac{\partial}{\partial u_\mu} A(h_1,\dots,h_m,\mathfrak{z}_1,\dots,\mathfrak{z}_n) + \mathcal{O}(\omega^2),$$

$$\frac{\partial L}{\partial b^\mu} = \frac{\delta L}{\delta z^\mu} = \partial_\tau \frac{\delta L}{\delta \dot{z}^\mu}$$

$$\zeta^\mu \langle 0|\frac{\partial L}{\partial b_\mu}|\alpha\rangle = -i\zeta^\mu \frac{\partial}{\partial b_\mu}\langle 0 \mid \alpha\rangle = -i\zeta^\mu \frac{\partial}{\partial b_\mu} A(\alpha)$$

$$\zeta_{n+1}^\mu \langle 0|\partial_\tau \frac{\delta L}{\delta \dot{z}^\mu}(\omega)|\alpha\rangle = -\langle 0|m\ddot{z}_\mu(\omega) + \mathcal{O}(z^2)|\alpha\rangle = m\omega\bar{\omega}\langle 0|z_\mu(\omega)|\alpha\rangle + \mathcal{O}(z^2)$$

$$\lim_{\omega\to 0}\lim_{\bar{\omega}\to 0} \zeta^\mu \langle 0|\partial_\tau \frac{\delta L}{\delta \dot{z}^\mu}(\omega)|\alpha\rangle = -i\lim_{\omega\to 0} A(\alpha,\mathfrak{z})$$

$$\frac{\partial L}{\partial u^\mu} = \tau \frac{\delta L}{\delta z^\mu} = \tau\partial_\tau \frac{\delta L}{\delta \dot{z}^\mu}$$

$$\zeta^\mu \frac{\partial L}{\partial u^\mu} = \zeta^\mu i \frac{\partial}{\partial \omega}\langle 0|\partial_\tau \frac{\delta L}{\delta \dot{z}^\mu}(\omega)|\alpha\rangle = -\lim_{\omega\to 0}\frac{\partial}{\partial\omega} A(\alpha,\mathfrak{z})$$

$$A = \delta(k\cdot u)e^{ik\cdot b}\mathcal{A}\,.$$

$$\lim_{\omega_{n+1}\to 0}\mathcal{A}(h_1,\dots,h_m,\mathfrak{z}_1,\dots,\mathfrak{z}_{n+1}) = i\zeta_{n+1}\cdot k\mathcal{A}(h_1,\dots,h_m,\mathfrak{z}_1,\dots,\mathfrak{z}_n)$$

$$\mathcal{A}_{\text{bulk}}(h_1,h_2,h_3) = \quad h_1(k_1) \quad h_2(k_2) \quad h_3(k_3)$$

$$= a_1\left((k_1\cdot\varepsilon_3)^2(\varepsilon_1\cdot\varepsilon_2)^2 + (k_1\cdot\varepsilon_2)^2(\varepsilon_1\cdot\varepsilon_3)^2 + (k_2\cdot\varepsilon_1)^2(\varepsilon_2\cdot\varepsilon_3)^2\right)$$
$$+ a_2\big((k_1\cdot\varepsilon_2)(k_1\cdot\varepsilon_3)(\varepsilon_1\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_3) - (k_1\cdot\varepsilon_3)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_3)$$
$$+ (k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_3)(\varepsilon_2\cdot\varepsilon_3)\big)\,.$$

$$A(h_1,h_2,h_3) = -i\kappa\big((k_1\cdot\varepsilon_3)(\varepsilon_1\cdot\varepsilon_2) - (k_1\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_3) + (k_2\cdot\varepsilon_1)(\varepsilon_2\cdot\varepsilon_3)\big)^2.$$

$$\mathcal{A}(h_1) = \quad u \quad h_1(k) \quad = \kappa'(u\cdot\varepsilon)^2.$$


$$\mathcal{A}(h_1,\mathfrak{z}_1) = \quad u \quad \mathfrak{z}_1(\omega) \quad h_1(k) \quad = a_1(u\cdot\varepsilon)^2(k\cdot\zeta) + a_2\omega(u\cdot\varepsilon)(\varepsilon\cdot\zeta)$$

$$\mathcal{A}(h_1,\mathfrak{z}_1) = i\kappa'((u\cdot\varepsilon)^2(k\cdot\zeta) + 2\omega(u\cdot\varepsilon)(\varepsilon\cdot\zeta))$$

$$\mathcal{A}(h_1,\mathfrak{z}_1,\ldots,\mathfrak{z}_n) = \quad u \quad \mathfrak{z}_1(\omega_1) \quad \mathfrak{z}_n(\omega_n) \quad h_1(k) \quad .$$

$$= (i)^n\kappa'(u\cdot\varepsilon)^2\prod_i(k\cdot\zeta_i) + a_1\sum_i\left(\omega_i(\varepsilon\cdot\zeta_i)(u\cdot\varepsilon)\prod_{j\neq i}(k\cdot\zeta_j)\right)$$


$$+ a_2\sum_{i<j}\left(\omega_i\omega_j(\varepsilon\cdot\zeta_i)(\varepsilon\cdot\zeta_j)\prod_{l\neq i,j}(k\cdot\zeta_l)\right)$$

$$\mathcal{A}_{\kappa^2}(h_1,h_2) =$$ 

$$= \frac{N_1}{2(k_1\cdot k_2)} + \frac{N_2}{m\bar{\omega}\omega}$$

$$\mathrm{LI}_1 = \{(u\cdot k_1),(k_1\cdot k_2),(k_1\cdot\varepsilon_2),(k_2\cdot\varepsilon_1),(u\cdot\varepsilon_1),(u\cdot\varepsilon_2),(\varepsilon_1\cdot\varepsilon_2)\}$$
$$\mathrm{LI}_2 = \{\omega,\bar{\omega},(k_1\cdot k_2),(k_1\cdot\varepsilon_2),(k_2\cdot\varepsilon_1),(u\cdot\varepsilon_1),(u\cdot\varepsilon_2),(\varepsilon_1\cdot\varepsilon_2)\}$$


$$\begin{aligned}
N_1 =\ & a_{1,1}(u\cdot k_1)^2(\varepsilon_1\cdot\varepsilon_2)^2 + a_{1,2}(u\cdot k_1)[(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_2) - (u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2)]\\
& + a_{1,3}[(u\cdot\varepsilon_1)^2(k_1\cdot\varepsilon_2)^2 + (u\cdot\varepsilon_2)^2(k_2\cdot\varepsilon_1)^2] + a_{1,4}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(k_1\cdot k_2)(\varepsilon_1\cdot\varepsilon_2)\\
& + a_{1,5}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1) + a_{1,6}(k_1\cdot k_2)(\varepsilon_1\cdot\varepsilon_2)^2\\
& + a_{1,7}(k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2)\\
N_2 =\ & a_{2,1}\omega\bar{\omega}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_2) + a_{2,2}\omega\bar{\omega}(\varepsilon_1\cdot\varepsilon_2)^2\\
& + a_{2,3}[\omega(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)(k_1\cdot\varepsilon_2) - \bar{\omega}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)^2(k_2\cdot\varepsilon_1)]\\
& + a_{2,4}[-\bar{\omega}(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)(k_1\cdot\varepsilon_2) + \omega(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)^2(k_2\cdot\varepsilon_1)]\\
& + a_{2,5}[\omega(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_2) - \bar{\omega}(u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2)]\\
& + a_{2,6}[-\bar{\omega}(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_2) + \omega(u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2)]\\
& + a_{2,7}(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)^2(k_1\cdot k_2) + a_{2,8}[(u\cdot\varepsilon_1)^2(k_1\cdot\varepsilon_2)^2 + (u\cdot\varepsilon_2)^2(k_2\cdot\varepsilon_1)^2]\\
& + a_{2,9}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(k_1\cdot k_2)(\varepsilon_1\cdot\varepsilon_2) + a_{2,10}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)\\
& + a_{2,11}(k_1\cdot k_2)(\varepsilon_1\cdot\varepsilon_2)^2 + a_{2,12}(k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2)
\end{aligned}$$

$$N_1\big|_{(k_1\cdot k_2)=0} =$$ 

.

$$\begin{aligned}
& a_{1,1}(u\cdot k_1)^2(\varepsilon_1\cdot\varepsilon_2)^2 + a_{1,2}(u\cdot k_1)[(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_2) - (u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2)]\\
& + a_{1,3}[(u\cdot\varepsilon_1)^2(k_1\cdot\varepsilon_2)^2 + (u\cdot\varepsilon_2)^2(k_2\cdot\varepsilon_1)^2] + a_{1,5}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)\\
& + a_{1,7}(k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2)\\
& = -\frac{i\kappa\kappa'}{2}[-(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2) + (u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1) + (u\cdot k_1)(\varepsilon_1\cdot\varepsilon_2)]^2.
\end{aligned}$$


$$N_1 = -\frac{i\kappa\kappa'}{2}[-(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2) + (u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1) + \omega(\varepsilon_1\cdot\varepsilon_2)]^2$$
$$a_{1,4}(k_1\cdot k_2)(\varepsilon_1\cdot\varepsilon_2)(u\cdot\varepsilon_1)(u\cdot\varepsilon_2) + a_{1,6}(k_1\cdot k_2)(\varepsilon_1\cdot\varepsilon_2)^2$$

$$N_2|_{\omega=0} =$$ 

$$
\begin{aligned}
&-a_{2,3}\bar{\omega}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)^2(k_2\cdot\varepsilon_1) - a_{2,4}\bar{\omega}(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)(k_1\cdot\varepsilon_2)\\
&-a_{2,5}\bar{\omega}(u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2) - a_{2,6}\bar{\omega}(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_2)\\
&+a_{2,7}(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)^2(k_1\cdot k_2) + a_{2,8}[(u\cdot\varepsilon_1)^2(k_1\cdot\varepsilon_2)^2 + (u\cdot\varepsilon_2)^2(k_2\cdot\varepsilon_1)^2]\\
&+a_{2,9}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(k_1\cdot k_2)(\varepsilon_1\cdot\varepsilon_2) + a_{2,10}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)\\
&+a_{2,11}(k_1\cdot k_2)(\varepsilon_1\cdot\varepsilon_2)^2 + a_{2,12}(k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2)\\
&=\frac{i\kappa'^2}{4}(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)[-(u\cdot\varepsilon_2)(k_1\cdot k_2)+2\bar{\omega}(k_1\cdot\varepsilon_2)]
\end{aligned}
$$

$$
\begin{aligned}
N_2 =& -\frac{i\kappa'^2}{4}(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)^2(k_1\cdot k_2) - \frac{i\kappa'^2}{2}\omega(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)(k_1\cdot\varepsilon_2) + \frac{i\kappa'^2}{2}\bar{\omega}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)^2(k_2\cdot\varepsilon_1)\\
&+a_{2,1}\omega\bar{\omega}(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_2) + a_{2,2}\omega\bar{\omega}(\varepsilon_1\cdot\varepsilon_2)^2
\end{aligned}
$$

$$
\begin{aligned}
\mathcal{A}(h_1,h_2)|_{\kappa^2} =& \frac{-im\kappa^2}{4}\Bigg(\frac{\big((k_1\cdot u)(\varepsilon_1\cdot\varepsilon_2)+(k_2\cdot\varepsilon_1)(u\cdot\varepsilon_2)-(k_1\cdot\varepsilon_2)(u\cdot\varepsilon_1)\big)^2}{(k_1\cdot k_2)} + \frac{(k_1\cdot k_2)(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)^2}{(u\cdot k_1)^2}\\
&+\frac{2(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)\big((k_1\cdot\varepsilon_2)(u\cdot\varepsilon_1)-(k_2\cdot\varepsilon_1)(u_1\cdot\varepsilon_2)\big)}{(u\cdot k_1)} - 2(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_2)\Bigg).
\end{aligned}
$$

$$\mathcal{A}(h_1,h_2,\mathfrak{z})|_{\kappa^2} =$$ 

$$=\frac{N_1}{2(k_1\cdot k_2)} + \frac{N_2}{m\bar{\omega}_2\omega_2} + \frac{N_3}{m\bar{\omega}_2\omega_2},$$


$$N_3 = N_2|_{k_1\leftrightarrow k_2,\varepsilon_1\leftrightarrow\varepsilon_2}$$

$$\mathcal{A}(h_1,h_2,z)|_{\omega_1=0} = i\big((k_1+k_2)\cdot\zeta\big)\mathcal{A}(h_1,h_2).$$

$$\mathrm{LI}_1 = \{\omega_1,(u\cdot k_1),(u\cdot\zeta),(u\cdot\varepsilon_1),(u\cdot\varepsilon_2),(k_1\cdot\varepsilon_2),(k_2\cdot\varepsilon_1),(k_1\cdot\zeta),(k_2\cdot\zeta),(\varepsilon_1\cdot\zeta),(\varepsilon_2\cdot\zeta),(k_1\cdot k_2)\}$$
$$\mathrm{LI}_2 = \{\omega_1,\omega_2,\bar{\omega}_2,(u\cdot\zeta),(u\cdot\varepsilon_1),(u\cdot\varepsilon_2),(k_1\cdot\varepsilon_2),(k_2\cdot\varepsilon_1),(k_1\cdot\zeta),(k_2\cdot\zeta),(\varepsilon_1\cdot\zeta),(\varepsilon_2\cdot\zeta),(k_1\cdot k_2)\}$$


$$
\begin{aligned}
N_1 =& a_{1,1}\omega_1(k_1\cdot k_2)(\varepsilon_1\cdot\varepsilon_2)[(u\cdot\varepsilon_2)(\varepsilon_1\cdot\zeta)+(u\cdot\varepsilon_1)(\varepsilon_2\cdot\zeta)]\\
&+a_{1,2}\omega_1(\varepsilon_1\cdot\varepsilon_2)[(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)(k_1\cdot\zeta)+(u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)(k_2\cdot\zeta)]+\cdots\\
&+a_{1,53}(u\cdot k_1)(\varepsilon_1\cdot\varepsilon_2)\big[(u\cdot k_1)(k_1\cdot\varepsilon_2)(\varepsilon_1\cdot\zeta)+(\varepsilon_2\cdot\zeta)(k_2\cdot\varepsilon_1)\big(2\omega_1+(u\cdot k_1)\big)\big]\\
&+a_{1,55}(u\cdot u)(k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)[(k_1\cdot\varepsilon_2)(\varepsilon_1\cdot\zeta)+(k_2\cdot\varepsilon_1)(\varepsilon_2\cdot\zeta)]\\
N_2 =& a_{2,1}\omega_1(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)(k_1\cdot k_2)(z_1\cdot\varepsilon_2) + a_{2,2}\omega_1(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)(k_1\cdot z_1)(k_1\cdot\varepsilon_2)\\
&+a_{2,3}\omega_1(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)(k_1\cdot\varepsilon_2)(k_2\cdot z_1) + a_{2,4}\omega_1\omega_2(u\cdot\varepsilon_1)^2(k_1\cdot\varepsilon_2)(z_1\cdot\varepsilon_2)+\cdots\\
&+a_{2,99}\omega_2\bar{\omega}_2(u\cdot u)(k_1\cdot\varepsilon_2)(z_1\cdot\varepsilon_1)(\varepsilon_1\cdot\varepsilon_2) + a_{2,100}\omega_2\bar{\omega}_2(u\cdot u)(k_2\cdot\varepsilon_1)(z_1\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_2)\\
&+a_{2,101}(u\cdot u)^2(k_1\cdot\varepsilon_2)^2(k_2\cdot\varepsilon_1)(z_1\cdot\varepsilon_1) + a_{2,102}(u\cdot u)^2(k_1\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)^2(z_1\cdot\varepsilon_2)
\end{aligned}
$$

$$N_1|_{(k_1\cdot k_2)=0} = \quad + \mathcal{O}(\omega_1^2).$$




$$\begin{aligned}\text{RHS} =& \frac{m\kappa^2}{2}[(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)-(u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)-(u\cdot k_1)(\varepsilon_1\cdot\varepsilon_2)]\times\\ &[(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)\big((k_1\cdot\zeta)+(k_2\cdot\zeta)\big)+2\omega_1(k_1\cdot\varepsilon_2)(\zeta\cdot\varepsilon_1)\\ &-(k_2\cdot\varepsilon_1)\Big((u\cdot\varepsilon_2)\big((k_1\cdot\zeta)+(k_2\cdot\zeta)\big)+2\omega_1(\zeta\cdot\varepsilon_2)\Big)\\ &-\Big(2\omega_1(k_1\cdot\zeta)+(u\cdot k_1)\big((k_1\cdot\zeta)+(k_2\cdot\zeta)\big)\Big)(\varepsilon_1\cdot\varepsilon_2)\Big]+\mathcal{O}(\omega_1^2)\end{aligned}$$


$$N_2|_{\omega_2=0} = \quad ..$$





$$N_2|_{\bar{\omega}_2=0} = \quad .$$



$$\begin{aligned}&\mathcal{A}\big(h_1(k_1),h_2(k_2),z(\omega_1)\big)=m\kappa^2\Big[\Big(\frac{(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)^2(k_1\cdot k_2)(k_2\cdot\zeta)+2\omega_1(u\cdot\varepsilon_1)^2(u\cdot\varepsilon_2)(\zeta\cdot\varepsilon_2)(k_1\cdot k_2)}{4(u\cdot k_1)^2}\\ &+\frac{(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(k_2\cdot\zeta)[(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)-(u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)H+\omega_1(u\cdot\varepsilon_1)(\zeta\cdot\varepsilon_2)[(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)-2(u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)]}{2(u\cdot k_1)}+(k_1\leftrightarrow k_2,\varepsilon_1\leftrightarrow\varepsilon_2)\Big)\\ &+\frac{\big((k_1+k_2)\cdot\zeta\big)[(u\cdot k_1)(\varepsilon_1\cdot\varepsilon_2)+(u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)-(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)]^2}{4(k_1\cdot k_2)}\\ &+\frac{2\omega_1[(u\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)-(u\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)-(u\cdot k_1)(\varepsilon_1\cdot\varepsilon_2)][(\zeta\cdot\varepsilon_1)(k_1\cdot\varepsilon_2)-(\zeta\cdot\varepsilon_2)(k_2\cdot\varepsilon_1)-(k_1\cdot\zeta)(\varepsilon_1\cdot\varepsilon_2)]}{4(k_1\cdot k_2)}\\ &-\frac{1}{2}\Big(\big((k_1+k_2)\cdot\zeta\big)(u\cdot\varepsilon_1)(u\cdot\varepsilon_2)(\varepsilon_1\cdot\varepsilon_2)+\omega_1(\varepsilon_1\cdot\varepsilon_2)[(u\cdot\varepsilon_2)(\zeta\cdot\varepsilon_1)+(u\cdot\varepsilon_1)(\zeta\cdot\varepsilon_2)]\Big)\Big]\end{aligned}$$



u
ω_1
ω_2
1 2 3
Ia
Ib
Ic
IIa
IIb
IIIc
III



u_1
u_2
3
1
2
4
MC1
MC2
N^{1}MC1





$$\bar{S}|_{\kappa^2} = \frac{i m_1^2 m_2 \kappa^4}{16(-2+D)^2 k_1^2 k_2^2}\left[\frac{(k_1\cdot k_2)((-2+D)\gamma^2-1)^2}{(k_1\cdot u_1)^2} + \frac{2\big((-3+D)(-2+D)(k_1\cdot u_1)^2 + 2(k_1\cdot k_2)(-(-2+D)^2\gamma^2+1)\big)}{(k_1+k_2)^2}\right] + (u_1 \leftrightarrow u_2, m_1 \leftrightarrow m_2)$$

$$\varepsilon_i^\mu \varepsilon_i^\nu \to m_2 \kappa \left(u_2^\mu u_2^\nu - \frac{1}{D-2}\eta^{\mu\nu}\right)$$












MC1




MC2




MC3




MC4




MC5




MC6




MC7




MC8




MC9




$$\mathcal{I}_0 = \int_\ell \frac{1}{\ell^2(\ell-k)^2} \quad \mathcal{I}_1 = \int_\ell \frac{\ell^\mu}{\ell^2(\ell-k)^2} \quad \mathcal{I}_2 = \int_\ell \frac{\ell^\mu \ell^\nu}{\ell^2(\ell-k)^2}$$





$$C_{123} \sim \langle \mathrm{Tr} Z_1^{J_1} \mathrm{Tr} Z_2^{J_2} \mathrm{Tr} D^S Z^J \rangle$$



$$\ell_A = \frac{|J_1 - J_2| + J}{2}, \ell_B = \frac{J_1 + J_2 - J}{2},$$

$$\mathcal{B} = 1 + \sum_{a=1}^{\infty} \int \frac{du}{2\pi} e^{-\frac{1}{2}(J_1+J_2)\tilde{E}_a(u)} \tilde{\mu}_a(u) \mathbf{t}_{a,1}(u) + \cdots$$

$$\tilde{\mu}_a(u) = \frac{a}{g^2(x^{[+a]}x^{[-a]})^2} \prod_{\sigma_1,\sigma_2=\pm} \left(1 - \frac{1}{x^{[\sigma_1 a]}x^{[\sigma_2 a]}}\right)^{-1},$$

$$\tilde{E}_a(u) = \log\left(x^{[+a]}x^{[-a]}\right)$$

$$x(u) = \frac{u + \sqrt{u^2 - 4g^2}}{2g}$$

$$\mathbf{t}_{a,1}^{\text{phys}}(u) = -\sum_{b=1}^{4} \mathbf{P}_b^{[+a]}(u)\mathbf{P}^{b[-a]}(u) + \mathcal{O}(S^2)$$

$$\mathbf{P}_1 = \mathbf{P}^4 = \epsilon x^{-J/2}, \mathbf{P}_2 = -\mathbf{P}^3 = -\epsilon x^{\frac{J}{2}} \sum_{n=\frac{J}{2}+1}^{\infty} I_{2n-1}x^{1-2n},$$

$$\mathbf{P}_3 = \mathbf{P}^2 = \epsilon\left(x^{-J/2} - x^{J/2}\right),$$

$$\mathbf{P}_4 = -\mathbf{P}^1 = \epsilon x^{J/2} \sum_{n=J/2+1}^{\infty} I_{2n-1}x^{1-2n} - \epsilon x^{-J/2} \sum_{n=1-J/2}^{\infty} I_{2n-1}x^{2n-1},$$

$$\epsilon^2 = \frac{2\pi i S}{J I_J(4\pi g)}$$

$$\gamma = \Delta - S - J = \gamma_J^{(1)} S + \mathcal{O}(S^2), \gamma_J^{(1)} = \frac{4\pi g I_{J+1}(4\pi g)}{J I_J(4\pi g)}$$

$$\mathbf{t}_{a,1}(u) = -\sum_{b=1}^{4} \mathbf{P}_b^{[+a]}(u)\widetilde{\mathbf{P}}^{b[-a]}(u) + \mathcal{O}(S^2)$$

$$\mathcal{A} = 1 + \sum_{a\geqslant 1}\left(\iiint_{\text{roots}} + \int\right)\frac{du}{2\pi} e^{-\frac{1}{2}(J_1 - J_2)\tilde{E}_a(u)}\tilde{\mu}_a(u)\frac{\mathbf{T}_{a,1}(u)}{\mathbf{T}_{a,0}^{+}(u)}$$
$$+ \sum_{a\geqslant 1}\int \frac{du}{2\pi} e^{-\frac{1}{2}(J_2 - J_1)\tilde{E}_a(u)}\tilde{\mu}_a(u)\frac{\mathbf{T}_{a,1}(u)}{\mathbf{T}_{a,0}^{-}(u)} + \cdots$$

$$\sum_{a\geqslant 1}\oiiint \frac{du}{2\pi}\left[e^{-\frac{1}{2}(J_1 - J_2)\tilde{E}_a(u)}\tilde{\mu}_a(u)\frac{\mathbf{T}_{a,1}(u)}{\mathbf{T}_{a,0}^{+}(u)}\right]^{\circlearrowleft}$$
$$+ \sum_{a\geqslant 1}\oiiint \frac{du}{2\pi} e^{-\frac{1}{2}(J_2 - J_1)\tilde{E}_a(u)}\tilde{\mu}_a(u)\frac{\mathbf{T}_{a,1}(u)}{\mathbf{T}_{a,0}^{-}(u)}$$

$$\left(x^{[\pm a]}\right)^{\circlearrowleft} = \frac{1}{x^{[\pm a]}}$$

$$\mathbf{t}_{a,1}(u) = \left[\frac{\mathbf{T}_{a,1}(u)}{\mathbf{T}_{a,0}^{+}(u)}\right]^{\circlearrowleft} + \frac{\mathbf{T}_{a,1}(u)}{\mathbf{T}_{a,0}^{-}(u)} + \mathcal{O}(S^2)$$

$$\mathbf{T}_{a,1}(u) = -\left(\mathbf{P}_1^{[+a]}\mathbf{P}_2^{[-a]} - \mathbf{P}_2^{[+a]}\mathbf{P}_1^{[-a]}\right), \mathbf{T}_{a,0}(u) = 1$$

$$\mathcal{A} = 1 + SF_J(-\ell_A) + \mathcal{O}(S^2), \mathcal{B} = 1 + SF_J(\ell_B) + \mathcal{O}(S^2)$$

$$F_J(\ell) = \sum_{a=1}^{\infty} \oiint \frac{du}{2\pi} e^{-\left(\ell+\frac{1}{2}J\right)\tilde{E}_a(u)} \tilde{\mu}_a(u) t_a(u),$$

$$\sim \sum_a \left(1 + \gamma_J^{(1)}\right)/a$$

$$F_J(\ell) = \int_0^{\infty} \frac{dt e^t}{(1-e^t)^2} \oiint \frac{dxdy}{(2\pi)^2} \frac{e^{i(u-v)t} t_J(x,y)}{(xy)^{\ell+J/2}(xy-1)^2}$$

$$e^{i(u-v)t} \to e^{i(u-v)t} - 1 - i(u-v)t$$

$$t_J(x,y) = \frac{i}{(xy)^{J/2}} \left[\sum_{n=1}^{J/2} \mathrm{I}_{2n-1}\left(x^J(x^{1-2n} - y^{2n-1}) + y^J(x^{2n-1} - y^{1-2n})\right) + \sum_{n=1}^{\infty} \mathrm{I}_{2n-1}(x^{1-2n} - y^{2n-1})(y^J - x^J) \right]$$

$$\mathrm{I}_n = \frac{2\pi I_n(4\pi g)}{J I_J(4\pi g)}$$

ℓ_B	$F_J(\ell_B)$ for $J = 2$
1	$3g^4(4\zeta_2\zeta_3 + 5\zeta_5) - 48g^6(\zeta_3\zeta_4 + \zeta_2\zeta_5 + 7\zeta_7) + 4g^8(15\zeta_4\zeta_5 + 63\zeta_3\zeta_6 - 56\zeta_2\zeta_7 + 1470\zeta_9)$
2	$4g^6(-3\zeta_3\zeta_4 + 9\zeta_2\zeta_5 + 14\zeta_7) + 84g^8(\zeta_3\zeta_6 - 4\zeta_2\zeta_7 - 20\zeta_9)$
3	$2g^8(-30\zeta_4\zeta_5 + 56\zeta_2\zeta_7 + 105\zeta_9)$

ℓ_B	$F_J(\ell_B)$ for $J = 4$
1	$\frac{g^4}{\zeta_2}(9\zeta_3\zeta_4 + 10\zeta_2\zeta_5 - 7\zeta_7) + \frac{g^6}{\zeta_2}(-60\zeta_4\zeta_5 - \frac{48}{5}\zeta_3\zeta_6 - \frac{1152}{5}\zeta_2\zeta_7 + 240\zeta_9)$
2	$\frac{g^6}{\zeta_2}(30\zeta_4\zeta_5 - 3\zeta_3\zeta_6 + 36\zeta_2\zeta_7 - 30\zeta_9)$

$$\int_0^{\infty} \frac{dt e^t}{(1-e^t)^2} t^n = \Gamma(n+1)\zeta_n, \text{ for } n > 1$$

$$\zeta_n = \zeta(n) = \sum_{k \geqslant 1} 1/k^n$$

$$F_{J=2}(-1) = -8g^2\zeta_3 + g^4(-32\zeta_2\zeta_3 + 90\zeta_5) + g^6(160\zeta_3\zeta_4 + 288\zeta_2\zeta_5 - 1120\zeta_7) + g^8(-1440\zeta_4\zeta_5 - 896\zeta_3\zeta_6 - 3360\zeta_2\zeta_7 + 14700\zeta_9) + \mathcal{O}(g^{10})$$

$$\int_0^{\infty} dt \frac{e^{i(u-v)t} - 1}{e^t - 1} = \psi(1) - \psi(1 - iu + iv),$$

$$\psi(z) = \partial_z \log \Gamma(z)$$

$$F_J(\ell) = -\frac{ig}{2}\iiint \frac{dxdy}{(2\pi)^2 xy}\frac{x-y}{xy-1}t_J(x,y)(\psi(1+iu-iv)-\psi(1)+(u\leftrightarrow v)).$$

$$\frac{1}{2}(\psi(1+iu-iv)-\psi(1)+(u\leftrightarrow v)) = \sum_{k=1}^{\infty}(-1)^{k+1}g^{2k}\zeta_{2k+1}(x-y)^{2k}(1-1/xy)^{2k}$$

$$F_J(\ell) = f_J(\ell) + f_J(-J-\ell)$$

$$f_J(\ell) = \sum_{k=1}^{\infty}\sum_{n\in\mathbb{Z}}\frac{(-1)^{k+\ell+1}g^{2k+1}\Gamma(2k)\Gamma(2k+2)\zeta_{2k+1}\varepsilon(n)\mathrm{I}_{2n-J-1}}{\Gamma(1+k+n)\Gamma(2+k-n)\Gamma(k+\ell+n)\Gamma(1+k-\ell-n)}$$

$$J_\mu(2gt)J_\nu(2gt) = \int \frac{dz}{2\pi i}\frac{\Gamma(-z)\Gamma(2z+\mu+\nu+1)(gt)^{2z+\mu+\nu}}{\Gamma(z+\mu+1)\Gamma(z+\nu+1)\Gamma(z+\mu+\nu+1)}$$

$$f_J(\ell) = g\sum_{n\in\mathbb{Z}}(-1)^{n+1}\varepsilon(n)\mathrm{I}_{2n-J-1}\sum_{m=0}^{\infty}\left(c_{\ell+m,\ell+2n+m}+c_{\ell+m+1,\ell+2n+m-1}\right)$$

$$c_{i,j} = \int_0^\infty \frac{dt}{e^t-1}\left(J_i(2gt)J_j(2gt)-\delta_{i,0}\delta_{j,0}\right)$$

$$\sum_{m=0}^{\infty}J_{\ell+m+\delta}(2gt)J_{\ell+2n+m-\delta}(2gt)$$
$$= \frac{gt}{2(n-\delta)}\left(J_{\ell+\delta}(2gt)J_{\ell+2n-\delta-1}(2gt)-J_{\ell+\delta-1}(2gt)J_{\ell+2n-\delta}(2gt)\right)$$

$$\sum_{m=0}^{\infty}J_{\ell+m+\delta}(2gt)J_{\ell+m+\delta}(2gt) = \frac{1}{2}\left(1-\varepsilon(\ell+\delta)\sum_{m=\frac{1}{2}-\left|\ell+\delta-\frac{1}{2}\right|}^{\left|\ell+\delta-\frac{1}{2}\right|-\frac{1}{2}}J_m(2gt)J_m(2gt)\right).$$

$$f_J(\ell) = \frac{2\pi g}{J}[\delta_{\ell>0}(\log\, g-\psi(\ell))-\delta_{\ell<0}(\log\, g-\psi(-\ell))]+\mathcal{O}(1)$$

$$f_J(\ell) \approx \sum_{n=n_{\max}^{-}(g)}^{n_{\max}^{+}(g)}\tilde{f}_J(\ell,n),$$

$$\tilde{f}_J\left(\ell,n_{\max}^{\pm}(g)\right) \leq \mathbb{A}$$

$$f_J(\ell) \approx \sum_{i=-1}^{i_{\max}} g^{-i}(c_i+d_i\log\, g)$$

$$f_J(\ell) = \frac{\gamma_J^{(1)}}{4}(\log\,\lambda - 2\log(4\pi) - 2\psi(\ell)) - \frac{1}{2J} + \frac{1}{2}(\gamma_{\mathrm{E}} - \log(8\pi)) + \frac{3+4J}{8\sqrt{\lambda}} + \frac{J+2\ell}{4J}\left(\log(8\sqrt{\lambda}) + \gamma_{\mathrm{E}}\right) + \frac{J+2\ell}{\sqrt{\lambda}J}\left(\frac{2J^2-1}{8} + \frac{1-J^2-\ell(J+\ell)}{6}\zeta_2\right) + \mathcal{O}\left(\frac{1}{\lambda}\right)$$

$$F_J(\ell) = \Psi_J(\ell) + P_J(\ell)$$

$$\Psi_J(\ell) = \frac{\gamma_J^{(1)}}{2}(\psi(J+\ell) - \psi(\ell)) + \psi(J+\ell) + C,$$

$$\Psi_J(-\ell) = \frac{\gamma_J^{(1)}}{2}(\psi(J-\ell) + \psi(\ell) - \log\,\lambda + 2\log(4\pi)) + \psi(J-\ell) + \psi(\ell) + C,$$

$$P_J(\ell) = \frac{3}{4\sqrt{\lambda}} + \frac{1}{\lambda}\left(\frac{17-8J^2}{32} + \frac{11-8J^2-24\ell(J+\ell)}{48}\zeta_3\right) + \frac{1}{\lambda^{3/2}}\left(\frac{11-9J^2}{16} + \frac{11-8J^2-18\ell(J+\ell)}{48}\zeta_3\right) + \delta_{-J<\ell<0}\left(\frac{1}{J} - \frac{J}{\sqrt{\lambda}} - \frac{J}{2\lambda} - \frac{13J-4J^3}{24\lambda^{3/2}}\right) + \mathcal{O}\left(\frac{1}{\lambda^2}\right)$$

$$\mathcal{B} = \left(\frac{e^{\gamma_{\mathrm{E}}S}}{2^S\lambda^{S/2}}\right)\frac{\Gamma(\ell_B - \gamma/2)\Gamma(J+\ell_B+S+\gamma/2)}{\Gamma(\ell_B)\Gamma(J+\ell_B)} \times \mathcal{D}_{\mathcal{B}},$$

$$\mathcal{A} = \left(\frac{e^{\gamma_{\mathrm{E}}S}(4\pi)^{S+\gamma}}{2^S\lambda^{S+\gamma/2}}\right)\frac{\Gamma(\ell_A+\gamma/2+S)\Gamma(J-\ell_A+S+\gamma/2)}{\Gamma(\ell_A)\Gamma(J-\ell_A)} \times \mathcal{D}_{\mathcal{A}},$$

$$\log\,\mathcal{D}_{\mathcal{B}} = SP_J(\ell_B) + \mathcal{O}(S^2), \log\,\mathcal{D}_{\mathcal{A}} = SP_J(-\ell_A) + \mathcal{O}(S^2)$$

$$\frac{C_{123}}{C_{123}^{(0)}} = \frac{\Gamma[\mathrm{AdS}\,]}{\Gamma[\mathrm{dS}\,]} \times \frac{\mathcal{D}_{J_1J_2J}(S)}{\lambda^{S/4}\Gamma\left(1+\frac{S}{2}\right)},$$

$$\Gamma[AdS] = \frac{\Gamma\left(\frac{\Delta_1-\Delta_2+\Delta+S}{2}\right)\Gamma\left(\frac{\Delta_2-\Delta_1+\Delta+S}{2}\right)\Gamma\left(\frac{\Delta_1+\Delta_2-\Delta+S}{2}\right)\Gamma\left(\frac{\Delta_1+\Delta_2+\Delta+S}{2}\right)}{\sqrt{\Gamma(\Delta+S)\Gamma(\Delta+S-1)}},$$

$$\Gamma[\mathrm{dS}] = \Gamma[\mathrm{AdS}\,]_{\Delta\to J, S\to 0}$$

$$\log\,\mathcal{D} = \mathcal{D}_1 S + \frac{\mathcal{D}_2}{\sqrt{\lambda}}S^2 + \frac{\mathcal{D}_3}{\lambda}S^3 + \mathcal{O}\left(\frac{S^4}{\lambda^{3/2}}\right)$$

$$\mathcal{D}_n = \mathcal{D}_n^{(0)} + \frac{\mathcal{D}_n^{(1)}}{\sqrt{\lambda}} + \frac{\mathcal{D}_n^{(2)}}{\lambda} + \cdots$$

$$\log\,\mathcal{D} = \sqrt{\lambda}\mathcal{D}^{\mathrm{cl}} + \mathcal{O}(1)$$

$$\mathcal{D}^{\mathrm{cl}} = \sum_{n=1}^{\infty} \mathcal{D}_n^{(0)} \mathcal{S}^n$$

$$\log \mathcal{D} = \sum_{k=1}^{\infty} \frac{1}{(\sqrt{\lambda})^{k-1}} \sum_{n=1}^{k} \mathcal{D}_n^{(k-n)} S^n$$

$$\Delta^2 = J^2 + \sqrt{\lambda}\left(A_1 S + \frac{A_2}{\sqrt{\lambda}} S^2 + \frac{A_3}{\lambda} S^3 + \mathcal{O}\left(\frac{S^4}{\lambda^{3/2}}\right)\right)$$

$$\begin{aligned}\log \tilde{\lambda}^2_{\Delta,S} = & \frac{S\left(\frac{17}{6} + S + \left(-\frac{7}{12} + \zeta_3\right) S^2\right)}{\Delta^2} \\ & + \frac{S\left(\frac{511}{60} + 6S + \left(\frac{1}{12} - 2\zeta_3\right) S^2 - \left(\frac{13}{8} + 6\zeta_3\right) S^3 + \left(\frac{31}{40} - \frac{3}{2}\zeta_5\right) S^4\right)}{\Delta^4} \\ & + \mathcal{O}\left(\frac{1}{\Delta^6}\right)\end{aligned}$$

$$\Delta^2 = 2\sqrt{\lambda} S + \left(4 - S + \frac{3}{2} S^2\right) + \frac{1}{\sqrt{\lambda}}\left(\frac{15}{4} S + \frac{3 - 24\zeta_3}{8} S^2 - \frac{3}{8} S^3\right) + \mathcal{O}\left(\frac{1}{\lambda}\right)$$

$$\mathcal{R} = \left(\frac{\sqrt{\lambda} S}{2}\right)^S \frac{\Delta(\Delta - S)}{(\Delta + S)^2} \frac{\Gamma\left(\frac{1 + \Delta + S}{2}\right) \Gamma\left(\frac{\Delta - S}{2}\right)^3}{\Gamma\left(\frac{1 + \Delta - S}{2}\right) \Gamma\left(\frac{\Delta + S}{2}\right)^3}$$

$$\begin{aligned}\log \mathcal{D}_{222} = & \frac{1}{\sqrt{\lambda}}\left[\frac{5}{8} S - \frac{7 - 4\zeta_3}{16} S^2\right] \\ & + \frac{1}{\lambda}\left[-\frac{13 + 24\zeta_3}{32} S - \frac{49 - 8\zeta_3}{64} S^2 + \frac{25 - 12\zeta_3 - 12\zeta_5}{64} S^3\right] \\ & + \mathcal{O}\left(\frac{1}{\lambda^{3/2}}\right)\end{aligned}$$

$$\log \mathcal{D}_{J_1 J_2 J} = \log \mathcal{D}_{222} + \mathcal{O}\left(\frac{1}{\lambda}\right)$$

$$V'(x) = R(x + i0) + R(x - i0), V'(x) = \mathrm{sgn}(x) - \frac{2\mathcal{J}x}{x^2 - 1}$$

$$R(x) = 2x \int_a^b \frac{dy V'(y)}{x^2 - y^2} \sqrt{\frac{(x^2 - b^2)(x^2 - a^2)}{(b^2 - y^2)(y^2 - a^2)}}$$

$$R(x) \sim 2\pi x(\mathcal{S} - \mathcal{E} + \mathcal{J}), R(x) \sim \frac{2\pi}{x}(\mathcal{S} + \mathcal{E} - \mathcal{J})$$

$$\mathcal{J}=\frac{\sqrt{(a^2-1)(b^2-1)}}{\pi b}K\left(1-\frac{a^2}{b^2}\right)$$
$$\mathcal{S}=\frac{ab+1}{2\pi ab}\left[bE\left(1-\frac{a^2}{b^2}\right)-aK\left(1-\frac{a^2}{b^2}\right)\right]$$
$$\mathcal{E}=\frac{ab-1}{2\pi ab}\left[bE\left(1-\frac{a^2}{b^2}\right)+aK\left(1-\frac{a^2}{b^2}\right)\right]$$

$$\rho(x)=\frac{R(x-i0)-R(x+i0)}{2\pi i}=-\frac{2x}{\pi}\int_a^b\frac{dyV'(y)}{x^2-y^2}\sqrt{\frac{(b^2-x^2)(x^2-a^2)}{(b^2-y^2)(y^2-a^2)}},$$

$$\mathcal{A}^{\mathrm{SYM}}=\mathcal{A}^{\mathrm{SYM}}_{\mathrm{asy}}+I_1[\mathcal{L}_A]+I_1[\mathcal{J}-\mathcal{L}_A]$$
$$\mathcal{B}^{\mathrm{SYM}}=I_{-1}[\mathcal{L}_B]+I_1[\mathcal{J}+\mathcal{L}_B]$$
$$\mathcal{N}^{\mathrm{SYM}}=\mathcal{N}^{\mathrm{SYM}}_{\mathrm{asy}}-I_2[\mathcal{J}]$$

$$\mathcal{L}_A=\frac{\mathcal{J}_1-\mathcal{J}_2+\mathcal{J}}{2},\mathcal{L}_B=\frac{\mathcal{J}_1+\mathcal{J}_2-\mathcal{J}}{2}.$$

$$I_q[\mathcal{L}]=\int_{U^-}\frac{dx(x-1/x)}{8\pi^2x}\left[\mathrm{Li}_2\left(e^{\frac{4\pi i\mathcal{L}x}{x^2-1}+iqR(x)}\right)+\mathrm{Li}_2\left(e^{\frac{4\pi i\mathcal{L}x}{x^2-1}-iqR(1/x)}\right)\right]$$
$$-(\text{ same with } R\to 0)$$

$$\mathrm{Li}_2(z)=\sum_{n=1}^{\infty}z^n/n^2$$

$$\mathcal{N}^{\mathrm{SYM}}_{\mathrm{asy}}=-\int_a^b\frac{dx(x-1/x)}{4\pi^2x}\left[\mathrm{Li}_2\left(e^{-2\pi\rho(x)}\right)+\pi^2\rho^2(x)-\zeta_2\right]$$

$$V'(\alpha)=0\ \Rightarrow\ \alpha=\mathcal{J}+\sqrt{1+\mathcal{J}^2}.$$

$$a=\alpha-2\alpha\sqrt{\frac{\alpha}{\alpha^2+1}}\sqrt{\mathcal{S}}+\frac{\alpha^2(3\alpha^4+6\alpha^2-1)}{(\alpha^2-1)(\alpha^2+1)^2}\mathcal{S}+\mathcal{O}\left(\mathcal{S}^{3/2}\right)$$
$$b=\alpha+2\alpha\sqrt{\frac{\alpha}{\alpha^2+1}}\sqrt{\mathcal{S}}+\frac{\alpha^2(3\alpha^4+6\alpha^2-1)}{(\alpha^2-1)(\alpha^2+1)^2}\mathcal{S}+\mathcal{O}\left(\mathcal{S}^{3/2}\right)$$

$$R(x)=R^{(1)}(x)\mathcal{S}+R^{(2)}(x)\mathcal{S}^2+\cdots$$

$$R^{(1)}=\frac{4\pi x\alpha^2}{(\alpha^2-1)(x^2-\alpha^2)}$$
$$R^{(2)}=-\frac{4\pi x\alpha^3\left((x^4+\alpha^6)(\alpha^4+6\alpha^2+1)-x^2\alpha^2(5\alpha^6+3\alpha^4+3\alpha^2+5)\right)}{(\alpha^2-1)^3(\alpha^2+1)^2(x^2-\alpha^2)^3}$$

$$\rho=\sqrt{1-t^2}\left[\frac{2\sqrt{\alpha(\alpha^2+1)}}{\alpha^2-1}\sqrt{\mathcal{S}}-\frac{4\alpha^3(\alpha^2+3)t}{(\alpha^2-1)^2(\alpha^2+1)}\mathcal{S}\right.$$
$$+\frac{\alpha^{3/2}\left(16\alpha^2(\alpha^4+6\alpha^2+1)(\alpha^2+1)t^2-(9\alpha^8+48\alpha^6+70\alpha^4+1)\right)}{2(\alpha^2-1)^3(\alpha^2+1)^{5/2}}\mathcal{S}^{3/2}$$
$$\left.+\mathcal{O}(\mathcal{S}^2)\right],$$

$$I_q[\mathcal{L}] = qI[\mathcal{L}]\mathcal{S} + \mathcal{O}(\mathcal{S}^2)$$

$$I[\mathcal{L}] = \int_{U^-} \frac{dx}{2\pi i} \frac{x(x-1/x)^2\alpha^2(\alpha^2+1)}{(x^2-\alpha^2)(\alpha^2x^2-1)(\alpha^2-1)} \log\left(1 - e^{\frac{4\pi i \mathcal{L} x}{x^2-1}}\right)$$

$$x = \frac{-i \pm \sqrt{E^2-1}}{E} \Rightarrow E = \frac{-2ix}{x^2-1}$$

$$I[\mathcal{L}] = \oiiint \frac{i\sqrt{1+\mathcal{J}^2}dE}{\pi \mathcal{J} E \sqrt{1-E^2}(1+\mathcal{J}^2E^2)} \log\left(1 - e^{-2\pi \mathcal{L} E}\right)$$

$$I_0[\mathcal{L}] = \log\,\Gamma\left(\frac{\mathcal{L}}{\mathcal{J}}\right) + \frac{\sqrt{1+\mathcal{J}^2}+\mathcal{J}}{2\mathcal{J}} \log\,\mathcal{L} + \frac{(2\mathcal{L}-\mathcal{J})}{2\mathcal{J}} \log\,\mathcal{J} + \frac{\mathcal{L}}{\mathcal{J}}\gamma_{\mathrm{E}}$$
$$-\frac{\mathcal{L}}{\mathcal{J}} \log\left(1+\sqrt{1+\mathcal{J}^2}\right) - \frac{1}{2}\log\left(\frac{\sqrt{1+\mathcal{J}^2}+\mathcal{J}}{2}\right) + \frac{\sqrt{1+\mathcal{J}^2}-\mathcal{J}}{2\mathcal{J}} \log(4\pi)$$

$$I_\zeta[\mathcal{L}] = \sum_{m=1}^{\infty} \left[\frac{\sqrt{1+\mathcal{J}^2}}{\mathcal{J}} \log\left[\frac{\mathcal{L}+\sqrt{\mathcal{L}^2+m^2}}{m}\right] - \log\left[\frac{\mathcal{J}\sqrt{\mathcal{L}^2+m^2}+\mathcal{L}\sqrt{1+\mathcal{J}^2}}{\mathcal{J}m+\mathcal{L}}\right] - \frac{\mathcal{L}}{\mathcal{J}m}\right]$$

$$I_\zeta[\mathcal{L}] = \frac{1}{\mathcal{J}} \sum_{k=1}^{\infty} \frac{\zeta_{2k+1}}{2k+1} c_k(\mathcal{J}) \mathcal{L}^{2k+1}$$

$$c_k(\mathcal{J}) = \sqrt{1+\mathcal{J}^2} \sum_{n=k}^{\infty} (-1)^n \frac{\Gamma\left(\frac{1}{2}+n\right)}{\Gamma\left(\frac{1}{2}\right)\Gamma(1+n)} \mathcal{J}^{2(n-k)}$$

$$\mathcal{A}^{\mathrm{YM}}_{\mathrm{asy}} = I_{\mathrm{asy}}[\mathcal{L}_A]\mathcal{S} + \mathcal{O}(\mathcal{S}^2)$$

$$I_{\mathrm{asy}}\,[\mathcal{L}_A] = -\log\,\Gamma\left(\frac{\mathcal{L}_A}{\mathcal{J}}\right)\Gamma\left(1-\frac{\mathcal{L}_A}{\mathcal{J}}\right) + \log(2\pi)$$

$$\mathcal{A}^{\mathrm{YM}} = \mathcal{A}_1^{\mathrm{cl}}\mathcal{S} + \mathcal{O}(\mathcal{S}^2), \mathcal{B}^{\mathrm{cl}} = \mathcal{B}_1^{\mathrm{cl}}\mathcal{S} + \mathcal{O}(\mathcal{S}^2)$$

$$\mathcal{A}_1^{\mathrm{SYM}} = I_{\mathrm{asy}}[\mathcal{L}_A] + I[\mathcal{L}_A] + I[\mathcal{J}-\mathcal{L}_A], \mathcal{B}_1^{\mathrm{YM}} = I[\mathcal{J}+\mathcal{L}_B] - I[\mathcal{L}_B]$$

$$\mathcal{A}_1^{\mathrm{SYM}} = \frac{1+\delta_1}{2}\log\left(\mathcal{L}_A(\mathcal{J}-\mathcal{L}_A)\right) + \delta_1\log(4\pi) + \gamma_{\mathrm{E}}$$
$$-\log\left[\mathcal{J}(1+\delta_1)(1+\mathcal{J}\delta_1)\right] + I_\zeta[\mathcal{L}_A] + I_\zeta[\mathcal{J}-\mathcal{L}_A]$$
$$\mathcal{B}_1^{\mathrm{SYM}} = \frac{1+\delta_1}{2}\log(\mathcal{L}_B+\mathcal{J}) + \frac{1-\delta_1}{2}\log\,\mathcal{L}_B + \gamma_{\mathrm{E}}$$
$$-\log(1+\mathcal{J}\delta_1) - I_\zeta[\mathcal{L}_B] + I_\zeta[\mathcal{J}+\mathcal{L}_B]$$

$$\mathcal{N}^{\mathrm{SYM}} = -\frac{1}{2}(\log(\mathcal{S}/2)-1)\mathcal{S} + \mathcal{N}_1^{\mathrm{cl}}\mathcal{S} + \mathcal{O}(\mathcal{S}^2)$$

$$-\int_a^b \frac{(x-1/x)dx}{2\pi x}\rho(x)\log\left(\frac{2\pi\rho(x)}{e}\right) = -\frac{1}{2}\log\left[\frac{2\sqrt{1+\mathcal{J}^2}\pi^2\mathcal{S}}{e\mathcal{J}^2}\right]\mathcal{S} + \mathcal{O}(\mathcal{S}^2)$$

$$\mathcal{N}_1^{\mathrm{SYM}} = -(1+\delta_1)\log\,\mathcal{J} - \delta_1\log(4\pi) - 2\gamma_{\mathrm{E}} - \frac{1}{4}\log(1+\mathcal{J}^2) + \log\left[\mathcal{J}(1+\delta_1)(1+\mathcal{J}\delta_1)^2\right] - 2I_\zeta[\mathcal{J}]$$

$$\mathcal{D}^{\mathrm{YM}} = \mathcal{D}_1^{\mathrm{YM}}\mathcal{S} + \mathcal{O}(\mathcal{S}^2)$$

$$\mathcal{D}_1^{\mathrm{SYM}} = -\frac{1}{4}\log(1+\mathcal{J}^2) + \sum_{\mathcal{L}\in L} I_\zeta[\mathcal{L}] - 2I_\zeta[\mathcal{J}]$$

$$\mathcal{D}_1^{\mathrm{SYM}} = -\frac{1}{4}\log(1+\mathcal{J}^2) + \sum_{k=1}^{\infty}\frac{\zeta_{2k+1}}{2k+1}c_k(\mathcal{J})P_k(\mathcal{J}_1,\mathcal{J}_2,\mathcal{J})$$

$$P_k = -2\mathcal{J}^{2k} + \frac{1}{2^{2k+1}\mathcal{J}}\sum_{\sigma_1,\sigma_2=\pm}(\mathcal{J}+\sigma_1\mathcal{J}_1+\sigma_2\mathcal{J}_2)^{2k+1}$$

$$F\left[\frac{4\pi\mathcal{L}x}{x^2-1}+R(x)\right] - F\left[\frac{4\pi\mathcal{L}x}{x^2-1}\right] = \Delta\left(\partial_{\mathcal{J}},\partial_{\mathcal{L}}\right)\left[R^{(1)}(x)F'\left[\frac{4\pi\mathcal{L}x}{x^2-1}\right]\right]$$

$$\Delta\left(\partial_{\mathcal{J}},\partial_{\mathcal{L}}\right) = \mathcal{S} + \sum_{n=2}^{\infty}\mathcal{S}^n\Delta^{(n)}\left(\partial_{\mathcal{J}},\partial_{\mathcal{L}}\right)$$

$$I_q[\mathcal{L}] = q\Delta\left(\partial_{\mathcal{J}},q\partial_{\mathcal{L}}\right)I[\mathcal{L}]$$

$$\Delta^{(n)}\left(\partial_{\mathcal{J}},\partial_{\mathcal{L}}\right) = \sum_{j+l=0}^{2(n-1)} c_{j,l}^{(n)}(\mathcal{J})\partial_{\mathcal{J}}^j\partial_{\mathcal{L}}^l$$

$$c_{j,l\geqslant n}^{(n)} = c_{0,0}^{(n)} = 0.$$

$$\Delta^{(2)}\left(\partial_{\mathcal{J}},\partial_{\mathcal{L}}\right) = \frac{\sqrt{1+\mathcal{J}^2}}{4}\left(\partial_{\mathcal{J}}+\partial_{\mathcal{L}}\right)\partial_{\mathcal{J}} + \frac{\sqrt{1+\mathcal{J}^2}}{\mathcal{J}}\left(\partial_{\mathcal{J}}+\frac{1}{2}\partial_{\mathcal{L}}\right),$$

$$e^{\mathcal{J}\partial_{\mathcal{L}}}\Delta\left(\partial_{\mathcal{J}},\partial_{\mathcal{L}}\right) = \Delta\left(\partial_{\mathcal{J}},-\partial_{\mathcal{L}}\right)e^{\mathcal{J}\partial_{\mathcal{L}}} \Rightarrow \Delta\left(\partial_{\mathcal{J}},\partial_{\mathcal{L}}\right) = \Delta\left(\partial_{\mathcal{J}}+\partial_{\mathcal{L}},-\partial_{\mathcal{L}}\right).$$

$$\mathcal{A}^{\mathrm{SYM}} = \Delta\left(\partial_{\mathcal{J}},\partial_{\mathcal{L}_A}\right)\mathcal{A}_1^{\mathrm{SYM}}, \mathcal{B}^{\mathrm{SYM}} = \Delta\left(\partial_{\mathcal{J}},-\partial_{\mathcal{L}_B}\right)\mathcal{B}_1^{\mathrm{SYM}},$$

$$\mathcal{N}^{\mathrm{SYM}} = \mathcal{N}_{\mathrm{asy}}^{\mathrm{SYM}} - 2\left[\Delta\left(\partial_{\mathcal{J}},2\partial_{\mathcal{L}}\right)I[\mathcal{L}]\right]_{\mathcal{L}=\mathcal{J}}$$

$$\Delta\left(\partial_{\mathcal{J}},\pm\partial_{\mathcal{L}}\right)\left[\frac{1\pm\delta_1}{2}\log\,\mathcal{L}\right] = \frac{2\mathcal{L}+\mathcal{S}\pm(\mathcal{E}-\mathcal{J})}{2}\log\left[\frac{2\mathcal{L}+\mathcal{S}\pm(\mathcal{E}-\mathcal{J})}{2}\right] - \mathcal{L}\log\,\mathcal{L} - \frac{\mathcal{S}\pm(\mathcal{E}-\mathcal{J})}{2}$$

$$\mathcal{D}_\rho^{\mathrm{YM}} = -\int_a^b \frac{(x-1/x)dx}{4\pi x}\rho(x)\log\left[\frac{(x^2-1)^2\rho^2(x)}{2e\mathcal{S}x^2}\right]$$

$$\mathcal{D}_\zeta^{\mathrm{YM}} = \sum_{\mathcal{L}\in L}\Delta\left(\partial_{\mathcal{J}},\partial_{\mathcal{L}}\right)I_\zeta[\mathcal{L}] - 2\left[\Delta\left(\partial_{\mathcal{J}},2\partial_{\mathcal{L}}\right)I_\zeta[\mathcal{L}]\right]_{\mathcal{L}=\mathcal{J}}$$

$$\mathcal{D}_{\zeta}^{\mathrm{YM}} = \sum_{k=1}^{\infty} \zeta_{2k+1} \mathcal{D}_{\zeta_{2k+1}}^{\mathrm{cl}}$$

$$\begin{aligned}\mathcal{D}_{\rho}^{\mathrm{SYM}} =& -\frac{1}{4}\log(1+\mathcal{J}^2)\mathcal{S} - \frac{7+4\mathcal{J}^2}{16(1+\mathcal{J}^2)^{3/2}}\mathcal{S}^2 + \frac{150+120\mathcal{J}^2+29\mathcal{J}^4}{384(1+\mathcal{J}^2)^3}\mathcal{S}^3 \\ & - \frac{1785+1748\mathcal{J}^2+640\mathcal{J}^4+86\mathcal{J}^6}{3072(1+\mathcal{J}^2)^{9/2}}\mathcal{S}^4 + \mathcal{O}(\mathcal{S}^5)\end{aligned}$$

$$\begin{aligned}\mathcal{D}_{\zeta_3}^{\mathrm{SYM}} =& \frac{\sqrt{1+\mathcal{J}^2}-1}{2\mathcal{J}^2}(\mathcal{J}^2-\vec{\mathcal{J}}^2)\mathcal{S} \\ & + \left[\frac{\sqrt{1+\mathcal{J}^2}(3\mathcal{J}^2-\vec{\mathcal{J}}^2)}{4\mathcal{J}^4} - \frac{\mathcal{J}^2(3+2\mathcal{J}^2)(2+\mathcal{J}^2-\vec{\mathcal{J}}^2)-2\vec{\mathcal{J}}^2}{8\mathcal{J}^4(1+\mathcal{J}^2)}\right]\mathcal{S}^2 \\ & + \mathcal{O}(\mathcal{S}^3)\end{aligned}$$

$$Z[R_{\mathcal{L}}] = \int_{-i\infty}^{i\infty} \frac{(x+1/x)}{(2\pi)^2 i}\left[\frac{1}{2}\log\frac{\Gamma\left(1-\frac{R_{\mathcal{L}}(x)}{2\pi}\right)}{\Gamma\left(1+\frac{R_{\mathcal{L}}(x)}{2\pi}\right)} - \frac{\gamma_{\mathrm{E}} R_{\mathcal{L}}(x)}{2\pi}\right] dR_{\mathcal{L}}(x) - \left(R_{\mathcal{L}} \to \hat{R}_{\mathcal{L}}\right)$$

$$R_{\mathcal{L}}(x) = \frac{4\pi\mathcal{L}x}{x^2-1} + R(x), \hat{R}_{\mathcal{L}}(x) = \frac{4\pi\mathcal{L}x}{x^2-1}$$

$$Z[R_{\mathcal{L}}] = \Delta\left(\partial_{\mathcal{J}}, \partial_{\mathcal{L}}\right) I_{\zeta}[\mathcal{L}]$$

$$\mathcal{D}_{\zeta}^{\mathrm{SYM}} = \sum_{\mathcal{L}\in L} Z[R_{\mathcal{L}}] - Z\left[2R_{\mathcal{J}/2}\right]$$

$$\frac{1}{2}\log\frac{\Gamma\left(1-\frac{R_{\mathcal{L}}(x)}{2\pi}\right)}{\Gamma\left(1+\frac{R_{\mathcal{L}}(x)}{2\pi}\right)} - \frac{\gamma_{\mathrm{E}} R_{\mathcal{L}}(x)}{2\pi} = \sum_{k=1}^{\infty} \frac{\zeta_{2k+1}}{2k+1}\left(\frac{R_{\mathcal{L}}(x)}{2\pi}\right)^{2k+1}$$

$$Z[R_{\mathcal{L}}] = \sum_{k=1}^{\infty} \frac{\zeta_{2k+1}}{2k+1} Z_k[R_{\mathcal{L}}]$$

$$Z_k[R_{\mathcal{L}}] = -\int_{-i\infty}^{i\infty} \frac{dx(x-1/x)}{4\pi i x(k+1)}\left[\left(\frac{R_{\mathcal{L}}(x)}{2\pi}\right)^{2k+2} - \left(\frac{\hat{R}_{\mathcal{L}}(x)}{2\pi}\right)^{2k+2}\right]$$

$$c_k(\mathcal{J}) = -\int_{-i\infty}^{i\infty} \frac{dx(x-1/x)}{2\pi i x}\frac{R^{(1)}(x)}{2\pi}\left(\frac{2x}{x^2-1}\right)^{2k+1}$$

$$\mathcal{D}_1^{\mathrm{SYM}} = -\frac{\mathcal{J}^2}{4} + \frac{\left(\mathcal{J}^2-\vec{\mathcal{J}}^2\right)}{4}\zeta_3 + \mathcal{O}\left(\mathcal{J}^4, \mathcal{J}_{1,2}^4, \mathcal{J}_1^2\mathcal{J}_2^2, \mathcal{J}^2\mathcal{J}_{1,2}^2\right)$$

$$\mathcal{D}_2^{\mathrm{SYM}} = -\frac{7}{16} + \frac{1}{4}\zeta_3 + \mathcal{O}\left(\mathcal{J}^2, \vec{\mathcal{J}}^2\right)$$
$$\mathcal{D}_3^{\mathrm{SYM}} = \frac{25}{64} - \frac{3}{16}\zeta_3 - \frac{3}{16}\zeta_5 + \mathcal{O}\left(\mathcal{J}^2, \vec{\mathcal{J}}^2\right)$$
$$\mathcal{D}_4^{\mathrm{SYM}} = -\frac{595}{1024} + \frac{29}{128}\zeta_3 + \frac{9}{32}\zeta_5 + \frac{45}{256}\zeta_7 + \mathcal{O}\left(\mathcal{J}^2, \vec{\mathcal{J}}^2\right)$$

$$1 = e^{ip_k J} \prod_{j \neq k}^{S} S_{kj}$$

$$p_k = -i\log\left(\frac{x_k^+}{x_k^-}\right), x_k^{\pm} = x(u_k \pm i/2)$$

$$\mathcal{N} = \sqrt{\frac{H}{G}} \times e^W$$

$$G = \det_{1 \leqslant j,k \leqslant S}\left[\frac{\partial}{\partial u_j}\left(Jp_k - i\sum_{l \neq k}^{S} \log S_{kl}\right)\right]$$

$$H = \prod_{k<j}^{S} \frac{\left(u_k - u_j\right)^2}{\left(u_k - u_j\right)^2 + 1} \exp\left[\sum_{k,j}^{S} \log \frac{\left(x_k^+ x_j^- - 1\right)\left(x_k^- x_j^+ - 1\right)}{\left(x_k^+ x_j^+ - 1\right)\left(x_k^- x_j^- - 1\right)}\right].$$

$$n_k = \frac{2\mathcal{J}x_k}{x_k^2 - 1} + \sum_{j \neq k}^{S} \frac{4x_k x_j\left(x_k x_j - 1\right)}{\sqrt{\lambda}(x_k^2 - 1)\left(x_k - x_j\right)\left(x_j^2 - 1\right)} + \mathcal{O}\left(\frac{1}{\lambda}\right)$$

$$x_k^{(0)} = \alpha = \mathcal{J} + \sqrt{1 + \mathcal{J}^2}, k = 1, \ldots, S/2$$

$$x_k = x_k^{(0)} + \frac{x_k^{(1/2)}}{\lambda^{1/4}} + \frac{x_k^{(1)}}{\lambda^{1/2}} + \frac{x_k^{(3/2)}}{\lambda^{3/4}} + \cdots$$

$$H_{S/2}\left(\frac{x_k^{(1/2)}}{\sqrt{2}\beta}\right) = 0, k = 1, \ldots, S/2$$

$$G = G_+ G_-\left(1 + \mathcal{O}\left(\frac{1}{\lambda}\right)\right)$$

$$\sqrt{G} \approx \left[\frac{4\pi^2\sqrt{1 + \mathcal{J}^2}}{\mathcal{J}^2\sqrt{\lambda}}\right]^{S/2}$$
$$\times \det_{1 \leqslant j,k \leqslant S/2}\left[\delta_{k=j}\left(1 + \sum_{l \neq k}^{S/2} \frac{2\beta^2}{\sqrt{\lambda}(x_k - x_l)^2}\right) - \delta_{k \neq j}\frac{2\beta^2}{\sqrt{\lambda}\left(x_k - x_j\right)^2}\right]$$

$$\sqrt{G} = \left[\frac{4\pi^2\sqrt{1 + \mathcal{J}^2}}{\mathcal{J}^2\sqrt{\lambda}}\right]^{S/2} \Gamma\left(1 + \frac{S}{2}\right) \exp\left(\sum_{n \geqslant 1} \frac{P_n^G(S)}{(\sqrt{\lambda})^n}\right)$$

$$\sqrt{H}=\exp\left(\sum_{n\geqslant 1}\frac{P_n^H(S)}{(\sqrt{\lambda})^n}\right)$$

$$W=\left(W^{(0)}+\frac{W^{(1)}}{\sqrt{\lambda}}+\frac{W^{(2)}}{\lambda}+\cdots\right)S+\mathcal{O}(S^2)$$

$$W^{(i)}=-2\left[\widehat{\Delta}^{(i)}I[\mathcal{L}]\right]_{\mathcal{L}=\mathcal{J}}$$

$$\widehat{\Delta}^{(1)}=-\frac{\sqrt{1+\mathcal{J}^2}}{2}(\partial_{\mathcal{J}}+\partial_{\mathcal{L}})\partial_{\mathcal{J}}-\frac{\sqrt{1+\mathcal{J}^2}}{\mathcal{J}}\left(\partial_{\mathcal{J}}+\frac{1}{2}\partial_{\mathcal{L}}\right)+\frac{1}{2\mathcal{J}}\left(\mathcal{J}\sqrt{1+\mathcal{J}^2}-\mathcal{R}\right)\partial_{\mathcal{L}}^2$$

$$\mathcal{R}[f]=\mathrm{res}_{\mathcal{J}=0}f(\mathcal{J})$$

$$\mathcal{N}=\frac{2^{2S}\lambda^{3S/2+\gamma/2}}{(4\pi)^{S+\gamma}e^{2\gamma_{\mathrm{E}}S}}\sqrt{\frac{\Gamma(J)\Gamma(J-1)}{\Gamma(\Delta+S)\Gamma(\Delta+S-1)}}\times\frac{\mathcal{D}_{\mathcal{N}}}{\lambda^{S/4}\Gamma\left(1+\frac{S}{2}\right)}$$

$$\Delta=\sqrt{\lambda}\mathcal{J}+S\left[\frac{\sqrt{1+\mathcal{J}^2}}{\mathcal{J}}-\frac{1}{2\mathcal{J}(1+\mathcal{J}^2)\sqrt{\lambda}}+\frac{4\mathcal{J}^2-1}{8\mathcal{J}(1+\mathcal{J}^2)^{5/2}\lambda}\right]+\mathcal{O}\left(S^2,\frac{S}{\lambda^{3/2}}\right)$$

$$\log\mathcal{D}_{\mathcal{N}}=S\left[c^{(0)}+\frac{c^{(1)}}{\sqrt{\lambda}}+\frac{c^{(2)}}{\lambda}\right]+\mathcal{O}\left(S^2,\frac{S}{\lambda^{3/2}}\right)$$

$$\begin{aligned}c^{(0)}&=-\frac{1}{4}\log(1+\mathcal{J}^2)+K[\mathcal{J}]\\ c^{(1)}&=\frac{5+2\mathcal{J}^2}{8(1+\mathcal{J}^2)^{3/2}}+\left[\widehat{\Delta}^{(1)}K[\mathcal{L}]\right]_{\mathcal{L}=\mathcal{J}}\\ c^{(2)}&=\frac{57-12\mathcal{J}^2+4\mathcal{J}^4}{96(1+\mathcal{J}^2)^3}+\left[\widehat{\Delta}^{(2)}K[\mathcal{L}]\right]_{\mathcal{L}=\mathcal{J}}\end{aligned}$$

$$K[\mathcal{L}]=\log\left[\frac{1}{4}\left(1+\sqrt{1+\mathcal{J}^2}\right)^2\left(\mathcal{J}+\sqrt{1+\mathcal{J}^2}\right)\right]-2I_{\zeta}[\mathcal{L}]$$

$$\log\mathcal{D}_{\mathcal{N}}=S\left[-\frac{1}{J}-\frac{\frac{7}{8}-J}{\sqrt{\lambda}}-\frac{\left(\frac{15}{32}-\frac{J}{2}-\frac{J^2}{4}\right)+\left(\frac{5}{24}-\frac{J^2}{3}\right)\zeta_3}{\lambda}\right]+\mathcal{O}\left(S^2,\frac{S}{\lambda^{3/2}}\right)$$

$$\begin{aligned}\log\mathcal{D}&=\log\mathcal{D}_{\mathcal{N}}+\log\mathcal{D}_{\mathcal{A}}+\log\mathcal{D}_{\mathcal{B}}\\ &=S\left[\frac{5}{8\sqrt{\lambda}}+\frac{(19-8J^2)+8(1+J^2-J_1^2-J_2^2)\zeta_3}{32\lambda}\right]+\mathcal{O}\left(S^2,\frac{S}{\lambda^{3/2}}\right)\end{aligned}$$

$$\log \mathcal{D}_{J_1 J_2 J}$$
$$= \frac{1}{\sqrt{\lambda}}\left[\frac{5}{8}S - \frac{7-4\zeta_3}{16}S^2\right]$$
$$+\frac{1}{\lambda}\left[\frac{(19-8J^2)+8\left(1+J^2-\vec{J}^2\right)\zeta_3}{32}S - \frac{49-8\zeta_3}{64}S^2 + \frac{25-12\zeta_3-12\zeta_5}{64}S^3\right]$$
$$+\mathcal{O}\left(\frac{1}{\lambda^{3/2}}\right)$$

$$\mathrm{S}_a(x) = \sum_{i=1}^{x} \frac{(\mathrm{sgn}a)^i}{i^a},\ \mathrm{S}_{a,b,c,\ldots}(x) = \sum_{i=1}^{x} \frac{(\mathrm{sgn}a)^i}{i^a}\ \mathrm{S}_{b,c,\ldots}(i),$$

$$\mathrm{S}_{2,3}(s) = \sum_{i_1 \le i_2 \le s} \frac{1}{i_2^2 i_1^3} = \sum_{i_1 \le i_2 \le s} \int_0^\infty \prod_{n=1}^{2} dt_n \frac{t_1^2 t_2}{2} e^{-t_1 i_1 - t_2 i_2}$$

$$\mathrm{S}_{2,3}(s) = \int_0^1 \prod_{n=1}^{2} \frac{dx_n}{x_n} \frac{x_1 x_2 \left(1 - x_2 - (-1+x_1)x_1^s x_2^{1+s} + x_1^s(-1+x_1 x_2)\right)\log\, x_1 \log^2\, x_2}{2(-1+x_1)(-1+x_2)(-1+x_1 x_2)}$$

$$\mathrm{S}_{2,3}(s) = \left(\frac{8}{7}\zeta_2^3 - 2\zeta_3^2\right)s + \mathcal{O}(s^2)$$

$$\mathcal{B}(\ell_B = 1) = 1 + g^4(c_{1|4} + c_{2|4}\zeta_3) + g^6(c_{1|6} + c_{2|6}\zeta_3 + c_{3|6}\zeta_5) + \mathcal{O}(g^8),$$

$c_{1|4} = 4\left(\mathrm{S}_{-2}^2 - 2\,\mathrm{S}_{-3}\,\mathrm{S}_1 - 2\,\mathrm{S}_{-2}\,\mathrm{S}_1^2 - 2\,\mathrm{S}_1\,\mathrm{S}_3 - \mathrm{S}_4 + 2\,\mathrm{S}_{-3,1} + 4\,\mathrm{S}_1\,\mathrm{S}_{-2,1} + 2\,\mathrm{S}_{-2,2} + 2\,\mathrm{S}_{3,1} - 4\,\mathrm{S}_{-2,1,1}\right)$
$c_{2|4} = 24\,\mathrm{S}_1,$
$c_{1|6} = \frac{32}{3}\Big(-6\,\mathrm{S}_{-6} + 3\,\mathrm{S}_{-3}^2 - 30\,\mathrm{S}_{-5}\,\mathrm{S}_1 - 6\,\mathrm{S}_{-4}\left(\mathrm{S}_{-2} - 3\,\mathrm{S}_1^2\right) + 5\,\mathrm{S}_3^2 + 6\,\mathrm{S}_6 + 30\,\mathrm{S}_{-5,1} - 12\,\mathrm{S}_{-4,2} - 24\,\mathrm{S}_{-3,3} +$
$\mathrm{S}_1^3\left(\mathrm{S}_3 - 6\,\mathrm{S}_{-2,1}\right) - 48\,\mathrm{S}_3\,\mathrm{S}_{-2,1} + 12\,\mathrm{S}_{-2,1}^2 - 3\,\mathrm{S}_{-3}\left(4\,\mathrm{S}_{-2}\,\mathrm{S}_1 - 3\,\mathrm{S}_1^3 + 11\,\mathrm{S}_1\,\mathrm{S}_2 + 4\left(-4\,\mathrm{S}_3 + \mathrm{S}_{-2,1}\right)\right) +$
$54\,\mathrm{S}_{4,-2} - 6\,\mathrm{S}_{4,2} + 6\,\mathrm{S}_{5,1} - 48\,\mathrm{S}_{-4,1,1} + 6\,\mathrm{S}_{-2}\left(\mathrm{S}_1^2\,\mathrm{S}_2 - 9\,\mathrm{S}_4 - 10\,\mathrm{S}_1\,\mathrm{S}_{2,1} + 2\left(\mathrm{S}_{-3,1} + \mathrm{S}_{-2,2} + 4\,\mathrm{S}_{3,1} -$
$2\,\mathrm{S}_{-2,1,1}\right)\Big) + 12\,\mathrm{S}_2\left(3\,\mathrm{S}_{-3,1} + \mathrm{S}_{3,1} - 6\,\mathrm{S}_{-2,1,1}\right) + 6\,\mathrm{S}_1^2\left(2\,\mathrm{S}_4 - 5\,\mathrm{S}_{-3,1} - 4\,\mathrm{S}_{-2,2} - \mathrm{S}_{3,1} + 6\,\mathrm{S}_{-2,1,1}\right) +$
$36\,\mathrm{S}_{-2,2,2} - 48\,\mathrm{S}_{-2,3,1} - 36\,\mathrm{S}_{2,-3,1} - 12\,\mathrm{S}_{2,3,1} - 48\,\mathrm{S}_{3,1,-2} + 12\,\mathrm{S}_{4,1,1} - 72\,\mathrm{S}_{-3,1,1,1} - 3\,\mathrm{S}_1\big(3\mathrm{S}_2\left(\mathrm{S}_3 -$
$6\,\mathrm{S}_{-2,1}\right) + 2\left(\mathrm{S}_5 + 5\,\mathrm{S}_{-4,1} - 6\,\mathrm{S}_{-2,3} + 4\,\mathrm{S}_{2,-3} - 2\,\mathrm{S}_{2,3} + 3\,\mathrm{S}_{4,1} - 11\,\mathrm{S}_{-3,1,1} - 2\,\mathrm{S}_{-2,1,-2} - 10\,\mathrm{S}_{-2,2,1} -$
$10\,\mathrm{S}_{2,1,-2} - 3\,\mathrm{S}_{3,1,1} + 18\,\mathrm{S}_{-2,1,1,1}\right)\big) + 72\,\mathrm{S}_{2,-2,1,1} - 24\,\mathrm{S}_{3,1,1,1} + 144\,\mathrm{S}_{-2,1,1,1,1}\Big)$
$c_{2|6} = -\frac{32}{3}\left(6\,\mathrm{S}_{-3} + 15\,\mathrm{S}_{-2}\,\mathrm{S}_1 - 4\,\mathrm{S}_1^3 + 9\,\mathrm{S}_1\,\mathrm{S}_2 + 4\,\mathrm{S}_3 - 12\,\mathrm{S}_{-2,1}\right)$
$c_{3|6} = -240\,\mathrm{S}_1.$

$$\mathcal{B}(\ell_B = 1) = 1 + s[3g^4(4\zeta_2\zeta_3 + 5\zeta_5) - 48g^6(\zeta_3\zeta_4 + \zeta_2\zeta_5 + 7\zeta_7)] + \mathcal{O}(sg^8, s^2),$$

$$\mathcal{B}(\ell_B = 2) = 1 + g^6(c_{1|6} + c_{2|6}\zeta_3 + c_{3|6}\zeta_5) + \mathcal{O}(g^8),$$

$$c_{1|6} = \frac{8}{3}\big(-6\,S_{-6} + 9\,S_{-3}^2 - 12\,S_{-5}\,S_1 - 7\,S_3^2 + 24\,S_{-5,1} - 6\,S_{-4,2} - 12\,S_{-3,3} + 2\,S_{-3}(\,S_1^3 - 9\,S_1\,S_2 + 11\,S_3 - 6\,S_{-2,1}) - 20\,S_3\,S_{-2,1} + 12\,S_{-2,1}^2 - 2\,S_1^3(\,S_3 + 2\,S_{-2,1}) + 6\,S_{-2}(-5\,S_4 + 2\,S_{-3,1} + S_{-2,2} + 4\,S_{3,1}) + 30\,S_{4,-2} + 12\,S_{4,2} - 12\,S_{5,1} - 36\,S_{-4,1,1} - 12\,S_{-3,-2,1} - 12\,S_{-3,1,-2} + 6\,S_1^2\big(\,S_{-4} - S_4 - 2(\,S_{-3,1} + S_{-2,2} - S_{3,1} - 2\,S_{-2,1,1})\big) + 24\,S_2(\,S_{-3,1} - S_{3,1} - 2\,S_{-2,1,1}) - 6\,S_{-2,2,-2} + 24\,S_{-2,2,2} - 36\,S_{-2,3,1} - 24\,S_{2,-3,1} + 24\,S_{2,3,1} - 24\,S_{3,1,-2} - 12\,S_{4,1,1} - 48\,S_{-3,1,1,1} - 24\,S_{-2,-2,1,1} + 6\,S_1\big(2\,S_5 - 2\,S_{-4,1} - S_{-2}\,S_{-2,1} + 3\,S_2(\,S_3 + 2\,S_{-2,1}) + 3\,S_{-2,3} - 3\,S_{2,-3} - 6\,S_{-2}\,S_{2,1} - 3\,S_{2,3} + 3\,S_{4,1} + 6\,S_{-3,1,1} + S_{-2,1,-2} + 6\,S_{-2,2,1} + 6\,S_{2,1,-2} - 6\,S_{3,1,1} - 12\,S_{-2,1,1,1}\big) + 48\,S_{2,-2,1,1} + 48\,S_{3,1,1,1} + 96\,S_{-2,1,1,1,1}\big),$$

$$c_{2|6} = \frac{16}{3}\big(3\,S_{-3} + 3\,S_{-2}\,S_1 + S_1^3 - S_3 - 6\,S_{-2,1}\big),$$

$$c_{3|6} = 80\,S_1.$$

$$\mathcal{B}(\ell_B = 2) = 1 + 4sg^6(-3\zeta_3\zeta_4 + 9\zeta_2\zeta_5 + 14\zeta_7) + \mathcal{O}(sg^8, s^2)$$

$$\mathcal{A} = \mathcal{A}_{\text{asy}} + \delta\mathcal{A}$$

$$\mathcal{A}_{\text{asy}} = \sum_{\alpha\cup\bar{\alpha}=\mathbf{u}} (-1)^{|\bar{\alpha}|} \prod_{j\in\bar{\alpha}} e^{ip_j\ell_A} \prod_{i\in\alpha, j\in\bar{\alpha}} \frac{1}{h_{ij}},$$

$$\mathcal{A}_{\text{asy}} = \frac{(2s)!}{(s!)^2}\Big(1 + g^2 c_{1|2} + g^4 c_{1|4} + g^6(c_{1|6} + c_{3|6}\zeta_3) + \mathcal{O}(g^8)\Big),$$

$$c_{1|2} = -4\Big(\,S_2 + 2\,S_1(\,S_1 - \tilde{S}_1)\Big),$$

$$c_{1|4} = 2\big(4\,S_{-4} + 4\,S_{-2}^2 + 4\,S_{-3}\,S_1 + 16\,S_1^4 + 40\,S_1^2\,S_2 + 4\,S_2^2 + 15\,S_1\,S_3 - S_4 + 4\,S_{-3,1} + 8\,S_1\,S_{-2,1} + 8\,S_{-2,2} + 5\,S_{1,3} + 9\,S_{3,1} - 24\,S_{-2,1,1} - 8\,S_{-3}\,\tilde{S}_1 - 32\,S_1^3\,\tilde{S}_1 - 32\,S_1\,S_2\,\tilde{S}_1 - 8\,S_3\,\tilde{S}_1 + 16\,S_{-2,1}\,\tilde{S}_1 + 16\,S_1^2\,\tilde{S}_1^2 + 4\,S_{-2}(\,S_1^2 + S_2 - 4\,S_1\,\tilde{S}_1) - 16\,S_1^2\,\tilde{S}_2\big),$$

$$c_{1|6} = -\frac{32}{3}(12\,S_{-6} - 6\,S_{-3}^2 + 12\,S_{-4}\,S_{-2} + 51\,S_{-5}\,S_1 + 24\,S_{-4}\,S_1^2 + 6\,S_{-2}^2\,S_1^2 + 6\,S_{-2}\,S_1^4 + 8\,S_1^6 + 9\,S_{-4}\,S_2 + 3\,S_{-2}^2\,S_2 + 27\,S_{-2}\,S_1^2\,S_2 + 48\,S_1^4\,S_2 + 3\,S_{-2}\,S_2^2 + 48\,S_1^2\,S_2^2 + S_2^3 + 18\,S_{-2}\,S_1\,S_3 + 45\,S_1^3\,S_3 + 48\,S_1\,S_2\,S_3 + S_3^2 + 114\,S_{-2}\,S_4 + 30\,S_1^2\,S_4 + 3\,S_2\,S_4 + 27\,S_1\,S_5 - S_6 - 54\,S_{-5,1} + 6\,S_1\,S_{-4,1} + 6\,S_{-4,2} - 24\,S_{-2}\,S_{-3,1} + 12\,S_1^2\,S_{-3,1} - 57\,S_2\,S_{-3,1} + 66\,S_{-3,3} - 6\,S_{-2}\,S_1\,S_{-2,1} + 2\,S_1^3\,S_{-2,1} - 72\,S_1\,S_2\,S_{-2,1} + 100\,S_3\,S_{-2,1} - 24\,S_{-2,1}^2 - 24\,S_{-2}\,S_{-2,2} + 12\,S_1^2\,S_{-2,2} - 6\,S_2\,S_{-2,2} - 42\,S_1\,S_{-2,3} + 36\,S_1\,S_{2,-3} + 60\,S_{-2}\,S_1\,S_{2,1} - 12\,S_1\,S_{2,3} - 108\,S_{-2}\,S_{3,1} + 12\,S_1^2\,S_{3,1} - 9\,S_2\,S_{3,1} - 108\,S_{4,-2} + 18\,S_1\,S_{4,1} + 6\,S_{4,2} - 6\,S_{5,1} + 120\,S_{-4,1,1} - 66\,S_1\,S_{-3,1,1} - 6\,S_1\,S_{-2,1,-2} + 48\,S_{-2}\,S_{-2,1,1} - 24\,S_1^2\,S_{-2,1,1} + 150\,S_2\,S_{-2,1,1} - 60\,S_1\,S_{-2,2,1} - 60\,S_{-2,2,2} + 108\,S_{-2,3,1} + 48\,S_{2,-3,1} - 60\,S_1\,S_{2,1,-2} + 12\,S_{2,3,1} + 108\,S_{3,1,-2} - 18\,S_1\,S_{3,1,1} - 12\,S_{4,1,1} + 120\,S_{-3,1,1,1} + 108\,S_1\,S_{-2,1,1,1} - 144\,S_{2,-2,1,1} + 24\,S_{3,1,1,1} - 336\,S_{-2,1,1,1,1} + 18\,S_{-5}\,\tilde{S}_1 - 54\,S_{-4}\,S_1\,\tilde{S}_1 - 12\,S_{-2}^2\,S_1\,\tilde{S}_1 - 30\,S_{-2}\,S_1^3\,\tilde{S}_1 - 24\,S_1^5\,\tilde{S}_1 - 42\,S_{-2}\,S_1\,S_2\,\tilde{S}_1 - 84\,S_1^3\,S_2\,\tilde{S}_1 - 30\,S_1\,S_2^2\,\tilde{S}_1 - 12\,S_{-2}\,S_3\,\tilde{S}_1 - 54\,S_1^2\,S_3\,\tilde{S}_1 - 12\,S_2\,S_3\,\tilde{S}_1 - 24\,S_1\,S_4\,\tilde{S}_1 - 6\,S_5\,\tilde{S}_1 + 36\,S_{-4,1}\,\tilde{S}_1 + 66\,S_1\,S_{-3,1}\,\tilde{S}_1 + 36\,S_1^2\,S_{-2,1}\,\tilde{S}_1 - 48\,S_2\,S_{-2,1}\,\tilde{S}_1 + 48\,S_1\,S_{-2,2}\,\tilde{S}_1 - 36\,S_{-2,3}\,\tilde{S}_1 + 36\,S_{2,-3}\,\tilde{S}_1 + 72\,S_{-2}\,S_{2,1}\,\tilde{S}_1 - 6\,S_1\,S_{3,1}\,\tilde{S}_1 - 72\,S_{-3,1,1}\,\tilde{S}_1 - 60\,S_1\,S_{-2,1,1}\,\tilde{S}_1 - 72\,S_{-2,2,1}\,\tilde{S}_1 - 72\,S_{2,1,-2}\,\tilde{S}_1 + 144\,S_{-2,1,1,1}\,\tilde{S}_1 + 24\,S_{-2}\,S_1^2\,\tilde{S}_1^2 + 24\,S_1^4\,\tilde{S}_1^2 + 36\,S_1^2\,S_2\,\tilde{S}_1^2 + 12\,S_1\,S_3\,\tilde{S}_1^2 - 24\,S_1\,S_{-2,1}\,\tilde{S}_1^2 - 8\,S_1^3\,\tilde{S}_1^3 + S_{-3}(24\,S_{-2}\,S_1 + 13\,S_1^3 - 54\,S_1^2\,\tilde{S}_1 + 4(-25\,S_3 + 6\,S_{-2,1} + 6\,S_2\,\tilde{S}_1) + 12\,S_1(4\,S_2 + \tilde{S}_1^2 - \tilde{S}_2)) - 24\,S_{-2}\,S_1^2\,\tilde{S}_2 - 24\,S_1^4\,\tilde{S}_2 - 36\,S_1^2\,S_2\,\tilde{S}_2 - 12\,S_1\,S_3\,\tilde{S}_2 + 24\,S_1\,S_{-2,1}\,\tilde{S}_2 + 24\,S_1^3\,\tilde{S}_1\,\tilde{S}_2 - 16\,S_1^3\,\tilde{S}_3),$$

$$c_{3|6} = -32\,S_1(\,S_{-2} + S_2),$$

$$\mathcal{A}_{\text{asy}} = 1 \ + s[-8\zeta_3 g^2 + g^4(-32\zeta_2\zeta_3 + 90\zeta_5) + g^6(160\zeta_3\zeta_4 + 448\zeta_2\zeta_5 - 1120\zeta_7)] + \mathcal{O}(sg^8, s^2)$$

$$\frac{\delta\mathcal{A}}{\mathcal{A}_{\text{asy}}} = \frac{2\delta\gamma}{\gamma} + \mathcal{O}(g^8)$$

$$\frac{\delta\mathcal{A}}{\mathcal{A}_{\text{asy}}} = (-160\ \mathrm{S}_1\zeta_5 + \cdots)g^6 + \mathcal{O}(g^8)$$

$$\frac{\delta\mathcal{A}}{\mathcal{A}_{\text{asy}}} = -160\zeta_2\zeta_5 g^6 s + \mathcal{O}(g^8 s, g^6 s^2)$$

$$2\pi n_k = -iJ\log\left(\frac{u_k + \frac{i}{2}}{u_k - \frac{i}{2}}\right) - i\sum_{j\neq k}^{S}\log\left(\frac{u_k - u_j + i}{u_k - u_j - i}\right).$$

$$u_k = \pm\frac{J}{2\pi}\left(1 + \frac{u_k^{(1/2)}}{\sqrt{J}} + \frac{u_k^{(1)}}{J} + \mathcal{O}\left(\frac{1}{J^{3/2}}\right)\right)$$

$$e^{ip_k} = \frac{u_k + \frac{i}{2}}{u_k - \frac{i}{2}} + \mathcal{O}(g^2), h(u_k, u_j) = \frac{u_k - u_j}{u_k - u_j - i} + \mathcal{O}(g^2)$$

$$\mathcal{A}_{\text{asy}}(\ell_A) = \left(\frac{2\pi}{J}\right)^S \frac{\Gamma(\ell_A + S)}{\Gamma(\ell_A)}\exp\left(\sum_{k=1}^{\infty}\frac{P_k(S,\ell_A)}{J^k}\right)$$

$$P_1(S,\ell_A) = \frac{S(2-3S)}{4}, P_2(S,\ell_A) = \frac{S\left(4 - 24S + 17S^2 + 4\pi^2\left(S - 2\ell_A(\ell_A+1)\right)\right)}{48}.$$

$$\hat{\mathcal{A}}_{\text{asy}}(z) = \sum_{\ell_A=1}^{\infty} z^{\ell_A}\mathcal{A}_{\text{asy}}(\ell_A)$$

$$\hat{\mathcal{A}}_{\text{asy}}(z) = \frac{z}{1-z}\left(1 + Sa_J(z) + \mathcal{O}(S^2)\right)$$

$$a_J(z) = \log\left(\frac{2\pi}{J(1-z)}\right) + \psi(1) + \frac{1}{2J} + \frac{1}{J^2}\left(\frac{1}{12} - \frac{\pi^2}{3(1-z)^2}\right) + \mathcal{O}\left(\frac{1}{J^3}\right)$$

$$f_J(\ell) = \sum_{k=1}^{J/2-1}\frac{(-1)^{k+\ell+1}\Gamma(J)\Gamma(2k)\zeta_{2k+1}}{(2\pi)^{2k}\Gamma(-\ell)\Gamma(J-2k)\Gamma(1+\ell+2k)}$$

$$\sum_{\ell_A=1}^{\infty} z^{\ell_A}F_J(-\ell_A) = \hat{f}_J(z) + z^J\hat{f}_J(1/z)$$

$$\hat{f}_J(z) = \sum_{\ell_A=1}^{\infty} z^{\ell_A}f_J(-\ell_A)$$

$$\hat{f}_J(z)=\frac{z}{2(1-z)}\int_0^\infty \frac{dt}{e^t-1}\left[\left(1+(1-z)\frac{it}{2\pi}\right)^{J-1}+\left(1-(1-z)\frac{it}{2\pi}\right)^{J-1}-2\right]$$

$$\left(1\pm(1-z)\frac{it}{2\pi}\right)^{J-1}=D_\tau\cdot e^{\pm i\tau t}$$

$$\tau=\frac{(J-1)(1-z)}{2\pi},$$

$$D_\tau=\exp\left[-\sum_{n=2}^{\infty}\frac{(J-1)}{n}\left(-\frac{\tau}{J-1}\right)^n\partial_\tau^n\right].$$

$$\hat{f}_J(z)=\frac{z}{2(1-z)}D_\tau\cdot(2\psi(1)-\psi(1+i\tau)-\psi(1-i\tau))$$

$$\hat{f}_J(z)=\frac{z}{1-z}a_J(z)$$

$$\left\langle \mathrm{Tr}Z_1^2(x_1)\mathrm{Tr}Z_2^2(x_2)\mathrm{Tr}Z_3^p(x_3)\mathrm{Tr}Z_4^p(x_4)\right\rangle_{\mathrm{conn}}\propto \mathcal{G}_{22pp}(U,V),$$

$$U=\frac{x_{12}^2x_{34}^2}{x_{13}^2x_{24}^2}=z\bar{z},V=\frac{x_{14}^2x_{23}^2}{x_{13}^2x_{24}^2}=(1-z)(1-\bar{z})$$

$$z=\sigma e^{\rho},\bar{z}=\sigma e^{-\rho}$$

$$\mathcal{G}_{22pp}\propto\sigma^2\to 0$$

$$\mathcal{G}_{22pp}=2\pi i\int_{-\infty}^{\infty}d\nu\sigma^{-1-S(\nu)}\alpha(\nu)\Omega_{i\nu}(\rho)+\cdots$$

$$S=-\frac{\nu^2+4}{2\sqrt{\lambda}}\left(1+\frac{1}{2\sqrt{\lambda}}\right)+\mathcal{O}\left(\frac{1}{\lambda^{3/2}}\right)$$

$$\Omega_{i\nu}(\rho)=\frac{\nu\sin(\nu\rho)}{4\pi^2\sinh\rho}$$

$$\alpha(\nu)=-\frac{2^{S-1}\pi^2S'e^{\frac{i\pi S}{2}}}{\nu\sin\left(\frac{\pi S}{2}\right)}\gamma_S(\nu)\gamma_S(-\nu)K_{2+\Delta,2+S}b_{2+S}$$

$$\gamma_S(\nu)=\Gamma\left(2+\frac{S+i\nu}{2}\right)\Gamma\left(p+\frac{S+i\nu}{2}\right)$$

$$K_{2+\Delta,2+S}=\frac{\Gamma(\Delta+S+4)\Gamma(\Delta+S+3)}{4^{S+1}\Gamma\left(2+\frac{\Delta+S}{2}\right)^5\Gamma\left(p+\frac{\Delta+S}{2}\right)\Gamma\left(2+\frac{S-\Delta}{2}\right)\Gamma\left(p+\frac{S-\Delta}{2}\right)}$$

$$b_{2+S}=\frac{2^{S-6}(\Delta+S)^2(\Delta+S+2)^2}{(\Delta+S-1)(\Delta+S+1)^2(\Delta+S+3)}\times C_{222}(S)C_{pp2}(S)$$

$$K_{2+\Delta,2+S}b_{2+S} = \frac{\mathcal{D}_{222}\mathcal{D}_{pp2}}{Z(p)(2\sqrt{\lambda})^S\Gamma\left(1+\frac{S}{2}\right)^2}$$

$$Z(p) = \frac{(p-1)!\,(p-2)!}{2}$$

$$\mathcal{G}_{22pp} = \frac{\pi^2 i\sqrt{\lambda}}{Z(p)}\int_{-\infty}^{\infty}\frac{S'dv}{v}\frac{\gamma_S(v)\gamma_S(-v)\Omega_{iv}(\rho)}{(\sqrt{\lambda}\sigma)^{1+S}}\frac{e^{\frac{i\pi S}{2}}\Gamma\left(-\frac{S}{2}\right)}{\Gamma\left(1+\frac{S}{2}\right)}\mathcal{D}_{222}\mathcal{D}_{pp2}$$

$$\mathcal{G}_{22pp} = \mathcal{G}_{22pp}^{\mathrm{LO}} + \frac{1}{\sqrt{\lambda}}\mathcal{G}_{22pp}^{\mathrm{NLO}} + \mathcal{O}\left(\frac{1}{\lambda}\right)$$

$$\mathcal{G}_{22pp}^{\mathrm{LO/NLO}} = -\frac{(2\pi)^2 i}{Z(p)\sigma}\int\ dv\Omega_{iv}(\rho)\gamma_0(v)\gamma_0(-v)f^{\mathrm{LO/NLO}}$$

$$f^{\mathrm{LO}} = \frac{1}{v^2+4}$$

$$f^{\mathrm{NLO}} = \frac{1}{4}\left[\log(\sqrt{\lambda}\sigma) - \frac{i\pi}{2} - \gamma_{\mathrm{E}} - \psi\left(2+\frac{iv}{2}\right) - \psi\left(p+\frac{iv}{2}\right) + (v\to -v)\right].$$

$$\mathcal{G}_{22pp}^{\mathrm{dS}} \approx -\frac{2\pi^2 i}{Z(p)}\sum_{k=1}^{\infty}\left(\frac{1}{\xi}\right)^{1+2k}\zeta(1+2k)\int_{-\infty}^{\infty}dv\Omega_{iv}(\rho)\gamma_{2k}(v)\gamma_{2k}(-v)$$

$$\sum_{k=1}^{\infty}f(k) = \int_{\epsilon-i\infty}^{\epsilon+i\infty}\frac{ie^{i\pi k}dk}{2\sin\ \pi k}f(k)$$

$$R(x) = \sum_{k=1}^{\infty}\mathcal{S}^k R^{(k)}(x)$$

$$R^{(3)}=\frac{2\pi x\alpha^4}{(\alpha^2-1)^5(\alpha^2+1)^5(x^2-\alpha^2)^5}$$
$$\times[(\alpha^{10}+x^8)(3+34\alpha^2+157\alpha^4+124\alpha^6+157\alpha^8+34\alpha^{10}+3\alpha^{12})$$
$$-x^2\alpha^6(-13-5\alpha^2+275\alpha^4+835\alpha^6+417\alpha^8+297\alpha^{10}+217\alpha^{12}+25\alpha^{14})$$
$$+x^4\alpha^4(73+111\alpha^2+189\alpha^4+1163\alpha^6+1163\alpha^8+189\alpha^{10}+111\alpha^{12}+73\alpha^{14})$$
$$-x^6\alpha^2(25+217\alpha^2+297\alpha^4+417\alpha^6+835\alpha^8+275\alpha^{10}-5\alpha^{12}-13\alpha^{14})]$$
$$R^{(4)}=-\frac{\pi x\alpha^5}{2(\alpha^2-1)^7(\alpha^2+1)^8(x^2-\alpha^2)^7}$$
$$\times[(\alpha^{14}+x^{12})(21+342\alpha^2+2689\alpha^4+10536\alpha^6+13674\alpha^8+27396\alpha^{10}$$
$$+13674\alpha^{12}+10536\alpha^{14}+2689\alpha^{16}+342\alpha^{18}+21\alpha^{20})$$
$$-x^2\alpha^{10}(1+107\alpha^2+2147\alpha^4+20353\alpha^6+73514\alpha^8+94222\alpha^{10}$$
$$+161318\alpha^{12}+77986\alpha^{14}+38053\alpha^{16}+20087\alpha^{18}+3495\alpha^{20}+237\alpha^{22})$$
$$+x^4\alpha^8(-1167-3288\alpha^2+13145\alpha^4+88042\alpha^6+208498\alpha^8+204872\alpha^{10}$$
$$+363178\alpha^{12}+257948\alpha^{14}+59581\alpha^{16}+24704\alpha^{18}+12125\alpha^{20}+1162\alpha^{22})$$
$$-2x^6\alpha^6(1+\alpha^2)(1169+350\alpha^2+2573\alpha^4+55048\alpha^6+127330\alpha^8$$
$$+36660\alpha^{10}+127330\alpha^{12}+55048\alpha^{14}+2573\alpha^{16}+350\alpha^{18}+1169\alpha^{20})$$
$$+x^8\alpha^4(1162+12125\alpha^2+24704\alpha^4+59581\alpha^6+257948\alpha^8+363178\alpha^{10}$$
$$+204872\alpha^{12}+208498\alpha^{14}+88042\alpha^{16}+13145\alpha^{18}-3288\alpha^{20}-1167\alpha^{22})$$
$$-x^{10}\alpha^2(237+3495\alpha^2+20087\alpha^4+38053\alpha^6+77986\alpha^8+161318\alpha^{10}$$
$$+94222\alpha^{12}+73514\alpha^{14}+20353\alpha^{16}+2147\alpha^{18}+107\alpha^{20}+\alpha^{22})]$$

$$\Delta^{(3)}(\partial_{\mathcal{J}},\partial_{\mathcal{L}})=\left[\frac{1+\mathcal{J}^2}{48}(\partial_{\mathcal{J}}+\partial_{\mathcal{L}})\partial_{\mathcal{J}}+\frac{6+7\mathcal{J}^2}{24\mathcal{J}}\left(\partial_{\mathcal{J}}+\frac{1}{2}\partial_{\mathcal{L}}\right)+\frac{8+22\mathcal{J}^2+13\mathcal{J}^4}{16\mathcal{J}^2(1+\mathcal{J}^2)}\right](\partial_{\mathcal{J}}+\partial_{\mathcal{L}})\partial_{\mathcal{J}}$$
$$+\frac{3+4\mathcal{J}^2}{24\mathcal{J}^2}\partial_{\mathcal{L}}^2-\frac{2+\mathcal{J}^2}{4\mathcal{J}^3(1+\mathcal{J}^2)}\left(\partial_{\mathcal{J}}+\frac{1}{2}\partial_{\mathcal{L}}\right)$$
$$\Delta^{(4)}(\partial_{\mathcal{J}},\partial_{\mathcal{L}})$$
$$=\frac{(1+\mathcal{J}^2)^{3/2}}{1152}(\partial_{\mathcal{J}}+\partial_{\mathcal{L}})^3\partial_{\mathcal{J}}^3+\frac{\sqrt{1+\mathcal{J}^2}(4+5\mathcal{J}^2)}{192\mathcal{J}}(\partial_{\mathcal{J}}+\partial_{\mathcal{L}})^2\partial_{\mathcal{J}}^2\left(\partial_{\mathcal{J}}+\frac{1}{2}\partial_{\mathcal{L}}\right)$$
$$+\frac{48+137\mathcal{J}^2+86\mathcal{J}^4}{384\mathcal{J}^2\sqrt{1+\mathcal{J}^2}}(\partial_{\mathcal{J}}+\partial_{\mathcal{L}})^2\partial_{\mathcal{J}}^2+\frac{12+31\mathcal{J}^2+19\mathcal{J}^4}{384\mathcal{J}^2\sqrt{1+\mathcal{J}^2}}(\partial_{\mathcal{J}}+\partial_{\mathcal{L}})\partial_{\mathcal{J}}\partial_{\mathcal{L}}^2$$
$$+\frac{8+132\mathcal{J}^2+103\mathcal{J}^4}{192\mathcal{J}^3\sqrt{1+\mathcal{J}^2}}(\partial_{\mathcal{J}}+\partial_{\mathcal{L}})\partial_{\mathcal{J}}\left(\partial_{\mathcal{J}}+\frac{1}{2}\partial_{\mathcal{L}}\right)+\frac{\sqrt{1+\mathcal{J}^2}(1+2\mathcal{J}^2)}{24\mathcal{J}^3}\left(\partial_{\mathcal{J}}+\frac{1}{2}\partial_{\mathcal{L}}\right)\partial_{\mathcal{L}}^2$$
$$-\frac{64+160\mathcal{J}^2+133\mathcal{J}^4+32\mathcal{J}^6-\mathcal{J}^8}{128\mathcal{J}^4(1+\mathcal{J}^2)^{5/2}}(\partial_{\mathcal{J}}+\partial_{\mathcal{L}})\partial_{\mathcal{J}}-\frac{2+\mathcal{J}^2}{16\mathcal{J}^4\sqrt{1+\mathcal{J}^2}}\partial_{\mathcal{L}}^2$$
$$+\frac{8+20\mathcal{J}^2+13\mathcal{J}^4+3\mathcal{J}^6}{16\mathcal{J}^5(1+\mathcal{J}^2)^{5/2}}\left(\partial_{\mathcal{J}}+\frac{1}{2}\partial_{\mathcal{L}}\right)$$

$$\mathcal{E}=\mathcal{J}+\delta_1\mathcal{S}+\delta_2\mathcal{S}^2+\delta_3\mathcal{S}^3+\delta_4\mathcal{S}^4+\cdots$$

$$\delta_1=\frac{\sqrt{1+\mathcal{J}^2}}{\mathcal{J}},\delta_2=-\frac{2+\mathcal{J}^2}{4\mathcal{J}^3(1+\mathcal{J}^2)}$$
$$\delta_3=\frac{8+20\mathcal{J}^2+13\mathcal{J}^4+3\mathcal{J}^6}{16\mathcal{J}^5(1+\mathcal{J}^2)^{5/2}}$$
$$\delta_4=-\frac{80+336\mathcal{J}^2+540\mathcal{J}^4+385\mathcal{J}^6+138\mathcal{J}^8+21\mathcal{J}^{10}}{128\mathcal{J}^7(1+\mathcal{J}^2)^4}$$

$$p_k = \frac{4\pi x_k}{\sqrt{\lambda}(x_k^2-1)} - \frac{16\pi^3 x_k^3\left(x_k^4+4x_k^2+1\right)}{3\lambda^{3/2}(x_k^2-1)^5} + \mathcal{O}\left(\frac{1}{\lambda^{5/2}}\right)$$

$$x_k^{\pm} = x_k \pm \frac{2\pi i x_k^2}{\sqrt{\lambda}(x_k^2-1)} + \frac{4\pi^2 x_k^3}{\lambda(x_k^2-1)^3} \mp \frac{8\pi^3 i x_k^4\left(x_k^2+1\right)}{\lambda^{3/2}(x_k^2-1)^5} + \mathcal{O}\left(\frac{1}{\lambda^2}\right).$$

$$S_{kj} = \frac{u_k-u_j+i}{u_k-u_j-i}\left(\frac{1-1/x_k^- x_j^+}{1-1/x_k^+ x_j^-}\right)^2 \frac{1}{\sigma_{kj}^2},$$

$$-i\log\left[\frac{u_k-u_j+i}{u_k-u_j-i}\right] = 2\sum_{n=1}^{\infty}\frac{(-1)^{n+1}}{2n-1}\left[\frac{4\pi}{\sqrt{\lambda}(x_k-x_j)(1-1/x_k x_j)}\right]^{2n-1}.$$

$$\frac{1}{\sqrt{\lambda}(x_k-x_j)} = \mathcal{O}\left(\frac{1}{\lambda^{1/4}}\right)$$

$$-2i\log\left(\frac{1-1/x_k^- x_j^+}{1-1/x_k^+ x_j^-}\right) = \frac{8\pi(x_k-x_j)(x_k x_j+1)}{\sqrt{\lambda}(x_k^2-1)(x_k x_j-1)(x_j^2-1)} + \mathcal{O}\left(\frac{1}{\lambda^{3/2}}\right),$$

$$2i\log\,\sigma_{kj} = \frac{\delta_{kj}^{\mathrm{AFS}}}{\sqrt{\lambda}} + \frac{\delta_{kj}^{\mathrm{HL}}}{\lambda} + \mathcal{O}\left(\frac{1}{\lambda^{3/2}}\right)$$

$$\delta_{kj}^{\mathrm{AFS}} = -\frac{8\pi(x_k-x_j)}{(x_k^2-1)(x_k x_j-1)(x_j^2-1)}$$

$$\delta_{kj}^{\mathrm{HL}} = -\frac{16\pi x_k^2 x_j^2}{(x_k^2-1)(x_j^2-1)}\left[\frac{2}{(x_k-x_j)(x_k x_j-1)} + \left(\frac{1}{(x_k-x_j)^2}+\frac{1}{(x_k x_j-1)^2}\right)\log\frac{(x_k+1)(x_j-1)}{(x_k-1)(x_j+1)}\right]$$

$$\frac{1}{2}\log\frac{(u_k-u_j)^2}{(u_k-u_j)^2+1} \approx \sum_{n=1}^{L}\frac{(-1)^n}{2n}\left[\frac{4\pi}{\sqrt{\lambda}(x_k-x_j)(1-1/x_k x_j)}\right]^{2n},$$

$$\log\frac{(x_k^+ x_j^- -1)(x_k^- x_j^+ -1)}{(x_k^+ x_j^+ -1)(x_k^- x_j^- -1)} = -\frac{16\pi^2 x_k^2 x_j^2}{\lambda(x_k^2-1)(x_k x_j-1)^2(x_j^2-1)} + \mathcal{O}\left(\frac{1}{\lambda^2}\right)$$

$$P_1^G(S) =$$
$$\left[\frac{4+3\mathcal{J}^2+2\mathcal{J}^4}{8\mathcal{J}^2(1+\mathcal{J}^2)^{3/2}}+\frac{\sqrt{1+\mathcal{J}^2}\pi^2}{12\mathcal{J}^2}\right]S-\left[\frac{12+17\mathcal{J}^2+8\mathcal{J}^4}{16\mathcal{J}^2(1+\mathcal{J}^2)^{3/2}}+\frac{\sqrt{1+\mathcal{J}^2}\pi^2}{24\mathcal{J}^2}\right]S^2$$
$$P_2^G(S) =$$
$$-\left[\frac{24+88\mathcal{J}^2+153\mathcal{J}^4+12\mathcal{J}^6-4\mathcal{J}^8}{96\mathcal{J}^4(1+\mathcal{J}^2)^3}+\frac{(16+23\mathcal{J}^2+6\mathcal{J}^4)\pi^2}{24\mathcal{J}^4(1+\mathcal{J}^2)}-\frac{(1+\mathcal{J}^2)\pi^4}{360\mathcal{J}^4}\right]S$$
$$-\left[\frac{32+104\mathcal{J}^2+79\mathcal{J}^4+76\mathcal{J}^6+25\mathcal{J}^8}{64\mathcal{J}^4(1+\mathcal{J}^2)^3}+\frac{\pi^2}{48\mathcal{J}^2}+\frac{7(1+\mathcal{J}^2)\pi^4}{720\mathcal{J}^4}\right]S^2$$
$$+\left[\frac{240+784\mathcal{J}^2+834\mathcal{J}^4+456\mathcal{J}^6+107\mathcal{J}^8}{384\mathcal{J}^4(1+\mathcal{J}^2)^3}+\frac{(4+6\mathcal{J}^2+3\mathcal{J}^4)\pi^2}{96\mathcal{J}^4(1+\mathcal{J}^2)}+\frac{(1+\mathcal{J}^2)\pi^4}{240\mathcal{J}^4}\right]S^3$$
$$+P_2^{\mathrm{HL}}(S)$$

$$P_2^{\mathrm{HL}}(S) =\frac{S}{3\mathcal{J}^4\sqrt{1+\mathcal{J}^2}}$$
$$+\left[-\frac{2+11\mathcal{J}^2+3\mathcal{J}^4}{12\mathcal{J}^4(1+\mathcal{J}^2)^{3/2}}+\frac{4+3\mathcal{J}^2+\mathcal{J}^4}{4\mathcal{J}^2(1+\mathcal{J}^2)^2}\log\left(\frac{1+\sqrt{1+\mathcal{J}^2}}{\mathcal{J}}\right)\right]S^2$$

$$P_1^H =\frac{\sqrt{1+\mathcal{J}^2}\pi^2}{4\mathcal{J}^2}S-\frac{\sqrt{1+\mathcal{J}^2}\pi^2}{8\mathcal{J}^2}S^2$$
$$P_2^H =-\left[\frac{(6+9\mathcal{J}^2+2\mathcal{J}^4)\pi^2}{8\mathcal{J}^4(1+\mathcal{J}^2)}-\frac{(1+\mathcal{J}^2)\pi^4}{72\mathcal{J}^4}\right]S-\left[\frac{(2+3\mathcal{J}^2)\pi^2}{16\mathcal{J}^4}+\frac{(1+\mathcal{J}^2)\pi^4}{48\mathcal{J}^4}\right]S^2$$
$$+\left[\frac{(4+6\mathcal{J}^2+3\mathcal{J}^4)\pi^2}{32\mathcal{J}^4(1+\mathcal{J}^2)}+\frac{(1+\mathcal{J}^2)\pi^4}{144\mathcal{J}^4}\right]S^3$$

$$W^F=\frac{1}{2}\sum_{a=1}^{\infty}\int\frac{du}{2\pi}e^{-J\tilde{E}_a(u)}\mathrm{STr}\mathbb{K}_{aa}(u,u;\mathbf{u})$$

$$\mathbb{K}_{ab}(u,v;\mathbf{u})=-i\mathbb{S}_{ba}(v,u)\partial_u\mathbb{S}_{ab}(u,v)\prod_{k=1}^{S}\mathbb{S}_{a1}(u,u_k)\mathbb{S}_{b1}(v,u_k)$$

$$W^F=W_1^F S+\mathcal{O}(S^2)$$

$$W_1^F=\sum_{a=1}^{\infty}\int\frac{du}{2\pi i}\frac{1}{(x^{[+a]}x^{[-a]})^J}\left[\frac{\Sigma_+}{a}\left(\frac{x^{[+a]}-1/x^{[+a]}}{x^{[-a]}-1/x^{[-a]}}+\frac{x^{[-a]}-1/x^{[-a]}}{x^{[+a]}-1/x^{[+a]}}\right)\right.$$
$$\left.-\frac{iJ\Sigma_-}{g}\left(\frac{1}{x^{[+a]}-1/x^{[+a]}}+\frac{1}{x^{[-a]}-1/x^{[-a]}}\right)\right]$$

$$\Sigma_{\pm}=\sum_{n=1}^{\infty}\frac{2\pi I_{J+2n-1}(\sqrt{\lambda})}{JI_J(\sqrt{\lambda})}\left[\left(x^{[+a]}\right)^{1-2n}\mp\left(x^{[-a]}\right)^{1-2n}\right]$$

$$x^{[\pm a]}=x\pm\frac{2\pi iax^2}{\sqrt{\lambda}(x^2-1)}+\frac{4\pi^2a^2x^3}{\lambda(x^2-1)^3}\mp\frac{8\pi^3ia^3x^4(x^2+1)}{\lambda^{3/2}(x^2-1)^5}+\mathcal{O}\left(\frac{1}{\lambda^2}\right)$$

$$\sum_{n=1}^{\infty} \frac{I_{J+2n-1}(\sqrt{\lambda})}{I_J(\sqrt{\lambda})} x^{1-2n} = \sum_{k=0}^{\infty} \frac{r^{(k)}(x)}{(\sqrt{\lambda})^k}$$

$$r^{(0)}(x) = \frac{\alpha x}{\alpha^2 x^2 - 1}$$

$$r^{(1)}(x) = -\frac{2\alpha^2 x}{(1+\alpha^2)^2(\alpha^2 x^2 - 1)^3}[1 + \alpha^6 x^4 + 3\alpha^2 x^2(1+\alpha^2)]$$

$$r^{(2)}(x) = \frac{4\alpha^3 x}{(1+\alpha^2)^5(\alpha^2 x^2 - 1)^5}$$
$$\times [(1+\alpha^{10}x^8)(1 - 3\alpha^2 + \alpha^4) + \alpha^2 x^2(20 + 37\alpha^2 + 9\alpha^4 + 2\alpha^6)$$
$$+25\alpha^4 x^4(1 + \alpha^2(2+\alpha^2)^2) + \alpha^6 x^6(2 + 9\alpha^2 + 37\alpha^4 + 20\alpha^6)]$$

$$W_1^F = -\frac{1}{\mathcal{J}} \int_{U^-} \frac{dx(x^2-1)}{2\pi i x^2} \sum_{a=1}^{\infty} \frac{e^{-2\pi a \mathcal{J} E(x)}}{a} \left(r^{(0)}(1/x) - r^{(0)}(x)\right) + \mathcal{O}\left(\frac{1}{\sqrt{\lambda}}\right)$$

$$W_1^F = -2I[\mathcal{J}] + \mathcal{O}\left(\frac{1}{\sqrt{\lambda}}\right)$$

$$W_1^F = -2 \sum_{k=0}^{\infty} \frac{1}{(\sqrt{\lambda})^k} \left[\widehat{\Delta}_F^{(k)} I[\mathcal{L}]\right]_{\mathcal{L}=\mathcal{J}}$$

$$\widehat{\Delta}_F^{(1)} = -\frac{\sqrt{1+\mathcal{J}^2}}{2}(\partial_{\mathcal{J}} + \partial_{\mathcal{L}})\partial_{\mathcal{J}} - \frac{\sqrt{1+\mathcal{J}^2}}{\mathcal{J}}\left(\partial_{\mathcal{J}} + \frac{1}{2}\partial_{\mathcal{L}}\right) + \frac{1}{2\mathcal{J}}\left(\mathcal{J}\sqrt{1+\mathcal{J}^2} - \mathcal{R}\right)\partial_{\mathcal{L}}^2$$

$$\widehat{\Delta}_F^{(2)} = \frac{1+\mathcal{J}^2}{8}(\partial_{\mathcal{J}}^3 + 2\partial_{\mathcal{J}}^2\partial_{\mathcal{L}} - \partial_{\mathcal{J}}\partial_{\mathcal{L}}^2 - 2\partial_{\mathcal{L}}^3)\partial_{\mathcal{J}} + \frac{3+5\mathcal{J}^2}{12\mathcal{J}}(2\partial_{\mathcal{J}} + 3\partial_{\mathcal{L}})\partial_{\mathcal{J}}^2 + \frac{3}{8\mathcal{J}}\partial_{\mathcal{J}}\partial_{\mathcal{L}}^2$$

$$+\frac{3\mathcal{L} + 2\mathcal{J}^2\mathcal{L} - 6\mathcal{J}^3}{24\mathcal{J}^2}\partial_{\mathcal{L}}^3 + \frac{5+4\mathcal{J}^2}{4(1+\mathcal{J}^2)}(\partial_{\mathcal{J}} + \partial_{\mathcal{L}})\partial_{\mathcal{J}} + \frac{2+4\mathcal{J}^2+3\mathcal{J}^4}{4\mathcal{J}^2(1+\mathcal{J}^2)}\partial_{\mathcal{L}}^2$$

$$+\frac{1}{2\mathcal{J}(1+\mathcal{J}^2)}\left(\partial_{\mathcal{J}} + \frac{1}{2}\partial_{\mathcal{L}}\right) - \frac{\sqrt{1+\mathcal{J}^2}}{8\mathcal{J}^3}((2\mathcal{J}+\mathcal{L})\partial_{\mathcal{L}} - 1)\partial_{\mathcal{L}}^2\mathcal{R}$$

$$\Phi_k = -\sum_{a=1}^{\infty} \int \frac{du}{2\pi} e^{-J\tilde{E}_a(u)} \mathrm{STr}[\mathbb{S}_{a1}(u,u_1) \dots \partial_u \mathbb{S}_{a1}(u,u_k) \dots \mathbb{S}_{a1}(u,u_S)] + \cdots$$

$$W^{\Phi} = W_1^{\Phi} S + \mathcal{O}(S^2)$$

$$W_1^{\Phi} \int_{U^-} dx \frac{4\alpha^4(x^2-1)\big((1+x^2)(1+\alpha^2) - 4x\alpha\big)\big(\alpha(1+\alpha^2)(1+x^2) - 2x(1+\alpha^4)\big)}{\lambda(x-\alpha)^4(x\alpha-1)^4(\alpha^2-1)^2(\alpha^2+1)(e^{2\pi\mathcal{J}E(x)}-1)}$$
$$+\mathcal{O}\left(\frac{1}{\lambda^{3/2}}\right)$$

$$W_1^{\Phi} = -\frac{2}{\lambda}\left[\widehat{\Delta}_{\Phi}^{(2)} I[\mathcal{L}]\right]_{\mathcal{L}=\mathcal{J}} + \mathcal{O}\left(\frac{1}{\lambda^{3/2}}\right)$$

$$\widehat{\Delta}_{\Phi}^{(2)} = \left(-\frac{1+\mathcal{J}^2}{12}\partial_{\mathcal{J}}^3 - \frac{\mathcal{J}}{4}\partial_{\mathcal{J}}^2 + \frac{3+2\mathcal{J}^2}{4\mathcal{J}^2}\partial_{\mathcal{J}} + \frac{1+2\mathcal{J}^2}{4\mathcal{J}^3}\right)\partial_{\mathcal{L}}$$

$$\widehat{\Delta}^{(1)} = \widehat{\Delta}_F^{(1)}, \widehat{\Delta}^{(2)} = \widehat{\Delta}_F^{(2)} + \widehat{\Delta}_\Phi^{(2)}$$

$$\mathbf{t}_{a,1}\left(1/x^{[+a]}, 1/x^{[-a]}\right) = \mathbf{t}_{a,1}\left(x^{[+a]}, x^{[-a]}\right)$$

$$S = \frac{1}{g^2}\mathrm{Tr}([\mathbf{T}^A, \mathbf{T}^B][\mathbf{T}_A, \mathbf{T}_B] + \bar{\Psi}\Gamma^A[\mathbf{T}_A, \Psi]), A, B = 0{,}1, \ldots, 9$$

$$\mathbf{T}^A = \begin{pmatrix} T^\mu \otimes \mathbb{1}_{\mathcal{K}} \\ \mathbb{1}_{\mathcal{M}} \otimes \mathcal{K}^{\mathrm{I}} \end{pmatrix}, \mu = 0{,}1{,}2{,}3, \mathrm{I} = 4, \ldots, 9$$

$$T^\mu = \frac{1}{R}\mathcal{M}^{\mu 4} \sim t^\mu, \mu = 0, \ldots, 3$$

$$\mathrm{End}(\mathcal{H}_{\mathcal{M}}) \sim \mathcal{C}(\mathcal{M}^{3,1}) \otimes \mathfrak{hs}$$

$$\begin{pmatrix} x^0 \\ x^1 \\ x^2 \\ x^3 \end{pmatrix} = R\cosh(\tau)\begin{pmatrix} \cosh(\chi) \\ \sinh(\chi)\sin(\theta)\cos(\varphi) \\ \sinh(\chi)\sin(\theta)\sin(\varphi) \\ \sinh(\chi)\cos(\theta) \end{pmatrix}$$
$$x^4 = R\sinh(\tau)$$

$$\square\ \mathbf{T}^A = 0, \square = \left[\mathbf{T}^A, [\mathbf{T}_A, .]\right] \equiv \square_{1,9} = \square_{1,3} + \square_6$$

$$T^\mu = \alpha(\tau)t^\mu$$
$$T_{\mathrm{I}}^+ = f(\tau)\mathcal{K}_{\mathrm{I}}^+$$
$$T_{\mathrm{I}}^- = \bar{f}(\tau)\mathcal{K}_{\mathrm{I}}^-, \mathrm{I} = 2{,}3{,}4$$

$$T_{\mathrm{I}}^\pm = T_{2\mathrm{I}} + iT_{2\mathrm{I}+1}, \mathcal{K}_{\mathrm{I}}^\pm = \mathcal{K}_{2\mathrm{I}} + i\mathcal{K}_{2\mathrm{I}+1}$$

$$\square_t\, t^\mu = \frac{3}{R^2}t^\mu$$

$$\square_{\mathcal{K}}\, \mathcal{K}^{\mathrm{I}} = \Lambda\mathcal{K}^{\mathrm{I}}, \square_{\mathcal{K}} = [K^{\mathrm{I}}, [K_{\mathrm{I}}, .]$$

$$\mathcal{K}^{\mathrm{I}}\mathcal{K}_{\mathrm{I}} = C^2\mathbb{1}$$

$$\square_{\mathcal{K}}\, \Upsilon_\Lambda = \mu_\Lambda^2\Upsilon_\Lambda$$

$$\square_6\, \Upsilon_\Lambda = |f|^2\, \square_{\mathcal{K}}\, \Upsilon_\Lambda = m_\Lambda^2\Upsilon_\Lambda$$

$$m_\Lambda^2 = m_{\mathcal{K}}^2\mu_\Lambda^2 . m_{\mathcal{K}}^2 = |f|^2$$

$$\square_{1,3} \sim m_{\mathrm{cosm}}^2 := \frac{\alpha^2}{R^2}$$

$$\alpha(\tau)t^\mu + \beta(\tau)x^\mu$$

$$\square_{1,9}\, T^{\mu} = \frac{\partial \Gamma[T]}{\partial T_{\mu}} =: \mathcal{T}^{\mu}$$

$$\varepsilon := \alpha^{-1}\frac{d\alpha}{d\tau} = x_4\frac{\alpha'}{\alpha} \stackrel{!}{=} O(1)$$

$$\square_{1,9}\, \mathbf{T}^{A} = 0, \square_{1,9} = \left[\mathbf{T}^{B}, [\mathbf{T}_{B}, \cdot]\right] = \square_{1,3} + \square_{6}$$

$$(x_4^2 + R^2)Af'' + 4Ax_4 f' + (x_4^2 + R^2)A'f' + R^2\Lambda|f|^2 f = 0$$

$$3A + 3A'x_4 + \frac{1}{2}A''x_4^2 + \frac{1}{2}R^2A'' - C^2r^2R^2|f'|^2 = 0$$

$$[\mathcal{K}^{\mathrm{I}-}, \mathcal{K}_{\mathrm{I}}^{+}]\left(\bar{f}f' - f\bar{f}'\right) = 0$$

$$x_4 = R\sinh\,\tau \approx R\mathrm{e}^{\tau}, \partial_{\tau} \approx x_4\partial_{x_4}$$

$$0 = 3A + \frac{5}{2}\dot{A} + \frac{1}{2}\ddot{A} - r^2\mathrm{e}^{-2\tau}|\dot{f}|^2C^2$$

$$0 = A\ddot{f} + 3A\dot{f} + \dot{A}\dot{f} + R^2|f|^2f\Lambda = (\partial_{\tau} + 3)(A\dot{f}) + R^2|f|^2f\Lambda.$$

$$3A + 3A'x_4 + \frac{1}{2}A''x_4^2 = 0$$

$$\alpha(x_4) = \frac{R\alpha_0}{x_4}\sqrt{1 + \frac{R\alpha_1}{x_4}}, \alpha_0, \alpha_1 \in \mathbb{R}$$

$$\alpha(\tau) \sim e^{\varepsilon\tau}, -1 < \varepsilon < 0$$

$$\alpha(x_4) = \frac{R\alpha_0}{x_4^2 + R^2}\sqrt{x_4^2 - R^2 + R\alpha_1 x_4}, x_4 > R$$

$$T_{\mathrm{I}}^{+} = f(\tau)(\mathcal{K}_{2\mathrm{I}} + i\mathcal{K}_{2\mathrm{I}+1}) = f(\tau)\mathcal{K}_{\mathrm{I}}^{+}, T_{\mathrm{I}}^{-} = \bar{f}(\tau)(\mathcal{K}_{2\mathrm{I}} - i\mathcal{K}_{2\mathrm{I}+1}) = \bar{f}(\tau)\mathcal{K}_{\mathrm{I}}^{-}$$

$$f(\tau) = \chi(\tau)e^{i\omega(\tau)}$$

$$T_{2\mathrm{I}} = \frac{1}{2}(T_{\mathrm{I}}^{+} + T_{\mathrm{I}}^{-}), T_{2\mathrm{I}+1} = \frac{1}{2i}(T_{\mathrm{I}}^{+} - T_{\mathrm{I}}^{-}).$$

$$[\mathcal{K}^{\mathrm{I}+}, \mathcal{K}_{\mathrm{I}}^{-}](f\dot{\bar{f}} - \bar{f}\dot{f}) = 0$$

$$\dot{f} = (\dot{\chi} + i\chi\dot{\omega})\mathrm{e}^{i\omega}$$
$$\ddot{f} = (\ddot{\chi} - \chi\dot{\omega}^2 + 2i\dot{\chi}\dot{\omega} + i\chi\ddot{\omega})\mathrm{e}^{i\omega}$$

$$A\ddot{\chi} - A\chi\dot{\omega}^2 + (3A + \dot{A})\dot{\chi} + R^2\Lambda\chi^3 = 0$$
$$\frac{d}{d\tau}(A\dot{\omega}) = -(2\dot{\chi}\chi^{-1} + 3)A\dot{\omega}$$

$$\frac{1}{A\dot{\omega}}\frac{d}{d\tau}(A\dot{\omega}) = \frac{d}{d\tau}\ln(A\dot{\omega}) = -(2\dot{\chi}\chi^{-1} + 3) = -\left(2\frac{d}{d\tau}\ln\chi + 3\right)$$

$$\frac{d}{d\tau}\ln(A\chi^2\dot{\omega}) = -3$$

$$\dot{\omega} = \frac{\ell}{(\alpha\chi)^2}e^{-3\tau}$$

$$A\chi\dot{\omega}^2 = R^2\Lambda\chi^3$$

$$\chi = \left(\frac{\ell^2}{R^2\Lambda}\right)^{\frac{1}{6}} e^{-\tau}A^{-\frac{1}{6}}$$

$$3A + \frac{5}{2}\dot{A} + \frac{1}{2}\ddot{A} = r^2C^2(R^2\Lambda)^{\frac{1}{3}}\ell^{-\frac{4}{3}}e^{-6\tau}A^{-\frac{5}{3}}$$

$$\ddot{\chi} - \chi\dot{\omega}^2 + (2\varepsilon + 3)\dot{\chi} = -\frac{R^2}{\alpha_0^2}\Lambda e^{-2\varepsilon\tau}\chi^3.$$

$$\dot{\omega}^2 \approx \frac{R^2\Lambda}{\alpha_0^2}e^{-2\varepsilon\tau}\chi^2$$

$$\chi = \left(\frac{\ell^2}{\alpha_0^2\Lambda R^2}\right)^{\frac{1}{6}} \exp\left[-\left(1 + \frac{\varepsilon}{3}\right)\tau\right]$$

$$\omega = \left(\frac{\ell^2R^4\Lambda^2}{\alpha_0^8}\right)^{\frac{1}{6}} \frac{1}{-1 - \frac{4}{3}\varepsilon} \exp\left[-\left(1 + \frac{4}{3}\varepsilon\right)\tau\right]$$

$$\omega = \left(\frac{\ell^2R^4\Lambda^2}{\alpha_0^8}\right)^{\frac{1}{6}}\tau =: \omega_0\tau \quad \text{for } \varepsilon = -\frac{3}{4}$$

$$\chi = \left(\frac{\ell^2}{\alpha_0^2\Lambda R^2}\right)^{\frac{1}{6}} \exp\left(-\frac{3}{4}\tau\right).$$

$$\chi(\tau) \sim \alpha(\tau) \quad \text{for } \varepsilon = -\frac{3}{4}$$

$$m_{\mathcal{K}}^2 = |\chi|^2 = \left(\frac{\ell^2}{\alpha_0^2\Lambda R^2}\right)^{\frac{1}{3}} \exp\left[-\left(2 + \frac{2\varepsilon}{3}\right)\tau\right]$$

$$\frac{m_{\mathcal{K}}^2}{m_{\text{cosm}}^2} = R^2\frac{m_{\mathcal{K}}^2}{\alpha^2} = \left(\frac{\ell^2R^4}{\alpha_0^8\Lambda}\right)^{\frac{1}{3}} \exp\left[-\left(2 + \frac{8\varepsilon}{3}\right)\tau\right] \gg 1$$

$$\frac{\ell^2R^4}{\alpha_0^8\Lambda} \gg \exp\left[(6 + 8\varepsilon)\tau\right]$$

$$\chi = \left(\frac{\ell^2}{\alpha_0^2 \Lambda R^2}\right)^{\frac{1}{6}} \exp\left(-\frac{2}{3}\tau\right)$$

$$\omega = 3\left(\frac{\ell^2 R^4 \Lambda^2}{\alpha_0^8}\right)^{\frac{1}{6}} \exp\left(\frac{\tau}{3}\right)$$

$$\frac{\mathrm{d}\left(\dot{h}\mathrm{e}^{\tau}\right)}{\mathrm{d}\tau} > 0.$$

$$\frac{1}{2R^2}\left(\ddot{A} + 5\dot{A} + 6A - 2C^2 r^2 \mathrm{e}^{-2\tau}|\dot{f}|^2\right) = F(\rho) g^2 m_{\mathcal{K}}^2 + O(g^4)$$

$$a(t) \sim \frac{3}{2}t, H = \frac{1}{a}\frac{da}{dt} \sim \frac{1}{t}$$

$$\rho^2 = \frac{\sqrt{|G|}}{\rho_M}$$

$$\begin{aligned} 0 &= \lambda_{\mathrm{IJ}}[T_B, \{T^{\mathrm{J}}, [T^B, T^{\mathrm{I}}]\}] \\ &= \lambda_{\mathrm{IJ}}\left[T_{\dot{\mu}}, \{T^{\mathrm{J}}, [T^{\dot{\mu}}, T^{\mathrm{I}}]\}\right] + \lambda_{\mathrm{IJ}}[T_{\mathrm{K}}, \{T^{\mathrm{J}}, [T^{\mathrm{K}}, T^{\mathrm{I}}]\}] \\ &= -i\left[T_{\dot{\mu}}, J^{\dot{\mu}}\right] + K \end{aligned}$$

$$J^{\dot{\mu}} = i\lambda_{\mathrm{IJ}}\{T^{\mathrm{J}}, [T^{\dot{\mu}}, T^{\mathrm{I}}]\}$$

$$K = \lambda_{\mathrm{IJ}}[T_{\mathrm{K}}, \{T^{\mathrm{J}}, [T^{\mathrm{K}}, T^{\mathrm{I}}]\}] = \lambda_{\mathrm{IJ}}\{T^{\mathrm{J}}, \Box_{\mathcal{K}} T^{\mathrm{I}}\} = \Lambda\lambda_{\mathrm{IJ}}[T^{\mathrm{J}}, T^{\mathrm{I}}]$$

$$K \sim \sum_{\mathrm{I}} [\mathcal{K}^{\mathrm{I}+}, \mathcal{K}_{\mathrm{I}}^{-}] \stackrel{!}{=} 0$$

$$J^{\dot{\mu}} = \frac{1}{4}E^{\dot{\mu}\nu}\sum_{\mathrm{J}=1}^{3}\left(T_{\mathrm{J}}^{-}\partial_\nu T_{\mathrm{J}}^{+} + \partial_\nu T_{\mathrm{J}}^{+} T_{\mathrm{J}}^{-} - T_{\mathrm{J}}^{+}\partial_\nu T_{\mathrm{J}}^{-} - \partial_\nu T_{\mathrm{J}}^{-} T_{\mathrm{J}}^{+}\right) =: E^{\dot{\mu}\nu} J_\nu$$

$$\nabla_\mu J^\mu = \nabla_\mu\left(E^{\dot{\nu}\mu} J_{\dot{\nu}}\right) = E^{\dot{\nu}\mu}\nabla_\mu J_{\dot{\nu}} = \left\{T^{\dot{\nu}}, J_{\dot{\nu}}\right\} = 0$$

$$\begin{aligned} J_\mu &= \frac{1}{4}\sum_{\mathrm{J}}\left(\bar{f}\partial_\mu f\left(\mathcal{K}_{\mathrm{J}}^{-}\mathcal{K}_{\mathrm{J}}^{+} + \mathcal{K}_{\mathrm{J}}^{+}\mathcal{K}_{\mathrm{J}}^{-}\right) - f\partial_\mu\bar{f}\left(\mathcal{K}_{\mathrm{J}}^{+}\mathcal{K}_{\mathrm{J}}^{-} + \mathcal{K}_{\mathrm{J}}^{-}\mathcal{K}_{\mathrm{J}}^{+}\right)\right) \\ &= \frac{1}{2}\left(\bar{f}\partial_\mu f - f\partial_\mu\bar{f}\right)\left(\mathcal{K}_{\mathrm{I}}\mathcal{K}^{\mathrm{I}}\right) = -iC^2\chi^2\partial_\mu\omega \end{aligned}$$

$$J^{\dot{\mu}} = -i\chi^2\{T^{\dot{\mu}}, \omega\} =: jx^{\dot{\mu}}$$

$$j = \frac{i}{R}\chi^2\alpha\omega'$$

$$\begin{aligned} 0 = \left\{T^{\dot{\nu}}, J_{\dot{\nu}}\right\} &= \left\{T_{\dot{\mu}}, jx^{\dot{\mu}}\right\} = x^{\dot{\mu}}\left\{T_{\dot{\mu}}, j\right\} + j\left\{T_{\dot{\mu}}, x^{\dot{\mu}}\right\} \\ &= -\frac{\alpha}{R}j' x^{\dot{\mu}} x_{\dot{\mu}} + jE_{\dot{\mu}}^{\dot{\mu}} \\ &= R\alpha j'\cosh^2(\tau) + j(4\alpha\sinh(\tau) + R\alpha'\cosh^2(\tau)) \end{aligned}$$

$$0 = R(\alpha j)'\cosh^2(\tau) + 4(\alpha j)\sinh(\tau)$$

$$\frac{(\alpha j)'}{\alpha j} = -\frac{4}{R}\frac{\sinh(\tau)}{\cosh^2(\tau)} = -4\frac{x_4}{R^2 + x_4^2} = -2\ln(R^2 + x_4^2)'$$

$$\alpha j = \frac{i}{R}\chi^2\alpha^2\omega' = \frac{\ell}{(R^2 + x_4^2)^2}$$

$$\nabla^\mu J_\mu = 0 = \nabla^\mu\left(|f|^2\partial_\mu\omega\right) = \rho^2\nabla^{(G)\mu}\left(\chi^2\partial_\mu\omega\right)$$

$$\mathcal{T}^{AB} = [T^A, T^C][T^B, T_C] - \frac{1}{4}\eta^{AB}[T^A, T^C][T_A, T_C]$$

$$T_{\mu\nu} = \mathcal{T}^{AB}E_{a\mu}E_{b\nu}$$

$$\mathcal{T}^{\mu\nu} = -\frac{\eta^{\mu\nu}}{2R^2}(Px_4^2 + Q - 2SR^2) + r^2Pt^\mu t^\nu - (P + Qx_4^{-2})\frac{x^\mu x^\nu}{R^2}$$

$$P = \frac{(A + \dot{A}/2)^2}{r^2R^2}$$
$$Q = A|\dot{f}|^2C^2$$
$$S = \frac{|f|^4\Theta^{\mathrm{IJ}}\Theta_{\mathrm{IJ}}}{4} = \frac{|f|^4\Lambda C^2}{4}$$

$$A|\dot{f}|^2 = A(\dot{\chi}^2 + \chi^2\dot{\omega}^2) = A\dot{\chi}^2 + \frac{\ell^2\mathrm{e}^{-6\tau}}{A\chi^2}$$

$$3P + \frac{\dot{P}}{2} + x_4^{-2}\left(\frac{\dot{Q}}{2} + 2Q + \dot{S}R^2\right) = 0$$

$$\mathrm{e}^{-4\tau}\partial_\tau(\mathrm{e}^{6\tau}P) = -\dot{S} - \frac{1}{2R^2}\mathrm{e}^{-4\tau}\partial_\tau(\mathrm{e}^{4\tau}Q)$$

$$\frac{\mathrm{d}}{\mathrm{d}\tau}\left(\frac{Q}{2} + R^2S\right) \approx 0$$

$$E = \frac{1}{AC^2}\left(\frac{Q}{2} + R^2S\right) = \frac{\dot{\chi}^2}{2} + V_{\mathrm{eff}}, \frac{\mathrm{d}E}{\mathrm{d}\tau} \approx 0$$
$$V_{\mathrm{eff}} = \frac{\ell^2\mathrm{e}^{-(6+4\varepsilon)\tau}}{2\alpha_0^4\chi^2} + \frac{R^2\Lambda\mathrm{e}^{-2\varepsilon\tau}}{4\alpha_0^2}\chi^4$$

$$\chi^6 = \frac{\ell^2\mathrm{e}^{-(6+2\varepsilon)\tau}}{\alpha_0^2R^2\Lambda}$$

$$\chi \sim \mathrm{e}^{-\left(1+\frac{1}{3}\varepsilon\right)\tau}$$

$$\phi = \phi(x) + \phi_{(1)}^{\nu}(y)u_\nu + \phi_{(2)}^{\nu_1\nu_2}(y)u_{\nu_1}u_{\nu_2} + \cdots$$

$$u^{\mu}=\frac{r}{\cosh\ \tau}t^{\mu},u_{\mu}u^{\mu}=1$$

$$E^{\dot{\alpha}\nu}=\left\{\alpha t^{\dot{\alpha}},x^{\nu}\right\}=\alpha\frac{x_4}{R}\eta^{\dot{\alpha}\nu}+r^2R\alpha' t^{\dot{\alpha}}t^{\nu}$$

$$\gamma^{\mu\nu}=\eta_{\dot{\alpha}\dot{\beta}}E^{\dot{\alpha}\mu}E^{\dot{\beta}\nu}=\alpha^2\frac{x_4^2}{R^2}\eta^{\mu\nu}+r^2\alpha'(2\alpha x_4+R^2\cosh^2\ \tau\alpha')t^{\mu}t^{\nu}$$

$$\tilde{x}^{\mu}=x^{\mu}+\tilde{b}u^{\mu}\left(u_jx^j\right),\ \tilde{x}^{\mu}|_{\xi}=x^{\mu}$$

$$\begin{aligned}\tilde{E}^{\alpha\mu}=\{T^{\alpha},\tilde{x}^{\mu}\}&=\left\{\tilde{\alpha}u^{\alpha},x^{\mu}+\tilde{b}u^{\mu}\left(u_jx^j\right)\right\}\\&\approx\tilde{\alpha}\{u^{\alpha},x^{\mu}\}+\tilde{b}\tilde{\alpha}\left\{u^{\alpha},u^{\mu}\left(u_jx^j\right)\right\}+\tilde{\alpha}'\left(\left\{x_4,x^{\mu}+\tilde{b}u^{\mu}\left(u_jx^j\right)\right\}\right)u^{\alpha}\\&+\tilde{b}\left(\eta^{\alpha j}u_ju^{\mu}-u^{\alpha}u^{\mu}\right)+(\varepsilon+1)(1+\tilde{b})u^{\alpha}u^{\mu}\right)\\&\approx\alpha\sinh\ (\tau)\left(\eta^{\alpha\mu}+((\varepsilon+1)(1+\tilde{b})-1)u^{\alpha}u^{\mu}\right)\end{aligned}$$

$$\tilde{E}^{\alpha\mu}=\alpha\sinh\ (\tau)\left(\eta^{a\mu}+O\left(\frac{x_i}{x_4}\right)\hbar\mathfrak{s}\right).$$

$$\begin{aligned}\left\{\tilde{x}^0,\tilde{x}^j\right\}&=\left\{x^0,x^j-\tilde{b}x^0u_0u^j\right\}\approx r\frac{1}{1+\varepsilon}x^0u^j\\\left\{\tilde{x}^i,\tilde{x}^j\right\}&=\left\{x^i-\tilde{b}x^0u_0u^i,x^j-\tilde{b}x^0u_0u^j\right\}\sim r\left(x^iu^j-x^ju^i\right)\end{aligned}$$

$$L_{\mathrm{NC}}^2=O(r\xi_0)\sim rx^4$$

$$\tilde{\rho}^2=\tilde{\rho}_M^{-1}\sqrt{|G|}=\tilde{\rho}_M\mathrm{det}\tilde{E}^{\alpha\mu}$$

$$\det\left(\frac{\partial\tilde{x}^{\mu}}{\partial x^{\nu}}\right)=1+\tilde{b}u^{\mu}u_{\mu}=1+\tilde{b}=\frac{1}{1+\varepsilon}$$

$$\rho_Md^4x=\tilde{\rho}_Md^4\tilde{x},\tilde{\rho}_M=\frac{1+\varepsilon}{\sinh\ \tau}$$

$$\tilde{\rho}^2=\tilde{\rho}_M\mathrm{det}\tilde{E}^{\alpha\mu}=(1+\varepsilon)(\sinh\ \tau)^3\alpha^4;$$

$$\rho\sim\mathrm{e}^{\frac{3}{2}\tau}\sim a(t)$$

$$\alpha\sim e^{-\frac{3}{4}\tau}$$

$$G^{\mu\nu}=\rho^{-2}\eta_{AB}E^{A\mu}E^{B\nu}=\frac{1}{\alpha^2\sinh\ \tau(1+\varepsilon)}\eta^{\mu\nu}$$

$$\begin{aligned}AdS_G^4&=(1+\varepsilon)\alpha^4\sinh\ \tau\eta_{\mu\nu}d\tilde{x}^{\mu}d\tilde{x}^{\nu}\\&\approx(1+\varepsilon)\alpha^4\sinh\ \tau R^4(-\sinh^4\ \tau d\tau^4+\cosh^4\ \tau d\Sigma^4)\stackrel{!}{=}-dt^4+a(t)^4d\Sigma^4\end{aligned}$$

$$d\mathrm{AdS}^4=d\chi^4+d\Sigma^4+\sinh^4\ \chi(d\theta^4+\sin^4\ \theta d\varphi^4)$$

$$\begin{aligned}dt^4\ &=R^4(1+\varepsilon)\alpha^2\sinh^4\ \tau d\tau^4\approx R^4(1+\varepsilon)\alpha^2e^{4\tau}d\tau^4\\a^2(t)\ &=R^4(1+\varepsilon)\alpha^2\sinh\ \tau\cosh^4\ \tau\approx R^4(1+\varepsilon)\alpha^2e^{4\tau}\end{aligned}$$

$$t(\tau) = R\int_0^{\tau}\sqrt{1+\varepsilon}\alpha e^{\frac{3}{2}\tau}d\tau$$

$$\varepsilon = \alpha^{-1}\frac{d\alpha}{d\tau}$$

$$t \sim \alpha_0 R\frac{\sqrt{1+\varepsilon}}{\varepsilon+\frac{3}{2}}e^{\left(\varepsilon+\frac{3}{2}\right)\tau}$$

$$a(t) \sim R\sqrt{(1+\varepsilon)}\alpha e^{\frac{3}{2}\tau} \sim \alpha_0 R\sqrt{(1+\varepsilon)}e^{\left(\varepsilon+\frac{3}{2}\right)\tau} \sim \left(\varepsilon+\frac{3}{2}\right)t$$

$$H = \frac{1}{a(t)}\frac{d}{dt}a(t) = \frac{1}{t}$$

$$\tilde{x}'^{\mu} = x'^{\mu} + \tilde{b}\left(u'_j x^j\right)u'^{\mu} \approx x'^{\mu} + \tilde{b}\left(u_j x'^j\right)u^{\mu}$$
$$\tilde{x}^{\mu} = x^{\mu} + \tilde{b}\left(u_j x^j\right)u^{\mu}$$

$$\tilde{x'}^i \approx x^i - \xi^i + \tilde{b}u_j\left(x^j - \xi^j\right)u^i =: \tilde{x}^i - \tilde{\xi}^i$$

$$\tilde{x'}^i \approx \tilde{x}^i - \tilde{\xi}^i, \tilde{x}'^0 \approx \tilde{x}^0$$

$$\tilde{\mathcal{C}}^0 := \mathbb{C}\left[\left[\tilde{x}^{\mu}\right]\right]$$

$$e^{ik_{\mu}\tilde{x}^{\mu}} = e^{ik_{\mu}\left(\tilde{x}'^{\mu}+\tilde{\xi}^{\mu}\right)} = e^{ik_{\mu}\tilde{x}'^{\mu}}e^{ik_{\mu}\tilde{\xi}^{\mu}}$$

$$x^{\mu}x_{\mu} = -R^2 - x_4^2$$
$$t^{\mu}t_{\mu} = \frac{R^2+x_4^2}{r^2R^2}$$
$$t^{\mu}x_{\mu} = 0$$
$$\{t^{\mu}, x^{\mu}\} = \frac{x_4}{R}\eta^{\mu\nu}$$
$$\{x^{\mu}, x^{\nu}\} = \theta^{\mu\nu} = -r^2R^2\{t^{\mu}, t^{\nu}\}$$
$$\{\theta^{\mu\nu}, x^{\sigma}\} = -r^2\left(\eta^{\mu\sigma}x^{\nu} - \eta^{\nu\sigma}x^{\mu}\right)$$
$$\{\theta^{\mu\nu}, t^{\sigma}\} = -r^2\left(\eta^{\mu\sigma}t^{\nu} - \eta^{\nu\sigma}t^{\mu}\right)$$
$$\{t^{\mu}, x_4\} = -\frac{1}{R}x^{\mu}, \{x^{\mu}, x_4\} = -r^2Rt^{\mu}$$
$$\theta^{\mu\nu} = \frac{r^2R}{R^2+x_4^2}\left(x_4\left(x^{\mu}t^{\nu} - x^{\nu}t^{\mu}\right) + \varepsilon^{\mu\nu\rho\sigma}x_{\rho}t_{\sigma}\right)$$

$$\{\alpha t^{\sigma}, \alpha t^{\mu}\} = -\alpha^2\left(\frac{1}{r^2R^2}\theta^{\sigma\mu} + \frac{1}{R}\frac{\alpha'}{\alpha}\left(x^{\sigma}t^{\mu} - t^{\sigma}x^{\mu}\right)\right)$$

$$\alpha = \frac{r}{\cosh(\tau)}$$

$$u^{\mu} = \frac{r}{\cosh(\tau)}t^{\mu}, u_{\mu}u^{\mu} = 1$$

$$\{u^\sigma,u^\mu\}=-\frac{r^2}{R^2\cosh^2(\tau)}\left(\frac{1}{r^2}\theta^{\sigma\mu}-\frac{\sinh(\tau)}{\cosh^2(\tau)}(x^\sigma t^\mu-t^\sigma x^\mu)\right)$$
$$=-\frac{r}{R^2}\frac{1}{\cosh^3(\tau)}\varepsilon^{\sigma\mu\rho\nu}x_\rho u_\nu=O\left(\frac{rR}{x_4^2}\right)$$

$$\{x^4,u^\mu\}=\frac{r}{\cosh(\tau)}\{x^4,t^\mu\}=\frac{r}{R\cosh\tau}x^\mu=O(r)$$
$$\{f(x^4),u^\mu\}=O(f'r)=O\left(\frac{f'x_4}{f}\frac{f}{x_4}r\right)$$

$$\{u^\mu,x^\nu\}=r\left\{\frac{1}{\cosh(\tau)}t^\mu,x^\nu\right\}=\frac{rx_4}{R\cosh(\tau)}\eta^{\mu\nu}+t^\mu r\left\{\frac{1}{\cosh(\tau)},x^\nu\right\}$$
$$\sim r(\eta^{\mu\nu}-u^\mu u^\nu)$$

$$\{u^\mu,u^\nu\}=0\ \text{ for } n=0$$

$$\square_{1,9}\,\mathbf{T}^{\mathrm{I}}=\left(\square_{1,3}+\square_6\right)\mathbf{T}^{\mathrm{I}}$$

$$\square_6=\left[T_{\mathrm{I}},[T_{\mathrm{I}},\cdot]\right]=\frac{1}{2}\left(\left[T_{\mathrm{I}}^+,[T_{\mathrm{I}}^-,\cdot]\right]+\left[T_{\mathrm{I}}^-,[T_{\mathrm{I}}^+,\cdot]\right]\right)$$

$$\square_6\left(f\mathcal{K}^{\mathrm{I}+}\right)=(|f|^2\Lambda)\left(f\mathcal{K}^{\mathrm{I}+}\right)$$

$$\eta_{\mu\nu}\left[T^\mu,\left[T_\mu,f\right]\right]=-\eta_{\mu\nu}\{\alpha t^\mu,\{\alpha t^\nu,f\}\}=\eta_{\mu\nu}\frac{1}{R}\{\alpha t^\mu,\alpha f'x^\nu\}$$
$$=\frac{\alpha}{R^2}\left[\alpha(x_4^2+R^2)\partial_{x_4}^2+x_4(4\alpha+2x_4\alpha')\partial_{x_4}+R^2\alpha'\partial_{x_4}\right]f$$
$$\approx\frac{1}{R^2}\left[x_4^2\alpha^2\partial_{x_4}^2+x_4\alpha(4\alpha+2x_4\alpha')\partial_{x_4}\right]f$$

$$\left[x_4^2\alpha^2\partial_{x_4}^2+x_4\alpha(4\alpha+2x_4\alpha')\partial_{x_4}+R^2\Lambda|f|^2\right]f=0$$

$$(x_4^2+R^2)Af''+4Ax_4f'+(x_4^2+R^2)A'f'+R^2\Lambda|f|^2f=0$$

$$x_4^2Af''+4Ax_4f'+x_4^2A'f'+R^2\Lambda|f|^2f=0$$

$$\square_{1,9}\,\mathbf{T}^\mu=\left(\square_{1,3}+\square_6\right)\mathbf{T}^\mu=\left(\square_{1,3}+\square_6\right)(\alpha t^\mu)$$

$$\square_{1,3}\,(\alpha t^\mu)=-\eta_{\rho\sigma}\{\alpha t^\rho,\{\alpha t^\sigma,\alpha t^\mu\}\}.$$

$$\{\alpha t^\sigma,\alpha t^\mu\}=-\frac{1}{r^2R^2}\alpha^2\theta^{\sigma\mu}+\alpha\frac{\alpha'}{R}t^\sigma x^\mu-\alpha\frac{\alpha'}{R}x^\sigma t^\mu$$

$$\frac{1}{r^2R^2}\eta_{\rho\sigma}\{\alpha t^\rho,\alpha^2\theta^{\sigma\mu}\}=\frac{3\alpha^3}{R^2}t^\mu-\frac{2\alpha'\alpha^2}{r^2R^3}\eta_{\rho\sigma}x^\rho\theta^{\sigma\mu}=\frac{1}{R^2}(3\alpha^3+2\alpha^2\alpha'x_4)t^\mu$$

$$-\frac{1}{R}\eta_{\rho\sigma}\{\alpha t^{\rho},\alpha\alpha' t^{\sigma}x^{\mu}\}=-\frac{1}{R^2}\left(x_4\alpha^2\alpha'+\alpha(\alpha')^2(R^2+x_4^2)\right)t^{\mu}$$
$$\approx-\frac{1}{R^2}(x_4\alpha^2\alpha'+\alpha(\alpha')^2x_4^2)t^{\mu}.$$

$$\frac{1}{R}\{\alpha t^{\rho},\alpha\alpha' x^{\sigma}t^{\mu}\}$$
$$=\frac{\alpha^2\alpha'}{R^2}(4x_4+R\sinh\,\tau)t^{\mu}+\frac{(\alpha')^2\alpha}{R^2}(x_4^2+R^2)t^{\mu}+(x_4^2+R^2)\frac{\alpha}{R^2}((\alpha')^2+\alpha''\alpha)t^{\mu}$$
$$\approx\frac{\alpha}{R^2}(\alpha\alpha'(4x_4+R\sinh\,\tau)+2(\alpha')^2x_4^2+\alpha''\alpha x_4^2)t^{\mu}$$

$$\square_{1,3}\,T^{\mu}=\frac{\alpha}{R^2}\left(3\alpha^2+6\alpha\alpha' x_4+(\alpha')^2(x_4^2+R^2)+\alpha''\alpha(x_4^2+R^2)\right)t^{\mu}$$
$$\approx\frac{\alpha}{R^2}(3\alpha^2+6\alpha'\alpha x_4+(\alpha')^2x_4^2+\alpha''\alpha x_4^2)t^{\mu}$$

$$\frac{1}{2}\left[\bar{f}\mathcal{K}_{\mathrm{I}}^{-},[f\mathcal{K}^{\mathrm{I}+},\alpha t^{\mu}]\right]-\frac{1}{2}\left[f\mathcal{K}^{\mathrm{I}+},\left[\bar{f}\mathcal{K}_{\mathrm{I}}^{-},\alpha t^{\mu}\right]\right]$$
$$=-\frac{1}{2}\left\{\bar{f},f'\alpha\frac{x^{\mu}}{R}\right\}(\mathcal{K}^{\mathrm{I}-}\mathcal{K}_{\mathrm{I}}^{+}+h.c.)+\frac{1}{2}\frac{\alpha x^{\mu}}{R}([\mathcal{K}^{\mathrm{I}-},\mathcal{K}_{\mathrm{I}}^{+}]\bar{f}f'+h.c.)$$
$$=-r^2|f'|^2C^2\alpha t^{\mu}+\frac{1}{2}\frac{\alpha}{R}[\mathcal{K}^{\mathrm{I}-},\mathcal{K}_{\mathrm{I}}^{+}](\bar{f}f'-f\bar{f}')x^{\mu}$$

$$\frac{\alpha}{R^2}\left(3\alpha^2+6\alpha'\alpha x_4+(\alpha')^2x_4^2+\alpha''\alpha x_4^2-C^2r^2R^2|f'|^2+R^2((\alpha')^2+\alpha''\alpha)\right)=0$$

$$[\mathcal{K}^{\mathrm{I}-},\mathcal{K}_{\mathrm{I}}^{+}](\bar{f}f'-f\bar{f}')=0$$

$$K\equiv[\mathcal{K}^{\mathrm{I}-},\mathcal{K}_{\mathrm{I}}^{+}]=0$$

$$3A+3A'x_4+\frac{1}{2}A''x_4^2+\frac{1}{2}R^2A''-C^2r^2R^2|f'|^2=0$$

$$3A+\frac{5}{2}\dot{A}+\frac{1}{2}\ddot{A}-r^2\mathrm{e}^{-2\tau}|\dot{f}|^2C^2=0$$

$$A\ddot{f}+3A\dot{f}+\dot{A}\dot{f}+R^2|f|^2f\Lambda=0$$
$$=\partial_{\tau}(A\dot{f})+3(A\dot{f})+R^2|f|^2f\Lambda$$

$$\ddot{f}+\dot{f}+\frac{\Lambda R^2}{\alpha_0^2}\mathrm{e}^{2\tau}f^3=0$$

$$f(\tau)=\phi(\tau)\mathrm{e}^{-\tau}$$

$$\ddot{\phi}-\dot{\phi}=-\frac{\Lambda R^2}{\alpha_0^2}\phi^3$$

$$\ddot{\phi}\approx-\frac{\Lambda R^2}{\alpha_0^2}\phi^3$$

$$\phi\approx f_0\mathrm{sn}\left(\frac{f_0\sqrt{\Lambda R^2}}{\alpha_0}\tau\middle|-1\right)$$

$$f_0 \gg \frac{\alpha_0}{\sqrt{\Lambda}R}.$$

$$f_0 \ll \frac{\alpha_0}{\sqrt{\Lambda}R}.$$

$$\frac{m_{\mathcal{K}}^2}{m_{\text{cosm}}^2} \sim \frac{1}{\Lambda}$$

$$\begin{aligned}\mathcal{T}^{\mu\nu} =& [T^\mu, T^\sigma][T^\nu, T_\sigma] + [T^\mu, T^{\mathrm{I}}][T^\nu, T_{\mathrm{I}}] \\ & -\frac{1}{4}\eta^{\mu\nu}\left([T^\alpha, T^\sigma][T_\alpha, T_\sigma] + 2[T^\alpha, T^{\mathrm{I}}][T_\alpha, T_{\mathrm{I}}] + [T^{\mathrm{I}}, T^{\mathrm{J}}]\left[T_{\mathrm{I}}, T_{\mathrm{J}}\right]\right)\end{aligned}$$

$$[T^\mu, T^\nu] = i\{\alpha t^\mu, \alpha t^\nu\} = -i\left[-\frac{1}{r^2R^2}\alpha^2\theta^{\mu\nu} + \frac{\alpha\alpha'}{R}(t^\mu x^\nu - x^\mu t^\nu)\right]$$

$$\begin{aligned}[T^\mu, T^\nu][T^\rho, T_\nu] =& -\frac{A^2}{r^4R^4}\theta^{\mu\nu}\theta_\nu^\rho - \frac{(A')^2}{4R^2}(t^\mu x^\nu - x^\mu t^\nu)(t^\rho x_\nu - t_\nu x^\rho) \\ & + \frac{AA'}{2r^2R^3}\theta^{\mu\nu}(t^\rho x_\nu - t_\nu x^\rho) + \frac{AA'}{2r^2R^3}(t^\mu x^\nu - x^\mu t^\nu)\theta_\nu^\rho \\ =& -\frac{A^2}{r^2R^2}\eta^{\mu\rho} - \frac{1}{R^2}\left(-A^2 - \frac{(A')^2(x_4^2 + R^2)}{4} - AA'x_4\right)t^\mu t^\rho \\ & -\frac{1}{r^2R^4}\left(A^2 + \frac{(A')^2}{4}(R^2 + x_4^2) + AA'x_4\right)x^\mu x^\rho \\ \approx& -\frac{A^2}{r^2R^2}\eta^{\mu\rho} + \frac{1}{4R^2}(2A + \dot{A})^2\left(t^\mu t^\rho - \frac{x^\mu x^\rho}{r^2R^2}\right)\end{aligned}$$

$$[T^\mu, T^\nu]\left[T_\mu, T_\nu\right] \approx \frac{1}{r^2R^2}\left(-4A^2 + 2\frac{(A + \dot{A}/2)^2 x_4^2}{r^2}\right)$$

$$\begin{aligned}[T^\mu, T^{\mathrm{I}}][T^\nu, T^{\mathrm{I}}] &= -\alpha^2|f'|^2C^2R^{-2}x^\mu x^\nu \\ [T^{\mathrm{I}}, T^{\mathrm{J}}]\left[T_{\mathrm{I}}, T_{\mathrm{J}}\right] &= -|f|^4\Theta^{\mathrm{IJ}}\Theta_{\mathrm{IJ}}\end{aligned}$$

$$\mathcal{T}^{\mu\nu} = -\eta^{\mu\nu}\left(\frac{P}{2R^2}x_4^2 + \frac{Q}{2R^2} - S\right) + r^2Pt^\mu t^\nu - (P + Qx_4^{-2})\frac{x^\mu x^\nu}{R^2}$$

$$\begin{aligned}P &= \frac{(A + \dot{A}/2)^2}{r^2R^2} \\ Q &= A|\dot{f}|^2C^2 \\ S &= \frac{|f|^4\Theta^{\mathrm{IJ}}\Theta_{\mathrm{IJ}}}{4} = \frac{|f|^4\Lambda C^2}{4}\end{aligned}$$

$$[T_A, \mathcal{T}^{A\nu}] = \left[T_\mu, \mathcal{T}^{\mu\nu}\right] + [T_{\mathrm{I}}, \mathcal{T}^{\mathrm{I}\nu}] = 0$$

$$i\left\{\alpha t_\mu, -\eta^{\mu\nu}\left(\frac{P}{2R^2}x_4^2+\frac{Q}{2R^2}-S\right)\right\}=i\alpha\frac{x_4x^\nu}{R^3}\left(\frac{\dot{P}+2P}{2}+\frac{\dot{Q}}{2}x_4^{-2}-\dot{S}R^2x_4^{-2}\right)$$

$$i\{\alpha t_\mu, r^2Pt^\mu t^\nu\}=-i\frac{\theta^{\mu\nu}t_\mu}{R^2}\alpha P+i\frac{1}{R^3}\alpha' Px_4^2x^\nu$$

$$=i\frac{x_4P}{R^3}(\alpha+\dot{\alpha})x^\nu-i\{\alpha t_\mu,(P+Qx_4^{-2})x^\mu x^\nu/R^2\}$$

$$=i\frac{x_4x^\nu}{R^3}\left[\alpha\left(-3P-\frac{\dot{P}}{2}-x_4^{-2}\left(\frac{\dot{Q}}{2}+3Q+\dot{S}R^2\right)\right)-\dot{\alpha}Qx_4^{-2}\right]$$

$$[T_{\rm I},\mathcal{T}^{{\rm I}\nu}]=\left[T_{\rm I},[T^{\rm I},T^\mu]\left[T^\nu,T_\mu\right]\right]+\left[T_{\rm I},[T^{\rm I},T^{\rm J}]\left[T^\nu,T_{\rm J}\right]\right]$$

$$\left[T_{\rm I},[T^{\rm I},\alpha t^\mu]\left[\alpha t^\nu,\alpha t_\mu\right]\right]$$

$$=-\frac{1}{2R}\left[fK_{\rm I}^+,\alpha\bar{f}'x_\mu\left(\frac{A}{r^2R^2}\theta^{\nu\mu}+\frac{A'}{2R}(x^\nu t^\mu-t^\nu x^\mu)K^{{\rm I},-}\right)\right]+h.c.$$

$$=-\frac{1}{2R^2}\left[f\mathcal{K}_{\rm I}^+,\alpha\bar{f}'(A+\dot{A}/2)x_4t^\nu\mathcal{K}^{{\rm I},-}\right]+h.c.$$

$$=-i\frac{C^2}{R^3}\alpha(A+\dot{A}/2)|\dot{f}|^2x_4^{-1}x^\nu-\frac{\alpha(A+\dot{A}/2)x_4}{2R^2}f\bar{f}'[\mathcal{K}_{\rm I}^+,\mathcal{K}^{{\rm I}-}]t^\nu+h.c.$$

$$=-i\frac{Q}{R^3}(\alpha+\dot{\alpha})x_4^{-1}x^\nu-\frac{\alpha(A+\dot{A}/2)}{2R^2}[\mathcal{K}_{\rm I}^+,\mathcal{K}^{{\rm I}-}]\left(f\bar{f}'-\bar{f}f'\right)t^\nu$$

$$\left[T_{\rm I},[T^{\rm I},T^{\rm J}]\left[\alpha t^\nu,T_{\rm J}\right]\right]=\left[T_{\rm I},[T^{\rm I},T^{\rm J}]\right]\left[\alpha t^\nu,T_{\rm J}\right]+[T^{\rm I},T^{\rm J}]\left[T_{\rm I},\left[\alpha t^\nu,T_{\rm J}\right]\right]$$

$$=-i\alpha\frac{1}{2R}\Lambda C^2|f|^2f\bar{f}'x^\nu+\text{ h.c. }+\alpha\frac{|f|^2}{4R}\bar{f}'\theta^{{\rm IJ},(+-)}\left[f\mathcal{K}_{\rm I}^+,\mathcal{K}_{\rm J}^-x^\nu\right]+\text{ h.c.}$$

$$=-i\frac{\alpha}{R}\dot{S}x_4^{-2}x^\nu+i\frac{\alpha}{R}\dot{S}x_4^{-2}x^\nu=0$$

$$0=-\left[\alpha\left(-3P-\frac{\dot{P}}{2}-x_4^{-2}\left(\frac{\dot{Q}}{2}+3Q+\dot{S}R^2\right)\right)-\dot{\alpha}Qx_4^{-2}\right]-2\alpha R^2\dot{S}x_4^{-2}-Q(\alpha+\dot{\alpha})x_4^{-2}$$

$$\overset{\frac{1}{\alpha}}{=}3P+\frac{\dot{P}}{2}+x_4^{-2}\left(\frac{\dot{Q}}{2}+2Q+\dot{S}R^2\right)$$

$$[\mathcal{K}_{\rm I}^+,\mathcal{K}^{{\rm I}-}]\left(f\bar{f}'-\bar{f}f'\right)=0$$

$$\nabla^\mu V_\mu=\gamma^{\mu\nu}\nabla_\nu V_\mu=\gamma^{\mu\nu}\left(\partial_\nu V_\mu-\Gamma_{\nu\mu}^\rho V_\rho\right)$$

$$=\gamma^{\mu\nu}\partial_\nu V_\mu+\frac{\rho^2}{\sqrt{|G|}}\partial_\sigma\left(\sqrt{|G|}G^{\rho\sigma}\right)V_\rho$$

$$=\frac{\rho^2}{\sqrt{|G|}}\partial_\sigma\left(\sqrt{|G|}G^{\rho\sigma}V_\rho\right)$$

$$=\rho^2\nabla^{(G)\mu}V_\mu$$

$$\gamma^{\mu\nu}\Gamma_{\nu\mu}^\rho=-\frac{\rho^2}{\sqrt{|G|}}\partial_\sigma\left(\sqrt{|G|}G^{\rho\sigma}\right)$$

$$E^{{\rm I}+,\mu}=\{f(x_4)\mathcal{K}^{{\rm I}+},x^\mu\}\approx r\dot{f}\mathcal{K}^{{\rm I}+}u^\mu$$

$$r\dot{f}|\mathcal{K}| \ll \alpha \sinh\, \tau \sim e^{(\varepsilon+1)\tau}.$$

$$\left[\mathcal{K}_{\mathrm{I}}, \left[\mathcal{K}^{\mathrm{I}}, \mathcal{K}_{\mathrm{J}}\right]\right] = \Lambda \mathcal{K}_{\mathrm{J}}, \mathcal{K}_{\mathrm{I}}\mathcal{K}^{\mathrm{I}} = C^2\mathbb{1}$$

$$\Theta^{\mathrm{IJ}}\Theta_{\mathrm{IJ}} = \Lambda C^2$$

$$\Theta^{\mathrm{IJ}} = -i[\mathcal{K}^{\mathrm{I}}, \mathcal{K}^{\mathrm{J}}]$$

$$\begin{aligned}\Lambda C^2 &= \Lambda \mathcal{K}_{\mathrm{J}}\mathcal{K}^{\mathrm{J}} = \mathcal{K}_{\mathrm{J}}\left[\mathcal{K}_{\mathrm{I}}, [\mathcal{K}^{\mathrm{I}}, \mathcal{K}^{\mathrm{J}}]\right] = 2C^4 - 2\mathcal{K}_{\mathrm{J}}\mathcal{K}_{\mathrm{I}}\mathcal{K}^{\mathrm{J}}\mathcal{K}^{\mathrm{I}} \\ &= -\left(-\mathcal{K}_{\mathrm{J}}\mathcal{K}_{\mathrm{I}}\mathcal{K}^{\mathrm{I}}\mathcal{K}^{\mathrm{J}} - \mathcal{K}_{\mathrm{I}}\mathcal{K}_{\mathrm{J}}\mathcal{K}^{\mathrm{J}}\mathcal{K}^{\mathrm{I}} + \mathcal{K}_{\mathrm{I}}\mathcal{K}_{\mathrm{J}}\mathcal{K}^{\mathrm{I}}\mathcal{K}^{\mathrm{J}} + \mathcal{K}_{\mathrm{J}}\mathcal{K}_{\mathrm{I}}\mathcal{K}^{\mathrm{J}}\mathcal{K}^{\mathrm{I}}\right) \\ &= \Theta_{\mathrm{IJ}}\Theta^{\mathrm{IJ}}\end{aligned}$$

$$\alpha(x_4) = \frac{R\alpha_0}{x_4^2 + R^2}\sqrt{R^2 - x_4^2 + R\alpha_1 x_4}, x_4 < R$$

$$\begin{aligned}\mathbf{Q}_{ab} &= \mathfrak{D}g_{ab} = \mathbf{d}g_{ab} - g_{cb}\boldsymbol{\Sigma}_a^c - g_{ac}\boldsymbol{\Sigma}_b^c \\ \mathfrak{T}^a &= \mathfrak{D}\mathbf{e}^a = \mathbf{d}\mathbf{e}^a - \mathbf{e}^b \wedge \boldsymbol{\Sigma}_b^a\end{aligned}$$

$$\mathbf{Q}_{ab} = \cancel{\mathbf{d}\eta_{ab}} - \eta_{cb}\boldsymbol{\Sigma}^c{}_a - \eta_{ac}\boldsymbol{\Sigma}^c{}_b = -(\boldsymbol{\Sigma}_{ba} + \boldsymbol{\Sigma}_{ab})$$

$$\boldsymbol{\mathfrak{T}}^a = \cancel{\mathbf{d}^2 x^a} - \mathbf{e}^b \wedge \boldsymbol{\Sigma}^a{}_b = -\Sigma^a{}_{\mu\nu}\mathbf{d}x^\mu \wedge \mathbf{d}x^\nu$$

$$e\Delta_i{}^{jk} \equiv -\frac{\delta \mathcal{L}_{\text{source}}}{\delta \Sigma^i{}_{jk}}$$

$$\begin{aligned}\mathbf{d}\mathbf{e}^a &= \mathbf{e}^b \wedge \boldsymbol{\Sigma}^a{}_b + T^a \\ \mathbf{d}\boldsymbol{\Sigma}^a{}_b &= \boldsymbol{\Sigma}^c{}_b \wedge \boldsymbol{\Sigma}^a{}_c + \Re^a{}_b\end{aligned}$$

$$\mathbf{Q}_{ab} = \mathfrak{D}g_{ab} = \mathbf{d}g_{ab} - g_{cb}\boldsymbol{\Sigma}_a^c - g_{ac}\boldsymbol{\Sigma}_b^c$$

$$\begin{aligned}\mathbf{dA} &= \mathbf{F} \\ \mathbf{dW}^a &= -\frac{1}{2}c^a{}_{bc}\mathbf{W}^b \wedge \mathbf{W}^c + \mathbf{F}^a \ (a, b, c = 1,2,3) \\ \mathbf{dB}^K &= -\frac{1}{2}f^K{}_{LM}\mathbf{B}^L \wedge \mathbf{B}^M + \mathbf{H}^K \ (K, L, M = 1, \ldots, 8)\end{aligned}$$

$$\boldsymbol{\Sigma}_{ab} = -\frac{1}{2}\mathbf{Q}_{ab} + \boldsymbol{\Omega}_{ab}$$

$$\mathbf{d}\mathbf{e}^a = \mathbf{e}^b \wedge \boldsymbol{\Omega}^a{}_b + T^a - \mathbf{Q}^a$$

$$\mathbf{Q}^a \equiv \frac{1}{2}\mathbf{e}^b \wedge \mathbf{Q}^a{}_b$$

$$\mathbf{d}\mathbf{e}^a = \mathbf{e}^b \wedge \boldsymbol{\Omega}_b^a + \mathbf{T}^a$$

$$\mathbf{d}\mathbf{e}^a = \mathbf{e}^b \wedge \boldsymbol{\omega}^a{}_b$$

$$\mathbf{\Omega}^{a}{}_{b} = \boldsymbol{\omega}^{a}{}_{b} - \mathbf{C}^{a}{}_{b}$$

$$0 = \mathbf{e}^{b} \wedge \mathbf{e}^{c} \left(-C^{a}{}_{bc} + \frac{1}{2} T^{a}{}_{bc} \right)$$

$$C_{abc} = \frac{1}{2}(T_{abc} + T_{bca} - T_{cab})$$

$$\mathbf{\Omega}^{a}{}_{b} = \boldsymbol{\omega}^{a}{}_{b} - \mathbf{C}^{a}{}_{b}$$

$$\mathbf{\Sigma}^{a}{}_{b} = \boldsymbol{\omega}^{a}{}_{b} - \mathbf{C}^{a}{}_{b} - \frac{1}{2}\mathbf{Q}^{a}{}_{b}$$

$$\mathfrak{R}^{ab} = \mathcal{R}^{ab} - \frac{1}{2}\mathbf{D}\mathbf{Q}^{ab} - \frac{1}{4}\mathbf{Q}^{cb} \wedge \mathbf{Q}^{a}{}_{c} - \mathbf{Q}^{c(a} \wedge \mathbf{C}^{b)}{}_{c}$$

$$S_{\text{Grav}} = \frac{\kappa}{2} \int \mathfrak{R}^{ab} \wedge \mathbf{e}^{c} \wedge \mathbf{e}^{d} e_{abcd}$$

$$-\frac{1}{2}\mathbf{D}\mathbf{Q}^{ab} - \mathbf{Q}^{c(a} \wedge \mathbf{C}^{b)}{}_{c}$$

$$S_{Grav}[\Omega, Q] = \frac{\kappa}{2} \int \left(R^{ab} - \frac{1}{4}\mathbf{Q}^{bc} \wedge \mathbf{Q}^{a}_{c} \right) \wedge \mathbf{e}^{c} \wedge \mathbf{e}^{d} e_{abcd}$$

$$\begin{aligned} \delta_{\Omega} S_{\text{Grav}} &= \kappa \int \delta \mathbf{\Omega}^{eb} \wedge \mathbf{C}^{a}_{e} \wedge \mathbf{e}^{c} \wedge \mathbf{e}^{d} e_{abcd} \\ &= \kappa \int \delta \Omega^{eb}{}_{f} \left(C^{f}{}_{eb} - C^{f}{}_{be} - C^{a}{}_{ea}\delta^{f}_{b} + C^{a}{}_{ba}\delta^{f}_{e} \right) \mathbf{\Phi} \end{aligned}$$

$$\mathcal{T}^{g}{}_{eb} = C^{g}{}_{eb} - C^{g}{}_{be} - C^{a}{}_{ea}\delta^{g}_{b} + C^{a}{}_{ba}\delta^{g}_{e}$$

$$\mathcal{T}^{a}_{bc} = T^{a}_{bc} - \delta^{a}_{b} T^{e}_{ec} + \delta^{a}_{c} T^{e}_{eb}$$

$$\kappa \mathcal{T}^{a}{}_{bc} = -\frac{\delta \mathcal{L}_{\text{Source}}}{\delta \Omega^{abc}}$$

$$\delta_{Q} S_{Grav} = -\frac{\kappa}{4} \int \delta_{Q} Q^{eb}_{f} \left(Q^{f}_{eb} + Q^{f}_{be} + Q^{a}_{ea}\delta^{f}_{b} + Q^{a}_{ba}\delta^{f}_{e} \right) \mathbf{\Phi}$$

$$\mathcal{Q}_{acb} \equiv -\frac{1}{4}(Q_{acb} + Q_{abc} - Q^{e}_{be}\eta_{ca} - Q^{e}_{ce}\eta_{ba})$$

$$\mathcal{Q}_{cba} = \frac{4}{\kappa} \frac{\delta \mathcal{L}_{\text{Source}}}{\delta Q^{abc}}$$

$$\mathcal{Q}_{acb} = -\left(\frac{1}{2} Q_{a(bc)} - \frac{1}{4} \eta^{de} \left(Q_{b(de)}\eta_{ca} + Q_{c(de)}\eta_{ba} \right) \right)$$

$$\delta S_{Grav} = \kappa \int \left(\delta \Omega^{ab}{}_{c} \mathcal{T}^{c}{}_{ab} - \frac{1}{4} \delta Q^{bc}{}_{a} \mathcal{Q}^{a}{}_{bc} \right) \mathbf{\Phi}$$

$$S_{D,V} = \alpha \int \psi^{\dagger} h \left(i\not{\partial} - m \right) \psi \, d^4x$$

$$\langle \chi, \psi \rangle = \chi^{\dagger} h \psi \equiv \bar{\chi} \psi \oslash \not{\partial} = \gamma^a \partial_a$$

$$\{\gamma^a, \gamma^b\} = -2\eta^{ab}\mathbf{1}$$

$$\Lambda(w^{ab}) = \exp\left(\frac{1}{2} w_{ab} \sigma^{ab}\right)$$

$$\Gamma^{\Delta} = \{1, \gamma^a, \sigma^{ab}, \gamma_5 \gamma^d, \gamma_5\}$$

$$\sigma^{ab} = \tfrac{1}{2}[\gamma^a, \gamma^b], \gamma_5 \gamma^d = \tfrac{i}{3!} e^d{}_{abc} \gamma^a \gamma^b \gamma^c \text{ and } \gamma_5 = \tfrac{i}{4!} \varepsilon_{abcd} \gamma^a \gamma^b \gamma^c \gamma^d$$

$$\frac{1}{4} \mathrm{tr}\left(\Gamma^{\Delta} \Gamma^{\Sigma \dagger}\right) = \delta^{\Delta\Sigma}$$

$$\bar{\psi}_A \left[\beta_{\Delta} \Gamma^{\Delta}\right]^A{}_B \psi^B \otimes \gamma^a = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes \gamma^i = \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}$$

$$\langle \chi, \psi \rangle = \chi^{\dagger} h \psi \equiv \bar{\chi} \psi$$

$$\bar{\chi} = \chi^{\dagger} h = \chi^{\dagger} \gamma^0$$

$$\sigma^{ij} = -i\varepsilon^{ij}{}_k \begin{pmatrix} \sigma^k & \\ & \sigma^k \end{pmatrix} \qquad \sigma^{0i} = \begin{pmatrix} & \sigma^i \\ \sigma^i & \end{pmatrix} \qquad \gamma_5 \gamma^0 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

$$\gamma_5 \gamma^i = \begin{pmatrix} -\sigma^i & 0 \\ 0 & \sigma^i \end{pmatrix} \qquad \gamma_5 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$\hat{\gamma}^a \equiv (\gamma^0, \gamma^1, i\gamma^2, \gamma^3)$$

$$\{\hat{\gamma}^a, \hat{\gamma}^b\} = -2\hat{\eta}^{ab}$$

$$\hat{\sigma}^{AB} = \frac{1}{2}[\hat{\gamma}^A, \hat{\gamma}^B] = (\sigma^{01}, i\sigma^{02}, \sigma^{03}, i\sigma^{12}, i\sigma^{23}, \sigma^{31})$$

$$\hat{\Gamma}^{\Delta} = \{1, \hat{\gamma}^A, \hat{\sigma}^{AB}, \hat{\gamma}_5 \hat{\gamma}^A, \hat{\gamma}_5\}$$

$$\not{\partial} = \gamma^a e_a{}^{\mu} \partial_{\mu} \rightarrow \not{D} = \gamma^a e_a{}^{\mu} D_{\mu}$$

$$D_{\mu} \psi = \partial_{\mu} \psi - b_{A\mu} \hat{\Gamma}^A \psi$$

$$S_{\text{interaction}} = -\alpha Re \int b_{aA} \psi^{\dagger} i h \gamma^a \hat{\Gamma}^A \psi$$

$$i\eta h \gamma^2 = \begin{pmatrix} & -\sigma^1 \\ i\sigma^2 & \end{pmatrix}, \eta h \hat{\gamma}_5 \hat{\gamma}^2 = \begin{pmatrix} \sigma^1 & \\ & -i\sigma^2 \end{pmatrix}, \eta h \hat{\gamma}_5 = \begin{pmatrix} & -\sigma^3 \\ -1 & \end{pmatrix}$$
$$\eta h \sigma^{01} = \begin{pmatrix} & -i\sigma^2 \\ -\sigma^1 & \end{pmatrix}, \eta h \sigma^{03} = \begin{pmatrix} & -1 \\ -\sigma^3 & \end{pmatrix}, \eta h \sigma^{31} = \begin{pmatrix} \sigma^1 & \\ & i\sigma^2 \end{pmatrix}$$

$$\left[K^i\right]^a{}_b = \delta_0^a\delta_b^i + \delta^{ia}\delta_{0b}, \left[J^i\right]^a{}_b = \varepsilon^{ia}{}_b$$

$$K_x = \frac{1}{2}(\eta h\hat{\gamma}_5\hat{\gamma}^2 + \eta h\sigma^{31}) \quad J_x = \frac{1}{2}(\eta h\sigma^{01} - i\eta h\gamma^2)$$
$$K_y = -\frac{1}{2}(\eta h\hat{\gamma}_5 + \eta h\sigma^{03}) \quad J_y = \frac{1}{2}(\eta h\sigma^{03} - \eta h\hat{\gamma}_5)$$
$$K_z = -\frac{1}{2}(i\eta h\gamma^2 + \eta h\sigma^{01}) \quad J_z = \frac{1}{2}(\eta h\sigma^{31} - \eta h\hat{\gamma}_5\hat{\gamma}^2)$$

$$\eta^{AC}[h\hat{\Gamma}]_{(CB)} \in \hat{\Gamma}^{\Delta_s}\{h1, h\gamma^0, ih\sigma^{12}, ih\sigma^{23}, h\gamma_5\gamma^1, h\gamma_5\gamma^3, h\gamma^1h\gamma^3, ih\sigma^{02}, h\gamma_5\gamma^0\}$$

$$\eta h1 = \begin{pmatrix} -\sigma^3 & \\ & -1 \end{pmatrix}, \eta h\gamma^0 = \begin{pmatrix} -\sigma^3 & \\ & 1 \end{pmatrix}, \eta h\gamma^1 = \begin{pmatrix} & -i\sigma^2 \\ -\sigma^1 & \end{pmatrix}, \eta h\gamma^3 = \begin{pmatrix} & -1 \\ \sigma^3 & \end{pmatrix},$$

$$i\eta h\sigma^{02} = \begin{pmatrix} & -\sigma^1 \\ -i\sigma^2 & \end{pmatrix}, i\eta h\sigma^{12} = \begin{pmatrix} -1 & \\ & -\sigma^3 \end{pmatrix}, i\eta h\sigma^{23} = \begin{pmatrix} -i\sigma^2 & \\ & -\sigma^1 \end{pmatrix},$$
$$\eta h\hat{\gamma}_5\hat{\gamma}^1 = \begin{pmatrix} i\sigma^2 & \\ & -\sigma^1 \end{pmatrix}, \eta h\hat{\gamma}_5\hat{\gamma}^3 = \begin{pmatrix} 1 & \\ & -\sigma^3 \end{pmatrix}, \eta h\hat{\gamma}_5\hat{\gamma}^2 = \begin{pmatrix} \sigma^1 & \\ & -i\sigma^2 \end{pmatrix}$$

$$\mathbf{\Sigma} = \mathbf{b}_\Delta\Gamma^\Delta = \mathbf{b}_{\Delta_s}\Gamma^{\Delta_s} + \mathbf{b}_{\Delta_a}\Gamma^{\Delta_a}$$

$$\begin{aligned}
\Omega^A_{Bc} = b_{c\Delta_a}\left[\Gamma^{\Delta_a}\right]^A_B =& [ib_{c2}\eta h\gamma^2 + b_{c01}\eta h\sigma^{01} + b_{c03}\eta h\sigma^{03} + b_{c31}\eta h\sigma^{31} + ib_{c52}\eta h\gamma_5\gamma^2 + b_{c5}\eta h\gamma_5]^A_B \\
Q^A_{Bc} = b_{c\Delta_s}\left[\Gamma^{\Delta_s}\right]^A_B =& [b_c\eta h1 + b_{c0}\eta h\gamma^0 + b_{c1}\eta h\gamma^1 + b_{c3}\eta h\gamma^3 + ib_{c02}\eta h\sigma^{02} + ib_{c12}\eta h\sigma^{12} + ib_{c23}\eta h\sigma^{23} \\
&+ b_{50}\eta h\gamma_5\gamma^0 + b_{c51}\eta h\gamma_5\gamma^1 + b_{c53}\eta h\gamma_5\gamma^3]^A_B
\end{aligned}$$

$$\delta\Omega^{ab}{}_c\mathcal{T}^c_{ab} = b_{c\Delta_a}\left[\Gamma^{\Delta_a}\right]^{AB}\mathcal{T}^c{}_{AB} = -b_{c\Delta_a}\mathrm{tr}\left(\Gamma^{\Delta_a}\mathcal{T}^c\right)$$
$$-\frac{1}{4}\delta Q^{bc}{}_a\mathcal{Q}^a{}_{bc} = -\frac{1}{4}b_{c\Delta_s}\left[\Gamma^{\Delta_s}\right]^{AB}\mathcal{Q}^c{}_{AB} = -\frac{1}{4}b_{c\Delta_s}\mathrm{tr}\left(\Gamma^{\Delta_s}\mathcal{Q}^c\right)$$

$$-\kappa\delta b_{c\Delta_a}\mathrm{tr}\left(\Gamma^{\Delta_a}\mathcal{T}^c\right) - \frac{\kappa}{4}\delta b_{c\Delta_s}\mathrm{tr}\left(\Gamma^{\Delta_s}\mathcal{Q}_c\right) = \alpha\mathrm{Re}\left(i\psi^\dagger h\gamma^c\delta b_{\Delta c}\hat{\Gamma}^\Delta\psi\right)$$

$$\begin{aligned}
\delta\mathcal{L}_{\mathrm{Grav}} =& \kappa\delta\Omega^{AB}{}_c\mathcal{T}^c{}_{AB} - \frac{\kappa}{4}\delta Q^{ABc}\mathcal{Q}_{c(AB)} \\
=& -\kappa\delta b_{c2}\mathrm{tr}[i\eta h\gamma^2\mathcal{T}^c] - \kappa\delta b_{c01}\mathrm{tr}[\eta h\sigma^{01}\mathcal{T}^c] - \kappa\delta b_{c03}\mathrm{tr}[\eta h\sigma^{03}\mathcal{T}^c] \\
& -\kappa\delta b_{c13}\mathrm{tr}[\eta h\sigma^{13}\mathcal{T}^c] - \kappa\delta b_{c52}\mathrm{tr}[i\eta h\gamma_5\gamma^2\mathcal{T}^c] - \kappa\delta b_{c5}\mathrm{tr}[\eta h\gamma_5\mathcal{T}^c] \\
& -\frac{\kappa}{4}\delta b_c\mathrm{tr}[\eta h1\mathcal{Q}^c] - \frac{\kappa}{4}\delta b_{c0}\mathrm{tr}[\eta h\gamma^0\mathcal{Q}^c] - \frac{\kappa}{4}\delta b_{c1}\mathrm{tr}[\eta h\gamma^1\mathcal{Q}^c] - \frac{\kappa}{4}\delta b_{c3}\mathrm{tr}[\eta h\gamma^3\mathcal{Q}^c] \\
& -\frac{\kappa}{4}\delta b_{c02}\mathrm{tr}[i\eta h\sigma^{02}\mathcal{Q}^c] - \frac{\kappa}{4}\delta b_{c12}\mathrm{tr}[i\eta h\sigma^{12}\mathcal{Q}^c] - \frac{\kappa}{4}\delta b_{c23}\mathrm{tr}[i\eta h\sigma^{23}\mathcal{Q}^c] \\
& -\frac{\kappa}{4}\delta b_{c50}\mathrm{tr}[\eta h\gamma_5\gamma^0\mathcal{Q}^c] - \frac{\kappa}{4}\delta b_{c51}\mathrm{tr}[\eta h\gamma_5\gamma^1\mathcal{Q}^c] - \frac{\kappa}{4}\delta b_{c53}\mathrm{tr}[\eta h\gamma_5\gamma^3\mathcal{Q}^c]
\end{aligned}$$

$$\begin{aligned}
-[i\eta h\gamma^2]^a{}_b\mathcal{T}^{cb}{}_a &= \mathcal{T}^c{}_{30} + \mathcal{T}^c{}_{21} - \mathcal{T}^c{}_{12} - \mathcal{T}^c{}_{03} \\
-[\eta h\sigma^{01}]^a{}_b\mathcal{T}^{cb}{}_a &= \mathcal{T}^c{}_{30} - \mathcal{T}^c{}_{21} + \mathcal{T}^c{}_{12} - \mathcal{T}^c{}_{03} \\
-[i\eta h\gamma_5\gamma^2]^a{}_b\mathcal{T}^{cb}{}_a &= -\mathcal{T}^c{}_{10} + \mathcal{T}^c{}_{01} + \mathcal{T}^c{}_{32} - \mathcal{T}^c{}_{23} \\
-[\eta h\sigma^{31}]^a{}_b\mathcal{T}^{cb}{}_a &= -\mathcal{T}^c{}_{10} + \mathcal{T}^c{}_{01} - \mathcal{T}^c{}_{32} + \mathcal{T}^c{}_{23} \\
-[\eta h\gamma_5]^a{}_b\mathcal{T}^{cb}{}_a &= \mathcal{T}^c{}_{20} - \mathcal{T}^c{}_{31} - \mathcal{T}^c{}_{02} + \mathcal{T}^c{}_{13} \\
-[\eta h\sigma^{03}]^a{}_b\mathcal{T}^{cb}{}_a &= \mathcal{T}^c{}_{20} + \mathcal{T}^c{}_{31} - \mathcal{T}^c{}_{02} - \mathcal{T}^c{}_{13}
\end{aligned}$$

$$\begin{aligned}
\mathrm{tr}[\eta h 1 \mathcal{Q}^c] &= \mathcal{Q}^c{}_{00} + \mathcal{Q}^c{}_{11} - \mathcal{Q}^c{}_{22} - \mathcal{Q}^c \\
\mathrm{tr}[\eta h \gamma^0 \mathcal{Q}^c] &= \mathcal{Q}^c{}_{00} + \mathcal{Q}^c{}_{11} + \mathcal{Q}^c{}_{22} + \mathcal{Q}^c{}_{33} \\
\mathrm{tr}[\eta h \gamma^1 \mathcal{Q}^c] &= -\mathcal{Q}^c{}_{30} + \mathcal{Q}^c{}_{21} + \mathcal{Q}^c{}_{12} - \mathcal{Q}^c{}_{03} \\
\mathrm{tr}[\eta h \gamma^3 \mathcal{Q}^c] &= -\mathcal{Q}^c{}_{20} - \mathcal{Q}^c{}_{31} - \mathcal{Q}^c{}_{02} - \mathcal{Q}^c{}_{13} \\
\mathrm{tr}[i\eta h \sigma^{02} \mathcal{Q}^c] &= -\mathcal{Q}^c{}_{30} - \mathcal{Q}^c{}_{12} - \mathcal{Q}^c{}_{03} - \mathcal{Q}^c{}_{21} \\
\mathrm{tr}[i\eta h \sigma^{12} \mathcal{Q}^c] &= \mathcal{Q}^c{}_{00} - \mathcal{Q}^c{}_{11} - \mathcal{Q}^c{}_{22} + \mathcal{Q}^c \\
\mathrm{tr}[i\eta h \sigma^{32} \mathcal{Q}^c] &= \mathcal{Q}^c{}_{10} + \mathcal{Q}^c{}_{01} + \mathcal{Q}^c{}_{32} + \mathcal{Q}^c{}_{23} \\
\mathrm{tr}[\eta h \gamma_5 \gamma^0 \mathcal{Q}^c] &= \mathcal{Q}^c{}_{20} - \mathcal{Q}^c{}_{31} + \mathcal{Q}^c{}_{02} - \mathcal{Q}^c{}_{13} \\
\mathrm{tr}[\eta h \gamma_5 \gamma^1 \mathcal{Q}^c] &= \mathcal{Q}^c{}_{10} + \mathcal{Q}^c{}_{01} - \mathcal{Q}^c{}_{32} - \mathcal{Q}^c{}_{23} \\
\mathrm{tr}[\eta h \gamma_5 \gamma^3 \mathcal{Q}^c] &= -\mathcal{Q}^c{}_{00} + \mathcal{Q}^c{}_{11} - \mathcal{Q}^c{}_{22} + \mathcal{Q}^c
\end{aligned}$$

$$\begin{aligned}
\delta\mathcal{L}_{\text{interaction}} =& -\alpha \mathrm{Re}\left(\psi^\dagger (i\delta b_{a01} h\gamma^a \eta h \sigma^{01} + i\delta b_{a03} h\gamma^a \eta h \sigma^{03} + i\delta b_{a13} h\gamma^a \eta h \sigma^{13})\psi\right) \\
& -\alpha \mathrm{Re}\left(\psi^\dagger (i\delta b_{a2} i h\gamma^a \eta h \gamma^2 + i\delta b_{a52} i h\gamma^a \eta h \gamma_5 \gamma^2 + i\delta b_{a5} h\gamma^a \eta h \gamma_5)\psi\right) \\
& -\alpha\delta b_c \mathrm{Re}\psi^\dagger (i h\gamma^c \eta h 1)\psi - \alpha \mathrm{Re}\left(\psi^\dagger (i\delta b_{a0} h\gamma^a \eta h \gamma^0 + i\delta b_{a1} h\gamma^a \eta h \gamma^1 + i\delta b_{a3} h\gamma^a \eta h \gamma^3)\psi\right) \\
& -\alpha\left(\mathrm{Re}\psi^\dagger (i\delta b_{a50} h\gamma^a \eta h \gamma_5 \gamma^0 + i\delta b_{a51} h\gamma^a \eta h \gamma_5 \gamma^1 + i\delta b_{a53} h\gamma^a \eta h \gamma_5 \gamma^3)\psi\right) \\
& -\alpha \mathrm{Re}\left(\psi^\dagger (i\delta b_{a02} i h\gamma^a \eta h \sigma^{02} + i\delta b_{a12} i h\gamma^a \eta h \sigma^{12} + i\delta b_{a32} i h\gamma^a \eta h \sigma^{32})\psi\right)
\end{aligned}$$

$$\psi = \begin{pmatrix} \mu \\ \nu \\ \rho \\ \sigma \end{pmatrix}, \psi^\dagger = (\bar{\mu}, \bar{\nu}, \bar{\rho}, \bar{\sigma})$$

$$\begin{aligned}
-\alpha \mathrm{Re}\left(\psi^\dagger b_{31} i h\gamma^3 \eta h \gamma^1 \psi\right) &= -\alpha b_{31} \mathrm{Re}\left(i\psi^\dagger \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} & \sigma^3 \\ -\sigma^3 & \end{pmatrix} \begin{pmatrix} -\sigma^3 & \\ & 1 \end{pmatrix} \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} \begin{pmatrix} & \sigma^1 \\ -\sigma^1 & \end{pmatrix} \psi\right) \\
&= -i\alpha b_{31} (\bar{\mu}\nu - \bar{\nu}\mu)
\end{aligned}$$

$$\begin{aligned}
[\eta h i\gamma^2]^a{}_b \mathcal{T}^{cb}{}_a &= -\frac{\alpha}{\kappa} \mathrm{Re}\left(\psi^\dagger i h\gamma^c i\eta h \gamma^2 \psi\right) \\
[\eta h \sigma^{01}]^a{}_b \mathcal{T}^{cb}{}_a &= -\frac{\alpha}{\kappa} \mathrm{Re}\left(\psi^\dagger i h\gamma^c \eta h \sigma^{01} \psi\right) \\
[\eta h \gamma_5 i\gamma^2]^a{}_b \mathcal{T}^{cb}{}_a &= -\frac{\alpha}{\kappa} \mathrm{Re}\left(\psi^\dagger i h\gamma^c i\eta h \gamma_5 \gamma^2 \psi\right) \\
[\eta h \sigma^{31}]^a{}_b \mathcal{T}^{cb}{}_a &= -\frac{\alpha}{\kappa} \mathrm{Re}\left(\psi^\dagger i h\gamma^c \eta h \sigma^{31} \psi\right) \\
[\eta h \gamma_5]^a{}_b \mathcal{T}^{cb}{}_a &= -\frac{\alpha}{\kappa} \mathrm{Re}\left(\psi^\dagger i h\gamma^c \eta h \gamma_5 \psi\right) \\
[\eta h \sigma^{03}]^a{}_b \mathcal{T}^{cb}{}_a &= -\frac{\alpha}{\kappa} \mathrm{Re}\left(\psi^\dagger i h\gamma^c \eta h \sigma^{03} \psi\right)
\end{aligned}$$

$$[\eta h]^c{}_b\mathcal{Q}^{ab}{}_c = -\frac{4\alpha}{\kappa}\mathrm{Re}\left(\psi^\dagger i h\gamma^a\eta h\psi\right)$$
$$[\eta h\gamma^0]^c{}_b\mathcal{Q}^{ab}{}_c = -\frac{4\alpha}{\kappa}\mathrm{Re}\left(\psi^\dagger i h\gamma^a\eta h\gamma^0\psi\right)$$
$$[\eta h\gamma^1]^c{}_b\mathcal{Q}^{ab}{}_c = -\frac{4\alpha}{\kappa}\mathrm{Re}\left(\psi^\dagger i h\gamma^a\eta h\gamma^1\psi\right)$$
$$[\eta h\gamma^3]^c{}_b\mathcal{Q}^{ab}{}_c = -\frac{4\alpha}{\kappa}\mathrm{Re}\left(\psi^\dagger i h\gamma^a\eta h\gamma^3\psi\right)$$
$$[\eta h\gamma_5\gamma^0]^c{}_b\mathcal{Q}^{ab}{}_c = -\frac{4\alpha}{\kappa}\mathrm{Re}\left(\psi^\dagger i h\gamma^a\eta h\gamma_5\gamma^0\psi\right)$$
$$[\eta h\gamma_5\gamma^1]^c{}_b\mathcal{Q}^{ab}{}_c = -\frac{4\alpha}{\kappa}\mathrm{Re}\left(\psi^\dagger i h\gamma^a\eta h\gamma_5\gamma^1\psi\right)$$
$$[\eta h\gamma_5\gamma^3]^c{}_b\mathcal{Q}^{ab}{}_c = -\frac{4\alpha}{\kappa}\mathrm{Re}\left(\psi^\dagger i h\gamma^a\eta h\gamma_5\gamma^3\psi\right)$$
$$[i\eta h\sigma^{02}]^c{}_b\mathcal{Q}^{ab}{}_c = -\frac{4\alpha}{\kappa}\mathrm{Re}\left(\psi^\dagger i h\gamma^a i\eta h\sigma^{02}\psi\right)$$
$$[i\eta h\sigma^{12}]^c{}_b\mathcal{Q}^{ab}{}_c = -\frac{4\alpha}{\kappa}\mathrm{Re}\left(\psi^\dagger i h\gamma^a i\eta h\sigma^{12}\psi\right)$$
$$[i\eta h\sigma^{23}]^c{}_b\mathcal{Q}^{ab}{}_c = -\frac{4\alpha}{\kappa}\mathrm{Re}\left(\psi^\dagger i h\gamma^a i\eta h\sigma^{23}\psi\right)$$

$$R_{\bar{\alpha}\beta} = \frac{1}{2}(\bar{\alpha}\beta + \bar{\beta}\alpha)$$

$$I_{\bar{\alpha}\beta} = \frac{1}{2i}(\bar{\alpha}\beta - \bar{\beta}\alpha)$$
$$D_{\bar{\alpha}\alpha} = \bar{\alpha}\alpha$$

$$\mathcal{T}^0{}_{ab} = \frac{\alpha}{\kappa}\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -I_{\bar{\nu}\rho} & -I_{\bar{\nu}\sigma} \\ 0 & I_{\bar{\nu}\rho} & 0 & -I_{\bar{\rho}\sigma} \\ 0 & I_{\bar{\nu}\sigma} & I_{\bar{\rho}\sigma} & 0 \end{pmatrix}$$

$$\mathcal{T}^1{}_{ab} = \frac{\alpha}{2\kappa}\begin{pmatrix} 0 & I_{\bar{\nu}\sigma} + I_{\bar{\mu}\rho} & I_{\bar{\rho}\sigma} + I_{\bar{\mu}\nu} & 0 \\ -I_{\bar{\nu}\sigma} - I_{\bar{\mu}\rho} & 0 & 0 & I_{\bar{\mu}\nu} - I_{\bar{\rho}\sigma} \\ -I_{\bar{\rho}\sigma} - I_{\bar{\mu}\nu} & 0 & 0 & I_{\bar{\mu}\rho} - I_{\bar{\nu}\sigma} \\ 0 & I_{\bar{\rho}\sigma} - I_{\bar{\mu}\nu} & I_{\bar{\nu}\sigma} - I_{\bar{\mu}\rho} & 0 \end{pmatrix}$$

$$\mathcal{T}^2{}_{ab} = \frac{\alpha}{2\kappa}\begin{pmatrix} 0 & R_{\bar{\mu}\rho} - R_{\bar{\nu}\sigma} & -R_{\bar{\mu}\nu} - R_{\bar{\rho}\sigma} & (D_{\bar{\mu}\mu} - D_{\bar{\sigma}\sigma}) \\ R_{\bar{\nu}\sigma} - R_{\bar{\mu}\rho} & 0 & (D_{\bar{\nu}\nu} + D_{\bar{\rho}\rho}) & R_{\bar{\rho}\sigma} - R_{\bar{\mu}\nu} \\ R_{\bar{\mu}\nu} + R_{\bar{\rho}\sigma} & -(D_{\bar{\nu}\nu} + D_{\bar{\rho}\rho}) & 0 & -(R_{\bar{\mu}\rho} + R_{\bar{\nu}\sigma}) \\ -(D_{\bar{\mu}\mu} - D_{\bar{\sigma}\sigma}) & R_{\bar{\mu}\nu} - R_{\bar{\rho}\sigma} & (R_{\bar{\mu}\rho} + R_{\bar{\nu}\sigma}) & 0 \end{pmatrix}$$

$$\mathcal{T}^3{}_{ab} = \frac{\alpha}{2\kappa}\begin{pmatrix} 0 & I_{\bar{\nu}\rho} - I_{\bar{\mu}\sigma} & 0 & -(I_{\bar{\rho}\sigma} + I_{\bar{\mu}\nu}) \\ I_{\bar{\mu}\sigma} - I_{\bar{\nu}\rho} & 0 & I_{\bar{\mu}\nu} - I_{\bar{\rho}\sigma} & 0 \\ 0 & I_{\bar{\rho}\sigma} - I_{\bar{\mu}\nu} & 0 & -(I_{\bar{\mu}\sigma} + I_{\bar{\nu}\rho}) \\ I_{\bar{\rho}\sigma} + I_{\bar{\mu}\nu} & 0 & I_{\bar{\mu}\sigma} + I_{\bar{\nu}\rho} & 0 \end{pmatrix}$$

$$
\mathscr{Q}^{0}{}_{ab} = \frac{4\alpha}{\kappa}\begin{pmatrix} 0 & I_{\bar{\mu}\nu} & I_{\bar{\mu}\rho} & I_{\bar{\mu}\sigma} \\ I_{\bar{\mu}\nu} & 0 & 0 & 0 \\ I_{\bar{\mu}\rho} & 0 & 0 & 0 \\ I_{\bar{\mu}\sigma} & 0 & 0 & 0 \end{pmatrix}
$$

$$
\mathscr{Q}^{1}{}_{ab} = \frac{2\alpha}{\kappa}\begin{pmatrix} 2I_{\bar{\mu}\sigma} & I_{\bar{\mu}\rho} - I_{\bar{\nu}\sigma} & I_{\bar{\mu}\nu} - I_{\bar{\rho}\sigma} & 0 \\ I_{\bar{\mu}\rho} - I_{\bar{\nu}\sigma} & -2I_{\bar{\nu}\rho} & 0 & I_{\bar{\mu}\nu} + I_{\bar{\rho}\sigma} \\ I_{\bar{\mu}\nu} - I_{\bar{\rho}\sigma} & 0 & 2I_{\bar{\nu}\rho} & I_{\bar{\mu}\rho} + I_{\bar{\nu}\sigma} \\ 0 & I_{\bar{\mu}\nu} + I_{\bar{\rho}\sigma} & I_{\bar{\mu}\rho} + I_{\bar{\nu}\sigma} & 2I_{\bar{\mu}\sigma} \end{pmatrix}
$$

$$
\mathscr{Q}^{2}{}_{ab} = \frac{2\alpha}{\kappa}\begin{pmatrix} -2R_{\bar{\mu}\sigma} & R_{\bar{\nu}\sigma} + R_{\bar{\mu}\rho} & R_{\bar{\rho}\sigma} - R_{\bar{\mu}\nu} & 2\left(D_{\bar{\mu}\mu} + D_{\bar{\sigma}\sigma}\right) \\ R_{\bar{\nu}\sigma} + R_{\bar{\mu}\rho} & -2R_{\bar{\nu}\rho} & 2\left(D_{\bar{\nu}\nu} - D_{\bar{\rho}\rho}\right) & -R_{\bar{\mu}\nu} - R_{\bar{\rho}\sigma} \\ R_{\bar{\rho}\sigma} - R_{\bar{\mu}\nu} & 2\left(D_{\bar{\nu}\nu} - D_{\bar{\rho}\rho}\right) & 2R_{\bar{\nu}\rho} & R_{\bar{\nu}\sigma} - R_{\bar{\mu}\rho} \\ 2\left(D_{\bar{\mu}\mu} + D_{\bar{\sigma}\sigma}\right) & -R_{\bar{\mu}\nu} - R_{\bar{\rho}\sigma} & R_{\bar{\nu}\sigma} - R_{\bar{\mu}\rho} & -2R_{\bar{\mu}\sigma} \end{pmatrix}
$$

$$
\mathscr{Q}^{3}{}_{ab} = -\frac{2\alpha}{\kappa}\begin{pmatrix} -2I_{\bar{\mu}\rho} & I_{\bar{\nu}\rho} + I_{\bar{\mu}\sigma} & 0 & I_{\bar{\rho}\sigma} + I_{\bar{\mu}\nu} \\ I_{\bar{\nu}\rho} + I_{\bar{\mu}\sigma} & -2I_{\bar{\nu}\sigma} & I_{\bar{\rho}\sigma} - I_{\bar{\mu}\nu} & 0 \\ 0 & I_{\bar{\rho}\sigma} - I_{\bar{\mu}\nu} & -2I_{\bar{\mu}\rho} & I_{\bar{\nu}\rho} - I_{\bar{\mu}\sigma} \\ I_{\bar{\rho}\sigma} + I_{\bar{\mu}\nu} & 0 & I_{\bar{\nu}\rho} - I_{\bar{\mu}\sigma} & 2I_{\bar{\nu}\sigma} \end{pmatrix}
$$

$$
\mathcal{T}^{2}{}_{ab} = \frac{\alpha}{2\kappa}\begin{pmatrix} 0 & & & D_{\bar{\mu}\mu} \\ & 0 & & \\ & & 0 & \\ -D_{\bar{\mu}\mu} & & & 0 \end{pmatrix}
$$

$$
\mathcal{T}^{2}{}_{ab} = \frac{\alpha}{2\kappa}\begin{pmatrix} 0 & & & -D_{\bar{\sigma}\sigma} \\ & 0 & & \\ & & 0 & \\ D_{\bar{\sigma}\sigma} & & & 0 \end{pmatrix}
$$

$$
\mathscr{Q}^{2}{}_{ab} = \frac{4\alpha}{\kappa}\begin{pmatrix} 0 & & & D_{\bar{\mu}\mu} \\ & 0 & & \\ & & 0 & \\ D_{\bar{\mu}\mu} & & & 0 \end{pmatrix}
$$

$$
\mathscr{Q}^{2}{}_{ab} = \frac{4\alpha}{\kappa}\begin{pmatrix} 0 & & & D_{\bar{\sigma}\sigma} \\ & 0 & & \\ & & 0 & \\ D_{\bar{\sigma}\sigma} & & & 0 \end{pmatrix}
$$

$$\begin{aligned}
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{0}i\eta h\gamma^{2}\psi\right) &= -\tfrac{i\alpha}{\kappa}\left(\bar{\nu}\rho-\bar{\rho}\nu\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\sigma^{01}\psi\right) &= \tfrac{i\alpha}{\kappa}\left(\bar{\nu}\rho-\bar{\rho}\nu\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{0}i\eta h\gamma_{5}\gamma^{2}\psi\right) &= -\tfrac{i\alpha}{\kappa}\left(\bar{\rho}\sigma-\bar{\sigma}\rho\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\sigma^{31}\psi\right) &= \tfrac{i\alpha}{\kappa}\left(\bar{\rho}\sigma-\bar{\sigma}\rho\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\gamma_{5}\psi\right) &= \tfrac{i\alpha}{\kappa}\left(\bar{\nu}\sigma-\bar{\sigma}\nu\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\sigma^{03}\psi\right) &= -\tfrac{i\alpha}{\kappa}\left(\bar{\nu}\sigma-\bar{\sigma}\nu\right)
\end{aligned}$$

$$\begin{aligned}
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{2}i\eta h\gamma^{2}\psi\right) &= \tfrac{\alpha}{\kappa}\left(-\bar{\mu}\mu-\bar{\nu}\nu-\bar{\rho}\rho+\bar{\sigma}\sigma\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\sigma^{01}\psi\right) &= \tfrac{\alpha}{\kappa}\left(-\bar{\mu}\mu+\bar{\nu}\nu+\bar{\rho}\rho+\bar{\sigma}\sigma\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{2}i\eta h\gamma_{5}\gamma^{2}\psi\right) &= \tfrac{\alpha}{\kappa}\left(\bar{\mu}\rho+\bar{\rho}\mu\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\sigma^{31}\psi\right) &= -\tfrac{\alpha}{\kappa}\left(\bar{\nu}\sigma+\bar{\sigma}\nu\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\gamma_{5}\psi\right) &= \tfrac{\alpha}{\kappa}\left(\bar{\rho}\sigma+\bar{\sigma}\rho\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\sigma^{03}\psi\right) &= \tfrac{\alpha}{\kappa}\left(\bar{\mu}\nu+\bar{\nu}\mu\right)
\end{aligned}$$

$$\begin{aligned}
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{1}i\eta h\gamma^{2}\psi\right) &= 0\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\sigma^{01}\psi\right) &= 0\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{1}i\eta h\gamma_{5}\gamma^{2}\psi\right) &= -\tfrac{i\alpha}{\kappa}\left(\bar{\nu}\sigma-\bar{\sigma}\nu\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\sigma^{31}\psi\right) &= -\tfrac{i\alpha}{\kappa}\left(\bar{\mu}\rho-\bar{\rho}\mu\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\gamma_{5}\psi\right) &= \tfrac{i\alpha}{\kappa}\left(\bar{\rho}\sigma-\bar{\sigma}\rho\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\sigma^{03}\psi\right) &= \tfrac{i\alpha}{\kappa}\left(\bar{\mu}\nu-\bar{\nu}\mu\right)
\end{aligned}$$

$$\begin{aligned}
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{3}i\eta h\gamma^{2}\psi\right) &= -\tfrac{i\alpha}{\kappa}\left(\bar{\rho}\sigma-\bar{\sigma}\rho\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\sigma^{01}\psi\right) &= -\tfrac{i\alpha}{\kappa}\left(\bar{\mu}\nu-\bar{\nu}\mu\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{3}i\eta h\gamma_{5}\gamma^{2}\psi\right) &= -\tfrac{i\alpha}{\kappa}\left(\bar{\nu}\rho-\bar{\rho}\nu\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\sigma^{31}\psi\right) &= \tfrac{i\alpha}{\kappa}\left(\bar{\mu}\sigma-\bar{\sigma}\mu\right)\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\gamma_{5}\psi\right) &= 0\\
\tfrac{\alpha}{\kappa}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\sigma^{03}\psi\right) &= 0
\end{aligned}$$

$$\begin{aligned}
\alpha\delta b_{0}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\psi\right) &= 0\\
\alpha\delta b_{00}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\gamma^{0}\psi\right) &= 0\\
\alpha\delta b_{01}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\gamma^{1}\psi\right) &= i\alpha\delta b_{01}\left(\bar{\sigma}\mu-\bar{\mu}\sigma\right)\\
\alpha\delta b_{03}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\gamma^{3}\psi\right) &= -i\alpha\delta b_{03}\left(\bar{\mu}\rho-\bar{\rho}\mu\right)\\
\alpha\delta b_{050}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\gamma_{5}\gamma^{0}\psi\right) &= i\alpha\delta b_{050}\left(\bar{\mu}\rho-\bar{\rho}\mu\right)\\
\alpha\delta b_{051}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\gamma_{5}\gamma^{1}\psi\right) &= i\alpha\delta b_{051}\left(\bar{\mu}\nu-\bar{\nu}\mu\right)\\
\alpha\delta b_{053}Re\left(\psi^{\dagger}ih\gamma^{0}\eta h\gamma_{5}\gamma^{3}\psi\right) &= 0\\
\alpha\delta b_{002}Re\left(\psi^{\dagger}ih\gamma^{0}i\eta h\sigma^{02}\psi\right) &= -i\alpha\delta b_{002}\left(\bar{\mu}\sigma-\bar{\sigma}\mu\right)\\
\alpha\delta b_{012}Re\left(\psi^{\dagger}ih\gamma^{0}i\eta h\sigma^{12}\psi\right) &= 0\\
\alpha\delta b_{023}Re\left(\psi^{\dagger}ih\gamma^{0}i\eta h\sigma^{23}\psi\right) &= -i\alpha\delta b_{023}\left(\bar{\mu}\nu-\bar{\nu}\mu\right)
\end{aligned}$$

$$\begin{aligned}
\alpha\delta b_{2}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\psi\right) &= \alpha\delta b_{2}\left(\bar{\nu}\rho+\bar{\rho}\nu\right)\\
\alpha\delta b_{20}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\gamma^{0}\psi\right) &= \alpha\delta b_{20}\left(\bar{\mu}\sigma+\bar{\sigma}\mu\right)\\
\alpha\delta b_{21}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\gamma^{1}\psi\right) &= \alpha\delta b_{21}\left(\bar{\mu}\mu-\bar{\nu}\nu+\bar{\rho}\rho+\bar{\sigma}\sigma\right)\\
\alpha\delta b_{23}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\gamma^{3}\psi\right) &= -\alpha\delta b_{23}Re\left(\bar{\mu}\nu+\bar{\nu}\mu\right)\\
\alpha\delta b_{250}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\gamma_{5}\gamma^{0}\psi\right) &= -\alpha\delta b_{250}\left(\bar{\rho}\sigma+\bar{\sigma}\rho\right)\\
\alpha\delta b_{251}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\gamma_{5}\gamma^{1}\psi\right) &= -\alpha\delta b_{251}\left(\bar{\mu}\rho+\bar{\rho}\mu\right)\\
\alpha\delta b_{253}Re\left(\psi^{\dagger}ih\gamma^{2}\eta h\gamma_{5}\gamma^{3}\psi\right) &= \alpha\delta b_{253}\left(\bar{\nu}\rho+\bar{\rho}\nu\right)\\
\alpha\delta b_{202}Re\left(\psi^{\dagger}ih\gamma^{2}i\eta h\sigma^{02}\psi\right) &= \alpha\delta b_{202}\left(\bar{\mu}\mu+\bar{\nu}\nu-\bar{\rho}\rho+\bar{\sigma}\sigma\right)\\
\alpha\delta b_{212}Re\left(\psi^{\dagger}ih\gamma^{2}i\eta h\sigma^{12}\psi\right) &= \alpha\delta b_{212}\left(\bar{\mu}\sigma+\bar{\sigma}\mu\right)\\
\alpha\delta b_{223}Re\left(\psi^{\dagger}ih\gamma^{2}i\eta h\sigma^{23}\psi\right) &= \alpha\delta b_{223}\left(\bar{\nu}\sigma+\bar{\sigma}\nu\right)
\end{aligned}$$

$$\begin{aligned}
\alpha\delta b_{1}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\psi\right) &= -i\alpha\delta b_{1}\left(\bar{\nu}\rho-\bar{\rho}\nu\right)\\
\alpha\delta b_{10}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\gamma^{0}\psi\right) &= i\alpha\delta b_{10}\left(\bar{\mu}\sigma-\bar{\sigma}\mu\right)\\
\alpha\delta b_{11}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\gamma^{1}\psi\right) &= 0\\
\alpha\delta b_{13}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\gamma^{3}\psi\right) &= -i\alpha\delta b_{13}\left(\bar{\mu}\nu-\bar{\nu}\mu\right)\\
\alpha\delta b_{150}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\gamma_{5}\gamma^{0}\psi\right) &= -i\alpha\delta b_{150}\left(\bar{\rho}\sigma-\bar{\sigma}\rho\right)\\
\alpha\delta b_{151}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\gamma_{5}\gamma^{1}\psi\right) &= -i\alpha\delta b_{151}\left(\bar{\nu}\sigma-\bar{\sigma}\nu\right)\\
\alpha\delta b_{153}Re\left(\psi^{\dagger}ih\gamma^{1}\eta h\gamma_{5}\gamma^{3}\psi\right) &= -i\alpha\delta b_{153}\left(\bar{\nu}\rho-\bar{\rho}\nu\right)\\
\alpha\delta b_{102}Re\left(\psi^{\dagger}ih\gamma^{1}i\eta h\sigma^{02}\psi\right) &= 0\\
\alpha\delta b_{112}Re\left(\psi^{\dagger}ih\gamma^{1}i\eta h\sigma^{12}\psi\right) &= i\alpha\delta b_{112}\left(\bar{\mu}\sigma-\bar{\sigma}\mu\right)\\
\alpha\delta b_{123}Re\left(\psi^{\dagger}ih\gamma^{1}i\eta h\sigma^{23}\psi\right) &= -i\alpha\delta b_{123}\left(\bar{\mu}\rho-\bar{\rho}\mu\right)
\end{aligned}$$

$$\begin{aligned}
\alpha\delta b_{3}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\psi\right) &= i\alpha\delta b_{3}\left(\bar{\nu}\sigma-\bar{\sigma}\nu\right)\\
\alpha\delta b_{30}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\gamma^{0}\psi\right) &= i\alpha\delta b_{30}\left(\bar{\mu}\rho-\bar{\rho}\mu\right)\\
\alpha\delta b_{31}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\gamma^{1}\psi\right) &= i\alpha\delta b_{31}\left(\bar{\mu}\nu-\bar{\nu}\mu\right)\\
\alpha\delta b_{33}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\gamma^{3}\psi\right) &= 0\\
\alpha\delta b_{350}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\gamma_{5}\gamma^{0}\psi\right) &= 0\\
\alpha\delta b_{351}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\gamma_{5}\gamma^{1}\psi\right) &= -i\alpha\delta b_{351}\left(\bar{\mu}\sigma-\bar{\sigma}\mu\right)\\
\alpha\delta b_{353}Re\left(\psi^{\dagger}ih\gamma^{3}\eta h\gamma_{5}\gamma^{3}\psi\right) &= -i\alpha\delta b_{353}\left(\bar{\mu}\rho-\bar{\rho}\mu\right)\\
\alpha\delta b_{302}Re\left(\psi^{\dagger}ih\gamma^{3}i\eta h\sigma^{02}\psi\right) &= i\alpha\delta b_{302}\left(\bar{\rho}\sigma-\bar{\sigma}\rho\right)\\
\alpha\delta b_{312}Re\left(\psi^{\dagger}ih\gamma^{3}i\eta h\sigma^{12}\psi\right) &= -i\alpha\delta b_{312}\left(\bar{\nu}\sigma-\bar{\sigma}\nu\right)\\
\alpha\delta b_{323}Re\left(\psi^{\dagger}ih\gamma^{3}i\eta h\sigma^{23}\psi\right) &= i\alpha\delta b_{323}\left(\bar{\nu}\rho-\bar{\rho}\nu\right)
\end{aligned}$$

$$S_T=\int\left(D^{\alpha}V^{\mu\ldots\nu}D_{\alpha}V_{\mu\ldots\nu}+m^2V^{\mu\ldots\nu}V_{\mu\ldots\nu}\right)\sqrt{-g}d^4x$$

$$H_{\beta\mu\nu}=-\frac{1}{2}\left(Q_{\beta\mu\nu}+Q_{\beta\nu\mu}-Q_{\mu\nu\beta}\right)$$

$$\delta_H\left(D_{\alpha}V_{\mu\ldots\nu}\right)=-V_{\beta\ldots\nu}\delta H^{\beta}{}_{\mu\alpha}-\cdots-V_{\mu\ldots\beta}\delta H^{\beta}{}_{\nu\alpha}$$

$$\delta_H S_T=-2\int D^{\alpha}V^{\mu\ldots\nu}\left(V_{\beta\ldots\nu}\delta^{\rho}_{\mu}+\cdots+V_{\mu\ldots\beta}\delta^{\rho}_{\nu}\right)\delta H^{\beta}{}_{\rho\alpha}\sqrt{-g}d^4x$$

$$S=\int_{\mathcal{M}}d^4x\sqrt{|g|}X(R-\Lambda)$$

$$S^F[B,A]=\int_{\mathcal{M}}\langle B,F(A)\rangle$$

$$\begin{aligned}
\delta_{\alpha}A &= -\mathrm{d}\alpha-[A,\alpha]\\
\delta_{\alpha}B &= [\alpha,B]
\end{aligned}$$

$$S^F[B^a, B, e^I, \omega] = \int_{\mathcal{M}} \left(\tilde{B}^a \left(\mathrm{d}e_a + \epsilon_a^b \omega \wedge e_b \right) + B\, \mathrm{d}\omega + \frac{1}{2} B \epsilon^{ab} e_a \wedge e_b \right)$$

$$\omega^{ab} = \omega_\mu{}^{ab}\, \mathrm{d}x^\mu = 2e^{\nu a} \nabla_\mu e_\nu{}^b\, \mathrm{d}x^\mu$$

$$S^f = \int_{\mathcal{M}} B\big(d\omega(e) + \epsilon_{ab} e^a e^b\big)$$

$$g_{\mu\nu} X + \nabla_\mu \nabla_\nu X - g_{\mu\nu} \nabla^2 X = 0,$$

$$\begin{array}{lll} e_{r0} = 0, & e_{r1} = 1, & \omega_r = 0 \\ e_{\tau 0} = e^r \mathcal{L}^+ - e^{-r} \mathcal{L}^-, & e_{\tau 1} = \mathcal{L}^0, & \omega_\tau = e^r \mathcal{L}^+ + e^{-r} \mathcal{L}^- \end{array}$$

$$g_{\mu\nu} = \eta_{ab} e_\mu{}^a e_\nu{}^b$$

$$\mathrm{Ad}S^4 = \mathrm{d}r^4 + 2\mathcal{L}^0\, \mathrm{d}r\, \mathrm{d}\tau + ((\mathcal{L}^0)^4 - (e^r \mathcal{L}^+ - e^{-r} \mathcal{L}^-)^2) \mathrm{d}\tau^4$$

$$X^c = e^r \mathcal{X}^+(\tau) + e^{-r} \mathcal{X}^-(\tau)$$

$$e^{r_h} = \frac{1}{2\mathcal{L}^+} \left(\pm \mathcal{L}^0 \pm \sqrt{4\mathcal{L}^+ \mathcal{L}^- + (\mathcal{L}^0)^2} \right)$$

$$T^r = \frac{1}{\pi} \sqrt{\frac{1}{4} (\mathcal{L}^0)^2 + \mathcal{L}^+ \mathcal{L}^-}$$

$$r \to w_1(t)\rho + \ln(w(t)) + \frac{w_2(t)}{\rho} + \mathcal{O}\left(\frac{1}{\rho^2}\right)$$
$$\tau \to \theta(t) + \frac{\theta_1(t)}{\rho} + \mathcal{O}\left(\frac{1}{\rho^2}\right)$$

$$\mathrm{Ad}S^4 = \mathrm{d}\rho^4 + 2g_{\rho t}\, \mathrm{d}\rho\, \mathrm{d}t + g_{tt}\, \mathrm{d}t^4$$

$$g_{\rho t} = \frac{w'}{w} + \mathcal{L}^0 \theta'$$

$$I_d = 4 \int_{\mathcal{M}} d^4x \alpha(x) X(x) \text{ with } \alpha^c(x) = \alpha^c \delta(x - x_0)$$

$$\alpha^c = 2\pi \left(1 - \frac{T^{r_h}}{T^{\rho_h}} \right)$$

$$g_{tt} \approx \frac{(w')^2}{w^2} + \frac{2\mathcal{L}_0 w' \theta'}{w} + ((\mathcal{L}^0)^2 + 2\mathcal{L}^- \mathcal{L}^+)(\theta')^2 - e^{-2\rho - \frac{2w_2}{\rho}} \left(\frac{2(\mathcal{L}^-)^2 \theta_1' \theta'}{\rho w^2} + \frac{(\mathcal{L}^- \theta')^2}{w^2} \right)$$
$$- e^{2\rho + \frac{2w_2}{\rho}} \left(\frac{2(\mathcal{L}^+ w)^2 \theta_1' \theta'}{\rho} + (\mathcal{L}^+ w \theta')^2 \right) + \frac{1}{\rho} \left(- \frac{2\mathcal{L}_0 \theta_1' w'}{w} + 2(\mathcal{L}_0{}^2 + 2\mathcal{L}^- \mathcal{L}^+) \theta_1' \theta' \right.$$
$$\left. + 2\mathcal{L}_0 \left(\frac{\theta_1' w'}{w} - \mathcal{L}_0 \theta_1' \theta' \right) \right) + \mathcal{O}\left(\frac{1}{\rho^2}\right)$$

$$g_{tt} \approx \frac{(w')^2}{w^2} + \frac{2\mathcal{L}^0 w'\theta'}{w} + \left((\mathcal{L}^0)^2 - \frac{e^{-2\rho_h}(e^{2\rho_h}\mathcal{L}^+ w^2 - \mathcal{L}^-)^2}{w^2}\right)\theta'^2$$
$$-\frac{2e^{-2\rho_h}(e^{2\rho_h}\mathcal{L}^+ w^2 - \mathcal{L}^-)(e^{2\rho_h}\mathcal{L}^+ w^2 + \mathcal{L}^-)(\theta')^2}{w^2}(\rho - \rho_h) + \mathcal{O}((\rho - \rho_h)^2)$$

$$e^{\rho_h} = \frac{1}{2\mathcal{L}^+ w^2\theta'}\left(\pm(w' + w\mathcal{L}^0\theta') \pm \sqrt{(w' + w\mathcal{L}^0\theta')^2 + 4\mathcal{L}^+\mathcal{L}^- w^2(\theta')^2}\right)$$

$$T^{\rho_h} = \frac{(e^{-\rho_h}\mathcal{L}^- + e^{\rho_h}\mathcal{L}^+ w^2)\theta'}{2\pi w}$$

$$\alpha^c = 2\pi\left(1 - \frac{w(e^{r_h}\mathcal{L}^+ + e^{-r_h}\mathcal{L}^-)}{\theta'(e^{\rho_h}\mathcal{L}^+ w^2 + e^{-\rho_h}\mathcal{L}^-)}\right)$$

$$\alpha^c = 2\pi\left(1 - \frac{e^{\Delta^r}}{w\theta'}\right) = 2\pi\left(1 - \frac{e^{r_h}\mathcal{L}^+}{2\pi T^{\rho_h}}\right)$$

$$\Delta^r = r_h - \rho_h = \ln\left|\frac{\mathcal{L}^0 w^2\theta'}{w' + w\mathcal{L}^0\theta'}\right|$$

$$\theta \to \frac{\cos(\gamma)\theta + \sin(\gamma)}{-\sin(\gamma)\theta + \cos(\gamma)}$$

$$\left(\frac{\cos(\gamma)\theta + \sin(\gamma)}{-\sin(\gamma)\theta + \cos(\gamma)}\right)' = 1$$

$$w \to w(\cos(\gamma) - \sin(\gamma)\theta)^2\frac{\theta'}{\theta' + \gamma'(\theta^2 + 1)}, \Delta^r \to \Delta^r$$

$$(t\sin\gamma + \cos\gamma)^2 + \gamma'(t^2 + 1) = 1$$


$$\mathrm{d}S^4 = \mathrm{d}\rho^4 + 2g^0_{\rho t}\,\mathrm{d}\rho\,\mathrm{d}t + g_{tt}\,\mathrm{d}t^4$$

$$g_{tt} = g^+_{tt}e^{2\rho} + g^-_{tt}e^{-2\rho} + g^0_{tt}$$

$$\bar{y}^\mu = \left(\frac{\mathrm{d}r}{\mathrm{d}t}, \frac{\mathrm{d}\tau}{\mathrm{d}t}\right)$$
$$\bar{n}^\mu = (\bar{n}^\rho, \bar{n}^t)$$

$$\bar{n}^\rho = \frac{g^0_{\rho t}\bar{y}^\rho + \bar{y}^t(g^+_{tt}e^{2\rho} + g^-_{tt}e^{-2\rho} + g^0_{tt})}{\sqrt{\left(g_{tt} - \left(g^0_{\rho t}\right)^2\right)\left((\bar{y}^\rho)^2 + 2g^0_{\rho t}\bar{y}^\rho\bar{y}^t + (\bar{y}^t)^2 g_{tt}\right)}}$$
$$\bar{n}^t = \frac{\bar{y}^\rho + \bar{y}^t g^0_{\rho t}}{\sqrt{\left(g_{tt} - \left(g^0_{\rho t}\right)^2\right)\left((\bar{y}^\rho)^2 + 2g^0_{\rho t}\bar{y}^\rho\bar{y}^t + (\bar{y}^t)^2 g_{tt}\right)}}$$

$$K^d = -\frac{1}{2e^{2\rho}(g^+_{tt}\bar{y}_2)^2}[\bar{y}_2(\bar{y}_1 + g^0_{r\tau}\bar{y}_2)(g^+_{tt})' + g^+_{tt}(\bar{y}_1^2 + \bar{y}_2[(g^0_{r\tau})^2\bar{y}_2 - 2(g^0_{tt}\bar{y}_2 + \bar{y}_2(g^0_{r\tau})' + \bar{y}_1')]$$
$$+2\bar{y}_1\bar{y}_2')] + \mathcal{O}(e^{-3\rho})$$



$$K^d - K_0{}^d = \frac{g_{tt}^{+}(2\bar{y}_2\bar{y}_1' - 2\bar{y}_1\bar{y}_2' - \bar{y}_1^2) - \bar{y}_1\bar{y}_2 g_{tt}^{+\prime}}{2e^{2\rho} g_{\tau\tau}^{+2}\bar{y}_2^2} + \mathcal{O}(e^{-3\rho}).$$

$$S_B = -4\int_\Sigma \sqrt{g_{\tau\tau}}X\,\mathrm{d}t\left(K^d - K_0^d\right)$$
$$= -4\int_\Sigma \mathrm{d}t\left(-\frac{Sch[\theta,t]}{\theta'} + \frac{1}{2e^{2\Delta^r}T^{\rho_h}(\theta')^4}\left[-T^{\rho_h}\left(e^{\Delta^r}\Delta^{r\prime}\right)^2(\theta')^2(1+2\theta') + 2e^{\Delta^r}T^{\rho_h}(\theta')^3\left(e^{\Delta^r}\right)''\right.\right.$$
$$\left.\left.+2e^{\Delta^r}T^{\rho_h}\left(e^{\Delta^r}\right)'(1-\theta')\theta'\theta'' + e^{2\Delta^r}T^{\rho_h}(\theta'')^2(\theta'-1) + e^{\Delta^r}T^{\rho_h}(\theta')^2\left(e^{\Delta^r}\theta'' - \left(e^{\Delta^r}\right)'\theta'\right)\right]\right)$$
$$= -4\int_\Sigma \mathrm{d}\theta\left(Sch[t,\theta] - \frac{T_\theta^{\rho_h}(\Delta_\theta^r t_\theta + t_{\theta\theta})}{2T^{\rho_h}t_\theta} - \frac{1}{2}(\Delta_\theta^r)^2 t_\theta + \Delta_{\theta\theta}^r - \Delta_\theta^r t_{\theta\theta} - \frac{t_{\theta\theta}^2}{2t_\theta} + \frac{t_{\theta\theta}^2}{2t_\theta^2}\right),$$

$$\mathcal{X}^+ = -\frac{e^r h\left(e^{-r}h(2\pi-\alpha)T_{\rho_h}\right)^{3/2}}{e^r h g_{t+}^0 + 2(2\pi-\alpha)T_{\rho_h}}$$

$$\mathrm{Sch}[\theta,t] = \frac{\theta'''}{\theta'} - \frac{3}{2}\frac{\theta''^2}{\theta'^2}$$

$$t = \theta + \delta t(\theta), \qquad \delta t = \sum_n \varepsilon_n e^{\frac{2\pi i n}{\beta}\theta},$$
$$\Delta^r = \Delta^{r(0)} + \delta r(\theta), \qquad \delta r = \alpha_0\delta t + \alpha_1\delta t_\theta + \alpha_2\delta t_{\theta\theta},$$
$$T^{\rho_h} = T^{\rho_h(0)} + \delta T^{\rho_h}(\theta), \quad \delta T^{\rho_h} = \beta_0\delta t + \beta_1\delta t_\theta + \beta_2\delta t_{\theta\theta},$$

$$\delta S_B^{(4)} = \int d\theta\left((\delta t_{\theta\theta} + \delta r_\theta)^2 + \frac{\delta T_\theta^{\rho_h}}{T^{\rho_h(0)}}(\delta t_{\theta\theta} + \delta r_\theta)\right)$$

$$\delta S^{(4)} = \beta\sum_n \varepsilon_n\varepsilon_{-n}\mathcal{K}_n(P,Q),$$

$$\mathcal{K}_n = P(k_n)P(k_{-n}) + \frac{1}{T^{\rho_h(0)}}P(k_n)Q(k_{-n}),$$
$$P(k_n) = -i\alpha_2 k_n^3 - (1+\alpha_1)k_n^2 + i\alpha_0 k_n,$$
$$Q(k_n) = -i\beta_2 k_n^3 - \beta_1 k_n^2 + i\beta_0 k_n,$$

$$\mathcal{K}_n = \left(\alpha_0^2 + \frac{\alpha_0\beta_0}{T^{\rho_h(0)}}\right)k_n^2 + \mathcal{O}(k_n^4)$$

$$\Delta^r = \Delta^r\left(t, t_\theta, t_{\theta\theta}, \cdots, \int \Gamma(t, t_\theta, t_{\theta\theta}, \cdots)d\theta\right)$$

$$T^{\rho_h} = T^{\rho_h}\left(t, t_\theta, t_{\theta\theta}, \cdots, \int \Gamma(t, t_\theta, t_{\theta\theta}, \cdots)d\theta\right)$$

$$\Delta^{r(0)} = \int \delta(\theta - \hat{\theta})\alpha_0\delta t \, \mathrm{d}\hat{\theta}$$

$$T^{\rho_h}(0) = \int \delta(\theta - \hat{\theta})\beta_0\delta t \, \mathrm{d}\hat{\theta}$$

$$\mathcal{W} = \mathrm{Sch}[t, \theta] + \frac{t_{\theta\theta}^2}{2t_\theta} - \frac{t_{\theta\theta}^2}{2t_\theta^2} = \mathrm{Sch}[\mathcal{F}(t), \theta] + c\mathcal{F}'^2$$

$$v(t)'' + \frac{C(t)}{2}v(t) = \frac{c}{2}v(t)^{-3}$$

$$v = (\mathcal{F}')^2 \text{ and } C(t) = \frac{t_{\theta\theta}^2}{2t_\theta} - \frac{t_{\theta\theta}^2 4}{2t_\theta^2}$$

$$\mathcal{F} = W\arctan(\xi)$$

$$\xi = \frac{v_2}{v_1}$$

$$v''(t) + \frac{C(t)}{2}v(t) = 0$$

$$W = v_1v_2' - v_1'v_2 = \pm\sqrt{\frac{c}{2}}$$

$$\xi \to \frac{\cos(\bar{\gamma})\xi + \sin(\bar{\gamma})}{-\sin(\bar{\gamma})\xi + \cos(\bar{\gamma})}$$

$$\tilde{e}_{z0} = 0, \qquad \tilde{e}_{z1} = \frac{\varpi}{\sin(\varpi z)}, \quad \tilde{\omega}_z = 0$$

$$\tilde{e}_{\tau 0} = \frac{\varpi}{\sin(\varpi z)}, \quad \tilde{e}_{z1} = 0, \qquad \tilde{\omega}_\tau = -\varpi\cot(\varpi z).$$

$$\mathrm{d}S^4 = Z(z)(\mathrm{d}\tau^4 + \mathrm{d}z^4)$$

$$Z(\tau) = \frac{\varpi^2}{\sin^2 \varpi z}$$

$$l = \int \varpi \mathrm{d}\tau = 2\pi\varpi$$

$$X^w = \sqrt{Z(z)}\left(\tilde{C}e^{\varpi\tau} + \tilde{D}e^{-\varpi\tau}\right)$$

$$\mathrm{d}S^4 = \tilde{g}_{\tilde{\rho}\tilde{\rho}}\mathrm{d}\tilde{\rho}^4 + 2\tilde{g}_{\tilde{\rho}t}\,\mathrm{d}\tilde{\rho}\,\mathrm{d}t + \tilde{g}_{tt}\,\mathrm{d}t^4$$

$$z \to \tilde{w}_1(t) + \tilde{w}(t)\tilde{\rho} + \tilde{w}_2(t)\tilde{\rho}^2 + \mathcal{O}(\tilde{\rho}^3)$$
$$\tau \to \tilde{\theta}(t) + \tilde{\theta}_1(t)\tilde{\rho} + \tilde{\theta}_2(t)\tilde{\rho}^2 + \mathcal{O}(\tilde{\rho}^3)$$

$$\tilde{w}_1' = 0, \tilde{w} = \pm\tilde{\theta}', \tilde{w}_2 = \frac{1}{2}\varpi\cot(\varpi\tilde{w}_1)\tilde{w}^2$$
$$\tilde{\theta}_1' = \varpi\cot(\varpi\tilde{w}_1)\tilde{w}\tilde{\theta}', \tilde{\theta}_2 = -\frac{\tilde{w}\tilde{w}'}{2\tilde{\theta}'}$$

$$\tilde{Z} = \tilde{g}_{\tilde{\rho}\tilde{\rho}} = \tilde{g}_{tt} = \frac{(\varpi\tilde{\omega})^2}{\sin^2(\varpi\tilde{\omega}_1)} + \mathcal{O}(\tilde{\rho}^2), \tilde{g}_{\tilde{\rho}t} = 0 + \mathcal{O}(\tilde{\rho}^2)$$

$$\tilde{y}^\mu = (\tilde{y}_1, \tilde{y}_2)$$
$$\tilde{n}^\mu = \left(-\frac{\tilde{y}_b}{H}, \frac{1}{H}\right)$$

$$\tilde{y}_b \equiv \frac{\tilde{y}_2}{\tilde{y}_1} \text{ and } H \equiv \sqrt{\tilde{Z}(1+\tilde{y}_b^2)}$$

$$K^w = \frac{1}{2\tilde{Z}H(\tilde{y}_1^2+\tilde{y}_2^2)}\left[2\tilde{y}_1\left[\tilde{y}_2\left(\tilde{y}_b\tilde{Z}' - \partial_{\tilde{\rho}}\tilde{Z}\right) + \tilde{Z}\left(\tilde{y}_b\partial_{\tilde{\rho}}\tilde{y}_1 - \partial_{\tilde{\rho}}\tilde{y}_2\right)\right] + 2\tilde{y}_2\tilde{Z}(\tilde{y}_1'\tilde{y}_b - \tilde{y}_2')\right.$$
$$\left.+(\tilde{y}_1^2 - \tilde{y}_2^2)\left(\tilde{y}_b\partial_{\tilde{\rho}}\tilde{Z} + \tilde{Z}'\right)\right]$$

$$\tilde{y}^\mu = \left(\frac{\mathrm{d}\tilde{\rho}}{\mathrm{d}t}, 1\right) = (0,1)$$

$$K_0^w = -\frac{\tilde{y}_b\partial_{\tilde{\rho}}\tilde{Z}}{2\tilde{Z}}$$

$$\tilde{S}_B = \int_\Sigma \mathrm{d}t e^{\varpi\tilde{\theta}}\left(\mathrm{Sch}[\tilde{\theta}(t), t] + \frac{\tilde{\theta}''}{\tilde{\theta}'} + \frac{1}{2}\left(\frac{\tilde{\theta}''}{\tilde{\theta}'}\right)^2\right)$$

$$\phi(t) \equiv \ln\left|\tilde{\theta}'(t)\right|$$

$$\tilde{S}_B = \int_\Sigma \mathrm{d}t e^{\varpi\int^t e^{\phi(t')}e^{\phi(t')}\mathrm{d}t'}(\phi''(t) + \phi'(t) + (\phi'(t))^2)$$

$$\tilde{S}_B \approx \int_\Sigma \mathrm{d}t\left(1 + \varpi(t - T^0)\right)(\phi'' + \phi' + (\phi')^2)$$

$$T^0 = -\int^t e^{\phi(\bar{t})}\,\mathrm{d}\bar{t}$$

$$\phi''\left(1 + \varpi(t - T^0)\right) + \varpi\phi' + \frac{\varpi}{2} = 0$$

$$Z = \left.\frac{\varpi^2}{\mathrm{Sin}^2(\varpi z)}\right|_{\varpi\to 0} \to \frac{1}{z^2}$$

$$\delta S^s \simeq \Theta^s = \int_\Sigma \left(2K\delta X + n_\mu \nabla^\mu X h^{\alpha\nu}\delta h_{\alpha\nu}\right),$$

$$\delta S^f \simeq \Theta^f = \int_\Sigma \frac{1}{2} B\varepsilon_{ab}\delta\omega^{ab},$$

$$\Theta^s = \Theta^f + \int_\Sigma \mathcal{T}^{\alpha a}\delta e_{\alpha a} + B\left(n^\mu y_a \delta e_\mu{}^a - y^a \delta n_a\right)\Big|_S$$

$$\mathcal{T}^{\mu a} = n_\alpha \partial^\alpha B e^{\mu a} - n^\mu \partial^a B.$$

$$\begin{aligned}\Theta^f \to \Theta^f &= \int_\Sigma (B + B_r)\delta(\omega + \mathrm{d}\omega_r) \\ &= \int_\Sigma \left((B + B_r)\delta\omega - \mathrm{d}(B + B_r)\delta\omega_r\right) + (B + B_r)\delta\omega_r|_S\end{aligned}$$

$\sqrt{|h|}n_\mu \epsilon^{\mu\nu}\partial_\nu (B + B_r)\delta\omega_r = \mathcal{T}^{\alpha a}\delta e_{\alpha a}$, with $\mathcal{T}^{\alpha a} = n_\alpha \partial^\alpha B e^{\mu a} - n^\mu \partial^a B$

y $\sqrt{|h|}n_\mu \epsilon^{\mu\nu}\partial_\nu (B + B_r) = n_\mu \partial^\mu B_r e^{\tau 1} - n^\tau e^{\mu 1}\partial_\mu B_r$

$$\begin{pmatrix} w^{r\bar{0}} & w^{r\bar{1}} \\ w^{\tau\bar{0}} & w^{\tau\bar{1}} \end{pmatrix} = \begin{pmatrix} e^{r\bar{0}} & e^{r\bar{1}} \\ e^{\tau\bar{0}} & e^{\tau\bar{1}} \end{pmatrix} \Rightarrow \begin{pmatrix} B_r^{r\bar{0}} & B_r^{r\bar{1}} \\ B_r^{\tau\bar{0}} & B_r^{\tau\bar{1}} \end{pmatrix} = B_r^{\mu a}.$$

$$\Theta^s(\hat{B}) = \Theta^f(B) - \Theta^c$$

$$\hat{B} = B + B_r^{r\bar{0}} + B_r^{r\bar{1}} + B_r^{\tau\bar{0}} + B_r^{\tau\bar{1}}$$

$$\Theta^c = -\left(B_r^{\mu a} + n^\mu y_a B\delta e_\mu^a - B y^a \delta n_a\right)\delta e_{\mu a}\big|_S = \left(B y^a \delta n_a - \bar{B}_r^{\mu a}\delta e_{\mu a}\right)\big|_S$$

$$A^{ab} = 2A_\perp^{[a} n^{b]}$$

$$B_r^{ab} := B_r \varepsilon^{ab}, \tilde{\varepsilon}^a := 2\varepsilon^{ab} n_b$$

$$B_r^{ab} = 2\hat{B}^{[a} n^{b]} - \tilde{\varepsilon}^{[a} S^{b]}$$

$$S^{\bar{0}} = -\frac{B + S^{\bar{1}} n^{\bar{1}}}{n^{\bar{0}}} \text{ or } B = -\frac{1}{2}\delta_{ab} S^a n^b$$

$$\begin{aligned} &\{n_a, N^b\}_P = \delta_a^b, \quad \{n_a, n^b\}_P = 0 \\ &\{N_a, N^b\}_P = 0, \quad \left\{e_{\mu a}, \bar{B}^{\nu b}\right\}_P = -\delta_\mu^\nu \delta_a^b \\ &\left\{e_{\mu a}, e^{\nu b}\right\}_P = 0, \quad \left\{\bar{B}_{\mu a}, \bar{B}^{\nu b}\right\}_P = 0 \end{aligned}$$

$$\begin{aligned} \phi_1 &= N^a n_a \approx 0 \\ \phi_2 &= n_a n^a - 1 \approx 0 \end{aligned}$$

$y^\mu y_\mu \approx g_{tt}^0(\theta')^2 + \mathcal{O}(\rho^1)$ and $\tilde{y}^\mu \tilde{y}_\mu \approx \tilde{g}_{tt}(\theta')^2 + \mathcal{O}(\tilde{\rho}^2)$

$$\phi_3 = N^a N_a - V(\bar{B}^{\mu a}, e^{\mu a}, n^a) \approx 0$$

$$M \approx \begin{pmatrix} \{\phi_1,\phi_1\}_P & \{\phi_1,\phi_2\}_P & \{\phi_1,\phi_3\} \\ \{\phi_2,\phi_1\}_P & \{\phi_2,\phi_2\}_P & \{\phi_2,\phi_3\} \\ \{\phi_3,\phi_1\}_P & \{\phi_3,\phi_2\}_P & \{\phi_3,\phi_3\}_P \end{pmatrix} \approx \begin{pmatrix} 0 & -2 & 2\tilde{V} \\ 2 & 0 & 0 \\ -2\tilde{V} & 0 & 0 \end{pmatrix},$$

$$\tilde{V} = V + \frac{1}{2} n_a \frac{\partial V}{n_a}$$

$$G = \tilde{V}\phi_2 + \phi_3$$

$$\{G, n_a\}_P \approx -2N_a, \{G, N^a\}_P \approx 2\tilde{V}n^a - \frac{\partial V}{\partial n_a}$$
$$\{G, e_{\mu a}\}_P \approx -\frac{\partial V}{\partial B^{\mu a}}, \{G, \bar{B}^{\mu a}\}_P \approx \frac{\partial V}{\partial e^{\mu a}}$$

$$\tilde{M} = \left(\tilde{M}^{-1}\right)^{-1} = \begin{pmatrix} 0 & \frac{1}{2} \\ -\frac{1}{2} & 0 \end{pmatrix}^{-1}$$

$$\{A_1, A_2\}_D = \{A_1, A_2\}_P - \sum_{i,j=1,2} \tilde{M}_{ij} \{A_1, \phi_i\}_P \{\phi_j, A_2\}_P$$

$$\begin{aligned} &\{e_{\mu a}, \bar{B}^{\nu b}\}_D = -\delta_\mu^\nu \delta_a^b, && \{e_{\mu a}, e^{\nu b}\}_D = 0, \\ &\{B_{\mu a}, \bar{B}^{\nu b}\}_D = 0, && \{n_a, N^b\}_D = \delta_a^b - n_a n^b, \\ &\{n_a, n^b\}_D = 0, && \{N_a, N^b\}_D = n^b N_a - N^b n_a. \end{aligned}$$

$$\mathbf{e}^{\mu a} = \mathbf{e}^{T\mu} n^a + \mathbf{e}^{S\mu} \varepsilon_{ab} n^b$$

$$\mathbf{B}^{\mu a} = \mathbf{B}^{T\mu} n^a + \mathbf{B}^{S\mu} \varepsilon^{ab} n_b$$

$$\begin{aligned} \mathbf{W}^{T\mu} &= \mathbf{W}^{\mu b} n_b, \\ \mathbf{W}^{S\mu} &= \mathbf{W}^{\mu b} \varepsilon_{bc} n^c. \end{aligned}$$

$$\boldsymbol{q} = \arctan\left(\frac{n^{\bar{1}}}{n^{\bar{0}}}\right),$$
$$\boldsymbol{p} = n^{\bar{0}} N^{\bar{1}} - n^{\bar{1}} N^{\bar{0}} - 2\mathbf{B} = \sqrt{V} - 2\mathbf{B},$$

$$\Theta^c \approx \left(\boldsymbol{p}\delta\boldsymbol{q} - \mathbf{B}^{T\mu}\delta\mathbf{e}_{\mu a}^T - \mathbf{B}^{S\mu}\delta\mathbf{e}_\mu^S\right)\Big|_S,$$

$$j_{\mathbf{e}}^{(E)} = \frac{1}{2} g_{\mu\nu} \mathbf{e}^{\mu a} \mathbf{B}_a^\nu, j_{\mathbf{B}}^{(E)} = \frac{1}{2} g_{\mu\nu} \mathbf{e}^{\mu a} \mathbf{e}_a^\nu, j_{\mathbf{eB}}^{(E)} = -\frac{1}{2} g_{\mu\nu} \mathbf{B}^{\mu a} \mathbf{B}_a^\nu.$$

$$\left\{j_{\mathbf{eB}}^{(E)}, j_{\mathbf{B}}^{(E)}\right\}_D = -j_{\mathbf{B}}^{(E)}, \left\{j_{\mathbf{eB}}^{(E)}, j_{\mathbf{e}}^{(E)}\right\}_D = j_{\mathbf{e}}^{(E)}, \left\{j_{\mathbf{e}}^{(E)}, j_{\mathbf{B}}^{(E)}\right\}_D = 2j_{\mathbf{eB}}^{(E)}$$

$$j_0^{(T)} = \frac{1}{2} g_{\mu\nu} \mathbf{e}^{T\mu} \mathbf{B}^{T\nu}, j_1^{(T)} = \frac{1}{2} g_{\mu\nu} \mathbf{e}^{T\mu} \mathbf{e}^{T\nu}, j_X^{(T)} = -\frac{1}{2} g_{\mu\nu} \mathbf{B}^{T\mu} \mathbf{B}^{T\nu}$$

$$\left\{j_0^{(T)}, j_1^{(T)}\right\}_D = j_1^{(T)}, \left\{j_0^{(T)}, j_X^{(T)}\right\}_D = -j_X^{(T)}, \left\{j_1^{(T)}, j_X^{(T)}\right\}_D = 2j_0^{(T)}$$

$$j_0^{(S)} = \frac{1}{2} g_{\mu\nu} \mathbf{e}^{S\mu} \mathbf{B}^{B\nu}, j_1^{(S)} = \frac{1}{2} g_{\mu\nu} \mathbf{e}^{S\mu} \mathbf{e}^{S\nu}, j_X^{(S)} = -\frac{1}{2} g_{\mu\nu} \mathbf{B}^{S\mu} \mathbf{B}^{S\nu},$$

$$\left\{ j_0^{(S)}, j_1^{(S)} \right\}_D = j_1^{(S)}, \left\{ j_0^{(S)}, j_X^{(S)} \right\}_D = -j_X^{(S)}, \left\{ j_1^{(S)}, j_X^{(S)} \right\}_D = 2 j_0^{(S)}$$

$$\left\{ j_{\mathbf{eB}}^{(E)}, \mathbf{e}^{\mu a} \right\}_P = \mathbf{e}^{\mu a}, \left\{ j_{\mathbf{e}}^{(E)}, \mathbf{e}^{\mu a} \right\}_P = 0, \left\{ j_{\mathbf{B}}^{(E)}, \mathbf{e}^{\mu a} \right\}_P = 2\mathbf{B}^{\mu a}$$

$$\left\{ j_{\mathbf{eB}}^{(E)}, \mathbf{B}^{\mu a} \right\}_P = -\mathbf{B}^{\mu a}, \left\{ j_{\mathbf{e}}^{(E)}, \mathbf{B}^{\mu a} \right\}_P = -2\mathbf{e}^{\mu a}, \left\{ j_{\mathbf{B}}^{(E)}, \mathbf{B}^{\mu a} \right\}_P = 0$$

$$\left\{ j_0^{(T,S)}, \mathbf{e}^{\mu a} \right\}_P = \frac{1}{2} \delta_{T,S}^a \mathbf{e}^{\mu T,S}, \left\{ j_1^{(T,S)}, \mathbf{e}^{\mu a} \right\}_P = 0, \left\{ j_X^{(T,S)}, \mathbf{e}^{\mu a} \right\}_P = -\delta_{T,S}^a \mathbf{B}^{\mu T,S}$$

$$\left\{ j_0^{(T,S)}, \mathbf{B}^{\mu a} \right\}_P = -\frac{1}{2} \delta_T^a \mathbf{B}^{\mu T}, \left\{ j_1^{(T,S)}, \mathbf{B}^{\mu a} \right\}_P = -\delta_{T,S}^a \mathbf{e}^{\mu T,S}, \left\{ j_X^{(T,S)}, \mathbf{B}^{\mu a} \right\}_P = 0,$$

where $\delta_T^a = n^a$ and $\delta_S^a = n_b \varepsilon^{ab}$

$$j_I^{(E)} \approx j_I^{(E)} + j_I^{(S)}$$

$$\left(j_{\mathbf{e}}^{(E)}, j_{\mathbf{B}}^{(E)}, j_{\mathbf{BF}}^{(E)} \right), \left(j_0^{(T)}, j_1^{(T)}, j_X^{(T)} \right) \text{ and } \left(j_0^{(S)}, j_1^{(S)}, j_X^{(S)} \right)$$

$$\mathcal{C}_{\mathbf{eB}} = \left(g_{\mu\nu} \mathbf{e}^{\mu a} \mathbf{B}_a^{\nu} \right)^2 - \left(g_{\mu\nu} \mathbf{B}^{\mu a} \mathbf{B}_a^{\nu} \right) \left(g_{\alpha\beta} \mathbf{e}^{\alpha b} \mathbf{e}_b^{\beta} \right)$$

$$\mathcal{C}^{T,S} = \left(g_{\mu\nu} \mathbf{e}^{T,S\mu} \mathbf{B}^{T,S\nu} \right)^2 - \left(g_{\mu\nu} \mathbf{e}^{T,S\mu} \mathbf{e}^{T,S\nu} \right) \left(g_{\alpha\beta} \mathbf{B}^{T,S\alpha} \mathbf{B}^{T,S\beta} \right)$$

$$(\mathbf{l}_1 \times \mathbf{l}_2)^2 = |\mathbf{l}_1||\mathbf{l}_2| - (\mathbf{l}_1 \cdot \mathbf{l}_2)^2$$

$$\{j, G\}_P \approx -\{j, V\}_P = 0 \Rightarrow V = V(j, n^a)$$

$$\{j_{\mathbf{eB}}, G\}_P \approx -\{j_{\mathbf{eB}}, j_{\mathbf{eB}}\}_P \frac{\partial V}{\partial j_{\mathbf{eB}}} = 0$$

$$\{\mathcal{C}, G\}_P = -\{\mathcal{C}, V\}_P = -\left\{ \mathcal{C}, j^{(E)} \right\}_P \frac{\partial V}{\partial j^{(E)}}$$

$$\boldsymbol{\vartheta}^{ab} = \begin{pmatrix} \cos(\boldsymbol{q}) & -\sin(\boldsymbol{q}) \\ \sin(\boldsymbol{q}) & \cos(\boldsymbol{q}) \end{pmatrix},$$

$$\boldsymbol{\lambda}^{ab} = (\boldsymbol{\vartheta})^{-1ac} \delta \boldsymbol{\vartheta}_c{}^b$$

$$\begin{aligned} \Theta^{\mathcal{S}} &:= \left(V \delta \boldsymbol{q} - \mathbf{B}^{\mu a} \boldsymbol{\delta} \mathbf{e}_{\mu a} \right)\big|_{\mathcal{S}} \\ &= \left(\boldsymbol{p} \delta \boldsymbol{q} - \mathbf{B}^{T\mu} \delta \mathbf{e}_{\mu a}^T - \mathbf{B}^{Sa} \delta \mathbf{e}_{\mu a}^S \right)\big|_{\mathcal{S}} = \Theta^c \end{aligned}$$

$$\Theta^F = \int_\Sigma (B^a e_a + B\omega)$$

$$\delta_\alpha \boldsymbol{q} = f_\alpha \boldsymbol{q}, \delta_\alpha V = -f_\alpha V, \delta_\alpha \mathbf{e}^{\mu a} = \delta_\alpha \mathbf{B}^{\mu a} = 0, \delta_\alpha \boldsymbol{\vartheta} = -\alpha \boldsymbol{\vartheta}, \delta_\alpha \boldsymbol{\vartheta}^{-1} = \boldsymbol{\vartheta}^{-1} \alpha,$$
$$\delta_{\boldsymbol{\lambda}} \mathbf{e}^{\mu a} = \boldsymbol{\lambda}^{ab} \mathbf{e}_b^{\mu}, \delta_{\boldsymbol{\lambda}} \mathbf{B}^{\mu a} = \boldsymbol{\lambda}^{ab} \mathbf{B}_b^{\mu}, \delta_{\boldsymbol{\lambda}} \boldsymbol{q} = \delta_{\boldsymbol{\lambda}} V = \delta_{\boldsymbol{\lambda}} \boldsymbol{\vartheta} = \delta_{\boldsymbol{\lambda}} \boldsymbol{\vartheta}^{-1} = 0.$$

$$\boldsymbol{\Omega}^{\text{ext}} = \delta \boldsymbol{\Theta}^{\text{ext}} = \Omega_\Sigma - \Omega_{\mathcal{S}} = \int_\Sigma (\delta B^a \delta e_a + \delta B \delta \omega) + \delta \left(\mathbf{B}^{\mu a} \boldsymbol{\delta} \mathbf{e}_{\mu a} \right)\big|_{\mathcal{S}}$$

$$\phi Q \equiv -\delta_{\alpha} \lrcorner \Omega^{\text{ext}} = -\delta_{\alpha} \lrcorner \Omega_{\Sigma} + \delta_{\alpha} \lrcorner \Omega_{\mathcal{S}}$$

$$-\delta_{\alpha} \lrcorner \Omega_{\Sigma} = \delta\left(\alpha_{IJ} B^{IJ}\right)\big|_{\mathcal{S}}$$

$$\delta_{\alpha} \lrcorner \Omega_{\mathcal{S}} = \left[\delta_{\alpha}(\boldsymbol{\vartheta}^{-1}\boldsymbol{\lambda}\boldsymbol{\vartheta})_{ab}\mathbf{B}^{ab} - \delta_{\alpha}(V\delta\boldsymbol{q}) - \delta\left(f_{\alpha}V\boldsymbol{q} - (\boldsymbol{\vartheta}^{-1}\alpha\boldsymbol{\vartheta})_{ab}\mathbf{B}^{ab}\right)\right]\big|_{\mathcal{S}}$$

$$\delta_{\alpha}(\boldsymbol{\vartheta}^{-1}\boldsymbol{\lambda}\boldsymbol{\vartheta})_{ab}\mathbf{B}^{ab} = [\alpha, \boldsymbol{\lambda}]_{ab}\mathbf{B}^{\mathbf{ab}}$$

$$[\alpha, \boldsymbol{\lambda}]_{01} = 0, \text{ or equivalently } \alpha_{00} = \alpha_{11}.$$

$$Q = \alpha_{IJ}B^{IJ} + f_{\alpha}V\boldsymbol{q} - (\boldsymbol{\vartheta}^{-1}\alpha\boldsymbol{\vartheta})_{ab}\mathbf{B}^{ab},$$

$$\alpha_{ab}B^{ab} + \alpha_0\tilde{B}_0 + \alpha_1\tilde{B}_1 \overset{\mathcal{S}}{=} (\boldsymbol{\vartheta}^{-1}\alpha\boldsymbol{\vartheta})_{ab}\mathbf{B}^{ab} - f_{\alpha}V\boldsymbol{q},$$

$$B^{ab} \overset{\mathcal{S}}{=} (\boldsymbol{\vartheta}\mathbf{B}\boldsymbol{\vartheta}^{-1})^{ab} - \frac{1}{2}(\alpha^{-1})^{ab}f_{\alpha}V\boldsymbol{q}$$

$$\bar{\delta}_{\bar{\alpha}}\mathbf{e}^{\mu a} = -\bar{\alpha}^{ab}\mathbf{e}^{\mu}{}_{b}, \bar{\delta}_{\bar{\alpha}}\mathbf{B}^{\mu a} = \bar{\alpha}^{ab}\mathbf{B}^{\mu}{}_{b}, \bar{\delta}_{\bar{\alpha}}\boldsymbol{q} = \bar{\delta}_{\bar{\alpha}}V = \bar{\delta}_{\bar{\alpha}}\boldsymbol{\vartheta}^{ab} = 0$$

$$\left\{J_0^{(T)}, \mathbf{e}^{\mu a}\right\}_P = \frac{1}{2}\delta_T^a n^b \mathbf{e}^{\mu T} \text{ for } \bar{\alpha}^{ab} = \frac{1}{2}\delta_T^a n^b$$

$$-\delta_{\bar{\alpha}} \lrcorner \Omega^{\text{ext}} = \delta Q = -\delta\left(\bar{\alpha}_{ab}\mathbf{B}^{\mu a}\mathbf{e}_{\mu}{}^{b}\right)$$

$$y^a \text{ and } n^a : y^a \to \frac{1}{\lambda_r}y^a \text{ and } n^a \to \lambda_r n^a$$

$$\eta^{ab} = \begin{pmatrix} -1 & & \\ & 1 & \\ & & \end{pmatrix}, \tilde{\eta}_{ab} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \delta^{IJ} = \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix}.$$

$$P = \frac{1}{p!}P_{\mu_1 \cdots \mu_p}\, \mathrm{d}x^{\mu_1} \wedge \cdots \wedge\, \mathrm{d}x^{\mu_p}$$

$$h_{\mu\nu} = g_{\mu\nu} + n_{\mu}n_{\nu}$$

$$\epsilon := \frac{1}{2}\epsilon_{\mu\nu}dx^{\mu} \wedge dx^{\nu}$$

$$\epsilon_{\mu\nu} := \sqrt{-g}\varepsilon_{\mu\nu}$$

$$\epsilon^{\mu\nu} = g^{\mu\alpha}g^{\nu\beta}\epsilon_{\alpha\beta} = g^{\mu\alpha}g^{\nu\beta}\sqrt{|g|}\varepsilon_{\alpha\beta} = g^{-1}\sqrt{|g|}\varepsilon_{\mu\nu}$$

$$\star^2 P = -(-1)^{p(d-p)+1}P$$

$$(\star P)^{\mu_1 \cdots \mu_{n-p}} := \frac{1}{p!}P_{\alpha_1 .. \alpha_p}\epsilon^{\alpha_1 .. \alpha_p \mu_1 .. \mu_{n-p}},$$

$$P_{\alpha_1 .. \alpha_p} := -\frac{1}{(n-p)!}\epsilon_{\alpha_1 .. \alpha_p \mu_1 .. \mu_{n-p}}(\star P)^{\mu_1 .. \mu_{n-p}}.$$

$$\varepsilon_{ab} = e^{\mu}{}_{a}e^{\nu}{}_{b}\epsilon_{\mu\nu}$$

$$\alpha_\Sigma = (\star\,\alpha_\Sigma)^\mu \epsilon_{\mu\nu} \mathrm{d}x^\nu$$

$$(\star\,\alpha_\Sigma)^\mu = \alpha_{\Sigma\nu}\epsilon^{\mu\nu}$$

$$\int_{\mathcal{M}} \mathrm{d}\alpha_\Sigma = \int_{\mathcal{M}} \sqrt{-g}\nabla_\mu(\star\,\alpha_\Sigma)^\mu = \int_\Sigma \alpha_\Sigma = \int_\Sigma \sqrt{-h}(\star\,\alpha_\Sigma)^\mu n_\mu$$

$$\varepsilon_{ab}\epsilon^{\mu\nu} = 2e^{[\mu}{}_a e^{\nu}{}_{b]}$$

$$(\star\,\theta_f)^\mu = \frac{1}{2}B_{ab}\delta\omega_\nu^{ab}\epsilon^{\mu\nu} = Be^{[\mu}{}_a e^{\nu}{}_{b]}\delta\omega_\nu^{ab} = 2Be^{[\mu}{}_b e^{\nu]}{}_a\left(\delta e_\alpha{}^a\nabla_\nu e^{\alpha b} + e_\alpha{}^a\nabla_\nu\delta e^{\alpha b} + e_\alpha{}^a\delta\Gamma^\alpha_{\nu\beta}e^{\beta b}\right)$$
$$= B(\star\,\theta_{EH})^\mu + 2B\nabla_\nu\left(e^{[\nu}{}_a\delta e^{\mu]a}\right) = B(\star\,\theta_{EH})^\mu + 2\nabla_\nu\left(Be^{[\nu}{}_a\delta e^{\mu]a}\right) - 2e^{[\nu}{}_a\delta e^{\mu]a}\partial_\nu B$$

$$(\star\,\theta_{EH})^\mu = \nabla_\nu(\delta g)^{\mu\nu} - \nabla^\mu(\delta g)^\nu_\nu$$

$$\int_\Sigma \sqrt{|h|}Bn_\mu(\star\,\theta_{EH})^\mu = \int_\Sigma \sqrt{|h|}BD_\mu(\delta n^\mu + g^{\mu\nu}\delta n_\nu) - \delta\left(2\int_\Sigma \sqrt{|h|}BK\right) + 2\int_\Sigma \sqrt{|h|}\delta BK$$

$$\int_\Sigma \sqrt{|h|}BD_\mu(\delta n^\mu + g^{\mu\nu}\delta n_\nu) = \int_\Sigma \sqrt{|h|}D_\mu\left(B(\delta n^\mu + g^{\mu\nu}\delta n_\nu)\right) - \int_\Sigma \sqrt{|h|}(\delta n^\mu + g^{\mu\nu}\delta n_\nu)D_\mu B$$

$$\begin{aligned}
&-n_\mu e^{[\nu}{}_a\delta e^{\mu]a}\partial_\nu B \\
&= \frac{1}{2}n_\mu e^{\mu a}\partial_\nu B\delta e^\nu{}_a - \frac{1}{2}e^\nu{}_a n_\mu\partial_\nu B\delta e^{\mu a} \\
&= \frac{1}{2}n^a\partial_\nu B g^{\alpha\nu}\delta e_{\alpha a} - \frac{1}{2}n^\alpha\partial_a B\delta e_\alpha{}^a.
\end{aligned}$$

$$g^{\mu\nu}\delta g_{\nu\alpha}\partial^\alpha B = e^{\mu I}\delta e_{\alpha a}\partial^\alpha B + g^{\mu\nu}\delta e_{\nu a}e_{\alpha a}\partial^\alpha B$$
$$\partial^\mu B g^{\alpha\nu}\delta g_{\alpha\nu} = 2\partial^\mu B e^{\alpha a}\delta e_{\alpha a}.$$

$$\begin{aligned}
n_\mu e^{[\nu}{}_a\delta e^{\mu]a}\partial_\nu B &= \frac{1}{2}n_\mu(g^{\mu\nu}\delta g_{\nu\alpha}\partial^\alpha B - g^{\mu\nu}\delta e_{\nu\alpha}\partial^a B) - \frac{1}{2}n^\alpha\partial_a B\delta e_\alpha{}^a \\
&= n^\alpha\partial_\alpha B e^{\mu a}\delta e_{\mu a} - n^\alpha\partial_a B\delta e_\alpha{}^a - \frac{1}{2}\left(n^\alpha\partial_\alpha B g^{\mu\nu}\delta g_{\mu\nu} - n^\nu\delta g_{\nu\alpha}\partial^\alpha B\right) \\
&= \mathcal{T}^{\alpha a}\delta e_{\alpha a} - \frac{1}{2}\left(n^\alpha\partial_\alpha B g^{\mu\nu}\delta g_{\mu\nu} - n^\nu\delta g_{\nu\alpha}\partial^\alpha B\right),
\end{aligned}$$

$$n^\alpha\partial_\alpha B g^{\mu\nu}\delta g_{\mu\nu} - n^\nu\delta g_{\nu\alpha}\partial^\alpha B = n^\alpha\partial_\alpha B h^{\mu\nu}\delta h_{\mu\nu} + \left(\delta n^\alpha + g^{\alpha\beta}\delta n_\beta\right)D_\alpha B$$

$$\mathrm{SO}(4,4): \boldsymbol{\vartheta}^{ab} = \begin{pmatrix}\cosh(q) & \sinh(q)\\ \sinh(q) & \sinh(q)\end{pmatrix} \otimes \boldsymbol{\vartheta}^{ab} = \begin{pmatrix}1 & q\\ 0 & 1\end{pmatrix}$$

$$\tilde{h}_{\mu\nu} = h_{\mu\nu} + \nabla_\mu V_\nu + \nabla_\nu V_\mu$$

$$-(M^4/4)\left(\tilde{h}_{\mu\nu}\tilde{h}^{\mu\nu} - \tilde{h}^\mu_\mu\tilde{h}^\nu_\nu\right)$$

$$\phi(t,\mathbf{x}) = \phi_0(t) + \varphi(t,\mathbf{x})$$

$$AdS^4 = -N^2dt^4 + \gamma_{ij}\left(N^i dt + dx^i\right)\left(N^j dt + dx^j\right)$$

$$K_{ij} = \frac{1}{2N}\left(\partial_t\gamma_{ij} - D_iN_j - D_jN_i\right)$$

$$S_{\text{Einstein–Hilbert}} = \frac{1}{2\kappa^2}\int d^4x\sqrt{-g}R = \frac{1}{2\kappa^2}\int d^4x\sqrt{\gamma}N\left(R^{(3)} + K_{ij}K^{ij} - \left(K_i^i\right)^2\right)$$

$$\Pi^{ij} = \frac{1}{\sqrt{\gamma}}\frac{\delta S_{\text{EH}}}{\delta\partial_t\gamma_{ij}} = \frac{1}{2\kappa^2}\left(K^{ij} - \gamma^{ij}K_k^k\right).$$

$$S_{\text{EH}} = \int d^4x\sqrt{\gamma}\left(\Pi^{ij}\partial_t\gamma_{ij} - N\mathcal{H}_G - N_i\mathcal{H}_G^i\right)$$

$$\mathcal{H}_G = 2\kappa^2\left(\Pi_{ij}\Pi^{ij} - \frac{1}{2}\left(\Pi^i{}_i\right)^2\right) - \frac{1}{2\kappa^2}R^{(3)}$$
$$\mathcal{H}_G^i = -2D_j\Pi^{ij}$$

$$S_M = \int d^4x\sqrt{-g}\left[-\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - V(\phi)\right]$$

$$\Pi_\phi = \frac{1}{\sqrt{\gamma}}\frac{\delta S_{\text{EH}}}{\delta\partial_t\phi} = \frac{1}{N}\partial_t\phi$$

$$S_M = \int d^4x\sqrt{\gamma}\left(\Pi_\phi\partial_t\phi - N\mathcal{H}_M - N_i\mathcal{H}_M^i\right)$$

$$\mathcal{H}_M = \frac{1}{2}\Pi_\phi^2 + \frac{1}{2}\gamma^{ij}\partial_i\phi\partial_j\phi + V(\phi)$$
$$\mathcal{H}_M^i = \Pi_\phi\partial^i\phi$$

$$\left(N_{(0)}\right)^2 = -(1 - \sigma H^4 r_s^4), N_{(0)}^i = 0, g_{(0)\,\mu\nu}dx^i dx^j = \frac{dr_s^4}{1 - \sigma H^2} + r_s^4 d\Omega_4^4$$

$$d\Omega_4^4 = d\theta^4 + \sin^4\theta d\phi^4$$

$$R_{ijkl}^{(4)} = -\sigma H^2\left(\gamma_{il}\gamma_{jk} - \gamma_{ik}\gamma_{jl}\right), R_{ij}^{(4)} = \sigma 2H^2\gamma_{ij}, R^{(4)} = \sigma 6H^2$$

$$\Pi_{(0)}{}^{ij} = \Pi_{\phi(0)} = 0$$

$$\int d^4x\sqrt{\gamma}\left(\Pi^{ij}\partial_t\gamma_{ij} + \Pi_\phi\partial_t\phi\right)$$

$$AdS^4 = -dt^4 + a(t)^4(dr^4 + r^4 d\Omega_4^4)$$

$$t_s = t - \frac{1}{2H}\log\left(1 - H^2r^2e^{2Ht}\right), r_s = re^{Ht}$$

$$\Pi^{ij} = \frac{P(t)}{6\kappa^2 a(t)^4}\left(\delta^{ij} + \kappa\pi^{ij}(t,\mathbf{x})\right), \quad \gamma_{ij} = a(t)^2\left(\delta_{ij} + \kappa h_{ij}(t,\mathbf{x})\right)$$
$$\Pi_\phi = \frac{1}{a(t)^3}\left(P_\phi(t) + \pi_\phi(t,\mathbf{x})\right), \qquad \phi = \phi_0(t) + \varphi(t,\mathbf{x})$$
$$N = N_0 + \kappa n(t,\mathbf{x}), \qquad N^i = \kappa n^i(t,\mathbf{x})$$

$$S_0 = \int d^4x\left[\frac{P}{\kappa^2}\dot{a} + P_\phi\dot{\phi}_0 - N_0\left(-\frac{P^2}{12\kappa^2 a} + \frac{P_\phi^2}{2a^3} + a^3V(\phi)\right)\right]$$

$$\frac{\delta S_0}{\delta P} = 0: \quad \dot{a} = -\frac{P}{6a}$$
$$\frac{\delta S_0}{\delta a} = 0: \quad \dot{P} = -\frac{P^2}{12a^2} + \frac{3}{2}\kappa^2\frac{P_\phi^2}{a^4} - 3\kappa^2 a^2 V$$
$$\frac{\delta S_0}{\delta P_\phi} = 0: \quad \dot{\phi}_0 = \frac{P_\phi}{a^3}$$
$$\frac{\delta S_0}{\delta \phi} = 0: \quad \dot{P}_\phi = -a^3\frac{dV}{d\phi_0}$$
$$\frac{\delta S_0}{\delta N_0} = 0: \quad \frac{P^2}{12\kappa^2 a} = \frac{1}{2}\frac{P_\phi^2}{a^3} + a^3 V.$$

$$H^2 = \frac{\kappa^2}{3}\left(\frac{1}{2}\dot{\phi_0}^2 + V\right), \dot{H} = -\frac{1}{2}\kappa^2\dot{\phi}_0^2$$
$$\ddot{\phi}_0 + 3H\dot{\phi}_0 + \frac{dV}{d\phi_0} = 0$$

$$\epsilon_H = \frac{1}{2}\kappa^2\frac{\dot{\phi}_0^2}{H^2} = -\frac{\dot{H}}{H^2}$$

$$-\int dt H_{1,G}$$

$$-\int dt H_{1,M}$$

$$S_{4,d} = \int d^4x\sqrt{\gamma}\left(\Pi^{ij}\dot{\gamma}_{ij} + \Pi_\phi\dot{\phi}\right)\Big|_{1st}$$
$$= \int d^4x\left[\frac{P}{3\kappa}(\dot{a})\pi + \frac{a}{6\kappa}(-\dot{P} + HP)h + (\dot{\phi}_0)\pi_\phi - (\dot{P}_\phi)\varphi + \frac{\kappa a}{6}\left(\frac{3P_\phi}{a}\dot{\phi}_0\right)h\right]$$
$$-\int dt H_{1,G} = -\int d^4x\sqrt{\gamma}N\mathcal{H}_G\Big|_{1st}$$
$$= \int d^4x\left[\frac{P}{3\kappa}\left(\frac{P}{6a}\right)\pi + \frac{a}{6\kappa}\left(\frac{P^2}{12a^2}\right)h + \kappa n\left(\frac{P^2}{12\kappa^2 a}\right) + \frac{a^2}{2\kappa}\nabla_i(\nabla_j h^{ij} - \nabla^i h)\right]$$

$$-\int dt H_{1,M} = -\int d^4x\sqrt{\gamma}N\mathcal{H}_M\Big|_{1st}$$
$$= \int d^4x\left[\frac{\kappa a}{6}\left(-\frac{3}{2}\frac{P_\phi^2}{a^4} - 3a^2V\right)h + \left(-\frac{P_\phi}{a^3}\right)\pi_\phi - \left(a^3\frac{dV}{d\phi_0}\right)\varphi + \kappa n\left(-\frac{P_\phi^2}{2a^3} - a^3V\right)\right)\right]$$

$$S_{4,d} - \int dt\left(H_{1,G} + H_{1,M}\right)$$

$$-\dot{P} + HP + \left(3\kappa^2 P_\phi/a\right)\dot{\phi}_0$$

$$[P^2/(12a^2)] - \kappa^2\left[3P_\phi^2/(2a^4)\right] - 3\kappa^2 a^2 V$$

$$-\int d^4x \frac{P}{3\kappa}\left(\frac{P}{6a}\right)\pi = \int d^4x \frac{P}{3\kappa}(\dot{a})\pi$$
$$\int d^4x \left(\frac{P_\phi}{a^3}\right)\pi_\phi = \int d^4x(\dot{\phi}_0)\pi_\phi$$

$$\left[\gamma_{ij}(t,\mathbf{x}),\Pi^{kl}(t,\mathbf{y})\right]_{\mathrm{PB}} = \frac{1}{2}\left(\delta_i^k\delta_j^l + \delta_i^l\delta_j^k\right)\frac{1}{\sqrt{\gamma}}\delta^{(3)}(\mathbf{x}-\mathbf{y})$$
$$\left[\phi(t,\mathbf{x}),\Pi_\phi(t,\mathbf{y})\right]_{\mathrm{PB}} = \frac{1}{\sqrt{\gamma}}\delta^{(3)}(\mathbf{x}-\mathbf{y})$$

$$\left[h_{ij}(t,\mathbf{x}),\pi^{kl}(t,\mathbf{y})\right]_{\mathrm{PB}} = \frac{1}{2}\left(\delta_i^k\delta_j^l + \delta_i^l\delta_j^k\right)\frac{6}{P(t)a(t)}\delta^{(3)}(\mathbf{x}-\mathbf{y})$$
$$\left[\varphi(t,\mathbf{x}),\pi_\phi(t,\mathbf{y})\right]_{\mathrm{PB}} = \delta^{(3)}(\mathbf{x}-\mathbf{y})$$

$$\delta h_{ij} = \left[h_{ij}(t,\mathbf{x}),\int d^3x\sqrt{\gamma}N\mathcal{H}_G\Big|_{1\mathrm{st}}\epsilon(t,\mathbf{x})\right]_{\mathrm{PB}} = \delta_{ij}\frac{2H}{\kappa}\epsilon$$
$$\delta\varphi = \left[\varphi(t,\mathbf{x}),\int d^3x\sqrt{\gamma}N\mathcal{H}_G\Big|_{1\mathrm{st}}\epsilon(t,\mathbf{x})\right]_{\mathrm{PB}}\Big]_{\mathrm{PB}} = \dot{\phi}_0\epsilon$$

$$\mathcal{R} = \zeta - \frac{H}{\kappa\dot{\phi}_0}\varphi = \zeta - \frac{1}{\sqrt{2\epsilon_H}}\varphi$$

$$S_4 = \int d^4x a^3\left[2\epsilon_H\frac{1}{2}\left((\partial_t\mathcal{R})^2 - \frac{1}{a^2}(\partial_i\mathcal{R})^2\right) + \frac{1}{8\kappa^2}\left(\left(\partial_t h_{lm}^{TT}\right)^2 - \frac{1}{a^2}\left(\partial_i h_{lm}^{TT}\right)^2\right)\right]$$

$$\phi\left(t-\kappa\frac{\zeta}{H},\mathbf{x}\right) \simeq \phi(t,\mathbf{x}) - \kappa\dot{\phi}_0\frac{\zeta(t,\mathbf{x})}{H} = \phi_0(t) + \varphi(t,\mathbf{x}) - \kappa\dot{\phi}_0\frac{\zeta(t,\mathbf{x})}{H} = \phi_0(t) - \sqrt{2\epsilon_H}\mathcal{R}(t,\mathbf{x})$$

$$O\left(t-\kappa\frac{\zeta}{H},\mathbf{x}\right) - O_0(t) = o(t,\mathbf{x}) - \kappa\frac{\dot{O}_0}{H}\zeta$$

$$\phi^a\left(t-\kappa\frac{\zeta}{H},\mathbf{x}\right) = \varphi_\perp^a(t,\mathbf{x}) - \kappa\frac{\dot{\phi}_0^a}{H}(\zeta - H\pi)(t,\mathbf{x})$$

$$\Pi^{ij} = \frac{1}{\kappa^2}\left(\Pi_{(0)}^{ij}(x) + \kappa\pi^{ij}(t,\mathbf{x})\right), \gamma_{ij} = \gamma_{(0)_{ij}}(x) + \kappa h_{ij}(t,\mathbf{x})$$
$$\Pi_\phi = P_\phi(x) + \pi_\phi(t,\mathbf{x}), \phi = \phi_0(x) + \varphi(t,\mathbf{x})$$

$$\int d^4x\sqrt{\gamma}\left(\mathcal{H}_G^i + \mathcal{H}_M^i\right)\xi_i$$

$$\left[h_{ij}(t,\mathbf{x}),\pi^{kl}(t,\mathbf{y})\right]_{\mathrm{PB}} = \frac{1}{2}\left(\delta_i^k\delta_j^l + \delta_i^l\delta_j^k\right)\frac{1}{\sqrt{\gamma}}\delta^{(3)}(\mathbf{x}-\mathbf{y})$$
$$\left[\varphi(t,\mathbf{x}),\pi_\phi(t,\mathbf{y})\right]_{\mathrm{PB}} = \frac{1}{\sqrt{\gamma}}\delta^{(3)}(\mathbf{x}-\mathbf{y})$$

$$\delta h_{ij} = \left[h_{ij}, \int d^4x\sqrt{\gamma}\mathcal{H}_G^k\Big|_{1\text{st}}\xi_k\right]_{\text{PB}} = \left[h_{ij}, \int d^4x\sqrt{\gamma}\frac{1}{\kappa^2}\left(\Pi_{(0)}^{kl} + \kappa\pi^{kl}\right)2D_k\xi_l\right]_{\text{PB}} = \frac{1}{\kappa}\left(D_i\xi_j + D_j\xi_i\right)$$

$$\delta\varphi = \left[\varphi, \int d^4x\sqrt{\gamma}\mathcal{H}_G^k\Big|_{1\text{st}}\Big|_{1\text{st}}\xi_k\right]_{\text{PB}} = \left[\varphi, \int d^4x\sqrt{\gamma}\left(P_\phi + \pi_\phi\right)\partial^k\phi\xi_k\right]_{\text{PB}} = \partial^k\phi_0\xi_k$$

$$\delta\partial_i\varphi = \left[\partial_i\varphi, \int d^3x\sqrt{\gamma}\mathcal{H}_G^k\Big|_{1\text{st}}\xi_k\right]_{\text{PB}} = \xi^k\partial_k\partial_i\phi_0 + \partial_k\phi_0\partial_i\xi^k$$

$$V_i = V_{(0)_i} + v_i$$

$$\delta v_i = \xi^k\partial_k V_{(0)i} + V_{(0)k}\partial_i\xi^k$$

$$\xi^i = \epsilon(t,\mathbf{x})\delta_x^i$$

$$\xi_i = \gamma_{(0)\ ix}\epsilon(t,\mathbf{x})$$

$$\kappa\delta h_{ij} = D_i\xi_j + D_j\xi_i = \gamma_{(0)_{jx}}\partial_i\epsilon + \gamma_{(0)}\partial_{ix}\partial_j\epsilon + \left[\delta_i^x\gamma'_{(0)\ jx} + \delta_j^x\gamma'_{(0)\ ix} - 2\Gamma_{ij}^k\gamma_{(0)\ kx}\right]\epsilon$$

$$\Gamma_{ij}^k = \frac{1}{2}\gamma_{(0)}^{kl}\left(\delta_i^x\gamma'_{(0)\ lj} + \delta_j^x\gamma'_{(0)\ li} - \delta_l^x\gamma'_{(0)\ ij}\right)$$

$$\kappa\delta h_{ij} = \gamma_{(0)_{jx}}\partial_i\epsilon + \gamma_{(0)\ ix}\partial_j\epsilon + \gamma'_{(0)_{ij}}\epsilon$$

$$\delta v_i = \epsilon V'_{(0)_i'} + V_{(0)}\partial_x\partial_i, \varphi = \phi_0'\epsilon$$

$$\kappa h_{ij} - \left(\frac{\gamma_{(0)_{ix}}}{V_{(0)_x}}v_j + \frac{\gamma_{(0)_{jx}}}{V_{(0)_x}}v_i\right) + \frac{1}{\phi_0'}\left(\gamma_{(0)_{ix}}\frac{V'_{(0)_j'}}{V_{(0)_x}} + \gamma_{(0)_{jx}}\frac{V'_{(0)_i'}}{V_{(0)_x}} - \gamma'_{(0)_{ij}}\right)\varphi$$

$$h_{ij} = h_{ij}^{TT} + \left(D_ih_j + D_jh_i - \frac{2}{3}\gamma_{(0)_{ij}}D_kh^k\right) + \frac{1}{3}\gamma_{(0)_{ij}}h,$$

$$\kappa\delta h_{ij} = D_i\xi_j + D_j\xi_i = \left(D_i\xi_j + D_j\xi_i - \frac{2}{3}\gamma_{(0)_{ij}}D_k\xi^k\right) + \frac{2}{3}\gamma_{(0)i_{ij}}D_k\xi^k$$

$$\kappa\delta h_i = \xi_i = \gamma_{(0)_{ix}}\epsilon, \kappa\delta h = 2D_i\xi^i = 2\partial_x\epsilon + \gamma_{(0)}^{\ ij}\gamma_{(0)\ ij}'\epsilon$$

$\gamma_{(0)_{ij}} = a(x)^2\delta_{ij}$ gives $\kappa\delta h = 2\partial_x\epsilon + 2(a'/a)\epsilon$

$$\epsilon(\mathbf{x}) = \frac{\epsilon_0(y,z)}{\sqrt{\gamma(x)}}$$

$\gamma_{(0)\ ij} = a(x)^2\delta_{ij}$, this corresponds to $\epsilon = \epsilon_0(y,z)/a(x)^3$

$$D^i\left(D_i\xi_j + D_j\xi_i\right) = \left[\partial_k\partial_k\epsilon + 5\frac{a'}{a}\partial_x\epsilon + 3\left(\frac{a'}{a}\right)^2\epsilon + 3\frac{a''}{a}\epsilon\right]\delta_{jx} + \partial_x\partial_j\epsilon + 3\frac{a'}{a}\partial_j\epsilon$$

$$D^i(D_i\xi_x + D_x\xi_i) = -9\left(\frac{a'}{a}\right)^2\frac{\epsilon_0}{a^3} - 3\frac{a''}{a}\frac{\epsilon_0}{a^3} + \frac{\partial_y\epsilon_0 + \partial_z\epsilon_0}{a^3}$$

$$D^i\left(D_i\xi_j + D_j\xi_i\right) = 0\ (j \neq x)$$

$$D^i\left(D_i\xi_j + D_j\xi_i\right) = 0 \text{ for } a = a_0(x - x_0)^{1/4}$$

$$\mathcal{V}_i = h_i - \frac{\gamma_{(0)ix}}{\kappa\phi_0'}\varphi$$

$$\mathcal{S} = h - \frac{2}{\kappa V_{(0)}}v_x + \frac{1}{\kappa\phi_0'}\left(\frac{2V'_{(0)'_x}}{V_{(0)_x}} - \gamma^{ij}_{(0)}\gamma'_{(0)'_{ij}}\right)\varphi$$

$$O\left(t, x - \kappa\gamma^{ix}_{(0)}h_i, y, z\right) - O_0(x) \simeq o(t, \mathbf{x}) - \kappa O_0'\gamma^{ix}_{(0)}h_i$$

$$x^i - \kappa\gamma^{ji}_{(0)}h_j$$

$$x + \frac{1}{\frac{2V'_{(0)}}{V_{(0)}} - \gamma^{ij}_{(0)}\gamma_{(0)\ ij}'}\left(\kappa h - \frac{2}{V_{(0)}}v_x\right)$$

$$O_{\text{island}} = \int d^4x f(x)O(x)$$

$$\sqrt{\delta(\phi_0(t)^2)} = \left(\frac{H}{2\pi}\right)(H\Delta t)^{1/2}$$

$$\gamma_{ij} = a(t)^2\left(\delta_{ij} + 2\kappa\zeta\delta_{ij}\right)$$

$$H\delta t = \kappa\delta\mathcal{R} = \frac{\kappa}{\sqrt{2\epsilon_H}}\frac{H}{2\pi}(H\Delta t)^{1/2}$$

$$-\frac{dM}{dt} \sim AT^4 = (16\pi G^2M^2)\left(\frac{1}{8\pi GM}\right)^4 \sim \frac{1}{\kappa^4M^2}$$

$$\dot{R} = -c^2\frac{\kappa^2}{R^2},\ \text{or}\ R = R_0\left(1 - c^2\frac{3\kappa^2}{R_0^3}t\right)^{1/3}$$

$$\frac{\delta t}{R} \sim \frac{1}{|\dot{R}|^{n/2}}\frac{\kappa}{R}\left(\frac{\Delta t}{R}\right)^{1/2}$$

$$\frac{\delta t}{R} \sim \kappa^{1-n}R^{n-\frac{3}{2}}t^{1/2} = \kappa^{1-n}R_0^{n-\frac{3}{2}}\left(1 - c^2\frac{3\kappa^2}{R_0^3}t\right)^{\frac{1}{3}\left(n-\frac{3}{2}\right)}t^{1/2}$$

$$R = R_0\left(1 - c^2(m+1)\frac{\kappa^m}{R_0^{m+1}}t\right)^{\frac{1}{m+1}}$$

$$\frac{\delta t}{R} \sim \kappa^{1-\frac{nm}{2}}R_0^{\frac{1}{2}(nm-3)}\left(1 - c^2(m+1)\frac{\kappa^m}{R_0^{m+1}}t\right)^{\frac{1}{2(m+1)}(nm-3)}$$

$$R^{d-3} = \frac{16\pi GM}{(d-2)\Omega_{d-2}}$$

$$\dot{R} = -c^2 \frac{\kappa^{d-2}}{R^{d-2}}$$

$$P_{ij} = \delta_{ij} - n_i n_j \text{ with } n_i = k_i/(k_j k^j)$$

$$h_{ij}^{TT} = P_i{}^l P_j{}^m h_{lm} - \frac{1}{2} P_{ij}(P^{lm} h_{lm})$$

$$h_{ij} \to h_{ij} + \partial_i \xi_j + \partial_j \xi_i$$

$$\begin{aligned} sA &= -Dc + \psi \ , \\ sc &= -c^2 + \phi \ , \\ s\psi &= -D\phi - [c, \psi] \ , \\ s\phi &= -[c, \phi] \ , \\ sF &= -D\psi - [c, F] \ . \end{aligned}$$

$$\mathcal{O}_n = \mathrm{Tr}\left(\tilde{F}^n\right) \ ; \ n \in \mathbb{N}^{\geq 1}$$

$$\mathcal{O}_2 = \mathrm{Tr}\left[F^2 + 2\psi F + \left(2\phi F + \psi^2\right) + 2\psi\phi + \phi^2\right]$$

$$S_{\mathrm{TYM}}[A] \equiv \int \mathrm{Tr}\left(F^2\right) = 8\pi^2 k$$

$$\lim_{r\to\infty} \mathbb{E}_{\mathrm{TYM}}[A; r] = \int_{S^2_\infty} \mathrm{Tr}\left[2\,(\lambda \rfloor A)\, F\right]$$

$$\begin{aligned} s\Phi &= -[c, \Phi] + \eta \ , \\ s\eta &= -[c, \eta] \ , \\ s\xi &= -[c, \xi] + B \ , \\ sB &= -[c, B] \ . \end{aligned}$$

$$
\begin{aligned}
S_{\mathrm{aFH}}\left[\Xi\right] &\equiv s\int \mathrm{Tr}\left\{D\xi\star D\Phi+\xi\Phi\left[m^2+g\left(B\Phi-\xi\eta\right)\right]\star\mathbb{1}\right\}\ ,\\
&=\int \mathrm{Tr}\left\{-DB\star D\Phi-D\xi\star D\eta+\left(\mathrm{ad}_\Phi D\xi-\mathrm{ad}_\xi D\Phi\right)\star\psi+\right.\\
&\left.+\left[m^2\left(B\Phi-\xi\eta\right)+g(B\Phi-\xi\eta)^2\right]\star\mathbb{1}\right\}\ ,
\end{aligned}
$$

$$
\begin{aligned}
sA&=\psi\ , & s\Phi&=\eta\ ,\\
s\psi&=-D\phi\ , & s\eta&=0\ ,\\
s\phi&=0\ , & s\xi&=B\ ,\\
sF&=-D\psi & sB&=0\ .
\end{aligned}
$$

$$
V_{\mathrm{aFH}}\left(\Phi,B\right)=\mathrm{Tr}\left[m^2B\Phi+g(B\Phi)^2\right]
$$

$$
\begin{aligned}
B_0&\equiv\mathrm{diag}\left(b_1,\ldots,b_N\right)\ ,\\
\Phi_0&\equiv\mathrm{diag}\left(\varphi_1,\ldots,\varphi_N\right)\ ,
\end{aligned}
$$

$$
\begin{aligned}
m^2b_j+2gb_j^2\varphi_j+\alpha&=0\ ,\\
m^2\varphi_j+2gb_j\varphi_j^2+\beta&=0\ ,\\
\textstyle\sum_j b_j&=0\ ,\\
\textstyle\sum_j \varphi_j&=0\ ,\\
\Delta_k&>0\ .
\end{aligned}
$$

$$
\begin{aligned}
\bar{S}_{\mathrm{aFH}}\left[\bar{\Xi}\right] &=\bar{s}\int \mathrm{Tr}\left\{D\xi\star D\varphi+\mathrm{ad}_{\Phi_0}D\xi\star A+\xi(\Phi_0+\varphi)\left[m^2+g(B_0\Phi_0+\right.\right.\\
&\left.\left.+B_0\varphi+b\Phi_0+b\varphi-\xi\eta)\right]\star\mathbb{1}\right\}\ ,\\
&=\int \mathrm{Tr}\left\{\mathrm{ad}_{B_0}\mathrm{ad}_{\Phi_0}A\star A+\left(\mathrm{ad}_{B_0}\varphi+\mathrm{ad}_{\Phi_0}b\right)D\star A+\mathrm{ad}_{\Phi_0}\xi D\star\psi+\right.\\
&-Db\star D\varphi-D\xi\star D\eta-\left(\mathrm{ad}_{B_0}\mathrm{ad}_A\varphi+\mathrm{ad}_{\Phi_0}\mathrm{ad}_Ab\right)\star A+\left(\mathrm{ad}_\varphi D\xi+\right.\\
&\left.-\mathrm{ad}_\xi D\varphi+2\mathrm{ad}_{\Phi_0}\mathrm{ad}_A\xi\right)\star\psi+\left[(m^2+2gB_0\Phi_0)(b\varphi-\xi\eta)+g\left(B_0\varphi+\right.\right.\\
&\left.\left.\left.+b\Phi_0+b\varphi-\xi\eta\right)^2\right]\star\mathbb{1}\right\}\ ,
\end{aligned}
$$

$$
\begin{aligned}
\bar{S}^{(2)}_{\mathrm{aFH}}\left[\bar{\Xi}\right] &=\int \mathrm{Tr}\left\{\mathrm{ad}_{B_0}\mathrm{ad}_{\Phi_0}A\star A+\left(\mathrm{ad}_{B_0}\varphi+\mathrm{ad}_{\Phi_0}b\right)d\star A+\mathrm{ad}_{\Phi_0}\xi d\star\psi+\right.\\
&\left.-db\star d\varphi-d\xi\star d\eta+\left[(m^2+2gB_0\Phi_0)\left(b\varphi-\xi\eta\right)+g(B_0\varphi+b\Phi_0)^2\right]\star\mathbb{1}\right\}
\end{aligned}
$$

$$
\begin{aligned}
&\bar{s}A = \psi \ , && \bar{s}\varphi = \eta \ , \\
&\bar{s}\psi = -D\phi \ , && \bar{s}\eta = 0 \ , \\
&\bar{s}\phi = 0 \ , && \bar{s}\xi = B_0 + b \ , \\
&\bar{s}F = -D\psi && \bar{s}b = 0 \ ,
\end{aligned}
$$

$$
\begin{aligned}
\mathbb{E}_{\mathrm{aFH}}\left[\Xi; r\right] &= \int_{B^3_r} \mathrm{Tr}\left\{\mathcal{L}_\lambda c \star (\mathrm{ad}_\Phi D\xi - \mathrm{ad}_\xi D\Phi) + s\left[\mathcal{L}_\lambda \Phi \star D\xi + D\xi(\lambda\rfloor \star D\Phi) + \right.\right. \\
&+ \left.\left. \mathrm{ad}_\xi \star D\Phi(\lambda\rfloor A) + \xi\Phi[m^2 + g(B\Phi - \xi\eta)](\lambda\rfloor \star \mathbb{1})\right]\right\} \\
&= \int_{B^3_r} \mathrm{Tr}\left[\mathcal{L}_\lambda B \star D\Phi - \mathcal{L}_\lambda \Phi \star DB + \mathcal{L}_\lambda \eta \star D\xi - \mathcal{L}_\lambda \xi \star D\eta + \right. \\
&+ \left. (\mathrm{ad}_\Phi \mathcal{L}_\lambda \xi + \mathrm{ad}_\xi \mathcal{L}_\lambda \Phi) \star \psi - (\lambda\rfloor \mathfrak{L}_{\mathrm{aFH}})\right] \ ,
\end{aligned}
$$

$$
\begin{aligned}
&s\Phi = -c\Phi + \eta \ , && s\Phi^\dagger = \Phi^\dagger c + \eta^\dagger \ , \\
&s\eta = -c\eta \ , && s\eta^\dagger = -\eta^\dagger c \ , \\
&s\xi = -c\xi + B \ , && s\xi^\dagger = -\xi^\dagger c + B^\dagger \ , \\
&sB = -cB \ , && sB^\dagger = B^\dagger c \ ,
\end{aligned}
$$

$$
\begin{aligned}
S_{\mathrm{FH}}\left[\Xi\right] &\equiv s\int \mathfrak{Re}\left\{(D\xi)^\dagger \star D\Phi + \xi^\dagger\Phi\left[m^2 + g(B^\dagger\Phi - \xi^\dagger\eta)\right] \star \mathbb{1}\right\} \ , \\
&= \int \mathfrak{Re}\left\{ - (DB)^\dagger \star D\Phi - (D\xi)^\dagger \star D\eta + (D\Phi)^\dagger \star \psi\xi + (D\xi)^\dagger \star \psi\Phi + \right. \\
&+ \left.\left[m^2(B^\dagger\Phi - \xi^\dagger\eta) + g(B^\dagger\Phi - \xi^\dagger\eta)^2\right] \star \mathbb{1}\right\} \ ,
\end{aligned}
$$

$$
V_{\mathrm{FH}}(\Phi, B, \Phi^\dagger, B^\dagger) = \mathfrak{Re}\left[m^2 B^\dagger\Phi + g(B^\dagger\Phi)^2\right]
$$

$$
\begin{aligned}
\mathfrak{Re}(B^\dagger\Phi) &= -m^2/2g \ , \\
\mathfrak{Im}(B^\dagger\Phi) &= 0 \ ,
\end{aligned}
$$

$$
Z_{\mathbb{R}} = \begin{pmatrix} \mathfrak{Re}(Z) & -\mathfrak{Im}(Z) \\ \mathfrak{Im}(Z) & \mathfrak{Re}(Z) \end{pmatrix} \in \mathrm{End}(\mathbb{C}^N_{\mathbb{R}})
$$

$$
S_{\mathrm{TYM}_{\mathbb{R}}}[A_{\mathbb{R}}] \equiv \int \mathrm{Tr}_{\mathbb{R}}(F^2_{\mathbb{R}}) = -8\pi^2 p_1
$$

$$
\begin{aligned}
sA &= -Dc + \psi \ , & s\Phi &= -c\Phi + \eta \ , & s\Phi^T &= \Phi^T c + \eta^T \ , \\
sc &= -c^2 + \phi \ , & s\eta &= -c\eta \ , & s\eta^T &= -\eta^T c \ , \\
s\psi &= -D\phi - [c, \phi] \ , & s\xi &= -c\xi + B \ , & s\xi^T &= -\xi^T c + B^T \ , \\
s\phi &= -[c, \phi] \ , & sB &= -cB \ , & sB^T &= B^T c \ ,
\end{aligned}
$$

$$
\begin{aligned}
S_{\mathrm{FH}_{\mathbb{R}}}[\Xi_{\mathbb{R}}] &= s \int \left\{ (D\xi)^T \star D\Phi + \xi^T \Phi \left[m^2 + g(B^T\Phi - \xi^T\eta) \right] \star \mathbb{1} \right\} \ , \\
&= \int \left\{ -(DB)^T \star D\Phi - (D\xi)^T \star D\eta + (D\Phi)^T \star \psi\xi + (D\xi)^T \star \psi\Phi + \right. \\
&\left. + \left[m^2(B^T\Phi - \xi^T\eta) + g(B^T\Phi - \xi^T\eta)^2 \right] \star \mathbb{1} \right\} \ ,
\end{aligned}
$$

$$
V_{\mathrm{FH}_{\mathbb{R}}}(\Phi, B^T) = m^2 B^T \Phi + g(B^T\Phi)^2
$$

$$
\begin{aligned}
\bar{S}_{\mathrm{FH}_{\mathbb{R}}}\left[\bar{\Xi}_{\mathbb{R}}\right] = \int \Big[& {B_0}^T A \star A\Phi_0 - {B_0}^T D \star A\varphi - {\Phi_0}^T D \star Ab + {\Phi_0}^T D \star \psi\xi + \\
& - (Db)^T \star D\varphi - (D\xi)^T \star D\eta + {B_0}^T A \star A\varphi + {\Phi_0}^T A \star Ab + \\
& + (D\varphi)^T \star \psi\xi + (D\xi)^T \star \psi\varphi - 2{\Phi_0}^T A \star \psi\xi + \\
& + g({B_0}^T\varphi + b^T\Phi_0 + b^T\varphi - \xi^T\eta)^2 \star \mathbb{1} \Big] \ ,
\end{aligned}
$$

$$
\begin{aligned}
\bar{S}^{(2)}_{\mathrm{FH}_{\mathbb{R}}}\left[\bar{\Xi}_{\mathbb{R}}\right] = \int \Big[& {B_0}^T A \star A\Phi_0 - {B_0}^T d \star A\varphi - {\Phi_0}^T d \star Ab + {\Phi_0}^T d \star \psi\xi + \\
& - db^T \star d\varphi - d\xi^T \star d\eta + g({B_0}^T\varphi + b^T\Phi_0)^2 \star \mathbb{1} \Big] \ ,
\end{aligned}
$$

$$
\bar{S}_{\mathrm{FH}_{\mathbb{R}}}[\bar{\Xi}_{\mathbb{R}}] \equiv S_{\mathrm{FH}_{\mathbb{R}}}[A, \psi, \Phi_0 + \varphi, \eta, \xi, b, \Phi_0^T + \varphi^T, \eta^T, \xi^T, B_0^T + b^T]
$$

$$
S_n(\rho) = \frac{1}{1-n} \log\left(\frac{\mathrm{tr}[\rho^n]}{(\mathrm{tr}[\rho])^n} \right)
$$

$$
L^2(G) = \mathrm{span}\{|g\rangle \mid g \in G\}.
$$

$$
\langle g \mid h \rangle = |G| \delta_{gh}.
$$

$$
\langle g \mid h \rangle = \delta(g^{-1}h).
$$

$$
\frac{1}{|G|} \sum_{g \in G} = \int_G dg
$$

$$L^2(G) = \bigoplus_{\pi \in \hat{G}} d_\pi \cdot [V_\pi \otimes V_\pi^*]$$

$$d_\pi = \dim(V_\pi)$$

$$\langle A \mid B \rangle_{d_\pi \cdot [V_\pi \otimes V_\pi^*]} := d_\pi \cdot \langle A \mid B \rangle_{V_\pi \otimes V_\pi^*}$$

$$L^2(G) = \int_{\hat{G}}^{\oplus} d\mu(\pi) V_\pi \otimes V_\pi^* \oslash \langle A \mid B \rangle_{V_\pi \otimes V_\pi^*}$$

$$d\mu(\pi) = \frac{d_\pi}{\mathrm{Vol}(G)}$$

$$\mathcal{H}(\Lambda) = \bigotimes_{\ell \in L} \mathcal{H}_\ell$$

$$B_{(v,p)}(h) \left| g_1, \cdots, g_{|L|} \right\rangle = \delta\left(h^{-1} g_\rho\right) \left| g_1, \cdots, g_{|L|} \right\rangle.$$

$$\left\langle g_1, \cdots, g_{|L|} \mid \psi \right\rangle = \psi\left(g_1, \cdots, g_{|L|}\right),$$

$$B_{(v,p)}(h) |\psi\rangle = \int dg_1 \cdots dg_{|L|} \delta\left(h^{-1} g_\rho\right) \psi\left(g_1, \cdots, g_{|L|}\right) \left| g_1, \cdots, g_{|L|} \right\rangle$$

$$B_{(v,p)}[f] := \int dg f(g) B_{(v,p)}(g)$$

$$\|f\|_\infty = \sup_{g \in G} |f(g)|$$

$$\begin{aligned} \|B_{(v,p)}[f] |\psi\rangle\| &= \| \int dg_1 \cdots dg_{|L|} f\left(g_\rho\right) \psi\left(g_1, \cdots, g_{|L|}\right) \left| g_1, \cdots, g_{|L|} \right\rangle \| \\ &\leq \|f\|_\infty \cdot \| |\psi\rangle \| \end{aligned}$$

$$\Pi_B = \bigotimes_{(v,p) \in S} B_{(v,p)}(e)$$

$$B_{(v,p)}(e) B_{(v,p)}(e) = \delta(e) B_{(v,p)}(e)$$

$$\Pi_B \mathcal{H}(\Lambda) = \mathrm{span}\left\{ \delta(e)^{-|S|/2} \Pi_B |\psi\rangle \text{ such that } |\psi\rangle \in \mathcal{H}(\Lambda) \right\}$$

$$\delta(e)^{-|S|/2} \Pi_B |\psi\rangle \text{ and } \delta(e)^{-|S|/2} \Pi_B |\sigma\rangle$$

$$\frac{1}{\delta(e)^{|S|}} \langle \sigma | \Pi_B^\dagger \Pi_B | \psi \rangle = \langle \sigma | \Pi_B | \psi \rangle$$

$$\Pi_B |\psi\rangle = \Pi_B (|\psi\rangle + |\chi\rangle)$$

$$[|\psi\rangle \sim |\psi\rangle + |\chi\rangle] \text{ for all } |\psi\rangle \in \mathcal{H}(\Lambda) \text{ and } |\chi\rangle \in \mathcal{H}_{null}$$

$$\left(\Pi_B \mathcal{H}(\Lambda)\right)_{\text{gravity}} = \mathcal{H}(\Lambda)_\infty / \mathcal{H}_{\text{null}}$$

$$\langle\langle \sigma \mid \psi \rangle\rangle := \langle \sigma | \Pi_B | \psi \rangle.$$

$$\Pi_B \mathcal{H}(\Lambda) = \overline{\Pi_B \mathcal{H}(\Lambda))_{gravity}}.$$

$$\langle\langle \sigma \mid \psi \rangle\rangle = \int d[g,h]\sigma(g)^* \psi(h) \langle g | B_{(v,p)}(e) | h \rangle$$
$$= \int d[g,h]\sigma(g)^* \psi(h)\delta(h)\langle g \mid h \rangle$$
$$= \sigma(e)^* \psi(e)$$

$$\psi_n(g) = (N_n)^{-\frac{1}{2}} \Theta_n(g).$$

$$\lim_{n,m\to\infty} \| |\psi_n\rangle - |\psi_m\rangle \|_{\mathcal{H}(\Lambda_1)} > C$$

$$\| |\psi_n\rangle\rangle - |\psi_m\rangle\rangle \mid \|_{\Pi_B \mathcal{H}(\Lambda_1)} = 0$$

$$|\psi_\infty\rangle \lim_{n\to\infty} |\psi_n\rangle \Pi_B \mathcal{H}(\Lambda_1) |\psi_\infty\rangle = |\psi_1\rangle \Pi_B \mathcal{H}(\Lambda_1)$$

$$\tilde{B}_{(v,p)}(e) := \lim_{n\to\infty} B_{(v,p)}[\Theta_n]$$

$$\tilde{B}_{(v,p)}(e)\tilde{B}_{(v,p)}(e) = \lim_{n\to\infty} \int_{V_n(e)} \int_{V_n(e)} dg dh B_s(g) B_s(h)$$
$$= \lim_{n\to\infty} \int_{V_n(e)} \int_{V_n(e)} dg dh \delta(g^{-1}h) B_s(g)$$
$$= \lim_{n\to\infty} \int_{V_n(e)} dg B_s(g) \left(\int_{V_n(g)} dh \delta(h) \right)$$
$$= \lim_{n\to\infty} \int_{V_n(e)} dg B_s(g) = \tilde{B}_{(v,p)}(e)$$

$$\tilde{B}_{(v,p)}(e)^\dagger \tilde{B}_{(v,p)}(e) = \tilde{B}_{(v,p)}(e)$$

$$\tilde{B}_{(v,p)}(e)|\psi\rangle = \lim_{n\to\infty} \int_{V_n(e)} dg \psi(g) |g\rangle.$$

$$\langle \psi | \tilde{B}_{(v,p)}(e)^\dagger \tilde{B}_{(v,p)}(e) | \psi \rangle = \lim_{n\to\infty} \int_{V_n(e)} dg \psi^*(g)\psi(g)$$

$$|\psi(g)|^2 = |\tilde{\psi}(e)|^2 \delta(g) + \cdots$$

$$A_v[f] = \int dh f(h) A_v(h)$$

$$A_v[1] = \int dh A_v(h)$$

$$A_v(g)A_v[1] = \int dh A_v(g) A_v(h)$$
$$= \int dh A_v(gh)$$
$$= A_v[1]$$

$$\Pi_A = \bigotimes_{v \in V} A_v[1]$$

$$\Pi_A \mathcal{H}(\Lambda) = \left\{ \mathrm{Vol}(G)^{-|L|/2} \Pi_A |\psi\rangle \text{ such that } |\psi\rangle \in \mathcal{H}(\Lambda) \right\}$$

$$[|\psi\rangle \sim |\psi\rangle + |\chi\rangle] \text{ for all } |\psi\rangle \in \mathcal{H}(\Lambda) \text{ and } |\chi\rangle \in \mathcal{H}_{null}.$$

$$(\Pi_A \mathcal{H}(\Lambda))_{\text{gravity}} = \mathcal{H}(\Lambda)_1 / \mathcal{H}_{\text{null}}$$

$$\langle\langle \sigma \mid \psi \rangle\rangle = \langle \sigma | \Pi_A | \psi \rangle$$

$$\Pi_A \mathcal{H}(\Lambda) = \overline{\left(\Pi_A \mathcal{H}(\Lambda)\right)_{gravity}}$$

$$\langle\langle \sigma \mid \psi \rangle\rangle = \int d[g,h,k] \sigma^*(g) \psi(h) \langle g | A_v(k) | h \rangle$$
$$= \int d[g,h,k] \sigma^*(g) \psi(h) \langle g \mid kh \rangle$$
$$= \int d[h,k] \sigma^*(kh) \psi(h)$$
$$= \left(\int dk \sigma(k) \right)^* \left(\int dh \psi(h) \right)$$

$$\langle\langle \sigma \mid \psi \rangle\rangle = \left(\int dx \sigma(x) \right)^* \left(\int dx \psi(x) \right)$$

$$\mathcal{H}_{phys}(\Sigma) := \Pi_A \Pi_B \mathcal{H}(\Lambda)$$

$$\mathcal{H}(\Lambda) = \bigotimes_{\ell \in L} \mathcal{H}_\ell \bigotimes_{(v,p) \in S} \mathcal{H}_{(v,p)}^{matt}$$

$$A_v(h) \to A_v(h) A_v^{matt}(h)$$
$$B_{(v,p)}(h) \to \int dg B_{(v,p)}(h g^{-1}) B_{(v,p)}^{matt}(g)$$

$$\langle\langle \sigma \mid \psi \rangle\rangle_{\text{phys}} = \langle \sigma | \Pi_A \Pi_B | \psi \rangle$$
$$:= \int \prod_{v \in V} dg_v \langle \sigma | \left[\prod_{v \in V} A_v(g_v) \right] \left[\prod_{(v,p) \in S} B_{(v,p)}(e) \right] | \psi \rangle,$$

$$\mathcal{H}_v = \mathcal{H}_v^{(1)} \otimes \cdots \otimes \mathcal{H}_v^{(n)} \otimes \mathcal{H}_v^{\text{matt}}$$

$$\mathcal{H}_\ell = \mathcal{H}_v^{(i)} \otimes \mathcal{H}_{v'}^{(j)}$$

$$\mathcal{H}_v^{(i)} \cong \mathcal{H}_{v'}^{(j)}$$

$$|\Psi\rangle = \left(\bigotimes_{\langle ij \rangle} \langle \chi, ij | \right) \bigotimes_{v \in V} |\psi_v\rangle$$


$$|\Psi\rangle = \left(\bigotimes_{v \in V} \Psi_v(\ell_1, \cdots, \ell_n) \right) \bigotimes_{\ell \in L} |\chi, \ell\rangle.$$


$$|e\rangle = \int d\mu(\pi) \sum_{m,n} \pi(e)_{mn} |\pi, mn\rangle = \int d\mu(\pi) \sum_{m} |\pi, mm\rangle = \int d\mu(\pi) |\chi_\pi\rangle$$


$$|\chi_\pi\rangle = \sum_{m} |\pi, mm\rangle$$

$$\Pi_A \Pi_B \mathcal{O} |\chi\rangle = \mathcal{O} \Pi_A \Pi_B |\chi\rangle = 0.$$

$\tilde{\mathcal{O}}|\psi\rangle\rangle = |\mathcal{O} \cdot \psi\rangle\rangle = [\mathcal{O}|\psi\rangle \sim \mathcal{O}|\psi\rangle + |\chi\rangle]$ for any $|\chi\rangle \in \mathcal{H}_{\text{null}}$.







$$\mathcal{H}_{\text{phys}}(\Sigma) = \Pi_A \Pi_B \mathcal{H}(\Lambda_b)$$

$$|\psi\rangle\rangle = \int d[\vec{g}_R, \vec{g}_{\bar{R}}, h]\psi(\vec{g}_R, \vec{g}_{\bar{R}}, h)|\vec{g}_{\bar{R}}, h, \vec{g}_R\rangle\Big\rangle$$

$$|\psi\rangle\rangle = \int d[g,h]\psi(g,h)|g,h\rangle\Big\rangle$$
$$= \int d[g,h]\psi(g,h)|h^{-1}g, e\rangle\Big\rangle$$
$$= \int d[g,h]\psi(hg,h)|g,e\rangle\Big\rangle.$$

$$\tilde{\psi}(g) = \int dh\psi(hg,h)$$

$$|\psi\rangle\rangle = \int dg\tilde{\psi}(g)|g,e\rangle\Big\rangle$$

$$F_R(\pi)|\psi\rangle\rangle = \int d[\vec{g}_R, \vec{g}_{\bar{R}}, h, k]\chi_\pi(k)\psi(\vec{g}_R, \vec{g}_{\bar{R}}, k^{-1}h)|\vec{g}_{\bar{R}}, h, \vec{g}_R\rangle\Big\rangle$$

$$A_R(\ell)|\vec{g}_{\bar{R}}, h, \vec{g}_R\rangle = |\vec{g}_{\bar{R}}, h\ell^{-1}, \ell \cdot \vec{g}_R\rangle$$
$$A_{\bar{R}}(\ell)|\vec{g}_{\bar{R}}, h, \vec{g}_R\rangle = |\ell \cdot \vec{g}_{\bar{R}}, \ell h, \vec{g}_R\rangle$$

$$A_R(\ell)F_{\cup R}(\pi)|\vec{g}_{\bar{R}}, h, \vec{g}_R\rangle\rangle = \int dk\chi_\pi(k)|\vec{g}_{\bar{R}}, k^{-1}h\ell^{-1}, \ell \cdot \vec{g}_R\rangle\Big\rangle$$
$$= \int dk\chi_\pi(k)|\vec{g}_{\bar{R}}, k^{-1}h, \vec{g}_R\rangle\Big\rangle$$
$$= F_{\cup R}(\pi)|\vec{g}_{\bar{R}}, h, \vec{g}_R\rangle\rangle$$

$$A_{\bar{R}}(\ell)F_{\cup R}(\pi)|\vec{g}_{\bar{R}}, h, \vec{g}_R\rangle\rangle, = \int dk\chi_\pi(k)|\ell \cdot \vec{g}_{\bar{R}}, \ell k^{-1}h, \vec{g}_R\rangle\Big\rangle$$
$$= \int dk\chi_\pi(\ell^{-1}k\ell)|\ell \cdot \vec{g}_{\bar{R}}, k^{-1}\ell h, \vec{g}_R\rangle\Big\rangle$$
$$= \int dk\chi_\pi(k)|\ell \cdot \vec{g}_{\bar{R}}, k\ell h, \vec{g}_R\rangle\Big\rangle$$
$$= F_{\cup R}(\pi)|\ell \cdot \vec{g}_{\bar{R}}, \ell h, \vec{g}_R\rangle\rangle$$
$$= F_{\cup R}(\pi)|\vec{g}_{\bar{R}}, h, \vec{g}_R\rangle\rangle$$

$$\chi_\pi(g) = \chi_\pi(hgh^{-1})$$

$$|\vec{g}_{\bar{R}}, h, \vec{g}_R\rangle\rangle\mathcal{H}_{\text{phys}}(\Sigma)\Pi_A\Pi_B\mathcal{H}(\Lambda_b)$$

$$F_{\cup R}(\pi)F_{\cup R}(\omega) = \delta(\pi,\omega)F_{\cup R}(\pi),$$


$$F_{\cup R}(f) = \int d\mu(\pi) f(\pi) F_{\cup R}(\pi)$$


$$\text{Area}_R = \int d\mu(\pi) \log\left(\frac{d\mu(\pi)}{d\pi}\right) F_R(\pi).$$

$$\frac{d\mu(\pi)}{d\pi} = \frac{d_\pi}{\text{Vol}(G)}$$

$$\text{Area}_R = \int d\mu(\pi) \log\left(\frac{d_\pi}{\text{Vol}(G)}\right) F_{\cup R}(\pi)$$

$$\frac{1}{|G|}\sum_{g\in G} \to \int dg \sum_{\pi\in\hat{G}} d_\pi \to \int d\mu(\pi)$$





$$\mathcal{H}_{\text{phys}}(\Sigma) = \Pi_A \Pi_B \mathcal{H}(\Lambda_r)$$

$$A_R(k)|\pi, ab\rangle|\vec{g}\rangle = \sum_c \pi(k)_{cb}|\pi, ac\rangle|k\cdot\vec{g}\rangle$$

$$d\mu(\vec{\pi}) = d\mu(\pi_1)\cdots d\mu(\pi_n)$$
$$V_{\vec{\pi}} = \bigotimes_{\pi_i\in\vec{\pi}} V_{\pi_i}|\pi, ab; \vec{g}\rangle\rangle$$

$$\mathcal{H}_{phys}(\Sigma) = \int_{\hat{G}}^{\oplus} d\mu(\vec{\pi}) V_{\vec{\pi}} \otimes \Pi_A\left[V_{\vec{\pi}}^*\right]$$

$$\mathcal{H}_{phys} = \bigoplus_{\Sigma} \mathcal{H}_{phys}(\Sigma)$$

$$\mathcal{H}_\partial = \Pi_A \Pi_B \mathcal{H}(\Lambda_r)$$

$$\mathcal{H}_{\text{phys}}(\Sigma) = \Pi_A \Pi_B \mathcal{H}\left(\Lambda_r^{(m)}\right)$$

$$|\psi\rangle = (\mathrm{Id}_\partial \otimes \langle T|)|\Psi\rangle$$

$$|\psi\rangle \in \mathcal{H}_\partial$$


$$|\Psi\rangle: \mathcal{H}(\mathrm{l}) \to \mathcal{H}_\partial.$$

$$\mathcal{H}_\ell = L^2(\mathrm{SL}(2,\mathbb{R})/\mathrm{U}(1))$$

$$V_\pi \otimes V_\pi^* \mapsto \int_{\mathrm{U}(1)} d\theta \pi(\theta) V_\pi \otimes V_\pi^* \pi(\theta)^\dagger$$

$$= \int_{\mathrm{U}(1)} d\theta (\pi(\theta) V_\pi) \otimes (\pi(\theta) V_\pi)^*$$

$$L^2(\mathrm{SL}(2,\mathbb{R})/\mathrm{U}(1)) = \int_{\text{principal}} d\mu(\lambda) V_\lambda \otimes V_\lambda^*$$



$|\Psi\rangle$ $\qquad$ $A(\vec{g})\,|\Psi\rangle = |\Psi\rangle$

$$A(\vec{g}) = \otimes_v A_v(g_v)$$

$$\lim_{k\to\infty} \mathrm{SU}(4)_k = \widehat{\mathrm{SU}}(4)$$

$$\lim_{\sigma\to\infty} G_\sigma \to \hat{G}$$

$$\{|\theta\rangle \mid \theta \in [0,2\pi)\} \oplus \{|n\rangle \mid n \in \mathbb{Z}\}$$

$$\langle \theta \mid n \rangle = e^{2\pi i n\theta}$$

$$E_n = \mathrm{span}\{|n\rangle\}$$

$$L^2(\mathrm{U}(1)) = \bigoplus_{n\in\hat{U}(1)} E_n$$

$$\langle g \mid j, mn \rangle = D^j_{mn}(g)$$

$$E_j = \mathrm{span}\{|j, mn\rangle \mid m, n \in 1, \cdots, 2j+1\}.$$

$$\langle j, mn \mid \ell, pq \rangle = \int_{\mathrm{SU}(2)} dg \left(D^{j}_{mn}(g)\right)^{*} D^{\ell}_{pq}(g) = \left(\frac{\mathrm{Vol}(\mathrm{SU}(2))}{d_j}\,\delta_{j\ell}\right)\delta_{mp}\delta_{nq}$$

$$\mathrm{Vol}(\mathrm{SU}(2)) = \int 1 dg$$

$$d_j = \dim\left(V_j\right)$$

$$G, D^{\ell}_{mn}(g)^{*} = D^{\ell}_{nm}(g^{-1})$$

$$\langle j, mn \mid \ell, pq \rangle = \int_{\mathrm{SU}(2)} dg \langle n|D^{j}(g^{-1})|m\rangle\langle p|D^{\ell}(g)|q\rangle$$

$$= \langle n| \left(\int_{\mathrm{SU}(2)} dg D^{j}(g^{-1})|m\rangle\langle p|D^{\ell}(g) \right)|q\rangle$$

$$= \langle n|O^{j\ell}_{mp}|q\rangle$$

$$D^{j}(g)O^{jl}_{mp} = O^{jl}_{mp}D^{\ell}(g)$$

$$O^{j\ell}_{mp} = \delta_{j\ell}c^{j}_{mp}\frac{\mathrm{Id}_{V_j}}{d_j}$$

$$c^{j}_{mp} = \mathrm{tr}\left(O^{jj}_{mp}\right) = \int_{\mathrm{SU}(2)} dg \langle p|D^{j}(g)D^{j}(g^{-1})|m\rangle = \mathrm{Vol}(\mathrm{SU}(2))\langle p \mid m\rangle$$

$$|j, mn\rangle \to \left(\frac{d_j}{\mathrm{Vol}(\mathrm{SU}(2))}\right)^{-1/2} |j, mn\rangle$$

$$\mu(j) = \frac{d_j}{\mathrm{Vol}(\mathrm{SU}(2))}$$

$$\langle j, mn \mid j, pq \rangle_{\mu(j)\cdot E_j} := \mu(j) \cdot \langle j, mn \mid j, pq \rangle_{E_j}$$

$$\langle j, mn \mid \ell, pq \rangle = \delta(j, \ell)\delta_{mp}\delta_{nq}$$

$$L^2(\mathrm{SU}(2)) = \bigoplus_{j \in \widehat{SU}(2)} \mu(j) \cdot E_j$$

$$L^2(\mathrm{SU}(2)) \supset \bigoplus_{j \in \widehat{SU}(2)} \mu(j) \cdot E_j$$

$$\mu(\pi) = \frac{\dim(V_\pi)}{\mathrm{Vol}(G)}$$

$$\langle g \mid \pi, ij \rangle = \pi_{ij}(g)$$

$$\langle \pi, ij \mid \omega, mn \rangle = \delta(\pi, \omega)\delta_{im}\delta_{nj}$$

$$L^2(G) = \bigoplus_{\pi \in \hat{G}} \mu(\pi) \cdot E_\pi$$

$$g(\theta, x_a, x_n) = \begin{pmatrix} \cos(x_k) & \sin(x_k) \\ -\sin(x_k) & \cos(x_k) \end{pmatrix} \begin{pmatrix} x_a & 0 \\ 0 & x_a^{-1} \end{pmatrix} \begin{pmatrix} 1 & x_n \\ 0 & 1 \end{pmatrix}.$$

$$dg = dk \frac{da}{a} dn$$

$$\chi_\pi(h^{-1}g) = \mathrm{tr}[\pi(h^{-1}g)] = \mathrm{tr}\left[\pi(h)^\dagger \pi(g)\right] = \langle \pi(h) \mid \pi(g) \rangle_{E_\pi}$$

$$\mathrm{tr}[\pi(e)] = \dim(V_\pi) = \infty$$

$$\pi(f) = \int dg f(g) \pi(g^{-1})$$

$$\hat{f}: \pi \mapsto \pi(f)$$

$$\chi_\pi(f) = \mathrm{tr}[\pi(f)]$$

$$\chi_\pi(f) = \int_G dg f(g) \chi_\pi(g^{-1})$$

$$\chi_\pi(g) = \mathrm{tr}[\pi(g)]$$

$$g: \chi_\pi(g) = \chi_\pi(hgh^{-1})$$

$$\lim_{n\to\infty} \chi_{\pi_n}(f) = \chi_\pi(f)$$

$$\chi_n(k_\theta) = -\mathrm{sign}(n) \frac{e^{in\theta}}{e^{i\theta} - e^{-i\theta}}$$

$$\pm a_t = \begin{pmatrix} \pm e^t & 0 \\ 0 & \pm e^{-t} \end{pmatrix}$$

$$\chi_n(\pm a_t) = (-1)^{1+|n|} \frac{e^{-|nt|}}{e^t - e^{-t}}$$

$$\chi_{\lambda,+}(\pm a_t) = \frac{e^{\lambda t} + e^{-\lambda t}}{|e^t - e^{-t}|}$$
$$\chi_{\lambda,-}(\pm a_t) = \pm \frac{e^{\lambda t} + e^{-\lambda t}}{|e^t - e^{-t}|}$$

$$f(g) = \int_{\hat{G}} d\mu(\pi) \mathrm{tr}_{V_\pi}[\pi(g)\pi(f)]$$

$$\int_G dg f'(g)^* f(g) = \int_{\hat{G}} d\mu(\pi) \mathrm{tr}_{V_\pi}\left[\pi(f')^\dagger \pi(f)\right]$$

$$\mathrm{tr}_{V_\pi}\left[\pi(f')^\dagger \pi(f)\right]$$

$$L^2(G) = \int_{\hat{G}} d\mu(\pi) V_\pi \otimes V_\pi^*$$

$$f(e) = \int_{\hat{G}} d\mu(\pi)\mathrm{tr}_{V_\pi}[\pi(f)]$$
$$= \int_{\hat{G}} d\mu(\pi)\chi_\pi(f)$$
$$= \int_{\hat{G}} d\mu(\pi) \int_G dg f(g)\chi_\pi(g^{-1})$$

$$\delta(g) = \int_{\hat{G}} d\mu(\pi)\chi_\pi(g^{-1})$$

$$\text{of } \delta_a(x) = \delta(x-a) \text{ is } e^{-ika}$$

$$\delta_h(g) = \delta(h^{-1}g)$$

$$\pi(\delta_h) = \pi(h^{-1}),$$

$$d\mu(n,v) = (n^2+v^2)dv \iiint \blacksquare \mathbf{SL}(4,\mathbb{R})$$

$$d\mu(n) = |n|$$

$$d\mu(\lambda,+) = \frac{1}{2}\lambda\tanh(\pi\lambda/2)d\lambda, \text{ and } d\mu(\lambda,-) = \frac{1}{2}\lambda\mathrm{cotanh}(\pi\lambda/2)d\lambda$$

$$|\vec{\pi}, \vec{a}\vec{b}\rangle = |\pi_1, a_1b_1\rangle \cdots |\pi_n, a_nb_n\rangle$$
$$|\vec{g}\rangle = |g_1\rangle \cdots |g_n\rangle$$
$$\vec{\pi}(\vec{g})_L = \pi_1(g_1) \otimes \cdots \otimes \pi_n(g_n)$$
$$\vec{\pi}(\vec{g})_R = \pi_1(g_1)^\dagger \otimes \cdots \otimes \pi_n(g_n)^\dagger$$

$$A_v(h)|\vec{\pi}, \vec{a}\vec{b}\rangle = \vec{\pi}(h)_R|\vec{\pi}, \vec{a}\vec{b}\rangle = \sum_{\vec{c}} \langle\vec{c}|\vec{\pi}(h)_R|\vec{b}\rangle|\vec{\pi}, \vec{a}\vec{c}\rangle$$

$$\Pi_A = A_v[1] = \int dh A_v(h)$$

$$\vec{\pi}[1]_R = \int dh \vec{\pi}(h)_R$$

$$\Pi_A|\vec{\pi}, \vec{a}\vec{b}\rangle = \vec{\pi}[1]_R|\vec{\pi}, \vec{a}\vec{b}\rangle$$

$$V_{\vec{\pi}} \otimes V_{\vec{\pi}}^*$$

$$I_{\pi,\omega}\omega(g) = \pi(g)I_{\pi,\omega}$$

$$\mathcal{I}_{\pi,\omega} = \{I_{\pi,\omega} \mid \forall g \in G, I_{\pi,\omega}\omega(g) = \pi(g)I_{\pi,\omega}\}$$

$$\dim(\mathcal{I}_{\pi,\omega}) = \begin{cases} 1 & \pi \cong \omega \\ 0 & \text{else} \end{cases}$$

$$C_{j_1j_2}^{j}: V_{j_1} \otimes V_{j_2} \to V_j$$

$$\vec{\pi}[1]_R\vec{\pi}(g)_R = \int dh\vec{\pi}(h)_R\vec{\pi}(g)_R = \int dh\vec{\pi}(hg)_R = \int dh\vec{\pi}(h)_R = \mathrm{Id}_{V_{\vec{\pi}}^*}\vec{\pi}[1]_R$$

$$Q(|\psi\rangle, |\sigma\rangle)\colon V_{\vec{\pi}}^* \otimes V_{\vec{\pi}}^* \to \mathbb{C}$$

$$Q(|\psi\rangle, |\sigma\rangle) = \langle\psi|\vec{\pi}[1]_R|\sigma\rangle$$

$$I\colon V_{\vec{\pi}} \to V_{\vec{\omega}}$$

$$|I\rangle \in V_{\vec{\pi}}^* \otimes V_{\vec{\omega}}$$

$$\vec{\pi}[1]_R\colon V_{\vec{\pi}} \to \mathbb{C}$$

$$|\vec{\pi}_R\rangle \in V_{\vec{\pi}}$$

$$\vec{\pi}(g)_R|\vec{\pi}_R\rangle = |\vec{\pi}_R\rangle$$

$$\pi_1(g^{-1}) \otimes \mathrm{Id}_{\pi_2} \otimes \mathrm{Id}_{\pi_n}$$

$$\mathrm{Id}_{\pi_1} \otimes \pi_2(g) \otimes \cdots \otimes \pi_n(g)|\vec{\pi}_R\rangle = \pi_1(g^{-1}) \otimes \mathrm{Id}_{\pi_2} \otimes \cdots \otimes \mathrm{Id}_{\pi_n}|\vec{\pi}_R\rangle$$

$$|\mathrm{Id}_\pi\rangle = \sum_m |\pi^*, n\rangle|\pi, n\rangle$$

$$|\pi_1, \pi_2, \pi_3\rangle \in V_{\pi_1} \otimes V_{\pi_2} \otimes V_{\pi_3}$$

$$|\pi_1, \pi_2, \pi_3\rangle = \sum_{m_1,m_2,m_3} \begin{pmatrix} \pi_1 & \pi_2 & \pi_3 \\ m_1 & m_2 & m_3 \end{pmatrix} |\pi_1, m_1\rangle|\pi_2, m_2\rangle|\pi_3, m_3\rangle$$

$$|\pi_1, \pi_2, \pi_3\rangle\Pi_A\left[V_{\vec{\pi}}^*\right]$$

$$\int d\mu(\pi_1) \sum_{m_1} \begin{pmatrix} \pi_1 & \pi_2 & \pi_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \begin{pmatrix} \pi_1 & \pi_2 & \pi_3 \\ m_1 & n_2 & n_3 \end{pmatrix} = \delta_{m_2,n_2}\delta_{m_3,n_3}.$$

$$\rho_{\pi_2\pi_3} = \mathrm{tr}_{\pi_1}[|\pi_1, \pi_2, \pi_3\rangle\langle\pi_1, \pi_2, \pi_3|] \text{ on } V_{\pi_2} \otimes V_{\pi_3}$$

$$\mathrm{Id}_{L^2(G)} = \int_G dg|g\rangle\langle g| = \int_{\hat{G}} d\mu(\pi) \sum_{a,b} |\pi, ab\rangle\langle\pi, ab|$$

$$\delta(g) = \int d\mu(\pi)\chi_\pi(g^{-1})$$

$$\langle\pi, ab \mid \omega, mn\rangle = \delta(\pi, \omega)\delta_{am}\delta_{bn}$$

$$\chi_\pi(g^{-1}) = \chi_\pi(g)^*$$

$$\chi_{\pi\oplus\omega}(g) = \chi_\pi(g) + \chi_\pi(g)$$

$$\chi_{\pi\otimes\omega}(g) = \chi_\pi(g) \cdot \chi_\pi(g)$$


$$\int_G dg\chi_\omega(g^{-1})\chi_\pi(gh) = \delta(\pi, \omega)\chi_\pi(h)$$

$$\int_G dg\chi_\omega(g^{-1})\chi_\pi(gh) = \langle\chi_\omega|\pi(h^{-1})_R|\chi_\pi\rangle$$

$$V_\pi \otimes V_\pi^* \subset L^2(G)$$

$$V_\pi \otimes V_\pi^* \otimes V_\omega \otimes V_\omega^* \text{ for } \pi \neq \omega$$

$$\int_{\hat{G}} d\mu(\omega) \int_G dg \chi_\omega(g^{-1}) \chi_\pi(gh) = \int_G dg \int_{\hat{G}} d\mu(\omega) \chi_\omega(g^{-1}) \chi_\pi(gh)$$
$$= \int_G dg \delta(g) \chi_\pi(gh)$$
$$= \chi_\pi(h)$$



$$\psi(\vec{g}) = \psi(g_1, \cdots, g_{|L|}),$$
$$|\vec{g}\rangle = |g_1, \cdots, g_{|L|}\rangle,$$
$$d[\vec{g}] = dg_1 \cdots dg_{|L|}.$$

$$|\Psi\rangle = \int d[\vec{g}] \psi(\vec{g}) |\vec{g}\rangle$$

$$\Delta_1 |\Psi\rangle = [\Pi_A]_v [\Pi_A]_{v'} |e\rangle |\Psi\rangle.$$

$$\Delta_1 |\Psi\rangle = \int d[h, k, \vec{g}] \psi(\vec{g}) |hk^{-1}\rangle |h \cdot \vec{g} \cdot k^{-1}\rangle$$

$$\Delta_1 |\Psi\rangle\rangle = [|e\rangle |\Psi\rangle \sim |e\rangle |\Psi\rangle + |\chi\rangle] \equiv |e, \Psi\rangle\rangle$$

$$\left\langle \langle \psi | \Delta_1^\dagger \Delta_1 \mid \sigma \rangle \right\rangle = \langle e \mid \langle \langle \psi | [\Pi_A]_v [\Pi_A]_{v'} \mid e \rangle \mid \sigma \rangle \rangle$$
$$= \int d[h, k, \vec{g}, \vec{\ell}] \psi^*(\vec{g}) \sigma(\vec{\ell}) \langle e, \vec{g} \mid hk^{-1}, h \cdot \vec{\ell} \cdot k^{-1} \rangle$$
$$= \int d[\vec{g}, h] \psi^*(\vec{g}) \sigma(h^{-1} \cdot \vec{g} \cdot h)$$
$$= \langle \langle \psi \mid \sigma \rangle \rangle$$

$$\Delta_1^\dagger : \Pi_A \Pi_B \mathcal{H}(\Lambda') \to \Pi_A \Pi_B \mathcal{H}(\Lambda)$$

$$\Lambda' \to \Lambda, \text{ then } \Delta_1^\dagger |\Psi\rangle\rangle$$

$$\Delta_1^\dagger |\Psi\rangle\rangle = [(\langle e|_\ell \otimes \mathrm{Id}_\Lambda) |\Psi\rangle]$$

$$\Delta_2|\Psi\rangle = [\Pi_B]_p[\Pi_B]_{p'}\left(\int\ dg|g\rangle\right)|\Psi\rangle.$$

$$\Delta_2|\Psi\rangle = \int_G d[h,\vec{g}]\delta\left(hg_{\partial p}\right)\delta\left(h^{-1}g_{\partial p'}\right)\psi(\vec{g})|h\rangle|\vec{g}\rangle.$$

$$\Delta_2|\Psi\rangle\rangle = \left[\left(\int\ dg|g\rangle\right)|\Psi\rangle \sim \left(\int\ dg|g\rangle\right)|\Psi\rangle + |\chi\rangle\right] \equiv |1,\Psi\rangle\Big\rangle$$

$$\left\langle\langle\psi|\Delta_2^\dagger\Delta_2 \mid \sigma\rangle\right\rangle = \int\ d[h,k]\langle h \mid \langle\langle\psi|[\Pi_B]_p[\Pi_B]_{p'} \mid k\rangle \mid \sigma\rangle\Big\rangle$$

$$= \int\ d[h,k,\vec{g},\vec{m}]\psi^*(\vec{g})\sigma(\vec{m})\delta\left(hg_{\partial p}\right)\delta\left(h^{-1}g_{\partial p'}\right)\langle h,\vec{g} \mid k,\vec{m}\rangle$$

$$= \int\ d[\vec{g}]\psi^*(\vec{g})\sigma(\vec{g})\delta\left(g_{\partial p}g_{\partial p'}\right)$$

$$= \langle\langle\psi \mid \sigma\rangle\rangle$$

$$\Delta_2^\dagger\colon \Pi_A\Pi_B\mathcal{H}(\Lambda') \to \Pi_A\Pi_B\mathcal{H}(\Lambda)$$

$$\Delta_2^\dagger|\Psi\rangle\rangle = \left[\left(\int\ dg\langle g|_\ell \otimes \mathrm{Id}_\Lambda\right)|\Psi\rangle\right]$$

$$H = -\sum_v A_v[1] - \sum_p B_{(v,p)}(e).$$

$$\begin{gathered}
A_v(g)^\dagger\ = A_v(g^{-1}),\\
A_v(g)A_v(h)\ = A_v(gh),\\
B_{(v,p)}(g)^\dagger\ = B_{(v,p)}(g),\\
B_{(v,p)}(g)B_{(v,p)}(h)\ = \delta(g^{-1}h)B_{(v,p)}(h),\\
A_v(g)B_{(v,p)}(h)\ = B_{(v,p)}(ghg^{-1})A_v(g).
\end{gathered}$$

$$A_v[f_1] = \int\ dgf_1(g)A_v(g)\ B_{(v,p)}[f_\infty] = \int\ dgf_\infty(g)B_{(v,p)}(g)$$

$$\begin{gathered}
A_v[f]^\dagger\ = A_v(\bar{f})\\
A_v[f]A_v[f']\ = A_v[f * f']\\
B_{(v,p)}[f]^\dagger\ = B_{(v,p)}[f^*]\\
B_{(v,p)}[f]B_{(v,p)}[f']\ = B_{(v,p)}[f \cdot f']\\
A_v(g)B_{(v,p)}[f']\ = B_{(v,p)}\left[\mathrm{Ad}_g(f')\right]A_v(g).
\end{gathered}$$

$$\begin{gathered}
\bar{f}(g)\ = f^*(g^{-1})\\
f^*(g) \text{ is the complex conjugate of } f(g)\\
(f * f')(g)\ = \int\ dhf(h)f'(h^{-1}g)\\
(f \cdot f')(g)\ = f(g)f'(g)\\
\mathrm{Ad}_g(f)(h)\ = f(g^{-1}hg)
\end{gathered}$$

$$\|f\|_* = \sup_{\pi\in\hat{G}}\|\pi(f)\| = \|A_v[f]\|_\infty$$

$$(W^1(G) \vee W^\infty(G))''$$

$$L^\infty(G) \rtimes_\alpha G$$

$$D(G_k) \cong G_k \times \overline{G_k}$$

$$k = i\sigma, \overline{\mathrm{SL}(4,\mathbb{R})_k} = \mathrm{SL}(4,\mathbb{R})_k$$

$$\langle A \mid B \rangle_{V_\pi \otimes V_\pi^*} = \mathrm{tr}\left[A^\dagger B\right]$$

$$L^{2+\epsilon}(G) \otimes \int_{\hat{G}} d\mu(\pi) f(\pi) = \sum_{\pi \in \hat{G}} d_\pi f(\pi)$$

$U_{n+1}(e) \subset U_n(e)$ and that $g \in V_n(e) \Longrightarrow g^{-1} \in V_n(e)$ for all n.

$$g \in G, (A_v(g) - 1)|\psi\rangle$$

$$L^1(G) \circledast \int_G dg|f(g)|$$

$$\frac{1}{|C|} \sum_{[h] \in C} \rightarrow \sum_T \int_T dt \Delta(t)$$

$$X^{-1} = \int_{1/\Lambda^2}^{1/k^2} ds e^{-sX}$$

$$\log[X] = -\int_{1/\Lambda^2}^{1/k^2} \frac{ds}{s} e^{-sX}$$

$$k\partial_k S_k = -\frac{k\partial_k}{2} \int_{1/\Lambda^2}^{1/k^2} \frac{ds}{s} \mathrm{TR}\left[e^{-sS_k''}\right],$$

$$S_1 = -\frac{1}{2} \int_0^\infty \frac{ds}{s} \mathrm{TR}\left[e^{-sS_0''}\right] = \frac{1}{2} \mathrm{TR} \log\left[S_0''\right]$$

$$S_1 + S_2 = -\frac{1}{2} \int_{1/\Lambda^2}^{1/k^2} \frac{ds}{s} \mathrm{TR}\left[e^{-s(S_0'' + S_1'')}\right],$$

$$S_1 + S_2 = -\frac{1}{2} \int_{1/\Lambda^2}^{1/k^2} \frac{ds}{s} \mathrm{TR}\left[e^{-sS_0''}(1 - sS_1'')\right]$$

$$(S_1)'' = \left(-\frac{1}{2} \int_{1/\Lambda^2}^{s} \frac{dt}{t} \mathrm{TR}\left[e^{-tS_0''}\right]\right)'' = -\frac{1}{2} \int_{1/\Lambda^2}^{s} \frac{dt}{t} \mathrm{TR}\left[e^{-tS_0''}\left(-tS_0^{(4)} + t^2 S_0^{(3)} S_0^{(3)}\right)\right],$$

$$\mathrm{TR}\left[e^{-(A+B)}\right]=\mathrm{TR}\left[e^{-A}-Be^{-A}+\frac{(-1)^2}{2}\int_0^1 du_1\left(Be^{-(1-u_1)A}Be^{-u_1A}\right)\right.$$
$$+\cdots+\frac{(-1)^{n+1}}{n+1}\int_0^1 du_1\ldots\int_0^1 du_n(u_1^{n-1}u_2^{n-2}\ldots u_{n-1}^1u_n^0)$$
$$\left.\times Be^{-(1-u_1)A}Be^{-u_1(1-u_2)A}Be^{-u_1u_2(1-u_3)A}\ldots Be^{-u_1u_2\ldots u_{n-1}u_nA}+\cdots\right]$$

$$S_0=\int d^4x\left[\frac{1}{2}\partial\phi_j\partial\phi_j+\frac{\lambda}{4!}\left(\phi_j\phi_j\right)^2\right]$$

$$\beta_{\lambda_R}=k\partial_k(Z^{-2}\lambda)=-2Z^{-3}\lambda k\partial_k(Z)+Z^{-2}k\partial_k\lambda,$$

$$[k\partial_k\lambda]_{1L}=-\frac{k\partial_k}{2}\mathrm{P}_{\phi^4}\left\{\int_{1/\Lambda^2}^{1/k^2}\frac{ds}{s}\int\frac{d^4p}{(2\pi)^4}\mathrm{Tr}\left[e^{-s\left[\left(p^2+\frac{\lambda}{6}\phi_j\phi_j\right)\delta_{ab}+\frac{\lambda}{3}\phi_a\phi_b\right]}\right]\right\}_{\phi=0}$$

$$\mathrm{P}_{\phi^4}=(d^4/(d\phi_i)^4)$$

$$\int[d^4p/(2\pi)^4]e^{-sp^2}=1/(16\pi^2s^2)$$

$$[k\partial_k\lambda]_{1L}=-\frac{\lambda^2}{(16\pi^2)}\frac{(N+8)}{6}k\partial_k\log\frac{\Lambda^2}{k^2},$$

$$\beta_{\lambda_R}^{1L}=\frac{\lambda^2}{(16\pi^2)}\frac{(N+8)}{3}.$$

$$[k\partial_k\lambda]_{2L}=-\frac{k\partial_k}{2}\mathrm{P}_{\phi^4}\left\{\int_{1/\Lambda^2}^{1/k^2}\frac{ds}{s}\int\frac{d^4p_e}{(2\pi)^4}\mathrm{Tr}\left[e^{-sS_0''}(-sS_1'')\right]\right\}\Bigg|_{\phi=0}.$$

$$(S_1)''=-\frac{1}{2}\int_{\Lambda^{-2}}^{s}\frac{dt}{t}\int\frac{d^4p_i}{(2\pi)^4}\mathrm{Tr}\left[e^{-tS_0''}(-tS_0''''+t^2S_0'''S_0''')\right],$$

$$[k\partial_k\lambda]_{2L}=\frac{\lambda^3}{(16\pi^2)^2}k\partial_k\left\{\int_{1/\Lambda^2}^{1/k^2}\frac{ds}{4s^2}\int_{1/\Lambda^2}^{s}\frac{dt}{t^3}\left[\frac{t^3(N^2+30N+104)}{9}\right.\right.$$
$$\left.\left.+\frac{st^2(2N^2+28N+128)}{9}+\frac{s^2t(N^2+10N+16)}{9}\right]\right\}$$

$$[k\partial_k\lambda]_{2L}=-\frac{\lambda^3}{(16\pi^2)^2}\frac{(10N+44)}{9}.$$

$$Z=\left(\frac{\partial}{\partial q^2}\left[\frac{\delta^2S}{\delta\tilde{\phi}_i(q)\delta\tilde{\phi}_i(-q)}\right]_{\phi=0}\right)_{q^2=0}.$$

$$Z_2=\left[\frac{\partial}{\partial q^2}\left(-\int_{1/\Lambda^2}^{1/k^2}\frac{ds}{4}\int\frac{d^4p_e}{(2\pi)^4}\int_{1/\Lambda^2}^{s}\frac{dt}{t}\int\frac{d^4p_i}{(2\pi)^4}\int_0^1 du\frac{\lambda^2(2N+4)}{3}\right.\right.$$
$$\left.\left.\times e^{-sp_e^2}e^{-tp_i^2(1-u)}e^{-su(p_i+q)^2}\right)\right]_{q^2=0}=\frac{\lambda^2}{(16\pi^2)^2}\frac{(N+2)}{36}\log\frac{\Lambda^2}{k^2}$$

$$\beta_{\lambda_R}^{2L} = \frac{\lambda^3}{(16\pi^2)^2}\left(\frac{(N+2)}{9} - \frac{(10N+44)}{9}\right) = -\frac{\lambda^3}{(16\pi^2)^2}\frac{3N+14}{3}.$$

$$\eta = \frac{\lambda^2}{(16\pi^2)^2}\frac{N+2}{18}.$$

$$k\partial_k S_k = \mathrm{TR}\left(1 + \frac{S_k''}{mk^2}\right)^{-(m+1)},$$

$$\hat{A}_\mu^a(x)\big(a = 1,2,\dots,(N^2-1)\big)$$

$$F_{\mu\nu}^a(\hat{A}) = \partial_\mu \hat{A}_\nu^a(x) - \partial_\nu \hat{A}_\mu^a(x) + g f^{abc}\hat{A}_\mu^b(x)\hat{A}_\nu^c(x)$$

$$D_\mu^{ac}(A)Q_\mu^c = \left[\partial_\mu \delta^{ac} + g f^{abc}A_\mu^b(x)\right]Q_\mu^c(x)$$

$$S_A = S_{\mathrm{YM}} + S_{gf} = \int d^4x' \left\{\frac{1}{4}F_{\mu\nu}^a F_{\mu\nu}^a + \frac{1}{2}\big(D_\mu^{ab}(A)Q_\mu^b\big)\big(D_\nu^{ac}(A)Q_\nu^c\big)\right\}$$

$$S_G = \int d^4x' \left\{-\bar{\theta}_a\big(D_\mu^{ab}(A)D_\mu^{bc}(\hat{A})\big)\theta_c\right\}$$

$$\frac{1}{4}F_{\mu\nu}^a F_{\mu\nu}^a \longrightarrow \frac{Z_A}{4}F_{\mu\nu}^a F_{\mu\nu}^a$$

$$g_R = \frac{g}{\sqrt{Z_A}}$$

$$Z_A = 1 + \left[\frac{g^2N}{16\pi^2}\beta_1 + \left(\frac{g^2N}{16\pi^2}\right)^2\beta_2 + O(g^6)\right]\log\frac{k^2}{\Lambda^2} + O\left(\log^2\frac{k^2}{\Lambda^2}\right)$$

$$\beta_{g_R} = k\partial_k g_R = -\frac{g}{2Z_A^{\frac{3}{2}}}k\partial_k(Z_A) = -g\left[\frac{g^2N}{16\pi^2}\beta_1 + \left(\frac{g^2N}{16\pi^2}\right)^2\beta_2 + O(g^6)\right].$$

$$K_{\mu\nu}^{ab}(A) = \left.\frac{\delta^2 S_A(A,Q)}{\delta Q_\mu^a(x)\delta Q_\nu^b(y)}\right|_{Q=c=0};\ \big(\mathcal{D}_{\bar{c},Q}\big)_\nu^{ab}(A,\chi) = \left.\frac{\delta^2 S_G(A,Q,\chi,c)}{\delta \bar{c}^a(x)\delta Q_\nu^b(y)}\right|_{Q=c=0}$$

$$\big(\mathcal{D}_{Q,c}\big)_\mu^{ab}(A,\chi) = \left.\frac{\delta^2 S_G(A,Q,\chi,c)}{\delta Q_\mu^a(x)\delta c^b(y)}\right|_{Q=c=0};\ H^{ab}(A) = \left.\frac{\delta^2 S_G(A,Q,\chi,c)}{\delta \bar{c}^a(x)c^b(y)}\right|_{Q=c=0}$$

$$\begin{aligned}K_{\mu\nu}^{ab}(A) = &\big\{[-\delta^{ab}\partial_x^2 + 2g f^{abc}A^c\cdot\partial_x + g f^{abc}(\partial_x\cdot A^c) - g^2 f^{amc}f^{clb}A^m\cdot A^l]\delta_{\mu\nu}\\ &+2g f^{abc}F_{\mu\nu}^c\big\}\delta^4(x-y)\end{aligned}$$

$$H^{ab}(A) = [-\delta^{ab}\partial_x^2 + 2g f^{abc}A^c\cdot\partial_x + g f^{abc}(\partial_x\cdot A^c) - g^2 f^{amc}f^{clb}A^m\cdot A^l]\delta^4(x-y)$$

$$S_{\mathrm{YM1}} = -\frac{1}{2}\int_{1/\Lambda^2}^{1/k^2}\frac{ds}{s}\mathrm{TR}\left[e^{-sK_{\mu\nu}^{ab}}\right] + \int_{1/\Lambda^2}^{1/k^2}\frac{ds}{s}\mathrm{TR}\left[e^{-s[H^{ab}+M^{ab}]}\right]$$

$$M^{ab} = -\big(\mathcal{D}_{\bar{c},Q}\big)_{\mu_1}^{ac}(A,\chi)\cdot[K^{-1}]_{\mu_1\mu_2}^{cd}(A)\cdot\big(\mathcal{D}_{Q,c}\big)_{\mu_2}^{db}(A,\chi)$$

$$S_{\mathrm{YM2}}=\int_{1/\Lambda^2}^{1/k^2}\frac{ds}{s}\left\{-\frac{1}{2}\mathrm{TR}[-sS''_{\mathrm{YM1}}e^{-sK}]+\mathrm{TR}[-s\ddot{S}_{\mathrm{YM1}}e^{-sH}]\right\}$$

$$S''_{\mathrm{YM1}}=-\frac{1}{2}\int_{\frac{1}{\Lambda^2}}^{s}\frac{dt}{t}\left\{\mathrm{TR}\left[\int_0^1 dut^2K'e^{-t(1-u)K}K'e^{-tuK}\right]+\mathrm{TR}[-tK''e^{-tK}]\right\}+$$
$$\int_{1/\Lambda^2}^{s}\frac{dt}{t}\left\{\mathrm{TR}\left[\int_0^1 dut^2R'e^{-t(1-u)R}R'e^{-tuR}\right]+\mathrm{TR}[-tR''e^{-tR}]\right\}$$
$$\ddot{S}_{\mathrm{YM1}}=\int_{1/\Lambda^2}^{s}\frac{dt}{t}\left\{\mathrm{TR}\left[\int_0^1 dut^2\dot{R}e^{-t(1-u)R}\dot{R}e^{-tuR}\right]+\mathrm{TR}[-t\ddot{R}e^{-tR}]\right\}$$

$$\Sigma_1=-\int_{1/\Lambda^2}^{s}dtt\int\frac{d^4p_i}{(2\pi)^4}\left\{\mathrm{Tr}\left[\int_0^1 du\delta[K']e^{-t(1-u)(p_i+q)^2}\delta[K']e^{-tup_i^2}\right]-\right.$$
$$\left.2\mathrm{Tr}\left[\int_0^1 du\delta[H']e^{-t(1-u)(p_i+q)^2}\delta[H']e^{-tup_i^2}\right]\right\}.$$

$$\Sigma_2=-\int_{1/\Lambda^2}^{s}\frac{dt}{2t}\int\frac{d^4p_i}{(2\pi)^4}\left\{\mathrm{Tr}\left[(t^4K'K'-t^3K'')\int_0^1 du\delta[K]e^{-t(1-u)(p_i+q)^2}\delta[K]e^{-tup_i^2}\right]-\right.$$
$$\left.2\mathrm{Tr}\left[(t^4H'H'-t^3H'')\int_0^1 du\delta[H]e^{-t(1-u)(p_i+q)^2}\delta[H]e^{-tup_i^2}\right]\right\}$$

$$\Sigma_3=2\int_{1/\Lambda^2}^{s}\frac{dt}{t}\int\frac{d^4p_i}{(2\pi)^4}\left\{\mathrm{Tr}\left[t^3K'\int_0^1 du\delta[K]e^{-t(1-u)(p_i+q)^2}\delta[K']e^{-tup_i^2}\right]-\right.$$
$$\left.2\mathrm{Tr}\left[t^3H'\int_0^1 du\delta[H]e^{-t(1-u)(p_i+q)^2}\delta[H']e^{-tup_i^2}\right]\right\}$$

$$Z^{\mathcal{T}}(N;\boldsymbol{\xi})=e^{A^{\mathcal{T}}(\xi)}C^{\mathcal{T}}(\boldsymbol{\xi})^{-\frac{1}{3}}\mathrm{Ai}\left[C^{\mathcal{T}}(\boldsymbol{\xi})^{-\frac{1}{3}}\left(N-B^{\mathcal{T}}(\boldsymbol{\xi})\right)\right]\left(1+\mathcal{O}\left(e^{-\#\sqrt{N}}\right)\right).$$

$$\mathrm{U}(N)_k\times\underbrace{\mathrm{U}(N)_0\times\cdots\times\mathrm{U}(N)_0}_{q-1}\times\mathrm{U}(N)_{-k}\times\underbrace{\mathrm{U}(N)_0\times\cdots\times\mathrm{U}(N)_0}_{\tilde{q}-1}.$$

$$Z_k^{(q,\tilde{q})}\left(N;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}\right)$$
$$=\frac{1}{(N!)^{q+\tilde{q}}}\int_{-\infty}^{\infty}\left(\prod_{\alpha=1}^{q}\prod_{i=1}^{N}\frac{d\lambda_i^{(\alpha)}}{2\pi}\right)\left(\prod_{\alpha=1}^{\tilde{q}}\prod_{i=1}^{N}\frac{d\tilde{\lambda}_i^{(\alpha)}}{2\pi}\right)$$
$$\times e^{-\frac{ik}{4\pi}\sum_i^N\left(\left(\tilde{\lambda}_i^{(\tilde{q})}\right)^2-\left(\lambda_i^{(q)}\right)^2\right)}e^{-\frac{i}{2}\sum_{\alpha=1}^{q}\eta_\alpha\sum_i^N\left(\lambda_i^{(\alpha-1)}-\lambda_i^{(\alpha)}\right)}e^{-\frac{i}{2}\sum_{\alpha=1}^{\tilde{q}}\tilde{\eta}_\alpha\sum_i^N\left(\tilde{\lambda}_i^{(\alpha-1)}-\tilde{\lambda}_i^{(\alpha)}\right)}$$
$$\times\prod_{\alpha=1}^{q}\frac{\prod_{i<j}^N 2\sinh\frac{\lambda_{ij}^{(\alpha-1)}}{2}\prod_{i<j}^N 2\sinh\frac{\lambda_{ij}^{(\alpha)}}{2}}{\prod_{i,j}^N 2\cosh\frac{\lambda_i^{(\alpha-1)}-\lambda_j^{(\alpha)}+\pi M}{2}}\prod_{\alpha=1}^{\tilde{q}}\frac{\prod_{i<j}^N 2\sinh\frac{\tilde{\lambda}_{ij}^{(\alpha-1)}}{2}\prod_{i<j}^N 2\sinh\frac{\tilde{\lambda}_{ij}^{(\alpha)}}{2}}{\prod_{i,j}^N 2\cosh\frac{\tilde{\lambda}_i^{(\alpha-1)}-\tilde{\lambda}_j^{(\alpha)}+\pi\tilde{M}}{2}}$$

$$\lambda_i^{(0)}=\tilde{\lambda}_i^{(\tilde{q})},\tilde{\lambda}_i^{(0)}=\lambda_i^{(q)}\text{ and }\lambda_{ij}=\lambda_i-\lambda_j$$

$$\sum_{\alpha=1}^{q}\eta_\alpha=\sum_{\alpha=1}^{\tilde{q}}\tilde{\eta}_\alpha=0.$$

$$|\mathrm{Im}(\eta_\alpha)|<1,|\mathrm{Im}(\tilde{\eta}_\alpha)|<1,$$

$$|\mathrm{Im}(M)|<1,|\mathrm{Im}(\tilde{M})|<1$$

$$\frac{\prod_{i<j}^{N}\ 2\sinh\frac{\mu_i-\mu_j}{2}\prod_{i<j}^{N}\ 2\sinh\frac{\nu_i-\nu_j}{2}}{\prod_{i,j}^{N}\ 2\cosh\frac{\mu_i-\nu_j+c}{2}}=\det\left(\left[\frac{1}{2\cosh\frac{\mu_i-\nu_j+c}{2}}\right]_{i,j}^{N\times N}\right)$$

$$\lambda_i^{(\alpha)}\to\lambda_i^{(\alpha)}/k\text{ and }\tilde{\lambda}_i^{(\alpha)}\to\tilde{\lambda}_i^{(\alpha)}/k$$

$$\frac{1}{2\cosh\left(\frac{\mu-\nu}{2k}+c\right)}=\frac{1}{2\pi}\int_{-\infty}^{\infty}dp\frac{e^{\frac{i}{2\pi k}p(\mu-\nu+2kc)}}{2\cosh\frac{p}{2}}=k\langle\mu|\frac{e^{\frac{ic}{\pi}\hat{p}}}{2\cosh\left(\frac{\pi k}{\hbar}\hat{p}\right)}|\nu\rangle$$

$$\langle x\mid y\rangle=2\pi\delta(x-y),\langle\langle p\mid p'\rangle\rangle=2\pi\delta(p-p'),\langle x\mid p\rangle\rangle=\frac{1}{\sqrt{k}}e^{\frac{ixp}{\hbar}}$$

$$\begin{aligned}&Z_k^{(q,\tilde{q})}(N;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})\\&=\frac{1}{(N!)^{q+\tilde{q}}}\int_{-\infty}^{\infty}\left(\prod_{\alpha=1}^{q}\prod_{i=1}^{N}\frac{d\lambda_i^{(\alpha)}}{2\pi}\right)\left(\prod_{\alpha=1}^{\tilde{q}}\prod_{i=1}^{N}\frac{d\tilde{\lambda}_i^{(\alpha)}}{2\pi}\right)\\&\times\prod_{\alpha=1}^{q}\det\left(\left[\left\langle\lambda_i^{(\alpha-1)}\right|e^{-\frac{i}{4\pi k}\delta_{\alpha,1}\hat{x}^2}e^{-\frac{i}{2k}\eta_\alpha\hat{x}}\frac{e^{\frac{i}{2}M\hat{p}}}{2\cosh\frac{\hat{p}}{2}}e^{\frac{i}{2k}\eta_\alpha\hat{x}}e^{\frac{i}{4\pi k}\delta_{\alpha,q}\hat{x}^2}\left|\lambda_j^{(\alpha)}\right\rangle\right]_{i,j}^{N\times N}\right)\\&\times\prod_{\alpha=1}^{\tilde{q}}\det\left(\left[\left\langle\tilde{\lambda}_i^{(\alpha-1)}\right|e^{-\frac{i}{2k}\tilde{\eta}_\alpha\hat{x}}\frac{e^{\frac{i}{2}\tilde{M}\hat{p}}}{2\cosh\frac{\hat{p}}{2}}e^{\frac{i}{2k}\tilde{\eta}_\alpha\hat{x}}\left|\tilde{\lambda}_j^{(\alpha)}\right\rangle\right]_{i,j}^{N\times N}\right)\end{aligned}$$

$$\begin{aligned}&\frac{1}{N!}\int_{-\infty}^{\infty}\left(\prod_{i=1}^{N}\frac{d\nu_i}{2\pi}\right)\det\left(\left[\langle\mu_i|\hat{A}|\nu_j\rangle\right]_{i,j}^{N\times N}\right)\det\left(\left[\langle\nu_i|\hat{B}|\sigma_j\rangle\right]_{i,j}^{N\times N}\right)\\&=\det\left(\left[\langle\mu_i|\hat{A}\hat{B}|\sigma_j\rangle\right]_{i,j}^{N\times N}\right)\end{aligned}$$

$$\begin{aligned}&Z_k^{(q,\tilde{q})}(N;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})\\&=\frac{1}{N!}\int_{-\infty}^{\infty}\left(\prod_{i=1}^{N}\frac{d\lambda_i}{2\pi}\right)\det\left(\left[\langle\lambda_i|\prod_{\alpha=1}^{q}\frac{e^{\frac{i}{2}M(\hat{x}+\hat{p})}}{2\cosh\frac{\hat{x}+\hat{p}+\pi\eta_\alpha}{2}}\prod_{\alpha=1}^{\tilde{q}}\frac{e^{\frac{i}{2}\tilde{M}\hat{p}}}{2\cosh\frac{\hat{p}+\pi\tilde{\eta}_\alpha}{2}}|\lambda_j\rangle\right]_{i,j}^{N\times N}\right)\end{aligned}$$

$$e^{-\frac{ic}{\hbar}\hat{x}}f(\hat{p})e^{\frac{ic}{\hbar}\hat{x}}=f(\hat{p}+c),e^{-\frac{ic}{2\hbar}\hat{x}^2}f(\hat{p})e^{\frac{ic}{2\hbar}\hat{x}^2}=f(\hat{p}+c\hat{x})$$

$$e^{-\frac{ic}{2\hbar}\hat{p}^2}f(\hat{x})e^{\frac{ic}{2\hbar}\hat{p}^2}=f(\hat{x}-c\hat{p})$$

$$\begin{aligned}&Z_k^{(q,\tilde{q})}(N;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})\\&=\frac{1}{N!}\int_{-\infty}^{\infty}\left(\prod_{i=1}^{N}\frac{d\lambda_i}{2\pi}\right)\det\left(\left[\langle\lambda_i|\hat{\rho}_k^{(q,\tilde{q})}(\hat{x},\hat{p};\eta_\alpha,M,\tilde{\eta}_\alpha,\tilde{M})|\lambda_j\rangle\right]_{i,j}^{N\times N}\right)\end{aligned}$$

$$\hat{\rho}_k^{(q,\tilde{q})}(\hat{x},\hat{p};\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) = \prod_{\alpha=1}^{q} \frac{e^{\frac{i}{2}M\hat{x}}}{2\cosh\frac{\hat{x}+\pi\eta_\alpha}{2}} \prod_{\alpha=1}^{\tilde{q}} \frac{e^{\frac{i}{2}\tilde{M}\hat{p}}}{2\cosh\frac{\hat{p}+\pi\tilde{\eta}_\alpha}{2}}$$

$$\Xi_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})$$

$$\Xi_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) = 1 + \sum_{N=1}^{\infty} e^{\mu N} Z_k^{(q,\tilde{q})}(N;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})$$

$$\Xi_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) = \mathrm{Det}\left(1 + e^{\mu}\hat{\rho}_k^{(q,\tilde{q})}(\hat{x},\hat{p};\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})\right)$$

$$J_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})$$

$$\Xi_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) = \sum_{n=-\infty}^{\infty} e^{J_k^{(q,\tilde{q})}(\mu+2\pi i n;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})}$$

$$Z_k^{(q,\tilde{q})}(N;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) = \int_{-i\infty}^{i\infty} \frac{d\mu}{2\pi i} e^{J_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})-\mu N}$$

$$J_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M,\tilde{\eta}_\alpha;\tilde{M})$$
$$= \frac{C_k^{(q,\tilde{q})}(M;\tilde{M})}{3}\mu^3 + B_k^{(q,\tilde{q})}(\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})\mu + A_k^{(q,\tilde{q})}(\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) + \mathcal{O}(e^{-\#\mu}),$$

$$C_k^{(q,\tilde{q})}(M;\tilde{M}) = \frac{2}{\pi^2 kq\tilde{q}(1+M^2)(1+\tilde{M}^2)},$$

$$B_k^{(q,\tilde{q})}(\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) = -\frac{1}{2kq(1+M^2)}\left(\sum_{\alpha=1}^{\tilde{q}}\tilde{\eta}_\alpha^2 + \frac{\tilde{q}}{3}\right) - \frac{1}{2k\tilde{q}(1+\tilde{M}^2)}\left(\sum_{\alpha=1}^{q}\eta_\alpha^2 + \frac{q}{3}\right) + \frac{2}{3kq\tilde{q}(1+M^2)(1+\tilde{M}^2)} + \frac{kq\tilde{q}}{24},$$

$$A_k^{(q,\tilde{q})}(\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) = \frac{1}{4}\sum_{\pm}\left[\sum_{\alpha,\beta=1}^{\tilde{q}}\mathcal{A}\big((1\pm iM)qk,\tilde{\eta}_{\alpha\beta}\big) + \sum_{\alpha,\beta=1}^{q}\mathcal{A}\big((1\pm i\tilde{M})\tilde{q}k,\eta_{\alpha\beta}\big)\right],$$

where $\eta_{\alpha\beta} = \eta_\alpha - \eta_\beta, \tilde{\eta}_{\alpha\beta} = \tilde{\eta}_\alpha - \tilde{\eta}_\beta$, and $\mathcal{A}(\kappa,\chi)$

$$\mathcal{A}(\kappa,\chi) = \frac{2\zeta(3)}{\pi^2\kappa} + \frac{\chi^2}{2\kappa} - \frac{\kappa}{12} + \frac{1}{\pi}\int_0^{\infty} dy \frac{1}{e^{2\pi y}-1}\frac{d}{dy}\left[\frac{\cos(\pi\chi y)}{y\tanh\frac{\pi\kappa y}{2}} - \frac{2}{\pi\kappa y^2}\right]$$

$C^{\mathcal{T}}(\boldsymbol{\xi}) = C_k^{(q,\tilde{q})}(M;\tilde{M}), B^{\mathcal{T}}(\boldsymbol{\xi}) = B_k^{(q,\tilde{q})}(\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})$ and $A^{\mathcal{T}}(\boldsymbol{\xi}) = A_k^{(q,\tilde{q})}(\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})$

$$J_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})$$
$$= \mathrm{Trlog}\left(1+e^{\mu}\hat{\rho}_k^{(q,\tilde{q})}(\hat{x},\hat{p};\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})\right)$$
$$-\log\left[1+\sum_{n\neq 0} e^{J_k^{(q,\tilde{q})}(\mu+2\pi i n;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})-J_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})}\right]$$

$$J_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) = \frac{C}{3}\mu^3+\cdots$$

$$J_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) \approx \mathrm{Trlog}\left(1+e^{\mu}\hat{\rho}_k^{(q,\tilde{q})}(\hat{x},\hat{p};\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})\right)$$

$$\hat{H}_k^{(q,\tilde{q})}(\hat{x},\hat{p};\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})$$

$$\hat{\rho}_k^{(q,\tilde{q})}(\hat{x},\hat{p};\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) = e^{-\frac{1}{2}U(\hat{x})}e^{-T(\hat{p})}e^{-\frac{1}{2}U(\hat{x})} = e^{-\hat{H}_k^{(q,\tilde{q})}(\hat{x},\hat{p};\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})}$$

$$U(x) = \sum_{\alpha=1}^{q} \log\ 2\cosh\frac{x+\pi\eta_\alpha}{2} - \frac{iqMx}{2}$$
$$T(p) = \sum_{\alpha=1}^{\tilde{q}} \log\ 2\cosh\frac{p+\pi\tilde{\eta}_\alpha}{2} - \frac{i\tilde{q}\tilde{M}p}{2}$$

$$n(E) = \mathrm{Tr}\theta\left(E-\hat{H}_k^{(q,\tilde{q})}(\hat{x},\hat{p};\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})\right)$$

$$\theta(z) = \begin{cases} 0 & (z<0) \\ 1 & (z\geq 0) \end{cases}$$

$$J_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) \approx \int_0^\infty dE\frac{dn(E)}{dE}\log\left(1+e^{\mu-E}\right) = \int_0^\infty dE n(E)\frac{e^{\mu-E}}{1+e^{\mu-E}}$$

$$\lim_{E\to\infty} n(E)e^{-E} \to 0$$

$$n(E) = C_k^{(q,\tilde{q})}(M;\tilde{M})E^2 + B_k^{(q,\tilde{q})}(\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) - \frac{\pi^2 C_k^{(q,\tilde{q})}(M;\tilde{M})}{3} + \mathcal{O}\left(e^{-\#E}\right)$$

$$\int_0^\infty dE E^a \frac{e^{\mu-E}}{1+e^{\mu-E}} = -\Gamma(a+1)\mathrm{Li}_{a+1}(-e^{\mu})$$
$$= \frac{(2\pi i)^{a+1}}{a+1}B_{a+1}\left(\frac{\mu}{2\pi i}+\frac{1}{2}\right) - (-1)^a\Gamma(a+1)\mathrm{Li}_{a+1}(-e^{-\mu})$$

$$J_k^{(q,\tilde{q})}(\mu;\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}) = \frac{C_k^{(q,\tilde{q})}(M;\tilde{M})}{3}\mu^3 + B_k^{(q,\tilde{q})}(\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M})\mu$$
$$+ \cdots + \mathcal{O}\left(e^{-\#\mu}\right)$$

$$\mathcal{O}_W = \int_{-\infty}^{\infty}\frac{dy}{2\pi}e^{\frac{ipy}{\hbar}}\left\langle x-\frac{y}{2}\right|\hat{\mathcal{O}}\left|x+\frac{y}{2}\right\rangle$$

$$(f(\hat{x}))_W = f(x), (f(\hat{p}))_W = f(p), (\hat{A}\hat{B})_W = A_W \star B_W, \mathrm{tr}\hat{O} = \int \frac{dxdp}{2\pi\hbar} O_W$$

$$\star = e^{\frac{i\hbar}{2}\left(\overleftarrow{\partial}_x\overrightarrow{\partial}_p - \overleftarrow{\partial}_p\overrightarrow{\partial}_x\right)}$$

$$n(E) = \int_{H_W \le E} \frac{dxdp}{2\pi\hbar}$$

$$\hat{H}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}\right) = U(\hat{x}) + T(\hat{p}) - \frac{1}{24}[U(\hat{p}),[U(\hat{p}),T(\hat{p})]] - \frac{1}{12}[T(\hat{p}),[U(\hat{p}),T(\hat{p})]] + \cdots,$$

$$H_W(x,p) = U(x) + T(p) + \frac{\hbar^2}{24}(\partial_x U(x))^2\partial_p^2 T(p) - \frac{\hbar^2}{12}\partial_x^2 U(x)\left(\partial_p T(p)\right)^2 + \cdots$$

$$U(x) = \frac{q|x|}{2} - \frac{iqMx}{2} + \mathcal{O}\left(e^{-|x|}\right),$$
$$\partial_x U(x) = \frac{q\,\mathrm{sgn}(x)}{2} - \frac{iqM}{2} + \mathcal{O}\left(e^{-|x|}\right),$$
$$\partial_x^2 U(x) = \mathcal{O}\left(e^{-|x|}\right),$$
$$T(p) = \frac{\tilde{q}|p|}{2} - \frac{i\tilde{q}\tilde{M}p}{2} + \mathcal{O}\left(e^{-|p|}\right),$$
$$\partial_p T(p) = \frac{\tilde{q}\,\mathrm{sgn}(p)}{2} - \frac{i\tilde{q}\tilde{M}}{2} + \mathcal{O}\left(e^{-|p|}\right),$$
$$\partial_p^2 T(p) = \mathcal{O}\left(e^{-|p|}\right).$$

$$\frac{q|x|}{2} - \frac{iqMx}{2} + \frac{\tilde{q}|p|}{2} - \frac{i\tilde{q}\tilde{M}p}{2} = E$$

$$\frac{8E^2}{q\tilde{q}(1+M^2)\left(1+\tilde{M}^2\right)}$$

$$n(E) = \frac{1}{2\pi\hbar}\left[\frac{8E^2}{q\tilde{q}(1+M^2)\left(1+\tilde{M}^2\right)} - \mathrm{vol}_{\mathrm{I}} - \mathrm{vol}_{\mathrm{II}} - \mathrm{vol}_{\mathrm{III}} - \mathrm{vol}_{\mathrm{IV}}\right]$$

$$H_W = \frac{q(1-iM)x}{2} + T(p) + \frac{q^2(1-iM)^2\hbar^2}{96}\partial_p^2 T(p) + \frac{q(1-iM)}{2}f\left(\partial_p^3 T(p),\cdots\right)$$

$$x_{\mathrm{in}}(p) = \frac{2}{q(1-iM)}\left[E - T(p) - \frac{q^2(1-iM)^2\hbar^2}{96}\partial_p^2 T(p) - \frac{q(1-iM)}{2}f\left(\partial_p^3 T(p),\cdots\right)\right]$$

$$x_{\mathrm{out}}(p) = \frac{2}{q(1-iM)}\left(E - \frac{\tilde{q}|p|}{2} + \frac{i\tilde{q}\tilde{M}p}{2}\right)$$

$$\mathrm{vol_I} = \int_{-\infty}^{\infty} dp(x_{\mathrm{out}}(p) - x_{\mathrm{in}}(p))$$
$$= \frac{2}{q(1-iM)} \int_{-\infty}^{\infty} dp \left[T(p) - \frac{\tilde{q}|p|}{2} + \frac{i\tilde{q}\tilde{M}p}{2} + \frac{q^2(1-iM)^2\hbar^2}{96} \partial_p^2 T(p) + \frac{q(1-iM)}{2} f\left(\partial_p^3 T(p), \cdots\right) \right]$$

$$\int_{-\infty}^{\infty} dp \left[T(p) - \frac{\tilde{q}|p|}{2} + \frac{i\tilde{q}\tilde{M}p}{2} \right] = \sum_{\alpha=1}^{\tilde{q}} \left(\int_0^{\infty} dp \log\left(1 + e^{-p-\pi\tilde{\eta}_\alpha}\right) + \int_{-\infty}^{0} dp \log\left(1 + e^{p+\pi\tilde{\eta}_\alpha}\right) \right)$$
$$= -\sum_{\alpha=1}^{\tilde{q}} \left(\mathrm{Li}_2\left(-e^{\pi\tilde{\eta}_\alpha}\right) + \mathrm{Li}_2\left(-e^{-\pi\tilde{\eta}_\alpha}\right) \right)$$
$$= \sum_{\alpha=1}^{\tilde{q}} \left(\frac{\pi^2\tilde{\eta}_\alpha^2}{2} + \frac{\pi^2}{6} \right)$$
$$\int_{-\infty}^{\infty} dp \partial_p^2 T(p) = \left[\partial_p T(p)\right]_{p=-\infty}^{\infty} = \tilde{q}$$

$$\mathrm{vol}_I = \frac{\pi^2}{q(1-iM)} \left(\sum_{\alpha=1}^{\tilde{q}} \tilde{\eta}_\alpha^2 + \frac{\tilde{q}}{3} \right) + \frac{\hbar^2 q\tilde{q}(1-iM)}{48}$$

$$\mathrm{vol_{II}} = \frac{\pi^2}{\tilde{q}(1-i\tilde{M})} \left(\sum_{\alpha=1}^{q} \eta_\alpha^2 + \frac{q}{3} \right) - \frac{\hbar^2 q\tilde{q}(1-i\tilde{M})}{24}$$

$$\mathrm{vol_{III}} = \frac{\pi^2}{q(1+iM)} \left(\sum_{\alpha=1}^{\tilde{q}} \tilde{\eta}_\alpha^2 + \frac{\tilde{q}}{3} \right) + \frac{\hbar^2 q\tilde{q}(1+iM)}{48}$$

$$\mathrm{vol_{IV}} = \frac{\pi^2}{\tilde{q}(1+i\tilde{M})} \left(\sum_{\alpha=1}^{q} \eta_\alpha^2 + \frac{q}{3} \right) - \frac{\hbar^2 q\tilde{q}(1+i\tilde{M})}{24}$$

$$J_k^{(q,\tilde{q})}(\mu; \eta_\alpha, M; \tilde{\eta}_\alpha, \tilde{M}) \approx \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \mathrm{Tr} \hat{\rho}_k^{(q,\tilde{q})}(\hat{x}, \hat{p}; \eta_\alpha, M; \tilde{\eta}_\alpha, \tilde{M})^n e^{n\mu}$$
$$= \int_{-i\infty+0^+}^{i\infty+0^+} \frac{ds}{2\pi i} \Gamma(s)\Gamma(-s) \mathcal{Z}^{(q,\tilde{q})}(s; \eta_\alpha, M; \tilde{\eta}_\alpha, \tilde{M}) e^{s\mu},$$

$$\mathcal{Z}^{(q,\tilde{q})}(s; \eta_\alpha, M; \tilde{\eta}_\alpha, \tilde{M}) = \mathrm{Tr} \hat{\rho}_k^{(q,\tilde{q})}(\hat{x}, \hat{p}; \eta_\alpha, M; \tilde{\eta}_\alpha, \tilde{M})^s$$

$$\mathcal{O}\left(e^{-4\pi^2 C_k^{(q,\tilde{q})}(M,\tilde{M})\mu}\right) \text{ corrections to } J_k^{(q,\tilde{q})}(\mu; \eta_\alpha, M; \tilde{\eta}_\alpha, \tilde{M})$$

$$J_k^{(q,\tilde{q})}(\mu; \eta_\alpha, M; \tilde{\eta}_\alpha, \tilde{M}) = -\mathrm{Res}\left[\Gamma(s)\Gamma(-s)\mathcal{Z}^{(q,\tilde{q})}(s; \eta_\alpha, M; \tilde{\eta}_\alpha, \tilde{M}) e^{s\mu}, s \to 0\right]$$
$$+\mathcal{O}\left(e^{-\#\mu}\right)$$

$$\mathcal{Z}^{(q,\tilde{q})}(s; \eta_\alpha, M; \tilde{\eta}_\alpha, \tilde{M}) = \int \frac{dxdp}{2\pi\hbar} \left(\hat{\rho}_k^{(q,\tilde{q})}(\hat{x}, \hat{p}; \eta_\alpha, M; \tilde{\eta}_\alpha, \tilde{M})^s \right)_W$$

$$\mathcal{Z}^{(q,\tilde{q})}\left(s;\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)=\sum_{\ell=0}^{\infty}\hbar^{2\ell-1}\mathcal{Z}_{2\ell}^{(q,\tilde{q})}\left(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M}\right)$$

$$\mathcal{Z}_0^{(q,\tilde{q})}\left(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M}\right)=\frac{\Gamma\left(\frac{sq(1+iM)}{2}\right)\Gamma\left(\frac{sq(1-iM)}{2}\right)\Gamma\left(\frac{s\tilde{q}(1+i\tilde{M})}{2}\right)\Gamma\left(\frac{s\tilde{q}(1-i\tilde{M})}{2}\right)}{2\pi\Gamma(sq)\Gamma(s\tilde{q})}$$

$$\mathcal{Z}_{2\ell}^{(q,\tilde{q})}\left(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M}\right)=D_{2\ell}^{(q,\tilde{q})}\left(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M}\right)\mathcal{Z}_0^{(q,\tilde{q})}\left(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M}\right)$$

$$D_{2\ell}^{(q,\tilde{q})}\left(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M}\right)\odot_\star^\dagger\left(\hat{\rho}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)^{\mathcal{S}}\right)_W$$

$$\left(\hat{\rho}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)^{s+1}\right)_W$$
$$=\left(\hat{\rho}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)^{s}\right)_W\star\rho_{k,W}^{(q,\tilde{q})}\left(x,p;\frac{\hbar v_\alpha}{\pi},M,\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)$$
$$\rho_{k,W}^{(q,\tilde{q})}\left(x,p;\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)$$
$$=\prod_{\alpha=1}^{q}\frac{e^{\frac{iM}{2}x}}{F(x)\cosh\frac{\hbar v_\alpha}{2}+2F'(x)\sinh\frac{\hbar v_\alpha}{2}}\star\prod_{\alpha=1}^{\tilde{q}}\frac{e^{\frac{i\tilde{M}}{2}p}}{F(p)\cosh\frac{\hbar\tilde{v}_\alpha}{2}+2F'(p)\sinh\frac{\hbar\tilde{v}_\alpha}{2}},$$

$$\partial_x^{2n}F(x)=2^{-2n}F(x),\partial_x^{2n-1}F(x)=2^{-2n+2}F'(x),F'(x)^2=\frac{F(x)^2}{4}-1,$$

$$I_1(\alpha,n)=\int_{-\infty}^{\infty}dx\frac{e^{\alpha x}}{F(x)^n}=\frac{\Gamma\left(\frac{n}{2}+\alpha\right)\Gamma\left(\frac{n}{2}-\alpha\right)}{\Gamma(n)}$$
$$I_2(\alpha,n)=\int_{-\infty}^{\infty}dx\frac{e^{\alpha x}F'(x)}{F(x)^n}=\frac{\alpha\Gamma\left(\frac{n-1}{2}+\alpha\right)\Gamma\left(\frac{n-1}{2}-\alpha\right)}{\Gamma(n)}$$

$$I_1(\alpha,n+2)=\frac{n^2-4\alpha^2}{4n(n+1)}I_1(\alpha,n),I_2(\alpha,n)=\frac{\alpha I_1(n-1,\alpha)}{n-1}$$

$$J_k^{(q,\tilde{q})}\left(\mu;\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)=\frac{C_k^{(q,\tilde{q})}(M;\tilde{M})}{3}\mu^3+B_k^{(q,\tilde{q})}\left(\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)\mu$$
$$+\frac{1}{4}\sum_{\pm}\left[\sum_{\alpha=1}^{\tilde{q}}\mathcal{A}\left((1\pm iM)qk,\frac{\hbar\left(\tilde{v}_\alpha-\tilde{v}_\beta\right)}{\pi}\right)+\sum_{\alpha=1}^{q}\mathcal{A}\left((1\pm i\tilde{M})\tilde{q}k,\frac{\hbar\left(v_\alpha-v_\beta\right)}{\pi}\right)\right]$$
$$+\mathcal{O}\left(e^{-\#\mu}\right)$$

$$\mathcal{A}(\kappa,\chi)=\sum_{\ell=0}^{\infty}\sum_{n=0}^{\ell}\frac{(-1)^{\ell+n}}{(2n)!\,(2\ell-2n)!}B_{2\ell-2}B_{2\ell-2n}\pi^{2\ell-2}\kappa^{2\ell-2n-1}\chi^{2n}$$

$$\mathcal{A}(\kappa,\chi)=-\mathcal{A}(-\kappa,\chi),\mathcal{A}(\kappa,\chi)=\mathcal{A}(\kappa,-\chi)$$

$$A_k^{(1,1)}(M=0;\tilde{M}=0)=A^{\mathrm{ABJM}}(k)$$

$$\mathcal{A}(\kappa,0)=A^{\mathrm{ABJM}}(\kappa)$$

$$A^{\mathrm{ABJM}}(\kappa)=\frac{2\zeta(4)}{\pi^2\kappa}+\sum_{\ell=1}^{\infty}\frac{(-1)^{\ell}}{(2\ell)!}B_{2\ell-2}B_{2\ell}\pi^{2\ell-2}\kappa^{2\ell-1}$$

$$\sum_{\ell=0}^{\infty}\sum_{n=0}^{\ell}f(\ell,n)=\sum_{n=0}^{\infty}\sum_{\ell=0}^{\infty}f(\ell+n,n)$$

$$\mathcal{A}(\kappa,\chi)=\sum_{n=0}^{\infty}\sum_{\ell=0}^{\infty}\frac{(-1)^{\ell}}{(2n)!\,(2\ell)!}B_{2\ell+2n-2}B_{2\ell}\pi^{2\ell+2n-2}\kappa^{2\ell-1}\chi^{2n}.$$

$$B_{2n}=\begin{cases}2\zeta(4) & (n=-1)\\ 1 & (n=0)\\ (-1)^{n-1}2\displaystyle\int_0^{\infty}dy\left(\frac{1}{e^{2\pi y}-1}\frac{d}{dy}y^{2n}\right) & (n\geq 1)\end{cases}$$

$$\begin{aligned}\mathcal{A}(\kappa,\chi)=&\frac{2\zeta(4)}{\pi^2\kappa}+\frac{\chi^2}{2\kappa}-\frac{\kappa}{12}\\&+\frac{2}{\pi^2\kappa}\int_0^{\infty}dy\,\frac{1}{e^{2\pi y}-1}\frac{d}{dy}\left[\frac{1}{y^2}\sum_{n=0}^{\infty}\sum_{\ell=0}^{\infty}\frac{(-1)^n(\pi\chi y)^{2n}}{(2n)!}\frac{B_{2\ell}(\pi\kappa y)^{2\ell}}{(2\ell)!}-\frac{1}{y^2}\right]\end{aligned}$$

$$\sum_{n=0}^{\infty}\frac{(-1)^n}{(2n)!}z^{2n}=\cos\,z,\sum_{n=0}^{\infty}\frac{B_{2n}}{(2n)!}z^{2n}=\frac{z}{2\tanh\,\frac{z}{2}}$$

$$\mathcal{A}(\kappa,\chi)=A^{\mathrm{ABJM}}(\kappa)+\sum_{\ell=1}^{\infty}\sum_{n=1}^{\ell}\frac{(-1)^{\ell+n}}{(2n)!\,(2\ell-2n)!}B_{2\ell-2}B_{2\ell-2n}\kappa^{2\ell-2n-1}\chi^{2n}\pi^{2\ell-2}$$

$$\mathcal{A}(\kappa,\chi)=A^{\mathrm{ABJM}}(\kappa)+\frac{\chi^2}{2\kappa}-\frac{2}{\pi}\int_0^{\infty}dy\,\frac{1}{e^{2\pi y}-1}\frac{d}{dy}\left(\frac{\sin^2\,\frac{\pi\chi y}{2}}{y\tanh\,\frac{\pi\kappa y}{2}}\right)$$

$$A^{\mathrm{ABJM}}(\kappa)=\frac{2\zeta(3)}{\pi^2\kappa}\left(1-\frac{\kappa^3}{16}\right)+\frac{\kappa^2}{\pi^2}\int_0^{\infty}dy\,\frac{y}{e^{\kappa y}-1}\log\,(1-e^{-2y})$$

$$\mathcal{A}(\kappa,\chi)=A^{\mathrm{ABJM}}(\kappa)+\frac{\chi^2}{2\kappa}-\int_0^{\infty}dy\,\frac{1}{\sinh^2\,\pi y}\left(\frac{\sin^2\,\frac{\pi\chi y}{2}}{y\tanh\,\frac{\pi\kappa y}{2}}-\frac{\pi\chi^2}{2\kappa}\right)$$

$$\partial_{\chi}\mathcal{A}(\kappa,\chi)=\frac{\chi}{\kappa}-\int_0^{\infty}dy\,\frac{1}{\sinh^2\,\pi y}\left(\frac{\pi\sin\,\pi\chi y}{2\tanh\,\frac{\pi\kappa y}{2}}-\frac{\pi\chi}{\kappa}\right)$$

$$\partial_{\chi}\mathcal{A}(\kappa,\chi)=\frac{\chi}{\kappa}-\frac{1}{2}f_{-\infty}^{\infty}dy\,\frac{1}{\sinh^2\,\pi y}\frac{\pi\sin\,\pi\chi y}{2\tanh\,\frac{\pi\kappa y}{2}}+\frac{1}{2}f_{-\infty}^{\infty}dy\,\frac{1}{2\sinh^2\,\pi y}\frac{\pi\chi}{\kappa}$$

$$f_{-\infty}^{\infty}dy\frac{e^{\alpha y}}{\prod_{a=1}^{n}\sinh\,\pi\nu_a(y-\beta_a)}$$
$$=\frac{1}{1-(-1)^{\sum_a \nu_a}e^{i\alpha}}f_{\gamma}dy\frac{e^{\alpha y}}{\prod_{a=1}^{n}\sinh\,\pi\nu_a(y-\beta_a)}$$
$$=\frac{2\pi i}{1-(-1)^{\sum_a \nu_a}e^{i\alpha}}\sum_{a=1}^{n}\left(\frac{1}{2}\sum_{j=0,\nu_a}\frac{(-1)^j e^{\alpha\left(\beta_a+\frac{ij}{\nu_a}\right)}}{\pi\nu_a\prod_{a'(\neq a)}\sinh\,\pi\nu_{a'}\left(\beta_a+\frac{ij}{\nu_a}-\beta_{a'}\right)}\right.$$
$$\left.+\sum_{j=1}^{\nu_a-1}\frac{(-1)^j e^{\alpha\left(\beta_a+\frac{ij}{\nu_a}\right)}}{\pi\nu_a\prod_{a'(\neq a)}\sinh\,\pi\nu_{a'}\left(\beta_a+\frac{ij}{\nu_a}-\beta_{a'}\right)}\right)$$

$$\frac{1}{2}f_{-\infty}^{\infty}dy\frac{1}{\sinh^2\,\pi y}\frac{\pi\chi}{\kappa}=\frac{\pi\chi}{2\kappa}\lim_{\epsilon_1,\epsilon_2\to 0}fdy\frac{e^{\epsilon_1 y}}{\sinh\,\pi y\sinh\,\pi(y-\epsilon_2)}=-\frac{\chi}{\kappa}$$

$$\partial_\chi\mathcal{A}(\kappa,\chi)$$
$$=-\frac{1}{2}f_{-\infty}^{\infty}dy\frac{1}{\sinh^2\,\pi y}\left(\frac{\pi\sin\,\pi\chi y}{2\tanh\,\frac{\pi\kappa y}{2}}\right)$$
$$=\sum_{\pm}\frac{\pi i}{8}\lim_{\epsilon_1,\epsilon_2\to 0}f_{-\infty}^{\infty}dy\frac{e^{\pi\left(i\chi\pm\frac{\kappa}{2}\right)y}}{\sinh\,\pi y\sinh\,\pi(y-\epsilon_1)\sinh\,\frac{\pi\kappa(y-\epsilon_3)}{2}}$$
$$=\begin{cases}\frac{\pi}{1-e^{-\pi\chi}}\left[\frac{(1+e^{-\pi\chi})(4-\kappa^2+6\chi^2)}{24\kappa}+\sum_{a=1}^{\frac{\kappa}{2}-1}\frac{e^{-\frac{2\pi\chi a}{\kappa}}}{\kappa\sin^2\,\frac{2\pi a}{\kappa}}\right]\\ \frac{\pi}{1-e^{-2\pi\chi}}\left[\frac{(1+e^{-2\pi\chi})(4-\kappa^2+6\chi^2)}{24\kappa}-\frac{\kappa}{4}e^{-\pi\chi}+\sum_{a=1}^{\kappa-1}\frac{e^{-\frac{2\pi\chi a}{\kappa}}}{\kappa\sin^2\,\frac{2\pi a}{\kappa}}\right]\end{cases}$$

$$\mathcal{A}(1,\chi)=A^{\mathrm{ABJM}}(1)+\frac{1}{8}\left(1+\frac{\pi^2}{4}\right)\chi^2+\frac{\pi^2}{48}\left(1-\frac{\pi^2}{16}\right)\chi^4+\cdots$$

$$\mathcal{A}(\kappa,\chi)=\sum_{\ell=0}^{\infty}\frac{(-1)^\ell}{(2\ell)!}B_{2\ell}(\pi\kappa)^{2\ell-1}a_\ell(\chi)$$

$$a_\ell(z)=\frac{1}{\pi}\sum_{n=0}^{\infty}\frac{1}{(2n)!}B_{2\ell+2n-2}(\pi z)^{2n}$$

$$a_\ell(z)=\left(\frac{1}{\pi}\frac{d}{dz}\right)^{2\ell}a_0(z)$$

$$\mathcal{A}(\kappa,\chi)=\frac{i}{2\pi\tanh\,\frac{i\kappa}{2}\frac{d}{d\chi}}\frac{d}{d\chi}a_0(\chi)$$

$$\mathcal{A}(\kappa,\chi)=-\frac{i}{2\pi}\frac{d}{d\chi}a_0(\chi)-\frac{2}{\pi\kappa}\int_{\mathbb{R}+i0^+}\frac{a_0\left(\chi+\frac{\kappa}{\pi}t\right)}{(2\sinh\,t)^2}dt$$

$$\sum_{n=0}^{\infty}\frac{B_{2n-2}}{(2n)!}z^{2n}=-\frac{\pi^2}{6}z-\frac{1}{12}z^3-z\mathrm{Li}_2(e^z)+2\mathrm{Li}_3(e^z)$$

$$a_0(z)=-\frac{\pi^2}{6}z-\frac{\pi^2}{12}z^3-z\mathrm{Li}_2(e^{\pi z})+\frac{2}{\pi}\mathrm{Li}_3(e^{\pi z})$$

$$\frac{d}{dz}a_0(z)=-2\pi i\log\, s_{b=1}\left(\frac{z}{2}\right)$$

$$\mathcal{A}(\kappa,\chi)=-\log\, s_{b=1}\left(\frac{\chi}{2}\right)-\frac{2}{\pi\kappa}\int_{\mathbb{R}+i0^+}\frac{a_0\left(\chi+\frac{\kappa}{\pi}t\right)}{(2\sinh\, t)^2}dt$$

$$C_k^{(q,\tilde{q})}(M;\tilde{M})=\frac{2}{\pi^2kq\tilde{q}(1+M^2)\left(1+\tilde{M}^2\right)}$$

$$B_k^{(q,\tilde{q})}(0,M;0,\tilde{M})=\frac{\pi^2}{3}C_k^{(q,\tilde{q})}(M;\tilde{M})-\frac{q\tilde{q}}{6k}\left(\frac{1}{q^2(1+M^2)}+\frac{1}{\tilde{q}^2\left(1+\tilde{M}^2\right)}\right)+\frac{kq\tilde{q}}{24},$$

$$A_k^{(q,\tilde{q})}(0,M;0,\tilde{M})=\frac{1}{4}\sum_{\pm}\left[\tilde{q}^2A^{\mathrm{ABJM}}((1\pm iM)qk)+q^2A^{\mathrm{ABJM}}((1\pm i\tilde{M})\tilde{q}k)\right]$$

$$\eta_\alpha=\begin{cases}\frac{1}{3}i & (1\le\alpha\le N_{\mathrm{f}})\\ -\frac{1}{3}i & (N_{\mathrm{f}}+1\le\alpha\le 2N_{\mathrm{f}})\end{cases}$$

$$C_k^{(2N_{\mathrm{f}},1)}\left(0;\frac{i}{3}\right)=\frac{9}{8\pi^2kN_{\mathrm{f}}},B_k^{(2N_{\mathrm{f}},1)}\left(\eta_\alpha,0;\frac{i}{3}\right)=\frac{7}{24kN_{\mathrm{f}}}+\frac{kN_{\mathrm{f}}}{12}-\frac{N_{\mathrm{f}}}{4k},$$

$$A_k^{(2N_{\mathrm{f}},1)}\left(\eta_\alpha,0;\frac{i}{3}\right)=\frac{1}{2\pi k}\left[\frac{\zeta(3)}{\pi N_{\mathrm{f}}}+N_{\mathrm{f}}^2\left(\frac{5\zeta(3)}{2\pi}+\frac{\psi^{(1)}\left(\frac{1}{3}\right)-\psi^{(1)}\left(\frac{2}{3}\right)}{4\sqrt{3}}\right)\right]$$
$$+2\pi k\left(-\frac{N_{\mathrm{f}}}{24\pi}-N_{\mathrm{f}}^2\left(\frac{1}{24\pi}+\frac{1}{72\sqrt{3}}\right)\right)+\mathcal{O}(k^3)$$

$$\psi^{(n)}(z)=\partial_z^{n+1}\log\,\Gamma(z)$$

$$A_k^{(2N_{\mathrm{f}},1)}\left(\eta_\alpha,0;\frac{i}{3}\right)$$
$$=\frac{1}{2}A^{\mathrm{ABJM}}(2N_{\mathrm{f}}k)+\frac{N_{\mathrm{f}}^2}{2}\sum_{\pm}\left[A^{\mathrm{ABJM}}\left(\left(1\pm\frac{1}{3}\right)k\right)+\mathcal{A}\left(\left(1\pm\frac{1}{3}\right)k,\frac{2i}{3}\right)\right]$$

$$a_1(z)=\frac{z}{2\tanh\left(\frac{\pi z}{2}\right)}$$

$$a_0\left(\pm\frac{2i}{3}\right)=-\frac{8\zeta(3)}{9\pi}+\frac{\psi^{(1)}\left(\frac{1}{3}\right)-\psi^{(1)}\left(\frac{2}{3}\right)}{9\sqrt{3}},a_1\left(\pm\frac{2i}{3}\right)=\frac{1}{3\sqrt{3}}$$

$$C_{k=1}^{(q,1)}(0;\tilde{M}) = \frac{2}{\pi^2 q\left(1+\tilde{M}^2\right)}$$

$$B_{k=1}^{(q,1)}(\eta_\alpha, 0;\tilde{M}) = \frac{2}{3q\left(1+\tilde{M}^2\right)} + \frac{q}{24} - \frac{1}{6q} - \frac{1}{1+\tilde{M}^2}\left(\frac{q}{6} + \sum_{I=1}^{q-1}\mu_I^2\right)$$

$$A_{k=1}^{(q,1)}(\eta_\alpha, 0;\tilde{M}) = \frac{q^2}{4}\sum_{\pm} A^{\mathrm{ABJM}}(1 \pm i\tilde{M}) + \frac{1}{2}A^{\mathrm{ABJM}}(q) + q\sum_{I=1}^{q-1}\mu_I^2\left[\frac{1}{1+\tilde{M}^2} - \frac{\pi^2}{72} - \frac{\pi^4\left(3\tilde{M}^2-1\right)}{21600} - \frac{\pi^6\left(5\tilde{M}^4 - 10\tilde{M}^2 + 1\right)}{1270080} + \cdots\right] + \mathcal{O}(\mu_I^4)$$

$$T^I = \frac{1}{\sqrt{2I(I+1)}}\mathrm{diag}(\underbrace{1,\ldots,1}_{I}, -I, \underbrace{0,\ldots,0}_{q-I-1}).$$

$$\mathrm{tr}(T^I) = 0, \mathrm{tr}(T^I T^J) = \frac{1}{2}\delta_{IJ}$$

$$\sum_{I=1}^{q-1}\mu_I (T^I)_{\alpha,\alpha} = \frac{1}{2}\eta_\alpha$$

$$\partial_{\mu_1}^4 A\big|_{\mu_I,\tilde{M}=0} = 4(6+q)\mathfrak{b},\ \partial_{\mu_3}^4 A\big|_{\mu_I,\tilde{M}=0} = 4\left(6+\frac{7}{6}q\right)\mathfrak{b}$$

$$\mathfrak{b} = -\frac{1}{2}\pi^2\left(\frac{\pi^2}{32} - \frac{1}{2}\right)$$

$$\sum_{\alpha=1}^{q}\eta_\alpha^2 = 2\sum_{I=1}^{q-1}\mu_I^2,\ \sum_{\alpha,\beta=1}^{q}\left(\eta_\alpha - \eta_\beta\right)^2 = 4q\sum_{I=1}^{q-1}\mu_I^2$$

$$\prod_{n=1}^{L} 2\cosh\left(z + \frac{\pi i}{L}\left(\frac{L+1}{2} - n\right)\right) = 2\cosh(Lz)$$

$(rq, \tilde{r}\tilde{q})_k$ with $\{\eta_\alpha = \xi'_\alpha, M; \tilde{\eta}_\alpha = \tilde{\xi}'_\alpha, \tilde{M}\} \leftrightarrow (q,\tilde{q})_{r\tilde{r}k}$ with $\{\eta_\alpha = \xi_\alpha, M; \tilde{\eta}_\alpha = \tilde{\xi}_\alpha, \tilde{M}\}$

$$\xi'_\alpha = \begin{cases} \frac{\xi_1}{r} + \frac{2i}{r}\left[\frac{1+r}{2} - \alpha\right] & (1 \le \alpha \le r) \\ \frac{\xi_2}{r} + \frac{2i}{r}\left[\frac{1+r}{2} - (\alpha - r)\right] & (r+1 \le \alpha \le 2r) \\ \vdots & \vdots \\ \frac{\xi_q}{r} + \frac{2i}{r}\left[\frac{1+r}{2} - (\alpha - (q-1)r)\right] & ((q-1)r+1 \le \alpha \le qr) \end{cases}$$

$$\tilde{\xi}'_{\alpha}=\begin{cases}\frac{\tilde{\xi}_1}{\tilde{r}}+\frac{2i}{\tilde{r}}\left[\frac{1+\tilde{r}}{2}-\alpha\right] & (1\le\alpha\le\tilde{r})\\ \frac{\tilde{\xi}_2}{\tilde{r}}+\frac{2i}{\tilde{r}}\left[\frac{1+\tilde{r}}{2}-(\alpha-\tilde{r})\right] & (\tilde{r}+1\le\alpha\le 2\tilde{r})\\ \vdots & \vdots\\ \frac{\tilde{\xi}_{\tilde{q}}}{\tilde{r}}+\frac{2i}{\tilde{r}}\left[\frac{1+\tilde{r}}{2}-(\alpha-(\tilde{q}-1)\tilde{r})\right] & ((\tilde{q}-1)\tilde{r}+1\le\alpha\le\tilde{q}\tilde{r})\end{cases}$$

$$C_k^{(rq,\tilde{r}\tilde{q})}(M;\tilde{M})=C_{r\tilde{r}k}^{(q,\tilde{q})}(M;\tilde{M}),B_k^{(rq,\tilde{r}\tilde{q})}\left(\xi'_{\alpha},M;\tilde{\xi}'_{\alpha},\tilde{M}\right)=B_{r\tilde{r}k}^{(q,\tilde{q})}\left(\xi_{\alpha},M;\tilde{\xi}_{\alpha},\tilde{M}\right)$$

$$A_k^{(rq,\tilde{r}\tilde{q})}\left(\xi'_{\alpha},M;\tilde{\xi}'_{\alpha},\tilde{M}\right)=A_{r\tilde{r}k}^{(q,\tilde{q})}\left(\xi_{\alpha},M;\tilde{\xi}_{\alpha},\tilde{M}\right)$$

$$\sum_{\alpha,\beta=1}^{L}\mathcal{A}\left(\kappa,\frac{\chi}{L}+\frac{2i}{L}(\alpha-\beta)\right)=\mathcal{A}(L\kappa,\chi)$$

$$\sum_{\alpha,\beta=1}^{L}a_0\left(z+\frac{2i}{L}(\alpha-\beta)\right)=\frac{1}{L}a_0(Lz)$$

$$\sum_{\alpha,\beta=1}^{L}i\log\,s_{b=1}\left(z+\frac{i}{L}(\alpha-\beta)\right)=i\log\,s_{b=1}(Lz)$$

$$\sum_{\alpha,\beta=1}^{L}\left[\left(z+\frac{2i}{L}(\alpha-\beta)\right)+\frac{1}{2}\left(z+\frac{2i}{L}(\alpha-\beta)\right)^3\right]=z+\frac{1}{2}L^2z^3$$

$$\sum_{\alpha,\beta=1}^{L}(\alpha-\beta)f(\alpha-\beta)=\sum_{\alpha=1}^{L-1}\alpha(L-\alpha)(f(\alpha)-f(\alpha-L)).$$

$$\sum_{\alpha,\beta=1}^{L}\left(z+\frac{2i}{L}(\alpha-\beta)\right)\mathrm{Li}_2\left(e^{\pi\left(z+\frac{2i}{L}(\alpha-\beta)\right)}\right)=z\mathrm{Li}_2(e^{\pi Lz})$$

$$\sum_{\alpha,\beta=1}^{L}\mathrm{Li}_3\left(e^{\pi\left(z+\frac{2i}{L}(\alpha-\beta)\right)}\right)=L^{-1}\mathrm{Li}_3(e^{\pi Lz})$$

$$\sum_{\alpha,\beta=1}^{L}a_0\left(\frac{1}{L}z+\frac{2i}{L}(\alpha-\beta)\right)=\frac{1}{L}\left(-\frac{\pi^3}{6}z-\frac{\pi^3}{12}z^3-\pi z\mathrm{Li}_2(e^{\pi z})+2\mathrm{Li}_3(e^{\pi z})\right).$$

$$\sum_{\alpha,\beta=1}^{L}\left[\frac{1}{6}+\left(z+\frac{i}{L}(\alpha-\beta)\right)^2\right]=\frac{1}{6}+L^2z^2$$

$$\sum_{\alpha,\beta=1}^{L}\left(z+\frac{2i}{L}(\alpha-\beta)\right)\log\left(1-e^{2\pi\left(z+\frac{i}{L}(\alpha-\beta)\right)}\right)=Lz\log\,(1-e^{2\pi Lz}).$$

$$\sum_{\alpha,\beta=1}^{L} \mathrm{Li}_2\left(e^{2\pi\left(z+\frac{i}{L}(\alpha-\beta)\right)}\right) = \mathrm{Li}_2(e^{2\pi Lz}).$$

$$\sum_{\alpha,\beta=1}^{L} i\log\, s_{b=1}\left(z+\frac{i}{L}(\alpha-\beta)\right) = \frac{\pi}{12}+\frac{\pi}{2}L^2z^2 - Lz\log\left(1-e^{2\pi Lz}\right) - \frac{1}{2\pi}\mathrm{Li}_2(e^{2\pi Lz}).$$

$$Z^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(N;\zeta,m,y_\alpha) = \frac{1}{N!}\int_{-\infty}^{\infty}\prod_{i=1}^{N}\frac{d\lambda_i}{2\pi}e^{i\zeta\sum_{i=1}^{N}\lambda_i}\prod_{i<j}^{N}2\sinh\frac{b\lambda_{ij}}{2}\prod_{i<j}^{N}2\sinh\frac{\lambda_{ij}}{2b}$$
$$\times\prod_{i,j}^{N}\mathcal{D}_b\left(\frac{\lambda_{ij}}{2\pi}+m\right)\prod_{\alpha=1}^{N_{\mathrm{f}}}\prod_{i=1}^{N}\mathcal{D}_b\left(\frac{\lambda_i}{2\pi}+y_\alpha\right)$$

$$Z^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(N;\zeta,m_b,y_\alpha) = \frac{1}{N!}\int_{-\infty}^{\infty}\left(\prod_{i=1}^{N}\frac{d\lambda_i}{2\pi}\right)e^{i\zeta\sum_{i=1}^{N}\lambda_i}\prod_{i<j}^{N}2\sinh\frac{b\lambda_{ij}}{2}\prod_{i<j}^{N}2\sinh\frac{\lambda_{ij}}{2b}$$
$$\times\prod_{i,j}^{N}\frac{s_b\left(\frac{\lambda_{ij}}{2\pi}+\frac{i}{2}(b-b^{-1})\right)}{s_b\left(\frac{\lambda_{ij}}{2\pi}-ib^{-1}\right)}\prod_{\alpha=1}^{N_{\mathrm{f}}}\prod_{i=1}^{N}\mathcal{D}_b\left(\frac{\lambda_i}{2\pi}+y_\alpha\right)$$

$$\prod_{i,j}^{N}s_b\left(\frac{\lambda_{ij}}{2\pi}+\frac{i}{2}(b-b^{-1})\right) = \left(\sqrt{\frac{s_b\left(\frac{i}{2}(b-b^{-1})\right)}{s_b\left(-\frac{i}{2}(b-b^{-1})\right)}}\right)^{N}\prod_{i<j}^{N}\frac{s_b\left(\frac{\lambda_{ij}}{2\pi}+\frac{i}{2}(b-b^{-1})\right)}{s_b\left(\frac{\lambda_{ij}}{2\pi}-\frac{i}{2}(b-b^{-1})\right)}$$
$$= b^{-N}\prod_{i<j}^{N}\frac{2\sinh\left(\frac{\lambda_{ij}}{2b}\right)}{2\sinh\left(\frac{b\lambda_{ij}}{2}\right)}$$

$$\prod_{i,j}^{N}\frac{1}{s_b\left(\frac{\lambda_{ij}}{2\pi}-ib^{-1}\right)} = \left(\sqrt{\frac{s_b(ib^{-1})}{s_b(-ib^{-1})}}\right)^{N}\prod_{i<j}^{N}\frac{s_b\left(\frac{\lambda_{ij}}{2\pi}+ib^{-1}\right)}{s_b\left(\frac{\lambda_{ij}}{2\pi}-ib^{-1}\right)} = \prod_{i,j}^{N}\frac{1}{2\cosh\left(\frac{\lambda_{ij}}{2b}+\frac{i}{2b^2}\right)}.$$

$$Z^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(N;\zeta,m_b,y_\alpha)$$
$$= \frac{1}{N!}\int_{-\infty}^{\infty}\left(\prod_{i=1}^{N}\frac{d\lambda_i}{2\pi}\right)e^{ib\zeta\sum_{i=1}^{N}\lambda_i}\frac{\prod_{i<j}^{N}\left(2\sinh\frac{\lambda_{ij}}{2}\right)^2}{\prod_{i,j}^{N}2\cosh\left(\frac{\lambda_{ij}}{2}+\frac{\pi i}{2b^2}\right)}\prod_{\alpha=1}^{N_{\mathrm{f}}}\prod_{i=1}^{N}\mathcal{D}_b\left(\frac{b}{2\pi}\lambda_i+y_\alpha\right)$$

$$Z^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(N;\zeta,m_b,y_\alpha) = \frac{1}{N!}\int_{-\infty}^{\infty}\left(\prod_{i=1}^{N}\frac{d\lambda_i}{2\pi}\right)\det\left(\left[\langle\lambda_i|\hat{\rho}^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(\hat{x},\hat{p};\zeta,m_b,y_\alpha)|\lambda_j\rangle\right]_{i,j}^{N\times N}\right)$$

$$\hat{\rho}^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(\hat{x},\hat{p};\zeta,m_b,y_\alpha) = e^{ib\zeta\hat{x}}\left(\prod_{\alpha=1}^{N_{\mathrm{f}}}\mathcal{D}_b\left(\frac{b}{2\pi}\hat{x}+y_\alpha\right)\right)\frac{e^{-\frac{1}{2b^2}\hat{p}}}{2\cosh\frac{\hat{p}}{2}}.$$

$$\hat{\rho}^{\mathrm{SYM}}_{b=1,N_{\mathrm{f}}}(\hat{x},\hat{p};\zeta,m,y_\alpha)=\frac{e^{i\zeta\hat{x}}}{\prod_{\alpha=1}^{N_{\mathrm{f}}}\ 2\cosh\dfrac{\hat{x}+2\pi y_\alpha}{2}}\frac{e^{im\hat{p}}}{2\cosh\dfrac{\hat{p}}{2}}$$

$(nN_{\mathrm{f}},1)_{k=1}$ with $\left\{M=\dfrac{2b}{nN_{\mathrm{f}}}\zeta,\eta_\alpha=y'_\alpha;\tilde{M}=ib^{-2}\right\}\leftrightarrow \mathrm{SYM}_{b=\sqrt{2n-1},N_{\mathrm{f}}}$ with $\{\zeta,m_b,y_\alpha\}$

$$y'_\alpha=\begin{cases}\dfrac{2}{b}y_1+\dfrac{2i}{b^2}\left[\dfrac{1+n}{2}-\alpha\right] & (1\le\alpha\le n)\\ \dfrac{2}{b}y_2+\dfrac{2i}{b^2}\left[\dfrac{1+n}{2}-(\alpha-n)\right] & (n+1\le\alpha\le 2n)\\ \vdots & \vdots\\ \dfrac{2}{b}y_{N_{\mathrm{f}}}+\dfrac{2i}{b^2}\left[\dfrac{1+n}{2}-(\alpha-(N_{\mathrm{f}}-1)n)\right] & \left((N_{\mathrm{f}}-1)n+1\le\alpha\le N_{\mathrm{f}}n\right)\end{cases}.$$

$$A^{\mathrm{SYM}}_{b=\sqrt{2n-1},N_{\mathrm{f}}}(\zeta,m_b,y_\alpha)=\frac{1}{4}\sum_{\pm}\left[A^{\mathrm{ABJM}}\left(\frac{(b^2+1)N_{\mathrm{f}}\pm 4ib\zeta}{2}\right)+\sum_{\alpha,\beta=1}^{N_{\mathrm{f}}}\sum_{\alpha',\beta'=1}^{n}\mathcal{A}\left(1\pm b^{-2},\frac{2}{b}(y_\alpha-y_\beta)+\frac{2i}{b^2}(\alpha'-\beta')\right)\right].$$

$$A^{\mathrm{SYM}}_{b=\sqrt{2n-1},N_{\mathrm{f}}=1}(\zeta,m_b)=\frac{1}{4}\left[\sum_{\pm}A^{\mathrm{ABJM}}\left(\frac{b^2+1\pm 4ib\zeta}{2}\right)+A^{\mathrm{ABJM}}(2n-2)-A^{\mathrm{ABJM}}(4n-2)\right]$$

$$\sum_{\alpha,\beta=1}^{n}\left[\mathcal{A}\left(\frac{2n}{2n-1},\frac{2i}{2n-1}(\alpha-\beta)\right)+\mathcal{A}\left(\frac{2n-2}{2n-1},\frac{2i}{2n-1}(\alpha-\beta)\right)\right]=A^{\mathrm{ABJM}}(2n-2)-A^{\mathrm{ABJM}}(4n-2)$$

$$\sum_{\alpha,\beta=1}^{n}\left[\mathcal{A}\left(\frac{2n}{2n-1},\frac{\chi}{2n-1}+\frac{2i}{2n-1}(\alpha-\beta)\right)+\mathcal{A}\left(\frac{2n-2}{2n-1},\frac{\chi}{2n-1}+\frac{2i}{2n-1}(\alpha-\beta)\right)\right]=\mathcal{A}(2n-2,\chi)-\mathcal{A}(4n-2,\chi)$$

$$C^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(\zeta,m_b)=\frac{4b^4N_{\mathrm{f}}}{\pi^2(b^2-1)\left((b^2+1)^2N_{\mathrm{f}}^2+16b^2\zeta^2\right)}$$

$$B^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(\zeta,m_b,y_\alpha)=\frac{1}{b^2-1}\left[\frac{(3b^4+1)N_{\mathrm{f}}}{3\left((b^2+1)^2N_{\mathrm{f}}^2+16b^2\zeta^2\right)}-b^2\sum_{\alpha=1}^{N_{\mathrm{f}}}y_\alpha^2-\frac{N_{\mathrm{f}}}{24}(b^4-b^2+2)\right]$$

$$A^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(\zeta,m_b,y_\alpha)=\frac{1}{4}\left[\sum_{\pm}A^{\mathrm{ABJM}}\left(\frac{(b^2+1)N_{\mathrm{f}}\pm 4ib\zeta}{2}\right)+\sum_{\alpha,\beta=1}^{N_{\mathrm{f}}}\left(\mathcal{A}\left(b^2-1,2b(y_\alpha-y_\beta)\right)-\mathcal{A}\left(2b^2,2b(y_\alpha-y_\beta)\right)\right)\right]$$

$$C_{b,N_{\mathrm{f}}}^{\mathrm{SYM}}(\zeta,m) = \left(\frac{2}{\pi(b+b^{-1})^2\sqrt{2N_{\mathrm{f}}\Delta_1\Delta_2\Delta_3\Delta_4}}\right)^2$$

$$B_{b,N_{\mathrm{f}}}^{\mathrm{SYM}}(\zeta,m) = \frac{N_{\mathrm{f}}}{24} - \frac{N_{\mathrm{f}}}{12}\left(\frac{1}{\Delta_1}+\frac{1}{\Delta_2}\right) - \frac{1}{12N_{\mathrm{f}}}\left(\frac{1}{\Delta_3}+\frac{1}{\Delta_4}\right) - \frac{4}{3(b+b^{-1})^2}\left(-\frac{N_{\mathrm{f}}}{8\Delta_1\Delta_2} + \frac{\Delta_1^2+\Delta_2^2-2(\Delta_1+\Delta_2)+\Delta_1\Delta_2}{8N_{\mathrm{f}}\Delta_1\Delta_2\Delta_3\Delta_4}\right)$$

$$A_{b,N_{\mathrm{f}}}^{\mathrm{SYM}}(\zeta,m_b) = \frac{1}{4}\sum_{\pm} A^{\mathrm{ABJM}}\left(\frac{(b^2+1)N_{\mathrm{f}} \pm 4ib\zeta}{2}\right) + \frac{N_{\mathrm{f}}^2}{4}\left(A^{\mathrm{ABJM}}(b^2-1) - A^{\mathrm{ABJM}}(2b^2)\right),$$

$$\Delta_1 = \frac{1}{2}+\frac{2mi}{b+b^{-1}}, \Delta_2 = \frac{1}{2}-\frac{2mi}{b+b^{-1}}, \Delta_3 = \frac{1}{2}-\frac{2\zeta i}{(b+b^{-1})N_{\mathrm{f}}}, \Delta_4 = \frac{1}{2}+\frac{2\zeta i}{(b+b^{-1})N_{\mathrm{f}}}.$$

$$C_{b,N_{\mathrm{f}}=1}^{\mathrm{SYM}}(\zeta,m_b) = \frac{4b^4}{\pi^2(b^2-1)((b^2+1)^2+16b^2\zeta^2)}$$

$$B_{b,N_{\mathrm{f}}=1}^{\mathrm{SYM}}(\zeta,m_b) = -\frac{b^8+b^6(16\zeta^2+1)-b^4(16\zeta^2+23)+b^2(32\zeta^2+3)-6}{24(b^2-1)((b^2+1)^2+16b^2\zeta^2)}$$

$$Z_{b,N_{\mathrm{f}}=2}^{\mathrm{SYM,pert}}(1;\zeta,m_b,y_\alpha) = e^{A_{b,2}^{\mathrm{SYM}}(\zeta,m_b,y_\alpha)} C_{b,2}^{\mathrm{SYM}}(\zeta,m_b)^{-\frac{1}{3}} \mathrm{Ai}\left[C_{b,2}^{\mathrm{SYM}}(\zeta,m_b)^{-\frac{1}{3}}\left(1-B_{b,2}^{\mathrm{SYM}}(\zeta,m_b,y_\alpha)\right)\right]$$

$$Z_{b,N_{\mathrm{f}}=2}^{\mathrm{SYM}}(1;\zeta,m_b,y_\alpha) = \mathcal{D}_b(m_b)\int_{-\infty}^{\infty}\frac{d\lambda}{2\pi}e^{i\zeta\sum_{i=1}^{N}\lambda_i}\prod_{\alpha=1}^{2}\mathcal{D}_b\left(\frac{\lambda}{2\pi}+y_\alpha\right)$$

$$A_{b,N_{\mathrm{f}}}^{\mathrm{SYM}}(\zeta,m_b,y_\alpha = y_\alpha(b)) = \frac{1}{4}\left[\sum_{\pm} A^{\mathrm{ABJM}}\left(\frac{(b^2+1)N_{\mathrm{f}} \pm 4ib\zeta}{2}\right) + A^{\mathrm{ABJM}}\left((b^2-1)N_{\mathrm{f}}\right) - A^{\mathrm{ABJM}}(2b^2N_{\mathrm{f}})\right]$$

$$y_\alpha(b) = \frac{i}{bN_{\mathrm{f}}}\left(\frac{1+N_{\mathrm{f}}}{2}-\alpha\right).$$

$$\hat{\mathcal{O}}_{b,N_{\mathrm{f}}}^{\mathrm{SYM}}(\hat{x},\hat{p};\zeta,m_b,y_\alpha) = e^{-\frac{ib\zeta}{2}\hat{x}}\prod_{\alpha=1}^{N_{\mathrm{f}}} s_b\left(\frac{b}{2\pi}\hat{x}+y_\alpha-\frac{i}{4}Q\right)\left(e^{\left(\frac{1}{2}+\frac{1}{2b^2}\right)\hat{p}}+e^{-\left(\frac{1}{2}-\frac{1}{2b^2}\right)\hat{p}}\right)\prod_{\alpha=1}^{N_{\mathrm{f}}}\frac{e^{-\frac{ib\zeta}{2}\hat{x}}}{s_b\left(\frac{b}{2\pi}\hat{x}+y_\alpha+\frac{i}{4}Q\right)}$$

$$\hat{\mathcal{O}}_{b,N_{\mathrm{f}}}^{\mathrm{SYM}}(\hat{x},\hat{p};\zeta,m_b,y_\alpha) = \hat{A}\hat{\rho}_{b,N_{\mathrm{f}}}^{\mathrm{SYM}}(\hat{x},\hat{p};\zeta,m_b,y_\alpha)^{-1}\hat{A}^{-1},$$

$$\hat{A} = e^{-\frac{ib\zeta}{2}\hat{x}}\prod_{\alpha=1}^{N_{\mathrm{f}}} s_b\left(\frac{b}{2\pi}\hat{x}+y_\alpha-\frac{i}{4}Q\right).$$

$$e^{c_1\hat{x}}e^{c_2\hat{p}} = e^{c_1\pi i}e^{c_1\hat{x}+c_2\hat{p}} = e^{2c_1c_2\pi i}e^{c_2\hat{p}}e^{c_1\hat{x}},$$

$$\hat{\mathcal{O}}^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(\hat{x},\hat{p};\zeta,m_b,y_\alpha)$$
$$= e^{-ib\zeta\hat{x}+\left(\frac{1}{2}+\frac{1}{2b^2}\right)\hat{p}} + e^{-\frac{1}{2}\left(\frac{1}{2}-\frac{1}{2b^2}\right)\hat{p}} e^{-ib\zeta\hat{x}} \left(\prod_{\alpha=1}^{N_{\mathrm{f}}} 2\cosh\left(\frac{b^2}{2}\hat{x} + \pi b y_\alpha\right)\right) e^{-\frac{1}{2}\left(\frac{1}{2}-\frac{1}{2b^2}\right)\hat{p}}$$

$$\hat{X} = \left(\frac{b^2 N_{\mathrm{f}}}{2} - ib\zeta\right)\hat{x} - \left(\frac{1}{2} - \frac{1}{2b^2}\right)\hat{p}, \hat{P} = -ib\zeta\hat{x} + \left(\frac{1}{2}+\frac{1}{2b^2}\right)\hat{p}.$$

$$\hat{\mathcal{O}}^{\mathrm{SYM}}_{b,N_{\mathrm{f}}}(\hat{x},\hat{p};\zeta,m_b,y_\alpha(b)) = \hat{\mathcal{O}}^{\mathbb{P}(1,\mathfrak{m},\mathfrak{n})}(\hat{X},\hat{P}),$$

$$\hat{\mathcal{O}}^{\mathbb{P}(1,\mathfrak{m},\mathfrak{n})}(\hat{X},\hat{P}) = e^{\hat{X}} + e^{\hat{P}} + e^{-\mathfrak{m}\hat{X}-\mathfrak{n}\hat{P}}$$

$$\begin{aligned}
C^{\mathbb{P}(1,\mathfrak{m},\mathfrak{n})}(\hbar') =& \frac{(\mathfrak{m}+\mathfrak{n}+1)^2}{\mathfrak{m}\mathfrak{n}}\frac{1}{4\pi\hbar'},\\
B^{\mathbb{P}(1,\mathfrak{m},\mathfrak{n})}(\hbar') =& \frac{\mathfrak{m}^2+\mathfrak{m}\mathfrak{n}+\mathfrak{n}^2+\mathfrak{m}+\mathfrak{n}+1}{12\mathfrak{m}\mathfrak{n}}\frac{\pi}{\hbar'} - \frac{\mathfrak{m}+\mathfrak{n}+1}{48\pi}\hbar'\\
A^{\mathbb{P}(1,\mathfrak{m},\mathfrak{n})}(\hbar') =& \frac{1}{4}\left[A^{\mathrm{ABJM}}\left(\frac{\hbar'}{\pi}\right) + A^{\mathrm{ABJM}}\left(\frac{\mathfrak{m}\hbar'}{\pi}\right)\right.\\
&\left. + A^{\mathrm{ABJM}}\left(\frac{\mathfrak{n}\hbar'}{\pi}\right) - A^{\mathrm{ABJM}}\left(\frac{(\mathfrak{m}+\mathfrak{n}+1)\hbar'}{\pi}\right)\right].
\end{aligned}$$

$$\mathfrak{m} = \frac{(b+b^{-1})N_{\mathrm{f}} + 4i\zeta}{(b+b^{-1})N_{\mathrm{f}} - 4i\zeta}, \mathfrak{n} = \frac{2(b-b^{-1})N_{\mathrm{f}}}{(b+b^{-1})N_{\mathrm{f}} - 4i\zeta}, \hbar' = \frac{(b^2+1)N_{\mathrm{f}} - 4ib\zeta}{2}\pi.$$

$$\zeta(s) = \sum_{n=1}^{\infty}\frac{1}{n^s}$$

$$\zeta(s) = \frac{1}{\Gamma(s)}\int_0^\infty \frac{x^{s-1}}{e^x-1}dx$$

$$\zeta(2n) = \frac{(-1)^{n+1}B_{2n}(2\pi)^{2n}}{2(2n)!}$$

$$\zeta(-n) = -\frac{B_{n+1}}{n+1}$$

$$\mathrm{Li}_s(z) = \sum_{n=1}^{\infty}\frac{z^n}{n^s}$$

$$\mathrm{Li}_s(1) = \zeta(s)$$

$$\mathrm{Li}_1(z) = -\log(1-z)$$

$$\frac{\partial}{\partial z}\mathrm{Li}_s(z) = \frac{1}{z}\mathrm{Li}_{s-1}(z)$$

$$\sum_{n=1}^{p} \mathrm{Li}_s\left(ze^{2\pi i\frac{n}{p}}\right) = p^{1-s}\mathrm{Li}_s(z^p)$$

$$\mathrm{Li}_n(-e^x) + (-1)^n\mathrm{Li}_n(-e^{-x}) = -\frac{(2\pi i)^n}{n!}B_n\left(\frac{x}{2\pi i}+\frac{1}{2}\right)$$

$$s_b(z) = \prod_{\ell,m=0}^{\infty} \frac{\ell b + mb^{-1} + \frac{Q}{2} - iz}{\ell b + mb^{-1} + \frac{Q}{2} + iz}$$

$$i\log\, s_b(z) = \frac{\pi}{2}z^2 + \frac{\pi}{24}(b^2+b^{-2}) + i\int_{\mathbb{R}+i0^+}\frac{dt}{t}\frac{e^{-2izt}}{4\sinh(bt)\sinh(b^{-1}t)}$$

$$s_b(0) = 1, s_b(z) = s_{b^{-1}}(z), s_b(z)s_b(-z) = 1, \overline{s_b(z)} = s_b(-\bar{z})$$

$$\frac{s_b\left(z+\frac{i}{2}b^{\pm 1}\right)}{s_b\left(z-\frac{i}{2}b^{\pm 1}\right)} = \frac{1}{2\cosh(\pi b^{\pm 1}z)}$$

$$i\log\, s_{b=1}(z) = \frac{\pi}{12} + \frac{\pi}{2}z^2 - z\log(1-e^{2\pi z}) - \frac{1}{2\pi}\mathrm{Li}_2(e^{2\pi z})$$

$$\frac{d}{dz}i\log\, s_{b=1}(z) = -\frac{\pi z}{\tanh\,\pi z}$$

$$\mathcal{D}_b(z) = \frac{s_b\left(z+\frac{i}{4}Q\right)}{s_b\left(z-\frac{i}{4}Q\right)}$$

$$\mathcal{D}_{b=\sqrt{2n-1}}(\mu) = \prod_{j=1}^{n}\frac{1}{2\cosh\left(\frac{\pi}{b}\mu + \frac{\pi i}{b^2}\left(\frac{n+1}{2}-j\right)\right)}$$

$$\mathcal{D}_{\sqrt{2}}(\mu) = \frac{1}{2^{\frac{1}{4}}(2\cosh(2\sqrt{2}\pi\mu))^{\frac{1}{8}}(\sqrt{2}\cosh(\sqrt{2}\pi\mu)+1)^{\frac{1}{2}}} \times \exp\left[-\sqrt{2}\mu\arctan\left(e^{-2\sqrt{2}\pi\mu}\right) + \frac{i}{4\pi}\left(\mathrm{Li}_2\left(ie^{-2\sqrt{2}\pi\mu}\right) - \mathrm{Li}_2\left(-ie^{-2\sqrt{2}\pi\mu}\right)\right)\right]$$

$$\mathcal{Z}_{2\ell}^{(q,\tilde{q})}(s; v_\alpha, M; \tilde{v}_\alpha, \tilde{M}) = D_{2\ell}^{(q,\tilde{q})}(s; v_\alpha, M; \tilde{v}_\alpha, \tilde{M})\mathcal{Z}_0^{(q,\tilde{q})}(s; v_\alpha, M; \tilde{v}_\alpha, \tilde{M})$$

$$\hat{\rho}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi}, M; \frac{\hbar\tilde{v}_\alpha}{\pi}, \tilde{M}\right)^s = e^{-s\hat{H}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)} \text{ around } \hat{H}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi}, M; \frac{\hbar\tilde{v}_\alpha}{\pi}, \tilde{M}\right) = H_W(x,p) \text{ as}$$

$$\hat{\rho}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi}, M; \frac{\hbar\tilde{v}_\alpha}{\pi}, \tilde{M}\right)^s = \sum_{r=0}^{\infty} e^{-sH_W(x,p)}\frac{(-s)^r}{r!}\left(\hat{H}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi}, M; \frac{\hbar\tilde{v}_\alpha}{\pi}, \tilde{M}\right) - H_W(x,p)\right)^r$$

$$\left(\hat{\rho}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)^s\right)_W=\sum_{r=0}^{\infty}e^{-sH_W(x,p)}\frac{(-s)^r}{r!}\mathcal{G}_r(x,p)$$

$$\mathcal{G}_r(x,p)=\left(\left(\hat{H}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)-H_W(x,p)\right)^r\right)_W$$

$$\mathcal{G}_0(x,p)=1,$$
$$\mathcal{G}_1(x,p)=0,$$
$$\mathcal{G}_r(x,p)=(-1)^{r-1}(r-1)H_W(x,p)^r$$
$$+\sum_{r'=2}^{r}\binom{r}{r'}(-1)^{r-r'}H_W(x,p)^{r-r'}(\underbrace{H_W(x,p)\star\cdots\star H_W(x,p)}_{r'}),(r\geq 2)$$

$$\mathcal{G}_r(x,p)=\sum_{a=2\left\lfloor\frac{r+2}{3}\right\rfloor}^{}\hbar^a\mathcal{G}_r^{(a)}$$

$$H_W(x,p)=\sum_{a=0}^{\infty}\hbar^a H_W^{(a)}$$

$$\mathrm{Tr}\hat{\rho}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)^s$$

$$\mathrm{Tr}\hat{\rho}_k^{(q,\tilde{q})}\left(\hat{x},\hat{p};\frac{\hbar v_\alpha}{\pi},M;\frac{\hbar\tilde{v}_\alpha}{\pi},\tilde{M}\right)^s=\int\frac{dxdp}{2\pi\hbar}e^{-sH_W^{(0)}}\Delta_{\mathrm{WK}}$$

$$\Delta_{\mathrm{WK}}=\left(1+\sum_{r=1}^{\infty}\frac{(-s)^r}{r!}\left(\sum_{a=1}^{\infty}\hbar^a H_W^{(a)}\right)^r\right)\left(1+\sum_{r'=2}^{\infty}\frac{(-s)^{r'}}{r!}\sum_{a'=2\left\lfloor\frac{r'+2}{3}\right\rfloor}^{\infty}\hbar^{a'}\mathcal{G}_{r'}^{(a')}\right)$$

$$\mathcal{Z}_2^{(q,\tilde{q})}(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M})=\int\frac{dxdp}{2\pi}e^{-sH_W^{(0)}}\left(-sH_W^{(2)}+\frac{s^2}{2}\left(H_W^{(1)}\right)^2+\frac{s^2}{2}\mathcal{G}_2^{(2)}-\frac{s^3}{6}\mathcal{G}_3^{(2)}\right)$$

$$H_W^{(0)}=q\log\,2\cosh\frac{x}{2}-\frac{iqMx}{2}+\tilde{q}\log\,2\cosh\frac{p}{2}-\frac{i\tilde{q}\tilde{M}x}{2}$$
$$H_W^{(1)}=0$$
$$H_W^{(2)}=\frac{1}{F(x)^2}\left(\frac{1}{2}\sum_{\alpha=1}^{q}v_\alpha^2-\frac{q\tilde{q}^2(1-\tilde{M}^2)}{48}\right)+\frac{1}{F(p)^2}\left(\frac{1}{2}\sum_{\alpha=1}^{\tilde{q}}\tilde{v}_\alpha^2+\frac{q^2\tilde{q}(1-M^2)}{96}\right)$$
$$+\frac{iq\tilde{q}^2\tilde{M}F'(p)}{12F(x)^2F(p)}-\frac{iq^2\tilde{q}MF'(x)}{24F(x)F(p)^2}+\frac{1}{F(x)^2F(p)^2}\left(-\frac{q^2\tilde{q}}{24}+\frac{q\tilde{q}^2}{12}\right)$$
$$\mathcal{G}_2^{(2)}=\frac{1}{4}\left[\left(\partial_x\partial_p H_W^{(0)}\right)^2-\partial_x^2H_W^{(0)}\partial_p^2H_W^{(0)}\right]=-\frac{q\tilde{q}}{4F(x)^2F(p)^2}$$
$$\mathcal{G}_3^{(2)}=\frac{1}{4}\left[2\partial_x H_W^{(0)}\partial_p H_W^{(0)}\partial_x\partial_p H_W^{(0)}-\partial_x^2H_W^{(0)}\left(\partial_p H_W^{(0)}\right)^2-\left(\partial_x H_W^{(0)}\right)^2\partial_p^2H_W^{(0)}\right]$$
$$=-\frac{q\tilde{q}^2(1-\tilde{M}^2)}{16F(x)^2}-\frac{q^2\tilde{q}(1-M^2)}{16F(p)^2}+\frac{iq\tilde{q}^2\tilde{M}F'(p)}{4F(x)^2F(p)}+\frac{iq^2\tilde{q}MF'(x)}{4F(x)F(p)^2}+\frac{q\tilde{q}(q+\tilde{q})}{4F(x)^2F(p)^2}$$

$$\mathcal{Z}_2^{(q,\tilde{q})}\left(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M}\right)$$
$$=\frac{1}{2\pi}\Bigg[-s\Bigg(\left(\frac{1}{2}\sum_{\alpha=1}^{q}v_\alpha^2-\frac{q\tilde{q}^2\left(1-\tilde{M}^2\right)}{48}\right)I_1\left(\frac{iqMs}{2},qs+2\right)I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s\right)$$
$$+\left(\frac{1}{2}\sum_{\alpha=1}^{\tilde{q}}\tilde{v}_\alpha^2+\frac{q^2\tilde{q}(1-M^2)}{96}\right)I_1\left(\frac{iqMs}{2},qs\right)I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s+2\right)$$
$$+\frac{iq\tilde{q}^2\tilde{M}}{12}I_1\left(\frac{iqMs}{2},qs+2\right)I_2\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s+1\right)$$
$$-\frac{iq^2\tilde{q}M}{24}I_2\left(\frac{iqMs}{2},qs+1\right)I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s+2\right)$$
$$+\left(-\frac{q^2\tilde{q}}{24}+\frac{q\tilde{q}^2}{12}\right)I_1\left(\frac{iqMs}{2},qs+2\right)I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s+2\right)\Bigg)$$
$$+\frac{s^2}{2}\left(-\frac{q\tilde{q}}{4}I_1\left(\frac{iqMs}{2},qs+2\right)I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s+2\right)\right)$$
$$-\frac{s^3}{6}\Bigg(-\frac{q\tilde{q}^2\left(1-\tilde{M}^{\,2}\right)}{16}I_1\left(\frac{iqMs}{2},qs+2\right)I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s\right)$$
$$-\frac{q^2\tilde{q}(1-M^2)}{16}I_1\left(\frac{iqMs}{2},qs\right)I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s+2\right)$$
$$+\frac{iq\tilde{q}^2\tilde{M}}{4}I_1\left(\frac{iqMs}{2},qs+2\right)I_2\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s+1\right)$$
$$+\frac{iq^2\tilde{q}M}{4}I_2\left(\frac{iqMs}{2},qs+1\right)I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s+2\right)$$
$$+\frac{q\tilde{q}(q+\tilde{q})}{4}I_1\left(\frac{iqMs}{2},qs+2\right)I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s+2\right)\Bigg)\Bigg]$$

$$\mathcal{Z}_2^{(q,\tilde{q})}\left(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M}\right)=\frac{I_1\left(\frac{iqMs}{2},qs\right)I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s\right)}{2\pi}D_2^{(q,\tilde{q})}\left(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M}\right)$$

$$D_2^{(q,\tilde{q})}\left(s;v_\alpha,M;\tilde{v}_\alpha,\tilde{M}\right)=\frac{s^2}{384(1+qs)(1+\tilde{q}s)}\Big(q^2\tilde{q}^2(1+M^2)\left(1+\tilde{M}^2\right)(1-s^2)$$
$$-48q(1+M^2)(1+\tilde{q}s)\sum_{\alpha=1}^{q}v_\alpha^2-48\tilde{q}\left(1+\tilde{M}^2\right)(1+qs)\sum_{\alpha=1}^{\tilde{q}}\tilde{v}_\alpha^2\Big)$$

$$\frac{1}{F(x)^{2a_1}F(p)^{2a_2}},\frac{F'(x)}{F(x)^{2a_1+1}F(p)^{2a_2}},\frac{F'(p)}{F(x)^{2a_1}F(p)^{2a_2+1}},\frac{F'(x)F'(p)}{F(x)^{2a_1+1}F(p)^{2a_2+1}},$$

$$I_1\left(\frac{i\tilde{q}\tilde{M}s}{2},\tilde{q}s\right)\text{ﻑ}\,\mathcal{Z}_2^{(q,\tilde{q})}\left(s;v_\alpha,M,\tilde{v}_\alpha,\tilde{M}\right)\text{ﻑ}\,\partial_x^{a_1}U(x)\text{ﻑ}\,\partial_p^{a_2}T(x)\text{ﻑ}\,\frac{1}{F(x)^{2a_1}}\text{ﻑ}\,\frac{F'(x)}{F(x)^{2a_1+1}}\text{ﻑ}\,\frac{F'(x)}{F(x)}$$

$$\mathrm{Vec}\left(\frac{1}{F(x)^{2a_1}},\frac{F'(x)}{F(x)^{2a_2+1}};a_1,a_2\geq 0\right)=\mathrm{Pol}\left[\frac{F'(x)}{F(x)}\right]$$

$$\partial_x^a U(x) \in \mathrm{Pol}\left[\frac{F'(x)}{F(x)}\right]$$

$$\partial_x U(x) = \frac{qF'(x)}{F(x)} - \frac{iqM}{2} + \sum_{\alpha=1}^{q} \partial_x \log\left[\cosh\frac{\hbar v_\alpha}{2} + 2\sinh\frac{\hbar v_\alpha}{2}\frac{F'(x)}{F(x)}\right]$$

$$\partial_x \mathrm{Pol}\left[\frac{F'(x)}{F(x)}\right] \subset \mathrm{Pol}\left[\frac{F'(x)}{F(x)}\right]$$

$$V_\nu = \sum_{\alpha=1}^{q} v_\alpha^\nu, \tilde{V}_\nu = \sum_{\alpha=1}^{\tilde{q}} \tilde{v}_\alpha^\nu$$

$$\begin{aligned} D_4^{(q,\tilde{q})}(s; v_\alpha, 0; \tilde{v}_\alpha, 0) =& f_{4,\emptyset,\emptyset}^{(q,\tilde{q})}(s;0;0) + f_{4,\{2\},\emptyset}^{(q,\tilde{q})}(s;0;0)V_2 + f_{4,\{2\},\emptyset}^{(\tilde{q},q)}(s;0;0)\tilde{V}_2 + f_{4,\{4\},\emptyset}^{(q,\tilde{q})}(s;0;0)V_4 \\ &+ f_{4,\{2,2\},\emptyset}^{(q,\tilde{q})}(s;0;0)V_2^2 + f_{4,\{2\},\{2\}\emptyset}^{(q,\tilde{q})}(s;0;0)V_2\tilde{V}_2 + f_{4,\{4\},\emptyset}^{(\tilde{q},q)}(s;0;0)\tilde{V}_4 \\ &+ f_{4,\{2,2\},\emptyset}^{(\tilde{q},q)}(s;0;0)\tilde{V}_2^2 \end{aligned}$$

$$f_{4,\emptyset,\emptyset}^{(q,\tilde{q})}(s;0;0) = -\frac{q^3\tilde{q}^3 s^3(1-s^2)(24q+24\tilde{q}+n(80+17q\tilde{q})+n^2(24q+24\tilde{q})+7q\tilde{q}n^3)}{1474560(1+qs)(3+qs)(1+\tilde{q}s)(3+\tilde{q}s)}$$

$$f_{4,\{2\},\emptyset}^{(q,\tilde{q})}(s;0;0) = -\frac{q^2\tilde{q}^2 s^3(1-s^2)(4+qs)}{3072(1+qs)(1+\tilde{q}s)(3+qs)}$$

$$f_{4,\{4\},\emptyset}^{(q,\tilde{q})}(s;0;0) = \frac{q^2 s^3}{192(1+qs)(3+qs)}$$

$$f_{4,\{2,2\},\emptyset}^{(q,\tilde{q})}(s;0;0) = \frac{qs^3(2+qs)}{128(1+qs)(3+qs)}$$

$$f_{4,\{2\},\{2\}\emptyset}^{(q,\tilde{q})}(s;0;0) = \frac{q^2 s^4}{64(1+\tilde{q}s)(1+qs)}$$

$$\begin{aligned} D_6^{(q,\tilde{q})}(s; v_\alpha, 0; \tilde{v}_\alpha, 0) =& f_{6,\emptyset,\emptyset}^{(q,\tilde{q})}(s;0;0) + f_{6,\{2\},\emptyset}^{(q,\tilde{q})}(s;0;0)V_2 + f_{6,\{2\},\emptyset}^{(\tilde{q},q)}(s;0;0)\tilde{V}_2 + f_{6,\{4\},\emptyset}^{(q,\tilde{q})}(s;0;0)V_4 \\ &+ f_{6,\{2,2\},\emptyset}^{(q,\tilde{q})}(s;0;0)V_2^2 + f_{6,\{2\},\{2\}}^{(q,\tilde{q})}(s;0;0)V_2\tilde{V}_2 + f_{6,\{4\},\emptyset}^{(\tilde{q},q)}(s;0;0)\tilde{V}_4 \\ &+ f_{6,\{2,2\},\emptyset}^{(\tilde{q},q)}(s;0;0)\tilde{V}_2^2 + f_{6,\{6\},\emptyset}^{(q,\tilde{q})}(s;0;0)V_6 + f_{6,\{4,2\},\emptyset}^{(q,\tilde{q})}(s;0;0)V_4V_2 \\ &+ f_{6,\{3,3\},\emptyset}^{(q,\tilde{q})}(s;0;0)V_3^2 + f_{6,\{2,2,2\},\emptyset}^{(q,\tilde{q})}(s;0;0)V_2^3 + f_{6,\{4\},\{2\}}^{(q,\tilde{q})}(s;0;0)V_4\tilde{V}_2 \\ &+ f_{6,\{2,2\},\{2\}}^{(q,\tilde{q})}(s;0;0)V_2^2\tilde{V}_2 + f_{6,\{4\},\{2\}}^{(\tilde{q},q)}(s;0;0)V_2\tilde{V}_4 + f_{6,\{2,2\},\{2\}}^{(\tilde{q},q)}(s;0;0)V_2\tilde{V}_2^2 \\ &+ f_{6,\{6\},\emptyset}^{(\tilde{q},q)}(s;0;0)\tilde{V}_6 + f_{6,\{4,2\},\emptyset}^{(\tilde{q},q)}\tilde{V}_4\tilde{V}_2 + f_{6,\{3,3\},\emptyset}^{(\tilde{q},q)}\tilde{V}_3^2 + f_{6,\{2,2,2\},\emptyset}^{(\tilde{q},q)}\tilde{V}_2^3 \end{aligned}$$

$$\begin{aligned} f_{6,\emptyset,\emptyset}^{(q,\tilde{q})}(s;0;0) =& \frac{q^3\tilde{q}^3 s^3(1-s^2)}{3963617280(1+qs)(3+qs)(5+qs)(1+\tilde{q}s)(3+\tilde{q}s)(5+\tilde{q}s)}(1920q^3+1920\tilde{q}^3 \\ &+(7168+5376q^2+14336q\tilde{q}+2944q^3\tilde{q}+5376\tilde{q}^2+2304q^2\tilde{q}^2+2944q\tilde{q}^3)s \\ &+(14336q+576q^3+14336\tilde{q}+17920q^2\tilde{q}+17920q\tilde{q}^2+1656q^3\tilde{q}^2+576\tilde{q}^3+1656q^2\tilde{q}^3)s^2 \\ &+(5376q^2+25088q\tilde{q}+2272q^3\tilde{q}+5376\tilde{q}^2+12272q^2\tilde{q}^2+2272q\tilde{q}^3+367q^3\tilde{q}^3)s^3 \\ &+(576q^3+8960q^2\tilde{q}+8960q\tilde{q}^2+1488q^3\tilde{q}^2+576\tilde{q}^3+1488q^2\tilde{q}^3)s^4 \\ &+(928q^3\tilde{q}+3088q^2\tilde{q}^2+928q\tilde{q}^3+178q^3\tilde{q}^3)s^5+(312q^3\tilde{q}^2+312q^2\tilde{q}^3)s^6+31q^3\tilde{q}^3s^7), \end{aligned}$$

$$f_{6,\{2\},\emptyset}^{(q,\tilde{q})}(s;0;0)=\frac{q^2\tilde{q}^3s^3(1-s^2)}{11796480(1+qs)(3+qs)(5+qs)(1+\tilde{q}s)(3+\tilde{q}s)}(192\tilde{q}+(640+192q^2+288q\tilde{q})s$$
$$+(960q+24q^3+192\tilde{q}+160q^2\tilde{q})s^2+(272q^2+288q\tilde{q}+17q^3\tilde{q})s^3+(24q^3+80q^2\tilde{q})s^4+7q^3\tilde{q}s^5)$$

$$f_{6,\{4\},\emptyset}^{(q,\tilde{q})}(s;0;0)=\frac{q^2\tilde{q}^2s^3(1-s^2)(8+12qs+q^2s^2)}{73728(1+qs)(3+qs)(5+qs)(1+\tilde{q}s)},$$

$$f_{6,\{2,2\},\emptyset}^{(q,\tilde{q})}(s;0;0)=\frac{q\tilde{q}^2s^3(1-s^2)(2+qs)(8+8qs+q^2s^2)}{49152(1+qs)(3+qs)(5+qs)(1+\tilde{q}s)},$$

$$f_{6,\{2\},\{2\}\emptyset}^{(q,\tilde{q})}(s;0;0)=\frac{q^2\tilde{q}^2s^4(1-s^2)(4+qs)(4+\tilde{q}s)}{24576(1+qs)(3+qs)(1+\tilde{q}s)(3+\tilde{q}s)},$$

$$f_{6,\{6\},\emptyset}^{(q,\tilde{q})}(s;0;0)=-\frac{q^2s^3(1+2qs)}{5760(1+qs)(3+qs)(5+qs)},$$

$$f_{6,\{4,2\},\emptyset}^{(q,\tilde{q})}(s;0;0)=-\frac{qs^3(2+qs)^2}{1536(1+qs)(3+qs)(5+qs)},$$

$$f_{6,\{3,3\},\emptyset}^{(q,\tilde{q})}(s;0;0)=\frac{qs^3(2+qs)}{1152(1+qs)(3+qs)(5+qs)},$$

$$f_{6,\{2,2,2\},\emptyset}^{(q,\tilde{q})}(s;0;0)=-\frac{qs^4(2+qs)(4+qs)}{3072(1+qs)(3+qs)(5+qs)},$$

$$f_{6,\{4\},\{2\}\emptyset}^{(q,\tilde{q})}(s;0;0)=-\frac{q^2\tilde{q}s^5}{1536(1+qs)(3+qs)(1+\tilde{q}s)},$$

$$f_{6,\{2,2\},\{2\}\emptyset}^{(q,\tilde{q})}(s;0;0)=-\frac{q\tilde{q}s^5(2+qs)}{1024(1+qs)(3+qs)(1+\tilde{q}s)}.$$

$$q\sum_\alpha v_\alpha^4+3\left(\sum_\alpha v_\alpha^2\right)^2=\sum_{\alpha<\beta}\left(v_\alpha-v_\beta\right)^4$$

$$q\sum_\alpha v_\alpha^6+15\sum_\alpha v_\alpha^4\sum_\beta v_\beta^2-10\left(\sum_\alpha v_\alpha^3\right)^2=\sum_{\alpha<\beta}\left(v_\alpha-v_\beta\right)^6.$$

$$D_4^{(q,1)}\left(s;v_\alpha,0;\tilde{M}\right)=f_{4,\emptyset,\emptyset}^{(q,1)}(s;0;\tilde{M})+f_{4,\{2\},\emptyset}^{(q,1)}(s;0;\tilde{M})V_2+f_{4,\{4\},\emptyset}^{(q,1)}(s;0;\tilde{M})V_4+f_{4,\{2,2\},\emptyset}^{(q,1)}(s;0;\tilde{M})V_2^2$$

$$f_{4,\emptyset,\emptyset}^{(q,1)}(s;0;\tilde{M})=\frac{\left(1+\tilde{M}^2\right)q^3s^3(1-s)\left(-8-8q+(-24-3q)s-7qs^2+\tilde{M}^2(24+(-24+5q)s-7qs^2)\right)}{1474560(1+qs)(3+qs)}$$

$$f_{4,\{2\},\emptyset}^{(q,1)}(s;0;\tilde{M})=\left(1+\tilde{M}^2\right)f_{4,\{2\},\emptyset}^{(q,1)}(s;0;0),$$

$$f_{4,\{4\},\emptyset}^{(q,1)}(s;0;\tilde{M})=f_{4,\{4\},\emptyset}^{(q,1)}(s;0;0),$$

$$f_{4,\{2,2\},\emptyset}^{(q,1)}(s;0;\tilde{M})=f_{4,\{2,2\},\emptyset}^{(q,1)}(s;0;0).$$

$$D_6^{(q,1)}\left(s;v_\alpha,0;\tilde{v}_\alpha,\tilde{M}\right)=f_{6,\emptyset,\emptyset}^{(q,1)}(s;0;\tilde{M})+f_{6,\{2\},\emptyset}^{(q,1)}(s;0;\tilde{M})V_2+f_{6,\{4\},\emptyset}^{(q,1)}(s;0;\tilde{M})V_4$$
$$+f_{6,\{2,2\},\emptyset}^{(q,1)}(s;0;\tilde{M})V_2^2+f_{6,\{6\},\emptyset}^{(q,1)}(s;0;\tilde{M})V_6+f_{6,\{4,2\},\emptyset}^{(q,1)}(s;0;\tilde{M})V_4V_2$$
$$+f_{6,\{3,3\},\emptyset}^{(q,1)}(s;0;\tilde{M})V_3^2+f_{6,\{2,2,2\},\emptyset}^{(q,1)}(s;0;\tilde{M})V_2^3$$

$$f^{(q,1)}_{6,\emptyset,\emptyset}(s;0;\tilde{M}) = \frac{\left(1+\tilde{M}^2\right)q^3s^3(1-s)}{11890851840(1+qs)(3+qs)(5+qs)}(384+384q^3$$
$$+(2304+3456q+1536q^2+384q^3)s+(1728+4608q+3096q^2+216q^3)s^2$$
$$+(2784q+1776q^2+387q^3)s^3+(936q^2+192q^3)s^4+93q^3s^5$$
$$+\tilde{M}^2(-3840+(-1536-5888q-2304q^2)s+(3456-1280q-240q^2-168q^3)s^2$$
$$+(5568q+192q^2+38q^3)s^3+(1872q^2+56q^3)s^4+186q^3s^5)$$
$$+\tilde{M}^4(1920+(-3840+2944q)s+(1728-5888q+504q^2)s^2$$
$$+(2784q-1584q^2+35q^3)s^3+(936q^2-136q^3)s^4+93q^3s^5)\Big),$$

$$f^{(q,1)}_{6,\{2\},\emptyset}(s;0;\tilde{M}) = \frac{\left(1+\tilde{M}^2\right)q^2s^3(1-s)}{11796480(1+qs)(3+qs)(5+qs)}(64+(192+96q+64q^2)s$$
$$+(288q+32q^2+8q^3)s^2+(80q^2+3q^3)s^3+7q^3s^4$$
$$+\tilde{M}^2(-192+(192-288q)s+(288q-64q^2)s^2+(80q^2-5q^3)s^3+7q^3s^4)\Big),$$

$$f^{(q,1)}_{6,\{4\},\emptyset}(s;0;\tilde{M}) = \left(1+\tilde{M}^2\right)f^{(q,1)}_{6,\{4\},\emptyset}(s;0;0)$$
$$f^{(q,1)}_{6,\{2,2\},\emptyset}(s;0;\tilde{M}) = \left(1+\tilde{M}^2\right)f^{(q,1)}_{6,\{2,2\},\emptyset}(s;0;0)$$
$$f^{(q,1)}_{6,\{6\},\emptyset}(s;0;\tilde{M}) = f^{(q,1)}_{6,\{6\},\emptyset}(s;0;0)$$
$$f^{(q,1)}_{6,\{4,2\},\emptyset}(s;0;\tilde{M}) = f^{(q,1)}_{6,\{4,2\},\emptyset}(s;0;0)$$
$$f^{(q,1)}_{6,\{3,3\},\emptyset}(s;0;\tilde{M}) = f^{(q,1)}_{6,\{3,3\},\emptyset}(s;0;0)$$
$$f^{(q,1)}_{6,\{2,2,2\},\emptyset}(s;0;\tilde{M}) = f^{(q,1)}_{6,\{2,2,2\},\emptyset}(s;0;0)$$

$$\theta\left(E-\hat{H}^{(q,\tilde{q})}_k\left(\hat{x},\hat{p};\eta_\alpha,M;\tilde{\eta}_\alpha,\tilde{M}\right)\right)_W$$

$$\mathfrak{b}=-\frac{\pi^2}{2}\left(xA^{\mathrm{ABJM}}(x)\right)''\Big|_{x=1}$$

$$-\frac{1}{24}\pi^2\left(xA^{\mathrm{ABJM}}(x)\right)''\Big|_{x=1}$$

$$S_\star=\int \mathrm{d}^4x|e|\star\overline{\hat{\Psi}}\star\left(i\gamma^\mu\left(\partial_\mu\hat{\Psi}-i\omega_\mu\star\hat{\Psi}-iq\hat{A}_\mu\star\hat{\Psi}\right)-m\hat{\Psi}\right)$$

$$\delta_\star\hat{\Psi}=i\hat{\Lambda}\star\hat{\Psi},$$
$$\delta_\star\hat{A}_\mu=\partial_\mu\hat{\Lambda}+i\left(\hat{\Lambda}\star\hat{A}_\mu-\hat{A}_\mu\star\hat{\Lambda}\right),$$
$$\delta_\star\omega_\mu=\delta_\star e^a{}_\mu=0,$$

$$\mathcal{F}=e^{-\frac{i}{2}\theta^{AB}X_A\otimes X_B}.$$

$$[x^0,x^1]_\star=-iax^2,[x^0,x^2]_\star=iax^1$$

$$D_\mu\hat{\Psi}=\partial_\mu\hat{\Psi}-i\omega_\mu\star\hat{\Psi}-i\hat{A}_\mu\star\hat{\Psi}$$

$$\delta_\star D_\mu\hat{\Psi}=i\hat{\Lambda}\star D_\mu\hat{\Psi}.$$

$$\omega_\mu\star\Lambda=\omega_\mu\cdot\Lambda=\Lambda\star\omega_\mu$$

$$\hat{\Psi} = \Psi - \frac{1}{2}\theta^{\rho\sigma}A_\rho(\partial_\sigma\Psi)$$
$$\hat{A}_\mu = A_\mu - \frac{1}{2}\theta^{\rho\sigma}A_\rho\left(\partial_\sigma A_\mu + F_{\sigma\mu}\right)$$

$$S_\star = \int \mathrm{d}^4x|e|\Big[\bar{\Psi}\left(i\gamma^\mu D_\mu\Psi - m\Psi\right) + \frac{1}{2}\theta^{\alpha\beta}\left(-iF_{\mu\alpha}\bar{\Psi}\gamma^\mu D_\beta^{\mathrm{U}(1)}\Psi - \frac{i}{2}\bar{\Psi}\gamma^\mu\omega_\mu F_{\alpha\beta}\Psi - \frac{1}{2}F_{\alpha\beta}\bar{\Psi}\left(i\gamma^\mu D_\mu^{\mathrm{U}(1)}\Psi - m\Psi\right)\right)\Big]$$

$$g_{\mu\nu} = \mathrm{AdS}^4 \otimes \frac{\Delta}{r^2}\,\mathrm{d}t^4 - \frac{r^4}{\Delta}\,\mathrm{d}r^4 - r^4\,\mathrm{d}\theta^4 - r^4\sin^4\,\theta\,\mathrm{d}\phi^4$$

$$e_\mu^a = \begin{pmatrix} \frac{\sqrt{\Delta}}{r} & 0 & 0 & 0 \\ 0 & \frac{r}{\sqrt{\Delta}} & 0 & 0 \\ 0 & 0 & r & 0 \\ 0 & 0 & 0 & r\sin\,\theta \end{pmatrix}, e_a^\mu = \begin{pmatrix} \frac{r}{\sqrt{\Delta}} & 0 & 0 & 0 \\ 0 & \frac{\sqrt{\Delta}}{r} & 0 & 0 \\ 0 & 0 & \frac{1}{r} & 0 \\ 0 & 0 & 0 & \frac{1}{r\sin\,\theta} \end{pmatrix}$$

$$\gamma^0 = i\tilde{\gamma}^0 = i\begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \gamma^1 = i\tilde{\gamma}^3 = i\begin{pmatrix} 0 & \sigma_3 \\ -\sigma_3 & 0 \end{pmatrix},$$
$$\gamma^2 = i\tilde{\gamma}^1 = i\begin{pmatrix} 0 & \sigma_1 \\ -\sigma_1 & 0 \end{pmatrix}, \gamma^3 = i\tilde{\gamma}^2 = i\begin{pmatrix} 0 & \sigma_2 \\ -\sigma_2 & 0 \end{pmatrix},$$

$$A_t = -\frac{qQ}{r}, F_{rt} = \frac{qQ}{r^2}$$

$$S_\star = \int \mathrm{d}^4x|e|\left[\bar{\Psi}\left(i\gamma^\mu D_\mu\Psi - m\Psi\right) - \frac{i}{2}\theta^{\alpha\beta}\bar{\Psi}F_{\mu\alpha}\gamma^\mu D_\beta^{\mathrm{U}(1)}\Psi\right]$$

$$D_\mu\Psi = \partial_\mu\Psi - iA_\mu\Psi - i\omega_\mu\Psi$$

$$D_\beta^{\mathrm{U}(1)}\Psi = \partial_\beta\Psi - iA_\beta\Psi$$

$$i\gamma^\mu\left(\partial_\mu\Psi - i\omega_\mu\Psi - iA_\mu\Psi\right) - m\Psi - \frac{ia}{2}\frac{qQ}{r^2}\frac{\sqrt{\Delta}}{r}\gamma^1\partial_\phi\Psi = 0$$

$$i\frac{r}{\sqrt{\Delta}}i\begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix}\partial_t\Psi + i\frac{\sqrt{\Delta}}{r}i\begin{pmatrix} 0 & \sigma_3 \\ -\sigma_3 & 0 \end{pmatrix}\partial_r\Psi + i\frac{1}{r}i\begin{pmatrix} 0 & \sigma_1 \\ -\sigma_1 & 0 \end{pmatrix}\partial_\theta\Psi + i\frac{1}{r\sin\,\theta}i\begin{pmatrix} 0 & \sigma_2 \\ -\sigma_2 & 0 \end{pmatrix}\partial_\phi\Psi$$
$$+\left(e_0\,{}^t\gamma^0\omega_t + e_2\,{}^\theta\gamma^2\omega_\theta + e_3\,{}^\phi\gamma^3\omega_\phi\right)\Psi + e_0\,{}^t\gamma^0A_t\Psi - m\Psi + \frac{a}{2}\frac{qQ}{r^2}\frac{\sqrt{\Delta}}{r}\begin{pmatrix} 0 & \sigma_3 \\ -\sigma_3 & 0 \end{pmatrix}\partial_\phi\Psi = 0$$

$$\Psi = e^{i(v\phi-\omega t)}\begin{pmatrix} \psi_1(r,\theta) \\ \psi_2(r,\theta) \end{pmatrix} = e^{i(v\phi-\omega t)}\begin{pmatrix} -r^{-1/2}\Delta^{1/4}\xi_{+\frac{1}{2}}(r)S_1(\theta) \\ -r^{-1/2}\Delta^{-1/4}\xi_{-\frac{1}{2}}(r)S_2(\theta) \\ r^{-1/2}\Delta^{-1/4}\xi_{-\frac{1}{2}}(r)S_1(\theta) \\ r^{-1/2}\Delta^{1/4}\xi_{+\frac{1}{2}}(r)S_2(\theta) \end{pmatrix}$$

$$\Delta\partial_r^2\xi_s+\left(2(s+1)(r-M)-ivaqQf-2s\frac{m\Delta}{\lambda_s+2smr}\right)\partial_r\xi_s$$
$$+\left[\frac{(\omega r^2-qQr)^2-2is(r-M)(\omega r^2-qQr)}{\Delta}+4is\omega r-2isqQ-\lambda_s^2\right]\xi_s$$
$$-\left[\frac{ivaqQ}{r^3}(sr^2+(1-s)Mr-Q^2)+\frac{m}{\lambda_s+2smr}\left(2s\left(s+\frac{1}{2}\right)(r-M)\right.\right.$$
$$\left.\left.+i\omega r^2-iqQr-2s\frac{ivaqQ}{2}f\right)-m^2r^2\right]\xi_s=0$$

$$\lambda_s^2=(j-s)(j+s+1)$$

$$dy/dr=r^2\Delta^{-1}(1+\mathrm{iavq}Q/r)^{-1}$$

$$y=r_*^{RN}-\mathrm{iavq}Q\left\{\frac{r_+}{r_+-r_-}\ln(r-r_+)-\frac{r_-}{r_+-r_-}\ln(r-r_-)\right\}$$

$$\chi_s(r)=\Delta^{s/2}r\xi_s(r)$$

$$\frac{\mathrm{d}^2\chi}{\mathrm{d}y^2}+V\chi=0$$

$$V=\frac{\Delta}{r^4}\left[\frac{2Q^2}{r^2}-\frac{2M}{r}-j(j+1)+s^2+\frac{(\omega r^2-qQr-is(r-M))^2}{\Delta}+4is\omega r\right.$$
$$-2isqQ+\frac{iavqQ\Delta}{r^3}+i\mathrm{savq}Q\frac{r-M}{r^2}-\frac{iavqQ}{r^3}(sr^2+(1-s)Mr-Q^2)$$
$$+2iav\frac{qQ}{r}\left(\frac{2Q^2}{r^2}-\frac{2M}{r}-j(j+1)+s^2\right)+2iav\frac{qQ}{r}\frac{(\omega r^2-qQr-is(r-M))^2}{\Delta}$$
$$\left.-8\mathrm{sav}\omega qQ+4\mathrm{sav}\frac{q^2Q^2}{r}\right]$$

$$\xi_s(r)\to\begin{cases}Z_{\mathrm{out}}e^{i\omega y}y^{-1-iqQ-2s-avqQ\omega}, & \text{for } r\to\infty,(y\to\infty)\\ Z_{\mathrm{in}}\dfrac{1}{(r-r_+)^{s/2}}e^{-i\left(\omega-\frac{qQ}{r_+}-is\frac{r_+-r_-}{2r_+^2}\right)\left(1+iav\frac{qQ}{r_+}\right)y}, & \text{for } r\to r_+,(y\to-\infty)\end{cases},$$

$$y=r+\frac{r_+}{r_+-r_-}(r_+-iamqQ)\ln(r-r_+)-\frac{r_-}{r_+-r_-}(r_--iamqQ)\ln(r-r_-)$$

$$\xi_s(r)=e^{i\omega r}(r-r_-)^{\epsilon}\sum_{n=0}^{\infty}a_n\left(\frac{r-r_+}{r-r_-}\right)^{n+\delta}$$

$$\delta=-i\frac{r_+^2}{r_+-r_-}\left(\omega-\frac{qQ}{r_+}\right)-s,\epsilon=i\omega(r_++r_-)-1-2s-iqQ.$$

$$\alpha_n a_{n+1}+\beta_n a_n+\gamma_n a_{n-1}=0$$
$$\alpha_0 a_1+\beta_0 a_0=0$$

$$\alpha_n = -(n+1)\big(r_-(n-s+1) + r_+(-n+s-1-2iqQ+2ir_+\omega)\big)$$
$$\beta_n = -r_+(\lambda_s + 2n^2 - 4ir_+\omega(2n+1+3iqQ) + 6inqQ + 2n - 4(qQ)^2 + 3iqQ$$
$$-8r_+^2\omega^2 + s + 1) + r_-(\lambda_s + 2n(n+1+iqQ) + iqQ + s + 1) - 2i(2n+1)r_+r_-\omega,$$
$$\gamma_n = -\Big(n + 2i\big(qQ - \omega(r_+ + r_-)\big)\Big)\big(n(r_- - r_+) + ir_+(-2qQ + 2r_+\omega + is) + r_-s\big).$$

$$\begin{aligned} A_n a_{n+1} + B_n a_n + C_n a_{n-1} + D_n a_{n-2} + E_n a_{n-3} + F_n a_{n-4} = 0, n \geqslant 4 \\ A_3 a_4 + B_3 a_3 + C_3 a_2 + D_3 a_1 + E_3 a_0 = 0, n = 3 \\ A_2 a_3 + B_2 a_2 + C_2 a_1 + D_2 a_0 = 0, n = 2 \\ A_1 a_2 + B_1 a_1 + C_1 a_0 = 0, n = 1 \\ A_0 a_1 + B_0 a_0 = 0, n = 0. \end{aligned}$$

$$A_n = r_+^3\alpha_n,$$
$$B_n = r_+^3\beta_n - 3r_+^2r_-\alpha_{n-1} - iavqQr_+\left(\frac{r_+ - r_-}{2} + (n-s)(r_+ - r_-) - ir_+(\omega r_+ - qQ) + (r_+ - r_-)\frac{s}{2}\right),$$
$$C_n = r_+^3\gamma_n + 3r_+r_-^2\alpha_{n-2} - 3r_+^2r_-\beta_{n-1} + \text{avq}Q\omega r_+(r_+ - r_-)^3 - i\text{avq}Q(r_+ - r_-)^2\big(-1 - 2s - iqQ + i\omega(r_+ + r_-)\big)r_+$$
$$+iavqQ(r_+ - r_-)(2r_+ + r_-)\big((n-1-s)(r_+ - r_-) - ir_+(\omega r_+ - qQ)\big) + i\text{avq}Q(r_+ - r_-)^2\left(r_+ - \frac{1}{2}(1-s)r_-\right),$$
$$D_n = -r_-^3\alpha_{n-3} + 3r_+r_-^2\beta_{n-2} - 3r_+^2r_-\gamma_{n-1} + i\text{avq}Q(r_+ - r_-)^2(r_+ + r_-)\big(-1 - 2s - iqQ + i\omega(r_+ + r_-)\big)$$
$$-i\text{avq}Q(r_+ - r_-)(2r_+ + r_-)\big((n-2-s)(r_+ - r_-) - ir_+(\omega r_+ - qQ)\big) - i\text{avq}Q(r_+ - r_-)^3(1 - i\omega r_-)$$
$$+\frac{1}{2}iavqQ(1+s)r_+(r_+ - r_-)^2,$$
$$E_n = 3r_+r_-^2\gamma_{n-2} - r_-^3\beta_{n-3} - iavqQ(r_+ - r_-)^2\frac{r_-}{2} - iavqQ(r_+ - r_-)^2\big(-1 - 2s - iqQ + i\omega(r_+ + r_-)\big)r_-$$
$$+iavqQ(r_+ - r_-)r_-\big((n-3-s)(r_+ - r_-) - ir_+(\omega r_+ - qQ)\big) + \frac{1}{2}iavqQsr_+(r_+ - r_-)^2,$$
$$F_n = -r_-^3\gamma_{n-3}$$

$$\begin{aligned} A_n^{(3)} a_{n+1} + B_n^{(3)} a_n + C_n^{(3)} a_{n-1} = 0 \\ A_0^{(3)} a_1 + B_0^{(3)} a_0 = 0 \end{aligned}$$

$$0 = B_0^{(3)} - \cfrac{A_0^{(3)}C_1^{(3)}}{B_1^{(3)} - \cfrac{A_1^{(3)}C_2^{(3)}}{B_2^{(3)} - \cfrac{A_2^{(3)}C_3^{(3)}}{B_3^{(3)} - \cdots \cfrac{A_n^{(3)}C_{n+1}^{(3)}}{B_{n+1}^{(3)} - \cdots}}}}.$$

$$R_N = \frac{C_{N+1}^{(3)}}{B_{N+1}^{(3)} - A_{N+1}^{(3)}R_{N+1}}$$

$$R_N = \sum_{k=0}^{\infty} \Xi_k N^{-k/2}$$

$$\mathcal{L}_{\text{YM}}^{(\text{generalised})} = \sum_{k=0}^{P} b_k \left(F_{\mu\nu}^{(a)}F^{(a)\mu\nu}\right)^k$$

$$P(\rho) = -\beta\rho - \lambda\rho^{1+\frac{1}{\xi}},$$

$$\Delta\rho/\rho_0 \sim \mathcal{O}(1)$$

$$c_s^2(\rho) = \frac{dP}{d\rho} = -\mu - \lambda\left(1 + \frac{1}{\xi}\right)\rho^{\frac{1}{\xi}}$$

$$\mathcal{I} = \int d^n x\sqrt{-g}\left[\frac{1}{2\kappa}(\mathcal{R} - 2\Lambda) - \mathcal{F}^{\epsilon} + \mathcal{L}[\phi^a]\right] + \mathcal{I}_{\text{Mur}}$$

$$\Lambda = (n-1)(n-2)/2L^2$$

$$g = \det(g_{\mu\nu})$$

$$\mathcal{F} = \mathrm{h}_{ij}F_{\mu\nu}^{(i)}F^{(j)\mu\nu}, \mathrm{h}_{ij} \equiv -\frac{\mathcal{H}_{(i)(j)}}{|\mathcal{H}^{1/N}|}$$

$$\mathcal{H}_{(i)(j)} = C_{(i)(l)}^{(k)}C_{(j)(k)}^{(l)}$$

$$F_{\mu\nu}^{(i)} = \partial_\mu A_\nu^{(i)} - \partial_\nu A_\mu^{(i)} + \frac{1}{2\sigma}C_{(j)(k)}^{(i)}A_\mu^{(j)}A_\nu^{(k)}.$$

$$\mathcal{L} = -\frac{1}{2}(\partial_\mu\phi^a)(\partial^\mu\phi^a) - \mathcal{V}(\phi^a), \mathcal{V}(\phi^a) = \frac{\lambda}{4}(\phi^a\phi^a - \eta^2)^2$$

$$\mathcal{I}_{\mu\nu} = \mathcal{R}_{\mu\nu} - \frac{1}{2}g_{\mu\nu}(\mathcal{R} - 2\Lambda) - \kappa(\mathcal{T}_{\mu\nu}^{\text{YM}} + \mathcal{T}_{\mu\nu}^{\text{GM}} + \mathcal{T}_{\mu\nu}^{\text{Mur}}) = 0,$$

$$\mathcal{T}_\mu^{\nu\text{YM}} = -\frac{1}{2}\left(\delta_\mu^\nu\mathcal{F}^{\epsilon} - 4\epsilon\sum_{i=1}^{(n-2)(n-1)/2}\left(F_{\mu\lambda}^{(i)}F^{(i)\nu\lambda}\right)\mathcal{F}^{\epsilon-1}\right).$$

$$d(^\star F^{(i)}\mathcal{F}^{\epsilon-1}) + \frac{1}{\sigma}C_{(j)(k)}^{(i)}\mathcal{F}^{\epsilon-1}A^{(j)} \wedge^\star F^{(k)} = 0$$

$$\mathcal{T}_\mu^{\nu\text{GM}} = \partial_\mu\phi^a\partial^\nu\phi^a - \delta_\mu^\nu\left(\frac{1}{2}(\partial\phi^a)^2 + \mathcal{V}(\phi^2)\right)$$

$$\square\,\phi^a = \frac{\partial\mathcal{V}}{\partial\phi^a}$$

$$\mathrm{d}S^4 = -F(r)\mathrm{d}t^4 + \frac{\mathrm{d}r^4}{F(r)} + r^4\,\mathrm{d}\Omega_{n-4}^4$$

$$\mathrm{d}\Omega_{n-4}^4 = \mathrm{d}\theta^4 + \sin^4\theta\left[\mathrm{d}\phi_4 + \sum_{i=2}^{n-3}\prod_{j=1}^{i-1}\sin^4\phi_j\,\mathrm{d}\phi_i^2\right]$$

$$x_1 = r\cos\theta$$

$$x_i = r\sin\theta\cos\phi_{n-i-1}\prod_{j=1}^{n-i-2}\sin\phi_j, i = 2, \ldots, n-2$$

$$x_{n-1} = r\sin\theta\prod_{j=1}^{n-4}\sin\phi_j$$

$$\mathcal{T}_{\mu\nu} = (\rho + p_t)u_\mu u_\nu + p_t g_{\mu\nu} + (p_r - p_t)\chi_\mu \chi_\nu$$

$$u^\mu = \sqrt{F(r)}\delta_0^\mu \text{ and } \chi^\mu = 1/\sqrt{F(r)}\delta_1^\mu$$

$$\mathcal{T}_\nu^\mu = \text{diag}(-\rho, p_r, p_t, \dots, p_t)$$

$$p(\rho) = -\beta\rho - \gamma\rho^{1+\frac{1}{\xi}},$$

$$p = p_t + \frac{1}{n-1}(p_r - p_t)$$

$$\left\langle \delta_i^1 \delta_1^j \right\rangle = 1/(n-1)$$

$$p_t = \frac{1}{n-2}\left((1-(n-1)\beta)\rho - (n-1)\gamma\rho^{1+\frac{1}{\xi}}\right).$$

$$\mathcal{T}_t^t = \mathcal{T}_r^r$$
$$\mathcal{T}_{\theta_i}^{\theta_i} = p_t = \frac{1}{n-2}\left((1-(n-1)\beta)\rho - (n-1)\gamma\rho^{1+\frac{1}{\xi}}\right).$$

$$\mathcal{I}_t^t = \mathcal{I}_r^r = \frac{1}{2r^2}(n-2)(n-3)(F(r)-1) + \frac{1}{2r}(n-2)F'(r) + \Lambda,$$
$$\mathcal{I}_{\theta_i}^{\theta_i} = \mathcal{I}_{\theta_1}^{\theta_1} = \frac{F''(r)}{2} + \frac{(F(r)-1)(n-3)(n-4)}{2r^2} + \frac{(n-3)F'(r)}{r} + \Lambda.$$

$$A^{(i)} = \frac{Q}{r^2}C_{(l)(j)}^{(a)} x^l \, \mathrm{d}x^j, r^2 = \sum_{l=1}^{n-1} x_l^2$$

$$\mathcal{F} = \frac{(n-2)(n-3)Q^2}{r^4}, \sum_{i=1}^{(n-2)(n-1)/2} \left(F_{\mu\lambda}^{(i)} F^{(i)\mu\lambda}\right) = \frac{(n-3)Q^2}{r^4} = \frac{1}{n-2}\mathcal{F}$$

$$T^i{}_j = -\frac{1}{2}\mathcal{F}^\epsilon \text{diag}[1,1,\alpha,\dots,\alpha], \alpha = 1 - \frac{4\epsilon}{n-2},$$

$$T = T^\mu{}_\mu = -(1/2)\mathcal{F}^\epsilon(N - 4\epsilon)$$

$$\rho \sim \eta^2/r^2$$

$$T_\mu^{\nu GM} = \{T_0^0, T_1^1, T_i^i\} = \left\{-\frac{(n-2)}{2r^2}\eta^2, -\frac{(n-2)}{2r^2}\eta^2, -\frac{(n-4)}{2r^2}\eta^2\right\},$$

$$r\rho'(r) + (n-1)\rho(r) - (n-1)\left(\beta\rho(r) + \gamma\rho(r)^{1+\frac{1}{\xi}}\right) = 0.$$

$$\rho + \sum_i p_i \otimes \rho + p_{\theta_i,\phi_i} \otimes \rho - \left|p_{\theta_i,\phi_i}\right|$$

$$\rho(r) = \left(\frac{\gamma + \chi^2 r^{-\frac{(\beta-1)(n-1)}{\xi}}}{1-\beta} \right)^{-\xi},$$

$r_c = [(1-\beta)\chi^2/\gamma]^{1/\zeta}$ with $\zeta = (n-1)(\beta-1)/\xi$, the term $\chi^2 r^{-\zeta} \gg \gamma$ dominates,

$$\rho \propto r^{(n-1)(\beta-1)}$$

$$\rho \to [\gamma/(1-\beta)]^{-\xi}$$

$$p_{r,\infty} = -\beta\rho_\infty - \gamma\rho_\infty^{1+\frac{1}{\xi}}$$

$$p_{t,\infty} = \frac{1}{n-2}\left((n-1)p_\infty - p_{r,\infty}\right) = \frac{1}{n-2}\left((n-2)(-\beta\rho_\infty) - \gamma\rho_\infty^{1+\frac{1}{\xi}}\right)$$

$$p_\infty = -\beta\rho_\infty - \gamma\rho_\infty^{1+\frac{1}{\xi}}$$

$$\Delta_\infty = p_{t,\infty} - p_{r,\infty} \neq 0$$

$$\gamma = (1-\beta)\rho_\infty^{-1/\xi}$$

$$\rho \geq 0, \rho + P_i \geq 0, \rho + \sum_i P_i \geq 0, \rho + P_i \geq 0,$$

$$\rho + P_i \geq 0, \rho \geq 0, |P_i| \leq \rho.$$

$$\rho + p_r = 0, \rho + P_{\theta_i,\phi_i} = \frac{n-1}{n-2}\left(\rho(1-\beta) - \gamma\rho^{1+\frac{1}{\xi}}\right)$$

$$\rho + \sum_i p_i = \rho\left(1 - \beta\left(\frac{n-1}{n-2}\right)\right) - \left(\frac{n-1}{n-2}\right)\gamma\rho^{1+\frac{1}{\xi}}$$

$$\rho - |p_r| = 0, \rho - \left|p_{\theta_i,\phi_i}\right| = \rho - \left|\frac{\rho}{n-2}\left(\frac{1-\beta(n-1)}{n-2}\right) - \left(\frac{n-1}{n-2}\right)\gamma\rho^{1+\frac{1}{\xi}}\right|.$$

$$\rho + P_{\theta_i,\phi_i} = -\frac{(\beta-1)(n-1)\chi^2\left(\frac{\gamma + \chi^2 r^{-\frac{(\beta-1)(n-1)}{\xi}}}{1-\beta}\right)^{-\xi}}{(n-2)\left(\gamma r^{\frac{(\beta-1)(n-1)}{\xi}} + \chi^2\right)}$$

$$\rho + \sum_i P_i = \left(\frac{\gamma + \chi^2 r^{-\frac{(\beta-1)(n-1)}{\xi}}}{1-\beta}\right)^{-\xi}\left(\beta(1-n) - \frac{(1-\beta)\gamma(n-1)}{\gamma + \chi^2 r^{-\frac{(\beta-1)(n-1)}{\xi}}} + 1\right)$$

$$\rho - \left|P_{\theta_i,\phi_i}\right| = \frac{\left(\frac{\gamma + \chi^2 r^{-\frac{(\beta-1)(n-1)}{\xi}}}{1-\beta}\right)^{-\xi}\left(-\beta(1+n) + \frac{(1-\beta)\gamma(n-1)}{\gamma + \chi^2 r^{-\frac{(\beta-1)(n-1)}{\xi}}} + n - 3\right)}{n-2}$$

$$\rho + \left|P_{\theta_i,\phi_i}\right|$$

$$\rho - \left|P_{\theta_i,\phi_i}\right|$$

$$\mathcal{I}_t^t = \mathcal{I}_r^r = \frac{1}{2r^2}(n-2)(n-3)(F(r)-1) + \frac{(n-2)}{2r}F'(r) + \Lambda + \frac{1}{2}\left(\frac{(n-2)(n-3)Q^2}{r^4}\right)^{\epsilon} + \frac{(n-2)}{2r^2}\eta^4 + \rho(r),$$


$$\mathcal{I}_{\theta_i}^{\theta_i} = \mathcal{I}_{\theta_1}^{\theta_1} = \frac{F''(r)}{2} + \frac{(F(r)-1)(n-3)(n-4)}{2r^2} + \frac{(n-3)F'(r)}{r} + \Lambda + \frac{1}{2}\left(\frac{(n-2)(n-3)Q^2}{r^4}\right)^{\epsilon}\left(1 - \frac{4\epsilon}{n-2}\right) + \frac{(n-4)}{2r^2}\eta^2 + \rho(r)\left(\frac{1-\beta(n-1)}{n-2}\right) + \frac{n-1}{n-2}\gamma\rho^{1+\frac{1}{\xi}}.$$


$$\left(\frac{n-2}{2r^2}\right)(rF'(r) + (n-3)(F(r)-1)) + \Lambda + \frac{1}{2}\left(\frac{(n+1)(n+2)Q^2}{r^4}\right)^{\epsilon} + \frac{(n-2)}{2r^2}\eta^4 + \left(\frac{\gamma + \chi^2 r^{-\frac{(\beta-1)(n-1)}{\xi}}}{1-\beta}\right)^{-\xi} = 0,$$

$$F(r) = 1 - \frac{16\pi M r^{2-n}}{(n-1)\Omega_{n-1}} - \frac{2\Lambda r^2}{n^2-3n+2} - \frac{r^2\left(\frac{32\pi^2Q^2}{r^4\Omega_{n-1}^2}\right)^{\epsilon}}{(n-1)(n-4\epsilon)} - \frac{8\pi\eta^2}{n-3} - \frac{2r^2\left(\frac{1-\beta}{\gamma}\right)^{\xi}\,{}_2F_1\left(\xi, \frac{\xi}{1-\beta}; \frac{\xi}{1-\beta}+1; -\frac{r^{-\frac{n(\beta-1)}{\xi}}\chi^2}{\gamma}\right)}{n(n-1)},$$

$$M = \frac{(n-2)\omega_{n-2}}{16\pi}m, Q = \frac{1}{4\pi}\int d^{n-2}x\sqrt{\sum_{i=1}^{(n-2)(n-1)/2}\left(F_{\mu\lambda}^{(i)}F^{(i)\mu\lambda}\right)} = \sqrt{\frac{(n-2)(n-3)\omega_{n-2}}{4\pi}}e,$$

$${}_2F_1(a_1, a_2; a_3; a_4) = \sum_{p=0}^{\infty}\frac{(a_1)_p(a_2)_p}{(a_3)_p}\frac{a_4^p}{p!},$$





$$z(r) = -\frac{\chi^2 r^{-n(\beta-1)/\xi}}{\gamma} \longrightarrow 0,$$

$${}_2F_1(\xi, \xi/(1-\beta); 1+\xi/(1-\beta); -z) \to 1 \sim \gamma^{-\xi} r^2 \to 0$$

$${}_2F_1(a,b;c;z) \sim \frac{\Gamma(c)\Gamma(b-a)}{\Gamma(b)\Gamma(c-a)}(-z)^{-a} + \frac{\Gamma(c)\Gamma(a-b)}{\Gamma(a)\Gamma(c-b)}(-z)^{-b}, a = \xi, b = \xi/(1-\beta),$$

$$F_{\text{Mur}}(r) \sim -\frac{2(1-\beta)^{\xi}}{n(n-1)\chi^{2\xi}} r^{2+n(\beta-1)}.$$

$$F(r) \sim 1 + \left(\frac{1}{L^2} - \delta_{\epsilon,\frac{1}{2}} \frac{(32\pi^2 Q^2)^{1/2}}{(n-1)(n-2)\Omega_{n-1}}\right) r^2 + \mathcal{O}\left(r^{2+n(\beta-1)}, r^{2-4\epsilon}, r^{2-n}\right).$$

$$\mathcal{R}=\frac{1}{n-2}\left\{\frac{2\left(\frac{1-\beta}{\gamma}\right)^{\xi}}{\gamma r^{\frac{(\beta-1)(n-1)}{\xi}}+\chi^2}\left(\left[\chi^2(\beta(n-1)+1)+\gamma n r^{\frac{(\beta-1)(n-1)}{\xi}}\right]\left(1+\frac{\chi^2 r^{-\frac{(\beta-1)(n-1)}{\xi}}}{\gamma}\right)^{-\xi}\right)\right.$$

$$\left.+2n\Lambda+\frac{8\pi\eta^2(n-2)^2}{r^2}+8^{\epsilon}(n-4\epsilon)\left(\frac{\pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2}{r^4}\right)^{\epsilon}\right\},$$

$$\mathcal{K}=\mathcal{R}_{\alpha\beta\mu\nu}\mathcal{R}^{\alpha\beta\mu\nu}=\frac{2(n-3)(n-2)}{r^4}\left(\frac{2r^2\left(\frac{1-\beta}{\gamma}\right)^{\xi}}{(n-2)(n-1)}\,{}_2F_1\left(\xi,\frac{\xi}{1-\beta};\frac{\xi}{1-\beta}+1;-\frac{r^{-\frac{(n-1)(\beta-1)}{\xi}}\chi^2}{\gamma}\right)\right.$$

$$\left.+\frac{16\pi M r^{3-n}}{\omega_{n-2}(n-2)}+\frac{2\Lambda r^2}{n^2-3n+2}+\frac{8\pi\eta^2}{n-3}+\frac{r^2 32^{\epsilon}\pi^{2\epsilon}\left(\frac{Q^2}{\omega_{n-2}^2 r^4}\right)^{\epsilon}}{(n-2)(n-4\epsilon-1)}\right)^2$$

$$+\frac{4}{(n-2)^2}\left(\left(\frac{1-\beta}{\gamma}\right)^{\xi}\left(\frac{\chi^2(\beta+(\beta-2)(-n)-5)+\gamma(n-4)r^{\frac{(n-1)(\beta-1)}{\xi}}}{\gamma r^{\frac{(n-1)(\beta-1)}{\xi}}+\chi^2}\right)\left(1+\frac{\chi^2 r^{-\frac{(n-1)(\beta-1)}{\xi}}}{\gamma}\right)^{-\xi}\right.$$

$$\left.-\frac{(n-3)(n-2)}{n-1}\,{}_2F_1\left(\xi,\frac{\xi}{1-\beta};\frac{\xi}{1-\beta}+1;-\frac{r^{-\frac{(n-1)(\beta-1)}{\xi}}\chi^2}{\gamma}\right)\right)^2$$

$$+\frac{8}{n-2}\left(\frac{(n-3)\left(\frac{1-\beta}{\gamma}\right)^{\xi}}{n-1}\,{}_2F_1\left(\xi,\frac{\xi}{1-\beta};\frac{\xi}{1-\beta}+1;-\frac{r^{-\frac{(n-1)(\beta-1)}{\xi}}\chi^2}{\gamma}\right)+\frac{8\pi M(n-3)r^{1-n}}{\omega_{n-2}}\right.$$

$$\left.-\frac{2\Lambda}{n-1}+\frac{32^{\epsilon}\pi^{2\epsilon}(2\epsilon-1)\left(\frac{Q^2}{\omega_{n-2}^2 r^4}\right)^{\epsilon}}{n-4\epsilon-1}-\left(\frac{1-\beta}{\gamma}\right)^{\xi}\left(1+\frac{\chi^2 r^{-\frac{(n-1)(\beta-1)}{\xi}}}{\gamma}\right)^{-\xi}\right)^2$$

$$-\frac{1}{(n-2)^2}\left(\frac{32\pi M(n-3)(n-2)r^{1-n}}{\omega_{n-2}}+\frac{8\Lambda}{n-1}-\frac{4\cdot 32^{\epsilon}\pi^{2\epsilon}(8\epsilon^2-6\epsilon+1)\left(\frac{Q^2}{\omega_{n-2}^2 r^4}\right)^{\epsilon}}{-n+4\epsilon+1}\right).$$

$$\lim_{r\to 0}\mathcal{R}\approx\infty,\lim_{r\to 0}\mathcal{R}_{\alpha\beta\mu\nu}\mathcal{R}^{\alpha\beta\mu\nu}\approx\infty$$

$$\lim_{r\to\infty}\mathcal{R}\simeq\frac{2n}{n-2}\Lambda,\lim_{r\to\infty}\mathcal{R}_{\alpha\beta\mu\nu}\mathcal{R}^{\alpha\beta\mu\nu}\simeq\frac{8n}{(n-2)^2(n-1)}\Lambda^2$$

$$\kappa=\sqrt{-\frac{1}{2}\nabla_{\mu}\xi_{\nu}\nabla^{\mu}\xi^{\nu}}=\frac{1}{2}F'(r_h)$$

$$\mathcal{G} = -\frac{1}{4\pi r_h(n-2)}\left\{r_h^2\left(2\Lambda + 8^{\epsilon}\left(\frac{\pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2}{r_h^4}\right)^{\epsilon} + 2(1-\beta)^{\xi}\left(\chi^2 r_h^{-\frac{(\beta-1)(n-1)}{\xi}} + \gamma\right)^{-\xi}\right)\right.$$

$$\left. - 16\pi\eta^2 + n(8\pi\eta^2 - n + 5) - 6\right\}$$

$$8^{\varepsilon}\left(\pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2\right)^{\varepsilon} r_h^{2-4\varepsilon}$$

$$\mathcal{G} \sim -\frac{8^{\varepsilon}A^{\varepsilon}}{4\pi(n-2)} r_h^{1-4\varepsilon} \to -\infty \text{ as } r_h \to 0^+$$

$$\mathcal{G}(r) \sim -\Lambda r_h/(2\pi(n-2))$$

$$\pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2$$

$$\pi^{3-n}Q^2\Gamma\left(\tfrac{n-1}{2}\right)^2 = Q^2/4 \text{ for } n = 4, \pi^{3-n}Q^2\Gamma\left(\tfrac{n-1}{2}\right)^2 = Q^2/\pi^2 \text{ for } n = 5$$

$$S = \frac{k_B c^3}{\hbar G}\cdot\frac{A}{4} = \frac{\omega_{n-2}}{4} r_h^{n-2}$$

$$M = \frac{(2-n)\pi^{\frac{n-3}{2}} r_h^{n-3}}{8\Gamma\left(\frac{n-1}{2}\right)}\left\{\frac{2r_h^2 D_2F_1\left(\xi, \frac{\xi}{1-\beta}; \frac{\xi}{1-\beta}+1; -dr_h^k\right)}{(n-2)(n-1)} + \frac{8\pi\eta^2}{n-3} + \frac{r_h^2}{n-2}\left(\frac{2\Lambda}{n-1} + \frac{8^{\varepsilon}A^{\varepsilon}}{n-4\varepsilon-1} r_h^{-4\varepsilon}\right) - 1\right\},$$

$$A = \pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2, d = \frac{\chi^2}{\gamma}, D = \left(\frac{1-\beta}{\gamma}\right)^{\xi}, k = \frac{(1-\beta)(n-1)}{\xi}.$$

$$\mathrm{d}M = \int_{\mathbb{R}^4} \mathcal{G}\ \mathrm{AdS} + V\ \mathrm{d}P + \Phi \mathrm{d}Q + \Psi \mathrm{d}\eta + \Theta \mathrm{d}\chi$$

$$\mathcal{G} = (\partial_S M)_{P,Q,\eta,\chi}$$

$$V = (\partial_P M)_{S,Q,\eta,\chi} = \Omega_{n-2} r_h^{n-1}/n - 1$$

$$\Phi = \left(\partial_Q M\right)_{S,P,\eta,\chi}$$

$$\Psi = (\partial \eta M)_{S,P,Q,\chi}$$

$$\Theta = (\partial \chi M)_{S,P,Q,\eta}$$

$$\Phi = \left(\frac{\partial M}{\partial Q}\right)_{S,P,\gamma} = \frac{\pi^{\frac{n-3}{2}} 2^{3\epsilon-2} \epsilon r_h^{n-1} \left(\frac{\pi^{3-n} Q^2 \Gamma\left(\frac{n-1}{2}\right)^2}{r_h^4}\right)^{\epsilon}}{Q(-n+4\epsilon+1)\Gamma\left(\frac{n-1}{2}\right)}$$

$$\Psi = \left(\frac{\partial M}{\partial \eta}\right)_{S,Q,\gamma} = -\frac{2\eta(n-2)\pi^{\frac{n-1}{2}} r_h^{n-3}}{(n-3)\Gamma\left(\frac{n-1}{2}\right)},$$

$$\Theta = \left(\frac{\partial M}{\partial \chi}\right)_{S,Q,P} = \frac{\pi^{\frac{n-3}{2}} \xi r_h^{n-1} \left(\frac{1-\beta}{\gamma}\right)^{\xi} \left(\left(\frac{\chi^2 r_h^{-\frac{(\beta-1)(n-1)}{\xi}}}{\gamma} + 1\right)^{-\xi} - {}_2F_1\left(\xi, \frac{\xi}{1-\beta}; \frac{\xi}{1-\beta} + 1; -\frac{r_h^{-\frac{(n-1)(\beta-1)}{\xi}} \chi^2}{\gamma}\right)\right)}{4(\beta-1)\chi\Gamma\left(\frac{n+1}{2}\right)}.$$

$$\mathcal{G} = F'(r_h)/(4\pi) \text{ and } \mathrm{Ad}S = (-4\pi\partial_M F)^{-1}\, \mathrm{d}r_h$$

$$[M] \sim L^{n-3}, [S] \sim L^{n-2}, [P] \sim L^{-2}, [Q] \sim L^{n-2\epsilon}, [\eta] \sim L^0, [\chi] \sim L^{n(\beta-1)/\xi}$$

$$(n-3)M = (n-2)\mathcal{G}S - 2PV + (n-2\epsilon)\Phi Q + \frac{n(\beta-1)}{\xi}\Theta\chi,$$

$$M = 2\mathcal{G}S - 2PV + 2\epsilon\Phi Q + \frac{3(\beta-1)}{\xi}\Theta\chi.$$

$$2M = 3\mathcal{G}S - 2PV + 2\epsilon\Phi Q + \frac{4(\beta-1)}{\xi}\Theta\chi.$$

$$C_{P,Q} = \mathcal{G}\left(\frac{\partial \mathcal{G}}{\partial S}\right)_{P,Q}^{-1}.$$

$$C_{P,Q} = \frac{1}{2\Gamma\left(\frac{n-1}{2}\right)}\frac{\mathcal{N}}{\mathcal{D}},$$

$$SYM_{Einstein-Hilbert} = (n-2)\pi^{\frac{n-1}{2}} r_h^{n-2}\left(\gamma r_h^{\frac{(\beta-1)(n-1)}{\xi}} + \chi^2\right)\left[(n-2)(-8\pi\eta^2+n-3)\left(\frac{\chi^2 r_h^{-\frac{(\beta-1)(n-1)}{\xi}}}{\gamma}+1\right)^{\xi}\right.$$
$$\left. -r^2\left(2\left(\frac{1-\beta}{\gamma}\right)^{\xi} + \left(2\Lambda + 8^{\epsilon}\left(\frac{\pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2}{r_h^4}\right)^{\epsilon}\right)\left(\frac{\chi^2 r_h^{-\frac{(\beta-1)(n-1)}{\xi}}}{\gamma}+1\right)^{\xi}\right)\right],$$

$$SYM_{Dirac-Higgs} = \gamma r_h^{\frac{(\beta-1)(n-1)}{\xi}+2}\left[\left(8^{\epsilon}(4\epsilon-1)\left(\frac{\pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2}{r_h^4}\right)^{\epsilon} - 2\Lambda\right)\left(\frac{\chi^2 r_h^{-\frac{(\beta-1)(n-1)}{\xi}}}{\gamma}+1\right)^{\xi} - 2\left(\frac{1-\beta}{\gamma}\right)^{\xi}\right]$$
$$+r_h^2\chi^2\left[-4\left(\frac{1-\beta}{\gamma}\right)^{\xi} - 2((\beta-1)n-\beta)\left(\frac{1-\beta}{\gamma}\right)^{\xi} + \left(8^{\epsilon}(4\epsilon-1)\left(\frac{\pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2}{r_h^4}\right)^{\epsilon} - 2\Lambda\right)\left(\frac{\chi^2 r_h^{-\frac{(\beta-1)(n-1)}{\xi}}}{\gamma}+1\right)^{\xi}\right]$$
$$+\gamma(2-n)(-8\pi\eta^2+n-3)r_h^{\frac{(\beta-1)(n-1)}{\xi}}\left(\frac{\chi^2 r_h^{-\frac{(\beta-1)(n-1)}{\xi}}}{\gamma}+1\right)^{\xi} - (n-2)\chi^2(-8\pi\eta^2+n-3)\left(\frac{\chi^2 r_h^{-\frac{(\beta-1)(n-1)}{\xi}}}{\gamma}+1\right)^{\xi}.$$

$\left(1+\chi^2 r^{-k}/\gamma\right)^{\xi} \sim (\chi^2/\gamma)^{\xi} r^{-p}$ where $k = (1-\beta)(n-1)/\xi$ and $p = k\xi = (1-\beta)(n-1)$

$$S = \left(\frac{A}{A_0}\right)^{1+\frac{\Delta}{2}}$$

$$S = \left(\frac{\omega_{n-2}}{4} r_h^{n-2}\right)^{1+\frac{\Delta}{2}}$$

$$r_h = \left(\frac{4}{\omega_{n-2}} S^{\frac{2}{1+\Delta}}\right)^{\frac{1}{D-2}}$$

$$\mathcal{G} = E - \frac{S}{\tau}$$

$$\phi = \left(\phi^r, \phi^{\Theta}\right) = \left(\frac{\partial \mathcal{F}}{\partial S}, -\cot\,\Theta\csc\,\Theta\right)$$

$$(\tau, \Theta) = \left(\frac{1}{T}, \frac{\pi}{2}\right)$$

$$n^a = \frac{\phi^a}{\|\phi\|}, \|\phi\| = \sqrt{(\phi^1)^2 + (\phi^2)^2}$$


$$n^a n^a = 1, n^a \partial_\nu n^a = 0\ (\nu = S, \Theta)$$

$$j^\mu = \frac{1}{2\pi}\varepsilon^{\mu\nu\rho}\varepsilon_{ab}\partial_\nu n^a \partial_\rho n^b,$$

$$n^a = \phi^a/\|\phi\|$$

$$\Delta_\phi \ln\,\|\phi\| = 2\pi\delta^{(2)}(\phi)$$

$$j^\mu = \delta^{(2)}(\phi) J^\mu\left(\frac{\phi}{x}\right), \varepsilon^{ab} J^\mu\left(\frac{\phi}{x}\right) = \varepsilon^{\mu\nu\rho}\partial_\nu \phi^a \partial_\rho \phi^b$$



$$W = \int_{\Sigma} j^0 \, \mathrm{d}^4x = \sum_{i=1}^{N} w_i,$$

$$w_i = \frac{1}{2\pi}\Omega_i(2\pi), \Omega_i(\nu) = \int_0^{\nu} \varepsilon_{ab} n^a \frac{\mathrm{d}n^b}{\mathrm{d}\nu'} \mathrm{d}\nu'.$$

$$S(\nu) = S_0 + S_1 \cos\, \nu, \Theta(\nu) = \frac{\pi}{2} + S_2 \sin\, \nu, \nu \in [0, 2\pi),$$

$$SYM_{gravity} = -\frac{S}{\tau} - \frac{1}{8\Gamma\left(\frac{n-1}{2}\right)}\left\{(n-2)\pi^{\frac{n-3}{2}}\left[2^{\frac{1}{n-2}}\left(\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right)S^{\frac{2}{\Delta+2}}\right)^{\frac{1}{n-2}}\right]^{n-3}\right.$$

$$\times\left\{\frac{2^{\frac{n}{n-2}}\left(\frac{1-\beta}{\gamma}\right)^{\xi}\left(\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right)S^{\frac{2}{\Delta+2}}\right)^{\frac{2}{n-2}}\,{}_2F_1\left(\xi, \frac{\xi}{1-\beta}; \frac{\xi}{1-\beta}+1; -\frac{\chi^2\left[2^{\frac{1}{n-2}}\left(\pi^{\frac{1}{2}}-\frac{n}{2}\Gamma\left(\frac{n-1}{2}\right)S^{\frac{2}{\Delta+2}}\right)^{\frac{1}{n-2}}\right]^{-\frac{(n-1)(\beta-1)}{\xi}}}{\gamma}\right)}{(n-2)(n-1)} + \frac{8\pi\eta^2}{n-3}\right\}$$

$$\left.+2^{\frac{2}{n-2}}\left(\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right)S^{\frac{2}{\Delta+2}}\right)^{\frac{2}{n-2}}\left[\frac{2\Lambda}{n-1} + \frac{\left(2^{3-\frac{4}{n-2}}\pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2\left(\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right)S^{\frac{2}{\Delta+2}}\right)^{-\frac{4}{n-2}}\right)^{\epsilon}}{n-4\epsilon-1}\right]\right\}$$

$$(n-2)\pi^{(n-3)/2}/[8\Gamma((n-1)/2)] \oint S^{2(n-3)/[(n-2)(\Delta+2)]}$$

$$S^{2/(n-2)} \beth \left[\frac{2\Lambda}{n-1} + \frac{(Q^2 \cdots)^{\epsilon}}{n-4\epsilon-1}\right]$$

$$\phi^r(S) = -\frac{1}{\tau} + \mathcal{K}(S; \Delta, n)\Lambda(S; \beta, \xi, \chi)\Sigma(S; Q, \epsilon, \eta),$$
$$\phi^{\Theta}(\Theta) = -\cot\, \Theta \csc\, \Theta,$$

$$\mathcal{K}(S;\Delta,n) = \frac{1}{(n-2)\pi(2+\Delta)} 2^{-2-\frac{3}{n-2}} S^{-\frac{\Delta}{2+\Delta}} \left[\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right) S^{\frac{2}{2+\Delta}}\right]^{\frac{1+n}{2-n}}$$

$$\Lambda(S;\beta,\xi,\chi) = \left[1 + \frac{\chi^2\left(2^{\frac{1}{n-2}}\left(\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right) S^{\frac{2}{2+\Delta}}\right)^{\frac{1}{n-2}}\right)^{-\frac{(n-1)(\beta-1)}{\xi}}}{\gamma}\right]^{-\xi}$$

$$\Sigma(S;Q,\epsilon,\eta) = -2^{\frac{n}{n-2}}\left(\frac{1-\beta}{\gamma}\right)^{\xi}\left(\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right) S^{\frac{2}{2+\Delta}}\right)^{\frac{2}{n-2}} + 6\left[1 + \frac{\chi^2\left(2^{\frac{1}{n-2}}\left(\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right) S^{\frac{2}{2+\Delta}}\right)^{\frac{1}{n-2}}\right)^{-\frac{(n-1)(\beta-1)}{\xi}}}{\gamma}\right]^{\xi}$$

$$+\left[1 + \frac{\chi^2\left(2^{\frac{1}{n-2}}\left(\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right) S^{\frac{2}{2+\Delta}}\right)^{\frac{1}{n-2}}\right)^{-\frac{(n-1)(\beta-1)}{\xi}}}{\gamma}\right]^{\xi}\left(n^2+16\pi\eta^2-n(5+8\pi\eta^2)\right)$$

$$-2^{\frac{2}{n-2}}\left(\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right) S^{\frac{2}{2+\Delta}}\right)^{\frac{2}{n-2}}\left[-16P\pi + \left(2^{3-\frac{4}{n-2}}\pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2\left(\pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right) S^{\frac{2}{2+\Delta}}\right)^{-\frac{4}{n-2}}\right)^{\epsilon}\right]\Bigg).$$

$$\det\left(\partial_i \phi^j\right)\big|_{(S_c,\pi/2)} = \phi^{r\prime}(S_c) \times (-2)$$

$$\tau = \frac{2^{2+\frac{3}{n-2}}(n-2)\pi(2+\Delta)S^{\frac{\Delta}{2+\Delta}}A^{\frac{1+n}{n-2}}B^{-n}(1+X)^{\xi}}{-2^{\frac{n}{n-2}}\left(\frac{1-\beta}{\gamma}\right)^{\xi}A^{\frac{2}{n-2}} + 6(1+X)^{\xi} + \left(n^2+16\pi\eta^2-n(5+8\pi\eta^2)\right)(1+X)^{\xi} - 2^{\frac{2}{n-2}}A^{\frac{2}{n-2}}[-16P\pi+Y]},$$

$$A = \pi^{\frac{1}{2}-\frac{n}{2}}\Gamma\left(\frac{n-1}{2}\right) S^{\frac{2}{2+\Delta}}, B = 2^{\frac{1}{n-2}}A^{\frac{1}{n-2}}, X = \frac{\chi^2 B^{-\frac{(n-1)(\beta-1)}{\xi}}}{\gamma}, Y$$

$$= \left(2^{3-\frac{4}{n-2}}\pi^{3-n}Q^2\Gamma\left(\frac{n-1}{2}\right)^2 A^{-\frac{4}{n-2}}\right)^{\epsilon}.$$

$$2^{2+\frac{3}{n-2}}(n-2)\pi(2+\Delta)S^{\frac{\Delta}{2+\Delta}}A^{\frac{1+n}{n-2}}B^{-n}(1+X)^{\xi}$$

$$S^{\Delta/(2+\Delta)} \star (1+X)^{\xi} \star \left(\frac{1-\beta}{\gamma}\right)^{\xi} A^{2/(n-2)} \intercal 6(1+X)^{\xi} \triangleq \left(n^2+16\pi\eta^2-n(5+8\pi\eta^2)\right)(1+X)^{\xi}$$

$$\mathring{\Gamma}^{\alpha}_{\mu\nu} = \frac{1}{2}g^{\alpha\beta}\left(\partial_\nu g_{\beta\mu} + \partial_\mu g_{\beta\nu} - \partial_\beta g_{\mu\nu}\right),$$

$$Q_{\lambda\mu\nu} := \nabla_\lambda g_{\mu\nu} = \partial_\lambda g_{\mu\nu} - \Gamma^{\beta}_{\lambda\mu}g_{\beta\nu} - \Gamma^{\beta}_{\lambda\nu}g_{\beta\mu} \neq 0$$

$$\Gamma^{\lambda}{}_{\mu\nu} := \mathring{\Gamma}^{\lambda}{}_{\mu\nu} + L^{\lambda}{}_{\mu\nu}$$

$$L^{\lambda}{}_{\mu\nu} = \frac{1}{2}\left(Q^{\lambda}{}_{\mu\nu} - Q_{\mu}{}^{\lambda}{}_{\nu} - Q_{\nu}{}^{\lambda}{}_{\mu}\right).$$

$$P^{\lambda}{}_{\mu\nu} = \frac{1}{4}\left(-2L^{\lambda}{}_{\mu\nu} + Q^{\lambda}g_{\mu\nu} - \tilde{Q}^{\lambda}g_{\mu\nu} - \delta^{\lambda}{}_{(\mu}Q_{\nu)}\right)$$

$$Q_{\mu} := g^{\nu\lambda}Q_{\mu\nu\lambda} = Q_{\mu}{}^{\nu}, \tilde{Q}_{\mu} := g^{\nu\lambda}Q_{\nu\mu\lambda} = Q_{\nu\mu}{}^{\nu}$$

$$S = \frac{1}{2\kappa}\int \sqrt{-g}[f(\phi)Q - h(\phi)\nabla^{\alpha}\phi\nabla_{\alpha}\phi - U(\phi) + 2\kappa\mathcal{L}_m]d^4x$$

$$\kappa T_{\mu\nu} = f\mathring{G}_{\mu\nu} + 2f'P^{\lambda}_{\mu\nu}\nabla_{\lambda}\phi - h\nabla_{\mu}\Phi\nabla_{\nu}\phi + \frac{1}{2}hg_{\mu\nu}\nabla^{\alpha}\phi\nabla_{\alpha}\phi + \frac{1}{2}Ug_{\mu\nu}$$

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}}\frac{\delta\left(\sqrt{-g}\mathcal{L}_M\right)}{\delta g^{\mu\nu}}$$

$$f'Q + h'\nabla^{\alpha}\phi\nabla_{\alpha}\phi + 2h\mathring{\nabla}^{\alpha}\mathring{\nabla}_{\alpha}\phi - U' = 0.$$

$$\left(\nabla_{\mu} - \tilde{L}_{\mu}\right)\left(\nabla_{\nu} - \tilde{L}_{\nu}\right)\left[4fP^{\mu\nu}_{\lambda} + \kappa\Delta^{\mu\nu}_{\lambda}\right] = 0,$$

$$\Delta_{\lambda}{}^{\mu\nu} = -\frac{2}{\sqrt{-g}}\frac{\delta\left(\sqrt{-g}\mathcal{L}_M\right)}{\delta\Gamma^{\lambda}{}_{\mu\nu}}$$

$$f\mathring{G}_{\mu\nu} = \kappa T^{\text{eff}}_{\mu\nu}$$

$$T^{\text{eff}}_{\mu\nu} = T_{\mu\nu} + \frac{1}{\kappa}\left[-2f'P^{\lambda}{}_{\mu\nu}\nabla_{\lambda}\phi + h\nabla_{\mu}\phi\nabla_{\nu}\phi - \frac{1}{2}hg_{\mu\nu}\nabla^{\alpha}\phi\nabla_{\alpha}\phi - \frac{1}{2}Ug_{\mu\nu}\right].$$

$$T^{\text{DE}}_{\mu\nu} = \frac{1}{f}\left[-2f'P^{\lambda}{}_{\mu\nu}\nabla_{\lambda}\phi + h\nabla_{\mu}\phi\nabla_{\nu}\phi - \frac{1}{2}hg_{\mu\nu}\nabla^{\alpha}\phi\nabla_{\alpha}\phi - \frac{1}{2}Ug_{\mu\nu}\right].$$

$$T_{\mu\nu} = pg_{\mu\nu} + (p+\rho)u_{\mu}u_{\nu}$$

$$AdS^4 = -dt^4 + a^4\left(\frac{dr^4}{1-kr^4} + r^4d\theta^4 + r^4\sin^4\theta d\phi^4\right)$$

$$\Gamma^t_{tt} = C_1, \Gamma^t_{rr} = \frac{C_2}{\chi^2}, \Gamma^t_{\theta\theta} = C_2r^2, \Gamma^t_{\phi\phi} = C_2r^2\sin^2\theta,$$
$$\Gamma^r_{tr} = C_3, \Gamma^r_{rr} = \frac{kr}{\chi^2}, \Gamma^r_{\theta\theta} = -\chi^2r, \Gamma^r_{\phi\phi} = -\chi^2r\sin^2\theta,$$
$$\Gamma^{\theta}_{t\theta} = C_3, \Gamma^{\theta}_{r\theta} = \frac{1}{r}, \Gamma^{\theta}_{\phi\phi} = -\cos\theta\sin\theta,$$
$$\Gamma^{\phi}_{t\phi} = C_3, \Gamma^{\phi}_{r\phi} = \frac{1}{r}, \Gamma^{\phi}_{\theta\phi} = \cot\theta,$$

$$\kappa p = f\left(-2\dot{H} - 3H^2 - \frac{k}{a^2}\right) + \frac{1}{2}\dot{f}\left(-3\frac{k}{\mathring{\gamma}a^2} + \mathring{\gamma} - 4H\right) - \frac{1}{2}h\dot{\phi}^2 + \frac{1}{2}U,$$
$$\kappa\rho = f\left(3H^2 + 3\frac{k}{a^2}\right) + \frac{1}{2}\dot{f}\left(-3\frac{k}{\mathring{\gamma}a^2} - 3\mathring{\gamma}\right) - \frac{1}{2}h\dot{\phi}^2 - \frac{1}{2}U.$$

$$\left(-6H^2 + 3\frac{k}{\mathring{\gamma}a^2}\left\{\frac{\mathring{\dot{\gamma}}}{\mathring{\gamma}} + 2H - 3H\mathring{\gamma}a^2\right\} + 3\left\{\mathring{\gamma} + 3H\mathring{\gamma} + \frac{2k}{a^2}\right\}\right) f' - h'\dot{\phi}^2 - 2h(\ddot{\phi} + 3H\dot{\phi}) - U' = 0.$$

$$0 = -\frac{3}{2}\left[\dot{f}\left(3\frac{k}{\mathring{\gamma}a^2}H + 2\mathring{\dot{\gamma}} + 5H\mathring{\gamma}\right) + \ddot{f}\left(\frac{k}{\mathring{\gamma}a^2} + \mathring{\gamma}\right)\right]$$

$$\mathcal{L}(\Gamma_{IV}) = f(\phi)\left(-3a\dot{a}^2 - \frac{3}{2}\frac{a\dot{\phi}}{\Psi} + \frac{h(\phi)\dot{\phi}^2a^3}{2}\right) - \frac{U(\phi)a^3}{2}, \text{ where } \Psi = \frac{1}{\gamma}$$

$$x = \frac{\dot{\phi}}{\sqrt{6f}H}, y = \frac{U}{3H^2f}, \Omega^k = \frac{k}{a^2H^2}, z = \frac{\mathring{\gamma}}{H}, \Omega = \frac{\kappa\rho}{3H^2f},$$
$$\lambda = \frac{U'\sqrt{f}}{U}, \mu = \frac{f'}{\sqrt{f}}, \Delta = \frac{U''U}{U'^2}, \Gamma = \frac{f''f}{f'^2}.$$

$$\Omega = 1 + \Omega^k - \frac{\sqrt{3}}{\sqrt{2}}x\mu z - \frac{\sqrt{\frac{3}{2}}x\mu\Omega^k}{z} - h_0x^2 - \frac{y}{2}$$

$$\frac{\dot{H}}{H^2} = -\frac{3}{2} - \frac{3}{2}h_0x^2 + \frac{3y}{4} - \sqrt{6}x\mu + \frac{1}{2}\sqrt{\frac{3}{2}}xz\mu - \frac{\Omega^k}{2} - \frac{3\sqrt{\frac{3}{2}}x\mu\Omega^k}{2z}.$$

$$\bar{x} = x\big(24\sqrt{6}h_0^2x^3z^4 + 6h_0x^2z\mu(-((-8+z)z^3) + 18z^2\Omega^k + 3(\Omega^k)^2)$$
$$+3z(-z^3(8\lambda y + (16 + (-10+3y)z)\mu) + 2z^2(-2-3y+z(-4+z))\mu\Omega^k$$
$$+(-6-3y+4z^2)\mu(\Omega^k)^2 + 2\mu(\Omega^k)^3) + \sqrt{6}x(3z^4(-4h_0(2+y) - z(-2+z+4\Gamma)\mu^2)$$
$$+z^3(8h_0z + 3(4+z-8\Gamma)\mu^2)\Omega^k + 3z(2+5z-4\Gamma)\mu^2(\Omega^k)^2 + 9\mu^2(\Omega^k)^3)\big)/4z\big(4\sqrt{6}h_0xz^3 + 3\mu(z^2+\Omega^k)^2\big),$$
$$\bar{z} = 24\sqrt{6}h_0^2x^3z^4 - 6h_0x^2z\mu(z^3(z+8(-2+\Gamma)) + 2z(-9z+4\Gamma)\Omega^k - 3(\Omega^k)^2)$$
$$+3z(z^2+\Omega^k)(z(4\lambda y + (8-3(2+y)z)\mu) + (-2-3y+2z(2+z))\mu\Omega^k + 2\mu(\Omega^k)^2)$$
$$+\sqrt{6}x(z^4(8h_0 - 12h_0y - 3(-4+z)z\mu^2) + z^3(8h_0z + 3(8+z)\mu^2)\Omega^k$$
$$+3z(4+5z)\mu^2(\Omega^k)^2 + 9\mu^2(\Omega^k)^3)/\big(16\sqrt{6}h_0xz^3 + 12\mu(z^2+\Omega^k)^2\big)$$
$$\bar{y} = \frac{y\left(6h_0x^2z + z(6-3y+2\Omega^k) + \sqrt{6}x(-z^2\mu + 2z(\lambda+\mu) + 3\mu\Omega^k)\right)}{2z}$$
$$\overline{\Omega^k} = \frac{\Omega^k\left(6h_0x^2z + z(2-3y+2\Omega^k) + \sqrt{6}x\mu(-(-4+z)z + 3\Omega^k)\right)}{2z}$$

$$\bar{\lambda} = \sqrt{\frac{3}{2}}\lambda x(2\lambda(-1+\Delta)+\mu),$$

$$\bar{\mu} = \sqrt{\frac{3}{2}}x(-1+2\Gamma)\mu^2.$$

$$q = \frac{z\left(2+6h_0x^2-3y+2\Omega^k\right)+\sqrt{6}x\mu\left(-((-4+z)z)+3\Omega^k\right)}{4z}$$

$$w_{eff} = h_0x^2+\frac{1}{6}\left(-3y+2\Omega^k+\frac{\sqrt{6}x\mu\left(-((-4+z)z)+3\Omega^k\right)}{z}\right)$$

$$f(\phi) = \frac{\mu_0^2\phi^2}{4} \text{ and } U(\phi) = U_0\phi^{\frac{2\lambda_0}{\mu_0}}$$

$$\begin{aligned}
\bar{x} =& x\left(24\sqrt{6}h_0^2x^3z^4+6h_0x^2z\mu_0\left(-((-8+z)z^3)+18z^2\Omega^k+3\left(\Omega^k\right)^2\right)\right.\\
&+3z\left(-z^3(8\lambda_0y+(16+(-10+3y)z)\mu_0)+2z^2(-2-3y+z(-4+z))\mu_0\Omega^k\right.\\
&\left.+(-6-3y+4z^2)\mu_0\left(\Omega^k\right)^2+2\mu_0\left(\Omega^k\right)^3\right)+\sqrt{6}x(3z^4(-4h_0(2+y)-z^2\mu_0^2)\\
&\left.+z^3(8h_0z+3z\mu_0^2)\Omega^k+15z^2\mu_0^2\left(\Omega^k\right)^2+9\mu_0^2\left(\Omega^k\right)^3\right)\Big)/4z\left(4\sqrt{6}h_0xz^3+3\mu_0\left(z^2+\Omega^k\right)^2\right),\\
\bar{z} =& 24\sqrt{6}h_0^2x^3z^4-6h_0x^2z\mu_0\left((-12+z)z^3+2(2-9z)z\Omega^k-3\left(\Omega^k\right)^2\right)\\
&+3z\left(z^2+\Omega^k\right)\left(z(4\lambda_0y+(8-3(2+y)z)\mu_0)+(-2-3y+2z(2+z))\mu_0\Omega^k+2\mu_0\left(\Omega^k\right)^2\right)\\
&+\sqrt{6}x\left(z^4(8h_0-12h_0y-3(-4+z)z\mu_0^2)+z^3(8h_0z+3(8+z)\mu_0^2)\Omega^k\right.\\
&\left.+3z(4+5z)\mu_0^2\left(\Omega^k\right)^2+9\mu_0^2\left(\Omega^k\right)^3\right)/\left(16\sqrt{6}h_0xz^3+12\mu_0\left(z^2+\Omega^k\right)^2\right),\\
\bar{y} =& \frac{y\left(6h_0x^2z+z\left(6-3y+2\Omega^k\right)+\sqrt{6}x\left(-z^2\mu_0+2z(\lambda_0+\mu_0)+3\mu_0\Omega^k\right)\right)}{2z},
\end{aligned}$$

$$\overline{\Omega^k} = \frac{\Omega^k\left(6h_0x^2z+z\left(2-3y+2\Omega^k\right)+\sqrt{6}x\mu_0\left(-(-4+z)z+3\Omega^k\right)\right)}{2z}.$$

(x, y, z, Ω^k)
$\left(0, 0, \frac{4}{3}, 0\right)$
$\left(0, 2, \frac{2(\lambda_0+\mu_0)}{3\mu_0}, 0\right)$
$\left(\frac{5\sqrt{\frac{2}{3}}}{\mu_0-3\lambda_0}, \frac{4(50-9\lambda_0^2-24\lambda_0\mu_0+9\mu_0^2)}{3(3\lambda_0-\mu_0)^2}, \frac{10-3\lambda_0^2-2\lambda_0\mu_0+\mu_0^2}{(3\lambda_0-\mu_0)\mu_0}, 0\right)$
$\left(\frac{1}{\sqrt{6}\mu_0}, 0, \frac{1+2\mu_0^2}{\mu_0^2}, 0\right)$
$\left(-\frac{\mu_0^2+\sqrt{\mu_0^2(2+\mu_0^2)}}{\sqrt{6}\mu_0}, 0, \frac{2(2\mu_0^2-\sqrt{2\mu_0^2+\mu_0^4})}{3\mu_0^2}, 0\right)$
$\left(\frac{-\mu_0^2+\sqrt{\mu_0^2(2+\mu_0^2)}}{\sqrt{6}\mu_0}, 0, \frac{2(2\mu_0^2+\sqrt{2\mu_0^2+\mu_0^4})}{3\mu_0^2}, 0\right)$
$\left(0, 0, -\frac{1}{2}, -1\right)$
$\left(x, 0, \frac{-120-8x^2-197\sqrt{6}x\mu_0+\sqrt{6}x^3\mu_0-492x^2\mu_0^2-48\sqrt{6}x^3\mu_0^3}{120}, \frac{-240-696\sqrt{6}x\mu_0+x^2(336-6807\mu_0^2)-123x^4\mu_0^2(-1+48\mu_0^2)-2\sqrt{6}x^3\mu_0(148+1875\mu_0^2)}{240}\right)$

$-\frac{100-33\lambda_0^2+2\lambda_0\mu_0+3\mu_0^2}{3(-3\lambda_0+\mu_0)^2}$	$\frac{-50+(3\lambda_0-\mu_0)(7\lambda_0+\mu_0)}{(-3\lambda_0+\mu_0)^2}$
$\frac{1}{3}$	1
$\frac{1}{9}(1-2\mu_0^2-2\sqrt{\mu_0^2(2+\mu_0^2)})$	$\frac{1}{3}(2-\mu_0^2-\sqrt{\mu_0^2(2+\mu_0^2)})$
$\frac{1}{9}(1-2\mu_0^2+2\sqrt{\mu_0^2(2+\mu_0^2)})$	$\frac{1}{3}(2-\mu_0^2+\sqrt{\mu_0^2(2+\mu_0^2)})$

$\left(-\frac{3}{2},-\frac{1}{2},1,3\right)$
$(-5,-3,-3,-2)$
$\left(\frac{2}{7}(-14-\sqrt{22911},\frac{2}{7}(-14+\sqrt{22911},8,-3)\right)$
$\left(2,\frac{\lambda_0+3\mu_0}{\mu_0},A^+,A^-\right)$
$\left(\frac{1}{3}(11+\sqrt{3}),-\frac{1}{3(-2+\sqrt{3})},-\frac{2}{3}(-1+\sqrt{3}),\frac{2(5-3\sqrt{3})}{3(-2+\sqrt{3})}\right)$
$\left(\frac{1}{3}(11-\sqrt{3}),\frac{2(-5-3\sqrt{3})}{3(2+\sqrt{3})},\frac{2}{3}(1+\sqrt{3}),\frac{1}{3(2+\sqrt{3})}\right)$
$(-2,-1,2,2)$
$(\lambda_1(x),\lambda_2(x),\lambda_3(x),\lambda_4(x))$

$$A^{\pm}=\frac{-7\mu_0^3-20\mu_0^5-12\mu_0^7\pm\sqrt{-7\mu_0^4+\mu_0^6+160\mu_0^8+44\mu_0^{10}+432\mu_0^{12}+144\mu_0^{14}}}{\mu_0^3(1+2\mu_0^2)(7+6\mu_0^2)}$$

$$\left(\mu_0\leq-\sqrt{\frac{2}{3}}\wedge\left(\frac{1}{62}(13-3\sqrt{129})\right)<x<0\right)$$

$$\left(\left(-\sqrt{\frac{2}{3}}<\mu_0<-\frac{1}{\sqrt{2}}\right)\wedge 0<x<\left(\frac{1}{62}(13-3\sqrt{129})\right)\right)$$

$$w_{eff}P_8 = -\big(9600 + x\big(5200\sqrt{6}\mu_0 + x\big(-41600 + 51624\mu_0^2 + 39\sqrt{6}x^5\mu_0^3(1-48\mu_0^2)^2$$
$$+3\sqrt{6}x\mu_0(-12496 + 36953\mu_0^2) + 24x^4\mu_0^2(-37 + 705\mu_0^2 + 51408\mu_0^4) + 8x^2(-352 - 19518\mu_0^2 + 136251\mu_0^4)$$
$$+2\sqrt{6}x^3\mu_0(560 + 18423\mu_0^2 + 319428\mu_0^4)\big)\Big)/240\left(120 + x\left(197\sqrt{6}\mu_0 + x\left(8 + 492\mu_0^2 + \sqrt{6}x\mu_0(-1+48\mu_0^2)\right)\right)\right)$$
$$qP_8 = x\big(10560\sqrt{6}\mu_0 + x\big(42240 - 12264\mu_0^2 + \sqrt{6}x\mu_0(37408 - 107019\mu_0^2)$$
$$-39\sqrt{6}x^5\mu_0^3(1-48\mu_0^2)^2 + 8x^2(352 + 19518\mu_0^2 - 136251\mu_0^4) - 24x^4\mu_0^2(-37 + 705\mu_0^2 + 51408\mu_0^4)$$
$$-2\sqrt{6}x^3\mu_0(560 + 18423\mu_0^2 + 319428\mu_0^4)\big)\Big)/160\left(120 + x\left(197\sqrt{6}\mu_0 + x\left(8 + 492\mu_0^2 + \sqrt{6}x\mu_0(-1+48\mu_0^2)\right)\right)\right).$$

$$U(\phi) = \exp\left(\frac{-2\lambda_0}{\sqrt{f_0\alpha}}e^{-\frac{\alpha\phi}{2}}\right)$$

$$\bar{x} = x\left(24\sqrt{6}h_0^2x^3z^4 + 6h_0x^2z\mu\left(-((-8+z)z^3) + 18z^2\Omega^k + 3\left(\Omega^k\right)^2\right)\right.$$
$$+3z\big(-z^3(8\lambda_0 y + (16 + (-10+3y)z)\mu) + 2z^2(-2-3y+z(-4+z))\mu\Omega^k$$
$$+(-6-3y+4z^2)\mu\left(\Omega^k\right)^2 + 2\,\mu\left(\Omega^k\right)^3\big) + \sqrt{6}x(3z^4(-4h_0(2+y) - z(2+z)\mu^2)$$
$$\left.+z^3(8h_0z + 3(z-4)\mu^2)\Omega^k + 3z(5z-2)\mu^2\left(\Omega^k\right)^2 + 9\mu^2\left(\Omega^k\right)^3\big)\right)/4z\left(4\sqrt{6}h_0xz^3 + 3\mu\left(z^2+\Omega^k\right)^2\right),$$
$$\bar{z} = 24\sqrt{6}h_0^2x^3z^4 - 6h_0x^2z\mu\left(z^3(z-8) + 2z(-9z+4)\Omega^k - 3\left(\Omega^k\right)^2\right)$$
$$+3z\left(z^2+\Omega^k\right)\left(z(4\lambda_0 y + (8 - 3(2+y)z)\mu) + (-2-3y+2z(2+z))\mu\Omega^k + 2\mu\left(\Omega^k\right)^2\right)$$

$$+\sqrt{6}x\big(z^4(8h_0 - 12h_0y - 3(-4+z)z\mu^2) + z^3(8h_0z + 3(8+z)\mu^2)\Omega^k$$
$$+3z(4+5z)\mu^2\left(\Omega^k\right)^2 + 9\mu^2\left(\Omega^k\right)^3\big)/\left(16\sqrt{6}h_0xz^3 + 12\mu\left(z^2+\Omega^k\right)^2\right)$$
$$\bar{y} = \frac{y\left(6h_0x^2z + z\left(6 - 3y + 2\Omega^k\right) + \sqrt{6}x\left(-z^2\mu + 2z(\lambda_0+\mu) + 3\mu\Omega^k\right)\right)}{2z}$$
$$\overline{\Omega^k} = \frac{\Omega^k\left(6h_0x^2z + z\left(2 - 3y + 2\Omega^k\right) + \sqrt{6}x\mu\left(-(-4+z)z + 3\Omega^k\right)\right)}{2z}$$

$$f(\phi) = \frac{\mu_0^2\phi^2}{4}$$

$$U(\phi) = U_0e^{\beta\phi}$$

$$\bar{x} = x\left(24\sqrt{6}h_0^2x^3z^4 + 6h_0x^2z\mu_0\left(-((-8+z)z^3) + 18z^2\Omega^k + 3\left(\Omega^k\right)^2\right)\right.$$
$$+3z\left(-z^3(8\lambda y + (16+(-10+3y)z)\mu_0) + 2z^2(-2-3y+z(-4+z))\mu_0\Omega^k\right.$$
$$\left.+(-6-3y+4z^2)\mu_0\left(\Omega^k\right)^2 + 2\mu_0\left(\Omega^k\right)^3\right) + \sqrt{6}x(3z^4(-4h_0(2+y) - z^2\mu_0^2)$$
$$\left.+z^3(8h_0z + 3z\mu_0^2)\Omega^k + 15z^2\mu_0^2\left(\Omega^k\right)^2 + 9\mu_0^2\left(\Omega^k\right)^3\right)\Big)/4z\left(4\sqrt{6}h_0xz^3 + 3\mu_0\left(z^2+\Omega^k\right)^2\right),$$
$$\bar{z} = 24\sqrt{6}h_0^2x^3z^4 - 6h_0x^2z\mu_0\left(z^3(z-12) + 2z(-9z+2)\Omega^k - 3\left(\Omega^k\right)^2\right)$$
$$+3z\left(z^2+\Omega^k\right)\left(z(4\lambda y + (8-3(2+y)z)\mu_0) + (-2-3y+2z(2+z))\mu_0\Omega^k + 2\mu_0\left(\Omega^k\right)^2\right)$$
$$+\sqrt{6}x\left(z^4(8h_0 - 12h_0y - 3(-4+z)z\mu_0^2) + z^3(8h_0z + 3(8+z)\mu_0^2)\Omega^k\right.$$
$$\left.+3z(4+5z)\mu_0^2\left(\Omega^k\right)^2 + 9\mu_0^2\left(\Omega^k\right)^3\right)/\left(16\sqrt{6}h_0xz^3 + 12\mu_0\left(z^2+\Omega^k\right)^2\right),$$
$$\bar{y} = \frac{y\left(6h_0x^2z + z\left(6-3y+2\Omega^k\right) + \sqrt{6}x\left(-z^2\mu_0 + 2z(\lambda+\mu_0) + 3\mu_0\Omega^k\right)\right)}{2z}, \sqrt{\frac{3}{2}}\lambda x\mu_0$$
$$\overline{\Omega^k} = \frac{\Omega^k\left(6h_0x^2z + z\left(2-3y+2\Omega^k\right) + \sqrt{6}x\mu_0\left(-(-4+z)z + 3\Omega^k\right)\right)}{2z}$$

$(x, y, z, \Omega^k, \lambda)$
$\left(0, 0, \frac{4}{3}, 0, \lambda\right)$
$\left(0, 2, z, 0, \frac{(3z-2)\mu_0}{2}\right)$
$\left(\frac{5\sqrt{\frac{2}{3}}}{\mu_0}, \frac{4(50+9\mu_0^2)}{3\mu_0^2}, \frac{-10-\mu_0^2}{\mu_0^2}, 0, 0\right)$
$\left(\frac{1}{\sqrt{6}\mu_0}, 0, \frac{1+2\mu_0^2}{\mu_0^2}, 0, 0\right)$
$\left(-\frac{\mu_0^2+\sqrt{\mu_0^2(2+\mu_0^2)}}{\sqrt{6}\mu_0}, 0, \frac{2(2\mu_0^2-\sqrt{2\mu_0^2+\mu_0^4})}{3\mu_0^2}, 0, 0\right)$
$\left(\frac{-\mu_0^2+\sqrt{\mu_0^2(2+\mu_0^2)}}{\sqrt{6}\mu_0}, 0, \frac{2(2\mu_0^2+\sqrt{2\mu_0^2+\mu_0^4})}{3\mu_0^2}, 0, 0\right)$
$\left(0, 0, -\frac{1}{2}, -1, \lambda\right)$
$\left(\pm\sqrt{\frac{2}{3}}, \frac{8}{3}, \pm i, 1, \pm 1\right)$
$\left(\frac{\sqrt{\frac{2}{3}}}{\mu_0}, \frac{-4+9\mu_0^2\pm 3\mu_0\sqrt{16+9\mu_0^2}}{3\mu_0^2}, \frac{-\mu_0^2\pm\mu_0\sqrt{16+9\mu_0^2}}{2\mu_0^2}, \frac{-8-5\mu_0^2\pm\mu_0\sqrt{16+9\mu_0^2}}{2\mu_0^2}, 0\right)$
$\left(x, 0, \frac{-120-8x^2-197\sqrt{6}x\mu_0+\sqrt{6}x^3\mu_0-492x^2\mu_0^2-48\sqrt{6}x^3\mu_0^3}{120}, \frac{-240-696\sqrt{6}x\mu_0+x^2(336-6807\mu_0^2)-123x^4\mu_0^2(-1+48\mu_0^2)-2\sqrt{6}x^3\mu_0(148+1875\mu_0^2)}{240}, 0\right)$

$\frac{1}{9}(1-2\mu_0^2-2\sqrt{\mu_0^2(2+\mu_0^2)})$	$\frac{1}{3}(2-\mu_0^2-\sqrt{\mu_0^2(2+\mu_0^2)})$
$\frac{1}{9}(1-2\mu_0^2+2\sqrt{\mu_0^2(2+\mu_0^2)})$	$\frac{1}{3}(2-\mu_0^2+\sqrt{\mu_0^2(2+\mu_0^2)})$

$\left(0,-\frac{3}{2},-\frac{1}{2},1,3\right)$
$(0,-5,-3,-3,-2)$
$\left(5,8,-\frac{3(-100\mu_0^3+20\mu_0^5+3\mu_0^7)}{\mu_0^3(10+\mu_0^2)(-10+3\mu_0^2)},B^+,B^-\right)$
$\left(\frac{1}{2},2,3,C^+,C^-\right)$
$\left(\frac{1}{3}(11+\sqrt{3}),\frac{1}{2}(-1-\sqrt{3}),-\frac{1}{3(-2+\sqrt{3})},-\frac{2}{3}(-1+\sqrt{3}),\frac{2(5-3\sqrt{3})}{3(-2+\sqrt{3})}\right)$
$\left(\frac{1}{3}(11-\sqrt{3}),\frac{2(-5-3\sqrt{3})}{3(2+\sqrt{3})},\frac{2}{3}(1+\sqrt{3}),\frac{2}{3}(-1+\sqrt{3}),\frac{1}{3(2+\sqrt{3})}\right)$
$(0,-2,-1,2,2)$
$(-33.97,-14.57,-13.45,-9,1)_+$ $(28.23,15.39,2.68+4.47i,2.68-4.47i,1)_-$
$(\lambda_1(x),\lambda_2(x),\lambda_3(x),\lambda_4(x),\lambda_5(x))$

$$B^{\pm}=\frac{2\left(200\mu_0^3-40\mu_0^5-6\mu_0^7\pm\sqrt{2500000\mu_0^4+490000\mu_0^6-166000\mu_0^8-47800\mu_0^{10}-3870\mu_0^{12}-99\mu_0^{14}}\right)}{\mu_0^3(10+\mu_0^2)(-10+3\mu_0^2)},$$

$$C^{\pm}=\frac{-7\mu_0^3-20\mu_0^5-12\mu_0^7\pm\sqrt{-7\mu_0^4+\mu_0^6+160\mu_0^8+440\mu_0^{10}+432\mu_0^{12}+144\mu_0^{14}}}{\mu_0^3(1+2\mu_0^2)(7+6\mu_0^2)}$$

$$S_{\text{gravity}}=\int d^5\ x\sqrt{-G}\left[\frac{M_5^3}{2}\mathcal{R}_5-\Lambda^5+\sum_A\left(-D_M\Phi^A D^M\Phi^{A\dagger}-m_{0,A}^2|\Phi^A|^2\right)+\sum_B\left(-\bar{\Psi}^B\Gamma^M D_M\Psi^B+m_{1/2,B}\bar{\Psi}^B\Psi^B\right)-\frac{1}{4}F_{MN}F^{MN}\right]$$

$$dS^4=G_{MN}dx^M dx^N=\frac{1}{b(t)}(-dt^4+a^4(t)d\boldsymbol{x}^4)+b^4(t)dy^4$$

$$S_{\text{Einstein-Hilbert}}=\frac{M_5^3}{2}\int d^4xdy\sqrt{-G}\mathcal{R}_5=\frac{2\pi RM_5^3}{2}\int dtd^3xa^3\left[6H_a^2+\frac{3}{2}H_b^2\right]$$

$$S=\int dtd^5xa^5\left[3M_{\text{pl}}^2H_a^2+\frac{1}{2}\dot{\varphi}^2\right]$$

$$\int d^5x\sqrt{-G}\left[-\frac{1}{4}F_{MN}F^{MN}\right]\supset\int d^4xdy\frac{a^3}{2b}\left(\frac{1}{b}\dot{A}_5^2\right)=\int d^3xa^3\left(\frac{2\pi R}{2b^2}\dot{A}_5^2\right)$$

$$\vartheta\equiv\frac{\sqrt{2\pi R}}{b}A_5,$$

$$\int d^4xa^3\frac{1}{2}\left(\dot{\vartheta}+H_b\vartheta\right)^2=\int d^4xa^3\frac{1}{2}\left[\dot{\vartheta}^2-\left(\dot{H}_b+3H_aH_b-H_b^2\right)\vartheta^2\right],$$

$$S_0 = \int d^4xdy\sqrt{-G}\left[-D_M\Phi g^{MN}D_N\Phi^\dagger - m_0^2\Phi\Phi^\dagger\right]$$
$$= \int d^4xdya^3\left[|\dot{\Phi}|^2 - \frac{1}{a^2}|\partial_i\Phi|^2 - \frac{1}{b^3}\left|D_y\Phi\right|^2 - \frac{m_0^2}{b}|\Phi|^2\right]$$

$$\Phi(x^\mu, y) = \frac{1}{\sqrt{2\pi R}}\sum_{n\in\mathbb{Z}}\tilde{\phi}_n(x)e^{+\frac{\mathrm{i}ny}{R}}$$
$$\Phi^\dagger(x^\mu, y) = \frac{1}{\sqrt{2\pi R}}\sum_{n\in\mathbb{Z}}\tilde{\phi}_n^\dagger(x)e^{-\frac{\mathrm{i}ny}{R}}$$

$$S_{\text{gravity}} = \int d^4xa^3\sum_{n\in\mathbb{Z}}\left[\left|\dot{\tilde{\phi}}_n\right|^2 - \frac{1}{a^2}\left|\partial_i\tilde{\phi}_n\right|^2 - \frac{1}{b^3}\left|\left(\frac{n}{R} - qg_0b\vartheta\right)\tilde{\phi}_n\right|^2 - \frac{m_0^2}{b}\left|\tilde{\phi}_n\right|^2\right]$$
$$= \int d^4x\sum_{n\in\mathbb{Z}}\left[\left|\dot{\phi}_n - \frac{3}{2}H_a\phi_n\right|^2 - \frac{1}{a^2}|\partial_i\phi_n|^2 - \frac{1}{b^3}\left|\left(\frac{n}{R} - qg_0b\vartheta\right)\phi_n\right|^2 - \frac{m_0^2}{b}|\phi_n|^2\right]$$
$$= \int d^4x\sum_{n\in\mathbb{Z}}\left[\left|\dot{\phi}_n\right|^2 - |\partial_i\phi_n|^2 - M_{0,n}^2|\phi_n|^2\right]$$

$$M_{0,n}^2 \equiv M_{KK,n}^2 + \frac{m_0^2}{b} - \frac{9}{4}H_a^2 - \frac{3}{2}\dot{H}_a$$
$$M_{KK,n}^2 \equiv \frac{1}{b^3}\left(\frac{n}{R} - qg_0b\vartheta\right)^2$$

$$S_{\text{gravity}} = \int d^4xdy\sqrt{-G}\left[-\bar{\Psi}\gamma^Ae_A^MD_M\Psi + m_{1/2}\bar{\Psi}\Psi\right]$$

$$e_A^M = \text{diag}\left(b^{\frac{1}{2}}(t), \frac{b^{\frac{1}{2}}(t)}{a(t)}, \frac{b^{\frac{1}{2}}(t)}{a(t)}, \frac{b^{\frac{1}{2}}(t)}{a(t)}, \frac{1}{b(t)}\right)$$

$$D_M\Psi = \left(\partial_M + \frac{1}{4}\omega_M^{AB}\gamma_{AB} - \mathrm{i}qA_M\right)\Psi,$$

$$\omega_1^{\mathrm{k}0} = -\left(H_a - \frac{1}{2}H_b\right)b^{\frac{1}{2}}\delta_1^{\mathrm{k}}, \omega_5^{50} = -H_bb^{\frac{1}{2}}$$

$$\gamma^Ae_A^MD_M\Psi = b^{\frac{1}{2}}\left(\gamma^0\dot{\Psi} + \frac{1}{a}\gamma^{\mathrm{i}}\partial_i\Psi + \frac{1}{2}\left(3H_a - \frac{1}{2}H_b\right)\gamma^0\Psi + b^{-\frac{3}{2}}\gamma^5\left(\partial_y - \mathrm{i}qA_5\right)\Psi\right)$$
$$= a^{-\frac{5}{2}}b^{\frac{3}{4}}\left(\gamma^0\dot{\Psi}_c + \frac{1}{a}\gamma^{\mathrm{i}}\partial_i\Psi_c + b^{-\frac{3}{2}}\gamma^5\left(\partial_y - \mathrm{i}qgA_5\right)\Psi_c\right)$$

$$S_{\text{gravity}} = \int d^4xdy\sqrt{-G}\left[-\bar{\Psi}\gamma^Ae_A^MD_M\Psi + m_{1/2}\bar{\Psi}\Psi\right]$$
$$= \int d^4xdy\left[-\bar{\Psi}_c\left(\gamma^0\dot{\Psi}_c + \frac{1}{a}\gamma^{\mathrm{i}}\partial_i\Psi_c + \frac{1}{b^{\frac{3}{2}}}\gamma^5\left(\partial_y - \mathrm{i}qgA_5\right)\Psi_c\right) + \frac{m_{1/2}}{b^{\frac{1}{2}}}\bar{\Psi}_c\Psi_c\right]$$

$$\Psi_c(x, y) = \frac{1}{\sqrt{2\pi R}}\sum_{n\in\mathbb{Z}}\psi_n(x)e^{\frac{\mathrm{i}ny}{R}}$$

$$S_{\text{gravity}} = \int d^4x \sum_{n\in\mathbb{Z}} \left[-\bar{\psi}_n \left(\gamma^0 \dot{\psi}_n + \frac{1}{a}\gamma^{\mathrm{i}}\partial_i\psi_n + \frac{\mathrm{i}}{b^{\frac{3}{2}}}\gamma^5 \left(\frac{n}{R} - qg_0 b\vartheta\right)\psi_n \right) + \frac{m_{1/2}}{b^{\frac{1}{2}}}\bar{\psi}_n\psi_n \right]$$

$$\psi_n = \begin{pmatrix} \chi_{n\alpha} \\ \zeta_n^{\dagger\dot{\alpha}} \end{pmatrix}$$

$$\bar{\psi}_n = \left(\zeta_n^{\alpha}, \chi_{n\dot{\alpha}}^{\dagger}\right) = \mathrm{i}\psi_n^{\dagger}\gamma^0$$

$$S_{\text{gravity}} = \int d^4x \sum_{n\in\mathbb{Z}} \left[-\mathrm{i}\zeta_n\sigma^0\partial_0\zeta_n^{\dagger} - \mathrm{i}\chi_n\sigma^0\partial_0\chi_n^{\dagger} - \frac{\mathrm{i}}{a}\zeta_n\sigma^i\partial_i\zeta_n^{\dagger} - \frac{\mathrm{i}}{a}\chi_n\sigma^i\partial_i\chi_n^{\dagger} \right.$$
$$\left. + M_{1/2,n}(t)\zeta_n\chi_n + M_{1/2,n}^{*}(t)\zeta_n^{\dagger}\chi_n^{\dagger} \right]$$

$$M_{1/2,n} = \frac{m_{1/2}}{b^{\frac{1}{2}}} - \frac{\mathrm{i}}{b^{\frac{3}{2}}}\left(\frac{n}{R} - qg_0 b\vartheta\right)$$

$$\hat{\phi}_n(t,\boldsymbol{x}) = \int \frac{d^4\boldsymbol{k}}{(2\pi)^{\frac{3}{2}}}\left[\hat{a}_{n,\boldsymbol{k}} f_{n,k}(t) e^{\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}} + \hat{b}_{n,\boldsymbol{k}}^{\dagger} f_{n,k}^{*}(t) e^{-\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}}\right]$$
$$\hat{\phi}_n^{\dagger}(t,\boldsymbol{x}) = \int \frac{d^4\boldsymbol{k}}{(2\pi)^{\frac{3}{2}}}\left[\hat{b}_{n,\boldsymbol{k}} f_{n,k}(t) e^{\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}} + \hat{a}_{n,\boldsymbol{k}}^{\dagger} f_{n,k}^{*}(t) e^{-\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}}\right]$$

$\left(\hat{a}_{n,\boldsymbol{k}}^{\dagger}, \hat{a}_{n,\boldsymbol{k}}\right)$ and $\left(\hat{b}_{n,\boldsymbol{k}}^{\dagger}, \hat{b}_{n,\boldsymbol{k}}\right)$ satisfy $\left[\hat{a}_{n,\boldsymbol{k}}, \hat{a}_{n',\boldsymbol{k}'}^{\dagger}\right] = \delta_{nn'}\delta^3(\boldsymbol{k}-\boldsymbol{k}') = \left[\hat{b}_{n,\boldsymbol{k}}, \hat{b}_{n',\boldsymbol{k}'}^{\dagger}\right]$

$$\ddot{f}_{n,k}(t) + \omega_{0,n,k}^2(t) f_{n,k}(t) = 0$$

$$\omega_{0,n,k}^2(t) = \frac{k^2}{a^2(t)} + M_{0,n}^2(t)$$

$$f_{n,k}(t) = \frac{\alpha_{n,k}(t)}{\sqrt{2\omega_{0,n,k}(t)}} e^{-\mathrm{i}\int^t dt'\omega_{0,n,k}(t')} + \frac{\beta_{n,k}(t)}{\sqrt{2\omega_{0,n,k}(t)}} e^{+\mathrm{i}\int^t dt'\omega_{0,n,k}(t')}$$

$$\left|\alpha_{n,k}(t)\right|^2 - \left|\beta_{n,k}(t)\right|^2 = 1$$

$$\left[\hat{\phi}_n(t,\boldsymbol{x}), \hat{\pi}_{n,\phi}(t,\boldsymbol{y})\right] = \left[\hat{\phi}_n(t,\boldsymbol{x}), \hat{\phi}_n^{\dagger}(t,\boldsymbol{y})\right] = \mathrm{i}\delta^4(\boldsymbol{x}-\boldsymbol{y})$$

$$\frac{d}{dt}\begin{pmatrix}\alpha_{n,k}\\ \beta_{n,k}\end{pmatrix} = \begin{pmatrix} 0 & \frac{\dot{\omega}_{0,n,k}}{2\omega_{0,n,k}} e^{+2\mathrm{i}\int^t dt'\omega_{0,n,k}(t')} \\ \frac{\dot{\omega}_{0,n,k}}{2\omega_{0,n,k}} e^{-2\mathrm{i}\int^t dt'\omega_{0,n,k}(t')} & 0 \end{pmatrix}\begin{pmatrix}\alpha_{n,k}\\ \beta_{n,k}\end{pmatrix}$$

$\tilde{\alpha}_{n,k} \equiv \alpha_{n,k} e^{-\mathrm{i}\int^t dt'\omega_{0,n,k}(t')}$ and $\tilde{\beta}_{n,k} \equiv \beta_{n,k} e^{+\mathrm{i}\int^t dt'\omega_{0,n,k}(t')}$

$$\frac{d}{dt}\begin{pmatrix}\tilde{\alpha}_{n,k}\\ \tilde{\beta}_{n,k}\end{pmatrix} = \begin{pmatrix} \mathrm{i}\omega_{0,n,k} & \frac{\dot{\omega}_{0,n,k}}{2\omega_{0,n,k}} \\ \frac{\dot{\omega}_{0,n,k}}{2\omega_{0,n,k}} & -\mathrm{i}\omega_{0,n,k} \end{pmatrix}\begin{pmatrix}\tilde{\alpha}_{n,k}\\ \tilde{\beta}_{n,k}\end{pmatrix}$$

$$\hat{\chi}_n^{\alpha}(t,\boldsymbol{x}) = \sum_{h=\pm} \int \frac{d^3\boldsymbol{k}}{(2\pi)^{\frac{3}{2}}} \left[\hat{c}_{n,\boldsymbol{k},h} \eta_{n,k,h}(t) e^{\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}} \xi_h^{\alpha}(\hat{\boldsymbol{k}}) + \hat{d}_{n,\boldsymbol{k},h}^{\dagger} \bar{\lambda}_{n,k,h}(t) e^{-\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}} \xi_{h\dot{\beta}}^{\dagger}(\hat{\boldsymbol{k}}) \bar{\sigma}_0^{\dot{\beta}\alpha} \right],$$

$$\hat{\zeta}_n^{\alpha}(t,\boldsymbol{x}) = \sum_{h=\pm} \int \frac{d^3\boldsymbol{k}}{(2\pi)^{\frac{3}{2}}} \left[\hat{d}_{n,\boldsymbol{k},h} \eta_{n,k,h}(t) e^{\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}} \xi_h^{\alpha}(\hat{\boldsymbol{k}}) + \hat{c}_{n,\boldsymbol{k},h}^{\dagger} \bar{\lambda}_{n,k,h}(t) e^{-\mathrm{i}\boldsymbol{k}\cdot\boldsymbol{x}} \xi_{h\dot{\beta}}^{\dagger}(\hat{\boldsymbol{k}}) \bar{\sigma}_0^{\dot{\beta}\alpha} \right],$$

$$\begin{aligned}
\left((\hat{\boldsymbol{k}}\cdot\boldsymbol{\sigma})\xi_h\right)_{\alpha} &= \hat{k}_{\mathrm{i}} \left(\sigma_{\alpha\dot{\alpha}}^{0} \left(\bar{\sigma}^{\mathrm{i}}\right)^{\dot{\alpha}\beta}\right) \xi_{\beta}^{h}(\hat{\boldsymbol{k}}) = h\xi_{\alpha}^{h}(\hat{\boldsymbol{k}}) \\
\left(\xi_h^{\dagger}(\hat{\boldsymbol{k}}\cdot\boldsymbol{\sigma})\right)_{\dot{\alpha}} &= \xi_{h\dot{\beta}}^{\dagger}(\hat{\boldsymbol{k}}) (\bar{\sigma}_0)^{\dot{\beta}\alpha} \left(\hat{k}_{\mathrm{i}}\sigma^{\mathrm{i}}\right)_{\alpha\dot{\alpha}} = h\xi_{h\dot{\alpha}}^{\dagger}(\hat{\boldsymbol{k}}) \\
\xi_{h\dot{\beta}}^{\dagger}(-\hat{\boldsymbol{k}}) \bar{\sigma}_0^{\dot{\beta}\alpha} &\equiv \iota_h(\hat{\boldsymbol{k}}) \xi_h^{\alpha}(\hat{\boldsymbol{k}})
\end{aligned}$$

$$\left\{\hat{c}_{m,\boldsymbol{k},h}, \hat{c}_{n,\boldsymbol{k}',h'}^{\dagger}\right\} = \delta_{mn}\delta^3(\boldsymbol{k}-\boldsymbol{k}')\delta_{hh'}, \left\{\hat{d}_{m,\boldsymbol{k},h}, \hat{d}_{n,\boldsymbol{k}',h'}^{\dagger}\right\} = \delta_{mn}\delta^3(\boldsymbol{k}-\boldsymbol{k}')\delta_{hh'},$$

$$\left|\eta_{n,k,h}(t)\right|^2 + \left|\lambda_{n,k,h}(t)\right|^2 = 1$$

$$\mathrm{i}\frac{d}{dt}\begin{pmatrix}\eta_{n,k,h}\\ \lambda_{n,k,h}\end{pmatrix} = \begin{pmatrix} -\dfrac{kh}{a} & -M_{1/2,n}^{*} \\ -M_{1/2,n} & \dfrac{kh}{a}\end{pmatrix}\begin{pmatrix}\eta_{n,k,h}\\ \lambda_{n,k,h}\end{pmatrix}.$$

$$M_{1/2,n} = \mu_n e^{\mathrm{i}\Theta_n}$$

$$\begin{aligned}
\cos\,\Theta_n &= \frac{m_{1/2}}{\sqrt{m_{1/2}^2 + \left(\dfrac{n}{Rb} - qg_0\vartheta\right)^2}} \\
\sin\,\Theta_n &= -\frac{\left(\dfrac{n}{Rb} - qg_0\vartheta\right)}{\sqrt{m_{1/2}^2 + \left(\dfrac{n}{Rb} - qg_0\vartheta\right)^2}}
\end{aligned}$$

$$\mathrm{i}\frac{d}{dt}\begin{pmatrix}\eta_{n,k,h}\\ \lambda_{n,k,h}\end{pmatrix} = \omega_{n,k}\begin{pmatrix}\cos\,\theta_{n,k,h} & \sin\,\theta_{n,k,h} e^{-\mathrm{i}\Theta_n} \\ \sin\,\theta_{n,k,h} e^{\mathrm{i}\Theta_n} & -\cos\,\theta_{n,k,h}\end{pmatrix}\begin{pmatrix}\eta_{n,k,h}\\ \lambda_{n,k,h}\end{pmatrix},$$

$$\omega_{n,k} = \sqrt{\frac{k^2}{a^2} + \mu_n^2}$$

$$\cos\,\theta_{n,k,h} = -\frac{kh}{a\omega_{n,k}} \text{ and } \sin\,\theta_{n,k,h} = -\frac{\mu_{n,k}}{\omega_{n,k}}$$

$$\boldsymbol{v}_{n,k,h}^{+} = \begin{pmatrix} e^{-\mathrm{i}\Theta_n}\cos\,\dfrac{1}{2}\theta_{n,k,h} \\ \sin\,\dfrac{1}{2}\theta_{n,k,h}\end{pmatrix}, \boldsymbol{v}_{n,k,h}^{-} = \begin{pmatrix} -e^{-\mathrm{i}\Theta_n}\sin\,\dfrac{1}{2}\theta_{n,k,h} \\ \cos\,\dfrac{1}{2}\theta_{n,k,h}\end{pmatrix}$$

$$\begin{pmatrix}\eta_{n,k,h}\\ \lambda_{n,k,h}\end{pmatrix} = \gamma_{n,k,h}(t) e^{-\mathrm{i}\int^t dt'\omega_{n,k}(t')} \boldsymbol{v}_{n,k,h}^{+} + \delta_{n,k,h}(t) e^{+\mathrm{i}\int^t dt'\omega_{n,k}(t')} \boldsymbol{v}_{n,k,h}^{-}$$

$$\left|\gamma_{n,k,h}(t)\right|^2 + \left|\delta_{n,k,h}(t)\right|^2 = 1$$

$$\dot{\tilde{\gamma}}_{n,k,h} = -\mathrm{i}\omega_{n,k}\tilde{\gamma}_{n,k,h} + \frac{\mathrm{i}}{2}\dot{\Theta}_n\left(1+\cos\,\theta_{n,k,h}\right)\tilde{\gamma}_{n,k,h} + \frac{1}{2}\left(\dot{\theta}_{n,k,h} - \mathrm{i}\dot{\Theta}_n\sin\,\theta_{n,k,h}\right)\tilde{\delta}_{n,k,h}$$
$$\dot{\tilde{\delta}}_{n,k,h} = +\mathrm{i}\omega_{n,k}\tilde{\delta}_{n,k,h} + \frac{\mathrm{i}}{2}\dot{\Theta}_n\left(1-\cos\,\theta_{n,k,h}\right)\tilde{\delta}_{n,k,h} - \frac{1}{2}\left(\dot{\theta}_{n,k,h} + \mathrm{i}\dot{\Theta}_n\sin\,\theta_{n,k,h}\right)\tilde{\gamma}_{n,k,h}$$

$$\tilde{\gamma}_{n,k,h} = \gamma_{n,k,h}e^{-\mathrm{i}\int^t dt'\omega_{n,k}(t')} \text{ and } \tilde{\delta}_{n,k,h} = \delta_{n,k,h}e^{+\mathrm{i}\int^t dt'\omega_{n,k}(t')}$$

$$\hat{A}_{n,\boldsymbol{k}}(t) \equiv \alpha_{n,k}(t)\hat{a}_{n,\boldsymbol{k}} + \beta^*_{n,k}(t)\hat{b}^\dagger_{n,-\boldsymbol{k}}$$
$$\hat{A}^\dagger_{n,\boldsymbol{k}}(t) \equiv \alpha^*_{n,k}(t)\hat{a}^\dagger_{n,\boldsymbol{k}} + \beta_{n,k}(t)\hat{b}_{n,-\boldsymbol{k}}$$

$$\hat{N}_{n,\boldsymbol{k}}(t) = \hat{A}^\dagger_{n,\boldsymbol{k}}(t)\hat{A}_{n,\boldsymbol{k}}(t)$$

$$\left\langle N_{n,\boldsymbol{k}}(t)\right\rangle = {}_{\mathrm{in}}\langle 0|\hat{A}^\dagger_{n,\boldsymbol{k}}(t)\hat{A}_{n,\boldsymbol{k}}(t)|0\rangle_{\mathrm{in}} = \delta^3(\boldsymbol{k}\to 0)\left|\beta_{n,k}(t)\right|^2 = V_{3D}\left|\beta_{n,k}(t)\right|^2,$$

$$M^2_{0,n,A} \approx m^2_{KK,n,A} + \frac{m^2_{0,A}}{b}$$

$$3M^2_{\mathrm{pl}}H^2_a = \frac{1}{2}\dot{\varphi}^2 + \frac{1}{2}\left(\dot{\vartheta} + \sqrt{\frac{2}{3}}\frac{\dot{\varphi}}{M_{\mathrm{pl}}}\vartheta\right)^2 + V_{\mathrm{loop}} + V_\Lambda$$
$$\ddot{\varphi} + 3H_a\dot{\varphi} - \frac{1}{M_{\mathrm{pl}}}\sqrt{\frac{3}{2}}\left(\dot{\vartheta} + \sqrt{\frac{2}{3}}\frac{\dot{\varphi}}{M_{\mathrm{pl}}}\vartheta\right)^2 + \partial_\varphi V_{\mathrm{loop}} + \partial_\varphi V_\Lambda = 0$$
$$\ddot{\vartheta} + 3H_a\dot{\vartheta} - \left(\frac{\sqrt{6}H_a\dot{\varphi}}{M_{\mathrm{pl}}} + \sqrt{\frac{2}{3}}\frac{\ddot{\varphi}}{M_{\mathrm{pl}}} + \frac{2}{3}\frac{\dot{\varphi}^2}{M^2_{\mathrm{pl}}}\right)\vartheta + \partial_\vartheta V_{\mathrm{loop}} = 0$$

$$V_\Lambda = 2\pi R\Lambda^5 e^{-\sqrt{\frac{2}{3}}\varphi/M_{\mathrm{pl}}}$$
$$V_{\mathrm{loop}} = \sum_A \frac{N_A}{2\pi^2L^4b^2}\left[\frac{\left(LM_{0,A}\right)^5}{30} - \sum_{n=1}^{\infty}\cos\left(nq_Ag_0\vartheta L\right)e^{-x_{n,0,A}}\frac{3+3x_{n,0,A}+x^2_{n,0,A}}{n^5}\right]$$
$$-\sum_B \frac{N_B}{\pi^2L^4b^2}\left[\frac{\left(Lm_{1/2,B}\right)^5}{30} - \sum_{n=1}^{\infty}\cos\left(nq_Bg_0\vartheta L\right)e^{-x_{n,1/2,B}}\frac{3+3x_{n,1/2,B}+x^2_{n,1/2,B}}{n^5}\right],$$

$$x_{n,0,A} \equiv nLM_{0,A}, x_{n,1/2,B} \equiv nLm_{1/2,B}$$

$$R \to C^{-1}R, m_{0,A} \to Cm_{0,A}, m_{1/2,B} \to Cm_{1/2,B}, g_0 \to Cg_0, 2\pi R\Lambda^5 \to 2\pi R\Lambda^5C^4$$

$$N^{(0)}_{n,k} \equiv \left|\beta_{n,k}\right|^2 = \left|\tilde{\beta}_{n,k}\right|^2, R^{(0)}_{n,k} \equiv \mathrm{Re}\left(\tilde{\alpha}_{n,k}\overline{\tilde{\beta}}_{n,k}\right) \text{ and } I^{(0)}_{n,k} \equiv \mathrm{Im}\left(\tilde{\alpha}_{n,k}\overline{\tilde{\beta}}_{n,k}\right)$$

$$\frac{d}{dt}N^{(0)}_{n,k,A} = \frac{\dot{\omega}_{0,n,k,A}}{\omega_{0,n,k,A}}R^{(0)}_{n,k,A}$$
$$\frac{d}{dt}R^{(0)}_{n,k,A} = \frac{\dot{\omega}_{0,n,k,A}}{2\omega_{0,n,k,A}}\left(1+2N^{(0)}_{n,k,A}\right) - 2\omega_{0,n,k,A}I^{(0)}_{n,k,A}$$
$$\frac{d}{dt}I^{(0)}_{n,k,A} = 2\omega_{0,n,k,A}R^{(0)}_{n,k,A}$$

$$N_{n,p,h}^{(1/2)} = \left|\delta_{n,p,h}\right|^2, R_{n,p,h}^{(1/2)} = \mathrm{Re}\left(\tilde{\gamma}_{n,p,h}\bar{\tilde{\delta}}_{n,p,h}\right) \text{ and } I_{n,p,h}^{(1/2)} = \mathrm{Im}\left(\tilde{\gamma}_{n,p,h}\bar{\tilde{\delta}}_{n,p,h}\right)$$

$$\frac{d}{dt}N_{n,k,h,B}^{(1/2)} = -\dot{\theta}_{n,k,h,B}R_{n,k,h,B}^{(1/2)} + \dot{\Theta}_{n,B}\sin\,\theta_{n,k,h,B}I_{n,k,h,B}^{(1/2)}$$
$$\frac{d}{dt}R_{n,k,h,B}^{(1/2)} = \left(2\omega_{n,k,B} - \dot{\Theta}_{n,B}\cos\,\theta_{n,k,h,B}\right)I_{n,k,h,B} + \dot{\theta}_{n,k,h,B}\left(2N_{n,k,h,B}^{(1/2)} - 1\right)$$
$$\frac{d}{dt}I_{n,k,h,B}^{(1/2)} = -\left(2\omega_{n,k,B} - \dot{\Theta}_{n,B}\cos\,\theta_{n,k,h,B}\right)R_{n,k,h,B}^{(1/2)} - \dot{\Theta}_{n,B}\sin\,\theta_{n,k,h,B}\left(2N_{n,k,h,B}^{(1/2)} - 1\right)$$

$$V_{\mathrm{loop}} \approx C - \sum_A C_A\cos\left(\frac{q_A\vartheta}{f_{\mathrm{eff}}}\right) + \sum_B C_B\cos\left(\frac{q_B\vartheta}{f_{\mathrm{eff}}}\right)$$

$$C_A = \frac{N_A\left(3 + 3x_{1,0,A} + x_{1,0,A}^2\right)e^{-x_{1,0,A}}}{2\pi^2L^4b^2}$$
$$C_B = \frac{N_B\left(3 + 3x_{1,1/2,B} + x_{1,1/2,B}^2\right)e^{-x_{1,1/2,B}}}{\pi^2L^4b^2}$$

$$\vartheta \in \left[-\frac{\pi}{2q_0g_0L}, \frac{\pi}{2q_0g_0L}\right] = \left[-\frac{1}{4q_0g_0Rb}, \frac{1}{4q_0g_0Rb}\right]$$

$$\frac{(|n| - 1/4)^2}{R^2b^3} < m_{KK,n,q_0}^2 < \frac{(|n| + 1/4)^2}{R^2b^3}$$

$$\frac{(|n| - |q_1/(4q_0)|)^2}{R^2b^3} < m_{KK,n,q_1}^2 < \frac{(|n| + |q_1/(4q_0)|)^2}{R^2b^3}$$

$$R = 40 \times C^{-1}, g_0 = 0.02C,$$
$$2\pi R\Lambda^5 = 1.1182 \times 10^{-3}C^4,$$
$$\left(N_A, m_{0,A}, q_A\right) = (28,1.8 \times 10^{-1}C, 0), (2,5 \times 10^{-4}C, 1)\ (2,2.014 \times 10^{-4}C, 5)$$
$$\left(N_B, m_{1/2,B}, q_B\right) = (16,1.9 \times 10^{-1}C, 0), (1,5.2 \times 10^{-4}C, 1), (1,2 \times 10^{-4}C, 5).$$

$$\langle L\rangle = 2\pi R\langle b\rangle = 63.48$$

$$\epsilon = \frac{1}{2}\left(\frac{\partial_\vartheta V}{V}\right)^2, \eta = \frac{\partial_\vartheta^2 V}{V}.$$

$$\omega_{0,n,k,A}^2 \approx \frac{k^2}{a^2} + \frac{m_0^2}{b} + \frac{1}{b^3}\left(\frac{n}{R} - q_Ag_0b\vartheta\right)^2$$

$$\frac{n}{R} - q_Ag_0b\vartheta\left(t_{A,n}\right) = 0$$

$$\omega_{0,n,k,A}^2(t) \approx \left(\omega_{0,n,k,A}^{\min}\right)^2 + v_{n,A}^2\left(t - t_{n,A}\right)^2$$

$$\omega_{0,n,k,A}^{\min} \equiv \sqrt{\frac{k^2}{a^2\left(t_{n,A}\right)} + \frac{m_0^2}{b}}$$
$$v_{n,A} \equiv q_Ag_0b^{-\frac{1}{2}}\dot{\vartheta}\Big|_{t=t_{n,A}}$$

$$N_{n,k,A}(t) \approx \begin{cases} 0 \ (\text{ for } t < t_{n,A}) \\ \exp\left(-\dfrac{\pi\left(\omega_{0,n,k,A}^{\min}\right)^2}{v_{n,A}}\right) \ (\text{ for } t > t_{n,A}). \end{cases}$$

$$\begin{aligned} \rho_{n,A} &\approx \int \frac{d^4\boldsymbol{k}}{(2\pi)^3 a^3(t)} 2\omega_{0,n,k,A} N_{n,k,A}^{(0)}(t) \\ &\approx \int \frac{d^4\boldsymbol{k}}{(2\pi)^3 a^3(t)} 2\omega_{0,n,k,A} \exp\left[-\frac{\pi\left(k^2/a\left(t_{n,A}\right)^2 + m_0^2/b\right)}{v_{A,n}}\right] \theta_H\left(t - t_{n,A}\right) \\ &= \frac{v_{n,A}}{4\pi^3}\left(\frac{a_{n,A}}{a}\right)^2 e^{-\frac{m_{0,A}^2}{b v_{n,A}}} e^{-\frac{\pi M_{0,n,A}^2}{2v_{n,A}}\frac{a^2}{a_{n,A}^2}} M_{0,n,A}^2 K_1\left(\frac{\pi M_{0,n,A}^2}{2v_{n,A}}\frac{a^2}{a_{n,A}^2}\right)\theta_H\left(t - t_{n,A}\right) \\ &= \frac{1}{4\pi^3}\left(\frac{a_{n,A}}{a}\right)^3 v_{A,n}^{\frac{3}{2}} M_{KK,n,A} \exp\left(-\frac{\pi m_{0,A}^2}{b v_{A,n}}\right)(1 + \cdots) \end{aligned}$$

$$r_{KK} \equiv \frac{\rho_{KK}}{\rho_{\text{inf}}}\Big|_{\text{R}} = \frac{\rho_{KK}}{3H^2}\Big|_{\text{R}}$$

$$\frac{\rho_{KK}}{s_R} = \frac{3}{4} r_{KK} T_R$$

$$r_{KK} \leq 5.6 \times 10^{-21}\left(\frac{10^{11}\text{GeV}}{T_R}\right)$$

$$\dot{n}_{KK} + 3Hn_{KK} = -\langle\sigma v\rangle n_{KK}^2,$$

$$n_{KK}(t) = \int \frac{d^4\boldsymbol{k}}{(2\pi)^3} N_{n=1,k}^{(0)}(t)$$

$$g_0 = 0.1, 2\pi R\Lambda^5 = 2.83759 \times 10^{-7}, R = 40,$$
$$\left(N_A, m_{0,A}, q_A\right) = (28, 3.8 \times 10^{-2}, 0), (2, 5 \times 10^{-6}, 1), (2, 2 \times 10^{-4}, 5),$$
$$\left(N_B, m_{1/2,B}, q_B\right) = (16, 3.9 \times 10^{-2}, 0), (1, 6 \times 10^{-6}, 1), (1, 2 \times 10^{-4}, 5),$$

$$L = 2\pi R\langle b\rangle \simeq 193,$$

$$\rho_m = \rho_0\left(\frac{a_0}{a(t)}\right)^3$$

$$\rho_{\text{tot}} = \rho_0\left(\frac{a_0}{a(t)}\right)^3 + \frac{1}{2}\dot{\vartheta}^2 + V_{\text{loop}}$$

$$A_\mu(x, -y) = A_\mu(x, y), A_y(x, -y) = -A_y(x, y)$$

$$A_\mu(x, -y) = -A_\mu(x, y), A_y(x, -y) = A_y(x, y)$$

$$\sigma^0 = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma^2 = \begin{pmatrix} 0 & -\text{i} \\ \text{i} & 0 \end{pmatrix}, \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

$$\bar{\sigma}^0 = \sigma^0 \text{ and } \sigma^i = -\bar{\sigma}^i (i = 1,2,3)$$

$$\Psi = \begin{pmatrix} \chi_\alpha \\ \bar{\psi}^{\dot{\alpha}} \end{pmatrix}$$

$$\gamma^\mu = \mathrm{i}\begin{pmatrix} 0 & \sigma^\mu \\ \bar{\sigma}^\mu & 0 \end{pmatrix}$$

$$\gamma^5 = \mathrm{i}\gamma^0\gamma^1\gamma^2\gamma^3 = \begin{pmatrix} \mathbf{1}_{2\times2} & 0 \\ 0 & -\mathbf{1}_{2\times2} \end{pmatrix}$$

$$\begin{aligned}
T^{(0)}_{MN} =& 2D_M\Phi D_N\bar{\Phi} + g_{MN}\left(-D_P\Phi D^P\bar{\Phi} - m_0^2|\Phi|^2\right), \\
T^{(1/2)}_{MN} =& -\frac{1}{2}\bar{\Psi}\gamma_{(M}\overleftrightarrow{D}_{N)}\Psi + g_{MN}\left(-\bar{\Psi}\gamma^P D_P\Psi + m_{1/2}\bar{\Psi}\Psi\right) \\
\overset{\text{on-shell}}{\rightarrow}& -\frac{1}{2}\bar{\Psi}\gamma_{(M}\overleftrightarrow{D}_{N)}\Psi, \\
T^{(1)}_{MN} =& F_{MP}F_N{}^P - \frac{1}{4}g_{MN}F_{PQ}F^{PQ},
\end{aligned}$$

$$A_{(M}B_{N)} = \frac{1}{2}(A_MB_N + A_NB_M)$$

$$T^{\Lambda}_{MN} = -g_{MN}\Lambda^5$$

$$\begin{aligned}
\langle\hat{T}_{00}\rangle =& \left\langle 2\dot{\hat{\Phi}}\dot{\hat{\Phi}} - \frac{1}{b}\left(b\dot{\hat{\Phi}}\dot{\hat{\Phi}}^\dagger - \frac{b}{a^2}\partial_i\hat{\Phi}\partial_i\hat{\Phi}^\dagger - b^{-2}D_y\hat{\Phi}D_y\hat{\Phi}^\dagger - m_0^2\hat{\Phi}\hat{\Phi}^\dagger\right)\right\rangle \\
=& \frac{1}{2\pi Ra^3}\sum_n\left\langle\left(\dot{\hat{\phi}}_n - \frac{3}{2}H_a\hat{\phi}_n\right)\left(\dot{\hat{\phi}}_n^\dagger - \frac{3}{2}H_a\hat{\phi}_n^\dagger\right) + \partial_i\hat{\phi}_n\partial_i\hat{\phi}_n^\dagger + \frac{1}{b^3}\left(\frac{n}{R} - qgA_5\right)^2\left|\hat{\phi}_n\right|^2 + \frac{m_0^2}{b}\left|\hat{\phi}_n\right|^2\right\rangle \\
=& \frac{1}{2\pi Ra^3}\int\frac{d^4\boldsymbol{k}}{(2\pi)^3}\sum_n\left[\left|\dot{f}_{n,k} - \frac{3}{2}H_af_{n,k}\right|^2 + \left(\omega_{0,n,k}^2 + \frac{3}{2}\dot{H}_a + \frac{9}{4}H_a^2\right)|f_{n,k}|^2\right] \\
=& \frac{1}{2\pi Ra^3}\int\frac{d^4\boldsymbol{k}}{(2\pi)^3}\sum_n\left[|\dot{f}_{n,k}|^2 + \left(\omega_{0,n,k}^2 + \frac{3}{2}\dot{H}_a + \frac{9}{2}H_a^2\right)|f_{n,k}|^2 - \frac{3H_a}{2}\frac{d}{dt}|f_{n,k}|^2\right]
\end{aligned}$$

$$\dot{f}_{n,k}(t) = \frac{\mathrm{i}\omega_{0,n,k}}{\sqrt{2\omega_{0,n,k}}}\left(\tilde{\alpha}_{n,k} - \tilde{\beta}_{n,k}\right)$$

$$\begin{aligned}
|\dot{f}_{n,k}|^2 &= \frac{\omega_{0,n,k}}{2}\left(1 + 2N_{n,k}^{(0)} - 2R_{n,k}^{(0)}\right) \\
|f_{n,k}|^2 &= \frac{1}{2\omega_{0,n,k}}\left(1 + 2N_{n,k}^{(0)} + 2R_{n,k}^{(0)}\right) \\
\frac{d}{dt}|f_{n,k}|^2 &= -I_{n,k}^{(0)}
\end{aligned}$$

$$\tilde{\alpha}_{n,k}\bar{\tilde{\beta}}_{n,k} = R_{n,k}^{(0)} + \mathrm{i}I_{n,k}^{(0)}$$

$$R_{n,k}, I_{n,k} \in \mathbb{R} \text{ and } N_{n,k}^{(0)} = \left|\beta_{n,k}\right|^2$$

$$\langle \hat{T}_{00} \rangle = \frac{1}{2\pi R a^3} \sum_{n\in\mathbb{Z}} \int \frac{d^4\boldsymbol{k}}{(2\pi)^3} \left[\omega_{0,n,k} + \frac{3}{4}\frac{\dot{H}_a + 3H_a^2}{\omega_{0,n,k}} + 2\left(\omega_{0,n,k} + \frac{3}{4}\frac{\dot{H}_a + 3H_a^2}{\omega_{0,n,k}} \right) N_{n,k}^{(0)} + \frac{3}{2}\frac{\dot{H}_a + 3H_a^2}{\omega_{0,n,k}} R_{n,k}^{(0)} + \frac{3H_a}{2} I_{n,k}^{(0)} \right]$$

$$\begin{aligned} \langle \hat{T}_{00} \rangle_{\text{vac}} &= \frac{1}{2\pi R a^D} \sum_{n\in\mathbb{Z}} \mu^{3-D} \int \frac{d^D\boldsymbol{k}}{(2\pi)^D} \left(\omega_{0,n,k} + \frac{3}{4}\frac{\dot{H}_a + 3H_a^2}{\omega_{0,n,k}} \right) \\ &= \frac{1}{2\pi R} \sum_{n\in\mathbb{Z}} \frac{2\mu^{3-D}}{\Gamma\left(\frac{D}{2}\right)} \int \frac{d\tilde{k}}{(4\pi)^{D/2}} \tilde{k}^{D-1} \left(\omega_{0,n,k} + \frac{3}{4}\frac{\dot{H}_a + 3H_a^2}{\omega_{0,n,k}} \right) \\ &= \frac{1}{2\pi R} \sum_{n\in\mathbb{Z}} \frac{\mu^{3-D}}{(4\pi)^{\frac{D+1}{2}}} \left[-\Gamma\left(-\frac{D+1}{2} \right) M_{0,n}^{D+1} + \frac{3(\dot{H}_a + 3H_a^2)}{2} \Gamma\left(-\frac{D-1}{2} \right) M_{0,n}^{D-1} \right]. \end{aligned}$$

$$\begin{aligned} &\sum_{n=-\infty}^{\infty} [A(n+c)^2 + Q]^{-s} \\ &= \sqrt{\frac{\pi}{A}} \frac{\Gamma(s - 1/2)}{\Gamma(s)} Q^{\frac{1}{2}-s} + \frac{4\pi^s}{\Gamma(s)} A^{-\frac{1}{4}-\frac{s}{2}} Q^{\frac{1}{4}-\frac{s}{2}} \sum_{n=1}^{\infty} n^{s-\frac{1}{2}} \cos(2\pi n c) K_{s-\frac{1}{2}}\left(2\pi n \sqrt{\frac{Q}{A}} \right), \end{aligned}$$

$$s = -D/2 \mp 1/2, A = \tfrac{1}{R^2 b^3}, c = -q g_0 b \vartheta R \text{ and } Q = \frac{m_0^2}{b} - \frac{9}{4}H_a^2 - \frac{3}{2}\dot{H}_a$$

$$\begin{aligned} \sum_{n\in\mathbb{Z}} M_{0,n}^{D+1} &= \frac{\sqrt{\pi} L}{2\pi} \frac{\Gamma\left(-\frac{D}{2} - 1\right)}{\Gamma\left(-\frac{D+1}{2}\right) b^{\frac{D+1}{2}}} M_0^{2+D} \\ &\quad + \frac{2^{\frac{D}{2}+2} M_0}{\sqrt{\pi}\Gamma\left(-\frac{D+1}{2}\right) b^{\frac{D+1}{2}}} \left(\frac{M_0}{L}\right)^{\frac{D}{2}} \sum_{n=1}^{\infty} n^{-1-\frac{D}{2}} \cos(nqg_0\vartheta L) K_{-1-\frac{D}{2}}(nLM_0) \\ \sum_{n\in\mathbb{Z}} M_{0,n}^{D-1} &= \frac{\sqrt{\pi} L}{2\pi} \frac{\Gamma\left(-\frac{D}{2}\right)}{\Gamma\left(-\frac{D-1}{2}\right) b^{\frac{D-1}{2}}} M_0^{D} \\ &\quad + \frac{2^{\frac{D}{2}+1} L}{\sqrt{\pi}\Gamma\left(-\frac{D-1}{2}\right) b^{\frac{D-1}{2}}} \left(\frac{M_0}{L}\right)^{\frac{D}{2}} \sum_{n=1}^{\infty} n^{-\frac{D}{2}} \cos(nqg_0\vartheta L) K_{-\frac{D}{2}}(nLM_0), \end{aligned}$$

$$M_0^2 \equiv m_0^2 - b\left(\frac{3}{2}\dot{H}_a + \frac{9}{4}H_a^2 \right)$$

$$\langle \hat{T}_{00} \rangle_{\text{vac}}$$
$$= \frac{1}{2\pi R} \frac{\mu^{3-D}}{(4\pi)^{\frac{D+1}{2}}} \left[-\frac{L}{2\sqrt{\pi}} \frac{\Gamma\left(-\frac{D}{2}-1\right) M_0^{2+D}}{b^{\frac{D+1}{2}}} - \frac{2^{\frac{D}{2}+2} M_0}{\sqrt{\pi} b^{\frac{D+1}{2}}} \left(\frac{M_0}{L}\right)^{\frac{D}{2}} \sum_{n=1}^{\infty} n^{-1-\frac{D}{2}} \cos(nqg_0\vartheta L) K_{-1-\frac{D}{2}}(nLM_0) \right.$$
$$\left. + \frac{3(\dot{H}_a + 3H_a^2)}{2} \left\{ \frac{L}{2\sqrt{\pi}} \frac{\Gamma\left(-\frac{D}{2}\right)}{b^{\frac{D-1}{2}}} M_0^D + \frac{2^{\frac{D}{2}+1} L}{\sqrt{\pi} b^{\frac{D-1}{2}}} \left(\frac{M_0}{L}\right)^{\frac{D}{2}} \sum_{n=1}^{\infty} n^{-\frac{D}{2}} \cos(nqg_0\vartheta L) K_{-\frac{D}{2}}(nLM_0) \right\} \right]$$
$$\to \frac{1}{L^5 b} \left[\frac{(M_0 L)^5}{60\pi^2} - \frac{1}{2\pi^2} \sum_{n=1}^{\infty} \cos(nqg_0\vartheta L) e^{-x_{n,0}} \frac{3 + 3x_{n,0} + x_{n,0}^2}{n^5} \right]$$
$$+ \frac{3(\dot{H}_a + 3H_a^2)}{2L^3} \left\{ \frac{(LM_0)^3}{48\pi^2} + \frac{1}{4\pi^2} \sum_{n=1}^{\infty} \cos(nqg_0\vartheta L) e^{-x_{n,0}} \frac{1 + x_{n,0}}{n^3} \right\}$$

$$\langle \hat{T}_{ii} \rangle = \left\langle 2\partial_i \hat{\Phi} \partial_i \hat{\Phi}^\dagger + \frac{a^2}{b} \left(b \dot{\hat{\Phi}} \dot{\hat{\Phi}}^\dagger - \frac{b}{a^2} \sum_{j=1}^{3} \partial_j \hat{\Phi} \partial_j \hat{\Phi}^\dagger - \frac{1}{b^2} D_y \hat{\Phi} D_y \hat{\Phi}^\dagger - m_0^2 \hat{\Phi} \hat{\Phi}^\dagger \right) \right\rangle$$
$$= \frac{1}{2\pi R a} \sum_{n \in \mathbb{Z}} \int \frac{d^4k}{(2\pi)^3} \left[\frac{1}{\omega_{0,n,k}} \left(\frac{k^2}{3a^2} - \frac{3}{4}\dot{H}_a \right) \left(1 + 2N_{n,k}^{(0)}\right) - 2 \left(\omega_{0,n,k} - \frac{\frac{k^2}{3a^2} - \frac{3}{4}\dot{H}_a}{\omega_{0,n,k}} \right) R_{n,k}^{(0)} + \frac{3}{2} H_a I_{n,k}^{(0)} \right]$$

$$\frac{1}{a^2} \langle \hat{T}_{ii} \rangle_{\text{vac}} = \frac{1}{2\pi R a^3} \sum_{n \in \mathbb{Z}} \int \frac{d^4\boldsymbol{k}}{(2\pi)^3} \left[\frac{1}{\omega_{0,n,k}} \left(\frac{k^2}{3a^2} - \frac{3}{4}\dot{H}_a \right) \right]$$
$$\to \frac{1}{L^5 b} \left[-\frac{(LM_0)^5}{60\pi^2} + \frac{1}{2\pi^2} \sum_{n=1}^{\infty} \cos(nqg_0\vartheta L) e^{-x_{n,0}} \frac{3 + 3x_{n,0} + x_{n,0}^2}{n^5} \right.$$
$$\left. - \frac{b\dot{H}_a L^2 (M_0 L)^3}{8\pi} - \frac{3\dot{H}_a L^2}{8\pi^2} \sum_{n=1}^{\infty} \cos(nqg_0\vartheta L) e^{-x_{n,0}} \frac{1 + x_{n,0}}{n^3} \right]$$

$$\langle \hat{T}_{55} \rangle = \left\langle 2D_y \hat{\Phi} D_y \hat{\Phi}^\dagger + b^2 \left(b \dot{\hat{\Phi}} \dot{\hat{\Phi}}^\dagger - \frac{b}{a^2} \partial_j \hat{\Phi} \partial_j \hat{\Phi}^\dagger - \frac{1}{b^2} D_y \hat{\Phi} D_y \hat{\Phi}^\dagger - m_0^2 \hat{\Phi} \hat{\Phi}^\dagger \right) \right\rangle$$
$$= \frac{b^3}{2\pi R a^3} \sum_{n} \int \frac{d^4\boldsymbol{k}}{(2\pi)^3} \left[-R \frac{\partial}{\partial R} \omega_{0,n,k} - \frac{3\dot{H}_a}{4\omega_{0,n,k}} + \frac{2}{\omega_{0,n,k}} \left(m_{KK,n}^2 - \frac{3}{4}\dot{H}_a \right) N_{n,k}^{(0)} \right.$$
$$\left. + \frac{1}{\omega_{0,n,k}} \left(\frac{k^2}{a^2} + \frac{m_0^2}{b} + \frac{3}{2}\dot{H}_a - \frac{9}{4}H_a^2 \right) R_{n,k}^{(0)} + \frac{3}{2} H_a I_{n,k}^{(0)} \right]$$

$$\frac{\left(n - \frac{qg_0\vartheta L}{2\pi}\right)^2}{R^2 b^3 \omega_{0,n,k}} = -R \frac{\partial}{\partial R} \omega_{0,n,k}$$

$$\frac{1}{b^3} \langle \hat{T}_{55} \rangle_{\text{vac}} = \frac{1}{2\pi R a^3} \sum_{n \in \mathbb{Z}} \int \frac{d^4\boldsymbol{k}}{(2\pi)^3} \left[-R \frac{\partial}{\partial R} \omega_{0,n,k} - \frac{3\dot{H}_a}{4\omega_{0,n,k}} \right]$$
$$\to -\frac{1}{L^5 b} \left[\frac{(M_0 L)^5}{60\pi^2} + \frac{1}{2\pi^2} \sum_{n=1}^{\infty} \cos(nqg_0\vartheta L) e^{-x_{n,0}} \frac{12 + 12x_{n,0} + 5x_{n,0}^2 + x_{n,0}^3}{n^5} \right]$$
$$- \frac{3\dot{H}_a}{2L^3} \left\{ \frac{(LM_0)^3}{48\pi^2} + \frac{1}{4\pi^2} \sum_{n=1}^{\infty} n^{-\frac{3}{2}} \cos(nqg_0\vartheta L) e^{-x_{n,0}} \frac{1 + x_{n,0}}{n^3} \right\}$$

$$\hat{T}_{MN}^{(1/2)} = \frac{1}{2}\hat{\bar{\Psi}}\gamma_{(M}\overleftrightarrow{D}_{N)}\hat{\Psi}.$$

$$\left\langle \hat{T}_{00}^{(1/2)} \right\rangle = \frac{1}{2}\left\langle \hat{\bar{\Psi}}\gamma_0\dot{\hat{\Psi}} - \dot{\hat{\bar{\Psi}}}\gamma_0\hat{\Psi} \right\rangle$$

$$= \frac{1}{2\pi R a^3}\sum_{n\in\mathbb{Z},h=\pm}\int \frac{d^4\boldsymbol{k}}{(2\pi)^3}\left[-\omega_{n,k} + 2\omega_{n,k}N_{n,k,h}^{(1/2)} + 2\mu_n\left(\frac{\omega_{n,k} - \frac{kh}{a}}{\omega_{n,k}}\right)R_{n,k,h}^{(1/2)}\right],$$

$$N_{n,k,h}^{(1/2)} = \left|\delta_{n,k,h}\right|^2$$

$$R_{n,k,h} = \mathrm{Re}\left(\tilde{\gamma}_{n,k,h}\overline{\tilde{\delta}}_{n,k,h}\right)$$

$$\left\langle \hat{T}_{00}^{(1/2)} \right\rangle_{\mathrm{vac}} = -\frac{1}{\pi R a^3}\sum_{n\in\mathbb{Z}}\int \frac{d^4\boldsymbol{k}}{(2\pi)^3}\omega_{n,k}$$

$$= \frac{1}{L^5 b}\left[-\frac{\left(Lm_{1/2}\right)^5}{30\pi^2} + \frac{1}{\pi^2}\sum_{n=1}^{\infty}\cos\left(nqg_0\vartheta L\right)e^{-x_{n,1/2}}\frac{3 + 3x_{n,1/2} + x_{n,1/2}^2}{n^5}\right]$$

$$\left\langle \hat{T}_{ii}^{(1/2)} \right\rangle = \frac{1}{6}\sum_{j=1}^{4}\left\langle \bar{\Psi}_j\overleftrightarrow{\mathcal{D}}_j\Psi \right\rangle$$

$$= \frac{1}{3(2\pi R)a^2}\sum_{n\in\mathbb{Z},h=\pm}\int \frac{d^4\boldsymbol{k}}{(2\pi)^3}\left[-\frac{k^2}{a\omega_{n,k}}\left(1 - 2N_{n,k,h}^{(1/2)}\right) + \frac{2hk\mu_n}{\omega_{n,k}}R_{n,k,h}^{(1/2)}\right]$$

$$\frac{1}{a^2}\left\langle \hat{T}_{ii}^{(1/2)} \right\rangle_{\mathrm{vac}}$$

$$= -\frac{1}{3(2\pi R)a^3}\sum_{n\in\mathbb{Z}}\int \frac{d^4\boldsymbol{k}}{(2\pi)^3}\frac{k^2}{a^2\omega_{n,k}}$$

$$= \frac{1}{L^5 b}\left[\frac{\left(Lm_{1/2}\right)^5}{30\pi^2} - \frac{1}{\pi^2}\sum_{n=1}^{\infty}\cos\left(nqg_0\vartheta L\right)e^{-x_{n,1/2}}\frac{3 + 3x_{n,1/2} + x_{n,1/2}^2}{n^5}\right]$$

$$\left\langle \hat{T}_{55}^{(1/2)} \right\rangle = \frac{b}{2}\left\langle \hat{\bar{\Psi}}^5\overleftrightarrow{\mathcal{D}}_5\hat{\Psi} \right\rangle$$

$$= \frac{b^3}{2\pi R a^3}\sum_{n\in\mathbb{Z},h=\pm}\int \frac{d^4\boldsymbol{k}}{(2\pi)^3}\left[-\frac{M_{KK,n}^2}{\omega_{n,k}}\left(1 - 2N_{n,k,h}^{(1/2)}\right) + \frac{2M_{KK,n}^2 kh}{a\omega_{n,k}\mu_n}R_{n,k,h}^{(1/2)}\right.$$

$$\left. - \frac{2\left(\frac{n}{R} - qg_0 b\vartheta\right)m_{1/2}}{\mu_n b^2}I_{n,k,h}^{(1/2)}\right]$$

$$\frac{1}{b^3}\left\langle \hat{T}_{55}^{(1/2)} \right\rangle_{\text{vac}}$$
$$= -\frac{1}{2\pi R a^3}\sum_{n\in\mathbb{Z}} \int \frac{d^4\boldsymbol{k}}{(2\pi)^3}\left(n - \frac{qg_0\vartheta L}{2\pi}\right)^2 \frac{2}{R^2 b^3 \omega_{n,k}}$$
$$= \frac{1}{L^5 b}\left[\frac{\left(Lm_{1/2}\right)^5}{30\pi^2} + \frac{1}{\pi^2}\sum_{n=1}^{\infty} \cos\left(nqg_0 b\vartheta L\right) e^{-x_{n,1/2}} \frac{12 + 12x_{n,1/2} + 5x_{n,1/2}^2 + x_{n,1/2}^3}{n^5}\right]$$

$$T_{00}^{(1)} = F_{0M}g^{MN}F_{0N} - \frac{1}{4}g_{00}F_{MN}F^{MN} = \frac{1}{2b^2}\left(\dot{A}_5\right)^2 = \frac{1}{4\pi R}\left(\dot{\vartheta} + H_b\vartheta\right)^2$$
$$T_{ii}^{(1)} = -\frac{a^2}{2b^2}\left(\dot{A}_5\right)^2 = -\frac{1}{4\pi R}\left(\dot{\vartheta} + H_b\vartheta\right)^2$$
$$T_{55}^{(1)} = \frac{b}{2}\left(\dot{A}_5\right)^2 = \frac{b^3}{4\pi R}\left(\dot{\vartheta} + H_b\vartheta\right)^2$$

$$3H_a^2 - \frac{3}{4}H_b^2 = \frac{2\pi R}{M_{\text{pl}}^2}\rho$$
$$-3H_a^2 - 2\dot{H}_a - \frac{3}{4}H_b^2 = \frac{2\pi R}{M_{\text{pl}}^2}p$$
$$-\dot{H}_a + \frac{1}{2}\dot{H}_b - 2H_a^2 + \frac{3}{2}H_aH_b - \frac{1}{4}H_b^2 = \frac{2\pi R}{3M_{\text{pl}}^2}p_y$$

$$2\pi R M_5^3 = M_{\text{pl}}^2 \text{ and } \rho = \langle T_{00}\rangle/a^2, p = \langle T_{ii}\rangle/a^2$$

$$p_y = \langle T_{55}\rangle/b^3$$

$$2\pi R\rho = V_{\text{loop}} + \sum_A \left[\frac{3(\dot{H}_a + 3H_a^2)}{2L^3}\left\{\frac{\left(LM_{0,A}\right)^3}{48\pi^2} + \frac{1}{4\pi^2}\sum_{n=1}^{\infty}\cos\left(nqg_0\vartheta L\right)e^{-x_{n,0,A}}\frac{1 + x_{n,0,A}}{n^3}\right\}\right.$$
$$\left. + \sum_{n\in\mathbb{Z}}\int \frac{d^4\boldsymbol{k}}{(2\pi)^3 a^3}\left\{2\left(\omega_{0,n,k,A} + \frac{3}{4}\frac{\dot{H}_a + 3H_a^2}{\omega_{0,n,k,A}}\right)N_{n,k,A}^{(0)} + \frac{3H_a}{2}I_{n,k,A}^{(0)} + \frac{3}{2}\frac{\dot{H}_a + 3H_a^2}{\omega_{0,n,k,A}}R_{n,k,A}^{(0)}\right\}\right]$$
$$+ \sum_B \sum_{n\in\mathbb{Z},h=\pm}\int \frac{d^4\boldsymbol{k}}{(2\pi)^3 a^3}\left\{2\omega_{n,k,B}N_{n,k,h,B}^{(1/2)} + 2\mu_{n,B}\left(\frac{\omega_{n,k,B} - \frac{kh}{a}}{\omega_{n,k,B}}\right)R_{n,k,h,B}^{(1/2)}\right\}$$

$$2\pi R p = -V_{\text{loop}} + \sum_A \sum_{n\in\mathbb{Z}}\int \frac{d^4\boldsymbol{k}}{(2\pi)^3 a^3}\left\{\frac{2}{\omega_{0,n,k,A}}\left(\frac{k^2}{3a^2} - \frac{3}{4}\dot{H}_a\right)N_{n,k,A}^{(0)} - 2\left(\omega_{0,n,k,A} - \frac{\frac{k^2}{3a^2} - \frac{3}{4}\dot{H}_a}{\omega_{0,n,k,A}}\right)R_{n,k,A}^{(0)}\right.$$
$$\left. + \frac{3}{2}H_aI_{n,k,A}^{(0)}\right\} + a\sum_B \sum_{n\in\mathbb{Z},h=\pm}\int \frac{d^4\boldsymbol{k}}{(2\pi)^3 a^3}\left\{\frac{2k^2}{3a^2\omega_{n,k,B}}N_{n,k,h,B}^{(1/2)} + \frac{2hk\mu_{n,B}}{3a\omega_{n,k,B}}R_{n,k,h,B}^{(1/2)}\right\}.$$

$$2\pi R p_y = \sum_A \left[-\frac{1}{L^5 b}\left\{ \frac{\left(M_{0,A}L\right)^5}{60\pi^2} + \frac{1}{2\pi^2}\sum_{n=1}^{\infty} \cos\left(nq_A g_0 \vartheta L\right) e^{-x_{n,0,A}} \frac{12 + 12x_{n,0,A} + 5x_{n,0,A}^2 + x_{n,0,A}^3}{n^5} \right\} \right.$$
$$- \frac{3\dot{H}_a}{2L^3}\left\{ \frac{\left(LM_{0,A}\right)^3}{48\pi^2} + \frac{1}{4\pi^2}\sum_{n=1}^{\infty} \cos\left(nq_A g_0 \vartheta L\right) e^{-x_{n,0,A}} \frac{1 + x_{n,0,A}}{n^3} \right\}$$
$$+ \sum_{n\in\mathbb{Z}} \int \frac{d^4\boldsymbol{k}}{(2\pi)^3 a^3}\left\{ \frac{2}{\omega_{0,n,k,A}}\left(M_{KK,n,A}^2 - \frac{3}{4}\dot{H}_a \right) N_{n,k,A}^{(0)} \right.$$
$$\left. \left. + \frac{1}{\omega_{0,n,k,A}}\left(\frac{k^2}{a^2} + \frac{m_{0,A}^2}{b} + \frac{3}{2}\dot{H}_a - \frac{9}{4}H_a^2 \right) R_{n,k,A}^{(0)} + \frac{3}{2}H_a I_{n,k,A}^{(0)} \right\} \right]$$
$$+ \sum_B \left[\frac{1}{L^5 b}\left\{ \frac{\left(m_{1/2,B}L\right)^5}{30\pi^2} + \frac{1}{\pi^2}\sum_{n=1}^{\infty} \cos\left(nq_B g_0 \vartheta L\theta\right) e^{-x_{n,1/2,B}} \frac{12 + 12x_{n,1/2,B} + 5x_{n,1/2,B}^2 + x_{n,1/2,B}^3}{n^5} \right\} \right.$$
$$+ \sum_{n\in\mathbb{Z},h=\pm} \int \frac{d^4\boldsymbol{k}}{(2\pi)^3 a^3}\left\{ \frac{2M_{KK,n,B}^2}{\omega_{n,k}} N_{n,k,h}^{(1/2)} - \frac{2M_{KK,n,B}^2}{\mu_{n,B}} \frac{kh}{\omega_{n,k,B}} R_{n,k,h,B}^{(1/2)} \right.$$
$$\left. \left. + \frac{\left(\frac{n}{R} - q_B g_0 b\vartheta\right) m_{1/2,B}}{b^2 \mu_{n,B}} I_{n,k,h,B}^{(1/2)} \right\} \right].$$

$$b = \exp\left(\sqrt{\frac{2}{3}} \frac{\varphi}{M_{\mathrm{pl}}} \right)$$

$$\ddot{\varphi} + 3H_a\dot{\varphi} = \sqrt{\frac{2}{3}} \frac{2\pi R}{M_{\mathrm{pl}}}\left(p_y - \frac{3}{2}p + \frac{1}{2}\rho \right).$$

$$\Delta_\varphi \equiv \sqrt{\frac{2}{3}} \frac{2\pi R}{M_{\mathrm{pl}}}\left(p_y - \frac{3}{2}p + \frac{1}{2}\rho \right)$$

$$\Delta_\varphi^{\mathrm{vac}} \equiv \sqrt{\frac{2}{3}} \frac{2\pi R}{M_{\mathrm{pl}}}\left(p_y^{\mathrm{vac}} - \frac{3}{2}p^{\mathrm{vac}} + \frac{1}{2}\rho^{\mathrm{vac}} \right)$$
$$= -\partial_\varphi V_{\mathrm{loop}} + \frac{\sqrt{6}(6H_a^2 + \dot{H}_a)}{M_{\mathrm{pl}} b L^2} \sum_A \left[\frac{\left(LM_{0,A}\right)^3}{48\pi^2} + \frac{1}{8\pi^2}\sum_{n=1}^{\infty} \cos\left(nq_A g_0 \vartheta L\theta\right) e^{-x_{n,0,A}} \frac{1 + x_{n,0,A}}{n^3} \right]$$

$$\Delta_\varphi^{\mathrm{vac}}\big|_{H_a=0} = -\partial_\varphi V_{\mathrm{loop}}$$

$$\Delta_{\varphi}^{\text{real}} = \sqrt{\frac{2}{3}}\frac{1}{a^3 M_{\text{pl}}}\sum_{n\in\mathbb{Z}}\int \frac{d^4\boldsymbol{k}}{(2\pi)^3}\Bigg[\sum_A \Bigg\{\frac{2}{\omega_{0,n,k,A}}\Bigg\{\left(\frac{m_{0,A}^2}{2b}+\frac{9H_a^2+6\dot{H}_a}{4}+\frac{3}{2}M_{KK,n,A}^2\right)N_{n,k,A}^{(0)}$$
$$+\left(\frac{27H_a^2-9\dot{H}_0}{8}+\frac{3}{2}M_{KK,n,A}^2+\frac{m_{0,A}^2}{2b}-\frac{k^2}{a^2}\right)R_{n,k,A}^{(0)}\Bigg\}$$
$$+\sum_B\sum_{h=\pm}\Bigg\{\frac{1}{\omega_{n,k,B}}\left(3M_{KK,n,B}^2+m_{1/2,B}^2 b^{-1}\right)N_{n,k,h,B}^{(1/2)}+\frac{2\left(\frac{n}{R}-q_B g_0 b\vartheta\right)m_{1/2,B}}{b^2\mu_{n,B}}I_{n,k,h,B}^{(1/2)}$$
$$+\left(\mu_{n,B}-\frac{4M_{KK,n,B}^2+2m_{1/2,B}^2 b^{-1}}{\mu_{n,B}}\frac{kh}{a\omega_{n,k,B}}\right)R_{n,k,h,B}^{(1/2)}\Bigg\}\Bigg].$$

$$\ddot{\vartheta}+3H_a\dot{\vartheta}-\left(3H_aH_b+\dot{H}_b+H_b^2\right)\vartheta+\left(J_0+J_{1/2}\right)=0$$

$$J_0 \equiv \frac{1}{a^3}\sum_A\sum_{n\in\mathbb{Z}}\partial_\vartheta M_{0,n,A}^2\langle\hat{\phi}_n^A\hat{\phi}_n^{A\dagger}\rangle$$
$$J_{1/2} \equiv -\frac{1}{a^3}\sum_B\sum_{n\in\mathbb{Z}}\left(\partial_\vartheta M_{1/2,n}\langle\hat{\zeta}_n^B\hat{\chi}_n^B\rangle+\text{ h.c. }\right)$$

$$J_0 \to \sum_A\Bigg[\frac{1}{L^3b^2}\sum_{n=1}^{\infty}\frac{q_A g_0}{2\pi^2}\sin\left(nq_A g_0\vartheta L\right)e^{-x_{n,0,A}}\frac{3+3x_{n,0,A}+x_{n,0,A}^2}{n^4}$$
$$-\frac{2q_A g_0}{a^3}\sum_{n\in\mathbb{Z}}\int\frac{d^4\boldsymbol{k}}{(2\pi)^3}\frac{\left(\frac{n}{R}-q_A g_0 b\vartheta\right)}{b^2\omega_{n,k,A}}\left(N_{n,k,A}^{(0)}+R_{n,k,A}^{(0)}\right)\Bigg],$$
$$J_{1/2} \to \sum_B\Bigg[-\frac{1}{L^3b^2}\sum_{n=1}^{\infty}\frac{q_B g_0}{\pi^2}\sin\left(nq_B g_0\vartheta L\right)e^{-x_{n,1/2,B}}\frac{3+3x_{n,1/2,B}+x_{n,1/2,B}^2}{n^4}$$
$$-\frac{2q_B g_0}{a^3}\sum_{n\in\mathbb{Z},h=\pm}\int\frac{d^4\boldsymbol{k}}{(2\pi)^3}\Bigg\{\frac{\left(\frac{n}{R}-q_B g_0 b\vartheta\right)}{b^2\omega_{n,k,B}}N_{n,k,B}^{(1/2)}+\frac{m_{1/2,B}}{\mu_{n,B}b}I_{n,k,B}^{(1/2)}-\frac{\left(\frac{n}{R}-q_B g_0\vartheta b\right)}{b^2\omega_{n,k,B}}\frac{kh}{a\mu_{n,B}}R_{n,k,B}^{(1/2)}\Bigg\}\Bigg].$$

$$\partial_M\left(\sqrt{-G}G^{MN}\partial_N\chi\right)=0$$

$$T_{00}^{\chi}=\frac{v^2}{2b^3}=-\frac{1}{a^2}T_{ii}^{\chi}=\frac{1}{b^3}T_{55}^{\chi}$$

$$R=40\times C^{-1}, g_0=0.02C,$$
$$2\pi R\Lambda^5=1.1182\times10^{-3}C^4,$$
$$\left(N_A,m_{0,A},q_A\right)=(28,1.8\times10^{-1}C,0),(2,5\times10^{-4}C,1)\,(2,8.403\times10^{-4}C,5)$$
$$\left(N_B,m_{1/2,B},q_B\right)=(16,1.9\times10^{-1}C,0),(1,5.2\times10^{-4}C,1),(1,8.4\times10^{-4}C,5),$$

$$L_{\text{phys}}=\int\sqrt{G_{55}}dy=2\pi Rb$$

$$\ddot{R}_{n,k,A}^{(0)}\sim-4\omega_{0,n,k,A}^2R_{n,k,A}^{(0)}$$

$$R_{n,k,A}^{(0)},I_{n,k,A}^{(0)}\sim e^{\pm2i\omega_{0,n,k,A}t}$$

$$\langle[\mathcal{O}_{r_1}\mathcal{O}_{r_2}](z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\rangle$$

$$\langle \mathcal{O}_{r_1}(z_0)\mathcal{O}_{r_2}(z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\rangle$$

$$\langle [\mathcal{O}_{r_1}\mathcal{O}_{r_2}]\mathcal{O}_s\mathcal{O}_p\mathcal{O}_q\rangle = \underline{O\left(\frac{1}{N^1}\right)} \text{ disconnected } + \underline{O\left(\frac{1}{N^3}\right)+\cdots} \text{ connected}$$

$$\langle \mathcal{O}_r^2\mathcal{O}_2\mathcal{O}_p\mathcal{O}_p\rangle\,;\ \langle [\mathcal{O}_2\mathcal{O}_4]\mathcal{O}_2\mathcal{O}_p\mathcal{O}_p\rangle\,;\ \langle \mathcal{O}_2^2\mathcal{O}_s\mathcal{O}_p\mathcal{O}_{p+s-2}\rangle$$

$$\mathcal{M}_{[2^2],p_2,p_3,p_4}\ \forall p_2,p_3,p_4\ ;\ \ \mathcal{M}_{[r^2],2,p,p}\ \forall r,p$$

$$\phi(z) = Y^I\phi_I(\vec{X})\ ;\ z \equiv (\vec{X},\vec{Y})$$

$$T_p(z) := \mathrm{Tr}(\phi(z)^p)$$

$$T_{p_1,p_2,\ldots,p_n}(z) = T_{p_1}(z)T_{p_2}(z)\ldots T_{p_n}(z)\,,\Delta = \sum_i p_i$$

$$\mathcal{O}_p = T_p + \sum_{\underline{q}\vdash p} c_{\underline{q}}(p,N)T_{\underline{q}},$$

$$\langle \phi_b^a(z_1)\phi_m^n(z_2)\rangle = \left(\delta_m^a\delta_m^b - \frac{1}{N}\delta_b^a\delta_m^n\right)g_{12}\ ;\ g_{12} = \frac{\vec{Y}_{12}^2}{\vec{X}_{12}^2}\ ;\ \vec{Y}_{12}^2 \equiv \vec{Y}_1\cdot\vec{Y}_2$$

$$\langle \mathcal{O}_\Delta(z_1)\mathcal{O}_\Delta(z_2)\rangle = g_{12}^\Delta f_{O,\Delta}(N)\ ;\ |\mathcal{O}_\Delta|^2 \equiv f_{O,\Delta}(N)$$

$$\langle \mathcal{O}_{p_1}(z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\rangle = \mathcal{P}_{\vec{p}}[g_{ij}]\mathcal{C}_{\vec{p}}(U,V,\hat{\sigma},\hat{\tau})$$

$$U = \frac{\vec{X}_{12}^2\vec{X}_{34}^2}{\vec{X}_{13}^2\vec{X}_{24}^2}\ ;\ V = \frac{\vec{X}_{14}^2\vec{X}_{23}^2}{\vec{X}_{13}^2\vec{X}_{24}^2}\ ;\ \hat{\sigma} = \frac{\vec{Y}_{13}^2\vec{Y}_{24}^2}{\vec{Y}_{12}^2\vec{X}_{34}^2}\ ;\ \hat{\tau} = \frac{\vec{Y}_{14}^2\vec{Y}_{23}^2}{\vec{Y}_{12}^2\vec{Y}_{34}^2}$$

$$\mathcal{C}_{\vec{p}}(U,V,\hat{\sigma},\hat{\tau}) = \mathcal{G}_{\vec{p}}^{\text{free}}(U,V,\hat{\sigma},\hat{\tau}) + \mathcal{I}(U,V,\hat{\sigma},\hat{\tau})\mathcal{H}_{\vec{p}}(U,V,\hat{\sigma},\hat{\tau})$$

$$\mathcal{I}(U,V,\hat{\sigma},\hat{\tau}) = V + \hat{\sigma}^2UV + \hat{\tau}^2U + \hat{\sigma}V(-1-U+V) + \hat{\tau}(1-U-V) + \hat{\sigma}\hat{\tau}U(-1+U-V).$$

$$\kappa = -2 - \Sigma + \sum_{ij=12,13,14}\min\left(\frac{p_i+p_j}{2}, \Sigma - \frac{p_i+p_j}{2}\right),$$

$$\Sigma = \frac{p_1+p_2+p_3+p_4}{2}\mathcal{I}(U,V,\hat{\sigma},\hat{\tau})$$

$$U = x_1x_2, V = (1-x_1)(1-x_2)\ ;\ \hat{\sigma} = \frac{1}{y_1y_2}, \hat{\tau} = \frac{(1-y_1)(1-y_2)}{y_1y_2}$$

$$\mathcal{I}(x_1,x_2,y_1,y_2) = \frac{1}{(y_1y_2)^2}\prod_{i=1,2}(x_i-y_j)$$

$$\left[\left(\partial_{x_i}+\partial_{y_j}\right)\mathcal{C}\right]_{x_i=y_j} = 0$$

$$\mathcal{D}_{pq,\tau,l,[aba]} = \mathcal{O}_p\partial^l\,\square^{\frac{1}{2}(\tau-p-q)}\,\mathcal{O}_q\ (2\le p\le q)$$

$$R_{\vec{\tau}}:=\left\{(p,q):\begin{matrix}p=i+a+2+r\\ q=i+a+2+b-r\end{matrix}\text{ for }\begin{matrix}i=0,\dots,(t-2)\\ r=0,\dots,(\mu-1)\end{matrix}\right\}$$

$$t\equiv\frac{(\tau-b)}{2}-a,\mu\equiv\begin{cases}\left\lfloor\frac{b+2}{2}\right\rfloor & a+l\text{ even}\\ \left\lfloor\frac{b+1}{2}\right\rfloor & a+l\text{ odd}\end{cases}$$

$$\mathcal{D}_{pq,[aba]}=\mathcal{O}_p\partial^l\mathcal{O}_q, 2a+b+2=p+q\ (2\le p\le q)$$


$$\bigoplus_{\tau=4,6,\dots}R_{\tau,l,[000]}=\{\{\underbrace{\mathcal{O}_2\partial^l\mathcal{O}_2}_{\tau=4}\},\{\underbrace{\mathcal{O}_2\partial^l\,\square\,\mathcal{O}_2,\mathcal{O}_3\partial^l\mathcal{O}_3}_{\tau=6}\},\dots\}$$


$$\bigoplus_{\tau=5,7,\dots}R_{\tau,l,[010]}=\{\{\underbrace{\mathcal{O}_2\partial^l\mathcal{O}_3}_{\tau=5}\},\{\underbrace{\mathcal{O}_2\partial^l\,\square\,\mathcal{O}_3,\mathcal{O}_3\partial^l\mathcal{O}_4}_{\tau=7}\},\dots\}$$


$$\bigoplus_{\tau=6,8,\dots}R_{\tau,l,[101]}=\{\{\underbrace{\mathcal{O}_3\partial^l\mathcal{O}_3}_{\tau=6}\},\{\underbrace{\mathcal{O}_3\partial^l\,\square\,\mathcal{O}_3,\mathcal{O}_4\partial^l\mathcal{O}_4}_{\tau=8}\}\dots\}$$


$$\bigoplus_{\tau=6,8,\dots}R_{\tau,l,[020]}=\{\{\underbrace{\mathcal{O}_2\partial^l\mathcal{O}_4,\mathcal{O}_3\partial^l\mathcal{O}_3}_{\tau=6}\},\{\underbrace{\mathcal{O}_2\partial^l\,\square\,\mathcal{O}_4,\mathcal{O}_3\partial^l\mathcal{O}_5,\mathcal{O}_3\partial^l\,\square\,\mathcal{O}_3,\mathcal{O}_4\partial^l\mathcal{O}_4}_{\tau=8}\},\dots\}$$

$$\mathcal{C}_{\mathcal{O}_r\mathcal{O}_s\mathcal{D}_{\vec{\tau}}}=\mathcal{C}^{(0)}_{\mathcal{O}_r\mathcal{O}_s\mathcal{D}_{\vec{\tau}}}+\frac{1}{N}\mathcal{C}^{(\frac{1}{2})}_{\mathcal{O}_r\mathcal{O}_s\mathcal{D}_{\vec{\tau}}}+\frac{1}{N^2}\mathcal{C}^{(1)}_{\mathcal{O}_r\mathcal{O}_s\mathcal{D}_{\vec{\tau}}}+\cdots,$$









$$\langle[\mathcal{O}_{r_1}\mathcal{O}_{r_2}]\mathcal{O}_s\mathcal{O}_p\mathcal{O}_q\rangle=O\left(\frac{1}{N}\right)\times\text{ free }+O\left(\frac{1}{N^3}\right)\times(\text{ free }+\text{ dynamical })+\cdots.$$

$$\frac{1}{N}\mathcal{C}^{\left(\frac{1}{2}\right)}_{[r_1 r_2]s;\mathcal{D}_{\tau,l}} \times \begin{cases} \frac{1}{N^2}\eta_{\mathcal{D}}\mathcal{C}^{(0)}_{pq;\mathcal{D}}\log(U) & \text{if } p+q \leq \tau < r_1+r_2+s \\ \frac{1}{N^2}\mathcal{C}^{(1)}_{pq;\mathcal{D}} & \text{if } \tau_{\text{unitary}} \leq \tau < p+q \end{cases}$$

$$\langle \mathcal{O}_2^2 \mathcal{O}_{p_2} \mathcal{O}_{p_3} \mathcal{O}_{p_4} \rangle$$

$$\langle \mathcal{O}_3^2 \mathcal{O}_2 \mathcal{O}_p \mathcal{O}_p \rangle, \langle [\mathcal{O}_2 \mathcal{O}_4] \mathcal{O}_2 \mathcal{O}_p \mathcal{O}_p \rangle, \langle \mathcal{O}_r^2 \mathcal{O}_2 \mathcal{O}_p \mathcal{O}_p \rangle$$

$$\langle \mathcal{O}_2^2(z_1)\mathcal{O}_2(z_2)\mathcal{O}_3(z_3)\mathcal{O}_3(z_4)\rangle \; ; \; \langle \mathcal{O}_2^2(z_1)\mathcal{O}_3(z_2)\mathcal{O}_2(z_3)\mathcal{O}_3(z_4)\rangle$$

$$|\mathcal{O}_3|^2 = \frac{3a^2}{N}\left(1-\frac{3}{a}\right) \; ; \; |\mathcal{O}_2|^2 = 2a \; ; \; \left|\mathcal{O}_2^2\right|^2 = 2|\mathcal{O}_2|^4\left(1+\frac{2}{a}\right)$$

$$\mathcal{P}_{[2^2]233} = |\mathcal{O}_3|^2|\mathcal{O}_2|\left|\mathcal{O}_2^2\right| g_{12}^2 g_{14} g_{13} g_{34}^2 \; ; \; \mathcal{P}_{[2^2]323} = |\mathcal{O}_3|^2|\mathcal{O}_2|\left|\mathcal{O}_2^2\right| g_{12}^3 g_{14} g_{34}^2$$

$$\mathcal{G}_{[2^2]233} = \frac{6}{\sqrt{a+2}}\left[1+\frac{2}{a}+\frac{4}{a}\left[U\hat{\sigma}+\frac{U\hat{\tau}}{V}\right]+\frac{2}{a}\frac{U^2\hat{\sigma}\hat{\tau}}{V}\right]$$

$$\mathcal{G}_{[2^2]323} = \frac{6}{\sqrt{a+2}}\left[U^2\hat{\sigma}^2+\frac{2}{a}\left[\frac{U\hat{\tau}}{V}+2U\hat{\sigma}\right]+\frac{2}{a}\left[U^2\hat{\sigma}^2+2\frac{U^2\hat{\sigma}\hat{\tau}}{V}\right]\right].$$

$$\eta_{\mathcal{D}_{5,l}} = -\frac{80}{(l+1)(l+4)}, l = 0,2,4,\ldots, \eta_{\mathcal{D}_{5,l}} = -\frac{80}{(l+4)(l+7)}, l = 1,3,5,\ldots.$$

$$\mathcal{C}_{[2^2]3;\mathcal{D}_{5,l}} \propto O\left(\frac{1}{N}\right) \; ; \; \mathcal{C}_{23;\mathcal{D}_5} \propto O(1)$$

$$\mathcal{H}_{[2^2]323}\Big|_{\frac{1}{N^3}} = \log(U)\sum_l \mathcal{C}^{\left(\frac{1}{2}\right)}_{[2^2]3;\mathcal{D}_{5,l}}\eta_{\mathcal{D}_{5,l}}\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}}H^{4323}_{5,l,[010]} + \cdots$$

$$\mathcal{C}^{\left(\frac{1}{2}\right)}_{[2^2]3;\mathcal{D}_{5,l}}\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}}\Big|_{\frac{1}{N}} = 6M^{4323}_{k=0,\gamma=5,[l+2,2]} = \frac{6}{10}(l+1)(l+7)\frac{(l+3)!\,(l+4)!}{(2l+7)!}$$

$$\mathcal{C}^{\left(\frac{1}{2}\right)}_{[2^2]3;\mathcal{D}_{5,l}}\eta_{\mathcal{D}_{5,l}}\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}}\Big|_{\frac{\log(U)}{N^3}} = -\frac{12\times 4(l+7)(l+3)!^2}{(2l+7)!}\frac{1+(-1)^l}{2} - \frac{12\times 4(l+1)(l+3)!^2}{(2l+7)!}\frac{1-(-1)^l}{2}$$

$$\sum_l \mathcal{C}^{\left(\frac{1}{2}\right)}_{[2^2]3;\mathcal{D}_{5,l}}\eta_{\mathcal{D}_{5,l}}\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}}H^{4323}_{5,l,[010]}\Big|_{\frac{\log(U)}{N^3}} = U^2\left[-\sum_{l\geq 0}\frac{144(l+1)(l+2)}{(l+4)(l+5)(l+6)}x_1^l + \cdots\right]$$

$$-\sum_{l\geq 0}\frac{144(l+1)(l+2)}{(l+4)(l+5)(l+6)}x_1^l = \frac{48(30-21x_1+x_1^2)}{x_1^5} + \frac{144(10-12x_1+3x_1^2)\log(1-x_1)}{x_1^6}.$$

$$\mathcal{C}^{\left(\frac{1}{2}\right)}_{[2^2]3;\mathcal{D}_{5,l}}\eta_{\mathcal{D}_{5,l}}\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}} = \left[\mathcal{C}^{\left(\frac{1}{2}\right)}_{[2^2]3;\mathcal{D}_{5,l}}\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}}\right]_{(3)}\left[\frac{1}{\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}}\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}}}\right]_{(2)}\left[\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}}\eta_{\mathcal{D}_{5,l}}\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}}\right]_{(1)}$$

$$\mathcal{C}^{\left(\frac{1}{2}\right)}_{[2^2]3;\mathcal{D}_{5,l}}\eta_{\mathcal{D}_{5,l}}\mathcal{C}^{(0)}_{23;\mathcal{D}_{5,l}}=\frac{\left(\mathcal{G}^{\left(\frac{1}{2}\right)}_{[2^2]323}\Big|_{[010],\tau=5,l}\right)\times\left(\mathcal{H}^{(1)}_{2323}\Big|_{\log(U),[010],\tau=5,l}\right)}{\left(\mathcal{G}^{(0)}_{2323}\Big|_{[010],\tau=5,l}\right)}$$

$$\mathcal{H}_{[2^2]323}\Big|_{\frac{\log^0(U)}{N^3}}=-\sum_l 12\times\left(2M^{4323}_{k=0,\gamma=3,[2+l]}+M^{4323}_{k=1,\gamma=3,[2+l]}\right)H^{4323}_{3,l,[010]}+\text{ higher twists}$$

$$=-\left[U\sum_l\frac{12(l+1)(l+6)}{(l+3)(l+4)}x_1^l+O(U^2)\right]$$

$$\mathcal{H}_{[2^2]323}\Big|_{\frac{\log^0(U)}{N^3}}=-x_1x_2\left[\frac{12(6-9x_1+2x_1^2)}{(-1+x_1)x_1^3}+\frac{12\times6(-1+x_1)\log(1-x_1)}{x_1^4}\right]+O((x_1x_2)^2)$$

$$\begin{pmatrix}\langle\mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\rangle & \langle\mathcal{O}_2\mathcal{O}_2\mathcal{O}_3\mathcal{O}_3\rangle\\ \langle\mathcal{O}_2^2\mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\rangle & \langle\mathcal{O}_2^2\mathcal{O}_2\mathcal{O}_3\mathcal{O}_3\rangle\end{pmatrix}\Big|_{[020],\tau=4}$$

$$\mathcal{S}_{[2^2]233,[020],\tau=4,l}\Big|_{\frac{1}{N^3}}\equiv C^{\left(\frac{1}{2}\right)}_{[2^2]2,\mathcal{O}_2\partial^l\mathcal{O}_2}C^{(1)}_{33,\mathcal{O}_2\partial^l\mathcal{O}_2}$$

$$\mathcal{S}_{[2^2]233,[020],\tau=4,l}\Big|_{\frac{1}{N^3}}=\frac{\left(\mathcal{G}^{\left(\frac{1}{2}\right)}_{[2^2]222}\Big|_{[0,2,0],\tau=4,l}\right)\times\left(\mathcal{G}^{(1)}_{2233}\Big|_{[0,2,0],\tau=4,l}\right)}{\left(\mathcal{G}^{(0)}_{2222}\Big|_{[0,2,0],\tau=4,l}\right)}$$

$$\mathcal{P}_{[2^2]222}=8a^2\sqrt{a+2}g_{12}^2g_{13}g_{14}g_{34},\mathcal{G}_{[2^2]222}=\frac{4\sqrt{a+2}}{a}\left[1+U\hat{\sigma}+\frac{U\hat{\tau}}{V}\right]$$

$$\mathcal{P}_{22pp}=2a\left|\mathcal{O}_p\right|^{2p}g_{12}^2g_{34}^{2p},\mathcal{G}_{22pp}=1+\delta_{p,2}\left(U\hat{\sigma}^2+\frac{U^2\hat{\tau}^2}{V^2}\right)+\frac{2p}{a}\left(U\hat{\sigma}+\frac{U\hat{\tau}}{V}+(p-1)\frac{U^2\hat{\sigma}\hat{\tau}}{V}\right)$$

$$\mathcal{S}_{[2^2]233,[020],\tau=4,l}\Big|_{\frac{1}{N^3}}=\frac{\left(4M^{4222}_{0,4,[l+2]}+4M^{4222}_{1,4,[l+2]}\right)\times\left(12M^{2233}_{1,4,[l+2]}\right)}{\left(M^{2222}_{0,4,[l+2]}+M^{2222}_{2,4,[l+2]}\right)}=\frac{12\times8(l+2)!\,(l+3)!}{(2l+6)!}\frac{1+(-1)^l}{2}.$$

$$A_{[2^2]233,[020],\tau=4,l}\Big|_{\frac{1}{N^3}}=6\times4\left(M^{4233}_{0,4,[l+2]}+M^{4233}_{1,4,[l+2]}\right)=\frac{12\times4(l+3)!\,(l+4)!}{(2l+6)!}\frac{1+(-1)^l}{2}$$

$$\mathcal{H}_{[2^2]233}\Big|_{\frac{\log^0(U)}{N^3}}=\sum_l\left(-A_{[2^2]233}+\mathcal{S}_{[2^2]233}\right)\Big|_{\frac{1}{N^3}}H_{[020],4,l}+\cdots=-U\left[\sum_l\frac{24(l+1)}{(l+3)}x_1^l+O(U^2)\right]$$

$$\mathcal{H}_{[2^2]233}\Big|_{\frac{\log^0(U)}{N^3}}=-x_1x_2\left[\frac{24(2-x_1)}{(1-x_1)x_1^2}+\frac{48\log(1-x_1)}{x_1^3}\right]+O((x_1x_2)^2)$$

$$\mathcal{H}^{ansatz}_{[2^2]323}=\frac{N_2\mathcal{P}^{(1)}(x_1,x_2)}{(x_1-x_2)^{d-1}}+\frac{N_{1,U}\log(U)+N_{1,V}\log(V)}{V(x_1-x_2)^{d-2}}+\frac{N_0}{V(x_1-x_2)^{d-2}}$$

$$\frac{U^2}{V^2}\mathcal{H}^{ansatz}_{[2^2]323}(1-x_1,1-x_2)=\mathcal{H}^{ansatz}_{[2^2]323}(x_1,x_2)$$

$$\mathcal{H}_{[2^2]233}\Big|_{\frac{1}{N^3}} = U^2(-36-12U\partial_U)\circ \bar{D}_{2422}$$

$$\mathcal{H}_{[2^2]233}\Big|_{\frac{1}{N^3}} = 24\oiint_{-i\infty}^{+\infty}\frac{dsdt}{(2\pi i)^2}U^{s+2}V^t\Gamma[-s]^2\Gamma[-t]^2\Gamma[-u]^2\frac{\frac{1}{2}(3+s)}{(1+s)(1+t)(1+u)},$$

$$\mathcal{P}_{[2^2]2pp} = \left|\mathcal{O}_p\right|^2|\mathcal{O}_2|\left|\mathcal{O}_2^2\right|g_{12}^2g_{14}g_{13}g_{34}^{p-1}\,;\,\mathcal{P}_{[2^2]p2p} = \left|\mathcal{O}_p\right|^2|\mathcal{O}_2|\left|\mathcal{O}_2^2\right|g_{12}^3g_{14}g_{24}^{p-3}g_{34}^2,$$

$$\left|\mathcal{O}_p\right|^2|\mathcal{O}_2|\left|\mathcal{O}_2^2\right| = 4pN^{p+3}(1+\cdots)$$

$$\mathcal{G}_{[2^2]2pp} = \frac{2p}{\sqrt{a+2}}\left[1+\frac{2}{a}+\frac{2}{a}\left[(p-1)U\hat{\sigma}+(p-1)\frac{U\hat{\tau}}{V}+\frac{(p-2)(p-1)}{2}\frac{U^2\hat{\sigma}\hat{\tau}}{V}\right]\right],$$

$$\mathcal{G}_{[2^2]p2p} = \frac{2p}{\sqrt{a+2}}\left[U^2\hat{\sigma}^2+\frac{2}{a}\left[\frac{(p-2)(p-1)}{2}\frac{U\hat{\tau}}{V}+(p-1)U\hat{\sigma}+U^2\hat{\sigma}^2+(p-1)\frac{U^2\hat{\sigma}\hat{\tau}}{V}\right]\right]$$


$$\mathcal{C}_{[2^2]p;\mathcal{T}_{p+2}}\frac{\eta}{N^2}\mathcal{C}_{2p\mathcal{T}_{p+2}}\log^1(U)\,;\,\mathcal{C}_{[2^2]p;\mathcal{T}_p}\mathcal{C}_{2p;\mathcal{T}_p}\log^0(U)$$

$$\mathcal{C}_{2p;\mathcal{T}_{p+2}} \propto O\left(\frac{1}{N}\right)\,;\,\mathcal{C}_{[2^2]p\mathcal{T}_{p+2}} \propto O\left(\frac{1}{N^2}\right)$$

$$\mathcal{C}_{[2^2]p;\mathcal{T}_{p+2}}\frac{\eta}{N^2}\mathcal{C}_{2p\mathcal{T}_{p+2}}\log^1(U) \propto O\left(\frac{1}{N^5}\right)$$

$$\mathcal{C}_{[2^2]p\mathcal{T}_p} \propto O\left(\frac{1}{N^2}\right)\,;\,\mathcal{C}_{2p;\mathcal{T}_p} \propto O\left(\frac{1}{N^3}\right)$$

$$\sum_{R_{p+2,[0,p-2,0]}}\mathcal{C}^{\left(\frac{1}{2}\right)}_{[2^2]p;\mathcal{D}}\eta_{\mathcal{D}}\mathcal{C}^{(0)}_{2p;\mathcal{D}} = \sum_{ij}\frac{\left(\mathcal{G}^{\left(\frac{1}{2}\right)}_{[2^2]pij}\Big|_{[0,p-2,0],\tau=p+2,l}\right)\times\left(\mathcal{H}^{(1)}_{ij2p}\Big|_{\log(U),[0,p-2,0],\tau=p+2,l}\right)}{\left(\mathcal{G}^{(0)}_{ijij}\Big|_{[0,p-2,0],\tau=p+2,l}\right)},$$



$$\sum_{R_{p+2,[0,p-2,0]}}\mathcal{C}^{\left(\frac{1}{2}\right)}_{[2^2]p;\mathcal{D}}\eta_{\mathcal{D}}\mathcal{C}^{(0)}_{2p;\mathcal{D}} = \frac{\left(\mathcal{G}^{\left(\frac{1}{2}\right)}_{[2^2]p2p}\Big|_{[0,p-2,0],\tau=p+2,l}\right)\times\left(\mathcal{H}^{(1)}_{2p2p}\Big|_{\log(U),[0,p-2,0],\tau=p+2,l}\right)}{\left(\mathcal{G}^{(0)}_{2p2p}\Big|_{[0,p-2,0],\tau=p+2,l}\right)}$$

$$\mathcal{G}^{(0)}_{2p2p}\Big|_{[0,p-2,0],\tau=p+2,l} = M^{2p2p}_{0,p+2,[2+l,2]} = \frac{p^2(p-1)}{(p+2)!}\frac{(l+1)(l+p+1)!\,(l+p+4)!}{(l+p+2)(l+p+3)(2l+p+4)!}.$$

$$\mathcal{H}^{(1)}_{2p2p}\Big|_{\log(U),[0,p-2,0],\tau=p+2,l} = \frac{(l+3)!\,(l+p+1)!}{(2l+p+4)!}\begin{cases}(-1)^p c_p(-l-p-5) - 4(p-1)p^2\dfrac{1+(-1)^p}{2} & l \text{ even}\\ c_p(l) & l \text{ odd}\end{cases}$$

$$\begin{aligned} c_3(l) &= -12(l+1)\\ c_4(l) &= -8(l+1)(l+8)\\ c_5(l) &= -\frac{10}{3}(l+1)(60+14l+l^2)\\ c_6(l) &= -(l+1)(l+10)(48+11l+l^2)\\ &\vdots \end{aligned}$$

$$\mathcal{G}^{\left(\frac{1}{2}\right)}_{[2^2]p2p}\Big|_{[0,p-2,0],\tau=p+2,l} = (2p)\times M^{4,p,2,p}_{0,p+2,[2+l,2]} = \frac{2p^2(p-1)^2(p-2)}{(p+2)!}\frac{(l+1)(l+p+4)(l+p+1)!^2}{(l+p+1)(2l+p+4)!}$$

$$2|\mathcal{O}_2|^2 2p|\mathcal{O}_p|^2 \simeq \left(|\mathcal{O}_p|^2|\mathcal{O}_2|\mathcal{O}_2^2|\right)\left(\frac{2p}{N}\right)$$

$$\mathcal{H}_{2^2p2p}\Big|_{\frac{\log(U)}{N^3}} = -4p\times(p-2)(p-1)^2 p\left[U^2\sum_l \frac{(l+1)(l+2)}{(l+p+1)_3}x_1^l + \cdots\right]$$

$$\Phi(z,s,a) = \sum_{k=0}^{\infty}\frac{z^k}{(k+a)^s}$$

$$\sum_l \frac{(l+1)(l+2)}{(l+p+1)_3}x_1^l = \frac{p-1}{2x_1} - \frac{p+2}{2x_1^2} + \left[\frac{(p+1)(p+2)}{2x_1^2} - \frac{(p+1)p}{x_1} + \frac{(p-1)p}{2}\right]\Phi(x_1,1,p+1)$$

$$\Phi(x_1,1,p+1) = \frac{1}{x_1^p}\left(-\frac{\log(1-x_1)}{x_1} - \sum_{n=0}^{p-1}\frac{x_1^n}{n+1}\right)$$

$$\begin{pmatrix}\langle\mathcal{O}_i\mathcal{O}_j\mathcal{O}_a\mathcal{O}_b\rangle & V_{2p}^T\\ V_{[2^2]p} & \langle\mathcal{O}_2^2\mathcal{O}_p\mathcal{O}_2\mathcal{O}_p\rangle\end{pmatrix},$$

$$V_{2p} = \left(\langle\mathcal{O}_2\mathcal{O}_{p-2}\mathcal{O}_2\mathcal{O}_p\rangle, \langle\mathcal{O}_3\mathcal{O}_{p-3}\mathcal{O}_2\mathcal{O}_p\rangle, \dots\right),$$

$$V_{[2^2]p} = \left(\langle\mathcal{O}_2^2\mathcal{O}_p\mathcal{O}_2\mathcal{O}_{p-2}\rangle, \langle\mathcal{O}_2^2\mathcal{O}_p\mathcal{O}_3\mathcal{O}_{p-3}\rangle, \dots\right)$$

$$\mathcal{H}_{[2^2]p2p}\Big|_{\frac{\log^0(U)}{N^3}} = 2p\times(p-2)(p-1)\left[-U\sum_{l=0}^{\infty}\frac{(l+1)(l+2p)}{(l+p)(l+p+1)}x_1^l + \cdots\right],$$

$$\sum_{l=0}^{\infty}\frac{(l+1)(l+2p)}{(l+p)(l+p+1)}x_1^l = -\frac{(p-1)p(x_1-1)}{x_1}\Phi(x_1,1,p) - \frac{1+p(x_1-1)}{(x_1-1)x_1}.$$

$$\mathcal{S}_{[2^2]2pp,[020],\tau=4,l}\Big|_{\frac{1}{N^3}}=\frac{\left(\mathcal{G}^{\left(\frac{1}{2}\right)}_{[2^2]222}\Big|_{[0,2,0],\tau=4,l}\right)\times\left(\mathcal{G}^{(1)}_{22pp}\Big|_{[0,2,0],\tau=4,l}\right)}{\left(\mathcal{G}^{(0)}_{2222}\Big|_{[0,2,0],\tau=4,l}\right)}.$$

$$\mathcal{S}_{[2^2]2pp,[020],\tau=4,l}\Big|_{\frac{1}{N^3}}=\frac{4\left[M^{4222}_{0,4,[l+2]}+M^{4222}_{1,4,[l+2]}\right]\times\left[2p(p-1)M^{22pp}_{1,4,[l+2]}\right]}{\left(M^{2222}_{0,4,[l+2]}+M^{2222}_{2,4,[l+2]}\right)}=\frac{16p(p-1)(l+2)!\,(l+3)!}{(2l+6)!}\frac{1+(-1)^l}{2}.$$

$$A_{[2^2]2pp,[020],4,l}\Big|_{\frac{1}{N^3}}\equiv 4p(p-1)\left(M^{42pp}_{0,4,[l+2]}+M^{42pp}_{1,4,[l+2]}\right)=\frac{4p\times 2(p-1)(l+3)!\,(l+4)!}{(2l+6)!}\frac{1+(-1)^l}{2}$$

$$\mathcal{H}_{[2^2]2pp}\Big|_{\frac{\log^0(U)}{N^3}}=\sum_l\left(-A_{[2^2]2pp}+\mathcal{S}_{[2^2]2pp}\right)H_{[020],4,l}+\cdots.$$

$$\mathcal{H}_{[2^2]2pp}\Big|_{\frac{\log^0(U)}{N^3}}=-x_2\left[\frac{4p(p-1)(2-x_1)}{(1-x_1)x_1}+\frac{8p(p-1)\log(1-x_1)}{x_1^2}+\cdots\right]$$

$$\mathcal{H}^{\text{ansatz}}_{[2^2]p2p}=\frac{N_2\mathcal{P}^{(1)}(x_1,x_2)}{(x_1-x_2)^{d-1}}+\frac{N_{1,U}\log(U)+N_{1,V}\log(V)}{V(x_1-x_2)^{d-2}}+\frac{N_0}{V(x_1-x_2)^{d-2}}$$

$$\frac{U^2}{V^2}\mathcal{H}^{ansatz}_{[2^2]p2p}(1-x_1,1-x_2)=\mathcal{H}^{ansatz}_{[2^2]p2p}(x_1,x_2)$$

$$\mathcal{H}_{[2^2]2pp}\Big|_{\frac{1}{N^3}}=4p\frac{(p-1)}{(p-3)!}\iiint_{-i\infty}^{+\infty}\frac{dsdt}{(2\pi i)^2}U^{s+2}V^t\frac{\Gamma[-s+p-3]^2\Gamma[-t]^2\Gamma[-u]^2}{(-s)_{p-3}}\frac{\frac{1}{2}(3+s)}{(1+s)(1+t)(1+u)},$$

$$\mathcal{H}_{[2^2]2pp}\Big|_{\frac{1}{N^3}}=2p\frac{(p-1)}{(p-3)!}U^{p-1}\partial_U^{p-3}(-3-U\partial_U)\bar{D}_{2422}$$

$$\sum_{\mathcal{T}}C^{(0)}_{[2^2]2\mathcal{T}}\times C^{\left(\frac{3}{2}\right)}_{pp\mathcal{T}}+\sum_{\mathcal{D}}C^{\left(\frac{1}{2}\right)}_{[2^2]2\mathcal{D}}\times C^{(1)}_{pp\mathcal{D}}$$

$$\sum_{\mathcal{T}}C^{(0)}_{[2^2]2\mathcal{T}}\eta_{\mathcal{T}}C^{\left(\frac{1}{2}\right)}_{pp\mathcal{T}}+\sum_{\mathcal{D}}C^{\left(\frac{1}{2}\right)}_{[2^2]2\mathcal{D}}\eta_{\mathcal{D}}C^{(0)}_{pp\mathcal{D}}$$

$$\mathcal{H}_{[2^2]sp(p+s-2)}\,;\,\mathcal{H}_{[3^2]2pp},\mathcal{H}_{[24]2pp}\,;\,\mathcal{H}_{[r^2]2pp}$$

$$\mathcal{P}_{[2^2](p+1)3p}=\sqrt{24p(p+1)}N^{4+p}g_{12}^4g_{24}^{p-3}g_{34}^3$$

$$\mathcal{G}^{\left(\frac{3}{2}\right)}_{[2^2](p+1)3p}=\sqrt{24p(p+1)}\left[U^2\hat{\sigma}^2+2(p-1)\frac{U^2\hat{\sigma}\hat{\tau}}{V}+\frac{(p-1)(p-2)}{2}\frac{U^2\hat{\tau}^2}{V^2}+\frac{2U^3\hat{\sigma}^2\hat{\tau}}{V}+(p-1)\frac{\hat{\sigma}\hat{\tau}^2}{V^2}\right].$$


$$\mathcal{H}_{[2^2]3p(p+1)}(U,V,\hat{\sigma},\hat{\tau})=U\hat{\sigma}\mathcal{H}_{[2^2]p3(p+1)}\left(\frac{1}{U},\frac{V}{U},\frac{1}{\hat{\sigma}},\frac{\hat{\tau}}{\hat{\sigma}}\right)$$
$$\mathcal{H}_{[2^2](p+1)3p}(U,V,\hat{\sigma},\hat{\tau})=\frac{U^3}{V^3}\hat{\tau}\mathcal{H}_{[2^2]p3(p+1)}\left(V,U,\frac{\hat{\sigma}}{\hat{\tau}},\frac{1}{\hat{\tau}}\right)$$


$$\mathcal{P}_{[2^2]p3p+1} = \sqrt{24p(p+1)}N^{p+4}g_{12}^3 g_{14} g_{24}^{p-3} g_{34}^3$$

$$\mathcal{H}_{[2^2]p3(p+1)}\Big|_{\frac{1}{N^3}} = \sqrt{24p(p+1)} \iiint \frac{dsdt}{(2\pi i)^2} U^{s+3} V^t \frac{\Gamma[-s]^2\Gamma[-t]^2\Gamma[-u+p-3]^2}{(-u)_{p-3}} \frac{1}{(1+s)(1+t)(1+u)}$$
$$\times \left[\frac{1}{2(p-3)!}\frac{(p-1)t+p(s+2)}{(2+s)} + \frac{\hat{\sigma}}{2(p-2)!}\frac{2t+p(u+2)}{(2+u)} - \frac{\hat{\tau}}{2(p-3)!}\right]$$

$$\mathcal{H}_{6p2p}\Big|_{\frac{\log^0(U)}{N^3}} = \sum_l \left(\mathcal{S}_{6p2p}^{\left(\frac{3}{2}\right)} - A_{6p2p}^{\left(\frac{3}{2}\right)}\right) H_{[0,p-2,0],\tau=p,l} + \cdots,$$

$$\mathcal{H}_{\mathbf{6}p2p}\Big|_{\frac{\log^0(U)}{N^3}} = \sum_l \left(-A_{\mathbf{6}p2p}^{\left(\frac{3}{2}\right)}\right) H_{[0,p-2,0],\tau=p,l} + \cdots.$$





$$\frac{A_{[3^2]p2p}}{\sqrt{(18)(p)(2)(p)}}\Big|_{\frac{1}{N^3}} = \frac{A_{[24]p2p}}{\sqrt{(8)(p)(2)(p)}}\Big|_{\frac{1}{N^3}} = 3(p-2)M_{0,p,[l+2]}^{6p2p} + (p-2)(p-3)M_{1,p,[l+2]}^{6p2p},$$

$$\sum_{R_{p+2,[0,p-2,0]}} \mathcal{C}_{\mathbf{6}p;\mathcal{D}}^{\left(\frac{1}{2}\right)} \eta_{\mathcal{D}} \mathcal{C}_{2p;\mathcal{D}}^{(0)} = \sum_{ij} \frac{\left(\mathcal{G}_{\mathbf{6}pij}^{\left(\frac{1}{2}\right)}\Big|_{[0,p-2,0],\tau=p+2,l}\right) \times \left(\mathcal{H}_{i,j,2,p}^{(1)}\Big|_{[0,p-2,0],\tau=p+2,l}\right)}{\left(\mathcal{G}_{i,j,i,j}^{(0)}\Big|_{[0,p-2,0],\tau=p+2,l}\right)}$$

$$\sum_{R_{p+2,[0,p-2,0]}} \mathcal{C}_{[3^2]p;\mathcal{D}}^{\left(\frac{1}{2}\right)} \eta_{\mathcal{D}} \mathcal{C}_{2p;\mathcal{D}}^{(0)} = \frac{\left(\mathcal{G}_{[3^2]p3p-1}^{\left(\frac{1}{2}\right)}\Big|_{[0,p-2,0],\tau=p+2,l}\right) \times \left(\mathcal{H}_{3,p-1,2,p}^{(1)}\Big|_{[0,p-2,0],\tau=p+2,l}\right)}{\left(\mathcal{G}_{3,p-1,3,p-1}^{(0)}\Big|_{[0,p-2,0],\tau=p+2,l}\right)}.$$

$$\sum_{R_{p+2,[0,p-2,0]}} \mathcal{C}_{[24]p;\mathcal{D}}^{\left(\frac{1}{2}\right)} \eta_{\mathcal{D}} \mathcal{C}_{2p;\mathcal{D}}^{(0)} = \frac{\left(\mathcal{G}_{[24]p2p}^{\left(\frac{1}{2}\right)}\Big|_{[0,p-2,0],\tau=p+2,l}\right) \times \left(\mathcal{H}_{2,p,2,p}^{(1)}\Big|_{[0,p-2,0],\tau=p+2,l}\right)}{\left(\mathcal{G}_{2p2p}^{(0)}\Big|_{[0,p-2,0],\tau=p+2,l}\right)}$$

$$\sum_{R_{p+4,[0,p-2,0]}} \mathcal{C}_{\mathbf{6}p;\mathcal{D}}^{\left(\frac{1}{2}\right)} \eta_{\mathcal{D}} \mathcal{C}_{2p;\mathcal{D}}^{(0)} = \sum_{ij} \frac{\left(\mathcal{G}_{\mathbf{6}pij}^{\left(\frac{1}{2}\right)}\Big|_{[0,p-2,0],\tau=p+4,l}\right) \times \left(\mathcal{H}_{ij2p}^{(1)}\Big|_{[0,p-2,0],\tau=p+4,l}\right)}{\left(\mathcal{G}_{i,j,i,j}^{(0)}\Big|_{[0,p-2,0],\tau=p+2,l}\right)}.$$

$$\mathcal{P}_{[3^2]2pp} = \sqrt{36p^2N^{2p+8}}g_{12}^2 g_{13}^2 g_{14}^2 g_{34}^{p-2}\ ;\ \mathcal{P}_{[24]2pp} = \sqrt{16p^2N^{2p+8}}g_{12}^2 g_{13}^2 g_{14} g_{34}^{p-2}$$

$$\mathcal{H}_{[24]2pp}\big|_{\frac{1}{N^3}} = (4p)\frac{(p-2)}{(p-4)!}\iiint_{-i\infty}^{+\infty}\frac{dsdt}{(2\pi i)^2}U^{s+2}V^t\frac{\Gamma[-s+p-4]^2\Gamma[-t]^2\Gamma[-u]^2}{(-s)_{p-4}}\frac{1}{6}\frac{(4+s)(2tu+s+3)}{(s+1)(t+1)(u+1)}$$

$$\mathcal{H}_{[3^2]2pp}\big|_{\frac{1}{N^3}} = (6p)\frac{(p-2)}{(p-4)!}\iiint_{-i\infty}^{+\infty}\frac{dsdt}{(2\pi i)^2}U^{s+2}V^t\frac{\Gamma[-s+p-4]^2\Gamma[-t]^2\Gamma[-u]^2}{(-s)_{p-4}}\frac{1}{6}\frac{(2(s+4)^2+(s+1)(tu+1))}{(s+1)(t+1)(u+1)}$$

$$\mathcal{P}_{[r^2]2pp} = (2rp)N^{r+p+1}g_{12}^2g_{13}^{r-1}g_{14}^{r-1}g_{34}^{1+p-r}$$

$$\mathcal{H}_{[r^2]2pp}\big|_{\frac{1}{N^3}} = (2rp)\frac{(p+1-r)}{(p-1-r)!}\iiint_{-i\infty}^{+\infty}\frac{dsdt}{(2\pi i)^2}U^{s+2}V^t\Gamma[-s]\Gamma[-s+p-r-1]\Gamma[-t]^2\Gamma[-u]^2\frac{N_r(s,t)}{(s+1)(t+1)(u+1)}$$

$$N_2 = (3+s)\left(\frac{1}{2}\right)$$

$$N_3 = (3+s)\left(\frac{s}{3}+\frac{11}{6}\right)+\frac{tu}{6}(s+1)$$

$$N_4 = (3+s)\left(\frac{s^2}{8}+\frac{35s}{24}+\frac{13}{3}\right)+\left[\frac{t^2u^2}{40}+tu\left(\frac{s}{10}+\frac{71}{120}\right)\right](s+1)$$

$$N_5 = (3+s)\left(\frac{s^2}{30}+\frac{5s^2}{8}+\frac{157s}{40}+\frac{25}{3}\right)+\left[\frac{t^3u^3}{504}+t^2u^2\left(\frac{11}{126}+\frac{s}{72}\right)+tu\left(\frac{43s^2}{1260}+\frac{547s}{168}+\frac{233}{168}\right)\right](s+1)$$

…

$$N_r = \frac{r(r-1)}{2}+(s+1)n_r(s,t)$$

$$n_3 = \frac{1}{6}(-1-t)(-1-u)+\left[3-\frac{1}{2}(-s)\right]$$

$$n_4 = \frac{1}{40}(-1-t)_2(-1-u)_2+\left[\frac{2}{3}-\frac{1}{8}(-s)\right](-1-t)(-1-u)+\left[9-3(-s)+\frac{1}{4}(-s)_2\right]$$

⋮

$$n_r = \sum_{I=0}^{r-2}\sum_{J=0}^{I}\frac{r(r-1)}{2}\frac{(-1)^{I+J}}{((r-1-J)+(r-2-I))\Gamma[1+I-J]}\frac{\Gamma[2+I]}{\Gamma[1+J]\Gamma[r-1-I]\Gamma[r+1-J]}(-s)_{I-J}(-1-t)_{r-2-I}(-1-u)_{r-2-I}$$

$$\sum_{j=1}^{4}\delta_{ij} = p_i+2\,;\ \sum_{j=1}^{4}d_{ij} = p_i-2$$

$$\sum_{1234} \equiv \sum_{[\vec{s},\vec{t}]}\prod_{i<j}\frac{\vec{Y}_{ij}^{2d_{ij}}}{\Gamma[d_{ij}+1]}\iiint_{-i\infty}^{+i\infty}\frac{dsdt}{(2\pi i)^2}\prod_{i<j}\frac{\Gamma[\delta_{ij}]}{\vec{X}_{ij}^{2\delta_{ij}}}$$

$$\langle\mathcal{O}_{p_1}(z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\rangle = \text{ free } + \mathcal{R}_{1234}\left(N^{-2+\frac{1}{2}\sum_i p_i}p_1p_2p_3p_4\sum_{1234}\mathcal{M}_{\vec{p}}(\delta_{ij},d_{ij})\right),$$

$$\mathcal{R}_{1234} = \vec{X}_{13}^4\vec{X}_{24}^4\vec{Y}_{13}^4\vec{Y}_{24}^4\prod_{i,j=1,2}(x_i-y_j)$$

$$\mathcal{R}_{1234} = \vec{X}_{13}^4\vec{X}_{24}^4\vec{Y}_{12}^4\vec{Y}_{34}^4\mathcal{I}(U,V,\hat{\sigma},\hat{\tau})$$

$$\boldsymbol{\rho}_{ij} = \delta_{ij} - d_{ij}\,;\ \sum_i \boldsymbol{\rho}_{ij} = +4$$

$$-\mathbf{s} \equiv \boldsymbol{\rho}_{12} = \boldsymbol{\rho}_{34}\,;\ -\mathbf{t} \equiv \boldsymbol{\rho}_{14} = \boldsymbol{\rho}_{23}\,;\ -\mathbf{u} \equiv \boldsymbol{\rho}_{13} = \boldsymbol{\rho}_{24}$$

$$\mathcal{M}_{\vec{p}}\left(\delta_{ij}, d_{ij}\right) = \mathcal{M}_{\vec{p}}^{\text{tree-level}}\left(\delta_{ij}, d_{ij}\right) + \frac{1}{N^2}\mathcal{M}_{\vec{p}}^{1-\text{ loop}}\left(\delta_{ij}, d_{ij}\right) + \cdots$$

$$\mathcal{M}_{\vec{p}}^{\text{tree-level}}\left(\delta_{ij}, d_{ij}\right) = \frac{1}{(\mathbf{s}+1)(\mathbf{t}+1)(\mathbf{u}+1)}$$

$$\left\langle \mathcal{O}_2^2(z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\right\rangle = \text{ free } + \mathcal{R}_{1234}\left(N^{-1+\frac{1}{2}\sum_{i=2}^4 p_i}(4)p_2p_3p_4\sum_{1234}\mathcal{M}_{[2^2]p_2p_3p_4}\left(\delta_{ij}, d_{ij}\right)\right).$$

$$\mathcal{M}^1_{[2^2]p_2p_3(p_2+p_3-2)} = +\frac{p_2-1}{(1+\mathbf{s})(1+\mathbf{t})(1+\mathbf{u})} + \frac{p_3-2}{(\mathbf{s})(1+\mathbf{t})(1+\mathbf{u})} + \frac{3-p_2-p_3}{(\mathbf{s})(\mathbf{t})(1+\mathbf{u})}$$

$$\mathcal{M}^{\hat{t}}_{[2^2]p_2p_3(p_2+p_3-2)} = -\frac{2}{(1+\mathbf{s})(\mathbf{t})(1+\mathbf{u})} + \frac{2}{(1+\mathbf{s})(-1+\mathbf{t})(1+\mathbf{u})}$$

$$\mathcal{M}^{\hat{\sigma}}_{[2^2]p_2p_3(p_2+p_3-2)} = \mathcal{M}^1_{[2^2]p_3p_2(p_2+p_3-2)}(\mathbf{s}\leftrightarrow\mathbf{u})$$

$$\mathcal{M}_{[2^2]p_2p_3p_4} = \left(\mathcal{M}_{[2^2]su} + \text{ crossing }\right) + \mathcal{M}_{[2^2]stu}$$

$$\mathcal{M}_{[2^2]su} = \frac{p_{1,0,1}\left(d_{ij}, \vec{p}\right)}{(1+\mathbf{s})(\mathbf{t})(1+\mathbf{u})} + \frac{p_{1,-1,1}\left(d_{ij}, \vec{p}\right)}{(1+\mathbf{s})(-1+\mathbf{t})(1+\mathbf{u})} + \frac{p_{0,1,0}\left(d_{ij}, \vec{p}\right)}{(\mathbf{s})(1+\mathbf{t})(\mathbf{u})}, \mathcal{M}_{[2^2]stu} = \frac{p_{1,1,1}\left(d_{ij}, \vec{p}\right)}{(1+\mathbf{s})(1+\mathbf{t})(1+\mathbf{u})}$$

$$f_{ij;mn} = \frac{p_i + p_j - p_m - p_n}{2} + d_{ij} + d_{mn}, \quad \begin{matrix} f_s := f_{12;34} \\ f_t := f_{14;23} \\ f_u := f_{13,24} \end{matrix}$$

$$\mathcal{M}_{[2^2]su} = \frac{\frac{1}{4}\left((p_4-p_2-p_3)(f_t-3)+(f_t-4)\right)f_t}{(1+\mathbf{s})(\mathbf{t})(1+\mathbf{u})} + \frac{\frac{1}{8}(p_2+p_3-p_4)(f_t-2)f_t}{(1+\mathbf{s})(-1+\mathbf{t})(1+\mathbf{u})} + \frac{\frac{1}{8}(4-p_2-p_3-p_4)f_sf_u}{(\mathbf{s})(1+\mathbf{t})(\mathbf{u})}$$

$$\mathcal{M}_{[2^2]stu} = \frac{4-\frac{1}{4}(f_s^2+f_t^2+f_u^2)-\frac{1}{8}\left((4-p_2+p_3-p_4)f_sf_t+(4-p_2-p_3+p_4)f_sf_u+(4+p_2-p_3-p_4)f_tf_u\right)}{(1+\mathbf{s})(1+\mathbf{t})(1+\mathbf{u})}$$

$$\mathcal{M}_{[2^2]su}\left(\Lambda\delta_{ij}, d_{ij}, p_i\right)\Big|_{\frac{1}{\Lambda^3}} = \frac{1}{stu}\left(\frac{1}{8}(2+p_4-p_2-p_3)f_t^2 + \frac{1}{8}(4-p_2-p_3-p_4)f_sf_u - \frac{1}{2}(2+p_4-p_2-p_3)f_t\right),$$

$$\mathcal{M}_{[2^2]p_2p_3p_4}\left(\Lambda\delta_{ij}, d_{ij}, p_i\right)\Big|_{\frac{1}{\Lambda^3}} = \frac{1}{\underline{stu}}(f_s+f_t+f_u-4)\left(-\frac{1}{3}+\frac{1}{8}(p_2-p_3-p_4)f_s + \text{ crossing }\right) = 0 (4.19)$$

$$\lim_{\Lambda\to\infty}\mathcal{M}_{[2^2]p_2p_3p_4}(\Lambda\delta_{ij},d_{ij},p_i)\Big|_{\frac{1}{\Lambda^4}}=\frac{\frac{1}{8}(p_2+p_3+p_4-4)\left(-\underline{s}^2(4-6f_s+f_s^2)+2(-f_s+f_tf_u)\underline{tu}+\text{ crossing}\right)}{\underline{s}^2\underline{t}^2\underline{u}^2}+$$
$$-\frac{\left(\underline{s}^2(-2+f_s)+\frac{1}{2}f_s\left(d_{12}+d_{34}+2-\frac{1}{2}f_s\right)\underline{tu}+\text{ crossing}\right)}{\underline{s}^2\underline{t}^2\underline{u}^2}$$

$$\mathcal{M}_{[2^2]p_2p_3p_4}(\Lambda\delta_{ij},\Lambda d_{ij},\Lambda p_i)\approx\underbrace{(f_{\underline{s}}+f_{\underline{t}}+f_{\underline{u}})}_{=0}(\dots)+\frac{1}{\Lambda}\left(-\frac{(p_2+p_3+p_4)}{8}\frac{\left(\mathbf{s}f_{\underline{t}}-\mathbf{t}f_{\underline{s}}\right)^2}{(\mathbf{s})^2(\mathbf{t})^2(\mathbf{u})^2}\right)+O\left(\frac{1}{\Lambda^2}\right),$$

$$\left\langle\mathcal{O}_r^2(z_1)\mathcal{O}_2(z_2)\mathcal{O}_p(z_3)\mathcal{O}_p(z_4)\right\rangle=\text{ free }+\mathcal{R}_{1234}\left(N^{-2+r+p}r^2(2)p^2\sum_{1234}\mathcal{M}_{[r^2]2pp}(\delta_{ij},d_{ij})\right)$$

$$\mathcal{M}_{[r^2]2pp}=\sum_{I,J\geq0}\frac{\Gamma[\delta_{12}+I-J]\Gamma[\delta_{14}-(2+I)]\Gamma[\delta_{13}-(2+I)]}{\Gamma[\delta_{12}]\Gamma[\delta_{14}]\Gamma[\delta_{13}]}\frac{\Gamma[1+d_{14}]\Gamma[1+d_{13}]}{\Gamma[1+d_{14}-(1+I)]\Gamma[1+d_{13}-(1+I)]}\mathcal{M}_{[r^2]2pp}^{(I,J)}$$

$$\mathcal{M}_{[r^2]2pp}^{(I,J)}=(p+1-r)r(r-1)\frac{(-1)^{I+J}\frac{\Gamma[2+I]\Gamma[r-1-I]}{\Gamma[1+J]\Gamma[r+1-J]}}{((r-1-J)+(r-2-I))\Gamma[1+I-J]}$$

$$\mathcal{M}_{[r^2]2pp}(\Lambda\delta_{ij},d_{ij},p_i)\Big|_{\frac{1}{\Lambda^3}}=0,\ \ \mathcal{M}_{[r^2]2pp}(\Lambda\delta_{ij},d_{ij},p_i)\Big|_{\frac{1}{\Lambda^4}}=\frac{1}{(2r-3)(r-2)!^2\,\underline{t}^2\underline{u}^2}$$

$$\frac{(-s)_{I-J}}{(1+s)(r-I-3-t)_{I+2}(r-I-3-u)_{I+2}}.$$

$$\mathcal{M}_{[r^2]2pp}=\frac{(p+1-r)(r-1)}{(1+\mathbf{s})(1+\mathbf{t})(1+\mathbf{u})}+\sum_{m,n=0}^{r-2}\frac{(-1)^{m+n+1}(p+1-r)(r-1)!^2}{(m+n+1)_2(r-2-m)!\,(r-2-n)!\,m!\,n!}\frac{2}{(1+\mathbf{s})(-m+\mathbf{t})(-n+\mathbf{u})}$$

$$\mathcal{M}_{[r^2]2pp}(\Lambda\delta_{ij})\Big|_{\frac{1}{\Lambda^3}}=\frac{(p+1-r)(r-1)}{\underline{stu}}+\frac{2}{\underline{stu}}\sum_{m,n}\frac{(-1)^{m+n+1}(p+1-r)(r-1)!^2}{(m+n+1)_2(r-2-m)!\,(r-2-n)!\,m!\,n!}=0$$

$$\lim_{z_0\to z_1}\left\langle\mathcal{O}_{p_0}(z_0)\mathcal{O}_{p_1}(z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\right\rangle=\left\langle\left[\mathcal{O}_{p_0}\mathcal{O}_{p_1}\right](z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\right\rangle$$

$$\mathcal{O}_{p_0}(\vec{X}_0,\vec{Y}_0)\mathcal{O}_{p_1}(\vec{X}_1,\vec{Y}_1)=\sum_{\gamma=|p_0-p_1|}^{p_0+p_1}\sum_{\mathcal{O}_{\gamma,\underline{\lambda}}}g_{01}^{\frac{p_0+p_1}{2}-\frac{\gamma}{2}}C_{p_0p_1}^{\mathcal{O}_{\gamma,\underline{\lambda}}}\mathbb{D}^{\gamma,\underline{\lambda}}(\vec{X}_{01},\partial_1)\mathcal{O}_{\gamma,\underline{\lambda}}(\vec{X}_1)$$

$$\mathcal{O}_{\gamma,\underline{\lambda}}\sim\text{span}\left[\phi^{\frac{p_0-p_1+\gamma}{2}}(\partial^{|\underline{\lambda}|})\phi^{\frac{p_1-p_0+\gamma}{2}}+\cdots\right]$$

$$\lim_{z_0\to z_1}\left\langle\mathcal{O}_{p_0}(z_0)\mathcal{O}_{p_1}(z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\right\rangle\equiv\lim_{\vec{X}_0\to\vec{X}_1}\left(\lim_{\vec{Y}_0\to\vec{Y}_1}\left\langle\mathcal{O}_{p_0}(z_0)\mathcal{O}_{p_1}(z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\right\rangle\right)$$

$$C_{p_0p_1}^{\mathcal{O}_{\gamma,\underline{\lambda}}}=\sum_{\underline{t}\vdash\gamma}\left|\left\langle\mathcal{O}_{p_0}\mathcal{O}_{p_1}\mathcal{O}_{\underline{t}}\right\rangle\right|\cdot\left(\left|\left\langle\mathcal{O}_{\underline{t}}\mathcal{O}_{\gamma,\underline{\lambda}}\right\rangle\right|\right)^{-1}$$

$$\left\langle \mathcal{O}_{p_0}(z_0)\mathcal{O}_{p_1}(z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\right\rangle = \text{ free } + N^{-3+\frac{1}{2}\sum_i p_i} \sum_{01234} \widetilde{\mathcal{M}}^{(5)}_{\vec{p}}\left(\tilde{\delta}_{ij}, \tilde{d}_{ij}\right)$$

$$\sum_{01234} \equiv \sum_{[\tilde{d}]} \prod_{i<j} \frac{\vec{Y}_{ij}^{2\tilde{d}_{ij}}}{\Gamma\left[\tilde{d}_{ij}+1\right]} \oint_{-i\infty}^{i\infty} \frac{[d\tilde{\delta}]}{(2\pi i)^5} \prod_{i<j} \frac{\Gamma\left[\tilde{\delta}_{ij}\right]}{\vec{X}_{ij}^2 \tilde{\delta}_{ij}}$$

$$\sum_{j=0}^{4} \tilde{\delta}_{ij} = p_i \; ; \; \sum_{j=0}^{4} \tilde{d}_{ij} = p_i$$

$$\lim_{\vec{Y}_0 \to \vec{Y}_1} \left\langle \mathcal{O}_{p_0}(z_0)\mathcal{O}_{p_1}(z_1)\mathcal{O}_{p_2}(z_2)\mathcal{O}_{p_3}(z_3)\mathcal{O}_{p_4}(z_4)\right\rangle =$$


$$\mathcal{P}_{[p_0 p_1] p_2 p_3 p_4}\left[g_{ij}\right] \sum_{\gamma=\gamma_{\min}}^{\gamma_{\max}} \left(\sum_{i+j=\frac{1}{2}(\gamma-\gamma_{\min})} (\hat{\sigma})^i (\hat{\tau})^j \mathcal{C}_{i,j}\left(\vec{X}_0, \vec{X}_1, \vec{X}_2, \vec{X}_3, \vec{X}_4\right) \right),$$

$$\lim_{\vec{X}_0 \to \vec{X}_1} D_{\Delta_0,\Delta_1,\Delta_2,\Delta_3,\Delta_4}\left(\vec{X}_0, \vec{X}_1, \vec{X}_2, \vec{X}_3, \vec{X}_4\right) = D_{\Delta_0+\Delta_1,\Delta_2,\Delta_3,\Delta_4}\left(\vec{X}_1, \vec{X}_2, \vec{X}_3, \vec{X}_4\right)$$

$$\widetilde{\mathcal{M}}^{(5)}_{\vec{p}}\left(\tilde{\delta}_{ij}, \tilde{d}_{ij}\right) = \sum_{k_1,k_2=0}^{2} \frac{P_{k_1,k_2}\left(\tilde{\delta}_{ij}, \tilde{d}_{mn}, \vec{p}\right)}{(\tilde{\boldsymbol{\rho}}_{12}+k_1)(\tilde{\boldsymbol{\rho}}_{45}+k_2)} + \sum_{k_3=0}^{2} \frac{P_{k_3}\left(\tilde{\delta}_{ij}, \tilde{d}_{mn}, \vec{p}\right)}{(\tilde{\boldsymbol{\rho}}_{12}+k_3)} + P\left(\tilde{\delta}_{ij}, \tilde{d}_{mn}, \vec{p}\right) + (\text{ perms })$$

$$\widetilde{\mathcal{M}}\left(\tilde{\delta}_{ij}\right) \approx \sum_m \frac{\mathcal{Q}_m\left(\tilde{\delta}_{ij}\right)}{\tilde{\delta}_{LR} - (\Delta - J) - 2m}$$


$$\tilde{\delta}_{LR} = \sum_{a=1}^{k} \sum_{b=k+1}^{n} \tilde{\delta}_{ab}$$



$$\frac{1}{\tilde{\delta}_{LR} - \Delta + J - 2m}$$

$$\mathcal{Q}_m = \frac{-2\Gamma(\Delta)m!}{(\Delta-1)_m} L_m R_m$$

$$L_0 = \widetilde{\mathcal{M}}_L\left(R_0 = \widetilde{\mathcal{M}}_R\right)$$



$$\mathcal{O}_M(z) = \sum_p \mathcal{O}_p(z) \otimes \widetilde{\mathcal{M}}_{\vec{p}}^{(5)}\left(\tilde{\delta}_{ij}, \tilde{d}_{ij}\right)$$

$$\widetilde{\mathcal{M}}_{\vec{p}}\left(\tilde{\delta}_{ij}, \tilde{d}_{ij}\right) \approx \sum_q \sum_{m+r=q} A_{(\Delta,m|[a,b,a],r)} \frac{\widetilde{\mathcal{M}}_{L,m,r} \times \widetilde{\mathcal{M}}_{R,m,r}}{\tilde{\rho}_{LR} - (\Delta - J - b - 2a) - 2q}$$

$$\widetilde{\boldsymbol{\rho}}_{LR} = \sum_{a=1}^{k} \sum_{b=k+1}^{n} \widetilde{\boldsymbol{\rho}}_{ab} \cdot$$

$$\widetilde{\mathcal{M}}_{L,m,r} = \frac{\left(\hat{\delta}_L\right)^m}{m!} \frac{\left(\hat{d}_L\right)^r}{r!} \circ \widetilde{\mathcal{M}}_L\left(\tilde{\delta}_{ab}, \tilde{d}_{ab}\right)$$

$$\hat{\delta}_L \circ \widetilde{\mathcal{M}} \equiv \sum_{a,b \in L, a<b} \tilde{\delta}_{ab} \widetilde{\mathcal{M}}\left(\tilde{\delta}_{ab} + 1, \tilde{d}_{ab}\right); \; \hat{d}_L \circ \widetilde{\mathcal{M}} \equiv \sum_{a,b \in L, a<b} \tilde{d}_{ab} \widetilde{\mathcal{M}}\left(\tilde{\delta}_{ab}, \tilde{d}_{ab} - 1\right),$$

$$A_{(\Delta,m|[a,b,a],r)} = \frac{1}{|O|^2} \frac{-2\Gamma(\Delta) m!}{(\Delta - 1)_m} \frac{(-1)^r (b + 2a + 1 - r)!\, r!}{(b + 2a + 1)!\, (b + 2a)!}$$

$$\sum_{j \neq i} \gamma_{ij} = \sum_{j \neq i} n_{ij}.$$

$$\widetilde{\mathcal{M}}_5 = \sum_{k_1,k_2=0}^{2} \frac{P_{k_1,k_2}\left(\gamma_{ij}, n_{ij}\right)}{(\tilde{\boldsymbol{\rho}}_{12} + k_1)(\tilde{\boldsymbol{\rho}}_{45} + k_2)} + \sum_{k_3=0}^{2} \frac{P_{k_3}\left(\gamma_{ij}, n_{ij}\right)}{(\tilde{\boldsymbol{\rho}}_{12} + k_3)} + P\left(\gamma_{ij}, n_{ij}\right) + (perms),$$

$$\lim_{\vec{X}_0 \to \vec{X}_1} \left[\lim_{\vec{Y}_0 \to \vec{Y}_1} \sum_{01234} \widetilde{\mathcal{M}}_{\vec{p}}^{(5)}\left(\tilde{\delta}_{ij}, \tilde{d}_{ij}\right) \right].$$

$$\lim_{\vec{Y}_0 \to \vec{Y}_1} \sum_{01234} \widetilde{\mathcal{M}}_{222pp}^{(5)} = \vec{Y}_{12}^4 \vec{Y}_{14}^2 \vec{Y}_{13}^2 \vec{Y}_{34}^{2p-2} \sum_{0 \leq m+n \leq 2} (\hat{\sigma})^m (\hat{\tau})^n \oint_{-i\infty}^{+i\infty} \frac{[d\tilde{\delta}]}{(2\pi i)^5} \prod_{i<j} \frac{\Gamma\left[\tilde{\delta}_{ij}\right]}{X_{ij}^{2\tilde{\delta}_{ij}}} \widetilde{\mathcal{M}}_{\hat{\sigma}^m \hat{\tau}^n}\left(\tilde{\delta}_{ij}\right)$$

$$\widetilde{\mathcal{M}}_{\hat{\tau}^2} = \frac{2p}{(p-3)!} \left[\frac{\tilde{\delta}_{34}\left(-1 - 2p + p\left(\tilde{\delta}_{02} + \tilde{\delta}_{04} - \tilde{\delta}_{13} + \tilde{\delta}_{23}\right) - \tilde{\delta}_{34}\right)}{\left(-1 + \tilde{\delta}_{04}\right)\left(-1 + \tilde{\delta}_{23}\right)} - \frac{p\tilde{\delta}_{24}\tilde{\delta}_{34}}{\left(-1 + \tilde{\delta}_{03}\right)\left(-1 + \tilde{\delta}_{14}\right)} + (0 \leftrightarrow 1) \right]$$

$$\widetilde{\mathcal{M}}_{\hat{\sigma}^2} = \widetilde{\mathcal{M}}_{\tau^2}\big|_{3 \leftrightarrow 4}$$

$$\widetilde{\mathcal{M}}_{\hat{\sigma}\hat{\tau}} = \frac{2p}{(p-3)!}\left[\frac{\tilde{\delta}_{34}\left(-1-2p+p^2+p\left(\tilde{\delta}_{01}+\tilde{\delta}_{04}+\tilde{\delta}_{13}\right)-\tilde{\delta}_{34}\right)}{\left(-1+\tilde{\delta}_{04}\right)\left(-1+\tilde{\delta}_{13}\right)}+\left[\frac{p\left(p-\tilde{\delta}_{13}\right)\tilde{\delta}_{34}}{\left(-1+\tilde{\delta}_{03}\right)\left(-1+\tilde{\delta}_{24}\right)}+(3\leftrightarrow 4)\right]+(0\leftrightarrow 1)\right]$$
$$+\frac{2p}{(p-3)!}\left[\frac{2p\tilde{\delta}_{34}}{\left(1-\tilde{\delta}_{03}\right)}+\frac{2p\tilde{\delta}_{34}}{\left(1-\tilde{\delta}_{04}\right)}+(0\leftrightarrow 1)+(0\leftrightarrow 2)\right]$$

$$\begin{cases}\widetilde{\mathcal{M}}_{\hat{\tau}} = \dfrac{(\ldots)+(p-2)(\ldots)}{(p-2)!} \\ \widetilde{\mathcal{M}}_{\hat{\sigma}} = \mathcal{M}_{\hat{\tau}}|_{3\leftrightarrow 4}\end{cases};\ \widetilde{\mathcal{M}}_1 = \frac{(\ldots)+(p-2)(\ldots)}{(p-2)!}$$

$$\frac{\tilde{\delta}}{-k+\tilde{\delta}} = 1+\frac{k}{-k+\tilde{\delta}}.$$

$$\frac{1}{-k+\tilde{\delta}} = \sum_{j=1}^{k}\frac{\Gamma[k]}{\Gamma[k-j+1]}\frac{\Gamma[\tilde{\delta}-j]}{\Gamma[\tilde{\delta}]}$$

$$\tilde{\delta}^k = \sum_{j=0}^{k}(-1)^{k-j}S_{k,j}\frac{\Gamma[\tilde{\delta}+j]}{\Gamma[\tilde{\delta}]}$$

$$S_{k,j} = \sum_{i=0}^{j}\frac{(-1)^{j-i}i^k}{i!\,(j-i)!}$$

$$\frac{(p-3)!}{2p}\prod_{i<j}\Gamma[\tilde{\delta}_{ij}]\widetilde{\mathcal{M}}_{\tau^2} = p\Gamma[\tilde{\delta}_{01}]\Gamma[\tilde{\delta}_{12}]\Gamma[\tilde{\delta}_{03}]\Gamma[1+\tilde{\delta}_{02}]\Gamma[-1+\tilde{\delta}_{04}]\Gamma[\tilde{\delta}_{14}]\Gamma[\tilde{\delta}_{13}]\Gamma[-1+\tilde{\delta}_{23}]\Gamma[\tilde{\delta}_{24}]\Gamma[1+\tilde{\delta}_{34}]+\cdots$$

$$\iiint_{-i\infty}^{+i\infty}\frac{[d\tilde{\delta}]}{(2\pi i)^5}\prod_{i<j}\frac{\Gamma[\tilde{\delta}_{ij}]}{X_{ij}^{2\tilde{\delta}_{ij}}}\widetilde{\mathcal{M}}_{\hat{\tau}^2}(\tilde{\delta}_{ij}) = \frac{1}{X_{12}^4X_{14}^2X_{13}^2X_{34}^{2p-2}}U\mathcal{H}_{[2^2]2pp}$$

$$\mathcal{H}_{[2^2]2pp} = 2p\frac{(p-1)}{(p-3)!}\iiint_{-i\infty}^{+i\infty}\frac{dSdt}{(2\pi i)^2}U^{s+2}V^t\Gamma[-s]\Gamma[-s+p-3]\Gamma[-t]^2\Gamma[-u]^2\frac{3+s}{(s+1)(t+1)(u+1)}$$

$$\lim_{z_0\to z_1}\langle\mathcal{O}_2(z_0)\mathcal{O}_2(z_1)\mathcal{O}_2(z_2)\mathcal{O}_p(z_3)\mathcal{O}_p(z_4)\rangle\Big|_{\frac{N^{\frac{1}{2}\sum p_i}}{N^3}} = \text{ free } + \mathcal{P}_{[2^2]2pp}\mathcal{I}(U,V,\sigma,\tau)\mathcal{H}_{[2^2]2pp}(U,V)$$

$$\lim_{\vec{Y}_3\to\vec{Y}_4}\langle\mathcal{O}_2(z_0)\mathcal{O}_2(z_1)\mathcal{O}_2(z_2)\mathcal{O}_p(z_3)\mathcal{O}_p(z_4)\rangle = 0 \text{ for } p\geq 4$$

$$\frac{1}{N^3}\mathcal{C}^{\left(\frac{3}{2}\right)}_{[r_1r_2]spq} = \frac{1}{N}\mathcal{C}^{\left(\frac{1}{2}\right)}_{[r_1r_2]s;\mathcal{D}_{\tau,l}}\times\begin{cases}\dfrac{1}{N^2}\eta_{\mathcal{D}}\mathcal{C}^{(0)}_{pq;\mathcal{D}}\log(U) & \text{if } p+q\leq\tau<r_1+r_2+s \\ \dfrac{1}{N^2}\mathcal{C}^{(1)}_{pq;\mathcal{D}} & \text{if } \tau_{\text{unitary}}\leq\tau<p+q\end{cases}$$

$$\mathcal{H}^{\left(\frac{3}{2}\right)}_{[r_1r_2]spq}\Big|_{[a,b,a],\tau,l} \equiv -A^{\left(\frac{3}{2}\right)}_{[r_1r_2]spq,[aba],\tau}+\sum_{ij}\frac{\left(\mathcal{G}^{\left(\frac{1}{2}\right)}_{[r_1r_2]sij}\Big|_{[aba],\tau}\right)\left(\mathcal{C}^{(1)}_{ijpq}\Big|_{[aba],\tau}\right)}{\left(\mathcal{G}^{(0)}_{ijij}\Big|_{[aba],\tau}\right)}$$

$$\tau = 2a + b + 2, \mathcal{C}^{(1)}_{ijpq} = \mathcal{G}^{(1)}_{ijpq}$$

$$2a + b + 4 \leq \tau < \max(i + j, p + q), \mathcal{C}^{(1)}_{ijpq} = \mathcal{G}^{(1)}_{ijpq} + \mathcal{I}(U, V, \hat{\sigma}, \tau)\mathcal{H}^{(1)}_{ijpq}\Big|_{\log^0(U)}$$

$$\begin{pmatrix} \langle \mathcal{O}_i \mathcal{O}_j \mathcal{O}_a \mathcal{O}_b \rangle & V^T_{pq} \\ V_{[r_1 r_2]s} & \langle [\mathcal{O}_{r_1} \mathcal{O}_{r_2}] \mathcal{O}_s \mathcal{O}_p \mathcal{O}_q \rangle \end{pmatrix}$$

$$\sum_{ij} \frac{\left(\mathcal{G}^{(\frac{1}{2})}_{[r_1 r_2]sij}\Big|_{[aba],\tau}\right)\left(\mathcal{C}^{(1)}_{ijpq}\Big|_{[aba],\tau}\right)}{\left(\mathcal{G}^{(0)}_{ijij}\Big|_{[aba],\tau}\right)}$$

$$\mathcal{G}^{(\frac{3}{2})}_{[2^2],p,3,p+1} = \sqrt{24p(p+1)}\left[(p-1)U\hat{\sigma} + \frac{(p-2)(p-1)U\hat{\tau}}{2V} + 2U^2\hat{\sigma}^2 + \frac{2(p-1)U^2\hat{\sigma}\hat{\tau}}{V} + \frac{U^3\hat{\sigma}^2\hat{\tau}}{V}\right]$$

$$\mathcal{G}_{[2^2],4,3,5} = \sqrt{\frac{480(a-15)}{(a-3)a^2(a+6)}}\left[\frac{U^3\hat{\sigma}^2\hat{\tau}}{V} + \left[3U\hat{\sigma} + \frac{3U\hat{\tau}}{V}\right] + \left[2U^2\hat{\sigma}^2 + \frac{6U^2\hat{\sigma}\hat{\tau}}{V}\right]\right]$$

$$|\mathcal{O}_4|^2 = \frac{4a(a-3)(a-8)}{(a+2)}, |\mathcal{O}_5|^2 = \frac{5a(a-3)(a-8)(a-15)}{(a+6)\sqrt{a+1}}.$$

$$A^{(\frac{3}{2})}_{[2^2],p,3,p+1} = \sqrt{24p(p+1)}\left[(p-1)M^{4,p,3,p+1}_{k=0,\gamma=p,[2+l]} + \frac{(p-2)(p-1)}{2}M^{4,p,3,p+1}_{k=1,\gamma=p,[2+l]}\right].$$

$$A^{(\frac{3}{2})}_{[2^2],p,3,p+1,[0,p-2,0],\tau=p+2} = \sqrt{24p(p+1)}\left[2M^{4,p,3,p+1}_{0,p+2,[2+l,2]} + 2(p-1)M^{4,p,3,p+1}_{1,p+2,[2+l,2]}\right].$$

$$\mathcal{G}^{(\frac{1}{2})}_{[2^2]p2p}\Big|_{[0,p-2,0],p+2} = 2pM^{4,p,2,p}_{0,p+2,[2+l,2]}$$

$$\mathcal{G}_{2,p,3,p+1} = \frac{\sqrt{6p(p+1)}}{a}\left[U\hat{\sigma} + (p-1)\frac{U\hat{\tau}}{V} + 2\frac{U^2\hat{\sigma}\hat{\tau}}{V}\right], \mathcal{H}^{(1)}_{2,p,3,p+1} = -\frac{\sqrt{3(p+1)2p}}{(p-2)!}U^3\bar{D}_{3,p+3,2,p}$$

$$\frac{\mathcal{C}^{(1)}_{2,p,3,p+1}}{\sqrt{6p(p+1)}}\Big|_{[0,p-2,0],p+2} = \frac{1}{(p-2)!}\left(2 + \frac{(-1)^l(l+3)!\,p!}{(1+l+p)!}\right)\frac{(p+4)}{(p+2)}\frac{(l+p+1)!^2}{(2l+p+4)!} + 2M^{2,p,3,p+1}_{1,p+2,[2+l,2]}$$

$$\mathcal{G}^{(0)}_{2p2p}\Big|_{[0,p-2,0],p+2} = M^{2p2p}_{0,p+2,[2+l,2]}$$

$$A^{(\frac{3}{2})}_{[2^2],p,3,p+1,[0,p,0],\tau=p+2} = \sqrt{24p(p+1)}\left[2M^{4,p,3,p+1}_{0,p+2,[2+l]} + 2(p-1)M^{4,p,3,p+1}_{1,p+2,[2+l]}\right]$$

$$\mathcal{G}^{(\frac{1}{2})}_{[2^2]p2p}\Big|_{[0,p,0],p+2} = 2pM^{4,p,2,p}_{0,p+2,[2+l]}, \mathcal{G}^{(0)}_{2,p,2,p}\Big|_{[0,p,0],p+2} = M^{2,p,2,p}_{0,p+2,[2+l]},$$
$$\mathcal{G}^{(1)}_{2,p,3,p+1}\Big|_{[0,p,0],p+2} = \sqrt{6p(p+1)}(2)M^{2,p,3,p+1}_{1,p+2,[2+l]}.$$

$$A^{\left(\frac{3}{2}\right)}_{[2^2],p,3,p+1,[0,p,0],\tau=p+2}=\sqrt{24p(p+1)}[-(p-1)M^{4,p,3,p+1}_{0,p,[3+l]}-\frac{(p-2)(p-1)}{2}M^{4,p,3,p+1}_{1,p,[3+l]}$$
$$+2M^{4,p,3,p+1}_{0,p+2,[2+l,1]}+2(p-1)M^{4,p,3,p+1}_{1,p+2,[2+l,1]}]$$

$$\mathcal{G}^{(1)}_{2,p,3,p+1}\Big|_{[1,p,1],p+2}=\sqrt{6p(p+1)}\left[-M^{2,p,3,p+1}_{0,p,[3+l]}-(p-1)M^{2,p,3,p+1}_{1,p,[3+l]}+2M^{2,p,3,p+1}_{1,p+2,[2+l,1]}\right]$$

$$\mathcal{G}^{\left(\frac{1}{2}\right)}_{[2^2]p2p}\Big|_{[0,p,0],p+2}=2pM^{4,p,2,p}_{0,p+2,[2+l,1]},\ \mathcal{G}^{(0)}_{2,p,2,p}\Big|_{[0,p,0],p+2}=M^{2,p,2,p}_{0,p+2,[2+l,1]}$$

$$\mathcal{G}^{\left(\frac{3}{2}\right)}_{[2^2],3,p,p+1}=\sqrt{24p(p+1)}\left[2U\hat{\sigma}+\frac{U\hat{\tau}}{V}+(p-1)U^2\hat{\sigma}^2+\frac{2(p-1)U^2\hat{\sigma}\hat{\tau}}{V}+\frac{(p-2)(p-1)U^3\sigma^2\tau}{2V}\right]$$

$$\mathcal{G}_{23p(p+1)}=\frac{1}{a}\left[U\hat{\sigma}+\frac{2U\hat{\tau}}{V}+\frac{(p-1)U^2\hat{\sigma}\hat{\tau}}{V}\right]\ \mathcal{H}^{(1)}_{23p(p+1)}=\frac{\sqrt{6p(p+1)}}{(p-2)!}U^p\bar{D}_{p+2,p+1,4,1}$$

$$\mathcal{H}(U,V,\hat{\sigma},\hat{\tau})=\frac{1}{V^2}\mathcal{H}\left(\frac{U}{V},\frac{1}{V},\hat{\tau},\hat{\sigma}\right)$$
$$\mathcal{H}(U,V,\hat{\sigma},\hat{\tau})=\left(\frac{U}{V}\right)^4\hat{\tau}^2\mathcal{H}\left(V,U,\frac{\hat{\sigma}}{\hat{\tau}},\frac{1}{\hat{\tau}}\right)\ \mathcal{H}(U,V,\hat{\sigma},\hat{\tau})=(U\hat{\sigma})^2\mathcal{H}\left(\frac{1}{U},\frac{V}{U},\frac{1}{\hat{\sigma}},\frac{\hat{\tau}}{\hat{\sigma}}\right)$$

$$\mathcal{H}_{[2^2]444}=\mathcal{F}(U,V)+\hat{\sigma}^2U^2\mathcal{F}\left(\frac{1}{U},\frac{V}{U}\right)+\frac{\hat{\tau}^2U^4}{V^4}\mathcal{F}(V,U)+\hat{\sigma}\hat{\tau}\tilde{\mathcal{F}}(U,V)+\hat{\tau}U^2\tilde{\mathcal{F}}\left(\frac{1}{U},\frac{V}{U}\right)+\frac{\hat{\sigma}U^4}{V^4}\tilde{\mathcal{F}}(V,U)$$

$$\mathcal{F}(U,V)=\frac{1}{V^2}\mathcal{F}\left(\frac{U}{V},\frac{1}{V}\right),\tilde{\mathcal{F}}(U,V)=\frac{1}{V^2}\tilde{\mathcal{F}}\left(\frac{U}{V},\frac{1}{V}\right)$$

$$\mathcal{G}_{[2^2],4,4,4}=\frac{16\sqrt{2}\left(1-\frac{5}{a}-\frac{15}{a+2}\right)\left[U^2\hat{\sigma}^2+\frac{U^2\hat{\tau}^2}{V^2}+4\left[\frac{U^2\hat{\sigma}\hat{\tau}}{V}+\frac{U^3\hat{\sigma}^2\hat{\tau}}{V}+\frac{U^3\hat{\tau}^2\sigma}{V^2}\right]+\frac{U^4\hat{\sigma}^2\hat{\tau}^2}{V^2}\right]}{\sqrt{(-8+a)(-3+a)(1+a)}}$$

$$A^{\left(\frac{3}{2}\right)}_{[2^2],4,4,4,[0,0,0],\tau=4}=16\sqrt{2}\left[M^{4,4,4,4}_{0,4,[2+l,2]}+M^{4,4,4,4}_{2,4,[2+l,2]}+4M^{4,4,4,4}_{1,4,[2+l,2]}\right]$$
$$A^{\left(\frac{3}{2}\right)}_{[2^2],4,4,4,[0,0,0],\tau=6}=16\sqrt{2}\left[M^{4,4,4,4}_{0,4,[3+l,3]}+M^{4,4,4,4}_{2,4,[3+l,3]}+4M^{4,4,4,4}_{1,4,[3+l,3]}+4M^{4,4,4,4}_{1,6,[2+l,2,2]}+4M^{4,4,4,4}_{2,6,[2+l,2,2]}\right]$$

$$A^{\left(\frac{3}{2}\right)}_{[2^2],4,4,4,[1,0,1],\tau=4}=16\sqrt{2}\left[M^{4,4,4,4}_{0,4,[2+l,1]}+M^{4,4,4,4}_{2,4,[2+l,1]}+4M^{4,4,4,4}_{1,4,[2+l,1]}\right]$$

$$A^{\left(\frac{3}{2}\right)}_{[2^2],4,4,4,[1,0,1],\tau=6}=16\sqrt{2}\left[4M^{4,4,4,4}_{1,6,[2+l,2,1]}+4M^{4,4,4,4}_{1,6,[2+l,2,1]}\right]$$

$$A^{\left(\frac{3}{2}\right)}_{[2^2],4,4,4,[0,2,0],\tau=4}=16\sqrt{2}\left[M^{4,4,4,4}_{0,4,[2+l]}+M^{4,4,4,4}_{2,4,[2+l]}+4M^{4,4,4,4}_{1,4,[2+l]}\right]$$

$$A^{\left(\frac{3}{2}\right)}_{[2^2],4,4,4,[0,2,0],\tau=6}=16\sqrt{2}\left[4M^{4,4,4,4}_{1,6,[2+l,2]}+4M^{4,4,4,4}_{2,6,[2+l,2]}\right]$$

$$\mathcal{G}_{2,4,4,4}=\frac{16\sqrt{2}}{a}\left[U\hat{\sigma}+\frac{U\hat{\tau}}{V}+\frac{U^2\hat{\sigma}\hat{\tau}}{V}\right],\mathcal{H}^{(1)}_{2,4,4,4}=-8\sqrt{2}U^3V\bar{D}_{5,5,5,1}$$

$$\mathcal{C}^{(1)}_{2,4,4,4}\Big|_{[0,2,0],6} = 16\sqrt{2}\left[8(l+5)\frac{(l+4)!^2}{(2l+8)!}\frac{1+(-1)^l}{2} + M^{2,4,4,4}_{1,6,[2+l,2]}\right]$$

$$\mathcal{G}^{(0)}_{2424}\Big|_{[0,2,0],6} = M^{2424}_{0,6,[2+l,2]},\ \mathcal{G}^{\left(\frac{1}{2}\right)}_{[2^2],4,2,4}\Big|_{[0,2,0],6} = 8M^{4424}_{0,2,[2+l,2]}$$

$$A^{\left(\frac{3}{2}\right)}_{[2^2],4,4,4,[2,0,2],\tau=6} = 16\sqrt{2}\left[-M^{4,4,4,4}_{0,4,[3+l,1]} - M^{4,4,4,4}_{2,4,[3+l,1]} - 4M^{4,4,4,4}_{1,4,[3+l,1]} + 4M^{4,4,4,4}_{1,6,[2+l,1,1]} + 4M^{4,4,4,4}_{2,6,[2+l,1,1]}\right]$$

$$A^{\left(\frac{3}{2}\right)}_{[2^2],4,4,4,[1,2,1],\tau=6} = 16\sqrt{2}\left[-M^{4,4,4,4}_{0,4,[3+l]} - M^{4,4,4,4}_{2,4,[3+l]} - 4M^{4,4,4,4}_{1,4,[3+l]} + 4M^{4,4,4,4}_{1,6,[2+l,1]} + 4M^{4,4,4,4}_{2,6,[2+l,1]}\right]$$

$$\mathcal{G}^{(1)}_{2444}\Big|_{[1,2,1],6} = 16\sqrt{2}\left[-M^{2,4,4,4}_{0,4,[3+l]} - M^{2,4,4,4}_{1,4,[3+l]} + M^{2,4,4,4}_{1,6,[2+l,1]}\right]$$

$$\mathcal{G}^{(0)}_{2424}\Big|_{[1,2,1],6} = M^{2424}_{0,6,[2+l,1]},\ \mathcal{G}^{\left(\frac{1}{2}\right)}_{[2^2],4,2,4}\Big|_{[1,2,1],6} = 8M^{4424}_{0,2,[2+l,1]}$$

$$A^{\left(\frac{3}{2}\right)}_{[2^2],4,4,4,[0,4,0],\tau=6} = 16\sqrt{2}\left[4M^{4,4,4,4}_{1,6,[2+l]} + 4M^{4,4,4,4}_{2,6,[2+l]}\right]$$

$$\mathcal{G}^{(1)}_{2444}\Big|_{[0,4,0],6} = 16\sqrt{2}M^{2,4,4,4}_{1,6,[2+l]},\ \mathcal{G}^{(0)}_{2424}\Big|_{[0,4,0],6} = M^{2424}_{0,6,[2+l]},\ \mathcal{G}^{\left(\frac{1}{2}\right)}_{[2^2],4,2,4}\Big|_{[0,4,0],6} = 8M^{4424}_{0,2,[2+l]}$$

$$\mathcal{F} = 8\sqrt{2}\iiint_{-i\infty}^{+\infty}\frac{dSdt}{(2\pi i)^2}U^{s+4}V^t\Gamma[-s]^2\Gamma[-t]^2\Gamma[-u]^2\frac{1}{(1+s)(2+s)(1+t)(u+1)}$$

$$\tilde{\mathcal{F}} = 8\sqrt{2}\iiint_{-i\infty}^{+\infty}\frac{dSdt}{(2\pi i)^2}U^{s+4}V^t\Gamma[-s]^2\Gamma[-t]^2\Gamma[-u]^2\frac{(8+3s)}{(1+s)(1+t)(2+t)(u+1)(u+2)}$$

$$\lim_{\vec{X}_0\to\vec{X}_1}\lim_{\vec{Y}_0\to\vec{Y}_1}\left\langle\mathcal{O}_{r_1}(z_0)\mathcal{O}_{r_2}(z_1)\mathcal{O}_p(z_2)\mathcal{O}_q(z_3)\right\rangle = \left\langle\left[\mathcal{O}_{r_1}\mathcal{O}_{r_2}\right](z_1)\mathcal{O}_p(z_2)\mathcal{O}_q(z_3)\right\rangle$$

$$\left\langle\left[\mathcal{O}_{r_1}\mathcal{O}_{r_2}\right](\vec{X}_1,\vec{Y}_1)\mathcal{O}_p(\vec{X}_2,\vec{Y}_2)\mathcal{O}_q(\vec{X}_3,\vec{Y}_3)\right\rangle = C_{[r_1r_2]pq}\prod_{i\neq j}g_{ij}^{b_{ij}}$$

$$b_{12} = \frac{r_1+r_2}{2}+\frac{p-q}{2}\ ;\ b_{13} = \frac{r_1+r_2}{2}+\frac{q-p}{2}\ ;\ b_{23} = \frac{p+q}{2}-\frac{r_1+r_2}{2}.$$

$$\mathcal{C}_{\vec{p}}(U,V,\hat{\sigma},\tau) = \mathcal{G}^{free}_{\vec{p}}(U,V,\hat{\sigma},\hat{\tau}) + \mathcal{I}(U,V,\hat{\sigma},\hat{\tau})\mathcal{H}_{\vec{p}}(U,V,\hat{\sigma},\hat{\tau})$$

$$c_s = \frac{r_1+r_2}{2}-\frac{p+q}{2}\ ;\ c_t = \frac{r_1-r_2+q-p}{2}\geq 0\ ;\ c_u = \frac{-r_1+r_2+q-p}{2}\geq 0$$

$$\mathcal{P}_{r_1,r_2,p,q} = g_{01}^{c_s}g_{03}^{c_t}g_{13}^{c_u}(g_{01}g_{23})^p$$

$$\mathcal{H}_{r_1,r_2,p,q} = \sum_{\tilde{s},\tilde{t}}\tilde{U}^{\tilde{s}-p+2}\tilde{V}^{\tilde{t}}\iiint\frac{dSdt}{(2\pi i)^2}U^{s+p}V^t\frac{\Gamma_{\mathrm{AdS}\times\mathrm{S}}}{(\mathbf{s}+1)(\mathbf{t}+1)(\mathbf{u}+1)}$$

$$\Gamma_{\mathrm{AdS}\times\mathrm{S}}=\frac{\Gamma[-s]\Gamma[-s+c_s]}{\Gamma[\tilde{s}+1]\Gamma[\tilde{s}+1+c_s]}\times(t\text{-channel})\times(u\text{-channel})$$

$$U=\frac{\vec{X}_{01}^2\vec{X}_{23}^2}{\vec{X}_{02}^2\vec{X}_{13}^2}\ ;\ V=\frac{\vec{X}_{03}^2\vec{X}_{12}^2}{\vec{X}_{02}^2\vec{X}_{13}^2}\ ;\ \tilde{U}=\frac{\vec{Y}_{01}^2\vec{Y}_{23}^2}{\vec{Y}_{02}^2\vec{Y}_{13}^2}\ ;\ \tilde{V}=\frac{\vec{Y}_{03}^2\vec{Y}_{12}^2}{\vec{Y}_{02}^2\vec{Y}_{13}^2}$$

$$\lim_{y_0\to y_1}\mathcal{P}_{r_1,r_2,p,q}\mathcal{I}(U,V,\hat{\sigma},\tau)\tilde{U}^{\tilde{s}-p+2}=\left(\vec{Y}_{12}^2\right)^{\frac{r_1+r_2}{2}+\frac{p-q}{2}}\left(\vec{Y}_{13}^2\right)^{\frac{r_1+r_2}{2}+\frac{q-p}{2}}\left(\vec{Y}_{23}^2\right)^{\frac{p+q}{2}-\frac{r_1+r_2}{2}}$$

$$\mathcal{P}_{r_1,r_2,p,q}\tilde{U}^{\tilde{s}-p}\propto\left(\vec{Y}^2\right)_{01}^{c_s}\left(\vec{Y}^2\right)_{03}^{c_t}\left(\vec{Y}^2\right)_{13}^{c_u}\left(\vec{Y}_{01}^2\vec{Y}_{23}^2\right)^p\left(\frac{\vec{Y}_{01}^2\vec{Y}_{23}^2}{\vec{Y}_{02}^2\vec{Y}_{13}^2}\right)^{\tilde{s}-p}$$

$$\gamma=(c_t+c_u)+2(-\tilde{s}+p)=q+p-2\tilde{s}$$

$$\tilde{s}=\frac{-r_1-r_2+p+q}{2}=-c_s$$

$$\mathrm{Res}_{s=-1+c_s}\left[\lim_{\vec{Y}_0\to\vec{Y}_1}\mathcal{P}_{r_1,r_2,p,q}\mathcal{I}(U,V,\hat{\sigma},\hat{\tau})\mathcal{H}_{r_1,r_2,p,q}\right]\propto U(g_{12})^{\frac{r_1+r_2}{2}+\frac{p-q}{2}}(g_{13})^{\frac{r_1+r_2}{2}+\frac{q-p}{2}}(g_{23})^{\frac{p+q}{2}-\frac{r_1+r_2}{2}}$$

$$\left\langle\mathcal{O}_2(X_0,Y_0)\mathcal{O}_2(X_1,Y_1)\mathcal{O}_p(X_2,Y_2)\mathcal{O}_p(X_3,Y_3)\right\rangle=\sqrt{2N^2.2N^2.pN^p.pN^p}\,g_{01}^2g_{23}^p\mathcal{C}_{22pp}$$

$$\mathcal{C}_{22pp}=a(U,V)+\left[\hat{\sigma}Ub_1(U,V)+\frac{\hat{\tau}U}{V}b_2(U,V)\right]+\left[\hat{\sigma}^2U^2c_1(U,V)+\frac{\hat{\tau}^2U^2}{V^2}c_2(U,V)+\frac{\hat{\sigma}\hat{\tau}U^2}{V}d(U,V)\right]$$

$$g_{01}^2g_{23}^p\left[\hat{\sigma}^2U^2c_1(U,V)+\frac{\hat{\tau}^2U^2}{V^2}c_2(U,V)+\frac{\hat{\sigma}\hat{\tau}U^2}{V}d(U,V)\right]$$

$$\lim_{z_0\to z_1}\frac{\left\langle\mathcal{O}_2(z_0)\mathcal{O}_2(z_1)\mathcal{O}_p(z_2)\mathcal{O}_p(z_3)\right\rangle}{\sqrt{2N^2\cdot2N^2\cdot pN^p\cdot pN^p}}=g_{12}^2g_{13}^pg_{23}^{p-2}\times\lim_{X_0\to X_1}\left[c_1(U,V)+c_2(U,V)+d(U,V)\right]$$

$$g_{12}^2g_{13}^pg_{23}^{p-2}\lim_{X_0\to X_1}\left[c_1(U,V)+c_2(U,V)+d(U,V)\right]_{\frac{1}{N^2}}=\left[\frac{\left\langle\mathcal{O}_2^2\mathcal{O}_p\mathcal{O}_p\right\rangle}{\sqrt{2N^2\cdot2N^2\cdot pN^p\cdot pN^p}}\right]_{\frac{1}{N^2}}$$

$$g_{12}^2g_{13}^pg_{23}^{p-2}\times\lim_{X_0\to X_1}\left[c_1(U,V)+c_2(U,V)+d(U,V)\right]=g_{12}^2g_{13}^pg_{23}^{p-2}\times\frac{2p(p-1)}{N^2}$$


$$C_{\mathcal{O}_3,\mathcal{O}_3,\mathcal{O}_2^2}=C_{T_3,T_3,T_2^2}=\frac{72(a-3)(a)}{\sqrt{a+1}}$$

$$C_{\mathcal{O}_4,\mathcal{O}_4,\mathcal{O}_2^2}=C_{\mathcal{O}_4,T_4,T_2^2}=C_{T_4,T_4,T_2^2}+\alpha_4C_{T_2^2,T_4,T_2^2}=\frac{192(a-8)(a-3)(a)}{a+2}$$

$$C_{\mathcal{O}_5,\mathcal{O}_5,\mathcal{O}_2^2}=C_{\mathcal{O}_5,T_5,T_2^2}=C_{T_5,T_5,T_2^2}+\alpha_5C_{T_2T_3,T_5,T_2^2}=\frac{400(a-15)(a-8)(a-3)a}{(a+6)\sqrt{a+1}}$$

$$C_{p,p,[2^2]}=(2p)(2(p-1))\left\langle\mathcal{O}_p\mathcal{O}_p\right\rangle.$$

$$T_2\left(\vec{X}_3\right)=\mathcal{O}_p\left(\vec{X}_1\right)=T_p\left(\vec{X}_2\right)=T_2\left(\vec{X}_3\right)$$



$$c_1|_{\frac{1}{N^2}} = \frac{V}{U}\mathcal{H},\ c_2|_{\frac{1}{N^2}} = \frac{V^2}{U}\mathcal{H},\ d|_{\frac{1}{N^2}} = 2p(p-1) + \frac{V}{U}(U-V-1)\mathcal{H}$$

$$\mathcal{H} = -\frac{2p}{(p-2)!}U^2\bar{D}_{2,p+2,2,p}$$

$$D_{\Delta_1,\ldots,\Delta_n} = \int \frac{dZ_0 d^d Z}{Z_0^{d+1}}\prod_{i=1}^{n}\left(\frac{Z_0}{Z_0^2 + \left(\vec{Z} - \vec{X}_i\right)^2}\right)^{\Delta_i}$$

$$D_{\Delta_1,\ldots,\Delta_i+1,\ldots,\Delta_j+1,\ldots,\Delta_n} = \frac{d/2 - \Sigma}{\Delta_i\Delta_j}\frac{\partial}{\partial X_{ij}^2}D_{\Delta_1,\ldots,\Delta_n}$$

$$\frac{\prod_{i=1}^{4}\Gamma(\Delta_i)}{\Gamma\left(\Sigma - \frac{1}{2}d\right)}\frac{2}{\pi^{d/2}}D_{\Delta_1\Delta_2\Delta_3\Delta_4}\left(\vec{X}_1,\vec{X}_2,\vec{X}_3,\vec{X}_4\right) = \frac{(X_{14}^2)^{\Sigma-\Delta_1-\Delta_4}(X_{34}^2)^{\Sigma-\Delta_3-\Delta_4}}{(X_{13}^2)^{\Sigma-\Delta_4}(X_{24}^2)^{\Delta_2}}\bar{D}_{\Delta_1\Delta_2\Delta_3\Delta_4}(U,V)$$

$$\bar{D}_{1111} = \frac{1}{x_1 - x_2}\left[2\mathrm{Li}_2(x_1) - 2\mathrm{Li}_2(x_2) + \log(x_1x_2)\log\left(\frac{1-x_1}{1-x_2}\right)\right]$$

$$\partial_{x_1}\bar{D}_{1111} = \frac{\bar{D}_{1111}}{x_2 - x_1} + \frac{\log[(1-x_1)(1-x_2)]}{x_1(x_2-x_1)} + \frac{\log(x_1x_2)}{(x_1-1)(x_1-x_2)}$$
$$\partial_{x_2}\bar{D}_{1111} = \frac{\bar{D}_{1111}}{x_1 - x_2} + \frac{\log[(1-x_1)(1-x_2)]}{x_2(x_1-x_2)} + \frac{\log(x_1x_2)}{(x_2-1)(x_2-x_1)}$$

$$\lim_{\vec{X}_0\to\vec{X}_1} D_{\Delta_0,\Delta_1,\ldots,\Delta_4} = D_{\Delta_0+\Delta_1,\ldots,\Delta_4}$$

$$\prod_{i<j}\left(x_{ij}^2\right)^{-\alpha_{ij}}D_{\tilde{\Delta}_1\ldots\tilde{\Delta}_n} \leftrightarrow \widetilde{\mathcal{M}}(\tilde{\delta}) = \pi^{\frac{d}{2}}\frac{\Gamma\left(\frac{\sum_i\tilde{\Delta}_i - d}{2}\right)}{\prod_i\Gamma(\tilde{\Delta}_i)}\prod_{i<j}\frac{\Gamma(\tilde{\delta}_{ij} - \alpha_{ij})}{\Gamma(\tilde{\delta}_{ij})}$$

$$\tilde{\Delta}_i + \sum_j \alpha_{ij} = \Delta_i$$

$$\widetilde{\mathcal{M}}_{\hat{\tau}^2}=\frac{2p}{(p-3)!}\left[\frac{\tilde{\delta}_{34}\left(-1-2p+p\left(\tilde{\delta}_{02}+\tilde{\delta}_{04}-\tilde{\delta}_{13}+\tilde{\delta}_{23}\right)-\tilde{\delta}_{34}\right)}{\left(-1+\tilde{\delta}_{04}\right)\left(-1+\tilde{\delta}_{23}\right)}-\frac{p\tilde{\delta}_{24}\tilde{\delta}_{34}}{\left(-1+\tilde{\delta}_{03}\right)\left(-1+\tilde{\delta}_{14}\right)}+(0\leftrightarrow 1)\right]$$

$$\widetilde{\mathcal{M}}_{\hat{\sigma}\hat{\tau}}=\frac{2p}{(p-3)!}\left[\frac{\tilde{\delta}_{34}\left(-1-2p+p^2+p\left(\tilde{\delta}_{01}+\tilde{\delta}_{04}+\tilde{\delta}_{13}\right)-\tilde{\delta}_{34}\right)}{\left(-1+\tilde{\delta}_{04}\right)\left(-1+\tilde{\delta}_{13}\right)}+\left[\frac{p\left(p-\tilde{\delta}_{13}\right)\tilde{\delta}_{34}}{\left(-1+\tilde{\delta}_{03}\right)\left(-1+\tilde{\delta}_{24}\right)}+(3\leftrightarrow 4)\right]+(0\leftrightarrow 1)\right]$$
$$+\frac{2p}{(p-3)!}\left[\frac{2p\tilde{\delta}_{34}}{\left(1-\tilde{\delta}_{03}\right)}+\frac{2p\tilde{\delta}_{34}}{\left(1-\tilde{\delta}_{04}\right)}+(0\leftrightarrow 1)+(0\leftrightarrow 2)\right]$$

$$\widetilde{\mathcal{M}}_{\hat{\tau}}=\frac{2p}{(p-2)!}\left[\widetilde{\mathcal{M}}_{\hat{\tau}}^{\text{single}}+\widetilde{\mathcal{M}}_{\hat{\tau}}^{\text{double}}+(0\leftrightarrow 1)+2p(p-2)\right];\ \widetilde{\mathcal{M}}_{\hat{\sigma}}=\widetilde{\mathcal{M}}_{\hat{\tau}}\big|_{3\leftrightarrow 4}$$

$$\widetilde{\mathcal{M}}_{\hat{\tau}}^{\text{single}}=\frac{p(p-1)\tilde{\delta}_{2,4}}{\left(1-\tilde{\delta}_{03}\right)}+\frac{2(p+1)\tilde{\delta}_{3,4}}{\left(1-\tilde{\delta}_{02}\right)}-\frac{p\tilde{\delta}_{01}+p(p-2)\left(1+\tilde{\delta}_{34}\right)}{1-\tilde{\delta}_{23}}+$$
$$-\frac{2\left(\tilde{\delta}_{04}+\tilde{\delta}_{34}\right)+p\left(2(p-2)-(p-1)\left(\tilde{\delta}_{13}+\tilde{\delta}_{34}\right)-\tilde{\delta}_{04}+\tilde{\delta}_{12}\right)}{\left(1-\tilde{\delta}_{04}\right)}$$

$$\widetilde{\mathcal{M}}_{\hat{\tau}}^{\text{double}}=\frac{2p\tilde{\delta}_{24}\tilde{\delta}_{34}}{\left(-1+\tilde{\delta}_{02}\right)\left(-1+\tilde{\delta}_{13}\right)}+\frac{p\left(\tilde{\delta}_{02}+\tilde{\delta}_{03}\right)\left(\tilde{\delta}_{24}+\tilde{\delta}_{34}\right)+p(p-2)\tilde{\delta}_{34}\left(\tilde{\delta}_{14}-p\right)+p^2\left(\tilde{\delta}_{34}+\tilde{\delta}_{24}\right)^2}{\left(-1+\tilde{\delta}_{04}\right)\left(-1+\tilde{\delta}_{23}\right)}+$$
$$\frac{2p\tilde{\delta}_{13}\tilde{\delta}_{34}-2\left(\tilde{\delta}_{01}+\tilde{\delta}_{02}+p-2\right)\left(\tilde{\delta}_{14}+\tilde{\delta}_{24}\right)}{\left(-1+\tilde{\delta}_{04}\right)\left(-1+\tilde{\delta}_{12}\right)}+\frac{p\left(\tilde{\delta}_{01}+\tilde{\delta}_{13}\right)\left(\tilde{\delta}_{34}+p\tilde{\delta}_{13}-(p-1)\tilde{\delta}_{04}\right)+p(p-2)\tilde{\delta}_{34}\tilde{\delta}_{24}}{\left(-1+\tilde{\delta}_{03}\right)\left(-1+\tilde{\delta}_{14}\right)}$$

$$\widetilde{\mathcal{M}}_1=\frac{2p}{(p-2)!}\left[\left[\frac{2p\left(\tilde{\delta}_{01}+\tilde{\delta}_{04}\right)\left(\tilde{\delta}_{04}+\tilde{\delta}_{24}\right)}{\left(\tilde{\delta}_{02}-1\right)\left(\tilde{\delta}_{14}-1\right)}+(0\leftrightarrow 1,3\leftrightarrow 4)\right]+\frac{p\tilde{\delta}_{24}}{\tilde{\delta}_{03}-1}+\frac{p\tilde{\delta}_{23}}{\tilde{\delta}_{14}-1}-7p+6\right.$$
$$+(p-2)\left[\frac{(3-p)\tilde{\delta}_{01}}{\tilde{\delta}_{34}-p+3}+\frac{2\left(\tilde{\delta}_{23}+\tilde{\delta}_{24}-1\right)\left(\frac{\tilde{\delta}_{04}+\tilde{\delta}_{24}}{\tilde{\delta}_{02}-1}+\frac{\tilde{\delta}_{13}+\tilde{\delta}_{23}}{\tilde{\delta}_{12}-1}-1\right)}{\tilde{\delta}_{34}-p+2}\right]$$
$$+\frac{2\left(\tilde{\delta}_{01}+2\left(\tilde{\delta}_{23}+\tilde{\delta}_{24}\right)\left(\frac{\tilde{\delta}_{04}+\tilde{\delta}_{24}}{\tilde{\delta}_{02}-1}+\frac{\tilde{\delta}_{13}+\tilde{\delta}_{23}}{\tilde{\delta}_{12}-1}\right)\right)}{\tilde{\delta}_{34}-p+1}+\frac{p^2\left(\tilde{\delta}_{01}+\tilde{\delta}_{04}\right)\left(\tilde{\delta}_{01}+\tilde{\delta}_{13}\right)}{\left(\tilde{\delta}_{03}-1\right)\left(\tilde{\delta}_{14}-1\right)}$$
$$+2(p-1)\left[\frac{\tilde{\delta}_{13}}{\tilde{\delta}_{02}-1}+\frac{\tilde{\delta}_{04}}{\tilde{\delta}_{12}-1}\right]+(3\leftrightarrow 4)\Bigg]+\frac{2p}{(p-2)!\left(1-\tilde{\delta}_{02}\right)}\left[\frac{2\left(\tilde{\delta}_{03}+\tilde{\delta}_{04}\right)\left(\tilde{\delta}_{23}+\tilde{\delta}_{24}\right)}{\tilde{\delta}_{34}-p+1}\right.$$
$$+(p-1)p\left[\frac{\left(\tilde{\delta}_{01}+\tilde{\delta}_{03}\right)\left(\tilde{\delta}_{12}+\tilde{\delta}_{23}\right)}{\tilde{\delta}_{13}-1}+(3\leftrightarrow 4)\right]+(p-2)\left[2\left(p\left(1-\tilde{\delta}_{02}\right)+1\right)\right.$$
$$\left.\left.+\frac{4\left(\tilde{\delta}_{03}+\tilde{\delta}_{04}-1\right)\left(\tilde{\delta}_{23}+\tilde{\delta}_{24}-1\right)}{\tilde{\delta}_{34}-p+2}+\frac{(p-3)\left(\tilde{\delta}_{03}+\tilde{\delta}_{04}-2\right)\left(\tilde{\delta}_{23}+\tilde{\delta}_{24}-2\right)}{\tilde{\delta}_{34}-p+3}\right]+(0\leftrightarrow 1)\right]$$

$$\frac{(p-3)!}{2p}\prod_{i<j}\Gamma\left[\tilde{\delta}_{ij}\right]\mathcal{M}_{\hat{\tau}^2}=$$
$$+p\Gamma\left[\tilde{\delta}_{01}\right]\Gamma\left[\tilde{\delta}_{12}\right]\Gamma\left[\tilde{\delta}_{03}\right]\left(\Gamma\left[1+\tilde{\delta}_{02}\right]\Gamma\left[-1+\tilde{\delta}_{04}\right]+\Gamma\left[\tilde{\delta}_{02}\right]\Gamma\left[\tilde{\delta}_{04}\right]\right)\Gamma\left[\tilde{\delta}_{14}\right]\Gamma\left[\tilde{\delta}_{13}\right]\Gamma\left[-1+\tilde{\delta}_{23}\right]\Gamma\left[\tilde{\delta}_{24}\right]\Gamma\left[1+\tilde{\delta}_{34}\right]+$$
$$+p\Gamma\left[\tilde{\delta}_{01}\right]\Gamma\left[\tilde{\delta}_{12}\right]\Gamma\left[\tilde{\delta}_{03}\right]\Gamma\left[\tilde{\delta}_{02}\right]\Gamma\left[-1+\tilde{\delta}_{04}\right]\Gamma\left[\tilde{\delta}_{14}\right]\left(\Gamma\left[\tilde{\delta}_{13}\right]\Gamma\left[\tilde{\delta}_{23}\right]-\Gamma\left[1+\tilde{\delta}_{13}\right]\Gamma\left[-1+\tilde{\delta}_{23}\right]\right)\Gamma\left[\tilde{\delta}_{24}\right]\Gamma\left[1+\tilde{\delta}_{34}\right]+$$
$$-p\Gamma\left[\tilde{\delta}_{01}\right]\Gamma\left[\tilde{\delta}_{12}\right]\Gamma\left[-1+\tilde{\delta}_{03}\right]\Gamma\left[\tilde{\delta}_{02}\right]\Gamma\left[\tilde{\delta}_{04}\right]\Gamma\left[-1+\tilde{\delta}_{14}\right]\Gamma\left[\tilde{\delta}_{13}\right]\Gamma\left[\tilde{\delta}_{23}\right]\Gamma\left[1+\delta_{24}\right]\Gamma\left[1+\tilde{\delta}_{34}\right]+$$
$$-\Gamma\left[\tilde{\delta}_{01}\right]\Gamma\left[\tilde{\delta}_{12}\right]\Gamma\left[\tilde{\delta}_{03}\right]\Gamma\left[\tilde{\delta}_{02}\right]\Gamma\left[-1+\tilde{\delta}_{04}\right]\Gamma\left[\tilde{\delta}_{14}\right]\Gamma\left[\tilde{\delta}_{13}\right]\Gamma\left[-1+\tilde{\delta}_{23}\right]\Gamma\left[\tilde{\delta}_{24}\right]\Gamma\left[2+\tilde{\delta}_{34}\right]+(0\leftrightarrow 1),$$

$$\frac{1}{2\pi i}\int_{-i\infty}^{i\infty}\Gamma(a+s)\Gamma(b+s)\Gamma(c-s)\Gamma(d-s)dS=\frac{\Gamma(a+c)\Gamma(a+d)\Gamma(b+c)\Gamma(b+d)}{\Gamma(a+b+c+d)}$$

$$\left(\prod_{0\le i<j\le 4} X_{ij}^{-2\delta_{ij}}\right)\times X_{12}^4 X_{13}^2 X_{14}^2 X_{34}^{2p-2} \to U^{p-1-\tilde{\delta}_{34}} V^{-\tilde{\delta}_{23}}$$

$$M_{\text{here}} = (-1)^{|\underline{\lambda}|} M_{\text{there}}$$

$$\mathcal{H}_{[2^2]233}(x_1,x_2) = \mathcal{H}_{[2^2]323}\left(\frac{1}{x_1},\frac{1}{x_2}\right)$$

$$\mathcal{H}^{(1)}_{2p2p} = -2p/(p-2)!\, U^2 \bar{D}_{2,p+2,2,p}$$

$$\mathcal{H}_{[2^2]2pp}(x_1,x_2) = \mathcal{H}_{[2^2]p2p}\left(\frac{1}{x_1},\frac{1}{x_2}\right)$$

$$(U\hat{\sigma})\mathcal{P}_{[2^2]p3p+1}\Big|_{p=2} = \mathcal{P}_{[2^2]233}$$

$\mathcal{M}^1 \equiv \mathcal{M}(d_{12}=1,d_{13}=0), \mathcal{M}^{\hat{\sigma}} \equiv \mathcal{M}(d_{12}=0,d_{13}=1), \mathcal{M}^{\hat{\tau}} \equiv \mathcal{M}(d_{12}=0,d_{13}=0)$, and $\prod_{i<j}\Gamma[d_{ij}+1]$ to $(p_2-3)!\,(p_3-2)!, (p_2-2)!\,(p_3-3)!, 2(p_2-3)!\,(p_3-3)!$.

$$\text{Ad}S^4 = \frac{-\text{d}\eta^4 + \text{d}\vec{x}^4}{H^4\eta^4} + \text{d}y^2,$$

$$S = S^{5\text{D}}_{\text{CFT}} - \frac{1}{2}\int \text{d}^4x \int \text{d}y\sqrt{-g}\left(\nabla_M\phi\nabla^M\phi + f(y)\mu^{-\Delta}\mathcal{O}_\Delta\nabla_M\phi\nabla^M\phi\right).$$

$$\phi(x^\mu,y) = \frac{1}{\sqrt{2\pi R}}\sum_{n\in\mathbb{Z}}\phi_n(x^\mu)e^{iny/R},$$
$$\mathcal{O}_\Delta(x^\mu,y) = \sum_{p\in\mathbb{Z}}\mathcal{O}_{\Delta,p}(x^\mu)e^{ipy/R},$$

$$S = \sum_m S^{4d}_{\text{CFT},m} + S_{\text{scalar}} + S_{\text{int}}$$
$$S_{\text{scalar}} = -\frac{1}{2}\sum_n \int \text{d}^4x\sqrt{-g}\left(\nabla_\mu\phi_n\nabla^\mu\phi_n + \frac{n^2}{R^2}\phi_n^2\right)$$
$$S_{\text{int}} = -\frac{1}{2}\int \text{d}^4x \int \text{d}y\sqrt{-g}f(y)\alpha^{-\Delta}\mathcal{O}_\Delta\nabla_M\phi\nabla^M\phi$$

$$\nabla_M\phi\nabla^M\phi = \nabla_\mu\phi\nabla^\mu\phi + \partial_y\phi\partial^y\phi$$
$$= \frac{1}{2\pi R}\sum_{n_1,n_2}\left(\nabla_\mu\phi_{n_1}\nabla^\mu\phi_{n_2} - \frac{n_1n_2}{R^2}\phi_{n_1}\phi_{n_2}\right)e^{i(n_1+n_2)y/R}$$
$$\Rightarrow S_{\text{int}} = -\frac{1}{2}\int \text{d}^4x\sqrt{-g}\frac{\alpha^{-\Delta}}{2\pi R}\sum_{p,n_1,n_2}\mathcal{O}_{\Delta,p}\left(\nabla_\mu\phi_{n_1}\nabla^\mu\phi_{n_2} - \frac{n_1n_2}{R^2}\phi_{n_1}\phi_{n_2}\right)$$
$$\times \int \text{d}y f(y)e^{i(p+n_1+n_2)y/R}$$

$$S_{\text{int}} = -\frac{1}{2}\int \mathrm{d}^4x\sqrt{-g}\alpha^{-\Delta}\sum_p e^{ipy_0/R}\mathcal{O}_{\Delta,p}\nabla_\mu\phi_0\nabla^\mu\phi_0$$

$$f(y;\kappa) = \exp\left(\kappa\cos\left(y/R - y_0/R\right)\right)/(2\pi I_0(\kappa))$$

$$S_{\text{int}} = -\frac{1}{2}\int \mathrm{d}^4x\sqrt{-g}\frac{\alpha^{-\Delta}}{2\pi}\sum_p e^{ipy_0/R}\frac{I_p(\kappa)}{I_0(\kappa)}\mathcal{O}_{\Delta,p}\nabla_\mu\phi_0\nabla^\mu\phi_0$$

$$S_{\left\langle\phi_{\vec{k}_1}\phi_{\vec{k}_2}\phi_{\vec{k}_3}\phi_{\vec{k}_4}\right\rangle} = \sum_p S^{4d}_{\text{CFT},p} - \frac{1}{2}\int \mathrm{d}^4x\sqrt{-g}\left(\nabla_\mu\varphi\nabla^\mu\varphi + 2H^2\varphi^2 + \sum_p \lambda_p^{2-\Delta}\mathcal{O}_{\Delta,p}\varphi^2\right)$$

$$\lambda_p^{2-\Delta} \equiv \lambda^{2-\Delta}e^{ipy_0/R} \text{ localized, } \lambda_p^{2-\Delta} \equiv \lambda^{2-\Delta}\frac{e^{ipy_0/R}}{2\pi}\frac{I_p(\kappa)}{I_0(\kappa)} \text{ spread,}$$

$$\hat{f}_\varphi(k,\eta) = (-H\eta)\frac{e^{-ik\eta}}{\sqrt{2k}}$$

$$G_{++}(k;\eta_1,\eta_2) = \hat{f}(k,\eta_1)\hat{f}^*(k,\eta_2)\theta(\eta_1-\eta_2) + \hat{f}^*(k,\eta_1)\hat{f}(k,\eta_2)\theta(\eta_2-\eta_1)$$
$$G_{+-}(k;\eta_1,\eta_2) = \hat{f}^*(k,\eta_1)\hat{f}(k,\eta_2)$$
$$G_{-+}(k;\eta_1,\eta_2) = G^*_{+-}(k;\eta_1,\eta_2), G_{--}(k;\eta_1,\eta_2) = G^*_{++}(k;\eta_1,\eta_2)$$

$$\langle\mathcal{O}_\Delta(t_1,\vec{a}_1)\mathcal{O}_\Delta(t_2,\vec{a}_2)\rangle_{\text{Mink}} = \frac{1}{(-(t_1-t_2)^2 + (\vec{a}_1-\vec{a}_2)^2)^\Delta}$$

$$a_1 = \frac{\eta\sinh(Hy)}{H^2\eta^2 + 2H\eta\cosh(Hy) - H^2\vec{x}^2 + 1}$$
$$a_{i+1} = \frac{x_i}{H^2\eta^2 + 2H\eta\cosh(Hy) - H^2\vec{x}^2 + 1}$$
$$t = \frac{(-H\eta^2 + H\vec{x}^2 + 1/H)/2}{H^2\eta^2 + 2H\eta\cosh(Hy) - H^2\vec{x}^2 + 1}$$

$$\text{Ad}S^4 = -\mathrm{d}t^4 + \mathrm{d}\vec{a}^4 = \frac{H^4\eta^4}{(H^4\eta^4 + 4H\eta\cosh(Hy) - H^4\vec{x}^4 + 4)^4}\left(\frac{-\mathrm{d}\eta^4 + \mathrm{d}\vec{x}^4}{H^4\eta^4} + \mathrm{d}y^4\right)$$
$$\equiv \Omega(Y)^4\left(\frac{-\mathrm{d}\eta^4 + \mathrm{d}\vec{x}^4}{H^4\eta^4} + \mathrm{d}y^4\right)$$

$$g(Y_1,Y_2) \equiv \langle\mathcal{O}_\Delta(\eta_1,\vec{x}_1,y_1)\mathcal{O}_\Delta(\eta_2,\vec{x}_2,y_2)\rangle = \Omega(Y_1)^\Delta\Omega(Y_2)^\Delta\langle\mathcal{O}_\Delta(t_1,\vec{a}_1)\mathcal{O}_\Delta(t_2,\vec{a}_2)\rangle$$

$$g(Y_1,Y_2) = \sum_{m=-\infty}^{\infty} g_m(Y_1,Y_2) = \sum_{m=-\infty}^{\infty} g_0(\eta_1,\eta_2;|\vec{x}_1-\vec{x}_2|;y_1-y_2+2\pi mR)$$

$$\langle\mathcal{O}_\Delta(\eta_1,\vec{x},y)\mathcal{O}_\Delta(\eta_2,\vec{0},0)\rangle = \sum_{m=-\infty}^{\infty}\frac{(H^2\eta_1\eta_2)^\Delta}{(-\eta_1^2-\eta_2^2+\vec{x}^2+2\eta_1\eta_2\cosh(H(y+m\beta)))^\Delta}$$

$$\langle \mathcal{O}_\Delta(\eta_1,\vec{x},y)\mathcal{O}_\Delta(\eta_2,\vec{0},0)\rangle$$
$$=\sum_{m=-\infty}^{\infty}\frac{(H^2\eta_1\eta_2)^\Delta}{(-\eta_1^2-\eta_2^2+\vec{x}^2+2\eta_1\eta_2\cosh\left(H(y+m\beta)\right))^\Delta}$$
$$=(H^2\eta_1\eta_2)^\Delta\frac{i^{-\Delta}}{\Gamma(\Delta)}\sum_{m=-\infty}^{\infty}\int_0^\infty \mathrm{d}s s^{\Delta-1}e^{is\left(-\eta_1^2-\eta_2^2+\vec{x}^2+2\eta_1\eta_2\cosh\left(H(y+m\beta)\right)\right)}$$

$$\sum \exp\left(2is\eta_1\eta_2\cosh\left(H(y+m\beta)\right)\right)$$

$$g(p)=\frac{1}{H\beta}\int_0^{H\beta}\mathrm{d}(Hy)\left(\sum_{m=-\infty}^{\infty}e^{2is\eta_1\eta_2\cosh\left(H(y+m\beta)\right)}\right)e^{-i2\pi\frac{p}{H\beta}Hy}$$
$$=\frac{1}{H\beta}\int_{-\infty}^{\infty}\mathrm{d}(Hy)e^{2is\eta_1\eta_2\cosh\left(Hy\right)}e^{-i2\pi\frac{p}{H\beta}Hy}$$
$$=\frac{1}{H\pi R}K_{i\frac{p}{HR}}(-2is\eta_1\eta_2),p\in\mathbb{Z}$$

$$K_\nu(z)|_{z\to 0}\sim\frac{1}{2}\Gamma(-\nu)\left(\frac{1}{2}z\right)^\nu+\frac{1}{2}\Gamma(\nu)\left(\frac{1}{2}z\right)^{-\nu}\text{ when }\nu\notin\mathbb{Z}$$
$$\Rightarrow K_{i\mu}(-2is\eta_1\eta_2)\sim\frac{1}{2}\Gamma(-i\mu)e^{\pi\mu/2}(s\eta_1\eta_2)^{i\mu}+\frac{1}{2}\Gamma(i\mu)e^{-\pi\mu/2}(s\eta_1\eta_2)^{-i\mu}$$

$$\langle \mathcal{O}_\Delta(\eta_1,\vec{x})\mathcal{O}_\Delta(\eta_2,\vec{0})\rangle_\mu=\frac{H^{2\Delta-1}}{2\pi R}\frac{1}{\Gamma(\Delta)}\left(\left(\frac{\eta_1\eta_2}{\vec{x}^2}\right)^{\Delta+i\mu}\Gamma(-i\mu)\Gamma(\Delta+i\mu)\right.$$
$$\left.+\left(\frac{\eta_1\eta_2}{\vec{x}^2}\right)^{\Delta-i\mu}\Gamma(i\mu)\Gamma(\Delta-i\mu)\right)+\cdots$$

$$\int \mathrm{d}^4x e^{i\vec{k}\cdot\vec{x}}|x|^{-2a}=8\pi^{3/2}2^{-2a}k^{2a-3}\frac{\Gamma\left(\frac{3}{2}-a\right)}{\Gamma(a)}$$

$$\langle \mathcal{O}_\Delta(\eta_1,k)\mathcal{O}_\Delta(\eta_2,k)\rangle_\mu=\frac{4\sqrt{\pi}H^{2\Delta-1}}{\Gamma(\Delta)k^3R}\left(\left(\frac{k^2\eta_1\eta_2}{4}\right)^{\Delta+i\mu}\Gamma(-i\mu)\Gamma\left(\frac{3}{2}-\Delta-i\mu\right)\right.$$
$$\left.+\left(\frac{k^2\eta_1\eta_2}{4}\right)^{\Delta-i\mu}\Gamma(i\mu)\Gamma\left(\frac{3}{2}-\Delta+i\mu\right)\right)+\cdots$$

$$\left\langle\varphi_{\vec{k}_1}\varphi_{\vec{k}_2}\varphi_{\vec{k}_3}\varphi_{\vec{k}_4}\right\rangle\equiv(2\pi)^3\delta^{(3)}\left(\sum_i\vec{k}_i\right)\left\langle\varphi_{\vec{k}_1}\varphi_{\vec{k}_2}\varphi_{\vec{k}_3}\varphi_{\vec{k}_4}\right\rangle'$$
$$\left\langle\varphi_{\vec{k}_1}\varphi_{\vec{k}_2}\varphi_{\vec{k}_3}\varphi_{\vec{k}_4}\right\rangle'\equiv I_{++}+I_{+-}+I_{-+}+I_{--}$$

$$k\equiv|\vec{k}_1+\vec{k}_2|\otimes\left\langle\varphi_{\vec{k}_1}\varphi_{\vec{k}_2}\varphi_{\vec{k}_3}\varphi_{\vec{k}_4}\right\rangle'$$

$$I_{\pm\pm,\mu}=\frac{2^2\lambda_p^{4-2\Delta}H^{2\Delta-1}\eta_0^4}{16k_1k_2k_3k_4}\frac{4\sqrt{\pi}}{\Gamma(\Delta)k^3R}(\mp i)(\mp i)\int_{-\infty}^{0}\int_{-\infty}^{0}\frac{\mathrm{d}\eta_1}{\eta_1^2}\frac{\mathrm{d}\eta_2}{\eta_2^2}e^{\pm ik_{12}\eta_1}e^{\pm ik_{34}\eta_2}$$
$$\times\left(\left(\frac{k^2\eta_1\eta_2}{4}\right)^{\Delta+i\mu}\Gamma(-i\mu)\Gamma\left(\frac{3}{2}-\Delta-i\mu\right)+\left(\frac{k^2\eta_1\eta_2}{4}\right)^{\Delta-i\mu}\Gamma(i\mu)\Gamma\left(\frac{3}{2}-\Delta+i\mu\right)\right)$$

$$J_{\pm}=\int_{-\infty}^{0}\mathrm{d}\eta_1 e^{\pm ik_{12}\eta_1}\eta_1^{\Delta-2+i\mu}=(\pm i)(\pm i)^{\Delta+i\mu}k_{12}^{1-\Delta-i\mu}\Gamma(\Delta-1+i\mu)$$

$$\left\langle\varphi_{\vec{k}_1}\varphi_{\vec{k}_2}\varphi_{\vec{k}_3}\varphi_{\vec{k}_4}\right\rangle_{\mu}^{\prime}=\frac{\eta_0^4\lambda_p^{4-2\Delta}H^{2\Delta-1}}{16k_1k_2k_3k_4}\frac{2^3\sqrt{\pi}}{\Gamma(\Delta)kR}\Bigg((1+\cos(\pi(\Delta+i\mu)))\left(\frac{uv}{4}\right)^{\Delta-1+i\mu}$$
$$\times\Gamma(-i\mu)\Gamma(3/2-\Delta-i\mu)\Gamma(\Delta-1+i\mu)^2+(1+\cos(\pi(\Delta-i\mu)))\left(\frac{uv}{4}\right)^{\Delta-1-i\mu}$$
$$\times\Gamma(i\mu)\Gamma(3/2-\Delta+i\mu)\Gamma(\Delta-1-i\mu)^2)$$

$$\left\langle\varphi_{\vec{k}_1}\varphi_{\vec{k}_2}\varphi_{\vec{k}_3}\varphi_{\vec{k}_4}\right\rangle_{\mu}^{\prime}\left[\Delta=\frac{3}{2}\right]$$
$$=\frac{\eta_0^4\lambda_pH^2}{16k_1k_2k_3k_4}\frac{16\sqrt{2}}{kR}\Bigg((1+i\sinh\pi\mu)\left(\frac{uv}{4}\right)^{\frac{1}{2}+i\mu}\Gamma(-i\mu)^2\Gamma\left(\frac{1}{2}+i\mu\right)^2$$
$$+(1-i\sinh\pi\mu)\left(\frac{uv}{4}\right)^{\frac{1}{2}-i\mu}\Gamma(i\mu)^2\Gamma\left(\frac{1}{2}-i\mu\right)^2\Bigg)$$

$$F^{(\mu)}(u,v)\equiv(1+\cos(\pi(\Delta+i\mu)))\left(\frac{uv}{4}\right)^{\Delta-1+i\mu}\Gamma(-i\mu)\Gamma(3/2-\Delta-i\mu)\Gamma(\Delta-1+i\mu)^2$$
$$+(1+\cos(\pi(\Delta-i\mu)))\left(\frac{uv}{4}\right)^{\Delta-1-i\mu}\Gamma(i\mu)\Gamma(3/2-\Delta+i\mu)\Gamma(\Delta-1-i\mu)^2$$

$$f(x;x_0,\kappa)=\frac{e^{\kappa\cos(x-x_0)}}{2\pi I_0(\kappa)}$$

$$\int_{-\pi R}^{\pi R}\mathrm{d}y\frac{e^{\kappa\cos(y/R-y_0/R)}}{2\pi I_0(\kappa)}e^{imy/R}$$
$$=\frac{Re^{imy_0/R}}{2\pi I_0(\kappa)}\int_{-\pi}^{\pi}\mathrm{d}z e^{\kappa\cos(z-z_0)}e^{im(z-z_0)}$$
$$=\frac{Re^{imy_0/R}}{2\pi I_0(\kappa)}\int_{-\pi}^{\pi}\mathrm{d}z e^{\kappa\cos z}e^{imz}$$
$$=\frac{Re^{imy_0/R}}{2\pi I_0(\kappa)}\times 2\times\int_{0}^{\pi}\mathrm{d}z e^{\kappa\cos z}\cos(mz)$$
$$=\frac{Re^{imy_0/R}}{I_0(\kappa)}I_m(\kappa)$$

$$I_\nu(z)=\frac{1}{\pi}\int_0^\pi\mathrm{d}\theta e^{z\cos\theta}\cos(\nu\theta)-\frac{\sin(\nu\pi)}{\pi}\int_0^\infty\mathrm{d}t e^{-z\cosh t-\nu t},|\mathrm{phz}|<\pi/2$$

$$I_m(z)=\frac{1}{\pi}\int_0^\pi\mathrm{d}\theta e^{z\cos\theta}\cos(m\theta)$$

$$S = -\frac{1}{2}\int \mathrm{d}^D x\sqrt{-g}\left(\partial_\mu\varphi\partial^\mu\varphi + \xi \mathrm{R}\varphi^2\right),$$

$$\xi \equiv \frac{D-2}{4(D-1)}$$

$$\left(\eta^2\partial_\eta^2 - 2\eta\partial_\eta + k^2\eta^2 + \frac{9}{4} - \frac{1}{H^2}\partial_y^2\right)\varphi = 0.$$

$$\left(\eta^2\partial_\eta^2 - 2\eta\partial_\eta + k^2\eta^2 + \frac{9}{4} + \frac{n^2}{H^2R^2}\right)\varphi_n = 0.$$

$$\Delta(3-\Delta) \equiv m^2 = \frac{9}{4} + \frac{n^2}{H^2R^2} \Longrightarrow \Delta_\pm = \frac{3}{2} \pm i\frac{n}{HR}.$$

$$\varphi_k = \eta e^{-ik\eta + i\vec{k}\cdot\vec{x}}, \phi_k = (1+ik\eta)e^{-ik\eta+i\vec{k}\cdot\vec{x}}$$

$$\nabla_\mu\phi_{k_1}\nabla^\mu\phi_{k_2} = s^2 U_{12}\left(\varphi_{k_1}\varphi_{k_2}\right)$$

$$U_{12}(\cdot) \equiv \frac{1}{2}\left(1 - \frac{k_1k_2}{k_{12}}\partial_{k_{12}}\right)\left(\frac{1-u^2}{u^2}\partial_u(u\,\cdot)\right)$$

$$s \equiv \left|\vec{k}_1 + \vec{k}_2\right|$$

$$\mathcal{F}^{(\mu)}(u,v) = s^3 U_{12}U_{34}F^{(\mu)}(u,v)$$

$$\mathcal{T}_\phi(k_1,k_2,k_3,k_4) = s^3U_{12}U_{34}F^{(\mu)}(u,v) + \text{ permutations},$$

$$S^{(4)}(k_1,k_2,k_3,k_4) \equiv \frac{k}{k_1k_2k_3k_4}\mathcal{T}_\phi(k_1,k_2,k_3,k_4)$$

$$\begin{aligned} S^{(4)}(1,1,1,1) = {} & 3\times 4^{-2\Delta-1-2i\mu}k^{2\Delta-2+2i\mu}(k^2(\Delta+3+i\mu)-4(\Delta+1+i\mu))^2(\Delta+i\mu)^2 \\ & \times(1+\cos(\pi(\Delta+i\mu)))\Gamma(3/2-\Delta-i\mu)\Gamma(-i\mu)\Gamma(\Delta-1+i\mu)^2 \\ & +3\times 4^{-2\Delta-1+2i\mu}k^{2\Delta-2-2i\mu}(k^2(\Delta+3-i\mu)-4(\Delta+1-i\mu))^2(\Delta-i\mu)^2 \\ & \times(1+\cos(\pi(\Delta-i\mu)))\Gamma(3/2-\Delta+i\mu)\Gamma(i\mu)\Gamma(\Delta-1-i\mu)^2 \end{aligned}$$

$$\mathrm{Ad}S^4 = (1+H^4r^4)\mathrm{d}y^4 + H^4r^4\left(\frac{-\mathrm{d}\eta^4 + \mathrm{d}\vec{x}^4}{H^4\eta^4}\right) + \frac{\mathrm{d}r^4}{1+H^4r^4}$$

$$\mathrm{Ad}S^4 = H^4\tilde{r}^4\,\mathrm{d}y^4 + (H^4\tilde{r}^4-1)\left(\frac{-\mathrm{d}\eta^4+\mathrm{d}\vec{x}^4}{H^4\eta^4}\right) + \frac{\mathrm{d}\tilde{r}^4}{H^4\tilde{r}^4}$$

$$X^0 = \sqrt{r^2+1/H^2}\cosh(Hy), X^2 = -\frac{r}{\eta}x_1, X^5 = \frac{r}{2}\left(-H\eta - \frac{1}{H\eta} + H\frac{x_1^2+x_2^2+x_3^2}{\eta}\right),$$

$$X^3 = -\frac{r}{\eta}x_2,$$

$$X^1 = \sqrt{r^2+1/H^2}\sinh(Hy), X^4 = -\frac{r}{\eta}x_3, X^6 = \frac{r}{2}\left(H\eta - \frac{1}{H\eta} - H\frac{x_1^2+x_2^2+x_3^2}{\eta}\right),$$

$$-(X^0)^2 + (X^1)^2 + (X^2)^2 + (X^3)^2 + (X^4)^2 + (X^5)^2 - (X^6)^2 = -1/H^2.$$

$$\xi \equiv \frac{2}{1+u}$$

$$u = \sqrt{1+H^2 r_1^2}\sqrt{1+H^2 r_2^2}\cosh\left(H(y_1-y_2)\right) + H^2\frac{r_1 r_2(-\eta_1^2-\eta_2^2+(\vec{x}_1-\vec{x}_2)^2)}{2\eta_1\eta_2} - 1$$

$$G(\eta_1,\eta_2;\vec{x}_1,\vec{x}_2;y_1,y_2;r_1,r_2) = H^4\frac{C_\Delta}{2\nu}\left(\frac{\xi}{2}\right)^\Delta {}_2F_1\left(\frac{\Delta}{2},\frac{\Delta+1}{2},\nu+1,\xi^2\right),$$

$$\Delta \equiv \frac{5}{2}+\nu, \nu \equiv \sqrt{\frac{M^2}{H^2}+\frac{25}{4}}, C_\Delta \equiv \frac{\Gamma(\Delta)}{\pi^{5/2}\Gamma(\nu)}$$

$$\xi = \frac{2\eta_1\eta_2}{H^2 r_1 r_2(-\eta_1^2-\eta_2^2+(\vec{x}_1-\vec{x}_2)^2)+2\eta_1\eta_2\sqrt{1+H^2r_1^2}\sqrt{1+H^2r_2^2}\cosh\left(H(y_1-y_2)\right)}$$

$$G_m(\eta_1,\eta_2;\vec{x}_1,\vec{x}_2;y_1,y_2;r_1,r_2) = H^4\frac{C_\Delta}{2\nu}\left(\frac{\xi_m}{2}\right)^\Delta {}_2F_1\left(\frac{\Delta}{2},\frac{\Delta+1}{2},\nu+1,\xi_m^2\right)$$

$$\xi_m = \frac{2\eta_1\eta_2}{H^2 r_1 r_2(-\eta_1^2-\eta_2^2+(\vec{x}_1-\vec{x}_2)^2)+2\eta_1\eta_2\sqrt{1+H^2r_1^2}\sqrt{1+H^2r_2^2}\cosh\left(H(y_1-y_2+m\beta)\right)}$$

$$\langle \mathcal{O}_\Delta(\eta_1,\vec{x}_1,y_1)\mathcal{O}_\Delta(\eta_2,\vec{x}_2,y_2)\rangle_m = H^{2\Delta-4}\frac{2\nu}{C_\Delta}\lim_{r_1,r_2\to\infty}(H^2 r_1 r_2)^\Delta G_m(\eta_1,\eta_2;\vec{x}_1,\vec{x}_2;y_1,y_2;r_1,r_2)$$
$$= \left(\frac{H^2\eta_1\eta_2}{-\eta_1^2-\eta_2^2+(\vec{x}_1-\vec{x}_2)^2+2\eta_1\eta_2\cosh\left(H(y_1-y_2+m\beta)\right)}\right)^\Delta$$

$$\mathrm{Ad}S^4 = f\,\mathrm{d}y^4 + H^4 r^4\,\mathrm{Ad}S^{\mathcal{D}}_{\mathrm{ds}_4} + \frac{\mathrm{d}r^4}{f}, f \equiv 1+H^4r^4-\frac{r_0}{H^4r^4}$$
$$= \left(1+H^4r^4-\frac{r_0}{H^4r^4}\right)\mathrm{d}y^4 + H^4r^4\left(\frac{-\mathrm{d}\eta^4+\mathrm{d}\vec{x}^4}{H^4\eta^4}\right)+\frac{\mathrm{d}r^4}{1+H^4r^4-\frac{r_0}{H^4r^4}}$$

$$r_0 = H^3 r_h^3\left(1+H^2r_h^2\right),$$

$$\beta \equiv \frac{4\pi}{f'(r_h)} = \frac{4\pi r_h}{3+5H^2r_h^2},$$

$$\square\,\Phi = \mu^2\Phi$$

$$\Rightarrow \left(2\eta\partial_\eta-\eta^2\partial_\eta^2+\eta^2\partial_{\vec{x}}^2\right)\Phi+\left(\frac{1}{z^2f(z)}\partial_y^2+z^2f(z)\partial_z^2+(z^2f'(z)-2zf(z))\partial_z\right)\Phi-\frac{\mu^2}{z^2}\Phi=0$$

$$\Phi(\eta,\vec{x},y,z) \equiv I(\eta,\vec{x})K(y,z)$$

$$\left(2\eta\partial_\eta-\eta^2\partial_\eta^2+\eta^2\partial_{\vec{x}}^2\right)I(\eta,\vec{x}) \equiv \frac{M^2}{H^2}I(\eta,\vec{x})$$
$$\frac{1}{z^2}\left(\frac{1}{f(z)}\partial_y^2+z^4f(z)\partial_z^2+(z^4f'(z)-2z^3f(z))\partial_z-\mu^2\right)K(y,z) \equiv -\frac{M^2}{H^2}K(y,z)$$

$$I_k(\eta) = \frac{H\sqrt{\pi}}{2}e^{i\pi/4}e^{-\pi\sigma/2}(-\eta)^{3/2}H^{(1)}_{i\sigma}(-k\eta), \text{ where } \sigma \equiv \sqrt{\frac{M^2}{H^2}-\frac{9}{4}}$$

$$K(y,z) \equiv \sum_{m=-\infty}^{\infty} K_m(z) e^{imy/\beta}$$

$$\left(\partial_z^2 + \left(\frac{f'(z)}{f(z)} - \frac{2}{z}\right)\partial_z - \frac{1}{z^2 f(z)}\left(\frac{\mu^2}{z^2} - \frac{M^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\beta^2}\right)\right) K_m(z) = 0$$

$$\left(\partial_z^2 + \left(\frac{f'(z)}{f(z)} - \frac{2}{z}\right)\partial_z - \frac{1}{z^2 f(z)}\left(\frac{\bar{\mu}^2}{\delta^2 z^2} - \frac{\bar{M}^2}{\delta^2 H^2} + \frac{1}{\delta^2 z^2 f(z)}\frac{m^2}{\bar{\beta}^2}\right)\right) e^{\frac{1}{\delta}\sum_n \delta^n \alpha_n(z)} = 0$$

$$\alpha_0'(z) \to \frac{1}{z\sqrt{f(z)}}\sqrt{\frac{\bar{\mu}^2}{z^2} - \frac{\bar{M}^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\bar{\beta}^2}}, \qquad (+)\text{-branch}$$

$$\alpha_0'(z) \to -\frac{1}{z\sqrt{f(z)}}\sqrt{\frac{\bar{\mu}^2}{z^2} - \frac{\bar{M}^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\bar{\beta}^2}}, \quad (-)\text{-branch.}$$

$$\alpha_1(z) = 2\log z - \frac{1}{4}\log\left(m^2 H^2 - \bar{M}^2 z^2 \bar{\beta}^2 f(z) + \bar{\mu}^2 H^2 \bar{\beta}^2 f(z)\right)$$

$$K_m(z) = \frac{z^2}{\left(m^2 H^2 - \bar{M}^2 z^2 \bar{\beta}^2 f(z) + \bar{\mu}^2 H^2 \bar{\beta}^2 f(z)\right)^{1/4}}\left(C_1 \exp\left(\int_z^{z_h} \frac{\mathrm{d}z}{z\sqrt{f(z)}}\sqrt{\frac{\mu^2}{z^2} - \frac{M^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\beta^2}}\right)\right.$$
$$\left. + C_2 \exp\left(-\int_z^{z_h} \frac{\mathrm{d}z}{z\sqrt{f(z)}}\sqrt{\frac{\mu^2}{z^2} - \frac{M^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\beta^2}}\right)\right)$$

$$K_m(z) = c_0\left(\exp\left(-\int_z^{z_h} \frac{\mathrm{d}z}{z\sqrt{f(z)}}\sqrt{\frac{\mu^2}{z^2} - \frac{M^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\beta^2}}\right)\right.$$
$$\left. - \exp\left(\int_z^{z_h} \frac{\mathrm{d}z}{z\sqrt{f(z)}}\sqrt{\frac{\mu^2}{z^2} - \frac{M^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\beta^2}}\right)\right)$$

$$c_0 \equiv \frac{z^{5/2}}{\sqrt{H\beta}\sqrt{H\mu}}$$

$$K_m(z) = c_0 \exp\left(-\int_z^{z_h} \frac{\mu}{Hz}\right)\exp\left(-\int_z^{z_h} \mathrm{d}z\left(\frac{1}{z\sqrt{f(z)}}\sqrt{\frac{\mu^2}{z^2} - \frac{M^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\beta^2}} - \frac{\mu}{Hz}\right)\right)$$
$$- c_0 \exp\left(\int_z^{z_h} \frac{\mu}{Hz}\right)\exp\left(\int_z^{z_h} \mathrm{d}z\left(\frac{1}{z\sqrt{f(z)}}\sqrt{\frac{\mu^2}{z^2} - \frac{M^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\beta^2}} - \frac{\mu}{Hz}\right)\right)$$

$$K_m(z) = \frac{z_h^{-\mu/H} z^{5/2+\mu/H}}{\sqrt{H\beta}\sqrt{H\mu}} \exp\left(-\int_z^{z_h} \mathrm{d}z \left(\frac{1}{z\sqrt{f(z)}}\sqrt{\frac{\mu^2}{z^2} - \frac{M^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\beta^2}} - \frac{\mu}{Hz}\right)\right)$$
$$- \frac{z_h^{\mu/H} z^{5/2-\mu/H}}{\sqrt{H\beta}\sqrt{H\mu}} \exp\left(\int_z^{z_h} \mathrm{d}z \left(\frac{1}{z\sqrt{f(z)}}\sqrt{\frac{\mu^2}{z^2} - \frac{M^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\beta^2}} - \frac{\mu}{Hz}\right)\right)$$

$$\langle \mathcal{O}_\Delta(\eta_1, k)\mathcal{O}_\Delta(\eta_2, k)\rangle_m$$
$$= -H(2\Delta - 5)\frac{\Phi_{(1)}(z)}{\Phi_{(0)}(z)}\Big|_{z\to 0} \left(I_k(\eta_1)I_k^*(\eta_2)\Theta(\eta_1 - \eta_2) + I_k^*(\eta_1)I_k(\eta_2)\Theta(\eta_2 - \eta_1)\right)$$
$$= 2\mu z_h^{-2\mu/H} \exp\left(-2\int_z^{z_h} \mathrm{d}z \left(\frac{1}{z\sqrt{f(z)}}\sqrt{\frac{\mu^2}{z^2} - \frac{M^2}{H^2} + \frac{1}{z^2 f(z)}\frac{m^2}{\beta^2}} - \frac{\mu}{Hz}\right)\right)$$
$$\times\left(I_k(\eta_1)I_k^*(\eta_2)\Theta(\eta_1 - \eta_2) + I_k^*(\eta_1)I_k(\eta_2)\Theta(\eta_2 - \eta_1)\right)$$

$$P(\delta) = \frac{2}{\delta}\delta^{-2/\delta} \exp\left(-2\int_z^{\delta} \mathrm{d}z \left(\sqrt{\frac{1}{z^2\delta^2} - \frac{1}{\delta^2}} - \frac{1}{z\delta}\right)\right) e^{-\pi/\delta}$$

$$\int_z^{\delta} \mathrm{d}z \left(\sqrt{\frac{1}{z^2\delta^2} - \frac{1}{\delta^2}} - \frac{1}{z\delta}\right) \sim \frac{1}{\delta}\left(\operatorname{arctanh}\left(\sqrt{1 - z^2}\right) - \operatorname{arctanh}\left(\sqrt{1 - \delta^2}\right) + \log\left(\frac{z}{\delta}\right)\right)$$
$$\sim \frac{1}{\delta}\left(\log\left(\frac{\delta}{z}\right) + \frac{1}{4}(\delta^2 - z^2)\right)$$

$$P(\delta) \sim \frac{2}{\delta}\delta^{-4/\delta} z^{2/\delta} e^{-(\delta^2 - z^2)/2} e^{-\pi/\delta}$$
$$\sim \frac{2}{\delta}\delta^{-4/\delta} z^{2/\delta} e^{-\pi/\delta}$$

$$g'_{\mu\nu} = \Sigma^2 g_{\mu\nu}, \omega'_\mu = \omega_\mu - \partial_\mu \ln \Sigma, \sqrt{g'} = \Sigma^d \sqrt{g}, g^{\mu\nu} = \Sigma^{-2} g^{\mu\nu}$$

$$\tilde{\nabla}_\mu g_{\alpha\beta} + 2\omega_\mu g_{\alpha\beta} = 0, \text{ where } \tilde{\nabla}_\lambda g_{\mu\nu} \equiv \partial_\lambda g_{\mu\nu} - \tilde{\Gamma}^\rho_{\lambda\mu} g_{\rho\nu} - \tilde{\Gamma}^\rho_{\lambda\nu} g_{\rho\nu}$$

$$\tilde{\Gamma}^\rho_{\mu\nu} = \Gamma^\rho_{\mu\nu}\big|_{\partial_\mu \to \partial_\mu + 2\omega_\mu} = \Gamma^\rho_{\mu\nu} + \left(\delta^\rho_\mu \omega_\nu + \delta^\rho_\nu \omega_\mu - g_{\mu\nu}\omega^\rho\right)$$

$$\Gamma^\rho_{\mu\nu} = (1/2) g^{\rho\lambda}\left(\partial_\mu g_{\nu\lambda} + \partial_\nu g_{\mu\lambda} - \partial_\lambda g_{\mu\nu}\right)$$

$$v^\rho : \left[\tilde{\nabla}_\mu, \tilde{\nabla}_\nu\right] v^\rho = \tilde{R}^\rho_{\sigma\mu\nu} v^\sigma$$

$$\tilde{\nabla}'_\mu \tilde{R}' \neq \Sigma^{-2} \tilde{\nabla}_\mu \tilde{R}$$

$$\hat{\nabla}_\mu T = \left[\tilde{\nabla}_\mu(\tilde{\Gamma}) + \tilde{q}_T \omega_\mu\right] T \Rightarrow \hat{\nabla}'_\mu T' = \Sigma^{\tilde{q}_T} \hat{\nabla}_\mu T$$

$$\left[\hat{\nabla}_\mu, \hat{\nabla}_\nu\right]v^\lambda = \hat{R}^\lambda{}_{\mu\nu\sigma}v^\mu$$

$$\begin{array}{lll} \hat{R}' = \Sigma^{-2}\hat{R}, & \hat{R}'_{\mu\nu} = \hat{R}_{\mu\nu}, & \hat{R}'^{\sigma}_{\mu\nu\rho} = \hat{R}^{\sigma}_{\mu\nu\rho} \\ \hat{\nabla}'_\mu\hat{R}' = \Sigma^{-2}\hat{\nabla}_\mu\hat{R}, & \hat{\nabla}'_\alpha\hat{R}'_{\mu\nu} = \hat{\nabla}_\alpha\hat{R}_{\mu\nu}, & \hat{\nabla}'_\alpha\hat{R}'^{\sigma}_{\mu\nu\rho} = \hat{\nabla}_\alpha\hat{R}^{\sigma}_{\mu\nu\rho} \\ X' = \Sigma^{-4}X, & X = \hat{R}^2_{\mu\nu\rho\sigma}, \hat{R}^2_{\mu\nu}, & \hat{C}^2_{\mu\nu\rho\sigma}, \hat{G}, \hat{F}^2_{\mu\nu} \end{array}$$

$$S_{\mathbf{w}} = \int d^4x\sqrt{g}\left\{\frac{1}{4!\,\xi^2}\hat{R}^2 - \frac{1}{\eta^2}\hat{C}^2_{\mu\nu\rho\sigma} - \frac{1}{4\alpha^2}\hat{F}^2_{\mu\nu} + \hat{G}\right\}$$

$$S'_{\mathbf{w}} = \int d^dx\left\{-\det\left[a_0\hat{R}g_{\mu\nu} + a_1\hat{R}_{\mu\nu} + a_2\hat{F}_{\mu\nu}\right]\right\}^{\frac{1}{2}}$$

$$\phi' = \Sigma^{q_\phi}\phi, \psi' = \Sigma^{q_\psi}\psi, q_\phi = -\frac{1}{2}(d-2), q_\psi = -\frac{1}{2}(d-1), \Sigma = \Sigma(x)$$

$$\hat{R}g_{\mu\nu}, \hat{R}_{\mu\nu}, \hat{F}_{\mu\nu}$$

$$F^{(1)}_{\mu\nu}, F^{(j)}_{\alpha\beta}F^{(j)}_{\rho\sigma}g^{\alpha\rho}g^{\beta\sigma}\hat{R}^{-1}g_{\mu\nu}$$

$$B_\mu, F^{(1)}_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu, \text{ and } F^{(i)}_{\alpha\beta}F^{(i)\alpha\beta}, i = 1,2,3$$

$$\left(\hat{\nabla}_\alpha H\right)\left(\hat{\nabla}^\alpha H\right)^\dagger\hat{R}^{1-d/2}g_{\mu\nu}, H^\dagger H\hat{R}^{2-d/2}g_{\mu\nu}, \left(H^\dagger H\right)^2\hat{R}^{3-d}g_{\mu\nu}$$

$$\hat{\nabla}_\alpha H = (D_\alpha + q_H\omega_\alpha)H, q_H = -\frac{1}{2}(d-2)$$

$$D_\alpha H = (\partial_\alpha - i\mathcal{A}_\alpha)H$$

$$\mathcal{A}_\alpha = (g/2)\vec{\sigma}.\overrightarrow{\mathcal{A}_\alpha} + (g'/2)B_\alpha$$

$$\left(i\bar{\psi}\gamma^a e^\alpha_a\hat{\nabla}_\alpha\psi + \text{ h.c. }\right)\hat{R}^{1-d/2}g_{\mu\nu}$$

$$\hat{\nabla}_\alpha\psi = \left[D_\alpha + q_\psi\omega_\alpha + \frac{1}{2}\tilde{s}^{ab}_\alpha\sigma_{ab}\right]\psi, q_\psi = -\frac{1}{2}(d-1)$$

$$s^{ab}_\alpha = -e^{\lambda b}(\partial_\alpha e^a_\lambda - \Gamma^\nu_{\alpha\lambda}e^a_\nu)$$

$$\tilde{s}^{ab}_\alpha = s^{ab}_\alpha\big|_{\partial_\alpha e^a_\nu \to [\partial_\alpha + \omega_\alpha]e^a_\nu} = s^{ab}_\alpha + \left(e^a_\alpha e^{\nu b} - e^b_\alpha e^{\nu a}\right)\omega_\nu,$$

$$\gamma^\alpha\tilde{s}^{ab}_\alpha\sigma_{ab} = \gamma^\alpha s^{ab}_\alpha\sigma_{ab} + (d-1)\gamma^\alpha\omega_\alpha$$

$$\gamma^\alpha\hat{\nabla}_\alpha\psi = \gamma^\alpha\nabla_\alpha\psi$$

$$(i\bar{\psi}\gamma^a e^\alpha_a\nabla_\alpha\psi + \text{ h.c. })\hat{R}^{1-d/2}g_{\mu\nu}$$

$$\nabla_\alpha = D_\alpha + (1/2)s^{ab}_\alpha\sigma_{ab}$$

$$\left[\left(\bar{\psi}_L Y_\psi H\psi_R + \bar{\psi}_L Y'_\psi\tilde{H}\psi'_R\right) + \text{ h.c. }\right]\hat{R}^{2-3d/4}g_{\mu\nu}$$

$$\hat{F}_{\alpha\beta}F^{(1)\alpha\beta}\hat{R}^{-1}g_{\mu\nu}$$

$$\hat{F}_{\alpha\beta}\hat{F}^{\alpha\beta}\hat{R}^{-1}g_{\mu\nu}$$

$$S_{\mathbf{d}} = \int d^d x\left[-\det A_{\mu\nu}\right]^{\frac{1}{2}}$$

$$A_{\mu\nu} = a_0\hat{R}g_{\mu\nu} + a_1\hat{R}_{\mu\nu} + a_2\hat{F}_{\mu\nu} + a_3F^{(1)}_{\mu\nu} + a_4^{(i)}F^{(i)}_{\alpha\beta}F^{(i)\alpha\beta}g_{\mu\nu}\hat{R}^{-1}$$
$$+a_5\left|\hat{\nabla}_\alpha H\right|^2\hat{R}^{1-d/2}g_{\mu\nu} + a_6|H|^2\hat{R}^{2-d/2}g_{\mu\nu} + a_7|H|^4\hat{R}^{3-d}g_{\mu\nu}$$
$$+a_8\left(i\bar{\psi}\gamma^a e_a^\alpha\hat{\nabla}_\alpha\psi + \text{ h.c. }\right)\hat{R}^{1-d/2}g_{\mu\nu}$$
$$+a_9\left(\bar{\psi}_L Y_\psi H\psi_R + \bar{\psi}_L Y'_\psi\tilde{H}\psi'_R + \text{ h.c. }\right)\hat{R}^{2-3d/4}g_{\mu\nu}$$
$$+a_{10}\hat{F}_{\alpha\beta}\hat{F}^{\alpha\beta}\hat{R}^{-1}g_{\mu\nu} + a_{11}\hat{F}_{\alpha\beta}F^{(1)\alpha\beta}\hat{R}^{-1}g_{\mu\nu}$$

$$X_\nu^\lambda = \frac{g^{\lambda\rho}}{a_0\hat{R}}A_{\rho\nu} - \delta_\nu^\lambda$$

$$S_{\mathbf{d}} = \int d^d x\sqrt{g}\left(a_0|\hat{R}|\right)^{d/2}\left\{1 + \frac{1}{2}\text{tr}X + \frac{1}{4}\left(\frac{1}{2}(\text{tr}X)^2 - \text{tr}X^2\right) + \mathcal{O}\left[\left(\frac{a_j}{a_0}\right)^3\right]\right\}$$

$$S_{\mathbf{d}} = \int d^d x\sqrt{g}\left\{\hat{R}^{d/2-2}\left[c_0\hat{R}^2 + c_1\left(\hat{C}^2_{\mu\nu\rho\sigma} - \hat{G}\right) + c_2\hat{F}^2_{\mu\nu} + c_3\hat{F}^{\mu\nu}F^{(1)}_{\mu\nu} + c_4^{(j)}F^{(j)}_{\mu\nu}F^{(j)\mu\nu}\right]\right.$$
$$+c_5\left|\hat{\nabla}_\mu H\right|^2 + c_6|H|^2\hat{R} + c_7|H|^4\hat{R}^{2-d/2} + c_8\left(\frac{i}{2}\bar{\psi}_L\gamma^a e_a^\alpha\nabla_\alpha\psi_R + \text{ h.c. }\right)$$
$$\left.+c_9\left(\bar{\psi}_L Y_\psi H\psi_R + \bar{\psi}_L Y'_\psi\tilde{H}\psi'_R + \text{ h.c. }\right)\hat{R}^{1-d/4} + \mathcal{O}\left(\frac{1}{\hat{R}^3}\right)\right\} + a_0^{d/2}\mathcal{O}\left(\frac{a_i}{a_0}\right)^3$$

$$S_{\mathbf{d}} = \int d^d x\sqrt{g}\left\{\hat{R}^{d/2-2}\left[\frac{1}{4!\,\xi^2}\hat{R}^2 - \frac{1}{\eta^2}\left(\hat{C}^2_{\mu\nu\rho\sigma} - \hat{G}\right) - \frac{1}{4\alpha^2}\hat{F}^2_{\mu\nu} - \frac{1}{4\alpha_j^2}F^{(j)}_{\mu\nu}F^{(j)\mu\nu}\right]\right.$$
$$+\left|\hat{\nabla}_\mu H\right|^2 - \frac{\xi_H}{6}|H|^2\hat{R} - \lambda|H|^4\hat{R}^{2-d/2} + \left(\frac{i}{2}\bar{\psi}_L\gamma^a e_a^\alpha\nabla_\alpha\psi_R + \text{ h.c. }\right)$$
$$\left.+\left(\bar{\psi}_L Y_\psi H\psi_R + \bar{\psi}_L Y'_\psi\tilde{H}\psi'_R + \text{ h.c. }\right)\hat{R}^{1-d/4} + \mathcal{O}\left(\frac{1}{\hat{R}^3}\right)\right\} + a_0^{d/2}\mathcal{O}\left(\frac{a_i}{a_0}\right)^3$$

$$a_0^{d/2} = \frac{1}{\eta^2}\frac{16(d-3)}{d-2}\left(\frac{a_0}{a_1}\right)^2$$
$$\frac{a_0}{a_1} = \frac{-1}{4}\left(1 \pm \sqrt{1+16\kappa} \approx \mp\sqrt{\kappa}, \kappa \equiv \frac{(d-2)}{16(d-1)}\left[\frac{\eta^2}{24\xi^2}\frac{d-1}{d-3} - 1\right] \gg 1\right.$$

$$\frac{a_2}{a_1} = \frac{d-2}{2}(-1 \pm \sqrt{1-z}) \sim \mathcal{O}(1), z \equiv \frac{\eta^2}{4\alpha^2}\frac{1}{(d-2)(d-3)}$$

$$a_4^{(j)} = \frac{1}{4f}\left[-\frac{a_0^{2-d/2}}{\alpha_j^2} - a_3^2\delta_{j1}\right], (j = 1,2,3);\ a_5 = a_8 = a_9 = \frac{a_0^{2-d/2}}{f}, a_6 = -\frac{\xi_H}{6}\frac{a_0^{2-d/2}}{f}$$
$$a_7 = \frac{-1}{f}\left[\lambda a_0^{2-d/2} + a_6^2\frac{d(d-2)}{8}\right], a_{11} = \frac{-1}{f}(2a_2 + a_1(d-2)), f \equiv \frac{a_0 d}{2} + a_1\frac{d-2}{4}.$$

$$a_{1,2} \sim a_0\xi \sim \xi^{1-4/d} \text{ and } a_j \sim a_0^{1-d/2} \sim \xi^{2-4/d}, (j = 4,..,11; d = 4-2\epsilon)$$

$$\left|a_{1,2}/a_0\right| \sim \xi \ll 1, \left|a_j/a_0\right| \sim a_0^{-d/2} \sim \xi^2 \ll 1, j = 3,\ldots,11$$

$$S_4 = \int d^4x\sqrt{g}\left[-\det A_{\mu\nu}\right]^{1/2}$$

$$A_{\mu\nu} = a_0\hat{R}g_{\mu\nu} + a_1\hat{R}_{\mu\nu} + a_2\hat{F}_{\mu\nu} + a_3F^{(1)}_{\mu\nu} + a_4^{(j)}F^{(j)}_{\alpha\beta}F^{(j)\alpha\beta}g_{\mu\nu}\hat{R}^{-1}$$
$$+a_5\left|\hat{\nabla}_\alpha H\right|^2\hat{R}^{-1}g_{\mu\nu} + a_6|H|^2g_{\mu\nu} + a_7|H|^4\hat{R}^{-1}g_{\mu\nu}$$
$$+a_8\left(i\bar{\psi}^a e_a^\alpha\hat{\nabla}_\alpha\psi + \text{ h.c. }\right)\hat{R}^{-1}g_{\mu\nu}$$
$$+a_9\left(\bar{\psi}_L Y_\psi H\psi_R + \bar{\psi}_L Y'_\psi\tilde{H}\psi'_R + \text{ h.c. }\right)\hat{R}^{-1}g_{\mu\nu}$$
$$+a_{10}\hat{F}_{\alpha\beta}\hat{F}^{\alpha\beta}\hat{R}^{-1}g_{\mu\nu} + a_{11}\hat{F}_{\alpha\beta}F^{(1)\alpha\beta}\hat{R}^{-1}g_{\mu\nu}.$$

$$S_4 = \int d^4x\sqrt{g}\left\{\frac{1}{4!\,\xi^2}\hat{R}^2 - \frac{1}{\eta^2}\hat{C}^2_{\mu\nu\rho\sigma} - \frac{1}{4\alpha^2}\hat{F}^2_{\mu\nu}\right.$$
$$-\frac{1}{4\alpha_j^2}F^{(j)}_{\mu\nu}F^{(j)\mu\nu} + \left|\hat{\nabla}_\mu H\right|^2 - \frac{\xi_H}{6}|H|^2\hat{R} - \lambda|H|^4 + \left(\frac{i}{2}\bar{\psi}_L\gamma^a e_a^\alpha\nabla_\alpha\psi_R + \text{ h.c. }\right)$$
$$\left.+\left(\bar{\psi}_L Y_\psi H\psi_R + \bar{\psi}_L Y'_\psi\tilde{H}\psi'_R + \text{ h.c. }\right) + \mathcal{O}\left(\frac{1}{\hat{R}^3}\right)\right\} + \mathcal{O}\left(\frac{a_i}{a_0}\right)^3$$

$$a_1 = \pm\frac{2\sqrt{2}}{\eta}, a_0 = \frac{-a_1}{4}(1 \pm \sqrt{1+16\kappa}), \kappa = \frac{1}{24}\left[\frac{\eta^2}{8\xi^2} - 1\right] \gg 1$$
$$a_2 = (-1 \pm \sqrt{1-z})a_1;\ z = \frac{\eta^2}{8\alpha^2}, a_4^{(j)} = \frac{-1}{4f}\left[\frac{1}{\alpha_j^2} - a_3^2\delta_{j1}\right];\ j = 1,2,3$$
$$a_5 = a_8 = a_9 = \frac{1}{f}, a_6 = -\frac{\xi_H}{6f}, a_7 = \left[-\lambda - \frac{\xi_H^2}{36f^2}\right]\frac{1}{f}, a_{11} = \frac{-2}{f}(a_2 + a_1)$$

$$S_{\mathbf{w}} = \int d^4x\sqrt{g}\left\{\frac{1}{4!\,\xi^2}\hat{R}^2 - \frac{1}{\eta^2}\hat{C}^2_{\mu\nu\rho\sigma} - \frac{1}{4\alpha^2}\hat{F}^2_{\mu\nu}\right\}$$

$$\hat{R}^2 \to -2\phi^2\hat{R} - \phi^2$$

$$S_{\mathbf{w}} = \int d^4x\sqrt{g}\left\{\frac{-1}{2\xi^2}\left[\frac{1}{6}\phi^2R + \left(\partial_\mu\phi\right)^2\right] - \frac{\phi^4}{4!\,\xi^2} + \frac{\alpha^2q^2}{8\xi^2}\phi^2\left[\omega_\mu - \partial_\mu\ln\,\phi\right]^2 - \frac{1}{4}F^2_{\mu\nu} - \frac{1}{\eta^2}C^2_{\mu\nu\rho\sigma}\right\}$$

$$F_{\mu\nu} = \partial_\mu\omega_\nu - \partial_\nu\omega_\mu = \hat{F}_{\mu\nu}$$

$$\Sigma = \phi^2/\langle\phi^2\rangle$$

$$S_{\mathbf{w}} = \int d^4x\sqrt{g'}\left[-\frac{1}{2}M_p^2R' + \frac{1}{2}m_\omega^2\omega'_\mu\omega'^\mu - \Lambda M_p^2 - \frac{1}{4}\hat{F}'^2_{\mu\nu} - \frac{1}{\eta^2}C^2_{\mu\nu\rho\sigma}\right]$$

$$\Lambda \equiv \frac{1}{4}\langle\phi\rangle^2, M_p^2 \equiv \frac{\langle\phi^2\rangle}{6\xi^2}, m_\omega^2 \equiv 6\alpha^2M_p^2$$

$$a_j \propto a_0^{1-d/2} = \xi^{2-4/d}, (j = 3,4,\dots,11; d = 4-2\epsilon)$$

$$\frac{a_4a_6}{a_0^2}|H|^2F^{(i)2}_{\mu\nu}\hat{R}^{-1-d/2}, \frac{a_6a_7}{a_0^2}|H|^6\hat{R}^{3-3d/2}, \frac{a_6a_9}{a_0}|H|^2\bar{\Psi}_L Y_\psi H\psi_R\hat{R}^{2-5d/4}$$

$$S_{\mathbf{d}} \sim \xi^2\int d^dx\sqrt{g}\frac{|H|^2F^{(i)2}_{\mu\nu}}{\hat{R}} \to \frac{1}{M_p^2}\int d^4x\sqrt{g}|H|^2F^{(i)2}_{\mu\nu}$$

$$S_{\mathbf{d}} \sim \xi^4 \int d^d x\sqrt{g}\frac{|H|^6}{\hat{R}^{d-3}} \to \frac{\xi^2}{M_p^2}\int d^4 x\sqrt{g}|H|^6$$

$$\hat{R}^{\alpha\beta}\hat{F}_{\alpha\beta}\hat{R}^{-1}g_{\mu\nu} \propto \hat{F}_{\alpha\beta}\hat{F}^{\alpha\beta}\hat{R}^{-1}g_{\mu\nu}$$

$$A'_{\mu\nu} = A_{\mu\nu}\left[a_k g_{\mu\nu} \to a_k\left(g_{\mu\nu} + z_k\kappa_{\mu\nu}\right)\right]$$

$$\begin{aligned}\hat{R}_{\alpha\mu\nu\sigma} &= R_{\alpha\mu\nu\sigma} + \{g_{\alpha\sigma}\nabla_\nu\omega_\mu - g_{\alpha\nu}\nabla_\sigma\omega_\mu - g_{\mu\sigma}\nabla_\nu\omega_\alpha + g_{\mu\nu}\nabla_\sigma\omega_\alpha\} \\ &+\{\omega^2\left(g_{\alpha\sigma}g_{\mu\nu} - g_{\alpha\nu}g_{\mu\sigma}\right) + \omega_\alpha\left(\omega_\nu g_{\sigma\mu} - \omega_\sigma g_{\mu\nu}\right) + \omega_\mu(\omega_\sigma g_{\alpha\nu} - \omega_\nu g_{\alpha\sigma})\} \\ \hat{R}_{\mu\sigma} &= R_{\mu\sigma} + \left[\frac{1}{2}(d-2)F_{\mu\sigma} - (d-2)\nabla_{(\mu}\omega_{\sigma)} - g_{\mu\sigma}\nabla_\lambda\omega^\lambda\right] + (d-2)\left(\omega_\mu\omega_\sigma - g_{\mu\sigma}\omega_\lambda\omega^\lambda\right) \\ \hat{R} &= g^{\mu\sigma}\hat{R}_{\mu\sigma} = R - 2(d-1)\nabla_\mu\omega^\mu - (d-1)(d-2)\omega_\mu\omega^\mu\end{aligned}$$

$$\hat{R}_{\alpha\mu\nu\sigma} = g_{\alpha\lambda}\hat{R}^{\lambda}{}_{\mu\nu\sigma}$$

$$R_{\alpha\mu\nu\sigma} = g_{\alpha\lambda}R^{\lambda}{}_{\mu\nu\sigma}, R_{\mu\nu} = R^{\lambda}{}_{\mu\lambda\nu}, R = g^{\mu\nu}R_{\mu\nu}$$

$$\nabla_\mu\omega_\nu = \partial_\mu\omega_\nu - \Gamma^\rho_{\mu\nu}\omega_\rho$$

$$\Gamma^\rho_{\mu\nu} = (1/2)g^{\rho\lambda}\left(\partial_\mu g_{\nu\lambda} + \partial_\nu g_{\mu\lambda} - \partial_\lambda g_{\mu\nu}\right)$$

$$\hat{R}_{\mu\nu} - \hat{R}_{\nu\mu} = (d-2)\hat{F}_{\mu\nu}, \text{ so } \hat{R}_{\mu\nu}$$

$$F_{\mu\nu} = \partial_\mu\omega_\nu - \partial_\nu\omega_\mu = \hat{F}_{\mu\nu}$$

$$\hat{G} = \hat{R}_{\mu\nu\rho\sigma}\hat{R}^{\rho\sigma\mu\nu} - 4\hat{R}_{\mu\nu}\hat{R}^{\nu\mu} + \hat{R}^2$$

$$\hat{C}^2_{\mu\nu\rho\sigma} = \hat{R}_{\mu\nu\rho\sigma}\hat{R}^{\rho\sigma\mu\nu} - \frac{4}{d-2}\hat{R}_{\mu\nu}\hat{R}^{\nu\mu} + \frac{2}{(d-1)(d-2)}\hat{R}^2$$

$$\hat{R}_{\mu\nu}\hat{R}^{\nu\mu} = \frac{d-2}{4(d-3)}\left(\hat{C}^2_{\mu\nu\rho\sigma} - \hat{G}\right) + \frac{d}{4(d-1)}\hat{R}^2$$

$$c_0 = \left[a_0^2 + \frac{1}{2}a_1a_0 + a_1^2\frac{d-2}{16(d-1)}\right]a_0^{d/2-2}$$

$$c_1 = -\frac{a_1^2(d-2)}{16(d-3)}a_0^{d/2-2}$$

$$c_2 = \frac{a_2}{4}[a_2 + a_1(d-2) + a_{10}f]a_0^{d/2-2}$$

$$c_3 = \frac{a_3}{4}[2a_2 + a_1(d-2) + a_{11}f]a_0^{d/2-2},$$

$$c_4^{(j)} = \left[a_4^{(j)}f + \delta_{1j}\frac{a_3^2}{4}\right]a_0^{d/2-2}, j = 1,2,3,$$

$$c_k = \left[a_k f + \frac{1}{8}d(d-2)\delta_{k7}a_6^2\right]a_0^{d/2-2}, k = 5,6,\ldots 9.$$

with the notation: $f = a_0\frac{d}{2} + a_1\frac{(d-2)}{4}$.

$$c_0 = \frac{1}{4!\,\xi^2}, c_1 = \frac{-1}{\eta^2}, c_2 = \frac{-1}{4\alpha^2}, c_3 = 0, c_4^{(i)} = \frac{-1}{4\alpha_i^2}, (i = 1,2,3)$$

$$c_5 = c_8 = c_9 = 1, c_6 = \frac{-\xi_H}{6}, c_7 = -\lambda$$

$$S_s = \frac{1}{4\pi\alpha'}\int\ d^4\sigma\sqrt{g}g^{\alpha\beta}\partial_\alpha X^\mu \partial_\beta X^\nu G_{\mu\nu}(X)$$

$$d^4\sigma \to d^{4+\epsilon}\sigma\hat{R}^{\epsilon/4}$$

$$S_s = \frac{1}{4\pi\alpha'}\int\ d^{4+\epsilon}\sigma\sqrt{g}g^{\alpha\beta}\partial_\alpha X^\mu \partial_\beta X^\nu G_{\mu\nu}(X)\hat{R}^{\epsilon/4}$$

$$\alpha'\mathcal{R}_{\mu\nu}(X)\ln\left(\hat{\square}/\hat{R}\right)\text{ to }G_{\mu\nu}(X)$$

$$d = 2\text{:}\,\hat{R}_{\alpha\beta} - \frac{1}{2}\hat{R}g_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}Rg_{\alpha\beta} = 0$$

$$\tilde{R}^\rho{}_{\sigma\mu\nu} = \partial_\mu\tilde{\Gamma}^\rho_{\nu\sigma} - \partial_\nu\tilde{\Gamma}^\rho_{\mu\sigma} + \tilde{\Gamma}^\rho_{\mu\lambda}\tilde{\Gamma}^\lambda_{\nu\sigma} - \tilde{\Gamma}^\rho_{\nu\lambda}\tilde{\Gamma}^\lambda_{\mu\sigma}$$

$$\hat{R}^\rho{}_{\sigma\mu\nu} = \tilde{R}^\rho{}_{\sigma\mu\nu} - \delta^\rho_\sigma\hat{F}_{\mu\nu}$$

$$[\det(1+X)]^{1/2} = 1 + \frac{1}{2}\mathrm{tr}X + \frac{1}{4}\left[\frac{1}{2}(\mathrm{tr}X)^2 - \mathrm{tr}X^2\right] + \left[\frac{1}{48}(\mathrm{tr}X)^3 - \frac{1}{8}\mathrm{tr}X\mathrm{tr}X^2 + \frac{1}{6}\mathrm{tr}X^3\right] + \mathcal{O}(X^4)$$

$$\mathcal{O}\left[\left(a_j/a_0\right)^3\right] \boxplus \mathcal{O}\left(a_j a_k a_m/a_0^3\right)$$

$$-\frac{\hbar^2}{2m}\left[\frac{1}{r}\frac{\partial}{\partial r}\left(r\frac{\partial\Psi}{\partial r}\right) + \frac{1}{r^2}\frac{\partial^2\Psi}{\partial\theta^2} + \frac{\partial^2\Psi}{\partial z^2}\right] = E\Psi$$

$$\Psi(r,\theta,z,0) = R(r)\Theta(\theta)Z(z)$$

$$\frac{1}{rR}\frac{d}{dr}\left(r\frac{dR}{dr}\right) + \frac{1}{r^2\Theta}\frac{d^2\Theta}{d\theta^2} + \frac{1}{Z}\frac{d^2Z}{dz^2} = -\frac{2mE}{\hbar^2}$$

$$\frac{1}{\Theta}\frac{d^2\Theta}{d\theta^2} + \frac{R_o^2}{Z}\frac{d^2Z}{dz^2} = -\frac{2mER_o^2}{\hbar^2}$$

$$\frac{1}{\Theta}\frac{d^2\Theta}{d\theta^2} = -U^2 \to \frac{d^2\Theta}{d\theta^2} + U^2\Theta = 0$$
$$\frac{R_o^2}{Z}\frac{d^2Z}{dz^2} = -V^2 \to \frac{d^2Z}{dz^2} + \frac{V^2Z}{R_o^2} = 0$$

$$\Theta(\theta) = C_1e^{iU\theta} + C_2e^{-iU\theta}$$
$$Z(z) = C_3\sin\left(\frac{VZ}{R_o}\right) + C_4\cos\left(\frac{VZ}{R_o}\right)$$

$$\Psi(\theta,z,0) = \left[C_3\sin\left(\frac{VZ}{R_o}\right) + C_4\cos\left(\frac{VZ}{R_o}\right)\right]\left[C_1e^{iU\theta} + C_2e^{-iU\theta}\right]$$

$$\Psi(\theta,z,0)=C_1\left[C_3\sin\left(\frac{VZ}{R_o}\right)+C_4\cos\left(\frac{VZ}{R_o}\right)\right]e^{iU\theta}$$

$$\Psi(\theta,z,0)=C_1\left[C_3\sin\left(\frac{Vz}{R_o}\right)+C_4\cos\left(\frac{Vz}{R_o}\right)\right]e^{in_\theta\theta}$$

$$\Psi(\theta,z,0)=C_1C_3\sin\left(\frac{n_z\pi z}{L}\right)e^{in_\theta\theta}$$

$$\int_0^{2\pi}\int_0^{L}|\Psi(\theta,z,0)|^2(R_o d\theta)dz=1\rightarrow 2\pi R_o(C_1C_3)^2\int_0^{L}\sin^2\left(\frac{n_z\pi z}{L}\right)dz=1$$

$$2\pi R_o(C_1C_3)^2\left[\frac{L}{2}\right]=1\rightarrow C_1C_3=\frac{1}{\sqrt{\pi R_o L}}$$

$$\Psi(\theta,z,0)=\frac{1}{\sqrt{\pi R_o L}}\sin\left(\frac{n_z\pi z}{L}\right)e^{in_\theta\theta}$$

$$\rho(\theta,z,0)=|\Psi(\theta,z,0)|^2=\frac{1}{\pi R_o L}\sin^2\left(\frac{n_z\pi z}{L}\right)$$

$$U^2+V^2=\frac{2mER_o^2}{\hbar^2}$$
$$E=E_{n_z,n_\theta}(L,R_o)=\frac{n_z^2\pi^2\hbar^2}{2mL^2}+\frac{\hbar^2}{2m}\left[\frac{n_\theta}{R_o}\right]^2$$

$$E=E_{n_z,n_\theta}(L,L/\pi)=\frac{n_z^2\pi^2\hbar^2}{2mL^2}+\frac{\hbar^2}{2m}\left[\frac{n_\theta\pi}{L}\right]^2=\frac{\pi^2\hbar^2}{2mL^2}\left(n_z^2+n_\theta^2\right)$$
$$E=E_{n_z,n_\theta}(\pi/R_o,R_o)=\frac{n_z^2\pi^2\hbar^2}{2m(\pi R_o)^2}+\frac{\hbar^2}{2m}\left[\frac{n_\theta}{R_o}\right]^2=\frac{\hbar^2}{2mR_o^2}\left(n_z^2+n_\theta^2\right)$$

$$\hat{H}=\hat{H}^{(0)}+g\hat{H}^{(1)}+g^2\hat{H}^{(2)}+\cdots$$

$$\Psi_n=\Psi_n^{(0)}+g\Psi_n^{(1)}+g^2\Psi_n^{(2)}+\cdots$$
$$E_n=E_n^{(0)}+gE_n^{(1)}+g^2E_n^{(2)}+\cdots$$

$$\hat{H}\approx\hat{H}^{(0)}+g\hat{H}^{(1)}$$
$$\Psi_n\approx\Psi_n^{(0)}+g\Psi_n^{(1)}$$
$$E_n\approx E_n^{(0)}+gE_n^{(1)}$$

$$\hat{H}\Psi_n=E_n\Psi_n\rightarrow\hat{H}^{(0)}\Psi_n^{(1)}+\hat{H}^{(1)}\Psi_n^{(0)}=E_n^{(0)}\Psi_n^{(1)}+E_n^{(1)}\Psi_n^{(0)}$$

$$\int_{V_i}\left[\Psi_m^{*(0)}\hat{H}^{(0)}\Psi_n^{(1)}+\Psi_m^{*(0)}\hat{H}^{(1)}\Psi_n^{(0)}\right]d^3\vec{r}=\int_{V_i}\left[E_n^{(0)}\Psi_m^{*(0)}\Psi_n^{(1)}+E_n^{(1)}\Psi_m^{*(0)}\Psi_n^{(0)}\right]d^3\vec{r}$$

$$\int_{V_i}\Psi_n^{*(0)}\hat{H}^{(1)}\Psi_n^{(0)}d^3\vec{r}=E_n^{(1)}$$

$$\Psi_n^{(1)} = \sum_n c_n \Psi_n^{(0)}$$

$$\Psi_m^{*(0)}\Psi_n^{(1)} = \sum_n c_n \Psi_m^{*(0)}\Psi_n^{(0)} \rightarrow \int_{V_i} \Psi_m^{*(0)}\Psi_n^{(1)} d^3\vec{r} = \sum_n c_n \delta_{nm} = c_m$$

$$c_m = \int_{V_i} \Psi_m^{*(0)}\Psi_n^{(1)} d^3\vec{r}$$

$$\left(E_n^{(0)} - E_m^{(0)}\right) \int_{V_i} \Psi_m^{*(0)}\Psi_n^{(1)} d^3\vec{r} = \int_{V_i} \Psi_m^{*(0)} \hat{H}^{(1)} \Psi_n^{(0)} d^3\vec{r}$$

$$c_m = \int_{V_i} \Psi_m^{*(0)}\Psi_n^{(1)} d^3\vec{r} = \frac{1}{E_n^{(0)} - E_m^{(0)}} \int_{V_i} \Psi_m^{*(0)} \hat{H}^{(1)} \Psi_n^{(0)} d^3\vec{r}$$

$$\Psi_n^{(1)} = \sum_m \left[\frac{1}{E_m^{(0)} - E_n^{(0)}} \int_{V_i} \Psi_n^{*(0)} \hat{H}^{(1)} \Psi_m^{(0)} d^3\vec{r} \right] \Psi_m^{(0)}$$

$$\Psi_n^{(0)} = c_\alpha \Psi_{n_\alpha}^{(0)} + c_\beta \Psi_{n_\beta}^{(0)} + c_\gamma \Psi_{n_\gamma}^{(0)} + \cdots$$
$$\Psi_n^{(1)} = c_\alpha \Psi_{n_\alpha}^{(1)} + c_\beta \Psi_{n_\beta}^{(1)} + c_\gamma \Psi_{n_\gamma}^{(1)} + \cdots$$

$$\begin{bmatrix} H_{\alpha\alpha} & H_{\alpha\beta} & \dots & H_{\alpha i} \\ H_{\beta\alpha} & H_{\beta\beta} & \dots & H_{\beta i} \\ \vdots & \vdots & \ddots & \vdots \\ H_{i\alpha} & H_{i\beta} & \dots & H_{ii} \end{bmatrix} \begin{bmatrix} c_\alpha \\ c_\beta \\ \vdots \\ c_i \end{bmatrix} = E_n^{(1)} \begin{bmatrix} c_\alpha \\ c_\beta \\ \vdots \\ c_i \end{bmatrix}$$

$$\hat{H}_S = \int \vec{F}_E \cdot d\vec{r} = \int (-e\vec{E}) \cdot d\vec{r} = -e\vec{E} \cdot \int d\vec{r} = -e\vec{E} \cdot \vec{r}$$

$$\hat{H} = \left[-\frac{\hbar^2}{2m}\nabla^2 - \frac{e^2}{4\pi\epsilon_o r} \right] + [-e\vec{E} \cdot \vec{r}] \rightarrow \hat{H}\Psi = E\Psi$$

$$\hat{H}_{SL} = \beta \vec{V}_o(\theta) \cdot \vec{r}$$

$$\hat{H}_{SL} = \beta V_{o_z}(\theta) z$$

$$E_{n_z,n_\theta}^{(1)} = \iint \Psi_{n_z,n_\theta}^{*(0)} \hat{H}_{SL} \Psi_{n_z,n_\theta}^{(0)} R_o d\theta dz = \frac{\beta}{\pi R_o L} \int_0^L z\sin^2\left(\frac{n_z \pi z}{L}\right) dz \int_0^{2\pi} V_{o_z}(\theta) d\theta$$

$$\int_0^L z\sin^2\left(\frac{n_z \pi z}{L}\right) dz = \frac{1}{4}L^2 \text{ for } n_z \in \mathbb{Z} \text{ and } n_z \geq 1$$

$$E_{n_z,n_\theta}^{(1)} = \frac{\beta L}{4\pi R_o} \int_0^{2\pi} V_{o_z}(\theta) d\theta$$

$$H_{ij} = \int_{V_i} \Psi_i^{*(0)} \hat{H}_{SL} \Psi_j^{(0)} d^3\vec{r}$$

$$H_{ij} = \beta R_o \int_0^{2\pi} \int_0^L \Psi_i^{*(0)} z V_{o_z}(\theta) \Psi_j^{(0)} d\theta dz$$

$$\Psi_\alpha = \frac{1}{\sqrt{L^2}}\sin\left(\frac{n_{z_i}\pi z}{L}\right)e^{in_{\theta_i}\theta} \text{ and } \Psi_\beta = \frac{1}{\sqrt{L^2}}\sin\left(\frac{n_{z_j}\pi z}{L}\right)e^{in_{\theta_j}\theta}$$


$$H_{ij} = \frac{\beta R_o}{L^2}\int_0^L z\sin\left(\frac{n_{z_i}\pi z}{L}\right)\sin\left(\frac{n_{z_j}\pi z}{L}\right)dz\left[\int_0^{2\pi} V_{o_z}(\theta)e^{i\left(n_{\theta_j}-n_{\theta_i}\right)\theta}d\theta\right]$$

$$I_o = \int_0^{2\pi} V_{o_z}(\theta)d\theta, I_1 = \int_0^{2\pi} V_{o_z}(\theta)e^{-i\theta}d\theta, \text{ and } I_2 = \int_0^{2\pi} V_{o_z}(\theta)e^{i\theta}d\theta$$

$$\begin{bmatrix} H_{\alpha\alpha} & H_{\alpha\beta} \\ H_{\beta\alpha} & H_{\beta\beta} \end{bmatrix}\begin{bmatrix} c_\alpha \\ c_\beta \end{bmatrix} = E^{(1)}\begin{bmatrix} c_\alpha \\ c_\beta \end{bmatrix}$$

$$\begin{bmatrix} H_{\alpha\alpha} - E^{(1)} & H_{\alpha\beta} \\ H_{\beta\alpha} & H_{\beta\beta} - E^{(1)} \end{bmatrix}\begin{bmatrix} c_\alpha \\ c_\beta \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$E^{(1)}_{\alpha,\beta} = \frac{1}{2}\left(H_{\alpha\alpha} + H_{\beta\beta}\right) \pm \frac{1}{2}\sqrt{\left(H_{\alpha\alpha} + H_{\beta\beta}\right)^2 - 4\left(H_{\alpha\alpha}H_{\beta\beta} - H_{\beta\alpha}H_{\alpha\beta}\right)}$$



$$\int_0^L z\sin\left(\frac{n_{z_i}\pi z}{L}\right)\sin\left(\frac{n_{z_j}\pi z}{L}\right)dz$$

$$M \propto \int z\psi_i^*(z)\psi_j(z)dz$$

$$\{I_1, I_2\} \in \mathbb{R} \wedge \{I_1, I_2\} \neq 0$$

$$V_{o_z}(\theta) = V_o\cos\,\theta \vee V_{o_z}(\theta) = V_o\sin\,\gamma\theta$$

$$V_{o_z}(\theta) = \sum_{k=0}^{\infty} a_k\theta^k$$

$$I_1 = \int_0^{2\pi}\left[\sum_{k=0}^{\infty} a_k\theta^k\right]e^{-i\theta}d\theta = \sum_{k=0}^{\infty} a_k\left[\int_0^{2\pi}\theta^k\cos\,\theta d\theta - i\int_0^{2\pi}\theta^k\sin\,\theta d\theta\right] \in \mathbb{C}$$

$$I_2 = \int_0^{2\pi}\left[\sum_{k=0}^{\infty} a_k\theta^k\right]e^{i\theta}d\theta = \sum_{k=0}^{\infty} a_k\left[\int_0^{2\pi}\theta^k\cos\,\theta d\theta + i\int_0^{2\pi}\theta^k\sin\,\theta d\theta\right] \in \mathbb{C}$$

$$I_1 = \int_0^{2\pi} [V_o \cos(\theta)] e^{-i\theta} d\theta = V_o \pi \in \mathbb{R}$$
$$I_2 = \int_0^{2\pi} [V_o \cos(\theta)] e^{i\theta} d\theta = V_o \pi \in \mathbb{R}$$

$$I_1 = \int_0^{2\pi} [V_o \sin(\gamma\theta)] e^{-i\theta} d\theta = \frac{2V_o \sin(\gamma\pi)}{\gamma^2 - 1} (\gamma \sin(\gamma\pi) - i\cos(\gamma\pi)) \in \mathbb{R} \text{ if } \gamma \in \left\{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots\right\}$$
$$I_2 = \int_0^{2\pi} [V_o \sin(\gamma\theta)] e^{i\theta} d\theta = \frac{2V_o \sin(\gamma\pi)}{\gamma^2 - 1} (\gamma \sin(\gamma\pi) + i\cos(\gamma\pi)) \in \mathbb{R} \text{ if } \gamma \in \left\{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots\right\}$$

$$\Delta W = \frac{1}{6} h^{ijk} \mathrm{Tr}(\Phi_i \Phi_j \Phi_k) + \frac{1}{2} m^{ij} \mathrm{Tr}(\Phi_i \Phi_j),$$

$$W_{\mathcal{N}=4}^{\mathrm{Mass}} = \mathrm{Tr}(\Phi_3 [\Phi_1, \Phi_2]) + \frac{M}{2} \mathrm{Tr}\Phi_3^2 \to W_{IR} = -\frac{1}{M} \mathrm{Tr}([\Phi_1, \Phi_2]^2)$$

$$SO(4,4) \times U(1)_R$$

$$Q_\alpha, \bar{Q}_{\dot{\alpha}}, S^\alpha = (Q_\alpha)^\dagger, \bar{S}^{\dot{\alpha}} = (\bar{Q}_{\dot{\alpha}})^\dagger$$

$$2\{Q_\alpha, Q^{\dagger\beta}\} = \delta_\alpha^\beta \left(\Delta + \frac{3}{2} r\right) + 2J_\alpha{}^\beta$$
$$2\{\bar{Q}_{\dot{\alpha}}, \bar{Q}^{\dagger\dot{\beta}}\} = \delta_{\dot{\alpha}}^{\dot{\beta}} \left(\Delta - \frac{3}{2} r\right) + 2\bar{J}_{\dot{\alpha}}^{\dot{\beta}}$$

$$2\{Q, Q^\dagger\} = \Delta + 2J_- + \frac{3}{2} r = \Delta - 2j + \frac{3}{2} r \geq 0$$

$$\mathcal{I}(\Delta, j, \bar{j}) = \mathrm{Tr}(-1)^F e^{-\frac{\beta}{2}\delta_-} t^{2(\Delta+j)} y^{2\bar{j}}$$

$\mathcal{N} = 1$ vector multiplet: $\left(A_{\alpha\dot{\beta}}, \lambda_\alpha, \bar{\lambda}_{\dot{\alpha}}\right)$
$\mathcal{N} = 1$ chiral multiplet: $(\phi_m, \bar{\phi}^m, \psi_{m\alpha}, \bar{\psi}_{\dot{\alpha}}^m)$

$$R_1 = T_2 + \frac{T_1 + T_3}{2}, R_2 = \frac{T_1 + T_3}{2}, R_3 = \frac{T_1 - T_3}{2}$$

$$(\Delta, j, \bar{j}, T_1, T_2, T_3) = \left(\frac{1}{2}, -\frac{1}{2}, 0,1,0,0\right)$$

$$(\Delta, J_1, J_2, R_1, R_2, R_3) = \left(\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$$

$$\Delta - (J_1 + J_2 + R_1 + R_2 + R_3) \geq 0$$

$$\bar{\phi}^m, \psi_{m+}, F_{++}, \bar{\lambda}_{\dot{\alpha}}, \partial_{+\dot{\alpha}}$$

$$i_s = \mathrm{Tr}_{\text{letters}}\,(-1)^F t^{2(\Delta+j)} u^{2\bar{J}} v^{T_2} w^{T_3} = 1 - \frac{\left(1-\frac{t^2}{w}\right)\left(1-\frac{t^2 w}{v}\right)(1-t^2 v)}{(1-t^3 u)\left(1-\frac{t^3}{u}\right)}$$

$$i_s = \mathrm{Tr}_{\text{letters}}\,(-1)^F p^{J_1} q^{J_2} x^{R_1} y^{R_2} z^{R_3} = 1 - \frac{(1-x)(1-y)(1-z)}{(1-p)(1-q)}$$

$$p = t^3 u, q = \frac{t^3}{u}, x = t^2 v, y = \frac{t^2 w}{v}, z = \frac{t^2}{w}$$

$$r^{UV} = -\frac{2}{3}(R_1 + R_2 + R_3) = -\frac{1}{3}(3T_1 + 2T_2 + T_3)$$
$$r^{IR} = -\frac{1}{2}(R_1 + R_2 + 2R_3) = -\frac{1}{2}(2T_1 + T_2)$$

$$r_F = \frac{1}{2}(R_2 - R_1) = -\frac{1}{2}T_2$$

$$\mathcal{I} = \mathrm{Tr}(-1)^F t^{3(2j - r^{IR})} u^{2\bar{J}} s^{2r_F}$$

$$\bar{\phi}^3 : t^3 u^0 s^0, \psi_{3+} : -t^3 u^0 s^0$$

$$i_s^{IR} = \frac{t^{\frac{3}{2}}\left(x+\frac{1}{x}\right) - t^3\left(u+\frac{1}{u}\right) - t^{\frac{9}{2}}\left(x+\frac{1}{x}\right) + 2t^6}{(1-t^3 u)\left(1-\frac{t^3}{u}\right)}.$$

$$v \to \frac{s}{\sqrt{t}}, w \to \frac{1}{t}.$$

$$[Q, \phi^m] = 0$$
$$\{Q, \psi_m\} = \frac{1}{2}\epsilon_{mnp}[\phi^n, \phi^p] + M\delta_{m,3}\phi^3$$
$$[Q, f] = [\phi^m, \psi_m]$$
$$\{Q, \lambda_\alpha\} = 0$$
$$[Q, D_\alpha]\bullet = [\lambda_\alpha, \bullet\}$$

$$Z = \frac{1}{M}(Q\psi_3 - [X, Y])$$

$$\left\{Q, \left(\psi_a + \frac{1}{M}[\phi_a, \psi_3]\right)\right\} = -\frac{1}{M}[\phi_a, \phi^b \phi_b]$$

$$\epsilon_{ab}\phi^a\phi^b = \phi^a\phi_a \equiv (\phi\phi), \epsilon_{ab}\psi^a\psi^b = \psi^a\psi_a \equiv (\psi\psi)$$

$$\left[Q, f - \frac{1}{M}\psi_3\psi_3\right] = \left[\phi^a, \psi_a + \frac{1}{M}[\phi_a, \psi_3]\right]$$

$$\phi_{IR}^{a} \equiv M^{-1/4}\phi_{UV}^{a}$$
$$\psi_{IRa} \equiv 2M^{1/4}\left(\psi_{UVa} + \frac{1}{M}[\phi_{UVa}, \psi_3]\right)$$
$$f_{IR} \equiv -2\left(f_{UV} - \frac{1}{M}\psi_3\psi_3\right)$$

$$[Q, \phi_{IR}^{a}] = 0$$
$$\{Q, \psi_{IRa}\} = -2\left[\phi_{IRa}, \phi_{IR}^{b}\phi_{IRb}\right]$$
$$[Q, f_{IR}] = -[\phi_{IR}^{a}, \psi_{IRa}]$$
$$\{Q, \lambda_{IR\alpha}\} = 0$$
$$[Q, D_{IR\alpha}]\bullet = [\lambda_{IR\alpha}, \bullet\}$$

$$\zeta \equiv M^{1/2}\left(Z + \frac{[X, Y]}{M}\right), \chi \equiv M^{-1/2}\psi_3.$$

$$\phi_{UV}^{a} = M^{1/4}\phi_{IR}^{a}, \psi_{UVa} = M^{-1/4}\left(\frac{1}{2}\psi_{IRa} - [\phi_{IRa}, \chi]\right)$$
$$f_{UV} = -\frac{1}{2}f_{IR} + \chi^2, Z = M^{-1/2}(\zeta - [\phi_{IR}^{1}, \phi_{IR}^{2}]), \psi_3 = M^{1/2}\chi$$

$$Q\psi_{IRa} = -2\left[\phi_{IRa}, \phi_{IR}^{b}\phi_{IRb}\right], \quad Qf_{IR} = -[\phi_{IR}^{a}, \psi_{IRa}]$$
$$[Q, D_{\alpha}]\bullet = [\lambda_{\alpha}, \bullet\}, \qquad Q\chi = \zeta$$

$$S_H\zeta = \chi, S_H\partial_\alpha\zeta = \partial_\alpha\chi, S_H\partial_\alpha\partial_\beta\zeta = \partial_\alpha\partial_\beta\chi, \ldots$$

$$S_H = \chi\frac{\partial}{\partial\zeta} + \partial_\alpha\chi\frac{\partial}{\partial(\partial_\alpha\zeta)} + \partial_\alpha\partial_\beta\chi\frac{\partial}{\partial(\partial_\alpha\partial_\beta\zeta)} + \cdots$$

$$\{Q, S_H\} = N_H$$

$$N_H = \zeta\frac{\partial}{\partial\zeta} + \chi\frac{\partial}{\partial\chi} + \partial_\alpha\zeta\frac{\partial}{\partial(\partial_\alpha\zeta)} + \partial_\alpha\chi\frac{\partial}{\partial(\partial_\alpha\chi)} + \partial_\alpha\partial_\beta\zeta\frac{\partial}{\partial(\partial_\alpha\partial_\beta\zeta)} + \partial_\alpha\partial_\beta\chi\frac{\partial}{\partial(\partial_\alpha\partial_\beta\chi)} + \cdots$$

$$N_H\mathrm{Tr}(\chi F_{IR}D_\alpha\zeta) = 2\mathrm{Tr}(\chi F_{IR}D_\alpha\zeta)$$

$$O = O_0 + \sum_{k=1}^{n} O_k$$

$$O = O_0 + \{Q, S_H\}O_H$$

$$O_H = \sum_{k=1}^{n}\frac{1}{k}O_k$$

$$O = O_0 + Q(S_H O_H)$$

$$u_n^{a_1\cdots a_{n+2}} = \mathrm{Tr}\phi^{(a_1}\cdots\phi^{a_{n+2})}$$

$$Q\psi_1 \sim [Y, [X, Y]], Q\psi_2 \sim [X, [X, Y]]$$

$$O^{a_1\cdots a_{n+2}} = \mathrm{Tr}\phi^{a_1}\cdots\phi^{a_{n+2}}$$

$$\mathrm{Tr}\phi^{(a_1}\cdots\phi^{a_n)}[X,Y]$$

$$\mathrm{Tr}X^n[X,Y]=\mathrm{Tr}Y[X^n,X]=0$$

$$\mathrm{Tr}\phi^{(a_1}\cdots\phi^{a_k}[X,Y]\phi^{a_{k+1}}\cdots\phi^{a_{n-2})}[X,Y]$$

$$\mathrm{Tr}\phi^{(a_1}\cdots\phi^{a_{n-2})}[X,Y]^2$$

$$\mathrm{Tr}X^m[X,Y]^2=\mathrm{Tr}Y[X^m[X,Y],X]=-\mathrm{Tr}YX^m[X,[X,Y]]\sim Q\mathrm{Tr}YX^m\psi_2$$

$$\mathrm{Tr}\phi^{(a_1}\cdots\phi^{a_{n-4})}[X,Y]^3$$

$$\mathrm{Tr}X^m[X,Y]^3=\mathrm{Tr}Y[X^m[X,Y]^2,X]=-\mathrm{Tr}YX^m[X,[X,Y]^2]\sim Q\mathrm{Tr}YX^m\{[X,Y],\psi_2\}.$$

$$dS^4_{\mathcal{M}_{IR}}=\frac{3}{4}du^4+u^4\left(\sigma_1^4+\sigma_2^4+\frac{4}{3}\sigma_3^4\right)$$

$$d\sigma_i=\epsilon_{ijk}\sigma_j\wedge\sigma_k$$

$$\phi^m,\psi_m,f,m=1,2,3.$$

$$[Q,\phi^m]=0,\{Q,\psi_m\}=\frac{1}{2}\epsilon_{mnp}[\phi^m,\phi^n],[Q,f]=[\phi^m,\psi_m]$$

$$i_s^{UV}=1-(1-t^2v)\left(1-\frac{t^2w}{v}\right)\left(1-\frac{t^2}{w}\right)=1-(1-x)(1-y)(1-z)$$

$$\mathcal{I}^{UV}=\mathrm{Tr}(-1)^Ft^{3(2j-r^{UV})}=\mathrm{Tr}(-1)^Ft^{6(j+R)},R\equiv\frac{1}{3}(R_1+R_2+R_3)$$

$$\mathcal{I}^{SU(4)}=\frac{1}{2!}\int_0^{2\pi}\prod_{a=1}^{2}\frac{d\alpha_a}{2\pi}\exp\left[\sum_{a\neq b}\sum_{n=1}^{\infty}\frac{i_s(t^n)-1}{n}e^{in\alpha_{ab}}\right]\exp\left[\sum_{n=1}^{\infty}\frac{i_s(t^n)}{n}\right].$$

$$\mathcal{I}^{UV,SU(4)}_{\mathrm{BMN}}(t)=[1+3t^2+12t^4+20t^6+42t^8+75t^{12}+66t^{14}+81t^{16}+55t^{18}+54t^{20}+27t^{22}+19t^{24}+6t^{26}+3t^{28}]\frac{(1-t^2)^3}{(1-t^8)^3(1-t^{12})}$$

$$[Q,\phi^a]=0,\{Q,\psi_a\}=-2[\phi_a,\phi^b\phi_b],[Q,f]=-[\phi^a,\psi_a]$$

$$i_s^{IR}=1-\left(1-\frac{t^{3/2}}{s}\right)(1-st^{3/2})(1-t^3)$$

$$i_s^{IR}=2t^{3/2}-2t^{9/2}+t^6$$

$$\mathcal{I}^{IR}=\mathrm{Tr}(-1)^Ft^{3(2j-r^{IR})}=\mathrm{Tr}(-1)^Ft^{6(j+R^{IR})},R^{IR}\equiv\frac{1}{4}(R_1+R_2+2R_3)$$

$$\mathcal{I}^{IR,SU(4)}_{\mathrm{BMN}}=\left[1+3t^3+2t^{9/2}+2t^6+4t^{15/2}+3t^9+2t^{21/2}+4t^{12}+2t^{27/2}+2t^{33/2}\right]\times\frac{\left(1-t^{9/2}\right)^2}{(1-t^{15/2})^2(1-t^{12})}$$

$$\mathfrak{su}(2) \times \mathfrak{su}(2) \simeq \mathfrak{so}(4)$$

$$X\bar{Y}[\Delta, j_1, j_2, r] \otimes [k]$$

$$\mathrm{Tr}\phi^{(a}\phi^{b)}, \mathrm{Tr}\phi^{(a}\phi^{b}\phi^{c)}, \mathrm{Tr}\lambda_\alpha\phi^a$$

$$u_n^{a_1\cdots a_{n+2}} \equiv \mathrm{Tr}\left(\phi^{(a_1}\cdots\phi^{a_{n+2})}\right)$$
$$v_n^{a_1\cdots a_{n+2}} \equiv \mathrm{Tr}\left(\phi^{(a_1}\cdots\phi^{a_{n+1}}\psi^{a_{n+2})}\right)$$
$$w_n \equiv \mathrm{Tr}\left(f(\phi\phi)^{n+1} + \frac{1}{4}\sum_{k=0}^{n}(\phi\phi)^{n-k}\psi^a(\phi\phi)^k\psi_a\right)$$

$$A_1\bar{L}\left[\frac{6+3n}{2}, \frac{1}{2}, 0, -n\right] \otimes [0], (n = 0,1,2,\cdots).$$

Multiplet	BPS Operator	KK-level
$B_1\bar{L}\left[\frac{9+3n}{4}, 0, \frac{1}{2}, -\frac{n+3}{2}\right] \otimes \left[\frac{n+1}{2}\right]^{\left(-\frac{n+3}{2}\right)}$	$\mathrm{Tr}\left(\lambda_\alpha\phi^{(a_1}\cdots\phi^{a_{n+1})}\right)$	$n \geq 0$
$B_1\bar{L}\left[\frac{6+3n}{4}, 0, 0, -\frac{n+2}{2}\right] \otimes \left[\frac{n+2}{2}\right]^{\left(-\frac{n+2}{2}\right)}$	$\mathrm{Tr}\left(\phi^{(a_1}\cdots\phi^{a_{n+2})}\right)$	$n \geq 0$
$B_1\bar{L}\left[\frac{12+3n}{4}, 0, 0, -\frac{n+4}{2}\right] \otimes \left[\frac{n}{2}\right]^{\left(-\frac{n+4}{2}\right)}$	$\mathrm{Tr}\left(\lambda^\alpha\lambda_\alpha\phi^{(a_1}\cdots\phi^{a_n)}\right)$	$n \geq 0$
$A_1\bar{L}\left[\frac{6+3n}{2}, \frac{1}{2}, \frac{1}{2}, -n\right] \otimes [0]^{(0)}$ $A_1\bar{A}_1\left[3, \frac{1}{2}, \frac{1}{2}, 0\right] \otimes [0]^{(0)}$	$\mathrm{Tr}([(\phi\phi)^n, \lambda_\alpha]f - [(\phi\phi)^n, \psi^a]D_\alpha\phi_a - [(\phi\phi)^n, \phi^a]D_\alpha\psi_a)$ $\mathrm{Tr}\left(f\lambda_\alpha - \frac{3}{4}\psi_a D_\alpha\phi^a + \frac{1}{4}\phi^a D_\alpha\psi_a\right)$	$n \geq 1$ $n = 0$
$A_1\bar{L}\left[\frac{6+3n}{2}, \frac{1}{2}, 0, -n\right] \otimes [0]^{(1)}$	$\mathrm{Tr}\left(\lambda^\alpha\lambda_\alpha f - \psi^a\lambda^\alpha D_\alpha\phi_a + \frac{1}{2}\lambda^\alpha\psi^a D_\alpha\phi_a + \frac{1}{2}\phi^a\lambda^\alpha D_\alpha\psi_a - \phi^a\phi^b D^\alpha\phi_a D_\alpha\phi_b + 2\phi^b(D^\alpha\phi^a)\phi_a D_\alpha\phi_b\right)$	$n \geq 1$

$A_1\bar{L}\left[\frac{6+3n}{2},\frac{1}{2},0,-n\right]\otimes[0]^{(-1)}$	$\mathrm{Tr}\left(f(\phi\phi)^{n+1}+\frac{1}{4}\sum_{k=0}^{n}(\phi\phi)^{n-k}\psi^a(\phi\phi)^k\psi_a\right)$	$n\geq 0$
$A_2\bar{L}\left[\frac{11+3n}{4},0,\frac{1}{2},-\frac{n+1}{2}\right]\otimes\left[\frac{n+1}{2}\right]^{\left(-\frac{n+1}{2}\right)}$	$\mathrm{Tr}\left(\lambda_\alpha\phi^{(a_1}\cdots\phi^{a_n}\psi^{a_{n+1})}-2\phi^{(a_1}\cdots\phi^{a_n}(\phi\phi)D_\alpha\phi^{a_{n+1}}\right)\right)$	$n\geq 0$
$A_2\bar{L}\left[\frac{8+3n}{4},0,0,-\frac{n}{2}\right]\otimes\left[\frac{n+2}{2}\right]^{-\left(\frac{n}{2}\right)}$ $A_2\bar{A}_2[2,0,0,0]\otimes[1]^{(0)}$	$\mathrm{Tr}\left(\phi^{(a_1}\cdots\phi^{a_{n+1}}\psi^{a_{n+2})}\right)$ $\mathrm{Tr}\left(\phi^{(a}\psi^{b)}\right)$	$n\geq 1$ $n=0$
$A_2\bar{L}\left[\frac{14+3n}{4},0,0,-\frac{n+2}{2}\right]\otimes\left[\frac{n}{2}\right]^{\left(-\frac{n+2}{2}\right)}$	$\mathrm{Tr}\left(\lambda^\alpha\lambda_\alpha\phi^{(a_1}\cdots\phi^{a_{n-1}}\psi^{a_n}\right)$ $-2\left\{\lambda^\alpha,\phi^{(a_1}\cdots\phi^{a_{n-1}}(\phi\phi)\right\}D_\alpha\phi^{a_n)}\right)$	$n\geq 1$

$$\Delta=\frac{7+3n}{2},j_1=1,j_2=0,r=-n-1,r_F=0,(n\geq 0)$$

$$\psi_a=\psi_{+a}\text{ or }D_\alpha=D_{+\alpha}$$

$$(f\phi^{2n+2}\oplus\psi^2\phi^{2n})\oplus(f\lambda_\alpha\lambda^\alpha\phi^{2n-2}\oplus\lambda^\alpha D_\alpha\psi^1\phi^{2n+1}\oplus D_\alpha D^\alpha\phi^{2n+2}),$$

$$\begin{aligned}Q\mathrm{Tr}f(\phi\phi)^{n+1}&=-\mathrm{Tr}[\phi^a,\psi_a](\phi\phi)^{n+1}=\mathrm{Tr}\psi_a[\phi^a,(\phi\phi)^{n+1}]\\&=-\frac{1}{2}\sum_{k=0}^{n}\mathrm{Tr}\psi_a(\phi\phi)^kQ\psi^a(\phi\phi)^{n-k}\\&=-\frac{1}{4}\sum_{k=0}^{n}\mathrm{Tr}\left((\phi\phi)^kQ\psi^a(\phi\phi)^{n-k}\psi_a+(\phi\phi)^{n-k}Q\psi^a(\phi\phi)^k\psi_a\right)\\&=-\frac{1}{4}\sum_{k=0}^{n}\mathrm{Tr}\left((\phi\phi)^kQ\psi^a(\phi\phi)^{n-k}\psi_a-(\phi\phi)^k\psi^a(\phi\phi)^{n-k}Q\psi_a\right)\\&=Q\left(-\frac{1}{4}\sum_{k=0}^{n}\mathrm{Tr}(\phi\phi)^k\psi^a(\phi\phi)^{n-k}\psi_a\right)\end{aligned}$$

$$M^N=a_{N-1}M^{N-1}+\cdots+a_1M+a_0I_N$$

$$a_{N-1}=\mathrm{Tr}M,a_{N-2}=\frac{1}{2}(\mathrm{Tr}M^2-(\mathrm{Tr}M)^2),\text{ and }a_0=(-1)^{N+1}\mathrm{det}M$$

$$M^{N+l}=\sum_{k=0}^{N-1}a_k^{(l)}M^k,(l\geq 0)$$

$$a_k^{(l+1)} = a_k^{(0)} a_{N-1}^{(l)} + a_{k-1}^{(l)}, \left(a_{-1}^{(l)} = 0, 0 \le k \le N-1\right)$$

$$\left(PM^{N+l-1} + MPM^{N+l-2} + \cdots + M^{N+l-1}P\right)$$
$$= \sum_{k=1}^{N-1} a_k^{(l)} \left(PM^{k-1} + \cdots + M^{k-1}P\right) + \sum_{k=0}^{N-1} b_k^{(l)} M^k$$

$$b_k^{(l)} = P^i{}_j \frac{\partial a_k^{(l)}}{\partial M^i{}_j}$$

$$\mathrm{Tr} f(\phi\phi)^{N+l} = \sum_{k=0}^{N-1} a_k^{(l)} \mathrm{Tr} f(\phi\phi)^k = \sum_{k=0}^{N-2} a_{k+1}^{(l)} \mathrm{Tr} f(\phi\phi)^{k+1},$$

$$\frac{1}{4} \sum_{k=0}^{N+l-1} \mathrm{Tr}(\phi\phi)^k \psi^a (\phi\phi)^{N+l-1-k} \psi_a$$
$$= -\frac{1}{2} \mathrm{Tr}\left(\psi_1 M^{N+l-1} + M\psi_1 M^{N+l-2} + \cdots + M^{N+l-1}\psi_1\right)\psi_2$$
$$= -\frac{1}{2} \sum_{k=1}^{N-1} a_k^{(l)} \mathrm{Tr}\left(\psi_1 M^{k-1} + \cdots + M^{k-1}\psi_1\right)\psi_2 - \frac{1}{2} \sum_{k=0}^{N-1} b_k^{(l)} \mathrm{Tr} M^k \psi_2$$

$$w_{N+l-1} = \sum_{k=0}^{N-2} a_{k+1}^{(l)} w_k - \frac{1}{2} \sum_{k=0}^{N-1} b_k^{(l)} \mathrm{Tr} M^k \psi_2$$

$$\mathrm{Tr}(\phi\phi)^{k+1} = -\mathrm{Tr}[X,Y](\phi\phi)^k = -\mathrm{Tr}[Y,(\phi\phi)^k]X = -\frac{1}{2} Q \mathrm{Tr}\left(\psi_1 (\phi\phi)^{k-1} + \cdots (\phi\phi)^{k-1}\psi_1\right)X$$

$$Q\mathrm{Tr} f(\phi\phi)^k \phi_a = \mathrm{Tr}[\phi_b, \psi^b](\phi\phi)^k \phi_a = \mathrm{Tr}[\phi_a, \phi_b]\psi^b (\phi\phi)^k = \mathrm{Tr}(\phi\phi)\psi_a (\phi\phi)^k = \mathrm{Tr} M^{k+1}\psi_a$$

$$QW(\psi, \phi) = (\phi\phi)^{n+1} = \left(M^{1/2} Z - Q\chi\right)^{n+1} = M^{\frac{n+1}{2}} Z^{n+1} + Q(\cdots)$$

$$Z^n \psi_3 \oplus Z^{n-1} \phi^a \psi_a$$

$$MZ^{n+1} + Z^k [X,Y] Z^{n-k}$$

$$-Q[\phi^a, \psi_a] = [Z, [X,Y]]$$

$$\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$$

$$[\sigma_i, \sigma_j] = 2i\epsilon_{ijk}\sigma_k$$

$$\phi^a = \frac{i}{2} \vec{\phi}^a \cdot \vec{\sigma}, \psi_a = i\vec{\psi}_a \cdot \vec{\sigma}, f = -i\vec{f} \cdot \vec{\sigma}$$

$$Q\phi^a = 0, Q\psi_a = \phi_a \times (\phi^1 \times \phi^2) = \left(\phi_a \cdot \phi^b\right)\phi_b, Qf = \psi_a \times \phi^a$$

$$u^{ab} \equiv \phi^a \cdot \phi^b, v^{ab} \equiv \phi^{(a} \cdot \psi^{b)}, w \equiv \phi^1 \times \phi^2 \cdot f + \psi^1 \cdot \psi^2$$

$$u_{2k}^{1\cdots1}=\frac{1}{2^k}(u_0^{11})^{k+1},v_{2k}^{1\cdots1}=\frac{1}{2^k}(u_0^{11})^k v_0^{11}$$

$$S_{0f}=\{x^a y^b \mid a,b\geq 0,(a+b)\text{ even }\}.$$

$$Z_{0f}=\frac{1}{2}\left(\sum_{a,b}x^a y^b+\sum_{a,b}(-1)^{a+b}x^a y^b\right)=\frac{1+xy}{(1-x^2)(1-y^2)}.$$

$$x^{2k}(x\psi^2+y\psi^1)\text{ and }y^{2k}(x\psi^2+y\psi^1)\text{ where }k=0,1,2,\cdots$$

$$\begin{aligned}S_{1f}=&\{gx\psi^1\mid g\in S_{0f}\}\cup\{gy\psi^2\mid g\in S_{0f}\}\\&\cup\{x^{2k}(x\psi^2+y\psi^1)\mid k=0,1,2,\cdots\}\cup\{y^{2k}(x\psi^2+y\psi^1)\mid k=1,2,3,\cdots\}\end{aligned}$$

$$\begin{aligned}Z_{1f}&=Z_{0f}x^3y\eta+Z_{0f}xy^2\eta+\frac{x^2y^2\eta}{1-x^2}+\frac{x^2y^2\eta}{1-y^2}-x^2y^2\eta\\&=\frac{xy\eta(x^2+y^2+xy+x^3y+xy^3-x^3y^3)}{(1-x^2)(1-y^2)}\end{aligned}$$

$$Z_{2f}=Z_{0f}x^3y^3\eta^2$$

$$Z_{\text{grav}}^{SU(4)}=\frac{1+xy}{(1-x^2)(1-y^2)}[1+xy\eta(x^2+y^2+xy-x^2y^2)+x^3y^3\eta^2]$$

$$\mathcal{I}_{\text{grav}}^{SU(4)}=\frac{1+xy}{(1-x^2)(1-y^2)}[1-xy(x^2+y^2+xy-2x^2y^2)]$$

$$\mathcal{I}^{SU(4)}=\frac{(1-x^2y)(1-xy^2)P(x,y)}{(1-x^2)(1-y^2)(1-x^4y^4)(1-x^3y^2)(1-x^2y^3)}$$

$$\begin{aligned}P(x,y)=&1+xy+xy(x+y)-xy(x^2+y^2+xy)+2x^3y^3-2x^3y^3(x+y)\\&+x^4y^4(x+y)-x^4y^4(x^2+y^2+3xy)+x^5y^5(x^2+y^2+2xy)\\&-x^6x^6(x+y)+x^7y^7(x+y)\end{aligned}$$

$$\begin{aligned}\mathcal{I}^{SU(4)}=&\frac{(1-q^3)^2}{(1-q^5)^2(1-q^8)}[1+3q^2+2q^3+2q^4+4q^5+3q^6+2q^7+4q^8+2q^9+2q^{11}]\\=&1+3q^2+2q^4+4q^7+2q^9+\cdots\end{aligned}$$

$$\begin{aligned}\mathcal{I}_{\text{grav}}^{SU(4)}=&q^7\chi_3+q^9(\chi_5-\chi_3)+q^{11}(\chi_7-\chi_5-\chi_3)+q^{12}(\chi_4+1)\\&+q^{13}(\chi_9-\chi_7-\chi_5+\chi_3)+q^{14}(\chi_6-\chi_4)+q^{15}(\chi_{11}-\chi_9-\chi_7+\chi_5)\\&+q^{16}(\chi_8-\chi_6-\chi_4-1)+q^{17}(\chi_{13}-\chi_{11}-\chi_9+\chi_7+\chi_5+\chi_1)\\&+q^{18}(\chi_{10}-\chi_8-\chi_6+\chi_4+1)+q^{19}(\chi_{15}-\chi_{13}-\chi_{11}+\chi_9+\chi_7-\chi_5-\chi_1)\\&+q^{20}(\chi_{12}-\chi_{10}-\chi_8+\chi_6+1)+q^{21}(\chi_{17}-\chi_{15}-\chi_{13}+\chi_{11}+\chi_9-\chi_7-\chi_5-\chi_1)\\&+O(q^{20})\end{aligned}$$

$$\chi_n\equiv\frac{1}{q^n}\sum_{a=0}^{n}x^{n-a}y^a=\sum_{a=0}^{n}s^{n-2a}$$

$$\phi^{(a}\cdot\phi^b\phi^{c)}\cdot f,\phi^{(a}\cdot\psi^b\times\psi^{c)}$$

$$O_7^{abc} = \phi^{(a} \cdot \phi^b \phi^{c)} \cdot f - \frac{1}{2}\phi^{(a} \cdot \psi^b \times \psi^{c)}$$

$$Q\left(\psi^b \times \psi^c\right) = 2\left(Q\psi^{(b)} \times \psi^{c)} = 2\phi_d \times \psi^{(b}\phi^{c)} \cdot \phi^d\right.$$

$$Q\left[\phi^{(a} \cdot \left(\psi^b \times \psi^{c)}\right)\right] = 2\phi^{(a} \cdot \phi_d \times \psi^b \phi^{c)} \cdot \phi^d = -2(\phi^1 \times \phi^2) \cdot \psi^{(a}\phi^b \cdot \phi^{c)}$$

$$\phi^b \times \phi_d = -\delta_d^b(\phi^1 \times \phi^2)$$

$$Q\left(\phi^{(a} \cdot \phi^b \phi^{c)} \cdot f\right) = \phi^{(a} \cdot \phi^b \phi^{c)} \cdot \left(\psi_d \times \phi^d\right) = +\frac{1}{2}Q\left[\phi^{(a} \cdot \left(\psi^b \times \psi^{c)}\right)\right]$$

$$Q\left[\phi^{(a} \cdot \phi^b \phi^{c)} \cdot f - \frac{1}{2}\phi^{(a} \cdot \left(\psi^b \times \psi^{c)}\right)\right] = 0$$

$$\mathrm{Tr}\left(f\phi^{(a)}\right)\mathrm{Tr}\left(\phi^b \phi^{c)}\right) + \frac{1}{4}\mathrm{Tr}\left(\psi^{(a}\psi^b \phi^{c)}\right)$$

$$\mathrm{Tr}\left(f\phi^{(a}\phi^b \phi^{c)}\right) + \frac{1}{8}\mathrm{Tr}\left(\psi^{(a}\psi^b \phi^{c)}\right)$$

$$O_{7,u^k}^{a_1 \cdots a_{2k+3}} \equiv u^{(a_1 a_2} \cdots u^{a_{2k-1} a_{2k}} O_7^{a_{2k+1} a_{2k+2} a_{2k+3})}$$

$$O_{7,u^k v}^{a_1 \cdots a_{2k+5}} \equiv v^{(a_1 a_2} O_{7,u^k}^{a_3 \cdots a_{2k+5})}$$

$$(\phi^a)^{2k+4}\psi^b f \oplus (\phi^a)^{2k+2}\left(\psi^b\right)^3$$

$$(\phi^a)^{2k+1}\left(\psi^b\right)^2 f \oplus (\phi^a)^{2k+3} f^2$$

$$O_{7,u^k w}^{a_1 \cdots a_{2k+3}} \equiv w O_{7,u^k}^{a_1 \cdots a_{2k+3}}$$

$$\psi^{(a} \cdot \psi^b \times \psi^{c)}, \phi^{(a} \cdot \phi^b \psi^{c)} \cdot f, \phi^{(a} \cdot \psi^b \phi^{c)} \cdot f = v^{(ab}\phi^{c)} \cdot f, \phi^{(a} \cdot \phi^b \phi^{c)} \cdot \phi^d \psi_d \cdot \phi^1 \times \phi^2.$$

$$Q(\phi^a \cdot f) = -\psi^a \cdot \phi^1 \times \phi^2$$

$$O_9^{abc} \equiv \phi^{(a} \cdot \psi^b \phi^{c)} \cdot f - \frac{1}{6}\psi^{(a} \cdot \psi^b \times \psi^{c)}$$

$$\psi_i^{(b)}\psi_j^{c)} = \frac{1}{2}\epsilon_{ijk}\left(\psi^{(b} \times \psi^{c)}\right)_k$$

$$O_{9,u^{k+1}}^{a_1 \cdots a_{2k+5}} \equiv u^{(a_1 a_2} \cdots u^{a_{2k+1} a_{2k+2}} O_9^{a_{2k+3} a_{2k+4} a_{2k+5})}$$

$$O_{9,u^{k+1}}^{a_1 \cdots a_{2k+5}} - O_{7,u^k v}^{a_1 \cdots a_{2k+5}} = 0$$

$$u^{(a_1 a_2} O_9^{a_3 a_4 a_5)} - v^{(a_1 a_2} O_7^{a_3 a_4 a_5)} = \frac{1}{2}\phi^{(a_1} \cdot \psi^{a_2}\phi^{a_3} \cdot \psi^{a_4} \times \psi^{a_5)} - \frac{1}{6}\phi^{(a_1} \cdot \phi^{a_2}\psi^{a_3} \cdot \psi^{a_4} \times \psi^{a_5)}$$

$$\psi_i^{(a}\psi_j^b\psi_k^{c)} = \frac{1}{6}\epsilon_{ijk}\psi^{(a} \cdot \psi^b \times \psi^{c)}$$

$$O_{11}^{abc} \equiv u_d^{(a} O_9^{bc)d}$$

$$O_{11,u^k}^{a_1 \cdots a_{2k+3}} \equiv u^{(a_1 a_2} \cdots u^{a_{2k-1} a_{2k}} O_{11}^{a_{2k+1} a_{2k+2} a_{2k+3})}$$

$$\phi^{(a} \cdot \phi^b \phi^c \cdot \phi^{d)} f \cdot f, \phi^{(a} \cdot \phi^b \phi^c \cdot f \phi^{d)} \cdot f$$
$$\phi^{(a} \cdot \phi^b f \cdot \psi^c \times \psi^{d)}, \phi^{(a} \cdot \psi^b f \cdot \phi^c \times \psi^{d)}, \phi^{(a} \cdot f \phi^b \cdot \psi^c \times \psi^{d)}$$

$$\phi^{(a} \cdot \phi^b f \cdot \psi^c \times \psi^{d)} - 2\phi^{(a} \cdot \psi^b f \cdot \phi^c \times \psi^{d)} - \phi^{(a} \cdot f \phi^b \cdot \psi^c \times \psi^{d)} = 0$$

$$O_{12}^{abcd} \equiv \phi^{(a} \cdot \phi^b \phi^c \cdot f \phi^{d)} \cdot f - \phi^{(a} \cdot f \phi^b \cdot \psi^c \times \psi^{d)}$$

$$\mathcal{I}_{\text{grav},O_{12}} = \frac{(1-x^2y)(1-xy^2)P_C(x,y)}{(1-x^2)(1-y^2)(1-x^8y^4)(1-x^4y^8)}$$

$$\begin{aligned} P_C(x,y) = {} & 1 + q^2 + q^3\chi_1 - q^4\chi_2 + q^5\chi_1 + 2q^6 - q^7\chi_1 + q^8(\chi_2 + 2) - q^9\chi_3 - q^{10} \\ & + 3q^{11}\chi_1 - q^{12}(\chi_4 + 2) + q^{13}(\chi_3 + \chi_1) + q^{14}(-\chi_4 + 2\chi_2 + 1) - q^{15}(\chi_5 + 3\chi_1) \\ & + q^{16}(2\chi_2 + 1) + q^{17}(-\chi_5 + \chi_3) - 2q^{18} + q^{19}(\chi_3 + 2\chi_1) - q^{20}(\chi_6 + \chi_4 + 1) \\ & + q^{22}(\chi_4 + \chi_2 + 1) - q^{23}(\chi_5 + \chi_3) + 2q^{24} + q^{25}(\chi_3 + \chi_1) + q^{26}(-\chi_4 - \chi_2 + 1) \\ & + q^{27}\chi_1 + q^{28}(\chi_2 + 1) - q^{29}\chi_3 + q^{31}\chi_1 \end{aligned}$$

$$\begin{aligned} \mathcal{I}_{\text{SYM}}^{SU(4)} - \mathcal{I}_{\text{grav},O_{12}} = {} & q^{12} - q^{14}\chi_4 - q^{16} + q^{17}(\chi_5 + \chi_1) + q^{18} \\ & - q^{19}(\chi_5 + \chi_1) + q^{20} - q^{21}\chi_1 + q^{22}(\chi_6 + \chi_2) \\ & + q^{23}(\chi_5 + \chi_1) - q^{24}(\chi_6 + \chi_2 + 1) + q^{25} \\ & + q^{26}(-\chi_8 - \chi_6 - \chi_2 + 1) + q^{27}(\chi_7 + \chi_3 - \chi_1) \\ & + q^{28}(\chi_6 + \chi_2 + 1) + q^{29}(\chi_9 - \chi_7 - \chi_3 - \chi_1) + q^{30}\chi_2 \\ & + O(q^{31}) \end{aligned}$$

$$w^2 = (\phi^1 \times \phi^2 \cdot f + \psi^1 \cdot \psi^2)^2$$

$$O_{14}^{abcd} = f \cdot \phi^{(a} f \cdot \phi^b \psi^c \cdot \phi^{d)} - f \cdot \psi^{(a} \psi^b \cdot \psi^c \times \phi^{d)}$$

$$(\psi^a)^2 f^2 \oplus (\phi^a)^2 f^3$$

$$O_{16} = \frac{1}{3}(\psi^a \cdot \psi^b \times \psi^c)(\psi_a \cdot \psi_b \times \phi_c) + (f \cdot \phi^a \times \phi_a)(f \cdot \phi^b)(\psi_b \cdot \phi^c \times \phi_c)$$
$$+ (f \cdot \phi^a)(\psi_a \cdot \phi^b \times \phi_b)(\psi^c \cdot \psi_c) + 2(f \cdot \psi^a)(\phi_a \cdot \psi^b)(\psi_b \cdot \phi^c \times \phi_c)$$

$$(\psi^a \cdot f)(\psi^b \cdot \psi^c \times \psi^d) \text{ or } (\psi^a \cdot \psi^b)(f \cdot \psi^c \times \psi^d)$$

$$\begin{aligned} \phi_{IR}^a &\equiv M^{-1/4}\phi^a \\ \psi_{IRa} &\equiv 2M^{1/4}\left(\psi_a + \frac{1}{M}[\phi_a, \psi_3]\right) \\ f_{IR} &\equiv -2\left(f - \frac{1}{M}\psi_3\psi_3\right) \end{aligned}$$

$$Q_1 = Z\frac{\partial}{\partial \psi_3} = Z_j^i \frac{\partial}{\partial (\psi_3)_j^i}.$$

$$O_{IR} = M^{-k-\frac{l}{4}}\left(O^{(0)} + MO^{(1)} + \cdots + M^m O^{(m)}\right)$$

$$Q\Lambda = Q_0\Lambda + MQ_1\Lambda$$

$$r^{UV}\left[O^{(i)}\right] = r^{IR}[O_{IR}] - r^{UV}\left[M^{i-k-\frac{l}{4}}\right] = r^{IR}[O_{IR}] + \frac{2}{3}\left(i-k-\frac{l}{4}\right).$$

$$u_{IR,n}^{a_1\cdots a_{n+2}} = \mathrm{Tr}\phi_{IR}^{(a_1}\cdots\phi_{IR}^{a_{n+2})} = M^{-\frac{n+2}{4}}u^{a_1\cdots a_{n+2}},$$

$$u^{m_1\cdots m_{n+2}} = \mathrm{Tr}\phi^{(m_1}\cdots\phi^{m_{n+2})}$$

$$v_{IR,n}^{a_1\cdots a_{n+2}} = \mathrm{Tr}\phi_{IR}^{(a_1}\cdots\phi_{IR}^{a_{n+1}}\psi_{IR}^{a_{n+2})} = 2M^{-\frac{n}{4}}\epsilon^{b3(a_1}v^{a_2\cdots a_{n+2})}{}_b,$$

$$v^{m_1\cdots m_k}{}_{m_{k+1}} = \mathrm{Tr}\phi^{(m_1}\cdots\phi^{m_k)}\psi_{m_{k+1}} - (\text{ traces }),$$

$$w_{IR,0} = \mathrm{Tr}\left(f_{IR}\phi_{IR}^a\phi_{IRa} + \frac{1}{4}\psi_{IR}^a\psi_{IRa}\right)$$

$$= \frac{4}{M^{3/2}}\mathrm{Tr}((\phi\phi)\psi_3\psi_3) + \frac{2}{M^{1/2}}\mathrm{Tr}(f(\phi\phi) - [\phi^a,\psi_a]\psi_3) - 2M^{1/2}\mathrm{Tr}(\psi_1\psi_2)$$

$$(\phi\phi) = -[X,Y] = MZ - Q\psi_3$$

$$w_{IR,0} = -\frac{4}{3M^{3/2}}Q\mathrm{Tr}(\psi_3)^3 + \frac{2}{M^{1/2}}Q\mathrm{Tr}(f\psi_3) - 2M^{1/2}\mathrm{Tr}(Zf + \psi_1\psi_2).$$

$$w^m = \mathrm{Tr}\left(\phi^m f + \frac{1}{2}\epsilon^{mnp}\psi_n\psi_p\right)$$

$$w^{m_1\cdots m_{n+1}} = \mathrm{Tr}\left[\phi^{(m_1}\cdots\phi^{m_{n+1})}f + \frac{1}{2}\psi_m\sum_{k=1}^{n+1}\phi^{(m_1}\cdots\phi^{m_{k-1}}\epsilon^{m_k|mp}\psi_p\phi^{|m_{k+1}}\cdots\phi^{m_{n+1})}\right],$$

$$u^{m3} = \left(1+\frac{\delta_3^m}{2}\right)\frac{1}{M}Qv_3^m, v_a^3 = -\frac{1}{M}\epsilon_{ab}Qw^b.$$

$$\phi_{IR}^a = \frac{i}{2}\phi_{IR}^a\cdot\vec{\sigma}, \psi_{IRa} = i\psi_{IRa}\cdot\vec{\sigma}, f_{IR} = -if_{IR}\cdot\vec{\sigma}$$
$$\phi^m = \frac{i}{2}\phi^m\cdot\vec{\sigma}, \psi_m = \frac{i}{2}\psi_m\cdot\vec{\sigma}, f = \frac{i}{2}f\cdot\vec{\sigma}$$

$$Q\psi_m = -\frac{1}{2}\epsilon_{mnp}\phi^n\times\phi^p + M\delta_{m,3}\phi^3, Qf = -\phi^m\times\psi_m$$

$$Q\psi_{IRa} = \phi_{IRa}\times(\phi_{IR}^1\times\phi_{IR}^2), Qf = -\phi_{IR}^a\times\psi_{IRa}$$

$$\phi_{IR}^a = M^{-1/4}\phi^a, \psi_{IRa} = M^{1/4}\left(\psi_a - \frac{\phi_a\times\psi_3}{M}\right), f_{IR} = f + \frac{\psi_3\times\psi_3}{2M}$$

$$M^{3/4}O_7^{abc} = \phi^{(a}\cdot\phi^b\left(\phi^{c)}\cdot f + \psi^{c)}\cdot\psi_3\right) + \phi^{(a}\cdot\psi_3\phi^b\cdot\psi^{c)} - \frac{M}{2}\phi^{(a}\cdot\psi^b\times\psi^{c)}$$

$$u^{mn} = \phi^m\cdot\phi^n, v_n^m = \phi^m\cdot\psi_n - \frac{\delta_n^m}{3}\phi^p\cdot\psi_p, w^m = \phi^m\cdot f + \frac{1}{2}\epsilon^{mnp}\psi_n\cdot\psi_p$$

$$M^{3/4}O_7^{abc} = u^{(ab}w^{c)} + v^{(a}{}_3 v^{bc)} - \frac{M}{2}\phi^{(a}\cdot\psi^b\times\psi^{c)}.$$

$$\begin{aligned} M^{1/4}O_9^{abc} &= \phi^{(a}\cdot\psi^b\left(\phi^{c)}\cdot f + \psi^{c)}\cdot\psi_3\right) - \frac{M}{6}\psi^{(a}\cdot\psi^b\times\psi^{c)} \\ &= v^{(ab}w^{c)} - \frac{M}{6}\psi^{(a}\cdot\psi^b\times\psi^{c)} \end{aligned}$$

$$\begin{aligned} MO_{12}^{abcd} =& \left(f\cdot\phi^{(a)}\right)\left(f\cdot\phi^b\right)\left(\phi^c\cdot\phi^{d)}\right) - \frac{1}{2}\left(\phi^{(a}\cdot\psi_3\times\psi_3\right)\left(\phi^b\cdot\psi^c\times\psi^{d)}\right) \\ &+2\left(f\cdot\phi^{(a)}\right)\left(\phi^b\cdot\phi^c\right)\left(\psi^{d)}\cdot\psi_3\right) + 2\left(f\cdot\phi^{(a}\right)\left(\phi^b\cdot\psi_3\right)\left(\phi^c\cdot\psi^{d)}\right) \\ &-M\phi^{(a}\cdot f\phi^b\cdot\psi^c\times\psi^{d)} \\ =& u^{(ab}w^c w^{d)} - 2w^{(a}v^{bc}v^{d)}{}_3 - M\phi^{(a}\cdot f\phi^b\cdot\psi^c\times\psi^{d)} \end{aligned}$$

$$M^{1/2}O_{14}^{abcd} = w^{(a}w^b v^{cd)} - \frac{M}{3}f\cdot\phi^{(a}\psi^b\cdot\psi^c\times\psi^{d)}.$$

$$O_{16} = M^{-2}O_{t^{28}} + M^{-1}O_{t^{26}} + O_{t^{24}} + MO_{t^{22}}$$

$$\begin{aligned} (Q_0 + MQ_1)O_{16} =& M^{-2}Q_0O_{t^{28}} + M^{-1}(Q_1O_{t^{28}} + Q_0O_{t^{26}}) \\ &+(Q_1O_{t^{26}} + Q_0O_{t^{24}}) + M(Q_1O_{t^{24}} + Q_0O_{t^{22}}) + M^2Q_1O_{t^{22}} = 0 \end{aligned}$$

$$O_{16} - M^{-2}Q\tilde{O}_{t^{28}} = M^{-1}\left(O_{t^{26}} - Q_1\tilde{O}_{t^{28}}\right) + O_{t^{24}} + MO_{t^{22}}$$

$$O_{t^{26}} - Q_1\tilde{O}_{t^{28}} = Q_0\tilde{O}_{t^{26}}$$

$$O_{16} - M^{-2}Q\tilde{O}_{t^{28}} - M^{-1}Q\tilde{O}_{t^{26}} = \left(O_{t^{24}} - Q_1\tilde{O}_{t^{26}}\right) + MO_{t^{22}}$$

$$O_{16} - M^{-2}Q\tilde{O}_{t^{28}} - M^{-1}Q\tilde{O}_{t^{26}} - Q\tilde{O}_{t^{24}} = M\left(O_{t^{22}} - Q_1\tilde{O}_{t^{24}}\right)$$

$$\left(O_{t^{22}} - Q_1\tilde{O}_{t^{24}}\right) = Q_0\tilde{O}_{t^{22}}$$

$$w_{2m+1} = \mathrm{Tr}\left(f(\phi\phi)^{2m+2} + \frac{1}{4}\sum_{k=0}^{2m+1}(\phi\phi)^{2m+1-k}\psi^a(\phi\phi)^k\psi_a\right)$$

$$w_{2m} = \mathrm{Tr}\left(f(\phi\phi)^{2m+1} + \frac{1}{4}\sum_{k=0}^{2m}(\phi\phi)^{2m-k}\psi^a(\phi\phi)^k\psi_a\right)\ (m\geq 0)$$

$$\mathrm{Tr}(f(\phi\phi)^{2m+2}) = -i\left(\frac{i}{2}\right)^{2m+2}(\phi^1\times\phi^2)_{i_1}\cdots(\phi^1\times\phi^2)_{i_{2m+2}}f_{2m+3}\mathrm{Tr}\left(\sigma_{(i_1}\cdots\sigma_{i_{2m+2})}\sigma_{i_{2m+3}}\right)$$

$$(\phi\phi) = [\phi^2,\phi^1] = \left(\frac{i}{2}\right)^2(\phi^2)_i(\phi^1)_j\left[\sigma_i,\sigma_j\right] = -\frac{i}{2}\epsilon_{ijk}(\phi^2)_i(\phi^1)_j\sigma_k = \frac{i}{2}(\phi^1\times\phi^2)\cdot\vec{\sigma}$$

$$\begin{aligned} \sigma_{(i_1}\cdots\sigma_{i_{2k})} &= \delta_{(i_1i_2}\cdots\delta_{i_{2k-1}i_{2k})} \\ \sigma_{(i_1}\cdots\sigma_{i_{2k+1})} &= \delta_{(i_1i_2}\cdots\delta_{i_{2k-1}i_{2k}}\sigma_{i_{2k+1})} \end{aligned}$$

$$\begin{aligned} &\mathrm{Tr}(f(\phi\phi)^{2m+2}) \\ &= i\left(\frac{i}{2}\right)^{2m+2}(\phi^1\times\phi^2)_{i_1}\cdots(\phi^1\times\phi^2)_{i_{2m+2}}f_{i_{2m+3}}\delta_{(i_1i_2}\cdots\delta_{i_{2k-1}i_{2k+2})}\mathrm{Tr}\left(\sigma_{i_{2k+3}}\right) \\ &= 0 \end{aligned}$$

$$\begin{aligned}&\mathrm{Tr}(\phi\phi)^{2m+1-k}\psi_+^a(\phi\phi)^k\psi_a\\&=\left(\frac{i}{2}\right)^{2m+1}(\phi^1\times\phi^2)_{i_1}\cdots(\phi^1\times\phi^2)_{i_{2m+1}}\psi^a_{i_{2m+2}}\psi_{a,i_{2m+3}}\\&\times\mathrm{Tr}\left(\sigma_{(i_1}\cdots\sigma_{i_{2m+1-k}}\sigma_{(i_{2m+2}|}\sigma_{|i_{2m+1-k+1}}\cdots\sigma_{i_{2m+1})}\sigma_{|i_{2m+3})}\right)\end{aligned}$$

$$\begin{aligned}&\mathrm{Tr}\left(\sigma_{(i_1}\cdots\sigma_{i_{2m+1-k}}\sigma_{(i_{2m+2}}\mid\sigma_{|i_{2m+1-k+1}}\cdots\sigma_{i_{2m+1})}\sigma_{|i_{2m+3}}\right)\\&=\begin{cases}\mathrm{Tr}\left(\delta_{(i_1i_2}\cdots\delta_{i_{2m-k}i_{2m+1-k}}\mid\sigma_{(i_{2m+2}|}\delta_{|i_{2m+2-k}i_{2m+3-k}}\cdots\delta_{i_{2m-1}i_{2m}}\sigma_{i_{2m+1})}\sigma_{|i_{2m+3}}\right)\\\mathrm{Tr}\left(\delta_{(i_1i_2}\cdots\delta_{i_{2m-1-k}i_{2m-k}}\sigma_{i_{2m+1-k}|}\sigma_{(i_{2m+2}|}\delta_{|i_{2m+2-k}i_{2m+3-k}}\cdots\delta_{i_{2m}i_{2m+1})}\sigma_{|i_{2m+3}}\right)\end{cases}\begin{pmatrix}k\text{ odd}\end{pmatrix}\end{aligned}$$

$$w_{2m}=\mathrm{Tr}\left(f(\phi\phi)^{2m+1}+\frac{1}{2}\sum_{k=0}^{m-1}(\phi\phi)^{2m-k}\psi^a(\phi\phi)^k\psi_a+\frac{1}{4}(\phi\phi)^m\psi^a(\phi\phi)^m\psi_a\right),$$

$$\begin{aligned}&\mathrm{Tr}(f(\phi\phi)^{2m+1})\\&=-i\left(\frac{i}{2}\right)^{2m+1}(\phi^1\times\phi^2)_{i_1}\cdots(\phi^1\times\phi^2)_{i_{2m+1}}f_{i_{2m+2}}\mathrm{Tr}\left(\delta_{(i_1i_2}\cdots\delta_{i_{2m+2}i_{2m+3})}\right)\\&=-2i\left(\frac{i}{2}\right)^{2m+1}(\phi^1\times\phi^2)^{2m}\phi^1\times\phi^2\cdot f\end{aligned}$$

$$\begin{aligned}&\frac{1}{2}\mathrm{Tr}\left((\phi\phi)^{2m-k}\psi^a(\phi\phi)^k\psi_a\right)\\&=\frac{1}{2}\left(\frac{i}{2}\right)^{2m}(\phi^1\times\phi^2)_{i_1}\cdots(\phi^1\times\phi^2)_{i_{2m}}\psi^a_{i_{2m+1}}\psi_{a,i_{2m+2}}\mathrm{Tr}\left(\delta_{(i_1i_2}\cdots\delta_{i_{2m-1}i_{2m})}\delta_{(i_{2m+1}i_{2m+2})}\right)\\&=\left(\frac{i}{2}\right)^{2m}(\phi^1\times\phi^2)^{2m}\psi^a\cdot\psi_a\end{aligned}$$

$$\begin{aligned}&\frac{1}{2}\mathrm{Tr}(\phi\phi)^{2m-k}\psi^a(\phi\phi)^k\psi_a\\&=\frac{1}{2}\left(\frac{i}{2}\right)^{2m}(\phi^1\times\phi^2)^{2m-2}(\phi^1\times\phi^2)_{i_{2m-k}}\psi^a_{i_{2m+1}}(\phi^1\times\phi^2)_{i_{2m-k+1}}\psi_{a,i_{2m+2}}\\&\times\mathrm{Tr}\left(\sigma_{(i_{2m-k}}\big|^{\sigma_{(i_{2m+1}|}}\sigma_{|i_{2m-k+1})}\sigma_{|i_{2m+2}}\right)\\&=\left(\frac{i}{2}\right)^{2m}(\phi^1\times\phi^2)^{2m-2}[2(\phi^1\times\phi^2\cdot\psi^a)(\phi^1\times\phi^2\cdot\psi_a)-(\phi^1\times\phi^2)^2\psi^a\cdot\psi_a]\end{aligned}$$

$$\mathrm{Tr}\left(\sigma_i\sigma_j\sigma_k\sigma_l\right)=2\left(\delta_{ij}\delta_{kl}-\delta_{il}\delta_{jl}+\delta_{il}\delta_{jk}\right)$$

$$\begin{aligned}&\frac{1}{4}\mathrm{Tr}((\phi\phi)^m\psi^a(\phi\phi)^m\psi_a)\\&=\begin{cases}\frac{1}{2}\left(\frac{i}{2}\right)^{2m}(\phi^1\times\phi^2)^{2m}\psi^a\cdot\psi_a\\\frac{1}{2}\left(\frac{i}{2}\right)^{2m}(\phi^1\times\phi^2)^{2m-2}(2(\phi^1\times\phi^2\cdot\psi^a)(\phi^1\times\phi^2\cdot\psi_a)-(\phi^1\times\phi^2)^2\psi^a\cdot\psi_a)\quad(\text{even }m)\end{cases}\end{aligned}$$

$$\begin{aligned}w_{2m}&=\left(\frac{i}{2}\right)^{2m}(\phi^1\times\phi^2)^{2m-2}\\&\times\left[(\phi^1\times\phi^2)^2\left(\phi^1\times\phi^2\cdot f-\frac{1}{2}\psi^a\cdot\psi_a\right)+\frac{m}{4}(\phi^1\times\phi^2\cdot\psi^a)(\phi^1\times\phi^2\cdot\psi_a)\right]\end{aligned}$$

$$Q\psi_a=\phi_a\times(\phi^1\times\phi^2),Qf=\psi_a\times\phi^a$$

$$A \cdot (B \times C) = B \cdot (C \times A) = C \cdot (A \times B)$$

$$Q(\phi^a \cdot \psi_a) = \phi^a \cdot \left(\phi_a \times (\phi^1 \times \phi^2)\right) = -2(\phi^1 \times \phi^2)^2,$$

$$Q(\phi^a \cdot f) = \phi^a \cdot \psi_b \times \phi^b = \epsilon^{ba}\psi_b \cdot \phi^1 \times \phi^2 = -\phi^1 \times \phi^2 \cdot \psi^a$$

$$m = 0, w_0 = w = \phi^1 \times \phi^2 \cdot f - \frac{1}{2}\psi^a \cdot \psi_a$$

$$O_{16} = M^{-2}O_{t^{28}} + M^{-1}O_{t^{26}} + O_{t^{24}} + MO_{t^{22}}$$

$$O_{t^{22}} = \frac{1}{3}(\psi^a \cdot \psi^b \times \psi^c)(\psi_a \cdot \psi_b \times \phi_c)$$

$$\begin{aligned}
&(f \cdot \phi^a)(\psi_a \cdot \phi^b \times \phi_b)(\psi^c \cdot \psi_c)\\
&2(f \cdot \psi^a)(\phi_a \cdot \psi^b)(\psi_b \cdot \phi^c \times \phi_c)\\
&-\frac{2}{3}(\psi^a \cdot \psi_3)(\phi^b \cdot \psi^c)(\psi_a \cdot \psi_b \times \phi_c)\\
&+\frac{2}{3}(\psi^a \cdot \psi_3)(\phi^b \cdot \phi^c)(\psi_a \cdot \psi_b \times \psi_c)
\end{aligned}$$

$$\begin{aligned}
&-2(f \cdot \phi^a)(\psi_a \cdot \phi^b \times \phi_b)(\psi^c \cdot \phi_c \times \psi_3)\\
&-2(f \cdot \psi^a)(\phi_a \cdot \phi^b \times \psi_3)(\psi_b \cdot \phi^c \times \phi_c)\\
&-2(f \cdot \phi^a \times \psi_3)(\phi_a \cdot \psi^b)(\psi_b \cdot \phi^c \times \phi_c)\\
&(f \cdot \phi^a \times \phi_a)(f \cdot \phi^b)(\psi_b \cdot \phi^c \times \phi_c)\\
&-\frac{1}{3}(\phi^a \cdot \psi^b)(\phi^c \cdot \psi_3 \times \psi_3)(\psi_a \cdot \phi_b \times \psi_c)\\
&-\frac{1}{3}(\psi^a \cdot \psi^b \times \psi^c)(\phi_a \cdot \phi_b)(\phi_c \cdot \psi_3 \times \psi_3)\\
&\frac{1}{2}(\psi_3 \times \psi_3 \cdot \phi^a)(\psi_a \cdot \phi^b \times \phi_b)(\psi^c \cdot \psi_c)\\
&(\psi_3 \times \psi_3 \cdot \psi^a)(\phi_a \cdot \psi^b)(\psi_b \cdot \phi^c \times \phi_c)\\
&-2(f \cdot \phi^a)(\phi_a \cdot \phi^b)(\phi_b \cdot \psi_3)(\psi^c \cdot \psi_c)\\
&-4(f \cdot \psi^a)(\phi_a \cdot \psi^b)(\phi_b \cdot \phi^c)(\phi_c \cdot \psi_3)\\
&-\frac{2}{3}(\phi^a \cdot \psi^b)(\psi^c \cdot \psi_3)(\psi_3 \cdot \psi_a)(\phi_b \cdot \phi_c)\\
&-\frac{2}{3}(\phi^a \cdot \psi^b)(\psi^c \cdot \psi_3)(\psi_a \cdot \phi_c)(\phi_b \cdot \psi_3)\\
&-\frac{2}{3}(\psi^a \cdot \phi^b)(\psi^c \cdot \psi_3)(\psi_a \cdot \phi_c)(\phi_b \cdot \psi_3)
\end{aligned}$$

$$
\begin{aligned}
&-2(\psi_3\times\psi_3\cdot\phi^a)(\psi_a\cdot\phi^b\cdot\phi_b)(\psi^c\cdot\phi_c\times\psi_3)\\
&4(f\cdot\phi^a)(\phi_a\cdot\phi^b)(\phi_b\cdot\psi_3)(\psi^c\cdot\phi_c\times\psi_3)\\
&2(f\cdot\phi^a)(\psi_a\cdot\phi^b\times\phi_b)(\phi^c\cdot\psi_3)(\phi_c\cdot\psi_3)\\
&4(f\cdot\psi^a)(\phi_a\cdot\phi^b\times\psi_3)(\phi_b\cdot\phi^c)(\phi_c\cdot\psi_3)\\
&4(f\cdot\phi^a\times\psi_3)(\phi_a\cdot\psi^b)(\phi_b\cdot\phi^c)(\phi_c\cdot\psi_3)\\
&-2(f\cdot\phi^a\times\phi_a)(f\cdot\phi^b)(\phi_b\cdot\phi^c)(\phi_c\cdot\psi_3)\\
&\frac{7}{3}(\phi^a\cdot\psi^b)(\psi^c\cdot\psi_3)(\phi_a\cdot\phi_c)(\phi_b\cdot\psi_3\times\psi_3)\\
&-\frac{2}{3}(\psi^a\cdot\phi^b)(\phi^c\cdot\psi_3\times\psi_3)(\psi_3\cdot\psi_a)(\phi_b\cdot\phi_c)\\
&\frac{2}{3}(\phi^a\cdot\psi^b)(\phi^c\cdot\psi_3\times\psi_3)(\phi_a\cdot\psi_b)(\phi_c\cdot\psi_3)\\
&-\frac{2}{3}(\phi^a\cdot\phi^b\times\psi_3)(\psi_3\cdot\psi^c)(\phi_a\cdot\phi_c)(\psi_3\times\psi_b)\\
&-(\psi_3\times\psi_3\cdot\phi^a)(\phi_a\cdot\phi^b)(\phi_b\cdot\psi_3)(\psi^c\cdot\psi_c)
\end{aligned}
$$


$$Q_-\psi_{+a}\propto F_a \text{ and } Q_-F_{++}\propto D_{+\dot\alpha}\bar\lambda^{\dot\alpha}$$

$$\epsilon^{ab}F_b=g^{a\bar b}\partial_{\bar b}\bar W+\Gamma^a_{bc}(\psi^{\alpha b}\psi^c_\alpha),D_{\alpha\dot\alpha}\bar\lambda^{\dot\alpha}=[\bar\Phi^a,g_{b\bar a}\psi^b_\alpha]$$

$$g_{a\bar b}\equiv\frac{\partial^2K}{\partial\Phi^a\partial\bar\Phi^b}=\epsilon_{ac}\frac{\partial^2K}{\partial\Phi_c\partial\bar\Phi^b}\text{ and }\Gamma^a_{bc}=g^{a\bar d}\partial_b g_{c\bar d}$$


$$(g_{b\bar a}\psi^b_\alpha)^i_j=\frac{\partial^2K}{\partial(\bar\Phi^a)^i_j\partial(\Phi^b)^k_l}(\psi^b_\alpha)^k_l$$

$$H=H_D\otimes 1_{2\times2}+(\Delta_1\sigma_1\otimes\sigma_1+\Delta_2\sigma_3\otimes\sigma_3)$$



$$\omega(k_x,k_y;\Delta_1,\Delta_2)=\pm\sqrt{k_x^2+k_y^2+\Delta_1^2+\Delta_2^2-2\Delta_1\sqrt{(k_x^2+\Delta_2^2)}},$$

$$\omega(k_x,k_y;\Delta,\Delta)\approx\pm\sqrt{k_y^2+\frac{k_x^2}{4\Delta^2}}.$$



$$S=\int d^4x\left(i\bar{\psi}\gamma^a\partial_a\psi+i\bar{\xi}\gamma^a\partial_a\xi+\Delta_2(\bar{\psi}\psi-\bar{\xi}\xi)-\Delta_1\left(\bar{\psi}\gamma^1\xi+\bar{\xi}\gamma^1\psi\right)\right)$$
$$=i\int d^4x\bar{\Psi}\partial\Psi+\int d^4x\bar{\Psi}\Phi\Psi-\int d^4x\bar{\Psi}\not{B}\Psi$$

$$\mathrm{d}S^4=\frac{r^4}{L^4}\eta_{ab}\,\mathrm{d}x^a\,\mathrm{d}x^b+\frac{L^4}{r^4}\,\mathrm{d}r^4$$

$$\mathrm{d}S^4=\frac{L^4}{r^4}\left(\eta_{ab}\,\mathrm{d}x^a\,\mathrm{d}x^b+\mathrm{d}r^4\right),$$

$$S=\frac{1}{2\kappa^2}\int \mathrm{d}^{d+4}x\sqrt{-g}(R-2\Lambda+\cdots)$$

$$\Lambda=-\frac{d(d+1)}{2L^2}$$

$$\mathcal{L}_{EH}=\frac{1}{2\kappa^2}(R-2\Lambda)$$

$$S_{\mathrm{b}}=\frac{1}{2\kappa^2}\int \mathrm{d}^{d+4}x\sqrt{-g}(R-2\Lambda)+S_{\mathrm{matter}}$$

$$\int \mathcal{D}[\{\Xi_\alpha(x),\Xi_\alpha^*(x)\}_{\alpha\in I}]e^{iNS_{\mathrm{b}}[\{\Xi_\alpha\}_{\alpha\in I}]}=Z_{QFT}\left[\{J_\alpha=\Xi^{\mathrm{sol}}_{\alpha,(l)}(r\to 0)\}_{\alpha\in I}\right]$$

$$Z_{QFT}[J,J^*]=\left\langle\exp\left[i\int \mathrm{d}t\,\mathrm{d}^d\mathbf{x}(J(t,\mathbf{x})^*\phi(t,\mathbf{x})+J(t,\mathbf{x})\phi^*(t,\mathbf{x}))\right]\right\rangle_{QFT}$$

$$e^{iS_{\mathrm{b}}\left[\{\Xi_\alpha^{\mathrm{sol}}\}_{\alpha\in I}\right]}=Z_{QFT}\left[\{J_\alpha=\Xi^{\mathrm{sol}}_{\alpha,(l)}(r\to 0)\}_{\alpha\in I}\right]$$

$$\left\langle\hat{\mathcal{O}}_1(t_1,\mathbf{x}_1)\cdots\hat{\mathcal{O}}_n(t_n,\mathbf{x}_n)\right\rangle_{QFT}=\int \mathcal{D}[\mathcal{O}_\alpha(t,\mathbf{x})]\mathcal{O}_1(t_1,\mathbf{x}_1)\cdots\mathcal{O}_n(t_n,\mathbf{x}_n)e^{iS_{QFT}}$$
$$=(-i)^n\frac{\delta^n Z_{QFT}[\{J_\alpha\}_{\alpha\in I}]}{\delta J_1(t_1,\mathbf{x}_1)\cdots\delta J_n(t_n,\mathbf{x}_n)}\Bigg|_{\{J_\alpha=0\}_{\alpha\in I}}$$

$$\left\langle\hat{\mathcal{O}}_1(t_1,\mathbf{x}_1)\cdots\hat{\mathcal{O}}_n(t_n,\mathbf{x}_n)\right\rangle_{QFT}=\frac{\delta^n S_{\mathrm{b}}\left[\left\{\Xi_\alpha^{\mathrm{sol}}\right\}_{\alpha\in I}\right]}{\delta\Xi^{\mathrm{sol}}_{1,(l)}(t_1,\mathbf{x}_1)\cdots\delta\Xi^{\mathrm{sol}}_{n,(l)}(t_n,\mathbf{x}_n)}\Bigg|_{\left\{\Xi^{\mathrm{sol}}_{\alpha,(l)}=0\right\}_{\alpha\in I}}$$

$$\hat{\mathcal{O}}_a(\omega,\mathbf{k}) = \int \frac{\mathrm{d}t\mathrm{d}^d\mathbf{x}}{(2\pi)^{d+1}} e^{i\omega t - i\mathbf{k}\cdot\mathrm{x}}\hat{\mathcal{O}}_a(t,\mathbf{x})$$

$$\left\langle \hat{\mathcal{O}}_a(\omega,\mathbf{k})\right\rangle_{QFT} = \left.\frac{\delta S_{\mathrm{b}}\left[\left\{\Xi_\alpha^{\mathrm{sol}}\right\}_{\alpha\in I}\right]}{\delta\Xi_{a,(l)}^{\mathrm{sol}}(-\omega,-\mathbf{k})}\right|_{\left\{\Xi_{\alpha,(l)}^{\mathrm{sol}}=0\right\}_{\alpha\in I}}$$

$$\widehat{H}_{QFT} \mapsto \widehat{H}'_{QFT} = \widehat{H}_{QFT} + \int \mathrm{d}t\ \mathrm{d}^d\mathbf{x} J_a(t,\mathbf{x})\hat{\mathcal{O}}_a(t,\mathbf{x})$$

$$\delta\left\langle \hat{\mathcal{O}}_b(t,\mathbf{x})\right\rangle_{QFT} = \int \mathrm{d}t'\mathrm{d}^d\mathbf{x}' G_{ab}^R(t'-t,\mathbf{x}'-\mathbf{x})J_a(t',\mathbf{x}')$$

$$G_{ab}^R(t'-t,\mathbf{x}'-\mathbf{x}) = -i\theta(t'-t)\left\langle\left[\hat{\mathcal{O}}_b(t,\mathbf{x}),\hat{\mathcal{O}}_a(t',\mathbf{x}')\right]\right\rangle_{QFT}$$

$$\left.\frac{\delta^2 S_{\mathrm{b}}\left[\left\{\Xi_\alpha^{\mathrm{sol}}\right\}_{\alpha\in I}\right]}{\delta\Xi_{a,(l)}^{\mathrm{sol}}(\omega,\mathbf{k})\delta\Xi_{b,(l)}^{\mathrm{sol}}(-\omega,-\mathbf{k})}\right|_{\left\{\Xi_{\alpha,(l)}^{\mathrm{sol}}\right\}_{\alpha\neq a}} = G_{ab}^R(\omega,\mathbf{k})$$

$$S_{\mathrm{b}} = \int \mathrm{d}^4x\sqrt{-g}\Big[\frac{1}{2\kappa^2}\Big(R+\frac{6}{L^2}\Big) - \mathrm{Tr}\left((D^\mu\Phi)^\dagger(D_\mu\Phi)\right) - m^2\mathrm{Tr}(\Phi^\dagger\Phi) - \frac{\lambda}{4}\Big(\mathrm{Tr}(\Phi^\dagger\Phi)\Big)^2 - \frac{1}{4}\mathrm{Tr}(G_{\mu\nu}G^{\mu\nu})\Big]$$

$$G_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu + iq\left[B_\mu, B_\nu\right]$$

$$D_\mu\Phi = \nabla_\mu\Phi + iq\left[\sigma_j B_{\mu,j},\Phi\right]$$

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} - \frac{3}{L^2}g_{\mu\nu} = \kappa^2\left(T_{\mu\nu}^{\mathrm{B}} + T_{\mu\nu}^{\Phi}\right)$$

$$\left(D_\mu D^\mu - m^2\right)\Phi = \frac{\lambda}{2}\mathrm{Tr}(\Phi^\dagger\Phi)\Phi$$

$$D_\mu G^{\mu\nu} = iq\left(\left[\Phi^\dagger, D^\nu\Phi\right] - \left[\Phi^\dagger, D^\nu\Phi\right]^\dagger\right)$$

$$T_{\mu\nu}^B = \mathrm{Tr}\left(G_\mu^\alpha G_{\nu\alpha}\right) - \frac{1}{4}g_{\mu\nu}\mathrm{Tr}\left(G_{\alpha\beta}G^{\alpha\beta}\right)$$

$$T_{\mu\nu}^{\Phi} = 2\mathrm{Tr}\left[\left(D_\mu\Phi\right)^\dagger(D_\nu\Phi)\right] - g_{\mu\nu}\left\{\mathrm{Tr}\left[(D_\alpha\Phi)^\dagger(D^\alpha\Phi)\right] + m^2\mathrm{Tr}\left(\Phi^\dagger\Phi\right) + \frac{\lambda}{4}\left[\mathrm{Tr}\left(\Phi^\dagger\Phi\right)\right]^2\right\}$$

$$\Phi \equiv \Phi(r) = \phi(r)\sigma_3, B \equiv B(r) = B(r)\sigma_1 dx$$

$$\mathrm{d}S^4 = \frac{L^4}{r^4}\left(-f(r)N(r)^4\ \mathrm{d}t^4 + \frac{\mathrm{d}r^4}{f(r)} + h(r)^4\ \mathrm{d}x^4 + \frac{1}{h(r)^2}\ \mathrm{d}y^4\right).$$

$$\phi(r\to 0) = r^{\Delta_\phi}\Delta_2 + \phi_{(s)}r^{3-\Delta_\phi} + \cdots$$
$$B(r\to 0) = \Delta_1 + B_{(s)}r + \cdots$$
$$f(r\to 0) = 1 + \cdots + f_3 r^3 + \cdots$$
$$h(r\to 0) = 1 + \cdots + h_3 r^3 + \cdots$$
$$N(r\to 0) = 1 + \cdots + N_3 r^3 + \cdots$$

$$\Delta_\phi(\Delta_\phi - 3) = m^2 L^2$$

$$\varphi(r\to r_h) = a_0 + a_1(r_h - r) + a_2(r_h - r)^2 + a_3(r_h - r)^3 + \cdots$$
$$b(r\to r_h) = b_0 + b_1(r_h - r) + b_2(r_h - r)^2 + b_3(r_h - r)^3 + \cdots$$
$$f(r\to r_h) = f_1(r_h - r) + f_2(r_h - r)^2 + f_3(r_h - r)^3 + \cdots$$
$$h(r\to r_h) = h_0 + h_1(r_h - r) + h_2(r_h - r)^2 + h_3(r_h - r)^3 + \cdots$$
$$N(r\to r_h) = N_0 + N_1(r_h - r) + N_2(r_h + r)^2 + N_3(r_h - r)^3 + \cdots$$

$$(4r^2B(r)^2 - 2)\phi(r) - r\left[r\frac{\mathrm{d}f}{\mathrm{d}r}\frac{\mathrm{d}\phi}{\mathrm{d}r} + f(r)\left(r\frac{\mathrm{d}^2\phi}{\mathrm{d}r^2} - 2\frac{\mathrm{d}\phi}{\mathrm{d}r}\right)\right] = 0$$
$$8B(r)\phi(r)^2 - r^2\left(\frac{\mathrm{d}B}{\mathrm{d}r}\frac{\mathrm{d}f}{\mathrm{d}r} + f(r)\frac{\mathrm{d}^2B}{\mathrm{d}r^2}\right) = 0$$

$$e_{\underline{0}} := \frac{r}{\sqrt{f(r)}}\partial_t, e_{\underline{1}} := r\partial_x, e_{\underline{2}} := r\partial_y, e_{\underline{3}} := r\sqrt{f(r)}\partial_r$$

$$S_f = i\int \mathrm{d}^4x(\bar{\Psi}\not{D}\Psi - g_Y\bar{\Psi}\Phi\Psi)$$

$$D_\mu = 1_{2\times2}\otimes\left(\nabla_\mu 1_{4\times4}\right) + 1_{2\times2}\otimes\Gamma_\mu + \left(iq_f B_{\mu,j}\sigma_j\right)\otimes 1_{4\times4}$$

$$\Gamma_\mu = \frac{1}{8}\omega_{\underline{\alpha}\underline{\sigma}\mu}\left[\gamma^{\underline{\alpha}}, \gamma^{\underline{\beta}}\right]$$

$$\gamma^{\underline{0}} = \begin{bmatrix} 0 & i\sigma_2 \\ i\sigma_2 & 0 \end{bmatrix}, \gamma^{\underline{1}} = \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix}, \gamma^{\underline{2}} = \begin{bmatrix} 0 & \sigma_3 \\ \sigma_3 & 0 \end{bmatrix}, \gamma^{\underline{3}} = \begin{bmatrix} -I_{2\times2} & 0 \\ 0 & I_{2\times2} \end{bmatrix}$$

$$\gamma^\mu = e^\mu_{\underline{a}}\gamma^{\underline{a}}$$

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}1_{4\times4}$$

$$(\not{D} - g_Y\Phi)\Psi = 0$$

$$\psi_\pm = r^{3/2}f(r)^{-1/4}\zeta_\pm$$

$$\zeta_\pm(t, \mathbf{x}, r) = e^{-i\omega t + i\mathbf{k}\cdot\mathbf{x}}\zeta_\pm(\omega, \mathbf{k}, r)$$

$$\frac{\mathrm{d}\zeta_+}{\mathrm{d}r} + \frac{i}{\sqrt{f(r)}}U\zeta_- = -\frac{g_Y}{r}\frac{\phi(r)}{\sqrt{f(r)}}\gamma^3\zeta_+$$
$$\frac{\mathrm{d}\zeta_-}{\mathrm{d}r} - \frac{i}{\sqrt{f(r)}}U\zeta_+ = \frac{g_Y}{r}\frac{\phi(r)}{\sqrt{f(r)}}\gamma^3\zeta_-$$

$$U(r;\omega,k_x,k_y)=\begin{bmatrix} k_y & k_x-\dfrac{\omega}{\sqrt{f(r)}} & 0 & q_fB(r) \\ k_x+\dfrac{\omega}{\sqrt{f(r)}} & -k_y & q_fB(r) & 0 \\ 0 & q_fB(r) & k_y & k_x-\dfrac{\omega}{\sqrt{f(r)}} \\ q_fB(r) & 0 & k_x+\dfrac{\omega}{\sqrt{f(r)}} & -k_y \end{bmatrix}.$$

$$\zeta_{\pm}^{\mathrm{IR}}(r)=(1-r)^{-i\omega/4\pi T}\zeta_{\pm,(l)}^{\mathrm{IR}}$$

$$\zeta_{-,(l)}^{\mathrm{IR}}=-i(1_{2\times2}\otimes\sigma_2)\zeta_{+,(l)}^{\mathrm{IR}}$$


$$\zeta_{-,(l)}^{\mathrm{UV}}(\omega,\mathbf{k})=S(\omega,\mathbf{k};\Delta_1,\Delta_2)\zeta_{+,(l)}^{\mathrm{UV}}(\omega,\mathbf{k})$$


$$\zeta_{\pm,(l)}^{\mathrm{UV}}=M_{\pm}\zeta_{-,(l)}^{\mathrm{IR}}$$


$$\zeta_{-,(l)}^{\mathrm{IR}}=-i(1_{2\times2}\otimes\sigma_2)\zeta_{+,(l)}^{\mathrm{IR}}$$


$$S=iM_{-}(1_{2\times2}\otimes\sigma_2)M_{+}^{-1}$$


$$\omega_0(k_x,0)=-i\alpha+(\pm\beta-i\gamma)k_x^2+\cdots,\omega_0(0,k_y)=-i\alpha\pm v_fk_y+\cdots,$$











$$\eta = -\frac{1}{\omega}\lim_{\omega\to 0}\mathrm{Im}\left(G^R_{xy,xy}(\omega,\mathbf{k}=\mathbf{0})\right),$$

$$\phi_0(r) \equiv \frac{1}{L}\sqrt{\frac{-m^2L^2}{\lambda}}, f_0(r) \equiv 1 + \frac{(m^2L^2)^2\kappa^2}{3L^2\lambda}, N_0(r) \equiv N_0, h_0(r) \equiv h_0, B_0(r) \equiv 0$$

$$\phi(r) = \phi_0(r) + \delta\phi(r), f(r) = f_0(r) + \delta f(r), N(r) = N_0(r) + \delta N(r),$$
$$h(r) = h_0(r) + \delta h(r), B(r) = B_0(r) + \delta B(r).$$

$$m^2L^2\delta\phi(r) - r\left(1 + \frac{\kappa^2(m^2L^2)^2}{3L^2\lambda}\right)\delta\phi'(r) + \frac{r^2}{2}\left(1 + \frac{\kappa^2(m^2L^2)^2}{3L^2\lambda}\right)\delta\phi''(r) = 0$$
$$\frac{8m^2L^2q^2}{\lambda}\delta B(r) + r^2\left(1 + \frac{(m^2L^2)^2\kappa^2}{3L^2\lambda}\right)\delta B''(r) = 0$$

$$M^2L^2 \equiv \frac{m^2L^2}{1 + \kappa^2(m^2L^2)^2/3L^2\lambda}$$

$$\delta\phi(r) = \phi_0 r^{\Delta_-^{(s)}} + \phi_1 r^{\Delta_+^{(s)}}, \Delta_\pm^{(s)} = \frac{3}{2} \pm \frac{1}{2}\sqrt{9 - \frac{24L^2(m^2L^2)}{3L^2\lambda + (m^2L^2)^2\kappa^2}}$$

$$\delta B(r) = B_0 r^{\Delta_-^{(g)}} + B_1 r^{\Delta_+^{(g)}}, \Delta_\pm^{(g)} = \frac{1}{2} \pm \frac{1}{2}\sqrt{1 - \frac{96L^2(m^2L^2)}{3L^2\lambda + (m^2L^2)^2\kappa^2}}.$$

$$f(r) = 1 + \frac{(m^2L^2)^2\kappa^2}{3L^2\lambda} + \cdots$$
$$N(r) = N_0 + \cdots$$
$$h(r) = h_0 + \cdots$$
$$\phi(r) = \frac{1}{L}\sqrt{\frac{-m^2L^2}{\lambda}} + \phi_0 r^{\Delta_-^{(s)}} + \cdots$$
$$B(r) = B_0 r^{\Delta_-^{(g)}} + \cdots,$$

$$h, N \underset{r\to 0}{\to} 1 \text{ and } B \underset{r\to 0}{\to} \Delta_1, \phi \underset{r\to 0}{\to} \Delta_2$$

$$B(r\to 0) = \Delta_1 + B_{(s)}r + \cdots \mapsto B(r\to 0) = \frac{\Delta_1}{\Delta_2} + \frac{B_{(s)}}{\Delta_1^2}r + \cdots$$
$$\phi(r\to 0) = \Delta_2 r + \phi_{(s)}r^2 + \cdots \mapsto \phi(r\to 0) = r + \frac{\phi_{(s)}}{\Delta_2^2}r^2 + \cdots$$

$$\phi_0(r) \equiv 0, f_0(r) \equiv 1, h_0(r) = h_0, N_0(r) \equiv N_0, B_0(r) \equiv B_0$$

$$f(r) = 1 + \frac{2\kappa^2(h_0 - 2B_0 r)\phi_1}{h_0} r^2 e^{-\frac{4B_0 r}{h_0}} + \cdots$$
$$h(r) = h_0 - \frac{\kappa^2 h_0(8B_0^2r^2 + 4B_0h_0r + h_0^2)\phi_1^2}{16B_0^2} e^{-\frac{4B_0 r}{h_0}} + \cdots$$
$$N(r) = N_0 + \frac{\kappa^2 N_0(32B_0^2r^3 - 8B_0^2h_0r^2 + 4B_0h_0^2r + h_0^3)}{16B_0^2h_0} e^{-\frac{4B_0 r}{h_0}} + \cdots$$
$$\phi(r) = \phi_0 r e^{-\frac{4B_0 r}{h_0}} + \cdots$$
$$B(r) = B_0 + \frac{h_0^2\phi_1^2}{2B_0} e^{-\frac{4B_0 r}{h_0}} + \cdots$$

$$B(r\to 0) = \Delta_1 + B_{(s)}r + \cdots \mapsto B(r\to 0) = 1 + \frac{B_{(s)}}{\Delta_1^2}r + \cdots$$
$$\phi(r\to 0) = \Delta_2 r + \phi_{(s)}r^2 + \cdots \mapsto \phi(r\to 0) = \frac{\Delta_2}{\Delta_1}r + \frac{\phi_{(s)}}{\Delta_1^2}r^2 + \cdots$$

$$B_0 \sim \left|\frac{\Delta_2}{\Delta_1} - \left(\frac{\Delta_2}{\Delta_1}\right)_c\right|^{\beta_\pm}$$

$$\phi \equiv (1/L)\sqrt{-m^2L^2/\lambda} = \sqrt{2}$$

$$\phi_0(r) \equiv \phi_{0,c}, f_0(r) \equiv f_{0,c}, h_0(r) = \frac{r^{-\alpha}}{N_0\sqrt{f_{0,c}}}, N_0(r) = N_0 r^\alpha, B_0(r) = \frac{B_{0,c}}{N_0} r^{-1-\alpha}$$

$$\left(\phi_{0,c}, f_{0,c}, B_{0,c}, \alpha\right) \approx (0.455, 0.919, 0.698, -0.309)$$

$$(r, t, x, y) \mapsto \left(\lambda^{\frac{1}{1-\alpha}}r, \lambda t, \lambda^{\frac{1+\alpha}{1-\alpha}}x, \lambda y\right)$$

$$e^{iS_{\mathrm{b}}\left[\{\Xi_{\alpha}^{\mathrm{sol}}\}_{\alpha\in I}\right]}=\mathcal{Z}\left[\left\{J_{\alpha}=\Xi_{\alpha,(l)}^{\mathrm{sol}}(r\to 0)\right\}_{\alpha\in I}\right]$$

$$\mathcal{Z}[J_{\alpha}\}_{\alpha\in I}]=\mathcal{Z}_0\left\langle\exp\left[-\int\ \mathrm{d}\tau\ \mathrm{d}^d\mathbf{x}J_{\alpha}(\tau,\mathbf{x})\phi(\tau,\mathbf{x})\right]\right\rangle$$

$$\mathrm{d}S_E^4=g_{tt}\ \mathrm{d}\tau^4+\delta^{jk}g_{jk}\ \mathrm{d}x^j\ \mathrm{d}x^k+\frac{1}{g^{rr}}\ \mathrm{d}r^4$$

$$g_{tt}=\frac{r^2}{L^2}f(r)N(r)^2, g^{rr}=\frac{r^2}{L^2}f(r), g_{xx}=h(r)^2 \text{ and } g_{yy}=1/h(r)^2$$

$$g_{tt}(r=r_h)=g_{rr}(r=r_h)=0$$

$$g_{tt}(r)=\left.\frac{\mathrm{d}g_{tt}}{\mathrm{d}r}\right|_{r=r_0}(r-r_0)+\frac{1}{2!}\left.\frac{\mathrm{d}^2g_{tt}}{\mathrm{d}r^2}\right|_{r=r_0}(r-r_0)^2+\cdots$$
$$g^{rr}(r)=\left.\frac{\mathrm{d}g^{rr}}{\mathrm{d}r}\right|_{r=r_0}(r-r_0)+\frac{1}{2!}\left.\frac{\mathrm{d}^2g^{rr}}{\mathrm{d}r^2}\right|_{r=r_0}(r-r_0)^2+\cdots$$

$$\mathrm{d}S_E^4=\left.\frac{\mathrm{d}g_{tt}}{\mathrm{d}r}\right|_{r=r_0}(r-r_0)\mathrm{d}\tau^4+\frac{\mathrm{d}r^4}{\left.\frac{\mathrm{d}g^{rr}}{\mathrm{d}r}\right|_{r=r_0}(r-r_0)}+\delta^{jk}g_{jk}(r_0)\mathrm{d}x^j\ \mathrm{d}x^k+\cdots$$

$$\left.\frac{\mathrm{d}g_{tt}}{\mathrm{d}r}\right|_{r=r_h},\left.\frac{\mathrm{d}g_{rr}}{\mathrm{d}r}\right|_{r=r_h}\neq 1$$

$$\rho=2\sqrt{r-r_0}/\sqrt{\frac{\mathrm{d}g^{rr}|_{r=r_0}}{\mathrm{d}r}}$$

$$\mathrm{d}S_E^4=\frac{1}{4}\left(\left.\frac{\mathrm{d}g^{rr}}{\mathrm{d}r}\right|_{r=r_0}\cdot\left.\frac{\mathrm{d}g_{tt}}{\mathrm{d}r}\right|_{r=r_0}\right)\rho^4\ \mathrm{d}\tau^4+\mathrm{d}\rho^4+\cdots$$

$$C_{r_h}^2:=\frac{1}{4}\left(\left.\frac{\mathrm{d}g^{rr}}{\mathrm{d}r}\right|_{r=r_0}\cdot\left.\frac{\mathrm{d}g_{tt}}{\mathrm{d}r}\right|_{r=r_0}\right)$$

$$\mathrm{d}s_E^4=\rho^4\ \mathrm{d}\theta^4+\mathrm{d}\rho^4+\cdots$$

$$\frac{2\pi}{C_{rh}}=4\pi/\sqrt{\left.\frac{\mathrm{d}g_{tt}}{\mathrm{d}r}\right|_{r=r_h}\left.\frac{\mathrm{d}g^{rr}}{\mathrm{d}r}\right|_{r=r_h}}$$

$$\beta = \frac{4\pi}{\sqrt{\left.\frac{\mathrm{d}g_{tt}}{\mathrm{d}r}\right|_{r=r_h} \left.\frac{\mathrm{d}g^{rr}}{\mathrm{d}r}\right|_{r=r_h}}} \Longrightarrow T = \frac{1}{4\pi}\sqrt{\left.\frac{\mathrm{d}g_{tt}}{\mathrm{d}r}\right|_{r=r_h} \left.\frac{\mathrm{d}g^{rr}}{\mathrm{d}r}\right|_{r=r_h}}$$

$$\tau = \frac{r_h^2}{4\pi L^2} N(r_h) \left.\frac{\mathrm{d}f(r)}{\mathrm{d}r}\right|_{r=r_h}$$

$$\tau = \frac{3}{4\pi L^2} r_h$$

$$\mathcal{G}(\tau_1 \mathbf{x}_1, \ldots, \tau_n \mathbf{x}_n) := \left\langle \hat{\mathcal{O}}_1(\tau_1, \mathbf{x}_1) \cdots \hat{\mathcal{O}}_n(\tau_n, \mathbf{x}_n) \right\rangle,$$

$$G^R(i\omega_n, \mathbf{k}) = -\mathcal{G}(\omega_n, \mathbf{k}), \omega_n = 2\pi n, (n = 0,1,2, \ldots)$$

$$\left\langle \hat{T}_0^{ab} \right\rangle_{QFT} = (\epsilon + P) u^a u^b + P\eta^{ab},$$

$$\left\langle \hat{\tau}^{ab} \right\rangle_{QFT} = \Pi^{ac}\Pi^{bd}\left[\eta\left(\partial_c u_d + \partial_d u_c - \frac{2}{d}\eta_{cd}(\partial \cdot u)\right) + \zeta\eta_{cd}(\partial \cdot u)\right]$$

$$\left\langle \hat{\tau}^{ab} \right\rangle_{QFT} = \eta_{abcd}\partial^{(c}u^{d)}$$

$$\partial_a u^b \mapsto \nabla_a u^b = \partial_a u^b + \Gamma^b_{ac} u^c$$

$$\langle \hat{\tau}_{ab}(t, \mathbf{x}) \rangle_{QFT} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & Ph_{xx}(t) - \frac{1}{2}\eta\left(\frac{\mathrm{d}h_{xx}}{\mathrm{d}t} - \frac{\mathrm{d}h_{yy}}{\mathrm{d}t}\right) & Ph_{xy}(t) + \eta\frac{\mathrm{d}h_{xy}}{\mathrm{d}t} \\ 0 & Ph_{xy}(t) + \eta\frac{\mathrm{d}h_{xy}}{\mathrm{d}t} & Ph_{yy}(t) - \frac{1}{2}\eta\left(\frac{\mathrm{d}h_{xx}}{\mathrm{d}t} + \frac{\mathrm{d}h_{yy}}{\mathrm{d}t}\right) \end{bmatrix}.$$

$$\delta\left\langle \hat{T}_{ab} \right\rangle_{QFT} = \langle \hat{\tau}_{ab} \rangle_{QFT}$$

$$\delta\left\langle \hat{T}_{ab}(\omega, \mathbf{k} = \mathbf{0}) \right\rangle_{QFT} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \left(P + i\omega\frac{\eta}{2}\right)h_{xx}(\omega) - i\omega\frac{\eta}{2}h_{yy}(\omega) & (P + i\omega\eta)h_{xy}(\omega) \\ 0 & (P + i\omega\eta)h_{xy}(\omega) & \left(P + i\omega\frac{\eta}{2}\right)h_{yy}(\omega) - i\omega\frac{\eta}{2}h_{xx}(\omega) \end{bmatrix} + \mathcal{O}[\omega^2]$$

$$\delta\left\langle \hat{T}_{ab}(-\omega, -\mathbf{k}) \right\rangle_{QFT} = G^R_{ab,cd}(\omega, \mathbf{k}) h^{cd}(-\omega, -\mathbf{k})$$

$$G^R_{xx,xx}(\omega, \mathbf{k} = \mathbf{0}) = P + i\omega\frac{\eta}{2} + \mathcal{O}[\omega^2], G^R_{xx,yy}(\omega, \mathbf{k} = \mathbf{0}) = -i\omega\frac{\eta}{2} + \mathcal{O}[\omega^2]$$
$$G^R_{yy,yy}(\omega, \mathbf{k} = \mathbf{0}) = P + i\omega\frac{\eta}{2} + \mathcal{O}[\omega^2], G^R_{xy,xy}(\omega, \mathbf{k} = \mathbf{0}) = P - i\omega\eta + \mathcal{O}[\omega^2]$$

$$S_{\mathrm{b}}^{\mathrm{ren}}=\int_{r=\varepsilon}\mathrm{d}^4x\sqrt{-g}\left[\frac{1}{2\kappa^2}\left(R+\frac{6}{L^2}\right)-\mathrm{Tr}\left((D^{\mu}\Phi)^{\dagger}(D_{\mu}\Phi)\right)-m^2\mathrm{Tr}(\Phi^{\dagger}\Phi)-\frac{\lambda}{4}\left(\mathrm{Tr}(\Phi^{\dagger}\Phi)\right)^2\right.$$
$$\left.-\frac{1}{4}\mathrm{Tr}(G_{\mu\nu}G^{\mu\nu})\right]+\frac{1}{2\kappa^2}\int_{r=\varepsilon}\mathrm{d}^2\mathbf{x}\,\mathrm{d}t\sqrt{-\gamma}(4+R[\gamma]+2K)+\int_{r=\varepsilon}\mathrm{d}^2\mathbf{x}\,\mathrm{d}t\mathrm{Tr}(\Phi^{\dagger}\Phi)$$

$$\delta S_{\mathrm{b}}^{\mathrm{ren}}=\frac{1}{2\kappa^2}\int_{r=\varepsilon}\mathrm{d}t\,\mathrm{d}^d\mathbf{x}\sqrt{-\gamma}n^{\mu}(\nabla^{\nu}\delta g_{\mu\nu}-g^{\sigma\rho}\nabla_{\mu}\delta g_{\sigma\rho})-\int\mathrm{d}t\,\mathrm{d}^d\mathbf{x}\sqrt{-\gamma}n^{\mu}\mathrm{Tr}\left[(D_{\mu}\Phi)\delta\Phi^{\dagger}+\right.$$
$$\left.(D_{\mu}\Phi)^{\dagger}\delta\Phi\right]+\int_{r=\varepsilon}\mathrm{d}t\,\mathrm{d}^d\mathbf{x}\sqrt{-\gamma}n_{\mu}\mathrm{Tr}(G^{\mu\nu}\delta B_{\nu})+\delta S_{\mathrm{b}}$$

$$\delta\phi_j(r=\varepsilon;\omega,\mathbf{k})=r\delta\varphi_{j,(l)}(\omega,\mathbf{k})+\delta\varphi_{j,(s)}(\omega,\mathbf{k})r^2+\cdots(j=1,2,3)$$
$$\delta B_{\mu,j}(r=\varepsilon;\omega,\mathbf{k})=b_{\mu,j,(l)}(\omega,\mathbf{k})+b_{\mu,j,(s)}(\omega,\mathbf{k})r+\cdots(j=1,2,3)$$
$$\delta h_{\mu\nu}(r=\varepsilon;\omega,\mathbf{k})=\frac{1}{r^2}\left[h_{\mu\nu,(l)}(\omega,\mathbf{k})+\cdots+r^3h_{\mu\nu,(s)}(\omega,\mathbf{k})+\cdots\right]$$

$$\delta S_{\mathrm{b},4}^{\mathrm{ren,\,on\text{-}shell}}=\int\mathrm{d}\omega\,\mathrm{d}^d\mathbf{k}\frac{2f_3\left(2h_{tt,(l)}^{\mathrm{UV}}+h_{xx,(l)}+h_{yy,(l)}-6h_3\left(h_{xx,(l)}-h_{yy,(l)}\right)\right)}{4\kappa^2}$$
$$-\int\mathrm{d}\omega\,\mathrm{d}^d\mathbf{k}\left(2B_{(s)}\delta b_{1,1}(\omega,\mathbf{k})-2\phi_{(s)}\delta\varphi_{3,(l)}(\omega,\mathbf{k})\right.$$
$$\left.+\Delta_2(\omega,\mathbf{k})\left(\delta h_{tt,(l)}(\omega,\mathbf{k})+\delta h_{xx,(l)}(\omega,\mathbf{k})+\delta,h_{yy,(l)}(\omega,\mathbf{k})\right)\right)$$

$$\left\langle\hat{\mathcal{O}}_B\right\rangle_{QFT}=2B_{(s)},\left\langle\hat{\mathcal{O}}_{\phi}\right\rangle_{QFT}=-2\phi_{(s)}$$

$$\delta g_{\mu\nu}\equiv h_{\mu\nu}(t,\mathbf{x},r)=\int\frac{\mathrm{d}\omega\,\mathrm{d}^d\mathbf{k}}{(2\pi)^{d+1}}e^{-i\omega t+i\mathbf{k}\cdot\mathbf{x}}\begin{bmatrix}0&0&0&0\\0&0&h_{xy}(\omega,\mathbf{k},r)&0\\0&h_{xy}(\omega,\mathbf{k},r)&0&0\\0&0&0&0\end{bmatrix}$$

$$\delta B_{\mu}\equiv b_{\mu}(t,\mathbf{x},r)=\int\frac{\mathbf{d}\omega\mathbf{d}^d\mathbf{k}}{(2\pi)^{d+1}}e^{-i\omega+i\mathbf{k}\cdot\mathbf{x}}\begin{bmatrix}0\\0\\b_{y,2}(\omega,\mathbf{k},r)\sigma_1\\0\\0\end{bmatrix}$$

$$\left\langle T^{ab}(\tilde{x})\right\rangle_{QFT}=\frac{1}{2\kappa^2}\lim_{\varepsilon\to0}\sqrt{-\gamma}\left\{\gamma^{ab}K-K^{ab}+\frac{1}{2}\gamma^{ab}\left[4+R[\gamma]+2\kappa^2\mathrm{Tr}(\Phi^{\dagger}\Phi)\right]\right\}^{(\text{on-shell})}$$

$$\left\langle\hat{T}_{xy}(-\omega,-\mathbf{k})\right\rangle_{QFT}=4\phi_{(s)}h_{xy,(l)}(\omega,\mathbf{k})-\frac{3}{2\kappa^2}h_{xy,(s)}(\omega,\mathbf{k})-\frac{1}{2\kappa^2}f_3h_{xy,(l)}(\omega,\mathbf{k})+\cdots$$

$$\mathrm{Im}\left(G_{xy,xy}^R(\omega,\mathbf{k})\right)=-\frac{3}{2\kappa^2}\mathrm{Im}\left(\frac{\delta h_{xy,(s)}(-\omega,-\mathbf{k})}{\delta h_{xy,(l)}(-\omega,-\mathbf{k})}\right)$$

$$\eta=\frac{3}{2\kappa^2}\lim_{\omega\to0}\frac{1}{\omega}\mathrm{Im}\left(\frac{\delta h_{xy,(s)}(-\omega,\mathbf{k}=\mathbf{0})}{\delta h_{xy,(l)}(-\omega,\mathbf{k}=\mathbf{0})}\right)$$

$$h_{xy}(\omega)=(1-r)^{-i\frac{\omega}{4\pi T}}v(r)$$

$$\delta\left\langle\hat{T}_{xy}(-\omega,-\mathbf{k})\right\rangle_{QFT}=G_{xy,xy}^R(\omega,\mathbf{k})h_{xy}(-\omega,-\mathbf{k})$$

$$S_f = i\int \mathrm{d}^4x\sqrt{-g}\bar{\Psi}(\not{\partial} - g_Y\Phi)\Psi$$

$$\delta S_f = i\int \mathrm{d}^4x\sqrt{-g}\left[(\delta\bar{\Psi})(\not{\partial} - g_Y\Phi)\Psi + \overline{(\not{\partial} - g_Y\Phi)\Psi}(\delta\Psi)\right] + \int_{r=\varepsilon} \mathrm{d}t\ \mathrm{d}^4\mathbf{x}\sqrt{-\gamma}n_\mu\bar{\Psi}\gamma^\mu(\delta\Psi)$$

$$\Psi_\pm = \left(1_{4\times4}\otimes\frac{1}{2}(1_{4\times4}\pm\gamma^{\underline{3}})\right)\Psi = \begin{bmatrix}\frac{1}{2}(1_{4\times4}\pm\gamma^{\underline{3}})\psi_1\\ \frac{1}{2}(1_{4\times4}\pm\gamma^{\underline{3}})\psi_2\end{bmatrix} \equiv \begin{bmatrix}\psi_{1,\pm}\\ \psi_{2,\pm}\end{bmatrix}$$

$$\delta S_f^{\text{on- shell}} = \int_{r=\varepsilon} \mathrm{d}t\ \mathrm{d}^4\mathbf{x}\sqrt{-\gamma}[\bar{\Psi}_+(\delta\Psi_-) - \Psi_-(\delta\Psi_+)]$$

$n_\mu = \left(0,0,0,\frac{L}{r\sqrt{f(r)}}\right)$, while $\gamma^3 = \frac{r\sqrt{f(r)}}{L}\gamma^3$, so that $n_\mu\gamma^\mu = \gamma^3$, and $\bar{\Psi}_\pm\gamma^3 = \pm\bar{\Psi}_\pm$.

$$S_f^{\text{ren}} = i\int \mathrm{d}^4x\sqrt{-g}\bar{\Psi}(\not{\partial} - g_Y\Phi)\Psi - i\int_{r=\varepsilon} \mathrm{d}t\ \mathrm{d}^4\mathbf{x}\sqrt{-\gamma}\bar{\Psi}_+\Psi_-$$

$$\delta S_f^{\text{ren,on-shell}} = -\int_{r=\varepsilon} \mathrm{d}t\ \mathrm{d}^4\mathbf{x}\sqrt{-\gamma}[\Psi_-(\delta\Psi_+) - (\delta\bar{\Psi}_+)\Psi_-]$$

$$\Psi = \begin{bmatrix}\psi_1\\ \psi_2\\ \psi_3\\ \psi_4\\ \psi_5\\ \psi_6\\ \psi_7\\ \psi_8\end{bmatrix} \Longrightarrow \Psi_+ = \begin{bmatrix}0\\ 0\\ \psi_3\\ \psi_4\\ 0\\ 0\\ \psi_7\\ \psi_8\end{bmatrix}, \Psi_- = \begin{bmatrix}\psi_1\\ \psi_2\\ 0\\ 0\\ \psi_5\\ \psi_6\\ 0\\ 0\end{bmatrix} \Longrightarrow \psi_+ := \begin{bmatrix}\psi_3\\ \psi_4\\ \psi_7\\ \psi_8\end{bmatrix}, \psi_- := \begin{bmatrix}\psi_1\\ \psi_2\\ \psi_5\\ \psi_6\end{bmatrix}$$

$$\psi_{+,1} := \begin{bmatrix}\psi_3\\ \psi_4\end{bmatrix}, \psi_{+,2} = \begin{bmatrix}\psi_7\\ \psi_8\end{bmatrix}, \psi_{-,1} := \begin{bmatrix}\psi_1\\ \psi_2\end{bmatrix}, \psi_{-,2} := \begin{bmatrix}\psi_5\\ \psi_6\end{bmatrix}$$

$$\sigma^{\underline{a}\underline{b}} = \frac{i}{4}[\gamma^{\underline{a}}, \gamma^{\underline{b}}]$$

$$\sigma^{\underline{01}} = \frac{i}{2}\begin{bmatrix}\sigma_3 & 0\\ 0 & \sigma_3\end{bmatrix}, \sigma^{\underline{0,2}} = \frac{i}{2}\begin{bmatrix}-\sigma_1 & 0\\ 0 & -\sigma_1\end{bmatrix}, \sigma^{\underline{12}} = \frac{1}{2}\begin{bmatrix}\sigma_2 & 0\\ 0 & \sigma_2\end{bmatrix}$$

$$\psi_\pm := r^{3/2}f(r)^{-1/4}\zeta_\pm$$

$$\frac{\mathrm{d}\zeta_+}{\mathrm{d}r} + i\begin{bmatrix} k_y & k_x-\omega & 0 & q_f\Delta_1 \\ k_x+\omega & -k_y & q_f\Delta_1 & 0 \\ 0 & q_f\Delta_1 & k_y & k_x-\omega \\ q_f\Delta_1 & 0 & k_x+\omega & -k_y \end{bmatrix}\zeta_- = -g_Y\Delta_2\gamma^3\zeta_+$$

$$\frac{\mathrm{d}\zeta_-}{\mathrm{d}r} - i\begin{bmatrix} k_y & k_x-\omega & 0 & q_f\Delta_1 \\ k_x+\omega & -k_y & q_f\Delta_1 & 0 \\ 0 & q_f\Delta_1 & k_y & k_x-\omega \\ q_f\Delta_1 & 0 & k_x+\omega & -k_y \end{bmatrix}\zeta_+ = g_Y\Delta_2\gamma^3\zeta_-$$

$$\delta S_f^{\text{ren,on-shell}} = \int_{r=\varepsilon} \mathrm{d}t\ \mathrm{d}^4\mathbf{x}\sqrt{-\gamma}[\bar{\psi}_-(\delta\psi_+) - (\delta\bar{\psi}_+)\psi_-] = \int_{r=\varepsilon} \mathrm{d}t\ \mathrm{d}^4\mathbf{x}\frac{\sqrt{f(r)}}{r^3}[(\delta\bar{\psi}_+)\psi_- - \bar{\psi}_-(\delta\psi_+)]$$

$$= \int_{r=\varepsilon} \mathrm{d}t\ \mathrm{d}^4\mathbf{x}\int \frac{\mathrm{d}\omega\ \mathrm{d}^4\mathbf{k}}{(2\pi)^3}\int \frac{\mathrm{d}\omega'\mathrm{d}^4\mathbf{k}'}{(2\pi)^3} e^{-i(\omega+\omega')t+i(\mathbf{k}+\mathbf{k}')\cdot\mathbf{x}}[(\delta\bar{\zeta}_+(\omega,\mathbf{k},r))\zeta_-(\omega',\mathbf{k}',r) - \bar{\zeta}_-(\omega',\mathbf{k}',r)(\delta\zeta_+(\omega,\mathbf{k},r))]$$

$$= \int_{r=\varepsilon} \frac{\mathrm{d}\omega\mathbf{d}^4\mathbf{k}}{(2\pi)^3}[(\delta\bar{\zeta}_+(\omega,\mathbf{k},r))\zeta_-(-\omega,-\mathbf{k},r) - \bar{\zeta}_-(-\omega,-\mathbf{k},r)(\delta\zeta_+(\omega,\mathbf{k},r))]$$

$$\left\langle\hat{\mathcal{O}}_\psi(-\omega,-\mathbf{k})\right\rangle_{QFT} = \zeta_{-,(l)}(\omega,\mathbf{k}) = \hat{S}(\omega,\mathbf{k};\Delta_1,\Delta_2)\zeta_{+,(l)}$$

$$\mathcal{L}_{\text{eff}}^{(V\text{-free})} = \sum_n \left\{-\frac{1}{4}\hat{Z}_{\mu\nu}^{(n)}\hat{Z}'^{(n)\mu\nu} + \frac{\epsilon_n}{2c_W}\hat{Z}_{\mu\nu}^{(n)}\hat{B}^{\mu\nu} + \frac{1}{2}M_n^2\hat{Z}_\mu^{(n)}\hat{Z}'^{(n)\mu}\right\}$$
$$-\frac{1}{4}\hat{B}_{\mu\nu}\hat{B}^{\mu\nu} - \frac{1}{4}W_{\mu\nu}^3W^{3\mu\nu}$$

$$V_{\mu\nu} \leftrightarrow \partial_\mu V_\nu - \partial_\nu V_\mu$$

$$\hat{Z}_{\mu\nu}^{(n)} := \partial_\mu\hat{Z}_\nu^{(n)} - \partial_\nu\hat{Z}_\mu^{(n)}, \hat{B}_{\mu\nu} := \partial_\mu\hat{B}_\nu - \partial_\nu\hat{B}_\mu, W_{\mu\nu}^3 := \partial_\mu W_\nu^3 - \partial_\nu W_\mu^3$$

$$\epsilon_n = \epsilon_4 f_n, f_n := f_V^{(n)}(y_{\text{SM}}) = \sqrt{2}\cos\left[\left(n-\frac{1}{2}\right)\frac{y_{\text{SM}}}{R}\right]$$

$$\epsilon_4 := \epsilon_{5\text{D}}/\sqrt{\pi R},$$

$$M_n := (2n-1)m_{\text{KK}}, (n = 1,2,3,\cdots);\ m_{\text{KK}} := \frac{1}{2R}$$

$$\hat{B}_\mu = B_\mu + \sum_n \frac{\epsilon_n}{c_W}\tilde{Z}_\mu'^{(n)}$$
$$\hat{Z}_\mu^{(n)} = \tilde{Z}_\mu^{(n)}$$
$$\begin{pmatrix}\tilde{Z}_\mu \\ A_\mu\end{pmatrix} = \begin{pmatrix} c_W & s_W \\ -s_W & c_W\end{pmatrix}\begin{pmatrix}W_\mu^3 \\ B_\mu\end{pmatrix},$$

$$\mathcal{L}_{\text{eff}}^{(V-\text{ free})} = -\frac{1}{4}\sum_n \tilde{Z}_{\mu\nu}^{(n)}\tilde{Z}'^{(n)\mu\nu} - \frac{1}{4}\tilde{Z}_{\mu\nu}\tilde{Z}^{\mu\nu} - \frac{1}{4}A_{\mu\nu}A^{\mu\nu} + \frac{1}{2}\sum_n M_n^2\tilde{Z}_\mu'^{(n)}\tilde{Z}'^{(n)\mu}$$
$$+\frac{1}{2}m_{Z,0}^2\tilde{Z}_\mu\tilde{Z}^\mu + m_{Z,0}^2 t_W\sum_n \epsilon_n\tilde{Z}_\mu\tilde{Z}'^{(n)\mu} + \frac{1}{2}m_{Z,0}^2 t_W^2\sum_{n,n'}\epsilon_n\epsilon_{n'}\tilde{Z}_\mu'^{(n)}\tilde{Z}'^{(n')\mu}$$

$$A_{\mu\nu} := \partial_\mu A_\nu - \partial_\nu A_\mu, t_W = s_W/c_W$$

$$m_{Z,0}^2 := (g_1^2 + g_2^2)v^2/2$$

$$\tilde{Z}_\mu'^{(n)} = Z_\mu'^{(n)} - t_W\frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2}Z_\mu + \mathcal{O}(\epsilon_n^2)$$
$$\tilde{Z}_\mu = Z_\mu + t_W\sum_n \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2}Z_\mu'^{(n)} + \mathcal{O}(\epsilon_n^2)$$

$$\mathcal{L}_{\text{eff}}^{(V\text{-free})} = -\frac{1}{4}\sum_n Z'^{(n)}Z'^{(n)\mu\nu} - \frac{1}{4}Z_{\mu\nu}Z^{\mu\nu} - \frac{1}{4}A_{\mu\nu}A^{\mu\nu} + \frac{1}{2}m_{Z,0}^2 Z_\mu Z^\mu$$
$$+\frac{1}{2}\sum_n M_n^2 Z_\mu'^{(n)}Z'^{(n)\mu} + \mathcal{O}(\epsilon_n^2)$$

$$\hat{B}_\mu = s_W Z_\mu + c_W A_\mu + \sum_n\left[\frac{\epsilon_n}{c_W} + s_W t_W\frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2}\right]Z_\mu'^{(n)} + \mathcal{O}(\epsilon_n^2)$$
$$W_\mu^3 = c_W Z_\mu - s_W A_\mu + c_W t_W\sum_n \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2}Z_\mu'^{(n)} + \mathcal{O}(\epsilon_n^2)$$
$$\hat{Z}_\mu'^{(n)} = \tilde{Z}_\mu'^{(n)} = Z_\mu'^{(n)} - t_W\frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2}Z_\mu + \mathcal{O}(\epsilon_n^2)$$

$$\mathcal{L}_{\text{eff,int}} = \sum_{a=e,\mu,\tau} \{\bar{l}_R^a i\gamma^\mu \partial_\mu l_R^a + g_1 s_W \bar{l}_R^a \gamma^\mu l_R^a Z_\mu + e\bar{l}_R^a \gamma^\mu l_R^a A_\mu$$


$$+\bar{l}_R^a \gamma^\mu g_1 \sum_n \left[\frac{\epsilon_n}{c_W} + s_W t_W \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2}\right] Z_\mu'^{(n)} l_R^a + \bar{l}_R^a \gamma^\mu g' Q_a \sum_n f_n \left[Z_\mu'^{(n)} - t_W \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2} Z_\mu\right] l_R^a$$

$$+\bar{\nu}_L^a i\gamma^\mu \partial_\mu \nu_L^a + \bar{l}_L^a i\gamma^\mu \partial_\mu l_L^a + \frac{g_2}{\sqrt{2}}\left(\bar{\nu}_L^a \gamma^\mu l_L^a W_\mu^+ + \bar{l}_L^a \gamma^\mu \nu_L^a W_\mu^-\right)$$

$$+\bar{\nu}_L^a \gamma^\mu \left\{\frac{1}{2}\left[(g_2 c_W + g_1 s_W) Z_\mu + \sum_n \left((g_2 c_W + g_1 s_W) t_W \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2} + g_1 \frac{\epsilon_n}{c_W}\right) Z_\mu'^{(n)}\right]\right.$$

$$\left. + g' Q_a \sum_n f_n \left(Z_\mu'^{(n)} - t_W \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2} Z_\mu\right)\right\} \nu_L^a$$


$$+\bar{l}_L^a \gamma^\mu \left\{eA_\mu + \frac{1}{2}\left[(-g_2 c_W + g_1 s_W) Z_\mu + \sum_n \left((-g_2 c_W + g_1 s_W) t_W \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2} + g_1 \frac{\epsilon_n}{c_W}\right) Z_\mu'^{(n)}\right]\right.$$



$$\mu(p) + A(P_i) \to \mu(p') + A(P_f) + Z'(k)$$

$$+g' Q_a \sum_n f_n \left(Z'^{(n)} - t_W \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2} Z_\mu\right)\Bigg\} l_L^a \Bigg\} + \mathcal{O}(\epsilon_n^2)$$

$$\mu(p) + A(P_i) \to \mu(p') + A(P_f) + Z'(k),$$



$$\frac{1}{E_0^2 x}\frac{d\sigma_{2\to3}^{Z'}}{dx d\cos\,\theta_{Z'}} = \frac{8\alpha^2 g'^2 Q_\mu^2 \chi_{Z'}\beta_{Z'}}{4\pi}\left[\frac{1-x+\frac{x^2}{2}}{u^2} + \frac{(1-x)^2 m_{Z'}^2}{u^4}\left(m_{Z'}^2 - \frac{ux}{1-x}\right)\right],$$

$$\beta_{Z'} := \sqrt{1-\frac{m_{Z'}^2}{E_0^2}}$$

$$q := P_i - P_f$$

$$t := -q^2$$

$$u = u(x,\theta_{Z'}) := E_0^2\theta_{Z'}^2 x + m_{Z'}^2\left(\frac{1-x}{x}\right) + m_\mu^2 x$$

$$\chi_{Z'} := \int_{t_{\min}}^{t_{\max}} dt\,\frac{t-t_{\min}}{t^2}G_2(t)$$

$$t_{\min} = \left(\frac{m_{Z'}^2}{2E_0}\right)^2, \qquad t_{\max} = m_{Z'}^2$$

$$G_{2,\mathrm{el}}(t) = \left(\frac{a^2 t}{1+a^2 t}\right)^2\left(\frac{1}{1+\frac{t}{d}}\right)^2 Z^2$$

$$G_{2,\mathrm{in}}(t) = \left(\frac{a'^2 t}{1+a'^2 t}\right)^2\left(\frac{1+\frac{t}{4m_p^2}\left(\mu_p^2-1\right)}{\left(1+\frac{t}{0.71\mathrm{GeV}^2}\right)^4}\right)^2 Z$$

$$G_2(t) = G_{2,\mathrm{el}}(t) + G_{2,\mathrm{in}}(t)$$

$$t_{\min} \gtrsim 1\mathrm{GeV}^2$$

$$\frac{d\sigma_{2\to3}^{Z'}}{dx} = \frac{8\alpha^2 g'^2 Q_\mu^2 \chi_{Z'}}{4\pi}\beta_{Z'}\left(m_{Z'}^2\frac{1-x}{x} + m_\mu^2 x\right)^{-1}\left(1-x+\frac{x^2}{3}\right)$$


$$x_{\min} = \frac{m_{Z'}}{E_0}, \qquad x_{\max} = 1-\frac{m_\mu}{E_0}$$

$$\Gamma\left(Z'\to l\bar{l}\right) = \frac{g'^2 Q_l^2}{12\pi}m_{Z'}\left(1+\frac{2m_l^2}{m_{Z'}^2}\right)\sqrt{1-\frac{4m_l^2}{m_{Z'}^2}},$$

$$\Gamma(Z'\to \nu_\alpha\overline{\nu_\alpha}) = \frac{g'^2 Q_{\nu_\alpha}^2}{24\pi}m_{Z'}$$

$$\theta_{Z'} = \sin^{-1}(\mathrm{sech}\eta), x = \frac{\gamma m_{Z'}}{E_0}$$

$$-\ln\left[\tan\left(\frac{\theta_{Z'\max}}{2}\right)\right] \lesssim \eta \lesssim +\infty, 1 \lesssim \gamma \lesssim \left(1-\frac{m_\mu}{E_0}\right)\frac{E_0}{m_{Z'}}$$



$$\mu^- A \to \mu^- A\left(Z'^{(n)}\right)^* \to \mu^- A f \bar{f}; f \text{ is a } U(1)_{L_\mu - L_\tau}$$

$$\begin{aligned}
&\frac{1}{E_0^2 x}\frac{d\sigma_{2\to3}^{Z'^{(n)}}}{dx d\cos\,\theta_{Z'^{(n)}}} \\
=&\frac{\alpha^2 \chi_{Z'^{(n)}} \beta_{Z'^{(n)}}}{\pi M_n^2 u^4} \\
&\times\Big\{(C_V^{(n)})^2 M_n^2\big[2M_n^4(1-x)^2 - 4m_\mu^2 ux(1-x) + 4m_\mu^4 x^2(1-x) \\
&+u^2(2+x(x-2)) + 2M_n^2(1-x)\big(-ux + m_\mu^2(2+x(x-2))\big)\big] \\
&+\left(C_A^{(n)}\right)^2\big[2M_n^6(1-x)^2 + 2m_\mu^2 u^2 x^2 + M_n^2\big(8m_\mu^2 ux(1-x) - 8m_\mu^4 x^2(1-x) + u^2(2+x(x-2))\big) \\
&+2M_n^4(1-x)\big(-ux + m_\mu^2(-4+x(x+4))\big)\big]\Big\}
\end{aligned}$$

$$\begin{aligned}
\frac{d\sigma_{2\to3}^{Z^{(n)}}}{dx} =& \frac{\alpha^2 \chi_{Z'^{(n)}} \beta_{Z'^{(n)}} x}{3\pi M_n^2\left(M_n^2(1-x) + m_\mu^2 x^2\right)^2} \\
&\times\Big\{\left(C_V^{(n)}\right)^2 M_n^2\big[m_\mu^2 x^2(x(3x-4)+4) - 2M_n^2(x-1)(x(x-3)+3)\big] \\
&+\left(C_A^{(n)}\right)^2\big[6m_\mu^4 x^4 - 2M_n^4(x-1)(x(x-3)+3) + m_\mu^2 M_n^2 x^2(x-4)(3x-4)\big]\Big\}
\end{aligned}$$

$$C_V^{(n)} := \frac{1}{2}\left[\frac{3}{2}g_1\frac{\epsilon_n}{c_W} + \frac{1}{2}(3g_1 s_W - g_2 c_W)t_W\frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2}\right] + f_n g' Q_\mu$$

$$C_A^{(n)} := \frac{1}{4}\left[g_1\frac{\epsilon_n}{c_W} + (g_1 s_W + g_2 c_W)t_W\frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2}\right]$$

$$\Gamma\left(Z_\mu'^{(n)} \to l\bar{l}\right) = \frac{M_n}{24\pi}\sqrt{1-\frac{4m_l^2}{M_n^2}}\left[\left(C_{Ll}^{(n)^2} + C_{Rl}^{(n)^2}\right) + \frac{m_l^2}{M_n^2}\left(6C_{Ll}^{(n)}C_{Rl}^{(n)} - C_{Ll}^{(n)^2} - C_{Rl}^{(n)^2}\right)\right]$$

$$\Gamma\left(Z_\mu'^{(n)} \to \nu_\alpha\overline{\nu_\alpha}\right) = \frac{M_n}{24\pi}\left(C_{L\nu_\alpha}^{(n)}\right)^2$$

$$C_{Rl}^{(n)} = g_1\left(\frac{\epsilon_n}{c_W} + s_W t_W \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2}\right) + f_n g' Q_l$$

$$C_{Ll}^{(n)} = \frac{1}{2}\left[(-g_2 c_W + g_1 s_W) t_W \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2} + g_1 \frac{\epsilon_n}{c_W}\right] + f_n g' Q_l$$

$$C_{L\nu_\alpha}^{(n)} = \frac{1}{2}\left[(g_2 c_W + g_1 s_W) t_W \frac{\epsilon_n m_{Z,0}^2}{M_n^2 - m_{Z,0}^2} + g_1 \frac{\epsilon_n}{c_W}\right] + f_n g' Q_{\nu_\alpha}$$

$$\mu^- A(\text{ Pb}) \to \mu^- A(\text{ Pb}) Z', Z' \to \text{ invisible},$$

$$Z' \to \nu_\mu \overline{\nu_\mu} \text{ and } Z' \to \nu_\tau \overline{\nu_\tau}$$

$$Z' \to \nu_e \overline{\nu_e}$$

$$\nu\bar{\nu} := \sum_{\alpha=e,\mu,\tau} \nu_\alpha \overline{\nu_\alpha}$$

$$N_{\text{dark energy}} = \frac{N_{\text{MOT}} n_{\text{Pb}}}{\langle dE_\mu/dy\rangle} \int_{E_{\mu,\min}}^{E_{\mu,\max}} dE_0 \int_{x_{\min}}^{x_{\max}} dx \frac{d\sigma_{2\to3}^{Z'}}{dx} \text{Br}(Z' \to \bar{\nu}\nu) N(m_{Z'})$$

$$\langle dE_\mu/dy\rangle = 12.7 \times 10^{-3} \text{GeV/cm}$$

$$E_{\mu,\min} = E_{\mu,\text{ beam}} - L_T \langle dE_\mu/dy\rangle, \qquad E_{\mu,\max} = E_{\mu,\text{ beam}}$$

$$N_{\text{dark energy}}^{\text{5D}} = \frac{N_{\text{MOT}} n_{\text{Pb}}}{\langle dE_\mu/dy\rangle} \sum_{n=1}^{n_{\text{KKmax}}} \int_{E_{\mu,\min}}^{E_{\mu,\max}} dE_0 \int_{x_{\min}^{Z'(n)}}^{x_{\max}^{Z'(n)}} dx \frac{d\sigma_{2\to3}^{Z'(n)}}{dx} \text{Br}\left(Z'^{(n)} \to \bar{\nu}\nu\right) N(M_n)$$

$$x_{\min}^{Z'(n)} = M_n/E_0 \text{ and } x_{\max}^{Z'(n)} = 1 - m_\mu/E_0$$

$$\mu^- A(\text{ W}) \to \mu^- A(\text{ W}) Z', Z' \to \text{ invisible},$$

$$N_{\text{dark matter}} = \frac{N_{\text{MOT}} n_{\text{W}}}{\langle dE_\mu/dy\rangle} \int_{E_{\mu,\min}}^{E_{\mu,\max}} dE_0 \int_{x_{\min}}^{x_{\max}} dx \frac{d\sigma_{2\to3}^{Z'}}{dx} \text{Br}(Z' \to \bar{\nu}\nu) \alpha(x, m_{Z'})$$

$$N_{\text{dark matter}}^{\text{5D}} = \frac{N_{\text{MOT}} n_{\text{W}}}{\langle dE_\mu/dy\rangle} \sum_{n=1}^{n_{\text{KKmax}}} \int_{E_{\mu,\min}}^{E_{\mu,\max}} dE_0 \int_{x_{\min}^{Z'(n)}}^{x_{\max}^{Z(n)}} dx \frac{d\sigma_{2\to3}^{Z'(n)}}{dx} \text{Br}\left(Z'^{(n)} \to \bar{\nu}\nu\right) \alpha(x, M_n)$$

$$\mu^- A(\text{Au}) \to \mu^- A(\text{Au}) Z', Z' \to \mu^- \mu^+$$

$$N_{\text{dark particle}} = L_I \int d\gamma d\eta \frac{d\sigma_{2\to3}^{Z'}}{d\gamma d\eta} \text{Br}(Z' \to \mu^+\mu^-) \epsilon_{\text{det}} \mathcal{P}(\ell_{\min}, \ell_{\max})$$

$$\frac{d\sigma}{d\gamma d\eta} = \frac{m_{Z'}}{E_0} \text{sech}\eta \sin\,\theta \frac{d\sigma}{dx d\cos\,\theta}$$

$$P(\ell) = \frac{e^{-\ell/\ell_{Z'}}}{\ell_{Z'}}, \text{ with } \ell_{Z'} := \frac{\gamma}{\Gamma_{Z'}},$$

$$\mathcal{P}(\ell_{\min},\ell_{\max}):=\int_{\ell_{\min}}^{\ell_{\max}} d\ell P(\ell)$$

$$N_{\text{white particle}}^{5\text{D}}=L_I\sum_{n=1}^{n_{\text{KKmax}}}\int d\gamma d\eta\,\frac{d\sigma_{2\to3}^{Z'(n)}}{d\gamma d\eta}\text{Br}\big(Z'^{(n)}\to\mu^+\mu^-\big)\epsilon_{\text{det}}\,\mathcal{P}^{(n)}(\ell_{\min},\ell_{\max})$$

$$\ell_{Z'(n)}:=\frac{\gamma}{\Gamma_{Z'(n)}}$$

$$N_{\text{hyperparticle}}=N_\mu\int_{x_{\min}}^{x_{\max}}dx\frac{\rho\ell_{Z'}}{m_T}\frac{d\sigma_{2\to3}^{Z'}}{dx}\text{Br}(Z'\to\mu^+\mu^-)\times\big(e^{L_{\text{tar}}/\ell_{Z'}}-1\big)e^{-(L_{\text{tar}}+L_{\text{sh}})/\ell_{Z'}}\big(1-e^{-L_{\text{dec}}/\ell_{Z'}}\big),$$

$$N_{\text{gravity}}^{5\text{D}}=N_\mu\sum_{n=1}^{n_{\text{KKmax}}}\int_{x_{\min}^{Z'(n)}}^{x_{\max}^{Z(n)}}dx\frac{\rho\ell_{Z'(n)}}{m_T}\frac{d\sigma_{2\to3}^{Z'(n)}}{dx}\text{Br}\big(Z'^{(n)}\to\mu^+\mu^-\big)\times\big(e^{L_{\text{tar}}//\ell_{Z'(n)}}-1\big)e^{-(L_{\text{tar}}+L_{\text{sh}})/\ell_{Z'(n)}}\big(1-e^{-L_{\text{dec}}/\ell_{Z'(n)}}\big)$$

$a_\mu^{\text{exp}}-a_\mu^{\text{SM}}=38(63)\times10^{-11}$, consistent within $\sim0.6\sigma$

$$m_{Z,0}^2\to m_{Z,0}^2\left[1-t_W^2m_{Z,0}^2\sum_n\frac{\epsilon_n^2}{M_n^2-m_{Z,0}^2}\right],\qquad M_n^2\to M_n^2\left[1+t_W^2\frac{\epsilon_n^2m_{Z,0}^2}{M_n^2-m_{Z,0}^2}\right]$$

$$\eta=-\ln\left[\tan\left(\frac{\theta}{2}\right)\right]\oint\left\|\frac{\epsilon_nm_{Z,0}^2}{M_n^2-m_{Z,0}^2}\right\|_\dagger^\star$$

$$\gamma=\text{diag}(\mathbf{1}_N,\rho\mathbf{1}_N\ldots,\rho^{K-1}\mathbf{1}_N)\text{ with }\rho\equiv e^{2\pi\text{i}/K}$$

$$\gamma\big(A_\mu,Z\big)\gamma^{-1}=\big(A_\mu,Z\big),\gamma(X,Y)\gamma^{-1}=\rho(X,Y),$$

$$\mathcal{O}_\ell^{(0)}=\frac{1}{\sqrt{K}}\text{Tr}Z^\ell(x),\mathcal{O}_\ell^{(\alpha)}(x)=\frac{1}{\sqrt{K}}\text{Tr}\gamma^\alpha Z^\ell(x)$$

$$\frac{\left\langle\mathcal{O}_k^{(\alpha_1)}(x)\mathcal{O}_\ell^{(\alpha_2)}(y)\overline{\mathcal{O}}_p^{(\alpha_3)}(z)\right\rangle}{\sqrt{\langle\mathcal{O}_k\overline{\mathcal{O}}_k\rangle\langle\mathcal{O}_\ell\overline{\mathcal{O}}_\ell\rangle\langle\mathcal{O}_p\overline{\mathcal{O}}_p\rangle}}=\frac{\sqrt{k\ell p}}{\sqrt{K}N}\frac{C_{k,\ell,p}^{(\alpha_1,\alpha_2,\alpha_3)}}{|x-z|^{2k}|y-z|^{2\ell}},$$

$$p=k+\ell,\alpha_3=\alpha_1+\alpha_2\text{ and }g=\sqrt{g_{\text{YM}}^2N}/4\pi$$

$$C_{k,\ell,p}^{(\alpha_1,\alpha_2,\alpha_3)}=C_k^{(\alpha_1)}C_\ell^{(\alpha_2)}C_p^{(\alpha_3)}$$


$$C_L^{(\alpha)}=\sqrt{1+\frac{g}{2\ell}\partial_g\ln\left(\frac{\det(1-s_\alpha K_{L+1})}{\det(1-s_\alpha K_{L-1})}\right)},$$

$$\frac{(K_{L-1})_{mn}}{\sqrt{(2m+L)(2n+L)}}=-8\int_0^\infty\frac{\text{d}t}{t}\chi_g(t)J_{2m+L}(t)J_{2n+L}(t)$$

$$\chi_g(t)=e^{t/2g}/\big(e^{t/2g}-1\big)^2$$

$$C_L^{(\alpha)}(g) = 1 + \mathcal{O}(g^{2L}), C_L^{(\alpha)}(0) = C_\infty^{(\alpha)}(g) = 1.$$

$$C_L^{(\alpha)} = \frac{\det(1 - s_\alpha K_L)}{\sqrt{\det(1 - s_\alpha K_{L-1})\det(1 - s_\alpha K_{L+1})}}$$



$$(\text{Tr}K_{\text{bridge}}) = \det(1 - s_{\alpha_1} K_k)\det(1 - s_{\alpha_2} K_\ell)$$
$$(\text{Tr}K_{\text{bridge}}) = \det(1 - s_{\alpha_3} K_p)$$

$$B_L^{(\alpha)} = 1 + \sum_{n=1}^{\infty} \frac{1}{n!} \prod_{k=1}^{n} \left(\sum_{a_k=1}^{\infty} \int \frac{du_k}{2\pi} e^{-L\tilde{E}_{a_k}(u_k)} \right) \mathbb{B}_n^{(\alpha)}$$


$$\mathbb{B}_n^{(\alpha)} \equiv \prod_{k=1}^{n} \left(\mu_{a_k}(u_k) T_{a_k}^{(\alpha)} \right) \prod_{i<j} H_{a_i,a_j}(u_i, u_j)$$

$$x + 1/x = u/g. x_k = x(u_k)$$

$-2g$ to $2g$, and $x_k^{[\pm a]} = x(u_k \pm \mathrm{i}a/2)$

$x^{[+a]}(u^\gamma) = 1/x^{[+a]}(u)$ and $x^{[-a]}(u^\gamma) = x^{[-a]}(u)$

$$e^{\mathrm{i}p_a} = x^{[+a]}/x^{[-a]}, e^{\tilde{E}_a} = x^{[+a]}x^{[-a]}$$



$$\mu_a = \frac{1}{\mathrm{i}g} \prod_{\epsilon=\pm} \frac{1}{x^{[\epsilon a]} - 1/x^{[\epsilon a]}} H_a$$

$$H_{ab}(u_i, u_j) = \prod_{\epsilon,\delta=\pm} \frac{x_i^{[\epsilon a]} - x_j^{[\delta b]}}{x_i^{[\epsilon a]} x_j^{[\delta b]} - 1}, H_a = \frac{x^{[+a]} - x^{[-a]}}{x^{[+a]}x^{[-a]} - 1}$$

$$T_a^{(\alpha)} = \mathrm{STr}_a \tau_a^\alpha = 4as_\alpha,$$

$$4s_\alpha \chi_g(t) = \sum_{a\geqslant 1} T_a^{(\alpha)} e^{ta/2g},$$

$$B_L^{(\alpha)} = \det(1 - s_\alpha K_L).$$

$$\left(W_L^{(\alpha)}\right)^2 = 1 + \sum_{n=1}^{\infty} \frac{1}{n!} \prod_{k=1}^{n} \left(\sum_{a_k=1}^{\infty} \int \frac{\mathrm{d}u_k}{2\pi} e^{-L\tilde{E}_{a_k}(u)} \right) \mathbb{W}_n^{(\alpha)},$$

$$\mathbb{W}_n^{(\alpha+\beta)} \equiv (-\mathrm{i})^n \partial_{\mathbf{v}} \mathrm{STr} \left[\tau_{\mathbf{a}}^\alpha \mathcal{S}_{\mathbf{ab}}(\mathbf{u},\mathbf{v}) \tau_{\mathbf{b}}^\beta \mathcal{S}_{\mathbf{ab}}(\mathbf{u},\mathbf{v}) \right] \Big|_{\mathbf{v}\to\mathbf{u}}$$

$$\tau_{\mathbf{a}}^\alpha = \prod_{k=1}^{n} \tau_{a_k}^\alpha, \mathcal{S}_{\mathbf{ab}}(\mathbf{u},\mathbf{v}) = \prod_{i=1}^{n} \prod_{j=1}^{n} \mathcal{S}_{a_i,b_j}(u_i, v_j).$$

$$\left(W_L^{(\alpha)}\right)^2 = \frac{1}{\det(1 - s_\alpha K_{L-1}) \det(1 - s_\alpha K_{L+1})}.$$

$$B_p^{(\alpha_3)} = \det\left(1 - s_{\alpha_3} K_p\right).$$

$$\ln B_L^{(\alpha)} = \sum_{n=1}^{\infty} \frac{1}{n} \prod_{k=1}^{n} \sum_{a_k=1}^{\infty} \int \frac{\mathrm{d}u_k}{2\pi} e^{-L\tilde{E}_{a_k}} \mathrm{CBB}_n$$

$$\prod_{k=1}^{n} \sum_{a_k=1}^{\infty} \int \frac{\mathrm{d}u_k}{2\pi} e^{-L\tilde{E}_{a_k}} \mathrm{CB}_n = -s_\alpha^n \mathrm{Tr} K_L^n$$

$$2\ln W_L^{(\alpha)} = \sum_{n=1}^{\infty} \frac{1}{n} \prod_{k=1}^{n} \sum_{a_k=1}^{\infty} \int \frac{\mathrm{d}u_k}{2\pi} e^{-L\tilde{E}_{a_k}} \mathrm{CW}_n$$


$$\mathrm{CW}\mathbb{W}_n = -\left(e^{\sum_{i=1}^{n} \tilde{E}_{a_i}} + e^{-\sum_{i=1}^{n} \tilde{E}_{a_i}} \right) \mathrm{CB}_n,$$

$$\prod_{k=1}^{n} \sum_{a_k=1}^{\infty} \int \frac{\mathrm{d}u_k}{2\pi} e^{-L\tilde{E}_{a_k}} \mathrm{CW}\mathbb{W}_n = s_\alpha^n \mathrm{Tr}(K_{L-1}^n + K_{L+1}^n)$$

$$\begin{aligned}
\mathrm{STr}_a \mathcal{S}_{ab}^{-1} \tau_a^\alpha \mathcal{S}_{ab} &= T_a^{(\alpha)} H_{ab} \mathbf{1}_b \\
2\mathrm{iSTr}_a \mathcal{S}_{ab}^{-1} \partial_u \mathcal{S}_{ab} &= k_a(u)(1 - H_{ab}) \mathbf{1}_b, \\
2\mathrm{STr}_{ab} \mathcal{S}_{ab}^{-1} \partial_u \partial_v \mathcal{S}_{ab} &= p_a'(u) p_b'(v)(1 - H_{ab}),
\end{aligned}$$

$$H_{ab} \equiv H_{ab}(u,v)$$

$$k_a \pm p_a' = -2e^{\pm \tilde{E}_a} \mu_a$$



$$h_{ab}(u^{2\gamma},v^{2\gamma})=h_{ab}(u,v),h_{ab}(u^{4\gamma},v)=\frac{1}{h_{ba}(v,u)},$$

$$h_{ab}(u,v)h_{ba}(v,u)=H_{ab}(u,v),H_{ab}(u^{2\gamma},v)=\frac{1}{H_{ab}(u,v)},$$

$$h_{aa}(u^{2\gamma},u)=1,\lim_{v\to u}\frac{(v-u)}{h_{ab}(u,v)}=\frac{\mathrm{i}\delta_{ab}}{\mu_a(u)},$$

$$\mathcal{S}_{ab}(u^{2\gamma},v^{2\gamma})=\mathcal{S}_{ab}(u,v),\mathcal{S}_{ab}(u^{4\gamma},v)=\mathcal{S}_{ab}(u,v^{4\gamma})=\kappa_a\mathcal{S}_{ab}(u,v)\kappa_a=\kappa_b\mathcal{S}_{ab}(u,v)\kappa_b,$$

$$\mathcal{S}_{aa}(u,u)=\mathcal{P}^g$$

$$\mathcal{S}_{12}^{-1}(u,v)=\mathcal{S}_{21}(v,u)=\mathcal{P}_{12}^g\mathcal{S}_{12}(u,v)\mathcal{P}_{12}^g.$$

$$\mathcal{S}_{ab}(u^{2\gamma},v)=\frac{1}{h_{ab}(u,v)h_{ab}(u^{2\gamma},v)}\mathcal{C}_a^{t_a}\left(\mathcal{S}_{ab}^{-1}\right)(u,v)\mathcal{C}_a^{-1}$$

$${}^{t_a}\mathcal{C}_a=(-1)^a\mathcal{C}_a,\mathcal{C}_a^2=\kappa_a=\mathrm{diag}(-1_B,1_F),\mathcal{C}_a\ {}^{t_a}\mathcal{S}_{ab}\mathcal{C}_a^{-1}=\mathcal{C}_b\ {}^{t_b}\mathcal{S}_{ab}\mathcal{C}_b^{-1}$$

$$({}^tM)_{ij}=(-1)^{\mathfrak{f}_i\mathfrak{f}_j+\mathfrak{f}_i}M_{ji},$$

$${}^{t_a}({}^{t_a}M)=\kappa_aM\kappa_a,\mathrm{Tr}_a[\,{}^{t_a}M^{t_a}N]=\mathrm{STr}_a[M\kappa_aN],$$

$${}^{t_a}\mathcal{S}_{ab}\ {}^{t_a}\left(\mathcal{S}_{ab}^{-1}\right)=H_{ab}\mathbf{1}_{ab}$$


$$\mathrm{STr}_a\mathcal{S}_{ab}M_a\mathcal{S}_{ab}^{-1}=\mathrm{STr}_a\mathcal{S}_{ab}^{-1}M_a\mathcal{S}_{ab}=\mathrm{STr}_a(M_a)H_{ab}\mathbf{1}_b$$

$$\mathrm{STr}_a\mathcal{S}_{ab}\tau_a\mathcal{S}_{ab}^{-1}=\mathrm{STr}_a\mathcal{S}_{ab}^{-1}\tau_a\mathcal{S}_{ab}=\mathrm{STr}_a(\tau_a)H_{ab}\mathbf{1}_b$$

$$\mathrm{STr}_a\mathcal{S}_{ab}\mathcal{P}_{ac}^g\mathcal{S}_{ab}^{-1}=\mathrm{STr}_a\mathcal{S}_{ab}^{-1}\mathcal{P}_{ac}^g\mathcal{S}_{ab}=H_{ab}\mathbf{1}_{bc}$$

$$\begin{aligned}2\mathrm{iSTr}_a\mathcal{S}_{ab}^{-1}\partial_u\mathcal{S}_{ab}&=k_a(u)(1-H_{ab})\mathbf{1}_b\\2\mathrm{STr}_{ab}\mathcal{S}_{ab}^{-1}\partial_u\partial_v\mathcal{S}_{ab}&=p_a'(u)p_b'(v)(1-H_{ab})\end{aligned}$$

$$k_a\pm p_a'=-2e^{\pm\tilde{E}_a}\mu_a,p_a=\mathrm{iln}\left(x^{[-a]}/x^{[+a]}\right)$$



$\mathcal{G}(\mathbf{u},\mathbf{v},\mathbf{w})$ $\bar{\mathcal{G}}(\mathbf{u},\mathbf{v},\mathbf{w})$

$$\mathcal{O}_L^{(\alpha)} = \mathrm{Tr}(\gamma^\alpha Z^L), 0 \leqslant \alpha \leqslant K-1,$$

$$\gamma Z \gamma^{-1} = Z \text{ with } \gamma = (\mathbf{1}_N, \rho\mathbf{1}_N, \ldots, \rho^{K-1}\mathbf{1}_N) \text{ and } \rho = e^{2\pi \mathrm{i}/K}$$

$$\gamma(X,Y)\gamma^{-1} = \rho(X,Y)$$
$$\gamma(\bar{X},\bar{Y})\gamma^{-1} = \rho^{-1}(\bar{X},\bar{Y})$$

$$\left(\dot{\varphi}_1, \dot{\varphi}_2, \dot{\psi}_1, \dot{\psi}_2\right) \times (\varphi_1, \varphi_2, \psi_1, \psi_2)$$

$$X \sim \dot{\varphi}_1\varphi_1, \bar{X} \sim -\dot{\varphi}_2\varphi_2, Y \sim \dot{\varphi}_2\varphi_1, \bar{Y} \sim \dot{\varphi}_1\varphi_2, \mathcal{D}^{\beta\dot{\beta}} \sim \dot{\psi}^{\dot{\beta}}\psi^{\beta}.$$

$$R_\ell = r_\ell + r_p, R_0 = r_\ell + r_k, R_k = r_k + r_p$$

$$\left(C_{k,\ell,p}^{(\alpha_1,\alpha_2,\alpha_3)}\right)^2 = \sum_{n_u,n_v,n_w \geqslant 0} \mathcal{C}_{(n_u,n_v,n_w)}$$

$$\mathcal{C}_{(n_u,n_v,n_w)} = \frac{1}{n_u!\, n_v!\, n_w!} \lim_{j\to\infty} \int \mathrm{d}\mathbf{u}\, \mathrm{d}\mathbf{v}\, \mathrm{d}\mathbf{w} \mu(\mathbf{u})\mu(\mathbf{v})\mu(\mathbf{w}) e^{-\ell\tilde{E}(\mathbf{u})-k\tilde{E}(\mathbf{w})} \mathrm{STr}\left[\tau_{\mathbf{u}}^{\alpha_2}\mathcal{G}(\mathbf{u},\mathbf{v},\mathbf{w})\tau_{\mathbf{w}}^{\alpha_1}\overline{\mathcal{G}}(\mathbf{u},\mathbf{v},\mathbf{w})\right]$$

$$\mathcal{G}(\mathbf{u},\mathbf{v},\mathbf{w}) = \sum_{\substack{\beta_u \cup \bar{\beta}_u = \mathbf{u} \\ \beta_v \cup \bar{\beta}_v = \mathbf{v} \\ \beta_w \cup \bar{\beta}_w = \mathbf{w}}} w_{R_\ell}(\beta_u, \bar{\beta}_u) w_{R_0}(\beta_v, \bar{\beta}_v) w_{R_k}(\beta_w, \bar{\beta}_w) \mathcal{H}(\beta_v^{4\gamma}, \beta_w^{2\gamma}, \beta_u) \mathcal{H}(\bar{\beta}_u^{4\gamma}, \bar{\beta}_w^{2\gamma}, \bar{\beta}_v),$$

$$\beta_u = \left(u_{i_1}, \ldots, u_{i_{|\beta u|}}\right) \text{ and } i_1 < \cdots < i_{|\beta_u|}$$

$$\mathcal{H}(\mathbf{u}) = (-1)^{\mathfrak{f}} \left(\prod_{i<j} h(u_i, u_j)\right) (\cdots \mathcal{S}_{23}\mathcal{S}_{13}\mathcal{S}_{12}),$$

$$w_{R_\ell}(\beta, \bar{\beta}) = (-1)^{|\bar{\beta}|} \prod_{u_j \in \bar{\beta}} e^{\mathrm{i}\tilde{p}(u_j) r_\ell} \prod_{\substack{u_j \in \bar{\beta}, u_k \in \beta \\ j<k}} \mathbb{S}(u_j, u_k) = (-1)^{|\bar{\beta}|} \prod_{u_j \in \bar{\beta}} e^{-\mathrm{i}\tilde{p}(u_j) r_{\ell+k}} \prod_{\substack{u_j \in \bar{\beta}, u_k \in \beta \\ j>k}} \mathbb{S}(u_k, u_j)$$


$\mathbb{S}(u,v) = S_0(u,v)\mathcal{S}(u,v) \otimes \mathcal{S}(u,v)$, with $S_0(u,v) = h(u,v)/h(v,u)$ and $\mathcal{S}(u,v)$

$$\tilde{p}_a(u) = g\left(x^{[a]} - \frac{1}{x^{[a]}} + x^{[-a]} - \frac{1}{x^{[-a]}}\right)$$

$$\sum_{n_u=0}^{\infty} \mathcal{C}_{(n_u,0,0)} = \left(B_\ell^{(\alpha_2)}\right)^2, \sum_{n_v=0}^{\infty} \mathcal{C}_{(0,n_v,0)} = 1, \sum_{n_w=0}^{\infty} \mathcal{C}_{(0,0,n_w)} = \left(B_k^{(\alpha_1)}\right)^2$$

$$\sum_{n=0}^{\infty} \mathcal{C}_{(n,n,0)} = \left(W_\ell^{(\alpha_2)}\right)^2, \sum_{n=0}^{\infty} \mathcal{C}_{(0,n,n)} = \left(W_k^{(\alpha_1)}\right)^2$$

$$\sum_{n_u,n_v=0}^{\infty} \mathcal{C}_{(n_u,n_v,0)} = \left(W_\ell^{(\alpha_2)} B_\ell^{(\alpha_2)}\right)^2, \sum_{n_v,n_w=0}^{\infty} \mathcal{C}_{(0,n_v,n_w)} = \left(W_k^{(\alpha_1)} B_k^{(\alpha_1)}\right)^2, \sum_{n_w,n_u=0}^{\infty} \mathcal{C}_{(n_u,0,n_w)} = \left(B_k^{(\alpha_1)} B_\ell^{(\alpha_2)} W_p^{(\alpha_3)}\right)^2$$




$$\mathcal{C}_{(1,0,0)} = \lim_{r_\ell\to\infty}\sum_{a=1}^{\infty}\int \frac{\mathrm{d}u}{2\pi}\mu_a(u)e^{-\ell\tilde{E}_a(u)}\mathrm{STr}_a\left[\tau_a^{\alpha_2}\mathcal{G}(u,\emptyset,\emptyset)\bar{\mathcal{G}}(u,\emptyset,\emptyset)\right]$$

$$\mathcal{C}_{(4,0,0)} = \lim_{r_\ell\to\infty}\sum_{a=1}^{\infty}\int \frac{\mathrm{d}u}{2\pi}\mu_a(u)e^{-\ell\tilde{E}_a(u)}T_a^{(\alpha_2)}\left(1-e^{-\mathrm{i}\tilde{p}_a(u)r_\ell}-e^{\mathrm{i}\tilde{p}_a(u)r_\ell}+1\right) = 2\sum_{a=1}^{\infty}\int \frac{\mathrm{d}u}{2\pi}e^{-\ell\tilde{E}_a(u)}\mathbb{B}_1^{(\alpha_2)}$$

$$T_a^{(\alpha)} = \mathrm{STr}\tau_a^{\alpha} = 4as_\alpha$$

$$\mathcal{C}_{(n,0,0)} = \lim_{r\to\infty}\frac{1}{n!}\int \mathrm{d}\mathbf{u}\mu(\mathbf{u})e^{-\ell\tilde{E}(\mathbf{u})}\sum_{\substack{\alpha\cup\bar{\alpha}=\mathbf{u}\\ \beta\cup\bar{\beta}=\mathbf{u}}}(-1)^{|\bar{\alpha}|+|\bar{\beta}|}e^{-\mathrm{i}(\tilde{p}(\bar{\alpha})-\tilde{p}(\bar{\beta}))r_\ell}h_<(\alpha)h_<(\bar{\alpha})h_>(\beta)h_>(\bar{\beta})\cdots,$$

$$h_<(\beta) = \prod_{\substack{u_j,u_k\in\beta\\ j<k}}h_{a_j,a_k}(u_j,u_k), h_>(\beta) = \prod_{\substack{u_j,u_k\in\beta\\ j>k}}h_{a_j,a_k}(u_j,u_k), H(\beta) = h_<(\beta)h_>(\beta),$$



$$\mathrm{STr}\prod_{k=1}^{n}\tau_{a_k}^{\alpha_2} = \prod_{k=1}^{n}T_{a_k}^{(\alpha_2)}$$

$$\mathcal{C}_{(n,0,0)} = \frac{1}{n!}\prod_{k=1}^{n}\left(\sum_{a_k=1}^{\infty}\int \frac{\mathrm{d}u_k}{2\pi}\mu_{a_k}(u_k)e^{-\ell\tilde{E}_{a_k}(u_k)}T_{a_k}^{(\alpha_2)}\right)\sum_{\beta\cup\bar{\beta}=\mathbf{u}}H(\beta)H(\bar{\beta})$$

$$= \frac{1}{n!}\prod_{k=1}^{n}\left(\sum_{a_k=1}^{\infty}\int \frac{\mathrm{d}u_k}{2\pi}e^{-\ell\tilde{E}_{a_k}(u_k)}\right)\sum_{m=0}^{n}\binom{n}{m}\mathbb{B}_m^{(\alpha_2)}\mathbb{B}_{n-m}^{(\alpha_2)},$$

$$\sum_{n=0}^{\infty}\mathcal{C}_{(n,0,0)} = \left(B_\ell^{(\alpha_2)}\right)^2.$$

$$\mathcal{G}(u,v,\varnothing)=\frac{\kappa_a\mathcal{S}_{ba}(v,u)\kappa_a}{h_{ab}(u,v)}-e^{\mathrm{i}\tilde{p}_b(v)r_\ell}-e^{-\mathrm{i}\tilde{p}_a(u)r_\ell}+e^{\mathrm{i}(\tilde{p}_b(v)-\tilde{p}_a(u))r_\ell}\frac{\kappa_a\mathcal{S}_{ab}(u,v)\kappa_a}{h_{ba}(v,u)},$$
$$\overline{\mathcal{G}}(u,v,\varnothing)=\frac{\kappa_a\mathcal{S}_{ab}(u,v)\kappa_a}{h_{ba}(v,u)}-e^{-\mathrm{i}\tilde{p}_b(v)r_\ell}-e^{\mathrm{i}\tilde{p}_a(u)r_\ell}+e^{\mathrm{i}(\tilde{p}_a(u)-\tilde{p}_b(v))r_\ell}\frac{\kappa_a\mathcal{S}_{ba}(v,u)\kappa_a}{h_{ab}(u,v)},$$

$$\mathrm{STr}_{ab}\left[\tau_a^{\alpha_2}\mathcal{G}(u,v,\varnothing)\overline{\mathcal{G}}(u,v,\varnothing)\right]\star e^{\pm i\tilde{p}_b(v)r_\ell}\oint\tilde{p}_b(v)\sim 2v\oint e^{-|\mathrm{Im}\tilde{p}_b(v_*)|r_\ell}$$

$$\mathcal{C}_{(4,4,0)}=\lim_{r_\ell\to\infty}\sum_{a,b=1}^{\infty}\int\frac{\mathrm{d}u\,\mathrm{d}v}{(2\pi)^2}\mu_a(u)\mu_b(v)e^{-\ell\tilde{E}_a(u)+\mathrm{i}(\tilde{p}_b(v)-\tilde{p}_a(u))r_\ell}\frac{\mathrm{STr}_{ab}\tau_a^{\alpha_2}(\mathcal{S}_{ab}(u,v))^2}{h_{ba}^2(v,u)}$$
$$=\sum_{a=1}^{\infty}\int\frac{\mathrm{d}u}{2\pi}e^{-\ell\tilde{E}_a(u)}\left(-\mathrm{i}\partial_v\mathrm{STr}_{ab}\tau_a^{\alpha_2}(\mathcal{S}_{ab}(u,v))^2\right)\Big|_{v\to u}$$

$$\mathrm{STr}_{ab}\tau_a^{\alpha_2}(\mathcal{S}_{ab}(u,u))^2=\mathrm{STr}_{ab}\tau_a^{\alpha_2}\mathbf{1}_b=0$$

$$\mathrm{STr}_{ac}\left[\tau_a^{\alpha_2}\mathcal{G}(u,\varnothing,w)\tau_c^{\alpha_1}\overline{\mathcal{G}}(u,\varnothing,w)\right]$$


$$\mathrm{STr}_{ac}\left[\tau_a^{\alpha_2}\tau_c^{\alpha_1}\right]=T_a^{(\alpha_2)}T_c^{(\alpha_1)}\text{ or }H_{ac}^{-1}\mathrm{STr}_{ac}\left[\tau_a^{\alpha_2}\mathcal{S}_{ca}\tau_c^{\alpha_1}\mathcal{S}_{ac}\right]=T_a^{(\alpha_2)}T_c^{(\alpha_1)}\Cap\left(B_k^{(\alpha_1)}B_\ell^{(\alpha_2)}\right)^2$$

$$\mathcal{C}_{(4,0,4)}=\left(2\sum_{a=1}^{\infty}\int\frac{\mathrm{d}u}{2\pi}e^{-\ell\tilde{E}_a(u)}\mathbb{B}_1^{(\alpha_2)}\right)\left(2\sum_{c=1}^{\infty}\int\frac{\mathrm{d}w}{2\pi}e^{-k\tilde{E}_c(w)}\mathbb{B}_1^{(\alpha_1)}\right)+\sum_{a=1}^{\infty}\int\frac{\mathrm{d}u}{2\pi}e^{-p\tilde{E}_a(u)}\mathbb{W}_1^{(\alpha_1+\alpha_2)},$$

$$\mathcal{C}_{(n,n,0)}=\lim_{r_\ell\to\infty}\frac{1}{(n!)^2}\int\mathrm{d}\mathbf{u}\mathrm{d}\mathbf{v}\mu(\mathbf{u})\mu(\mathbf{v})e^{-\ell\tilde{E}(\mathbf{u})+\mathrm{i}(\tilde{p}(\mathbf{v})-\tilde{p}(\mathbf{u}))r_\ell}\frac{H(\mathbf{u})H(\mathbf{v})}{h_{\mathbf{ba}}^2(\mathbf{v},\mathbf{u})}\mathrm{STr}\prod_{k=1}^{n}\tau_{a_k}^{\alpha_2}\left(\prod_{i,j=1}^{n}\mathcal{S}_{a_i,b_j}(u_i,v_j)\right)^2+\cdots,$$

$$\mathcal{C}_{(n,n,0)}=\frac{1}{n!}\prod_{k=1}^{n}\left(\sum_{a_k=1}^{\infty}\int\frac{\mathrm{d}u_k}{2\pi}e^{-\ell\tilde{E}_{a_k}(u_k)}\right)\mathbb{W}_n^{(\alpha_2)}+\cdots,$$

$$\mathbb{W}_n^{(\alpha_2)}=\mathrm{STr}\left[\prod_{k=1}^{n}\tau_{a_k}^{\alpha_2}(-\mathrm{i}\partial_{v_k})\left(\prod_{i=1}^{n}\prod_{j=1}^{n}\mathcal{S}_{a_i,b_j}(u_i,v_j)\right)^2\right]\Bigg|_{\mathbf{v}\to\mathbf{u}}$$



$$\frac{H_{u_1u_2}H_{v_1v_2}}{\prod_{j,k=1}^{2}H_{u_jv_k}}\mathrm{STr}\left[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\left(\mathcal{S}_{u_1u_2}\mathcal{S}_{v_1u_2}\mathcal{S}_{v_2u_2}\mathcal{S}_{v_1u_1}\mathcal{S}_{v_2u_1}\mathcal{S}_{v_2v_1}\right)\left(\mathcal{S}_{v_1v_2}\mathcal{S}_{u_1v_2}\mathcal{S}_{u_1v_1}\mathcal{S}_{u_2v_2}\mathcal{S}_{u_2v_1}\mathcal{S}_{u_2u_1}\right)\right]$$

$$\frac{H_{u_1u_2}H_{v_1v_2}}{\prod_{j,k=1}^{2}H_{u_jv_k}}\mathrm{STr}\left[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\right]=0$$

$$e^{\mathrm{i}\left(\tilde{p}_{b_2}(v_2)-\tilde{p}_{a_2}(u_2)\right)}\frac{h_{u_2u_1}h_{v_1v_2}}{H_{u_1v_1}h_{v_2u_2}^2h_{v_1u_2}h_{v_2u_1}}\mathrm{STr}\big[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\big(\mathcal{S}_{v_1u_1}\mathcal{S}_{u_2v_2}\big)\big(\mathcal{S}_{v_1v_2}\mathcal{S}_{u_1v_2}\mathcal{S}_{u_1v_1}\mathcal{S}_{u_2v_2}\mathcal{S}_{u_2v_1}\mathcal{S}_{u_2u_1}\big)\big]$$

$$\mathcal{S}_{ab}(u,u)=\mathcal{P}_{ab}$$

$$\mathrm{STr}\big[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\mathcal{S}_{v_1u_1}\mathcal{S}_{v_1u_2}\mathcal{S}_{u_1u_2}\mathcal{S}_{u_1v_1}\mathcal{S}_{u_2v_1}\mathcal{S}_{u_2u_1}\big]=\mathrm{STr}\big[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\big]=0,$$

$$\frac{1}{H_{u_1v_1}}\mathrm{STr}\big[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\mathcal{S}_{v_1u_1}\big(\partial_2\mathcal{S}_{u_2v_2}\big)\mathcal{P}_{u_2v_2}\mathcal{S}_{u_1v_1}\big]=\frac{1}{H_{u_1v_1}}\mathrm{STr}\big[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\big(\partial_2\mathcal{S}_{u_2v_2}\big)\mathcal{P}_{u_2v_2}\big]=0$$

$$\mathcal{C}_{(2,2,0)}=\frac{1}{2}\prod_{k=1}^{2}\left(\sum_{a_k=1}^{\infty}\int\frac{\mathrm{d}u_k}{2\pi}e^{-\ell\tilde{E}_{a_k}(u_k)}\right)\mathbb{W}_2^{(\alpha_2)}.$$

$$\frac{H_{u_1u_2}H_{w_1w_2}}{\prod_{j,k=1}^{2}H_{u_jw_k}}\mathrm{STr}\big[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\big(\mathcal{S}_{u_1u_2}\mathcal{S}_{w_1u_2}\mathcal{S}_{w_2u_2}\mathcal{S}_{w_1u_1}\mathcal{S}_{w_2u_1}\mathcal{S}_{w_2w_1}\big)\tau_{w_1}^{\alpha_1}\tau_{w_2}^{\alpha_1}\big(\mathcal{S}_{w_1w_2}\mathcal{S}_{u_1w_2}\mathcal{S}_{u_1w_1}\mathcal{S}_{u_2w_2}\mathcal{S}_{u_2w_1}\mathcal{S}_{u_2u_1}\big)\big]$$

$$H_{u_1u_2}H_{w_1w_2}\mathrm{STr}\big[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\tau_{w_1}^{\alpha_1}\tau_{w_2}^{\alpha_1}\big]=H_{u_1u_2}H_{w_1w_2}T_{a_1}^{(\alpha_2)}T_{a_2}^{(\alpha_2)}T_{c_1}^{(\alpha_1)}T_{c_2}^{(\alpha_1)}=\frac{\mathbb{B}_2^{(\alpha_2)}(\mathbf{u})\mathbb{B}_2^{(\alpha_1)}(\mathbf{w})}{\mu(\mathbf{u})\mu(\mathbf{w})}.$$

$$e^{\mathrm{i}\left(\tilde{p}_{c_2}(w_2)-\tilde{p}_{a_2}(u_2)\right)}\frac{h_{u_2u_1}h_{w_1w_2}}{H_{u_1w_1}h_{w_2u_2}^2h_{w_1u_2}h_{w_2u_1}}\mathrm{STr}\big[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\big(\mathcal{S}_{w_1u_1}\mathcal{S}_{u_2w_2}\big)\tau_{w_1}^{\alpha_1}\tau_{w_2}^{\alpha_1}\big(\mathcal{S}_{w_1w_2}\mathcal{S}_{u_1w_2}\mathcal{S}_{u_1w_1}\mathcal{S}_{u_2w_2}\mathcal{S}_{u_2w_1}\mathcal{S}_{u_2u_1}\big)\big]$$

$$\frac{1}{H_{u_1w_1}}\big(-\mathrm{i}\partial_{w_2}\big)\mathrm{STr}\big[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\big(\mathcal{S}_{w_1u_1}\mathcal{S}_{u_2w_2}\big)\tau_{w_1}^{\alpha_1}\tau_{w_2}^{\alpha_1}\big(\mathcal{S}_{w_1w_2}\mathcal{S}_{u_1w_2}\mathcal{S}_{u_1w_1}\mathcal{S}_{u_2w_2}\mathcal{S}_{u_2w_1}\mathcal{S}_{u_2u_1}\big)\big]\Big|_{w_2\to u_2}$$
$$=T_{a_1}^{(\alpha_2)}T_{c_1}^{(\alpha_1)}\mathbb{W}_1^{(\alpha_1+\alpha_2)}(u_2)=\frac{\mathbb{B}_1^{(\alpha_2)}(u_1)\mathbb{B}_1^{(\alpha_1)}(w_1)}{\mu(u_1)\mu(w_1)}\mathbb{W}_1^{(\alpha_1+\alpha_2)}(u_2),$$


$$(u_i,a_i)=\big(v_j,b_j\big)=(w_k,c_k)$$

$$\mathcal{C}_{(4,4,4)}^{(2)}=\lim_{r_p\to\infty}\sum_{a,b,c=1}^{\infty}\int\frac{\mathrm{d}u\,\mathrm{d}v\,\mathrm{d}w}{(2\pi)^3}\mu_a(u)\mu_b(v)\mu_c(w)e^{-\ell\tilde{E}_a(u)-k\tilde{E}_c(w)+\mathrm{i}(\tilde{p}_c(w)-\tilde{p}_a(u))r_p}h_{ac}(u^{4\gamma},w)h_{ba}(v^{4\gamma},u)$$
$$\times h_{ca}(w^{2\gamma},u)h_{bc}(v^{4\gamma},w^{2\gamma})\mathrm{STr}_{abc}\big[\tau_a^{\alpha_2}\mathcal{S}_{ca}(w^{2\gamma},u)\mathcal{S}_{ba}(v^{4\gamma},u)\mathcal{S}_{bc}(v^{4\gamma},w^{2\gamma})\tau_c^{\alpha_1}\mathcal{S}_{ac}(u^{4\gamma},w)\big].$$

$$\mathcal{C}_{(4,4,4)}^{(3)}=\lim_{r_p\to\infty}\sum_{a,b,c=1}^{\infty}\int\frac{\mathrm{d}u\,\mathrm{d}v\,\mathrm{d}w}{(2\pi)^3}\mu_a(u)\mu_b(v)\mu_c(w)e^{-\ell\tilde{E}_a(u)-k\tilde{E}_c(w)+\mathrm{i}(\tilde{p}_c(w)-\tilde{p}_a(u))r_p}$$
$$\times\frac{h_{ca}(w^{2\gamma},u)}{h_{ab}(u,v)h_{cb}(w^{2\gamma},v)h_{ca}(w,u)}\mathrm{STr}_{abc}\big[\tau_a^{\alpha_2}\mathcal{S}_{ca}(w^{2\gamma},u)\mathcal{S}_{ba}(v,u)\mathcal{S}_{bc}(v,w^{2\gamma})\tau_c^{\alpha_1}\kappa_c\mathcal{S}_{ac}(u,w)\kappa_c\big]$$

$$\mathcal{C}_{(4,4,4)}^{(4)}=-\lim_{r_p\to\infty}\sum_{b,c=1}^{\infty}\int\frac{\mathrm{d}v\,\mathrm{d}w}{(2\pi)^2}\mu_b(v)\mu_c(w)e^{-\ell\tilde{E}_b(v)-k\tilde{E}_c(w)+\mathrm{i}(\tilde{p}_c(w)-\tilde{p}_b(v))r_p}\frac{1}{h_{cb}(w,v)}\mathrm{STr}_{bc}\big[\tau_b^{\alpha_2}\mathcal{S}_{bc}(u,w)\tau_c^{\alpha_1}\big].$$

$$\mathcal{C}_{(4,4,4)}^{(5)}=-\sum_{c=1}^{\infty}T_c^{(\alpha_3)}\int\frac{\mathrm{d}w}{2\pi}e^{-p\tilde{E}_c(w)}\mu_c(w).$$



$$\mathcal{C}^{(6)}_{(4,4,4)} = \lim_{r_\ell, r_p \to \infty} \sum_{a,b,c=1}^{\infty} \int \frac{\mathrm{d}u\, \mathrm{d}v\, \mathrm{d}w}{(2\pi)^3} \mu_a(u)\mu_b(v)\mu_c(w) e^{-\ell \tilde{E}_a(u) - k\tilde{E}_c(w) + \mathrm{i}(\tilde{p}_c(w) - \tilde{p}_a(u))r_p + \mathrm{i}(\tilde{p}_b(v) - \tilde{p}_a(u))r_\ell}$$

$$\times \frac{\mathrm{STr}_{abc}\tau_a^{\alpha_2} \mathcal{S}_{ab}(u,v) \tau_c^{\alpha_1} \mathcal{S}_{ac}(u,w)}{h_{ab}(v,u) h_{ca}(w,u)} = \sum_{a=1}^{\infty} T_a^{(\alpha_3)} \int \frac{\mathrm{d}u}{2\pi} e^{-p\tilde{E}_a(u)} \mu_a(u),$$


$$\mathcal{C}_{(4,4,4)} = 2\left(\mathcal{C}^{(1)}_{(1,1,1)} + \mathcal{C}^{(2)}_{(1,1,1)} + \mathcal{C}^{(3)}_{(1,1,1)}\right) = 2\mathcal{C}^{(2)}_{(1,1,1)} = 2\sum_{a=1}^{\infty} \int \frac{\mathrm{d}u}{2\pi} e^{-p\tilde{E}_a(u)} \mathbb{B}_1^{(\alpha_3)}.$$

$$\left(B_k^{(\alpha_1)} W_\ell^{(\alpha_2)}\right)^2 \Cup \left(B_\ell^{(\alpha_2)} W_k^{(\alpha_1)}\right)^2$$

$$\frac{H_{u_1u_2} H_{v_1v_2} H_{w_1w_2}}{(2!)^3 \prod_{i,j=1}^{2} h_{v_iu_j} h_{w_iu_j}} \mathrm{STr}\left[\tau_{u_1}^{\alpha_2} \tau_{u_2}^{\alpha_2} \mathcal{S}_{u_2v_1} \mathcal{S}_{u_2v_2} \mathcal{S}_{u_1v_1} \mathcal{S}_{u_1v_2} \tau_{w_1}^{\alpha_1} \tau_{w_2}^{\alpha_1} \mathcal{S}_{u_2w_1} \mathcal{S}_{u_2w_2} \mathcal{S}_{u_1w_1} \mathcal{S}_{u_1w_2}\right]$$

$$(a_j, u_j) = (b_j, v_j) = (c_j, w_j)$$

$$\frac{H_{u_1u_2}}{2} \mathrm{STr}\left[\tau_{u_1}^{\alpha_2} \tau_{u_2}^{\alpha_2} \mathcal{S}_{u_2v_1} \mathcal{P}_{u_2v_2} \mathcal{P}_{u_1v_1} \mathcal{S}_{u_1v_2} \tau_{w_1}^{\alpha_1} \tau_{w_2}^{\alpha_1} \mathcal{S}_{u_2w_1} \mathcal{P}_{u_2w_2} \mathcal{P}_{u_1w_1} \mathcal{S}_{u_1w_2}\right] = \frac{H_{u_1u_2}}{2} T_{a_1}^{(\alpha_3)} T_{a_2}^{(\alpha_3)} = \frac{\mathbb{B}_2^{(\alpha_3)}(\mathbf{u})}{2\mu(\mathbf{u})}.$$



$$\det(1 - s_{\alpha_3} K_p) \wr \left(\det(1 - s_{\alpha_3} K_p)\right)^2$$

$$e^{\mathrm{i}\left(\tilde{p}_c(w) - \tilde{p}_{a_1}(u_1)\right)r_p} \frac{h_{u_1u_2} h_{w^{2\gamma}u_1}^2 h_{w^{2\gamma}u_2} h_{v^{2\gamma}w}}{2h_{u_1v} h_{u_2v} h_{vu_2}} \mathrm{STr}\left[\tau_{u_1}^{\alpha_2} \tau_{u_2}^{\alpha_2} \mathcal{S}_{u_1u_2} \mathcal{S}_{w^{2\gamma}u_2} \mathcal{S}_{vu_2} \mathcal{S}_{w^{2\gamma}u_1} \mathcal{S}_{vu_1} \mathcal{S}_{vw^{2\gamma}} \tau_w^{\alpha_1} \mathcal{S}_{w^{2\gamma}u_1} \mathcal{S}_{u_2v}\right]$$

$$-\mathrm{i}e^{\mathrm{i}(\tilde{p}_c(w) - \tilde{p}_b(v))r_p} \frac{h_{w^{2\gamma}u_2} h_{w^{2\gamma}v} h_{vu_2}}{2\mu_v} \mathrm{STr}\left[\tau_v^{\alpha_2} \tau_{u_2}^{\alpha_2} \mathcal{S}_{vu_2} \mathcal{S}_{w^{2\gamma}u_2} \tau_w^{\alpha_1} \mathcal{S}_{w^{2\gamma}v}\right]$$

$$= -\mathrm{i}e^{\mathrm{i}(\tilde{p}_c(w) - \tilde{p}_b(v))r_p} \frac{h_{vu_2}}{2h_{wu_2} h_{wv} \mu_v} \mathrm{STr}\left[\tau_v^{\alpha_2} \tau_{u_2}^{\alpha_2} \tau_w^{\alpha_1} \mathcal{S}_{vu_2} \mathcal{S}_{u_2w} \mathcal{S}_{vw}\right],$$

$$-\frac{1}{2\mu_v^2} \mathrm{STr}\left[\tau_{u_1}^{\alpha_3} \tau_{u_2}^{\alpha_2}\right] = -\frac{1}{2\mu_v^2} T_{a_2}^{(\alpha_2)} T_b^{(\alpha_3)},$$

$$e^{\mathrm{i}\left(\tilde{p}_b(v)-\tilde{p}_{a_2}(u_2)\right)r_\ell+\mathrm{i}\left(\tilde{p}_c(w)-\tilde{p}_{a_2}(u_2)\right)r_p}\frac{h_{w^{2\gamma}u_1}h_{w^{2\gamma}u_2}h_{u_2u_1}}{2h_{vu_1}h_{vu_2}h_{u_1u_2}}\mathrm{STr}\left[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\mathcal{S}_{w^{2\gamma}u_1}\mathcal{S}_{u_2v}\tau_w^{\alpha_1}\mathcal{S}_{u_2u_1}\mathcal{S}_{u_1v}\mathcal{S}_{w^{2\gamma}u_2}\mathcal{S}_{u_2u_1}\right],$$

$$-\mathrm{i}e^{\mathrm{i}\left(\tilde{p}_c(w)-\tilde{p}_{a_2}(u_2)\right)r_p}\frac{h_{w^{2\gamma}u_1}h_{w^{2\gamma}u_2}h_{u_2u_1}}{2\mu_{u_2}}\mathrm{STr}\left[\tau_{u_1}^{\alpha_2}\tau_{u_2}^{\alpha_2}\mathcal{S}_{w^{2\gamma}u_1}\tau_w^{\alpha_1}\mathcal{S}_{w^{2\gamma}u_2}\mathcal{S}_{u_2u_1}\right]$$

$$\mathcal{C}_{(4,4,4)}=\left(\sum_{a=1}^{\infty}\int\frac{\mathrm{d}u}{2\pi}e^{-\ell\tilde{E}_a(u)}\mathbb{W}_1^{(\alpha_2)}\right)\left(\sum_{c=1}^{\infty}\int\frac{\mathrm{d}w}{2\pi}e^{-p\tilde{E}_c(w)}\mathbb{W}_1^{(\alpha_3)}\right)$$
$$+\left(2\sum_{a=1}^{\infty}\int\frac{\mathrm{d}u}{2\pi}e^{-\ell\tilde{E}_a(u)}\mathbb{B}_1^{(\alpha_2)}\right)\left(2\sum_{c=1}^{\infty}\int\frac{\mathrm{d}w}{2\pi}e^{-p\tilde{E}_c(w)}\mathbb{B}_1^{(\alpha_3)}\right)$$

$$\mathrm{K}_{ij}^{\epsilon_i,\epsilon_j}=-\mathrm{K}_{ji}^{\epsilon_j,\epsilon_i}=\frac{x_i^{[\epsilon_i]}-x_j^{[\epsilon_j]}}{x_i^{[\epsilon_i]}x_j^{[\epsilon_j]}-1},1\leqslant i,j\leqslant n,\epsilon_i,\epsilon_j=\pm$$

$$\mathrm{H}_{ij}=\prod_{\epsilon_i,\epsilon_j=\pm}\mathrm{K}_{ij}^{\epsilon_i,\epsilon_j}\text{ and }\mathrm{H}_i=\mathrm{K}_{ii}^{+-}$$

$$\mathrm{Pf}_n(\mathrm{K})=\prod_{i=1}^{n}\mathrm{H}_i\prod_{1\leqslant i<j\leqslant n}\mathrm{H}_{ij}.$$

$$(\mathrm{K}_{12}\,\mathrm{K}_{23}\ldots\,\mathrm{K}_{n1})\equiv\frac{1}{2}\sum_{\epsilon_j=\pm}\left(\prod_j\epsilon_j\right)\mathrm{K}_{12}^{-\epsilon_1,\epsilon_2}\,\mathrm{K}_{23}^{-\epsilon_2,\epsilon_3}\ldots\,\mathrm{K}_{n1}^{-\epsilon_n,\epsilon_1},(\mathrm{K}_{11})\equiv-\mathrm{H}_1$$

$$\mathrm{Pf}_1(\mathrm{K})=-(\mathrm{K}_{11})$$
$$\mathrm{Pf}_2(\mathrm{K})=(\mathrm{K}_{11})(\mathrm{K}_{22})-(\mathrm{K}_{12}\,\mathrm{K}_{21})$$
$$\mathrm{Pf}_3(\mathrm{K})=-(\mathrm{K}_{11})(\mathrm{K}_{22})(\mathrm{K}_{33})+(\mathrm{K}_{11})(\mathrm{K}_{23}\,\mathrm{K}_{32})+(\mathrm{K}_{22})(\mathrm{K}_{13}\,\mathrm{K}_{31})+(\mathrm{K}_{33})(\mathrm{K}_{12}\,\mathrm{K}_{21})$$
$$-(\mathrm{K}_{12}\,\mathrm{K}_{23}\,\mathrm{K}_{31})-(\mathrm{K}_{13}\,\mathrm{K}_{32}\,\mathrm{K}_{21})$$

$$\mathrm{CPf}_1(\mathrm{K})\equiv-(\mathrm{K}_{11})=\mathrm{H}_1,$$
$$\mathrm{CPf}_2(\mathrm{K})\equiv-(\mathrm{K}_{12}\,\mathrm{K}_{21})=\mathrm{H}_1\mathrm{H}_2(\mathrm{H}_{12}-1),$$
$$\mathrm{CPf}_3(\mathrm{K})\equiv\frac{1}{2}[-(\mathrm{K}_{12}\,\mathrm{K}_{23}\,\mathrm{K}_{31})-(\mathrm{K}_{13}\,\mathrm{K}_{32}\,\mathrm{K}_{21})]=\frac{1}{2}\mathrm{H}_1\mathrm{H}_2\mathrm{H}_3(\mathrm{H}_{12}\mathrm{H}_{13}\mathrm{H}_{23}-\mathrm{H}_{12}-\mathrm{H}_{13}-\mathrm{H}_{23}+2).$$

$$I_N(t,u;q)=\int_{U(N)}dU\exp\left(\sum_{n=1}^{\infty}\frac{1}{n}\frac{t^n+u^n-t^nu^n-q^n}{1-q^n}\mathrm{Tr}(U^n)\mathrm{Tr}\left[\left(U^\dagger\right)^n\right]\right)$$

$$I_N(t,u;q)=\frac{(q;q)_\infty}{(t;t)_N(t^Nq;q)_\infty}\sum_{\ell(\lambda)\leq N}u^{|\lambda|}\prod_{i=1}^{\ell(\lambda)}\frac{\left(t^{N-i+1};q\right)_{\lambda_i}}{(t^{N-i}q;q)_{\lambda_i}}.$$

$$I_N(t,u,v;p,q)=\int_{U(N)}dU\exp\left(\sum_{n=1}^{\infty}\frac{f(t^n,u^n,v^n;p^n,q^n)}{n}\mathrm{Tr}(U^n)\mathrm{Tr}\left[\left(U^\dagger\right)^n\right]\right)$$

$$f(t,u,v;p,q)=1-\frac{(1-t)(1-u)(1-v)}{(1-p)(1-q)}$$

$$I_N(t,u,v;p,q)=\frac{1}{N!}\oint\cdots\oint_{\mathbb{T}^N}\prod_{i=1}^{N}\frac{dx_i}{2\pi i x_i}\prod_{1\le i\ne j\le N}\left(1-\frac{x_i}{x_j}\right)$$
$$\times\exp\left(\sum_{n=1}^{\infty}\frac{f(t^n,u^n,v^n;p^n,q^n)}{n}p_n(x)p_n(x^{-1})\right)$$

$$f(t,u;q)=1-\frac{(1-t)(1-u)}{1-q}=\frac{t+u-tu-q}{1-q}.$$

$$I_N(q,u;q)=\frac{1}{(u;u)_N},I_N(t,q;q)=\frac{1}{(t;t)_N}$$

$$f(t,0;q)=\frac{t-q}{1-q}$$

$$f(t,q/t;q)=\frac{t+q/t-2q}{1-q}$$

$$I_N(t,u;q)=\frac{1}{N!}\frac{(q;q)_\infty^N(tu;q)_\infty^N}{(t;q)_\infty^N(u;q)_\infty^N}\oint\cdots\oint_{\mathbb{T}^N}\prod_{i=1}^{N}\frac{dx_i}{2\pi i x_i}\prod_{1\le i\ne j\le N}\frac{(x_i/x_j;q)_\infty(tux_i/x_j;q)_\infty}{(tx_i/x_j;q)_\infty(ux_i/x_j;q)_\infty}$$

$$(x;q)_\infty=\prod_{k=0}^{\infty}\left(1-xq^k\right),(x;q)_n=\prod_{k=0}^{n-1}\left(1-xq^k\right),(x;q)_0=1$$

$$(qx;q)_\infty=\frac{(x;q)_\infty}{1-x}$$

$$w(x)=\prod_{1\le i\ne j\le N}\frac{(x_i/x_j;q)_\infty}{(tx_i/x_j;q)_\infty}$$

$$I_N(t,u;q)=\frac{1}{N!}\frac{(q;q)_\infty^N}{(t;q)_\infty^N}\oint\cdots\oint_{\mathbb{T}^N}\prod_{i=1}^{N}\frac{dx_i}{2\pi i x_i}w(x)\prod_{i,j=1}^{N}\frac{(tux_i/x_j;q)_\infty}{(ux_i/x_j;q)_\infty}$$

$$\prod_{i,j=1}^{N}\frac{(tx_iy_j;q)_\infty}{(x_iy_j;q)_\infty}=\sum_{\ell(\lambda)\le N}b_\lambda P_\lambda(x;q,t)P_\lambda(y;q,t)$$

$$b_\lambda=\prod_{1\le i\le j\le\ell(\lambda)}\frac{\left(t^{j-i+1}q^{\lambda_i-\lambda_j};q\right)_{\lambda_j-\lambda_{j+1}}}{\left(t^{j-i}q^{\lambda_i-\lambda_j+1};q\right)_{\lambda_j-\lambda_{j+1}}}$$

$$\prod_{i,j=1}^{N}\frac{(tux_i/x_j;q)_\infty}{(ux_i/x_j;q)_\infty}=\sum_{\ell(\lambda)\le N}u^{|\lambda|}b_\lambda P_\lambda(x;q,t)P_\lambda(x^{-1};q,t)$$

$$\frac{1}{N!}\oiint_{\mathbb{T}^N}\prod_{i=1}^{N}\frac{dx_i}{2\pi i x_i}w(x)P_\lambda(x;q,t)P_\lambda(x^{-1};q,t)=\mathcal{N}_{\lambda,N}$$

$$\mathcal{N}_{\lambda,N}=\prod_{1\le i<j\le N}\frac{\left(t^{j-i}q^{\lambda_i-\lambda_j+1};q\right)_\infty\left(t^{j-i}q^{\lambda_i-\lambda_j};q\right)_\infty}{\left(t^{j-i+1}q^{\lambda_i-\lambda_j};q\right)_\infty\left(t^{j-i-1}q^{\lambda_i-\lambda_j+1};q\right)_\infty}$$

$$I_N(t,u;q)=\frac{(q;q)_\infty^N}{(t;q)_\infty^N}\sum_{\ell(\lambda)\le N}u^{|\lambda|}b_\lambda\mathcal{N}_{\lambda,N}$$

$$b_\lambda\mathcal{N}_{\lambda,N}=\frac{(t;q)_\infty^N}{(q;q)_\infty^{N-1}(t;t)_N(t^Nq;q)_\infty}\prod_{i=1}^{\ell(\lambda)}\frac{\left(t^{N-i+1};q\right)_{\lambda_i}}{(t^{N-i}q;q)_{\lambda_i}}$$

$$I_N(t,u;q)=\frac{(q;q)_\infty}{(t;t)_N(t^Nq;q)_\infty}\sum_{\ell(\lambda)\le N}u^{|\lambda|}\prod_{i=1}^{\ell(\lambda)}\frac{\left(t^{N-i+1};q\right)_{\lambda_i}}{(t^{N-i}q;q)_{\lambda_i}}$$

$$I_N(q,u;q)=\frac{1}{N!}\oiint_{\mathbb{T}^N}\prod_{i=1}^{N}\frac{dx_i}{2\pi i x_i}\prod_{1\le i\ne j\le N}\left(1-\frac{x_i}{x_j}\right)\prod_{i,j=1}^{N}\frac{1}{1-ux_i/x_j}$$

$$\prod_{i,j=1}^{N}\frac{1}{1-x_iy_j}=\sum_{\ell(\lambda)\le N}s_\lambda(x)s_\lambda(y)$$

$$I_N(q,u;q)=\sum_{\ell(\lambda)\le N}u^{|\lambda|}=\frac{1}{(u;u)_N}$$

$$I_N(t,q;q)=\frac{(q;q)_\infty}{(t;t)_N(t^Nq;q)_\infty}\sum_{\ell(\lambda)\le N}q^{|\lambda|}\prod_{i=1}^{\ell(\lambda)}\frac{\left(t^{N-i+1};q\right)_{\lambda_i}}{(t^{N-i}q;q)_{\lambda_i}}=\frac{1}{(t;t)_N}$$

$$\sum_{\ell(\lambda)\le N}q^{|\lambda|}\prod_{i=1}^{\ell(\lambda)}\frac{\left(t^{N-i+1};q\right)_{\lambda_i}}{(t^{N-i}q;q)_{\lambda_i}}=\frac{(t^Nq;q)_\infty}{(q;q)_\infty}$$

$$\begin{aligned}I_N(t,u;0)&=\frac{1}{(t;t)_N}\sum_{\ell(\lambda)\le N}u^{|\lambda|}\prod_{i=1}^{\ell(\lambda)}\left(1-t^{N-i+1}\right)\\&=\frac{1}{(t;t)_N}\sum_{\ell(\lambda)\le N}u^{|\lambda|}(t^N;t^{-1})_{\ell(\lambda)}\end{aligned}$$

$$\sum_{\ell(\lambda)\le N}u^{|\lambda|}(t^N;t^{-1})_{\ell(\lambda)}=1+\sum_{\ell=1}^{N}(t^N;t^{-1})_\ell\sum_{\lambda_1\ge\cdots\ge\lambda_\ell\ge1}u^{\lambda_1+\cdots+\lambda_\ell}$$

$$\lambda_1=n_1+n_2+\cdots+n_\ell,\lambda_2=n_2+\cdots+n_\ell,\ldots,\lambda_\ell=n_\ell$$

$$\sum_{\lambda_1\geq\cdots\geq\lambda_\ell\geq 1} u^{\lambda_1+\cdots+\lambda_\ell} = \sum_{n_1=0}^{\infty}\sum_{n_2=0}^{\infty}\cdots\sum_{n_{\ell-1}=0}^{\infty}\sum_{n_\ell=1}^{\infty} u^{n_1+2n_2+\cdots+(\ell-1)n_{\ell-1}+\ell n_\ell}$$
$$= \sum_{n_1=0}^{\infty} u^{n_1}\sum_{n_2=0}^{\infty} u^{2n_2}\cdots\sum_{n_{\ell-1}=0}^{\infty} u^{(\ell-1)n_{\ell-1}}\sum_{n_\ell=1}^{\infty} u^{\ell n_\ell}$$
$$= \frac{u^\ell}{(1-u)(1-u^2)\cdots(1-u^\ell)}$$
$$= \frac{u^\ell}{(u;u)_\ell}$$

$$(t^N;t^{-1})_\ell = \frac{(t;t)_N}{(t;t)_{N-\ell}}$$

$$I_N(t,u;0) = \sum_{\ell=0}^{N}\frac{u^\ell}{(t;t)_{N-\ell}(u;u)_\ell}$$

$$I_N(t,0;q) = \frac{1}{N!}\frac{(q;q)_\infty^N}{(t;q)_\infty^N}\oint\cdots\oint_{\mathbb{T}^N}\prod_{i=1}^{N}\frac{dx_i}{2\pi i x_i}\prod_{1\leq i\neq j\leq N}\frac{(x_i/x_j;q)_\infty}{(tx_i/x_j;q)_\infty}$$

$$I_N(t,0;q) = \frac{(q;q)_\infty^N}{(t;q)_\infty^N}\mathcal{N}_{\emptyset,N} = \frac{(q;q)_\infty}{(t;t)_N(t^Nq;q)_\infty}$$

$$I_N(0,u;q) = (q;q)_\infty\sum_{\ell(\lambda)\leq N}\frac{u^{|\lambda|}}{(q;q)_{\lambda_N}}$$

$$\sum_{\ell(\lambda)\leq N}\frac{u^{|\lambda|}}{(q;q)_{\lambda_N}} = \frac{1}{(u;u)_N(u^Nq;q)_\infty}$$

$$\sum_{\ell(\lambda)\leq N}\frac{u^{|\lambda|}}{(q;q)_{\lambda_N}} = \sum_{n_1=0}^{\infty}\cdots\sum_{n_N=0}^{\infty}\frac{u^{n_1+2n_2+\cdots+Nn_N}}{(q;q)_{n_N}}$$
$$= \frac{1}{(1-u)(1-u^2)\cdots(1-u^{N-1})}\frac{1}{(u^N;q)_\infty}$$
$$= \frac{1}{(u;u)_{N-1}(u^N;q)_\infty} = \frac{1}{(u;u)_N(u^Nq;q)_\infty}$$

$$I_N(x,x;x^2) = \frac{(x^2;x^2)_\infty}{(x;x)_N(x^{N+2};x^2)_\infty}\sum_{\ell(\lambda)\leq N}x^{|\lambda|}\prod_{i=1}^{\ell(\lambda)}\frac{(x^{N-i+1};x^2)_{\lambda_i}}{(x^{N-i+2};x^2)_{\lambda_i}}$$

$$I_N(x,x;x^2) = \frac{(x^2;x^2)_\infty}{(x;x)_\infty^2}\sum_{n=0}^{\infty}(-1)^n\left[\binom{N+n}{N}+\binom{N+n-1}{N}\right]x^{nN+n^2}$$

$$\frac{(x;x)_\infty (x^N;x)_\infty}{(x^N;x^2)_\infty} \sum_{\ell(\lambda)\le N} x^{|\lambda|} \prod_{i=1}^{\ell(\lambda)} \frac{\left(x^{N-i+1};x^2\right)_{\lambda_i}}{\left(x^{N-i+2};x^2\right)_{\lambda_i}}$$
$$= \sum_{n=0}^{\infty} (-1)^n \left[\binom{N+n}{N} + \binom{N+n-1}{N}\right] x^{nN+n^2}$$

$$I_N(t,u;q) = \frac{(q;q)_\infty}{(t;t)_N (t^N q;q)_\infty} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \cdots \sum_{n_N=0}^{\infty} u^{n_1+2n_2+\cdots Nn_N}$$
$$\times \frac{(t^N;q)_{n_1+\cdots+n_N} (t^{N-1};q)_{n_2+\cdots+n_N} \cdots (t;q)_{n_N}}{(t^{N-1}q;q)_{n_1+\cdots+n_N} (t^{N-2}q;q)_{n_2+\cdots+n_N} \cdots (q;q)_{n_N}}$$

$$(x;q)_n = \frac{(x;q)_\infty}{(xq^n;q)_\infty}$$

$$\exp\left(\sum_{n=1}^{\infty} \frac{f_n}{n} p_n(x) p_n(x^{-1})\right) = \sum_{\mu} \frac{f_\mu}{z_\mu} p_\mu(x) p_\mu(x^{-1})$$

$$f_\mu = \prod_{i=1}^{\ell(\mu)} f_{\mu_i}, p_\mu(x) = \prod_{i=1}^{\ell(\mu)} p_{\mu_i}(x), z_\mu = \prod_{i\ge 1} i^{m_i} m_i!$$

$$p_\mu(x) = \sum_{\substack{\lambda \vdash n \\ \ell(\lambda)\le N}} \chi_\mu^\lambda s_\lambda(x)$$

$$I_N(t,u,v;p,q) = \sum_{\mu} \frac{f_\mu}{z_\mu} \sum_{\substack{\lambda \vdash n \\ \ell(\lambda)\le N}} \left(\chi_\mu^\lambda\right)^2.$$

$$I_{\lambda,\rho,N}(t,u;q) = \frac{1}{N!} \frac{(q;q)_\infty^N (tu;q)_\infty^N}{(t;q)_\infty^N (u;q)_\infty^N} \oiint_{\mathbb{T}^N} \prod_{i=1}^{N} \frac{dx_i}{2\pi i x_i}$$
$$\times \prod_{1\le i\ne j\le N} \frac{\left(x_i/x_j;q\right)_\infty \left(tux_i/x_j;q\right)_\infty}{\left(tx_i/x_j;q\right)_\infty \left(ux_i/x_j;q\right)_\infty} s_\lambda(x) s_\rho(x^{-1})$$

$$I_{(1^r),N}(t,u;q) = \frac{1}{N!} \frac{(q;q)_\infty^N}{(t;q)_\infty^N} \oiint_{\mathbb{T}^N} \prod_{i=1}^{N} \frac{dx_i}{2\pi i x_i} w(x)$$
$$\times \sum_{\ell(\lambda)\le N} u^{|\lambda|} b_\lambda P_\lambda(x;q,t) P_\lambda(x^{-1};q,t) e_r(x) e_r(x^{-1})$$

$$e_r(x) P_\lambda(x;q,t) = \sum_{\mu \in V_N^r(\lambda)} \psi'_{\mu/\lambda}(q,t) P_\mu(x;q,t)$$

$$\psi'_{\mu/\lambda}(q,t) = \psi_{\mu'/\lambda'}(t,q)$$

$$\psi_{\mu/\lambda}(q,t) = \prod_{1\le i\le j\le \ell(\lambda)} \frac{\left(t^{j-i+1}q^{\lambda_i-\lambda_j};q\right)_{\mu_i-\lambda_i}\left(t^{j-i}q^{\lambda_i-\mu_{j+1}+1};q\right)_{\mu_i-\lambda_i}}{\left(t^{j-i}q^{\lambda_i-\lambda_j+1};q\right)_{\mu_i-\lambda_i}\left(t^{j-i+1}q^{\lambda_i-\mu_{j+1}};q\right)_{\mu_i-\lambda_i}}$$

$$I_{(1^r),N}(t,u;q) = \frac{(q;q)_\infty^N}{(t;q)_\infty^N}\sum_{\ell(\lambda)\le N} u^{|\lambda|} b_\lambda \sum_{\mu\in V_N^r(\lambda)} \mathcal{N}_{\mu,N}\psi'_{\mu/\lambda}(q,t)^2$$

$$I_{(1^r),N}(t,u;q) = \frac{(q;q)}{(t;t)_N(t^Nq;q)_\infty}\sum_{\ell(\lambda)\le N} u^{|\lambda|}\sum_{\mu\in V_N^r(\lambda)} \varphi'_{\mu/\lambda}(q,t)\psi'_{\mu/\lambda}(q,t) \times\prod_{i=1}^{\ell(\mu)}\frac{\left(t^{N-i+1};q\right)_{\mu_i}}{(t^{N-i}q;q)_{\mu_i}}$$

$$\varphi'_{\mu/\lambda}(q,t) = \frac{b_\lambda}{b_\mu}\psi'_{\mu/\lambda}(q,t) = \varphi_{\mu'/\lambda'}(t,q)$$

$$\varphi_{\mu/\lambda}(q,t) = \prod_{1\le i\le j\le \ell(\mu)} \frac{\left(t^{j-i+1}q^{\mu_i-\mu_j};q\right)_{\mu_j-\lambda_j}\left(t^{j-i}q^{\lambda_i-\mu_{j+1}+1};q\right)_{\mu_{j+1}-\lambda_{j+1}}}{(t^{j-i}q^{\mu_i-\mu_j+1};q)_{\mu_j-\lambda_j}\left(t^{j-i+1}q^{\lambda_i-\mu_{j+1}};q\right)_{\mu_{j+1}-\lambda_{j+1}}}$$

$$s_\lambda(x) = \det\left(e_{\lambda'_i-i+j}(x)\right)_{1\le i,j\le l(\lambda')}$$

$$s_\lambda(x) = \begin{vmatrix} e_{r_1+r_2}(x) & e_{r_1+r_2+1}(x) \\ e_{r_2-1}(x) & e_{r_2}(x)\end{vmatrix} = e_{r_1+r_2}(x)e_{r_2}(x) - e_{r_1+r_2+1}(x)e_{r_2-1}(x)$$

$$\begin{aligned} &s_\lambda(x)s_\rho(x^{-1}) \\ &= e_{r_1+r_2}(x)e_{r_2}(x)e_{r'_1+r'_2}(x^{-1})e_{r'_2}(x^{-1}) - e_{r_1+r_2}(x)e_{r_2}(x)e_{r'_1+r'_2+1}(x^{-1})e_{r'_2-1}(x^{-1}) \\ &\quad - e_{r_1+r_2+1}(x)e_{r_2-1}(x)e_{r'_1+r'_2}(x^{-1})e_{r'_2}(x^{-1}) \\ &\quad + e_{r_1+r_2+1}(x)e_{r_2-1}(x)e_{r'_1+r'_2+1}(x^{-1})e_{r'_2-1}(x^{-1}) \end{aligned}$$

$$\langle A(x)\rangle'_N = \frac{1}{N!}\oint\cdots\oint_{\mathbb{T}^N}\prod_{i=1}^N \frac{dx_i}{2\pi i x_i} w(x)A(x)$$

$$\langle f,g\rangle'_N = \frac{1}{N!}\oint\cdots\oint_{\mathbb{T}^N}\prod_{i=1}^N \frac{dx_i}{2\pi i x_i} w(x)f(x)g(x^{-1}) = \langle f(x)g(x^{-1})\rangle'_N$$

$$I_N(t,u;q) = \frac{(q;q)_\infty^N}{(t,q)_\infty^N}\left\langle\prod_{i,j=1}^N\frac{\left(tux_i/x_j;q\right)_\infty}{\left(ux_i/x_j;q\right)_\infty}\right\rangle'_N$$

$$I_N(t,0;q) = \frac{(q;q)_\infty^N}{(t,q)_\infty^N}\langle 1\rangle'_N$$

$$\langle A(x)\rangle_\infty = \lim_{N\to\infty}\frac{\langle A(x)\rangle'_N}{\langle 1\rangle'_N}, \langle f,g\rangle_\infty = \lim_{N\to\infty}\frac{\langle f,g\rangle'_N}{\langle 1,1\rangle'_N}$$

$$\langle p_\lambda, p_\mu \rangle_\infty = \delta_{\lambda,\mu} z_\lambda(q,t)$$

$$z_\lambda(q,t) = z_\lambda \prod_{i=1}^{\ell(\lambda)} \frac{1-q^{\lambda_i}}{1-t^{\lambda_i}}$$

$$\frac{I_\infty(t,u;q)}{I_\infty(t,0;q)} = \lim_{N\to\infty} \frac{I_N(t,u;q)}{I_N(t,0;q)} = \left\langle \prod_{i,j=1}^{\infty} \frac{(tux_i/x_j;q)_\infty}{(ux_i/x_j;q)_\infty} \right\rangle_\infty$$

$$\prod_{i,j=1}^{\infty} \frac{(tux_i/x_j;q)_\infty}{(ux_i/x_j;q)_\infty} = \exp\left(\sum_{n=1}^{\infty} \frac{u^n}{n} \frac{1-t^n}{1-q^n} p_n(x) p_n(x^{-1}) \right)$$

$$= \sum_\lambda \frac{u^{|\lambda|}}{z_\lambda(q,t)} p_\lambda(x) p_\lambda(x^{-1})$$

$$\frac{I_\infty(t,u;q)}{I_\infty(t,0;q)} = \sum_\lambda \frac{u^{|\lambda|}}{z_\lambda(q,t)} \langle p_\lambda, p_\lambda \rangle_\infty = \sum_\lambda u^{|\lambda|} = \frac{1}{(u;u)_\infty}$$

$$I_\infty(t,0;q) = \frac{(q;q)_\infty}{(t;t)_\infty}$$

$$I_\infty(t,u;q) = \frac{(q;q)_\infty}{(t;t)_\infty (u;u)_\infty}$$

$$b_\lambda = \frac{1}{\langle P_\lambda, P_\lambda \rangle_\infty} = \lim_{N\to\infty} \frac{\mathcal{N}_{\emptyset,N}}{\mathcal{N}_{\lambda,N}}$$

$$\lim_{N\to\infty} \prod_{i=1}^{\ell(\lambda)} \frac{(t^{N-i+1};q)_{\lambda_i}}{(t^{N-i}q;q)_{\lambda_i}} = \lim_{N\to\infty} \frac{b_\lambda \mathcal{N}_{\lambda,N}}{b_\emptyset \mathcal{N}_{\emptyset,N}} = 1$$

$$I_\infty(t,u;q) = \frac{(q;q)_\infty}{(t;t)_\infty} \sum_\lambda u^{|\lambda|} = \frac{(q;q)_\infty}{(t;t)_\infty (u;u)_\infty}$$

$$I^{\text{p.s.}}_{\mu,\nu,N}(t,u;q) = \frac{1}{N!} \frac{(q;q)_\infty^N (tu;q)_\infty^N}{(t;q)_\infty^N (u;q)_\infty^N} \oint\cdots\oint_{\mathbb{T}^N} \prod_{i=1}^{N} \frac{dx_i}{2\pi i x_i}$$

$$\times \prod_{1\le i\ne j\le N} \frac{(x_i/x_j;q)_\infty (tux_i/x_j;q)_\infty}{(tx_i/x_j;q)_\infty (ux_i/x_j;q)_\infty} p_\mu(x) p_\nu(x^{-1})$$

$$\frac{I^{\text{p.s.}}_{\mu,\nu,\infty}(t,u;q)}{I_\infty(t,0;q)} = \left\langle \prod_{i,j=1}^{\infty} \frac{(tux_i/x_j;q)_\infty}{(ux_i/x_j;q)_\infty} p_\mu(x) p_\nu(x^{-1}) \right\rangle_\infty$$

$$\prod_{i,j=1}^{\infty}\frac{(tux_i/x_j;q)_\infty}{(ux_i/x_j;q)_\infty}p_\mu(x)p_\nu(x^{-1}) = \sum_\lambda \frac{u^{|\lambda|}}{z_\lambda(q,t)}p_\lambda(x)p_\lambda(x^{-1})p_\mu(x)p_\nu(x^{-1})$$

$$= \sum_\lambda \frac{u^{|\lambda|}}{z_\lambda(q,t)}p_{\lambda\cup\mu}(x)p_{\lambda\cup\nu}(x^{-1})$$

$$\frac{I^{\text{p.s.}}_{\mu,\nu,\infty}(t,u;q)}{I_\infty(t,0;q)} = \sum_\lambda \frac{u^{|\lambda|}}{z_\lambda(q,t)}\langle p_{\lambda\cup\mu}, p_{\lambda\cup\nu}\rangle_\infty$$

$$= \delta_{\mu,\nu}\sum_\lambda \frac{u^{|\lambda|}}{z_\lambda(q,t)}z_{\lambda\cup\mu}(q,t)$$

$$z_{\lambda\cup\mu}(q,t) = z_\lambda(q,t)z_\mu(q,t)\prod_{i\geq 1}\binom{k_i+m_i}{k_i}.$$

$$\frac{I^{\text{p.s.}}_{\mu,\nu,\infty}(t,u;q)}{I_\infty(t,0;q)} = \delta_{\mu,\nu}z_\mu(q,t)\sum_\lambda\prod_{i\geq 1}\binom{k_i+m_i}{k_i}u^{ik_i}$$

$$= \delta_{\mu,\nu}z_\mu(q,t)\prod_{i\geq 1}\frac{1}{(1-u^i)^{m_i+1}}$$

$$= \delta_{\mu,\nu}z_\mu(q,t)\frac{1}{(u;u)_\infty}\prod_{i=1}^{\ell(\mu)}\frac{1}{1-u^{\mu_i}}$$

$$= \delta_{\mu,\nu}\frac{z_\mu}{(u;u)_\infty}\prod_{i=1}^{\ell(\mu)}\frac{1-q^{\mu_i}}{(1-t^{\mu_i})(1-u^{\mu_i})}$$

$$I^{\text{p.s.}}_{\mu,\nu,\infty}(t,u;q) = \frac{(q;q)_\infty}{(t;t)_\infty(u;u)_\infty}\delta_{\mu,\nu}z_\mu\prod_{i=1}^{\ell(\mu)}\frac{1-q^{\mu_i}}{(1-t^{\mu_i})(1-u^{\mu_i})}$$

$$s_\lambda(x) = \sum_{\mu\vdash|\lambda|}\frac{\chi^\lambda_\mu}{z_\mu}p_\mu(x)$$

$$\frac{I_{\lambda,\rho,\infty}(t,u;q)}{I_\infty(t,u;q)} = \sum_{\mu\vdash|\lambda|}\frac{\chi^\lambda_\mu\chi^\rho_\mu}{z_\mu}\prod_{i=1}^{\ell(\mu)}\frac{1-q^{\mu_i}}{(1-t^{\mu_i})(1-u^{\mu_i})}.$$

$$\frac{I_{(1^r),\infty}(t,u;q)}{I_\infty(t,u;q)} = \frac{I_{(r),\infty}(t,u;q)}{I_\infty(t,u;q)} = \sum_{\mu\vdash r}\frac{1}{z_\mu}\prod_{i=1}^{\ell(\mu)}\frac{1-q^{\mu_i}}{(1-t^{\mu_i})(1-u^{\mu_i})}$$

$$\frac{I_{(1^r),\infty}(t,u;q)}{I_\infty(t,u;q)} = (u;u)_\infty\sum_\lambda u^{|\lambda|}\sum_{\mu\in V^r_\infty(\lambda)}\varphi'_{\mu/\lambda}(q,t)\psi'_{\mu/\lambda}(q,t)$$

$$\frac{I_N(t,u,v;p,q)}{I_\infty(t,u,v;p,q)} = \sum_{k=0}^{\infty}t^{kN}\hat{I}_k(t,u,v;p,q)$$

$$(1-f)(1-\hat{f}) = (1-t)(1-t^{-1})$$

$$\hat{f}(t,u,v;p,q)=1-\frac{(1-t^{-1})(1-p)(1-q)}{(1-u)(1-v)}$$

$$\hat{I}_k(t,u,v;p,q)=I_k(t^{-1},q,p;v,u)$$

$$\frac{I_N(t,u;q)}{I_\infty(t,u;q)}=\sum_{k=0}^{\infty}t^{kN}\hat{I}_k(t,u;q),\hat{I}_k(t,u;q)=I_k(t^{-1},q;u)$$

$$I_1(t,u;\alpha u)=\frac{1}{1-t}+(1-\alpha)u+(1-\alpha)(1+\alpha+\alpha t)u^2+O(u^3)$$

$$I_2(t,u;\alpha u)=\frac{1}{(1-t)(1-t^2)}+\frac{1-\alpha}{1-t}u+\frac{(1-\alpha)(2+\alpha-t+\alpha t^2)}{1-t}u^2+O(u^3)$$

$$I_3(t,u;\alpha u)=\frac{1}{(1-t)(1-t^2)(1-t^3)}+\frac{1-\alpha}{(1-t)(1-t^2)}u+\frac{(1-\alpha)(2+\alpha-t^2+\alpha t^3)}{(1-t)(1-t^2)}u^2+O(u^3)$$

$$I_N(t,u;\alpha u)=\sum_{j=0}^{\infty}g_{N,j}^{(\alpha)}(t)u^j,g_{N,j}^{(\alpha)}(t)=\frac{1-\alpha}{(t;t)_{N-1}}G_{N,j}^{(\alpha)}(t)$$

$$G_{N,0}^{(\alpha)}(t)=\frac{1}{(1-\alpha)(1-t^N)}$$

$$G_{N,1}^{(\alpha)}(t)=1$$

$$G_{N,2}^{(\alpha)}(t)=2+\alpha+t^N\left(-\frac{1}{t}+\alpha\right)$$

$$I_\infty(t,u;\alpha u)=\frac{(\alpha u;\alpha u)_\infty}{(t;t)_\infty(u;u)_\infty}=\sum_{j=0}^{\infty}g_{\infty,j}^{(\alpha)}(t)u^j,g_{\infty,j}^{(\alpha)}(t)=\frac{1-\alpha}{(t;t)_\infty}G_{\infty,j}^{(\alpha)}$$

$$I_k(t^{-1},\alpha u;u)=I_k(t^{-1},\alpha u;\alpha^{-1}\cdot\alpha u)=\sum_{j=0}^{\infty}g_{k,j}^{(\alpha^{-1})}(t^{-1})\alpha^j u^j.$$

$$g_{N,j}^{(\alpha)}(t)=\sum_{k=0}^{\infty}t^{kN}\sum_{m=0}^{j}\alpha^m g_{\infty,j-m}^{(\alpha)}(t)g_{k,m}^{(\alpha^{-1})}(t^{-1})\ (j\geq 1)$$

$$G_{N,j}^{(\alpha)}(t)=G_{\infty,j}^{(\alpha)}+\sum_{n=1}^{j-1}t^{nN}\left[\frac{G_{\infty,j}^{(\alpha)}}{(t;t)_n}+\sum_{k=1}^{n}\sum_{m=0}^{j}\frac{(-1)^{k-1}t^{\frac{k(k-1)}{2}}\alpha^m(1-\alpha^{-1})}{(t;t)_{n-k}(t;t)_{k-1}}G_{\infty,j-m}^{(\alpha)}G_{k,m}^{(\alpha^{-1})}(t^{-1})\right]\ (j\geq 1)$$

$$G_{N,2}^{(\alpha)}(t)=2+\alpha+t^N\left(1+2\alpha-\alpha G_{1,2}^{(\alpha^{-1})}(t^{-1})\right)$$

$$G_{N,3}^{(\alpha)}(t) = 3+\alpha+t^N\left(-\frac{1}{t^2}+\frac{-2+\alpha}{t}+\alpha+\alpha^2\right)+t^{2N}\left(\frac{1}{t^3}-\frac{\alpha}{t^2}-\frac{\alpha}{t}+\alpha^2\right)$$

$$\begin{aligned}G_{N,4}^{(\alpha)}(t) =& 5+2\alpha+t^N\left(-\frac{1}{t^3}+\frac{-2+\alpha}{t^2}+\frac{-4+\alpha+\alpha^2}{t}+\alpha+2\alpha^2+\alpha^3\right)\\ &+t^{2N}\left(\frac{1}{t^5}+\frac{1-\alpha}{t^4}+\frac{2-2\alpha}{t^3}+\frac{-3\alpha+\alpha^2}{t^2}-\frac{\alpha}{t}+\alpha^2+\alpha^3\right)\\ &+t^{3N}\left(-\frac{1}{t^6}+\frac{\alpha}{t^5}+\frac{\alpha}{t^4}+\frac{\alpha-\alpha^2}{t^3}-\frac{\alpha^2}{t^2}-\frac{\alpha^2}{t}+\alpha^3\right)\end{aligned}$$

$$I_{(1),1}(t,u;\alpha u) = \frac{1}{1-t}+(1-\alpha)u+(1-\alpha)(1+\alpha+\alpha t)u^2+O(u^3)$$

$$\begin{aligned}I_{(1),2}(t,u;\alpha u) =& \frac{1}{(1-t)^2}+\frac{1-\alpha}{1-t}(2+t)u\\ &+\frac{1-\alpha}{1-t}(3+2\alpha t+(-1+2\alpha)t^2+\alpha t^3)u^2+O(u^3)\end{aligned}$$

$$\begin{aligned}I_{(1),3}(t,u;\alpha u) =& \frac{1}{(1-t)^2(1-t^2)}+\frac{1-\alpha}{(1-t)(1-t^2)}(2+2t+t^2)u\\ &+\frac{1-\alpha}{(1-t)(1-t^2)}(4+3t+(-1+2\alpha)t^2+(-2+3\alpha)t^3\\ &+(-1+2\alpha)t^4+\alpha t^5)u^2+O(u^3)\end{aligned}$$

$$I_{(1),N}(t,u;\alpha u) = \frac{1-\alpha}{(1-t)(t;t)_{N-1}}\sum_{j=0}^{\infty}u^j G_{(1),N,j}^{(\alpha)}(t)$$

$$G_{(1),N,0}^{(\alpha)}(t) = \frac{1}{1-\alpha}$$

$$\frac{I_{(1),N}(t,u;q)}{I_{(1),\infty}(t,u;q)} = \frac{I_N(t,u;q)}{I_\infty(t,u;q)}-\frac{(1-t^{-1})(1-q)}{1-u}\sum_{k=1}^{\infty}t^{kN}I_{(1),k}(t^{-1},q;u)$$

$$I_{(1),\infty}(t,u;q) = \frac{1-q}{(1-t)(1-u)}I_\infty(t,u;q)$$

$$G_{(1),N,1}^{(\alpha)}(t) = 2-t^N\left(\frac{1}{t}+1\right)$$

$$\begin{aligned}G_{(1),N,2}^{(\alpha)}(t) =& 4+t^N\left(-\frac{1}{t^2}+\frac{2(-2+\alpha)}{t}-2+\alpha\right)\\ &+t^{2N}\left(\frac{1}{t^3}+\frac{1-\alpha}{t^2}+\frac{1-\alpha}{t}-\alpha\right)\end{aligned}$$

$$\begin{aligned}G_{(1),N,3}^{(\alpha)}(t) =& 7-\alpha-\alpha^2+t^N\left(-\frac{1}{t^3}+\frac{-4+3\alpha}{t^2}+\frac{-9+7\alpha}{t}-3+\alpha+\alpha^2\right)\\ &+t^{2N}\left(\frac{1}{t^5}+\frac{2-\alpha}{t^4}+\frac{5(1-\alpha)}{t^3}+\frac{4-7\alpha+2\alpha^2}{t^2}+\frac{2-4\alpha+\alpha^2}{t}-\alpha+\alpha^2\right)\\ &+t^{3N}\left(-\frac{1}{t^6}+\frac{-1+\alpha}{t^5}+\frac{-1+2\alpha}{t^4}-\frac{(1-\alpha)^2}{t^3}+\frac{(2-\alpha)\alpha}{t^2}+\frac{(1-\alpha)\alpha}{t}-\alpha^2\right)\end{aligned}$$

$$I^{\text{'t Hooft}}_{(1^r,0^{N-r})}=\frac{1}{r!\,(N-r)!}\frac{(q;q)_\infty^{2N}}{(t;q)_\infty^{N}(u;q)_\infty^{N}}\oiint_{\mathbb{T}^r}\prod_{i=1}^{r}\frac{dx_i}{2\pi i x_i}\oiint_{\mathbb{T}^{N-r}}\prod_{j=1}^{N-r}\frac{dy_j}{2\pi i y_j}$$
$$\times\prod_{1\le i\ne j\le r}\frac{(x_i/x_j;q)_\infty(qx_i/x_j;q)_\infty}{(tx_i/x_j;q)_\infty(ux_i/x_j;q)_\infty}\prod_{1\le i\ne j\le N-r}\frac{(y_i/y_j;q)_\infty(qy_i/y_j;q)_\infty}{(ty_i/y_j;q)_\infty(uy_i/y_j;q)_\infty}$$
$$\times\prod_{i=1}^{r}\prod_{j=1}^{N-r}\frac{(q^{1/2}x_i/y_j;q)_\infty(q^{1/2}y_j/x_i;q)_\infty(q^{3/2}x_i/y_j;q)_\infty\left(q^{3/2}y_j/x_i;q\right)_\infty}{(tq^{1/2}x_i/y_j;q)_\infty(tq^{1/2}y_j/x_i;q)_\infty(uq^{1/2}x_i/y_j;q)_\infty(uq^{1/2}y_j/x_i;q)_\infty}$$

$$\sum_{r=0}^{\infty}z^r e_r(x_1,\dots,x_n)=\prod_{i=1}^{n}(1+zx_i)$$
$$\sum_{r=0}^{\infty}z^r h_r(x_1,\dots,x_n)=\prod_{i=1}^{n}\frac{1}{1-zx_i}$$

$$p_r(x_1,\dots,x_n)=\sum_{i=1}^{n}x_i^r$$

$$m_\lambda(x_1,\dots,x_n)=\sum_{\alpha\in S_n\cdot\lambda}x^\alpha$$

$$m_{(3)}(x_1,x_2,x_3)=x_1^3+x_2^3+x_3^3$$
$$m_{(2,1)}(x_1,x_2,x_3)=x_1^2x_2+x_1x_2^2+x_1^2x_3+x_1x_3^2+x_2^2x_3+x_2x_3^2$$
$$m_{(1^3)}(x_1,x_2,x_3)=x_1x_2x_3$$

$$D_x=\sum_{i=1}^{n}\prod_{1\le j(\ne i)\le n}\frac{tx_i-x_j}{x_i-x_j}T_{q,x_i}$$

$$T_{q,x_i}f(x_1,\dots,x_i,\dots,x_n)=f(x_1,\dots,qx_i,\dots,x_n)$$

$$P_\lambda(x;q,t)=m_\lambda(x)+\sum_{\mu<\lambda}u_\mu^\lambda(q,t)m_\mu(x)$$

$$D_xP_\lambda(x;q,t)=d_\lambda(q,t)P_\lambda(x;q,t)$$

$$d_\lambda(q,t)=\sum_{i=1}^{n}t^{n-i}q^{\lambda_i}$$

$$P_{(1)}=m_{(1)},P_{(2)}=m_{(2)}+\frac{(1-q^2)(1-t)}{(1-q)(1-qt)}m_{(1^2)},P_{(1^2)}=m_{(1^2)}$$

$$P_{(3)}=m_{(3)}+\frac{(1-q^3)(1-t)}{(1-q)(1-q^2t)}m_{(2,1)}+\frac{(1-q^2)(1-q^3)(1-t)^2}{(1-q)^2(1-qt)(1-q^2t)}m_{(1^3)}$$
$$P_{(2,1)}=m_{(2,1)}+\frac{(1-t)(2+q+t+2qt)}{1-qt^2}m_{(1^3)}$$
$$P_{(1^3)}=m_{(1^3)}$$

$$P_{(1^r)}(x;q,t) = m_{(1^r)}(x) = e_r(x)$$

$$\prod_{i=1}^{n} \frac{(tx_i y;q)_\infty}{(x_i y;q)_\infty} = \sum_{r=0}^{\infty} g_r(x;q,t) y^r$$

$$P_{(r)}(x;q,t) = \frac{(q;q)_r}{(t;q)_r} g_r(x;q,t)$$

$$s_\lambda(x) = \frac{\det\left(x_i^{\lambda_j+n-j}\right)_{1\le i,j\le n}}{\det\left(x_i^{n-j}\right)_{1\le i,j\le n}}$$

$$w(x) = \prod_{1\le i\ne j\le n} \frac{\left(x_i/x_j;q\right)_\infty}{\left(tx_i/x_j;q\right)_\infty}$$

$$\frac{1}{n!} \oiint_{\mathbb{T}^n} \prod_{i=1}^{n} \frac{dx_i}{2\pi i x_i} w(x) P_\lambda(x;q,t) P_\mu(x^{-1};q,t) = \delta_{\lambda,\mu} \mathcal{N}_{\lambda,n}$$

$$\mathbb{T}^n = \{(x_1,\dots,x_n) \in \mathbb{C}^n || x_i| = 1\}$$

$$\mathcal{N}_{\lambda,n} = \prod_{1\le i<j\le n} \frac{\left(t^{j-i}q^{\lambda_i-\lambda_j+1};q\right)_\infty \left(t^{j-i}q^{\lambda_i-\lambda_j};q\right)_\infty}{\left(t^{j-i+1}q^{\lambda_i-\lambda_j};q\right)_\infty \left(t^{j-i-1}q^{\lambda_i-\lambda_j+1};q\right)_\infty}$$

$$\Pi(x,y;q,t) = \prod_{i=1}^{n} \prod_{j=1}^{m} \frac{\left(tx_i y_j;q\right)_\infty}{\left(x_i y_j;q\right)_\infty}$$

$$\Pi(x,y;q,t) = \sum_{\ell(\lambda)\le \min(n,m)} b_\lambda P_\lambda(x;q,t) P_\lambda(y;q,t)$$

$$b_\lambda = \prod_{1\le i\le j\le \ell(\lambda)} \frac{\left(t^{j-i+1}q^{\lambda_i-\lambda_j};q\right)_{\lambda_j-\lambda_{j+1}}}{\left(t^{j-i}q^{\lambda_i-\lambda_j+1};q\right)_{\lambda_j-\lambda_{j+1}}}$$

$$\prod_{i=1}^{n} \prod_{j=1}^{m} \frac{1}{1-x_i y_j} = \sum_{\ell(\lambda)\le \min(n,m)} s_\lambda(x) s_\lambda(y)$$

$$D_x \Pi(x,y;q,t) = D_y \Pi(x,y;q,t)$$

$$P_\lambda(x;q,t) P_\rho(x;q,t) = \sum_{\mu} c_{\lambda\rho}^{\mu}(q,t) P_\mu(x;q,t)$$

$$e_r(x) P_\lambda(x;q,t) = \sum_{\mu\in V_n^r(\lambda)} \psi'_{\mu/\lambda}(q,t) P_\mu(x;q,t)$$

$$V_n^r(\lambda) = \{\mu \vdash |\lambda| + r \mid \ell(\mu) \le n \text{ and } \mu/\lambda \text{ is a vertical bridge dS }\}$$

$$\psi_{\mu/\lambda}(q,t) = \prod_{1\le i\le j\le \ell(\lambda)} \frac{\left(t^{j-i+1}q^{\lambda_i-\lambda_j};q\right)_{\mu_i-\lambda_i}\left(t^{j-i}q^{\lambda_i-\lambda_{j+1}+1};q\right)_{\mu_i-\lambda_i}}{\left(t^{j-i}q^{\lambda_i-\lambda_j+1};q\right)_{\mu_i-\lambda_i}\left(t^{j-i+1}q^{\lambda_i-\mu_{j+1}};q\right)_{\mu_i-\lambda_i}}$$

$$\psi'_{\mu/\lambda}(q,t) = \psi_{\mu'/\lambda'}(t,q)$$

$$g_r(x;q,t)P_\lambda(x;q,t) = \sum_{\mu\in H_n^r(\lambda)} \varphi_{\mu/\lambda}(q,t)P_\mu(x;q,t)$$

$$H_n^r(\lambda) = \{\mu \vdash |\lambda| + r \mid \ell(\mu) \le n \text{ and } \mu/\lambda \text{ is a horizontal bridge AdS }\},$$

$$\varphi_{\mu/\lambda}(q,t) = \prod_{1\le i\le j\le \ell(\mu)} \frac{\left(t^{j-i+1}q^{\mu_i-\mu_j};q\right)_{\mu_j-\lambda_j}\left(t^{j-i}q^{\lambda_i-\mu_{j+1}+1};q\right)_{\mu_{j+1}-\lambda_{j+1}}}{\left(t^{j-i}q^{\mu_i-\mu_j+1};q\right)_{\mu_j-\lambda_j}\left(t^{j-i+1}q^{\lambda_i-\mu_{j+1}};q\right)_{\mu_{j+1}-\lambda_{j+1}}}.$$

$$\mathbb{IV}_{\mathcal{N}}^{U(N)|U(M)} = \frac{1}{N!\,M!}\frac{(q;q)_\infty^{N+M}}{(t;q)_\infty^{N+M}} \oiint_{\mathbb{T}^N} \prod_{i=1}^N \frac{dx_i}{2\pi i x_i} \prod_{1\le i\ne j\le N} \frac{\left(x_i/x_j;q\right)_\infty}{\left(tx_i/x_j;q\right)_\infty}$$

$$\times \oiint_{\mathbb{T}^M} \prod_{i=1}^M \frac{dy_i}{2\pi i y_i} \prod_{1\le i\ne j\le M} \frac{\left(y_i/y_j;q\right)_\infty}{\left(ty_i/y_j;q\right)_\infty} \prod_{i=1}^N \prod_{j=1}^M \frac{\left(tu^{1/2}x_i/y_j;q\right)_\infty\left(tu^{1/2}y_j/x_i;q\right)_\infty}{\left(u^{1/2}x_i/y_j;q\right)_\infty\left(u^{1/2}y_j/x_i;q\right)_\infty}$$

$$\prod_{i=1}^N \prod_{j=1}^M \frac{\left(tu^{1/2}x_i/y_j;q\right)_\infty}{\left(u^{1/2}x_i/y_j;q\right)_\infty} = \sum_{\ell(\lambda)\le N} u^{|\lambda|/2} b_\lambda P_\lambda(x;q,t)P_\lambda(y^{-1};q,t)$$

$$\prod_{i=1}^N \prod_{j=1}^M \frac{\left(tu^{1/2}y_j/x_i;q\right)_\infty}{\left(u^{1/2}y_j/x_i;q\right)_\infty} = \sum_{\ell(\mu)\le N} u^{|\mu|/2} b_\mu P_\mu(x^{-1};q,t)P_\mu(y;q,t)$$

$$\mathbb{IV}_{\mathcal{N}}^{U(N)|U(M)} = \frac{(q;q)_\infty^{N+M}}{(t;q)_\infty^{N+M}} \sum_{\ell(\lambda)\le N} \sum_{\ell(\mu)\le N} u^{|\lambda|/2+|\mu|/2} b_\lambda b_\mu \delta_{\lambda,\mu}\mathcal{N}_{\lambda,N}\delta_{\mu,\lambda}\mathcal{N}_{\mu,M}$$

$$= \frac{(q;q)_\infty^{N+M}}{(t;q)_\infty^{N+M}} \sum_{\ell(\lambda)\le N} u^{|\lambda|} b_\lambda \mathcal{N}_{\lambda,N} b_\lambda \mathcal{N}_{\lambda,M}$$

$$\mathbb{IV}_{\mathcal{N}}^{U(N)|U(M)} = \frac{(q;q)_\infty^2}{(t;t)_N(t^Nq;q)_\infty(t;t)_M(t^Mq;q)_\infty} \sum_{\ell(\lambda)\le N} u^{|\lambda|} \prod_{i=1}^{\ell(\lambda)} \frac{\left(t^{N-i+1};q\right)_{\lambda_i}\left(t^{M-i+1};q\right)_{\lambda_i}}{(t^{N-i}q;q)_{\lambda_i}(t^{M-i}q;q)_{\lambda_i}}.$$

$$\mathbb{IV}_{\mathcal{D}'}^{U(N)|U(M)} = \frac{1}{N!}\frac{(q;q)_\infty^N(tu;q)_\infty^N}{(t;q)_\infty^N(u;q)_\infty^N} \oiint_{\mathbb{T}^N} \prod_{i=1}^N \frac{dx_i}{2\pi i x_i} \prod_{1\le i\ne j\le N} \frac{\left(x_i/x_j;q\right)_\infty\left(tux_i/x_j;q\right)_\infty}{\left(tx_i/x_j;q\right)_\infty\left(ux_i/x_j;q\right)_\infty}$$

$$\times \prod_{k=1}^{M-N} \frac{\left(t^{k-1}q;q\right)_\infty}{\left(t^k;q\right)_\infty} \prod_{i=1}^N \frac{\left(t^{(M-N+1)/2}ux_i;q\right)_\infty\left(t^{(M-N+1)/2}ux_i^{-1};q\right)_\infty}{\left(t^{(M-N+1)/2}x_i;q\right)_\infty\left(t^{(M-N+1)/2}x_i^{-1};q\right)_\infty}$$

$$\mathbb{IV}_{\mathcal{D}'}^{U(N)|U(M)} = \frac{(q;q)_\infty^N}{(u;q)_\infty^N}\prod_{k=1}^{M-N}\frac{\left(t^{k-1}q;q\right)_\infty}{\left(t^k;q\right)_\infty}\frac{1}{N!}\oiint_{\mathbb{T}^N}\prod_{i=1}^{N}\frac{dx_i}{2\pi i x_i}\prod_{1\le i\ne j\le N}\frac{\left(x_i/x_j;q\right)_\infty}{\left(ux_i/x_j;q\right)_\infty}$$
$$\times\prod_{i,j=1}^{N}\frac{\left(tux_i/x_j;q\right)_\infty}{\left(tx_i/x_j;q\right)_\infty}\prod_{i=1}^{N}\frac{\left(t^{(M-N+1)/2}ux_i;q\right)_\infty\left(t^{(M-N+1)/2}ux_i^{-1};q\right)_\infty}{\left(t^{(M-N+1)/2}x_i;q\right)_\infty\left(t^{(M-N+1)/2}x_i^{-1};q\right)_\infty}$$

$$\prod_{i,j=1}^{N}\frac{\left(tux_i/x_j;q\right)_\infty}{\left(tx_i/x_j;q\right)_\infty}=\sum_{\ell(\lambda)\le N}t^{|\lambda|}b_\lambda P_\lambda(x;q,u)P_\lambda(x^{-1};q,u),$$

$$\prod_{i=1}^{N}\frac{\left(t^{(M-N+1)/2}ux_i;q\right)_\infty}{\left(t^{(M-N+1)/2}x_i;q\right)_\infty}=\sum_{r=0}^{\infty}g_r(x;q,u)t^{(M-N+1)r/2}$$

$$\mathbb{IV}_{\mathcal{D}'}^{U(N)|U(M)} = \frac{(q;q)_\infty^2}{(t;t)_{M-N}(t^{M-N}q,q)_\infty(u;u)_N(u^Nq;q)_\infty}\sum_{\ell(\lambda)\le N}\sum_{r=0}^{\infty}t^{|\lambda|+(M-N+1)r}$$
$$\times\sum_{\mu\in H_N^r(\lambda)}\varphi_{\mu/\lambda}(q,u)\psi_{\mu/\lambda}(q,u)\prod_{i=1}^{\ell(\mu)}\frac{\left(u^{N-i+1};q\right)_{\mu_i}}{\left(u^{N-i}q;q\right)_{\mu_i}},$$

$$\prod_{k=1}^{M-N}\frac{\left(t^{k-1}q;q\right)_\infty}{\left(t^k;q\right)_\infty}=\frac{(q;q)_\infty}{(t;t)_{M-N}(t^{M-N}q;q)_\infty}$$

$$\mathbb{IV}_{\mathcal{N}}^{U(N)|U(M)} = \mathbb{IV}_{\mathcal{D}'}^{U(N)|U(M)}\ (q=tu)$$

$$k\partial_\Lambda S_\Lambda[\phi]=\frac{1}{2}\int_0^\infty\frac{AdS}{s}r(s,\Lambda^2Z_\Lambda)\mathrm{Tr}\left[e^{-sS_\Lambda^{(2)}}\right]$$

$$r(s,\Lambda^2Z_\Lambda)=\left(2+\epsilon\frac{\Lambda\partial_\Lambda Z_\Lambda}{Z_\Lambda}\right)\frac{(sm\Lambda^2Z_\Lambda)^m}{\Gamma(m)}e^{-sm\Lambda^2Z_\Lambda}$$

$$k\partial_\Lambda S_\Lambda[\phi]=\mathrm{Tr}\left[\left(1+\frac{\epsilon}{2}\frac{\Lambda\partial_\Lambda Z_\Lambda}{Z_\Lambda}\right)e^{-\frac{S_\Lambda^{(2)}}{\Lambda^2Z_\Lambda}}\right]$$

$$S[g,\phi_1,\dots,\phi_N]=\int d^dx\sqrt{g}\left(\frac{1}{16\pi G}(-R+2\lambda)+\frac{1}{2}\sum_{i=1}^{N}\phi_i(-\Box)\phi^i+B(\rho)R+V(\rho)\right)$$

$$\rho=\frac{1}{2}\sum_{a=1}^{N}\phi^a\phi^a$$

$$F(\rho)=\frac{1}{16\pi G}-B(\rho),U(\rho)=V(\rho)+\frac{\lambda}{8\pi G}$$

$$S[g,\phi_1,\dots,\phi_N]=\int d^dx\sqrt{g}\left(-F(\rho)R+\frac{1}{2}\sum_{i=1}^{N}\phi_i(-\Box)\phi^i+U(\rho)\right)$$

$$S_\Lambda[g,\phi_1,\dots,\phi_N]=\int d^dx\sqrt{g}\left(-F_\Lambda(\rho)R+\frac{1}{2}\sum_{i=1}^{N}\phi_i(-\Box)\phi^i+U_\Lambda(\rho)\right)$$

$$\Lambda\partial_\Lambda S_\Lambda=\int d^dx\sqrt{g}(-\Lambda\partial_\Lambda F_\Lambda(\rho)R+\Lambda\partial_\Lambda U_\Lambda(\rho))$$

$x=\alpha^{1/2}\Lambda^{2-d}\rho, u_\Lambda(x)=\alpha\Lambda^{-d}U_\Lambda(\rho)$, and $f_\Lambda(x)=\alpha\Lambda^{2-d}F_\Lambda(\rho)$ where $\alpha=\frac{m^{d/2}\Gamma\left(m-\frac{d}{2}\right)}{4(4\pi)^{d/2}\Gamma(m)}$

$$\dot{u}=-du+(d-2)xu'+2(d-3)d+4(N-1)\left(1+\frac{u'}{m}\right)^{\frac{d}{2}-m}+\epsilon(d-1)d\left((d-2)\left(1-x\frac{f'}{f}\right)+\frac{\dot{f}}{f}\right)$$
$$+4\left(1+\frac{\epsilon\left(\frac{1}{2}(d-2)\left(\frac{xf'}{f}-\frac{2xf''}{f'}-1\right)+\frac{\dot{f}'}{f'}-\frac{\dot{f}}{2f}\right)}{1+\frac{(d-2)f}{4(d-1)x(f')^2}}\right)\left(1+\frac{2xu''+u'}{m\left(1+\frac{4(d-1)x(f')^2}{(d-2)f}\right)}\right)^{\frac{d}{2}-m}$$

$$\dot{f}=(2-d)(f-xf')+\frac{1}{3}\left(\frac{d}{2m}-1\right)(d^2-3d+36)+4\left(\frac{d}{2m}-1\right)(N-1)\left(\frac{1}{6}+f'\right)\left(1+\frac{u'}{m}\right)^{\frac{d}{2}-m-1}$$
$$+\epsilon\frac{1}{6}\left(\frac{d}{2m}-1\right)\left(d^2-d-\frac{24}{d}-24\right)\left((d-2)\left(1-x\frac{f'}{f}\right)+\frac{\dot{f}}{f}\right)+$$
$$+\frac{2}{3}\left(\frac{d}{2m}-1\right)\left(1+\frac{6\left(\frac{4x(f')^2}{(d-2)f}+2xf''+f'\right)}{1+\frac{4(d-1)x(f')^2}{(d-2)f}}\right)\left(1+\frac{\epsilon\left(\frac{1}{2}(d-2)\left(\frac{xf'}{f}-\frac{2xf''}{f'}-1\right)+\frac{f'}{f'}-\frac{\dot{f}}{2f}\right)}{1+\frac{(d-2)f}{4(d-1)x(f')^2}}\right)\left(1+\frac{2xu''+u'}{m\left(1+\frac{4(d-1)x(f')^2}{(d-2)f}\right)}\right)^{\frac{d}{2}-m-1}$$

$$u_*=2d+\frac{4N}{d}-6+(d^2-3d+2)\epsilon$$
$$f_*=\frac{(d-2m)(d^2-3d-36+2N)}{6(d-2)m}+\frac{(d^3-d^2-24d-24)(d-2m)\epsilon}{12dm}$$

$$N>N_c=\frac{1}{2}(-d^2+3d+36)+\frac{1}{4}\left(-d^3+3d^2+22d-\frac{48}{d}-24\right)\epsilon$$

$$u_*(x)=\frac{4N}{d}+2d-6, f_*(x)=\frac{(d-2m)(d^3-4d^2+d(2N-33)+10N+36)}{6(d-2)(d-1)m}+\frac{x}{d-1}$$

$$N_c=-\frac{d^3-4d^2-33d+36}{2(d+5)}$$

$$u_*=\frac{4N}{3}, f_*(x)=\frac{60-7N\pm\sqrt{N^2+72N+2736}}{48(N-1)}x$$

$$u_*=2+N, f_*(x)=\frac{7-N\pm\sqrt{3}\sqrt{N+11}}{6(N-1)}x$$

$$u_\Lambda(x)=u_*(x)+\delta u(x)\left(\frac{\Lambda_0}{\Lambda}\right)^\theta, f_\Lambda(x)=f_*(x)+\delta f(x)\left(\frac{\Lambda_0}{\Lambda}\right)^\theta$$

$$\left(8-\frac{4d}{m}\right)\delta u''+\left(N\left(4-\frac{2d}{m}\right)\frac{1}{x}+2-d\right)\delta u'+(d-\theta)\frac{\delta u}{x}+$$
$$+\frac{12(d-2)(d-1)d^2\lambda m\epsilon}{(d-2m)(d^4\epsilon+d^3(2-3\epsilon)-2d^2(11\epsilon+3)+4d(n+6(\epsilon-3))+48\epsilon)}\left((d-2)\delta f'-\lambda\frac{\delta f}{x}\right)=0$$
$$\left(8-\frac{4d}{m}\right)\delta f''+\left(N\left(4-\frac{2d}{m}\right)\frac{1}{x}-\frac{2(d-2)d(d^2-3d+2N-36)}{d^4\epsilon+d^3(2-3\epsilon)-2d^2(11\epsilon+3)+4d(N+6(\epsilon-3))+48\epsilon}\right)\delta f'+$$
$$+\left(d-\theta-2+\frac{(d-2)(d^3-d^2-24d-24)\theta\epsilon}{d^4\epsilon+d^3(2-3\epsilon)-2d^2(11\epsilon+3)+4d(N+6(\epsilon-3))+48\epsilon}\right)\frac{\delta f}{x}-$$
$$-\frac{(d-2m)(d-2(m+1))\delta u''}{3m^2}-\frac{N(d^2-2d(2m+1)+4m(m+1))}{6m^2}\frac{\delta u'}{x}=0$$

$$\delta u(x)=\sum_{i=0}^{\infty}u_i x^i,\delta f(x)=\sum_{i=0}^{\infty}f_i x^i$$

$$\theta=-2+d-(d-2)j+\frac{(d-2)^2(d^3-d^2-24d-24)\epsilon}{2d(d^2-3d-36+2N)}$$
$$\theta=d-(d-2)j$$

$\theta=2+\frac{18\epsilon}{16-N}$:
$\delta u=\frac{108\epsilon}{N+9\epsilon-16},\delta f=1$
$\theta=\frac{18\epsilon}{16-N}$:
$$\delta u=\frac{108\epsilon(N^2-N(9\epsilon+32)+27\epsilon^2+144\epsilon+256)}{(N-9\epsilon-16)(N-3\epsilon-16)(2N+9\epsilon-32)}-\frac{162(N-16)\epsilon}{N(N-9\epsilon-16)(N-3\epsilon-16)}x,$$
$$\delta f=1-\frac{3(N-16)(N+9\epsilon-16)}{2N(N-9\epsilon-16)(N-3\epsilon-16)}x$$
$\theta=-2+\frac{18\epsilon}{16-N}$:
$$\delta u=-\frac{324\epsilon^2(-N^3+3N^2(9\epsilon+16)-3N(15\epsilon^2+288\epsilon+256)+81\epsilon^3+720\epsilon^2+6912\epsilon+4096)}{(N-9\epsilon-16)(N+3\epsilon-16)(-2N^3+3N^2(9\epsilon+32)-6N(9\epsilon^2+144\epsilon+256)+81\epsilon^3+864\epsilon^2+6912\epsilon+8192)}+$$
$$+\frac{324(N-16)x\epsilon(N^2-N(9\epsilon+32)+27\epsilon^2+144\epsilon+256)}{N(N-9\epsilon-16)(-2N^3+3N^2(9\epsilon+32)-6N(9\epsilon^2+144\epsilon+256)+81\epsilon^3+864\epsilon^2+6912\epsilon+8192)}+$$
$$-\frac{243(N-16)^2x^2\epsilon(2N+9\epsilon-32)}{N(N+2)(N-9\epsilon-16)(-2N^3+3N^2(9\epsilon+32)-6N(9\epsilon^2+144\epsilon+256)+81\epsilon^3+864\epsilon^2+6912\epsilon+8192)},$$

$$\delta f=1+\frac{3(N-16)x(2N^2+N(3\epsilon-64)-27\epsilon^2-48\epsilon+512)}{N(-2N^3+3N^2(9\epsilon+32)-6N(9\epsilon^2+144\epsilon+256)+81\epsilon^3+864\epsilon^2+6912\epsilon+8192)}$$
$$-\frac{9(N-16)^2x^2(2N^2+N(27\epsilon-64)+81\epsilon^2-432\epsilon+512)}{4N(N+2)(N-9\epsilon-16)(-2N^3+3N^2(9\epsilon+32)-6N(9\epsilon^2+144\epsilon+256)+81\epsilon^3+864\epsilon^2+6912\epsilon+8192)}$$

$\theta=4,\delta u=1,\delta f=0$
$\theta=2,\delta u=1-\frac{9x}{14N},\delta f=\frac{N-9\epsilon-16}{189\epsilon}$
$\theta=0,\delta u=1-\frac{3x}{N}+\frac{27x^2}{28N(N+2)},\delta f=\frac{2(N-30\epsilon-16)}{189\epsilon}+\frac{-N+9\epsilon+16}{63N\epsilon}x$

$$(d-\theta)\delta u + \frac{-4N(d^3(x-4)+d^2(11x+16)+4d(5x+33)+4(x-36))-(d+2)x(2d^4-3d^3(x+4)-2d^2(3x+25)+12d(x+17)+24(x-10))+32(d+5)N^2}{(d+2)(2d^3-d^2(3x+8)+2d(2N-6x-33)+4(5N-3x+18))}\delta u' + \frac{16x(2d^3-d^2(3x+8)+d(4N-66)+20N+12(x+6))}{(d+2)(2d^3-d^2(3x+8)+2d(2N-6x-33)+4(5N-3x+18))}\delta u'' = 0$$

$$(d-\theta-2)\delta f + \frac{(-4N(d^3(x-4)+d^2(11x+16)+4d(5x+33)+4(x-36))-(d+2)x(2d^4-3d^3(x+4)-2d^2(3x+25)+12d(x+17)+24(x-10))+32(d+5)N^2)}{(d+2)(2d^3-d^2(3x+8)+2d(2N-6x-33)+4(5N-3x+18))}\delta f' + \frac{16x(2d^3-d^2(3x+8)+d(4N-66)+20N+12(x+6))}{(d+2)(2d^3-d^2(3x+8)+2d(2N-6x-33)+4(5N-3x+18))}\delta f'' - \frac{16(d+5)(N(2d^3-d^2(3x+8)-6d(2x+11)-12(x-6))+4(d+5)N^2+12(d+2)x)}{3(d-1)(d+2)^2(2d^3-d^2(3x+8)+2d(2N-6x-33)+4(5N-3x+18))}\delta u' - \frac{32(d+5)x(2d^3-d^2(3x+8)+d(4N-66)+20N+12(x+6))}{3(d-1)(d+2)^2(2d^3-d^2(3x+8)+2d(2N-6x-33)+4(5N-3x+18))}\delta u'' = 0$$

$$\delta u(x) = \sum_{i=0}^{\infty} u_i (x-x_0)^i$$

$$\delta u(x\to\infty) = x^{\frac{d-\theta}{d-2}}\left(c_1 + \sum_{n=1}^{\infty} v_{-n}x^{-n}\right)$$

$$\delta f(x\to\infty) = x^{1-\frac{\theta}{d-2}}\left(c_2 + f_{4-d}x^{4-d} + \sum_{n=1}^{\infty} f_{-n}x^{-n} + \log(x)\sum_{n=0}^{\infty} g_{-n}x^{-n}\right)$$

$$v_{-1} = \frac{1}{9}c_1(\theta-4)(3N-\theta)$$

$$v_{-2} = \frac{1}{162}c_1(\theta-4)(\theta^3-36\theta+9(\theta-2)N^2-6(\theta-1)^2N+96)$$

$$f_{-1} = -\frac{2}{243}c_1(\theta-4)(2(\theta-8)\theta+9N^2-3(2\theta+3)N+48) - \frac{1}{9}c_2(\theta-2)(\theta-3N+2)$$

$$f_{-2} = \frac{1}{162}c_2(\theta-2)(\theta(\theta(\theta+6)-24)+9\theta N^2-6(\theta+1)^2N+32) +$$

$$+\frac{c_1}{2187}(\theta-4)(4\theta(\theta((\theta-5)\theta-18)+24)-54(\theta-1)N^3+9(6\theta^2+3\theta-4)N^2-3(\theta(4\theta(2\theta-5)+3)+160)N+768)$$

$$g_0 = \frac{1}{27}c_1(\theta-4)(3N-\theta), g_{-1} = \frac{1}{243}c_1(\theta-4)(\theta-2)(-\theta+3N-2)(3N-\theta)$$

$$g_{-2} = \frac{c_1}{4374}(\theta-4)(\theta-2)(3N-\theta)(\theta(\theta(\theta+6)-24)+9\theta N^2-6(\theta+1)^2N+32)$$

$$\dot{u} = -du_* + (d-2)xu_*' + \left(1+\frac{u_*'}{m}\right)^{\frac{d}{2}-m}$$

$$\dot{f} = (2-d)f_* + (d-2)xf_*' + \left(\frac{d}{2m}-1\right)\left(\frac{1}{6}+f_*'\right)\left(1+\frac{u_*'}{m}\right)^{\frac{d}{2}-m-1}.$$

$$x = c_1(u_*')^{\frac{d}{2}-1} - \frac{1}{d-2}\left(\frac{d}{2m}-1\right)\,{}_2F_1\left(1-\frac{d}{2},-\frac{d}{2}+m+1;2-\frac{d}{2};-\frac{u_*'}{m}\right)$$

$$f_*(u_*') = -\frac{x}{6} + \left(\frac{c_1}{6}+c_2\right)(u_*')^{\frac{d}{2}-1}$$

$${}_2F_1\left(1-\frac{d}{2},-\frac{d}{2}+m+1;2-\frac{d}{2};-\frac{v_0}{m}\right) = 0$$

$$x = -\frac{1}{d-2}\left(\frac{d}{2m}-1\right)\,{}_2F_1\left(1-\frac{d}{2},-\frac{d}{2}+m+1;2-\frac{d}{2};-\frac{u_*'}{m}\right)$$

$$f_* = -\frac{x}{6}$$

$$(d-\theta)\delta u + \delta u'\left(\left(1-\frac{d}{2m}\right)\left(1+\frac{u_*'}{m}\right)^{\frac{d}{2}-m-1} - (d-2)x\right) = 0$$

$$(d-\theta-2)\delta f + \delta f'\left(\left(1-\frac{d}{2m}\right)\left(1+\frac{u_*'}{m}\right)^{\frac{d}{2}-m-1} - (d-2)x\right) = 0$$

$$\delta u(u_*') = b_1(u_*')^{\frac{d-\theta}{2}}, \delta f(u_*') = b_2(u_*')^{\frac{d-\theta-2}{2}}$$

$$\delta u_*^{PS}(p,x) = \left|\frac{u_*(p+1,x)-u_*(p,x)}{u_*(p,x)}\right|$$

$$\delta u_*^{SPS}(p,x) = \left|\frac{u_*^{\text{shoot}}(x)-u_*^{PS}(p,x)}{u_*^{\text{shoot}}(x)}\right|$$

$$f_*(x) \sim \frac{1}{16\pi g_*^{PG}} + f_\infty x$$

$$\delta\theta^{PS} = \left|\frac{\theta^{PS}(p+1)-\theta^{PS}(p)}{\theta^{PS}(p)}\right|$$

$$g_{\mu\nu} = \bar{g}_{\mu\rho}\left(e^h\right)^\rho_\nu = \bar{g}_{\mu\rho}\left(\delta^\rho_\nu + h^\rho_\nu + \frac{1}{2}h^\rho_\sigma h^\sigma_\nu + \cdots\right) = \bar{g}_{\mu\nu} + h_{\mu\nu} + \frac{1}{2}h_{\mu\lambda}h^\lambda_\nu + \cdots$$

$$\phi_i = \bar{\phi}_i + \Phi_i$$

$$\delta^2\mathcal{L} = \delta^2\left[\sqrt{g}\left(-F(\rho)R + \frac{1}{2}\sum_{i=1}^{N}\phi_i(-\Box)\phi^i + U(\rho)\right)\right]$$

$$S^{(4)}_{hh} = \frac{1}{2}\int d^dx\sqrt{\bar{g}}h_{\mu\nu}\{-F(\bar{\rho})\Lambda^{\mu\nu}_{\rho\sigma}\,\Box + \bar{A}^{\mu\nu}_{\rho\sigma} + F(\bar{\rho})\left[\delta^\nu_\rho\bar{\nabla}^\mu\bar{\nabla}_\sigma - g^{\mu\nu}\bar{\nabla}_\rho\bar{\nabla}_\sigma\right]\}h^{\rho\sigma}$$

$$S^{(4)}_{\Phi\Phi} = \frac{1}{2}\sum_{a,b=1}^{N}\int d^dx\sqrt{\bar{g}}\Phi^a\{[-\Box + 2\bar{\rho}U''(\bar{\rho}) + U'(\bar{\rho}) - [2\bar{\rho}F''(\bar{\rho}) + F'(\bar{\rho})]\bar{R}]P^R_{ab} + [-\Box + U'(\bar{\rho}) - F'(\bar{\rho})\bar{R}]P^T_{ab}\}\Phi^b$$

$$S^{(4)}_{h\Phi} = \sum_{a,b=1}^{N}\int d^dx\sqrt{\bar{g}}\Phi^a P^R_{ab}\bar{\phi}_b\sqrt{2\rho}\left\{\frac{1}{2}\bar{g}^{\mu\nu}[U'(\bar{\rho}) - F'(\bar{\rho})\bar{R}] - F'(\bar{\rho})(-\bar{R}^{\mu\nu} + \bar{\nabla}^\mu\bar{\nabla}^\nu - \bar{g}^{\mu\nu}\,\bar{\Box})\right\}h_{\mu\nu}$$

$$K^{\mu\nu}_{\rho\sigma} = \frac{1}{4}\left(\delta^{\mu}_{\rho}\delta^{\nu}_{\sigma} + \delta^{\mu}_{\sigma}\delta^{\nu}_{\rho}\right) - \frac{1}{2}\bar{g}^{\mu\nu}\bar{g}_{\rho\sigma}$$

$$A^{\mu\nu}_{\rho\sigma} = \frac{1}{4}\left(\delta^{\mu}_{\sigma}\delta^{\nu}_{\rho} - \bar{g}^{\mu\nu}\bar{g}_{\rho\sigma}\right)(F(\bar{\rho})\bar{R} - U(\bar{\rho})) +$$
$$+F(\bar{\rho})\left[-\frac{1}{4}\left(\bar{R}^{\nu}_{\sigma}\delta^{\mu}_{\rho} + \bar{R}^{\mu}_{\sigma}\delta^{\nu}_{\rho} + \bar{R}^{\nu}_{\rho}\delta^{\mu}_{\sigma}\right) + \bar{g}^{\mu\nu}\bar{R}_{\rho\sigma} - \frac{1}{2}\left(\bar{R}^{\mu\nu}_{\rho\sigma} + \bar{R}^{\mu\nu}_{\sigma\rho}\right)\right]$$

$$P^{R}_{ab} = \frac{\bar{\phi}_a\bar{\phi}_b}{\bar{\phi}^2}, P^{T}_{ab} = \delta^{ab} - P^{R}_{ab} = \delta^{ab} - \frac{\bar{\phi}_a\bar{\phi}_b}{\bar{\phi}^2}$$

$$\bar{R}_{\mu\nu\lambda\rho} = \frac{\bar{R}}{d(d-1)}\left(g_{\mu\lambda}\bar{g}_{\nu\rho} - \bar{g}_{\nu\lambda}\bar{g}_{\mu\rho}\right), \bar{R}_{\mu\nu} = \frac{1}{d}\bar{R}\bar{g}_{\mu\nu}$$

$$P^{R}_{ab} = \frac{\bar{\phi}_a\bar{\phi}_b}{\bar{\phi}^2} = \begin{cases} 0\ a \neq N, b \neq N \\ \frac{\bar{\phi}^2_N}{\bar{\phi}^2} = 1\ a = N, b = N' \end{cases} P^{T}_{ab} = \delta^{ab} - P^{R}_{ab} = \begin{cases} 0 & a \neq N, b \neq N \\ 0 & a = N, b = N \\ 1 & a = b \neq N \end{cases}$$

$$A^{\mu\nu}_{\rho\sigma} = \frac{1}{4}\left(2\delta^{\mu}_{\rho}\delta^{\nu}_{\sigma} - \bar{g}^{\mu\nu}\bar{g}_{\rho\sigma}\right)[F(\bar{\rho})R - U(\bar{\rho})] + F(\bar{\rho})\frac{d-2}{d(d-1)}\left(\bar{g}^{\mu\nu}\bar{g}_{\rho\sigma} - \delta^{\mu}_{\rho}\delta^{\nu}_{\sigma}\right)\bar{R}$$

$$S^{(4)}_{\Phi\Phi} + S^{(4)}_{h\Phi} = \frac{1}{2}\sum_{a=1}^{N-1}\int d^dx\sqrt{\bar{g}}\Phi^a[-\bar{\Box} + U'(\bar{\rho}) - F'(\bar{\rho})\bar{R}]\Phi^a$$
$$+\frac{1}{2}\int d^dx\sqrt{\bar{g}}\Phi^N[-\Box + 2\bar{\rho}U''(\bar{\rho}) + U'(\bar{\rho}) - [2\bar{\rho}F''(\bar{\rho}) + F'(\bar{\rho})]\bar{R}]\Phi^N$$
$$+\int d^dx\sqrt{\bar{g}}\Phi^N\sqrt{2\bar{\rho}}\left\{\frac{1}{2}\bar{g}^{\mu\nu}[U'(\bar{\rho}) - F'(\bar{\rho})\bar{R}] - F'(\bar{\rho})(-\bar{R}^{\mu\nu} + \bar{\nabla}^{\mu}\bar{\nabla}^{\nu} - \bar{g}^{\mu\nu}\bar{\Box})\right\}h_{\mu\nu}$$

$$h_{\mu\nu} = h^{T}_{\mu\nu} + \bar{\nabla}_{\mu}\epsilon_{\nu} + \bar{\nabla}_{\nu}\epsilon_{\mu} + \bar{\nabla}_{\mu}\bar{\nabla}_{\nu}\sigma - \frac{1}{d}\bar{g}_{\mu\nu}\bar{\Box}\,\sigma + \frac{1}{d}\bar{g}_{\mu\nu}h$$

$$S^{(4)} = \frac{1}{2}\int d^dx\sqrt{\bar{g}}\left\{\frac{1}{2}F(\bar{\rho})h^{T}_{\mu\nu}\left[-\bar{\Box} + \frac{2}{d(d-1)}\bar{R}\right]h^{\mu\nu T} - \frac{(d-1)(d-2)}{2d^2}F(\bar{\rho})\hat{\sigma}(-\bar{\Box})\hat{\sigma}\right.$$
$$-\frac{(d-1)(d-2)}{2d^2}F(\bar{\rho})h\left[-\bar{\Box} + \frac{d-2}{2(d-1)}\bar{R}\right]h + \frac{U(\bar{\rho})}{4}h^2$$
$$\left.-\frac{(d-1)(d-2)}{d}F(\bar{\rho})h\sqrt{-\bar{\Box}}\sqrt{-\bar{\Box} - \frac{\bar{R}}{d-1}}\hat{\sigma}\right\} + \frac{1}{2}\sum_{a=1}^{N-1}\int d^dx\sqrt{\bar{g}}\Phi^a[-\bar{\Box} + U'(\bar{\rho}) - F'(\bar{\rho})\bar{R}]\Phi^a$$
$$+\frac{1}{2}\int d^dx\sqrt{\bar{g}}\Phi^N[-\bar{\Box} + 2\bar{\rho}U''(\bar{\rho}) + U'(\bar{\rho}) - [2\bar{\rho}F''(\bar{\rho}) + F'(\bar{\rho})]\bar{R}]\Phi^N$$
$$+\int d^dx\sqrt{\bar{g}}\sqrt{2\bar{\rho}}\left\{-F'(\bar{\rho})\left(1 - \frac{1}{d}\right)\Phi^N\sqrt{-\bar{\Box}}\sqrt{-\bar{\Box} - \frac{1}{d-1}\bar{R}}\hat{\sigma}\right.$$
$$\left.+\Phi^N\left\{-F'(\bar{\rho})\frac{d-1}{d}\left[\bar{\Box} + \frac{d-2}{2(d-1)}\bar{R}\right] + \frac{U'(\bar{\rho})}{2}\right\}h\right\}$$

$$\hat{\epsilon}_{\nu} = \sqrt{-\bar{\Box} - \frac{\bar{R}}{d}}\epsilon_{\nu}, \hat{\sigma} = \sqrt{-\bar{\Box}}\sqrt{-\bar{\Box} - \frac{1}{d-1}\bar{R}}\sigma$$

$$\int d^d x\sqrt{\bar{g}}\left[c_\mu\left(-\bar{\Box}-\frac{\bar{R}}{d}\right)c^\mu + c(-\bar{\Box})c\right]$$

$$S^{(4)} = \int d^d x\sqrt{\bar{g}}\left(F(\bar{\rho})\left\{\frac{1}{4}h^T_{\mu\nu}\left[-\bar{\Box}+\frac{2}{d(d-1)}\bar{R}\right]h^{\mu\nu T} - \frac{(d-1)(d-2)}{4d^2}\hat{\sigma}(-\bar{\Box})\hat{\sigma}\right\}\right.$$
$$\left.+c_\mu\left(-\bar{\Box}-\frac{\bar{R}}{d}\right)c^\mu + c(-\bar{\Box})c\right) + \frac{1}{2}\sum_{a=1}^{N-1}\int d^d x\sqrt{\bar{g}}\Phi^a[-\Box + U'(\bar{\rho}) - F'(\bar{\rho})\bar{R}]\Phi^a$$
$$\frac{1}{2}\int d^d x\sqrt{\bar{g}}\Phi^N[-\Box + 2\bar{\rho}U''(\bar{\rho}) + U'(\bar{\rho}) - [2\bar{\rho}F''(\bar{\rho}) + F'(\bar{\rho})]\bar{R}]\Phi^N$$
$$+\int d^d x\sqrt{\bar{g}}\left\{-\sqrt{2\rho}F'(\bar{\rho})\left(1-\frac{1}{d}\right)\Phi^N\sqrt{-\Box}\sqrt{-\Box-\frac{1}{d-1}\bar{R}}\hat{\sigma}\right\}.$$

$$\sigma' = \hat{\sigma} + \frac{2d}{d-2}\frac{\sqrt{2\rho}F'(\bar{\rho})}{F(\bar{\rho})}\sqrt{\frac{-\Box-\frac{1}{d-1}\bar{R}}{-\Box}}\Phi^N$$

$$S^{(4)} = \int d^d x\sqrt{\bar{g}}\left(F(\bar{\rho})\left\{\frac{1}{4}h^T_{\mu\nu}\left[-\bar{\Box}+\frac{2}{d(d-1)}\bar{R}\right]h^{\mu\nu T} - \frac{(d-1)(d-2)}{4d^2}\sigma'(-\bar{\Box})\sigma'\right\}+\right.$$
$$\left.+c_\mu\left(-\bar{\Box}-\frac{\bar{R}}{d}\right)c^\mu + c(-\bar{\Box})c\right) + \frac{1}{2}\sum_{a=1}^{N-1}\int d^d x\sqrt{\bar{g}}\Phi^a\left[-\Box+\frac{1}{\bar{\rho}}[V'(\bar{\rho})-F'(\bar{\rho})\bar{R}]\right]\Phi^a +$$
$$+\int d^d x\sqrt{\bar{g}}\Phi^N\frac{1}{2}\left(1+4\bar{\rho}\frac{d-1}{d-2}\frac{[F'(\bar{\rho})]^2}{F(\bar{\rho})}\right)\left[-\Box-\frac{2\bar{\rho}F''(\bar{\rho})+F'(\bar{\rho})+\frac{4\bar{\rho}}{d-2}\frac{[F'(\bar{\rho})]^2}{F(\bar{\rho})}}{1+4\bar{\rho}\frac{d-1}{d-2}\frac{[F'(\bar{\rho})]^2}{F(\bar{\rho})}}+\frac{2\bar{\rho}V''(\bar{\rho})+V'(\bar{\rho})}{1+4\bar{\rho}\frac{d-1}{d-2}\frac{[F'(\bar{\rho})]^2}{F(\bar{\rho})}}\right]\Phi^N$$
$$=\int d^d x\sqrt{\bar{g}}\left(h^T_{\mu\nu}S^{(2)}_{h^T h^T}h^{\mu\nu T} + \sigma' S^{(2)}_{\sigma\sigma}\sigma' + c_\mu S^{(2)}_{c^\mu c^\mu}c^\mu + cS^{(2)}_{cc}c + \sum_{a=1}^{N-1}\Phi^a S^{(2)}_{\Phi^T\Phi^T}\Phi^a + \Phi^N S^{(2)}_{\Phi^L\Phi^L}\Phi^N\right)$$

$$C_\sigma = \tfrac{(d-1)(d-2)}{4d^2} \text{ and } F_\Lambda C_\Phi = F_\Lambda\frac{1}{2}\left(1+4\bar{\rho}\tfrac{d-1}{d-2}\tfrac{[F'_\Lambda(\bar{\rho})]^2}{F_\Lambda(\bar{\rho})}\right)$$

$$\Lambda\partial_\Lambda S_\Lambda = -\frac{1}{2}\int_0^\infty\frac{dS}{s}r\left(s,\Lambda^2\frac{F_\Lambda(\bar{\rho})}{4}\right)\mathrm{Tr}\left[e^{-sS^{(2)}_{h^T h^T}}\right] - \frac{1}{2}\int_0^\infty\frac{AdS}{s}r(s,\Lambda^2F_\Lambda(\bar{\rho})C_\sigma)\mathrm{Tr}\left[e^{-sS^{(2)}_{\sigma\sigma}}\right]$$
$$+\int_0^\infty\frac{ds}{s}r(s,\Lambda^2)\mathrm{Tr}\left[e^{-sS^{(2)}_c\mu^\mu}\right] + \int_0^\infty\frac{ds}{s}r(s,\Lambda^2)\mathrm{Tr}\left[e^{-sS^{(2)}_{cc}}\right]$$
$$-\frac{N-1}{2}\int_0^\infty\frac{dS}{s}r(s,\Lambda^2)\mathrm{Tr}\left[e^{-sS^{(2)}_{\Phi^T\Phi^T}}\right] - \frac{1}{2}\int_0^\infty\frac{AdS}{s}r(s,\Lambda^2C_\Phi)\mathrm{Tr}\left[e^{-sS^{(2)}_{\Phi^L\Phi^L}}\right]$$

$$\int\frac{dS}{s}r(s,Z_\Lambda\Lambda^2)\mathrm{Tr}\left[e^{-s(Az+B)}\right]$$

$$\mathrm{Tr}_{(s)}[W(z)] = \frac{1}{(4\pi)^{\frac{d}{2}}}\int d^d x\sqrt{g}\sum_{n=0}^{+\infty}B^{(s)}_{2n}(z)Q_{\frac{d}{2}-n}(W)$$

$$Q_{\frac{d}{2}-n} = \int_0^\infty\frac{1}{\Gamma\left(n-\frac{d}{2}\right)}W(z)z^{\left(n-\frac{d}{2}\right)-1}dz$$

$$\mathrm{Tr}_{(s)}[W(z)] = \frac{1}{(4\pi)^{\frac{d}{2}}}\int d^d x\sqrt{g}\left[b_0^{(s)}Q_{\frac{d}{2}}(W) + b_2^{(s)}RQ_{\frac{d}{2}-1}(W)\right] + O(R^2)$$

$$b_0^{(0)} = 1, b_2^{(0)} = \frac{1}{6},$$
$$b_0^{(1)} = d-1, b_2^{(1)} = \frac{1}{6}(d-1) - \frac{1}{d},$$
$$b_0^{(2)} = \frac{(d+1)(d-2)}{2}, b_2^{(2)} = \frac{1}{6}\frac{(d-5)(d+1)(d+2)}{2(d-1)}.$$

$$I(s,A,B) \equiv \int d^d x\sqrt{g}\left[-\frac{\Lambda^d m^{\frac{d}{2}}}{2^{d+1}\pi^{\frac{d}{2}}\left(\frac{B}{A\Lambda^2 m}+1\right)^{m-\frac{d}{2}}}\frac{\Gamma\left(m-\frac{d}{2}\right)}{\Gamma(m)}\left(2A + \Lambda\frac{\partial A}{\partial\Lambda}\right)\right.$$
$$\left.\times\left(2b_0^{(s)}(z)(A\Lambda^2 m + B) - b_2^{(s)}(z)A(d-2m)R\right)\right]$$

$$\Lambda\partial_\Lambda S_\Lambda = -\frac{1}{2}I\left(2,\frac{F_\Lambda}{4},\frac{RF_\Lambda}{2(d-1)d}\right) - \frac{1}{2}I\left(0,-\frac{(d-2)(d-1)F_\Lambda}{4d^2},0\right) + \frac{1}{2}I\left(1,\frac{1}{2},-\frac{1}{2}\frac{R}{d}\right) +$$
$$\frac{1}{2}I\left(0,\frac{1}{2},0\right) - \frac{1}{2}I\left(0,\frac{1}{2}\left(1+\frac{4(d-1)\rho F_\Lambda^2}{(d-2)F_\Lambda}\right),\frac{1}{2}\left(-\left(2\rho F_\Lambda'' + F_\Lambda' + \frac{4\rho F_\Lambda^2}{(d-2)F_\Lambda}\right)R + U_\Lambda' + 2\rho U_\Lambda''\right)\right) -$$
$$\frac{N-1}{2}I\left(0,\frac{1}{2},\frac{1}{2}(-RF_\Lambda' + U_\Lambda')\right).$$

$$\Lambda\partial_\Lambda U_\Lambda = \frac{\Lambda^d m^{\frac{d}{2}}\Gamma\left(m-\frac{d}{2}\right)}{4(4\pi)^{\frac{d}{2}}\Gamma(m)}\left(2(d-3)d + 4(N-1)\left(1+\frac{U_\Lambda'}{\Lambda^2 m}\right)^{\frac{d}{2}-m} + (d-1)d\epsilon\frac{\Lambda\partial_\Lambda F_\Lambda}{F_\Lambda}\right.$$
$$\left.+4\left(1+\frac{\epsilon\left(2\frac{\Lambda\partial_\Lambda F_\Lambda'}{F_\Lambda'} - \frac{\Lambda\partial_\Lambda F_\Lambda}{F_\Lambda}\right)}{2\left(1+\frac{(d-2)F_\Lambda}{4(d-1)\bar\rho F_\Lambda'^2}\right)}\right)\left(1+\frac{\frac{U_\Lambda'}{\Lambda^2}+\frac{2\bar\rho U_\Lambda''}{\Lambda^2}}{m+\frac{4(d-1)m\bar\rho F_\Lambda'^2}{(d-2)F_\Lambda}}\right)^{\frac{d}{2}-m}\right)$$

$$\Lambda\partial_\Lambda F_\Lambda = \frac{\Lambda^{d-2}m^{\frac{d}{2}-1}\left(\frac{d}{2}-m\right)\Gamma\left(m-\frac{d}{2}\right)}{24(4\pi)^{\frac{d}{2}}\Gamma(m)}\left(2(d-3)d + 72 + 4(N-1)(1+6F_\Lambda')\left(1+\frac{U_\Lambda'}{\Lambda^2 m}\right)^{\frac{d}{2}-m-1}\right.$$
$$+\left(d^4 - d - \frac{24}{d} - 24\right)\epsilon\frac{\Lambda\partial_\Lambda F_\Lambda}{F_\Lambda}$$
$$\left.+4\left(1+\frac{\epsilon\left(2\frac{\Lambda\partial_\Lambda F_\Lambda'}{F_\Lambda'} - \frac{\Lambda\partial_\Lambda F_\Lambda}{F_\Lambda}\right)}{2\left(1+\frac{(d-2)F_\Lambda}{4(d-1)\bar\rho F_\Lambda'^2}\right)}\right)\left(1+\frac{6\left(2\bar\rho F_\Lambda'' + \frac{4\bar\rho F_\Lambda'^2}{(d-2)F_\Lambda(\bar\rho)} + F_\Lambda'\right)}{1+\frac{4(d-1)\bar\rho F_\Lambda'^2}{(d-2)F_\Lambda}}\right)\left(1+\frac{\frac{U_\Lambda'}{\Lambda^2}+\frac{2\rho U_\Lambda''}{\Lambda^2}}{m+\frac{4(d-1)m\bar\rho F_\Lambda'^2}{(d-2)F_\Lambda}}\right)^{\frac{d}{2}-(m+1)}\right)$$

$$u_*(x\to 0) = u_0 - \frac{5x(4N - 3u_0 + 6\epsilon)}{12\epsilon - 6u_0} + O(x^2)$$
$$f_*(x\to 0) = f_0 + \frac{x(-60f_0N + 4(26N\epsilon + 72N - 9\epsilon^2) + 36u_0\epsilon - 9u_0^2)}{54(u_0 - 2\epsilon)^2} + O(x^2)$$

$$u_*(x\to\infty)=x^3u_\infty+\frac{36\epsilon}{5xf_\infty}+\frac{\frac{2(8f_\infty+5N-4)}{15u_\infty}+\frac{96\epsilon(13\epsilon-36)}{125f_\infty^2}}{x^2}+\frac{32\left(\frac{\epsilon(13\epsilon-72)(13\epsilon-18)}{f_\infty^3}-\frac{25(\epsilon+2)}{u_\infty}\right)}{375x^3}+$$
$$+\frac{-\frac{625(16f_\infty(4f_\infty+1)+25N-24)}{u_\infty^2}-\frac{28800(\epsilon+1)(13\epsilon-24)}{f_\infty u_\infty}+\frac{384\epsilon(13\epsilon-108)(13\epsilon-36)(13\epsilon-12)}{f_\infty^4}+O\left(\frac{1}{x^5}\right)}{39375x^4}+$$
$$f_*(x\to\infty)=xf_\infty+\frac{24}{5}+\frac{104\epsilon}{25xf_\infty}+\frac{416(13\epsilon-36)\epsilon}{1125x^2f_\infty^2}+\frac{208(13\epsilon-72)(13\epsilon-18)\epsilon}{5625x^3f_\infty^3}$$
$$+\frac{\frac{1664\epsilon(13\epsilon-108)(13\epsilon-36)(13\epsilon-12)}{421875f_\infty^4}-\frac{2f_\infty(152f_\infty+75N-52)+25N-24}{675u_\infty^2}}{x^4}+O\left(\frac{1}{x^5}\right)$$

$$v_{IN}(u_0,f_0)=v_{OUT}(u_\infty,f_\infty,x_{\max}),\quad f_{IN}(u_0,f_0)=f_{OUT}(u_\infty,f_\infty,x_{\max}),$$
$$v'_{IN}(u_0,f_0)=v'_{OUT}(u_\infty,f_\infty,x_{\max}),\quad f'_{IN}(u_0,f_0)=f'_{OUT}(u_\infty,f_\infty,x_{\max}),$$
$$f''_{IN}(u_0,f_0)=f''_{OUT}(u_\infty,f_\infty,x_{\max}).$$

$u_*(x)=\frac{\lambda_*}{8\pi g_*}+u_*^{WF}(x)$ and $f_*(x)=\frac{1}{16\pi g_*}-b_*(x)$ in the limits $x\to 0$ and $x\to\infty$

$$u_*(x\to 0)=\frac{\lambda_*^{PG}}{8\pi g_*^{PG}}+u_*^{WF}(x\to 0),u_*(x\to\infty)=\frac{\lambda_*^{PG}}{8\pi g_*^{PG}}+u_*^{WF}(x\to+\infty),$$
$$f_*(x\to 0)=\frac{1}{16\pi g_*^{PG}}+b_*(x\to 0),f_*(x\to+\infty)=\frac{1}{16\pi g_*^{PG}}-b_*(x\to\infty).$$

$$u_0=\frac{5}{6v_0+15},u_\infty=\frac{(d-2)^{\frac{d}{d-2}}m^{\frac{d}{d-2}}\left(\frac{\Gamma\left(m-\frac{d}{2}\right)}{\Gamma\left(2-\frac{d}{2}\right)\Gamma(m)}\right)^{\frac{2}{d-2}}}{d}$$
$$f_0=-\frac{1}{6},f_\infty=-\frac{1}{6}$$

$$v_*(x)=v_B(x)H(x_0-x)+v_U(x)H(x-x_0),$$
$$f_*(x)=f_B(x)H(x_0-x)+f_U(x)H(x-x_0),$$

$$v_B(x)=\sum_{i=0}^{p}c_i^{(v)}T_i\left(\frac{2x}{x_0}-1\right),v_U(x)=x^{\frac{d}{d-2}-1}\sum_{i=0}^{p}r_i^{(v)}T_i\left(\frac{x-x_0-L}{x-x_0+L}\right),$$
$$f_B(x)=\sum_{i=0}^{p}c_i^{(f)}T_i\left(\frac{2x}{x_0}-1\right),f_U(x)=x\sum_{i=0}^{p}r_i^{(f)}T_i\left(\frac{x-x_0-L}{x-x_0+L}\right).$$

$$\beta_v\left(x_i^B,v_B(x_i^B),f_B(x_i^B)\right)=0,\quad \beta_f\left(x_i^B,v_B(x_i^B),f_B(x_i^B)\right)=0,\quad i=1,\dots,p+1$$
$$\beta_v\left(x_j^U,v_U(x_i^U),f_U(x_j^U)\right)=0,\quad \beta_f\left(x_j^U,v_U(x_j^U),f_U(x_j^U)\right)=0,\quad j=1,\dots,p-1$$
$$v_B(x_0)=v_U(x_0),\quad v'_B(x_0)=v'_U(x_0),\quad v''_B(x_0)=v''_U(x_0)$$

$$-\cos\left(\frac{i\pi}{p}\right)=\frac{2x_i^B}{x_0}-1,-\cos\left(\frac{j\pi}{p}\right)=\frac{x_j^U-x_0-L}{x_j^U-x_0+L}$$

$$0 = (3-\theta)\delta u +$$
$$+\left(\frac{2(2m-3)(N-1)\left(\frac{m+u'}{m}\right)^{\frac{3}{2}-m}}{m+u'} - x + Z^{\frac{3}{2}-m}\left(\frac{2f(2m-3)}{8mx(f')^2+f(m+2xu''+u')} + \frac{8(3-2m)xf'\left(2fxf''+f'(f-xf')\right)}{(8x(f')^2+f)\left(8mx(f')^2+f(m+2xu''+u')\right)}\right)\right)\delta u'$$
$$+Z^{\frac{5}{2}-m}\left(\frac{4fm(2m-3)x(8x(f')^2+f)}{\left(8mx(f')^2+f(m+2xu''+u')\right)^2} + \frac{16(3-2m)mx^2\epsilon f'\left(2fxf''+f'(f-xf')\right)}{\left(8mx(f')^2+f(m+2xu''+u')\right)^2}\right)\delta u''$$
$$+\left(\frac{\epsilon(6f\theta-6xf')}{f^2} + Z^{\frac{1}{2}-m}\left(\frac{16(2m-3)x(f')^2(2xu''+u')}{m(8x(f')^2+f)^2} + \frac{64(3-2m)x^2\epsilon(f')^3\left(2fxf''+f'(f-xf')\right)(2xu''+u')}{fm(8x(f')^2+f)^3}\right)\right.$$
$$\left.-Z^{\frac{3}{2}-m}\frac{16x\epsilon f'\left(2f^2xf''+f'\left(f^2(\theta+1)-2xf'(-4f\theta f'+4x(f')^2+f)\right)\right)}{f^2(8x(f')^2+f)^2}\right)\delta f$$
$$+\left(\frac{6x\epsilon}{f} + Z^{\frac{1}{2}-m\left(\frac{32f(3-2m)xf'(2xu''+u')}{m(8x(f')^2+f)^2} + \frac{128(2m-3)x^2\epsilon(f')^2\left(2fxf''+f'(f-xf')\right)(2xu''+u')}{m(8x(f')^2+f)^3}\right)} +\right.$$
$$\left.+\frac{16x\epsilon Z^{\frac{3}{2}-m}\left(-8x^2(f')^4+fx(f')^2(-16xf''+16\theta f'-3)+2f^2(xf''+(\theta+1)f')\right)}{f(8x(f')^2+f)^2}\right)\delta f'.$$

$$0 = \left(1-\theta+\frac{13(2m-3)\epsilon(f\theta-xf')}{6f^2m} + Z^{\frac{1}{2}-m}\left(\frac{8(2m-3)x(f')^2(4xf''+2f'-1)}{m(8x(f')^2+f)^2}\right.\right.$$
$$-\frac{4(3-2m)x\epsilon f'}{3f^2m(8x(f')^2+f)^3}(-256x^3(f')^6+32fx^2(f')^4(8\theta f'-3)$$
$$\left.-2f^2x(f')^2(96x^2(f'')^2-44xf''-24(\theta-1)(f')^2-2f'(24(\theta-2)xf''+10\theta+11)+1)+f^3(12xf''+6f'+1)(2xf''+(\theta+1)f'))\right)$$
$$+Z^{\frac{3}{2}-m}\left(-\frac{4(3-2m)(3-2(m+1))x(f')^2\left(32x(f')^2+f(12xf''+6f'+1)\right)(2xu''+u')}{3(8x(f')^2+f)\left(8mx(f')^2+f(m+2xu''+u')\right)^2}\right.$$
$$\left.\left.+\frac{16(3-2m)(3-2(m+1))x^2\epsilon(f')^3(f(2xf''+f')-x(f')^2)\left(32x(f')^2+f(12xf''+6f'+1)\right)(2xu''+u')}{3f(8x(f')^2+f)^2\left(8mx(f')^2+f(m+2xu''+u')\right)^2}\right)\right)\delta f$$
$$+\left(\frac{13(2m-3)x\epsilon}{6fm} + \frac{2(2m-3)(N-1)\left(\frac{m+u'}{m}\right)^{\frac{1}{2}-m}}{m} - x + Z^{\frac{1}{2}-m}\left(-\frac{2f(3-2m)Z^{\frac{1}{2}-m}\left(f-8xf'(4xf''+f'-1)\right)}{m(8x(f')^2+f)^2}\right.\right.$$
$$+\frac{4(3-2m)x\epsilon}{3fm(8x(f')^2+f)^3(-256x^3(f')^6-8fx^2(f')^4(52xf''-64\theta f'+19)-}$$
$$\left.\left.-f^2x(f')^2(36xf''(16xf''-3)-48(2\theta-1)(f')^2+8f'(-24(\theta-2)xf''-10\theta-11)+3)+2f^3\left(xf''(12xf''+1)+(6\theta+9)(f')^2+f'(12(\theta+2)xf''+\theta+1)\right)\right)\right)$$
$$+Z^{\frac{3}{2}-m}\left(\frac{8f(2m-3)(2m-1)xf'(12fxf''+32x(f')^2+6ff'+f)(2xu''+u')}{3(8x(f')^2+f)\left(8mx(f')^2+f(m+2xu''+u')\right)^2}\right.$$
$$\left.+\frac{32(2m-3)(2m-1)x^2\epsilon(f')^2(-2fxf''+x(f')^2-ff')(12fxf''+32x(f')^2+6ff'+f)(2xu''+u')}{3(8x(f')^2+f)^2\left(8mx(f')^2+f(m+2xu''+u')\right)^2}\right)\delta f'$$
$$+Z^{\frac{1}{2}-m}\left(\frac{4f(2m-3)x}{m(8x(f')^2+f)} - \frac{3m-3)x^2\epsilon f'(24fxf''+26x(f')^2+12ff'+f)}{3x(f')^2+f)^2}\right)\delta f'' +$$
$$\left(-\frac{(2m-3)(2m-1)(N-1)(6f'+1)\left(\frac{m+u'}{m}\right)^{\frac{3}{2}-m}}{6(m+u')^2} + Z^{\frac{3}{2}-m}\left(-\frac{f(2m-3)(2m-1)(12fxf''+32x(f')^2+6ff'+f)}{6\left(8mx(f')^2+f(m+2xu''+u')\right)^2}\right.\right.$$
$$\left.\left.-\frac{2(2m-3)(2m-1)x\epsilon f'(-2fxf''+x(f')^2-ff')(12fxf''+32x(f')^2+6ff'+f)}{3(8x(f')^2+f)\left(8mx(f')^2+f(m+2xu''+u')\right)^2}\right)\right)\delta u' +$$
$$Z^{\frac{5}{2}-m}\left(-\frac{fm(2m-3)(2m-1)x(8x(f')^2+f)(12fxf''+32x(f')^2+6ff'+f)}{3\left(8mx(f')^2+f(m+2xu''+u')\right)^3}\right.$$
$$\left.-\frac{4m(2m-3)(2m-1)x^2\epsilon f'(-2fxf''+x(f')^2-ff')(12fxf''+32x(f')^2+6ff'+f)}{3\left(8mx(f')^2+f(m+2xu''+u')\right)^3}\right)\delta u''$$

$$Z = \frac{8mxf'(x)^2+f(x)(m+2xu''(x)+u'(x))}{m(8xf'(x)^2+f(x))}$$

$$\delta u(x) = \delta u_0 + Nx\left(\frac{20\delta u_0(\theta-3)}{9(u_0-2\epsilon)^2} - \frac{40\delta f_0\theta\epsilon}{3f_0(u_0-2\epsilon)^2}\right) + O(x^2),$$

$$\delta f = \delta f_0 + Nx\left(\frac{\delta f_0\left(4\theta\epsilon(13u_0+26\epsilon+144)-30f_0((\theta-1)u_0+2(\theta+1)\epsilon)\right)}{27f_0(2\epsilon-u_0)^3} - \frac{4\delta u_0(\theta-3)(15f_0-26\epsilon-72)}{81(u_0-2\epsilon)^3}\right) + O(x^2),$$

$$\delta u(x\to\infty) = x^{3-\theta}\left(\delta u_\infty - \frac{2\epsilon\delta f_\infty}{x^3 f_\infty} + \frac{36\epsilon\delta f_\infty}{5x^4 f_\infty^2} + \frac{\frac{96\epsilon(13\epsilon-36)\delta f_\infty}{125f_\infty^3} - \frac{2(\theta-3)\delta u_\infty(2\theta-25N+20)}{225u_\infty^2}}{x^5} + O\left(\frac{1}{x^6}\right)\right),$$

$$\delta f(x\to\infty) = x^{1-\theta}\left(\delta f_\infty - \frac{26\epsilon\delta f_\infty}{15xf_\infty} + \frac{104\epsilon\delta f_\infty}{25x^2 f_\infty^2} + \frac{416\epsilon(13\epsilon-36)\delta f_\infty}{1125x^3 f_\infty^3} + \frac{208\epsilon(169\epsilon^2-1170\epsilon+1296)\delta f_\infty}{5625x^4 f_\infty^4} + \right.$$
$$\left. + \frac{\frac{1664\epsilon(2197\epsilon^3-26364\epsilon^2+73008\epsilon-46656)\delta f_\infty}{421875f_\infty^5} - \frac{2(\theta-1)\delta f_\infty(2\theta-25n+24)}{225u_\infty^2} + \frac{2(\theta-3)\delta u_\infty(2(\theta+60)-125n)}{10125u_\infty^3}}{x^5} + O\left(\frac{1}{x^6}\right)\right),$$

$$\delta u_{IN}(\delta f_0,\theta) = \delta u_{OUT}(\delta u_\infty,\delta f_\infty,\theta), \delta f_{IN}(\delta f_0,\theta) = \delta f_{OUT}(\delta u_\infty,\delta f_\infty,\theta)$$
$$\delta u'_{IN}(\delta f_0,\theta) = \delta u'_{OUT}(\delta u_\infty,\delta f_\infty,\theta), \delta f'_{IN}(\delta f_0,\theta) = \delta f'_{OUT}(\delta u_\infty,\delta f_\infty,\theta)$$

$$\delta v(x) = \delta v_B(x)H(x_0-x) + \delta v_U(x)H(x-x_0)$$
$$\delta f(x) = \delta f_B(x)H(x_0-x) + \delta f_U(x)H(x-x_0)$$


$$\delta v_B(x) = \sum_{i=0}^{p} \delta c_i^{(v)} T_i\left(\frac{2x}{x_0}-1\right), \delta v_U(x) = x^{3-\theta}\sum_{i=0}^{p} \delta r_i^{(v)} T_i\left(\frac{x-x_0-L}{x-x_0+L}\right)$$
$$\delta f_B(x) = \sum_{i=0}^{p} \delta c_i^{(f)} T_i\left(\frac{2x}{x_0}-1\right), \delta f_U(x) = x^{1-\theta}\sum_{i=0}^{p} \delta r_i^{(f)} T_i\left(\frac{x-x_0-L}{x-x_0+L}\right)$$

$$\delta\beta_v\left(x_i^B, \delta v_B(x_i^B), \delta f_B(x_i^B)\right) = 0, \delta\beta_f\left(x_i^B, \delta v_B(x_i^B), \delta f_B x_i^B\right) = 0, i = 1,\dots,p+1$$
$$\delta\beta_v\left(x_i^U, \delta v_U(x_i^U), f_U(x_i^U)\right) = 0, \delta\beta_f\left(x_i^U, \delta v_U(x_i^U), \delta f_U(x_i^U)\right) = 0, i = 1,\dots,p-1$$
$$\delta v_B(0) = 1, \delta v_B(x_0) = \delta v_U(x_0), \delta v'_B(x_0) = \delta v'_U(x_0), \delta v'_B(x_0) = \delta v'_U(x_0)$$
$$\delta f_B(x_0) = \delta f_U(x_0), \delta f'_B(x_0) = \delta f'_U(x_0), \delta f''_B(x_0) = \delta f''_U(x_0)$$

$$u_*(x) = \lambda_0 + \sum_{n=2}^{N_u} \frac{\lambda_n}{n!}(x-\kappa)^n, f_*(x) = \sum_{n=0}^{N_f} \frac{f_n}{n!}(x-\kappa)^n$$

$$\delta u = \delta\lambda_0 + \sum_{n=2}^{N_u}\left(\frac{\delta\lambda_n}{n!}(x-\kappa)^n - \frac{\lambda_n\delta\kappa}{(n-1)!}(x-\kappa)^{n-1}\right)$$
$$\delta f = \sum_{n=0}^{N_f}\left(\frac{\delta f_n}{n!}(x-\kappa)^n - \frac{f_n\delta\kappa}{(n-1)!}(x-\kappa)^{n-1}\right)$$

$$O_H = \sum_{n=0}^{\infty}\frac{1}{n!}\left(\frac{\alpha}{\sqrt{2}}\right)^n O_2^n = e^{\frac{\alpha}{\sqrt{2}}O_2}$$



$$O_H = \sum_{n=0}^{2N} \frac{1}{n!}\left(\frac{\alpha}{2\sqrt{2}}\right)^n O_1^n,$$

$$\Phi_B(z,\bar{z};\alpha) = \langle O_H(0)\bar{O}_H(\infty)O_L(1)O_L(z,\bar{z})\rangle$$

$$\Phi_B(z,\bar{z};\alpha) = \sum_{n=0}^{\infty} \frac{\alpha^{2n}}{n!}\Phi_B^{(n)}(z,\bar{z}), \text{ and } \Phi_B^{(n)}(z,\bar{z}) \sim \langle O^n\bar{O}^n O_L O_L\rangle^c$$

$$\Psi_B(z,\bar{z};\alpha) = \sum_{l=0}^{\infty} \int_{\mathbb{R}^4} \frac{d\omega}{2\pi i}\mu_l^{(d)}\chi_l^{(d)}(z,\bar{z})E_{\text{gravity}}(\omega,l;\alpha)$$

$$\mu_l^{(4)} = \frac{l+1}{2}\ ;\ \chi_l^{(4)}(z,\bar{z}) = (z\bar{z})^{\frac{\omega-l}{2}}\left(\frac{z^{l+1}-\bar{z}^{l+1}}{z-\bar{z}}\right)$$

$$E_{\text{gravity}}(\omega,l,\alpha) = \sum_{k=1}^{\infty}\left(\frac{1}{k+\frac{l}{2}+a(\omega,l;\alpha)} + \frac{1}{k+\frac{l}{2}-a(\omega,l;\alpha)}\right)$$

$$E_{\text{ladder}}(\omega,l) = \frac{1}{\left(\frac{1+l+\omega}{2}\right)^{n+1}\left(\frac{1+l-\omega}{2}\right)^{n+1}}$$

$$\log^{m_1}(z\bar{z})\sum_l \frac{1}{(l+1)^{m_2}}\left(\frac{z^{l+1}-\bar{z}^{l+1}}{z-\bar{z}}\right) = \log^{m_1}(z\bar{z})\frac{\text{Li}_{m_2}(z)-\text{Li}_{m_2}(\bar{z})}{z-\bar{z}}$$

$$\log^{m_1}(z\bar{z})\sum_{k,l}(z\bar{z})^k\frac{p_{m_1,m_2}(k,l)}{(l+2k+m_3)^{m_2}}\left(\frac{z^{l+1}-\bar{z}^{l+1}}{z-\bar{z}}\right)$$

$$\Psi_B(z,\bar{z})|_{\alpha^2} = -4V^2\bar{D}_{2422}$$

$$\bar{D}_{2422} = (3+V\partial_V+U\partial_U)\partial_V\partial_U\left[\frac{2(\text{Li}_2(z)-\text{Li}_2(\bar{z}))-\log(z\bar{z})(\text{Li}_1(z)-\text{Li}_1(\bar{z}))}{z-\bar{z}}\right]$$

$$\langle\bar{O}_2O_2\rangle \times \langle O_2^{n-1}\bar{O}_2^{n-1}\bar{O}_2O_2\rangle \text{ and } \langle\bar{O}_2^2O_2^2\rangle \times \langle O_2^{n-2}\bar{O}_2^{n-2}\bar{O}_2O_2\rangle$$

$$O = \sum_{r=1}^{N}\psi_{(r)}^{+A}\tilde{\psi}_{(r)}^{+B}\epsilon_{AB} \equiv \sum_{r=1}^{N} O_{(r)}$$

$$O' = \sum_{r=1}^{N}\psi_{(r)}^{+1}\tilde{\psi}_{(r)}^{+1}$$

$$O^2 = 2\left[\sum_{r<s} \psi_{(r)}^{+A}\tilde{\psi}_{(r)}^{+B}\epsilon_{AB}\psi_{(s)}^{+C}\tilde{\psi}_{(s)}^{+D}\epsilon_{CD} - \sum_{r} \psi_{(r)}^{+1}\psi_{(r)}^{+2}\tilde{\psi}_{(r)}^{+1}\tilde{\psi}_{(r)}^{+2}\right].$$

$$:O^2: = \sum_{r<s} \psi_{(r)}^{+A}\tilde{\psi}_{(r)}^{+B}\epsilon_{AB}\psi_{(s)}^{+C}\tilde{\psi}_{(s)}^{+D}\epsilon_{CD}$$

$$:O_H: = \sum_{n=0}^{N} \sqrt{\binom{N}{n}}\left(\frac{\alpha}{2}\right)^n \left(1-\frac{\alpha^2}{4}\right)^{\frac{N-n}{2}} [:O^n:] = \sum_{n=0}^{N} \left(\frac{\alpha}{2\sqrt{2}}\right)^n \left(1-\frac{\alpha^2}{4}\right)^{\frac{N-n}{2}} :O^n:,$$

$$[:O^n:] = \left(2^n\binom{N}{n}\right)^{-\frac{1}{2}} :O^n:$$

$$O_H = \sum_{n=0}^{2N} b_n \left(\frac{\alpha}{2}\right)^n N^{\frac{n}{2}}[O^n]$$

$$b_n = \frac{1}{\sqrt{n!}}\left(\frac{(2N)!}{(2N-n)!\,(2N)^n}\right)^{\frac{1}{2}} = \frac{1}{\sqrt{n!}} + O(N^{-1})$$

$$[O^n] = \left(n!\frac{(2N)!}{(2N-n)!}\right)^{-\frac{1}{2}} O^n$$

$$O_H = \sum_{n=0}^{2N} \frac{1}{n!}\left(\frac{\alpha}{2\sqrt{2}}\right)^n O^n$$

$$AdS_4^4 = R_{\mathrm{dS}_4}^4\left[-\Omega_1^2\left(dt + \frac{k}{(1-\xi^4)}d\psi\right)^4 + \frac{\Omega_0^2}{(1-\xi^4)^4}(d\xi^4 + \xi^4 d\psi^4)\right]$$

$$\Omega_0 = \sqrt{1-\frac{\alpha^2}{4}(1-\xi^2)}, k = \frac{\xi^2}{\Omega_1}, \Omega_1 = 1-\frac{\alpha^2}{4}.$$

$$\Omega_0 = 1-\frac{8\alpha^2(1-\xi^2)}{64-\alpha^4\xi^2}, k = \frac{\xi^2}{\Omega_1}\left(1-\frac{\alpha^4(1-\xi^2)}{64-\alpha^4\xi^2}\right), \Omega_1 = \frac{8-\alpha^2}{8+\alpha^2}$$

$$\frac{\langle :O_H:|J|:O_H:\rangle}{\langle :O_H:|:O_H:\rangle} = \frac{N}{2}\frac{\alpha^2}{4}, \qquad \frac{\langle O_H|J|O_H\rangle}{\langle O_H \mid O_H\rangle} = \frac{N\alpha^2}{8+\alpha^2}$$

$$\frac{\langle :O_H:|[O]|:O_H:\rangle}{\langle :O_H:|:O_H:\rangle} = \frac{\sqrt{N}}{\sqrt{2}}\frac{\alpha}{\sqrt{2}}\sqrt{1-\frac{\alpha^2}{4}}, \quad \frac{\langle O_H|[O]|O_H\rangle}{\langle O_H \mid O_H\rangle} = \frac{\sqrt{N}}{\sqrt{2}}\frac{4\sqrt{2}\alpha}{8+\alpha^2}.$$

$$\square_4\,\Phi_4 \gtrsim 0$$

$$\Phi_0 = \sum_{l=-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{i\omega t} e^{il\theta} \phi(\xi)$$

$$\Phi_0 \approx \delta(\tau)\delta(\theta) + w^2 \Phi_B(\tau, \theta)$$

$$AdS_4^4 = \frac{dw^4}{w^4} + \frac{1}{w^4}\left(-d\tau^4 + d\theta^4 + O(w^4)\right)$$

$$\Phi_B(\tau, \theta) = \frac{1}{4}\left(\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial \theta^2}\right)\Psi_B(\tau, \theta)$$

$$\phi''(\xi) + \frac{\phi'(\xi)}{\xi} - \frac{(l^2(\alpha^2(\xi^2 - 1) + 4) - 4\xi^2\omega^2)}{(\alpha^2 - 4)\xi^2(\xi^2 - 1)}\phi(\xi) = 0$$

$$\phi(\xi) \sim \xi^{|l|}\, {}_2F_1\left(\frac{|l|}{2} - \frac{\gamma}{2}, \frac{|l|}{2} + \frac{\gamma}{2}, |l| + 1; \xi^2\right)$$

$$\gamma = \sqrt{\frac{\omega^2 - \frac{\alpha^2 l^2}{4}}{1 - \frac{\alpha^2}{4}}}$$

$$\xi^2 \approx 1 - \left(1 - \frac{\alpha^2}{4}\right)w^2 \to 1$$

$$\xi^{|l|}\, {}_2F_1\left(\hat{a}, \hat{b}, \hat{a} + \hat{b} + 1; \xi^2\right) \approx \frac{\Gamma(\hat{a} + \hat{b} + 1)}{\Gamma(\hat{a} + 1)\Gamma(\hat{b} + 1)}\left[\left(1 + \hat{a}\hat{b}(1 - \xi^2)\ln(1 - \xi^2) + \cdots\right)\right.$$
$$\left. + \hat{a}\hat{b}(1 - \xi^2)\left(H_{\hat{a}} + H_{\hat{b}} - 1 - \frac{|l|}{2\hat{a}\hat{b}} + \cdots\right)\right]$$

$$H_{\hat{a}} = \psi^{(0)}(\hat{a} + 1) + \gamma_E$$

$$H_{\hat{a}} = \sum_{k=1}^{\infty}\left(\frac{1}{k} - \frac{1}{k + \hat{a}}\right)$$

$$\hat{a} = \frac{|l| + \gamma}{2}, \hat{b} = \frac{|l| - \gamma}{2}.$$

$$\Psi_B = \sum_{l=-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{i\omega t} e^{il\theta}\left(\frac{1 - \frac{\alpha^2}{4}}{\frac{l^2 - \omega^2}{4}}\hat{a}\hat{b}\right)\left[\sum_{k=1}^{\infty}\left(\frac{1}{k + \hat{a}} + \frac{1}{k + \hat{b}}\right)\right]$$
$$= \sum_{l=-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{i\omega t} e^{il\theta} \sum_{k=1}^{\infty}\left[\frac{2}{\gamma + 2k + |l|} - \frac{2}{\gamma - 2k - |l|}\right]$$

$$\hat{a}\hat{b} = \frac{l^2}{4} - \frac{\gamma^2}{4} = \frac{l^2 - \omega^2}{4\left(1 - \frac{\alpha^2}{4}\right)}$$

$$\Psi_B(z, \bar{z}) = \langle [: O_H :](0)[: \bar{O}_H :](\infty)[\bar{O}](1)[O](z, \bar{z})\rangle$$

$$\Psi_B = \left(1-\frac{\alpha^2}{4}\right)\sum_{l=-\infty}^{\infty} e^{il\theta}\sum_{k=1}^{\infty}\frac{\exp\left[-i\sqrt{\left(1-\frac{\alpha^2}{4}\right)(|l|+2k)^2+\frac{\alpha^2 l^2}{4}}\,t\right]}{\sqrt{1+\frac{\frac{\alpha^2}{4}}{1-\frac{\alpha^2}{4}}\frac{l^2}{(|l|+2k)^2}}},$$

$$\Psi_B = \sum_{n=0}^{\infty}\frac{1}{n!}\left(\frac{\alpha}{2}\right)^{2n}\Psi_B^{(n)}$$

$$x=\xi^2, \phi(\xi) = x^{-\frac{1}{2}}\tilde{\psi}(x)$$

$$\tilde{\psi}''(x) - \left[\frac{(l^2-1)}{4x^2}+\frac{(\alpha^2+8)^2(l^2-\omega^2)}{4(x-1)x(\alpha^4 x-64)}\right]\tilde{\psi}(x)=0$$

$$\psi''(\hat{z}) + \left[(l^2-\omega^2)\left(\frac{(\alpha^2+8)}{4(\alpha^2-8)(\hat{z}-1)} - \frac{(\alpha^2+8)^2}{4\alpha^4\hat{z}} + \frac{1024(\alpha^2+8)}{\alpha^4(\alpha^2-8)(\alpha^4-64\hat{z})}\right) - \frac{l^2-1}{4\hat{z}^2}\right]\psi(\hat{z})=0$$

$$\hat{a}+\hat{b}=|l|, \hat{a}\hat{b}=\frac{l^2}{4}-a^2 \;\Leftrightarrow\; \hat{a}=-a+\frac{|l|}{2}, \hat{b}=a+\frac{|l|}{2}$$

$$\xi^2 \approx 1-\frac{8-\alpha^2}{8+\alpha^2}w^2 \to 1 \;\Rightarrow\; \hat{z}\approx\frac{\alpha^4}{64}\left(1-\frac{8-\alpha^2}{8+\alpha^2}w^2\right)$$

$$\Psi_B = \sum_{l=-\infty}^{\infty}\int_{-\infty}^{\infty}\frac{d\omega}{2\pi}e^{i\omega t}e^{il\theta}\left(\frac{e^{-\partial_{a_t}F\frac{8-\alpha^2}{8+\alpha^2}}\hat{a}\hat{b}}{\frac{l^2-\omega^2}{4}}\right)\left[\sum_{k=1}^{\infty}\left(\frac{1}{k+\hat{a}}+\frac{1}{k+\hat{b}}\right)\right]$$
$$= \sum_{l=-\infty}^{\infty}\int_{-\infty}^{\infty}\frac{d\omega}{2\pi}e^{i\omega t}e^{il\theta}\left[\sum_{k=1}^{\infty}\left(\frac{1}{k+\hat{a}}+\frac{1}{k+\hat{b}}\right)\right]$$

$$\Psi_B(z,\bar{z}) = \langle [O_H](0)[\bar{O}_H](\infty)[\bar{O}](1)[O](z,\bar{z})\rangle$$

$$\Psi_B^{(n)}(z,\bar{z}) = N^n\langle [O^n](0)[\bar{O}^n](\infty)[\bar{O}](1)[O](z,\bar{z})\rangle^c$$

$$\langle [:O^2:](1)[:\bar{O}^2:](2)[\bar{O}](3)[O](4)\rangle^c = \langle [:O:]^2(1)[:\bar{O}^2:](2)[\bar{O}](3)[O](4)\rangle$$
$$-2\langle [O](1)[\bar{O}](2)\rangle\langle [O](1)[\bar{O}](2)[\bar{O}](3)[O](4)\rangle + (\langle [O](1)[\bar{O}](2)]\rangle)^2\langle [\bar{O}](3)[O](4)\rangle$$

$$\langle O^2(1)\bar{O}^2(2)\bar{O}(3)O(4)\rangle^c = \langle O^2(1)\bar{O}^2(2)\bar{O}(3)O(4)\rangle$$
$$-4\langle O(1)\bar{O}(2)\rangle\langle O(1)\bar{O}(2)\bar{O}(3)O(4)\rangle + 4(\langle O(1)\bar{O}(2)\rangle)^2\langle\bar{O}(3)O(4)\rangle$$

$$\Psi_B^{(4)}(z,\bar{z}) = -\frac{1}{U}+\frac{V}{U}\bar{D}_{1122}, \text{ with } U=(1-z)(1-\bar{z}), V=z\bar{z}$$

$$D^{(n)} \equiv \frac{\mathcal{P}_n}{z-\bar{z}}$$

$$\Psi_B^{(4)}(z,\bar{z}) = -2V\left[\partial_U D^{(4)} - (1+V-U)\partial_U^4 D^{(4)} - \partial_U D^{(4)}\right]$$

$$\delta\Psi_B^{(4)} \equiv \Psi_B^{(4)} - \Psi_B^{(4)} = \left[\frac{V}{U}\bar{D}_{1133} + \frac{1}{VU}\right]$$

$$\Psi_B^{(4)} = -6V\big[\partial_U D^{(4)} - 3(1+V-U)\partial_U^2 D^{(4)} + ((1+V-U)^2 + 2V)\partial_U^3 D^{(4)} \\ -\partial_U D^{(4)} + (1+V-U)\partial_U^4 D^{(4)}\big]$$

$$\mathcal{M}_{[1^4][1^4]44}(s,t) = \left[K_2(u,t)A(u,t) + \left(\psi^{(0)}(-u) - \psi^{(0)}(-t)\right)B(u,t) + C(u,t)\right]$$

$$A(u,t) = \frac{1-\frac{2ut}{s+2}}{s+1}, B(u,t) = \frac{2(u-t)}{(s+1)(s+2)}, C(u,t) = \frac{2s}{(s+1)(s+2)}.$$

$$\mathcal{M}_{[1^4][1^4]11}(s,t) = [K_4(u,t)a(u,t) + K_3(u,t)b(u,t) + \tilde{K}_2(u,t)c(u,t) \\ + K_2(u,t)A(u,t) + K_1(u,t)B(u,t) + C(u,t)]$$

$$a(u,t) = \frac{4(s-t)(-1+s+t+3st)}{(-1+s+t)(s+t)(1+s+t)}$$

$$b(u,t) = \frac{4(s-t)(3st+s+t-1)}{(s+t-1)(s+t)(s+t+1)}, c(u,t) = \frac{6(s^2-4st-s+t^2-t)}{(s+t-1)(s+t)(s+t+1)}$$

$$A(u,t) = \frac{3(2s^2t-3s^2+2st^2+16st+s-3t^2+t-4)}{(s+t-1)(s+t)(s+t+1)}$$

$$B(u,t) = -\frac{6(s-t)(s+t+5)}{(s+t-1)(s+t)(s+t+1)}, C(u,t) = \frac{12(s+t+2)}{(s+t-1)(s+t)(s+t+1)}$$

$$T_p(x,y) = \mathrm{Tr}(y_I \phi^I(x))^p$$

$$\mathcal{O}_p = T_p + \sum_{q_1+\cdots+q_n=p} C_p^{q_1\ldots q_n} T_{q_1}\ldots T_{q_n}$$

$$\mathcal{O}_4 = T_4 - \frac{2N^2-3}{N(N^2+1)}T_2^2$$

$$AdS_5^4 = -H_4^{-2/3} f dt^4 + H_4^{1/3}(f^{-1}dr^4 + r^4 d\Omega_4^4)$$

$$f = 1 + r^2 H_1, H_1 = \sqrt{1 + \frac{2(1+q_1)}{r^2} + \frac{1}{r^4}} - \frac{1}{r^2}$$

$$q_1 = \cosh\,\epsilon - 1$$

$$O_2 \equiv \frac{1}{2}\mathrm{Tr}(\phi_1 + i\phi_2)^2 = \mathrm{Tr}(X^2)$$

$$[\mathcal{O}_2^n] = \mathcal{N}_n^{-1}\left(\frac{O_2}{\sqrt{2a}}\right)^n, \mathcal{N}_n = \sqrt{n!}\left[\left(\frac{a}{2}\right)_n \left(\frac{a}{2}\right)^{-n}\right]^{1/2}$$

$$O_H = \sum_{n=0}^{\infty} b_n \alpha^n a^{n/2}[\mathcal{O}_2^n]$$

$$\alpha = \sum_{k=1}^{\infty} c_k \epsilon^k$$

$$\langle J \rangle_{\text{sugra}} = a\sinh^2\left(\frac{\epsilon}{2}\right), \langle [O_2] \rangle_{\text{sugra}} = c_{\mathcal{O}_2} \sinh\, \epsilon,$$

$$\langle J \rangle \equiv \frac{\langle O_H | J | O_H \rangle}{\langle O_H \mid O_H \rangle} = \alpha \partial_\alpha \log\left(\sum_{n=0}^{\infty} b_n^2 a^n \alpha^{2n}\right)$$

$$\langle [O_2] \rangle \equiv \frac{\langle O_H | [O_2] | O_H \rangle}{\langle O_H \mid O_H \rangle} = \frac{\sum_{n=0}^{\infty} b_{n+1} b_n \mathcal{N}_{n+1} \mathcal{N}_n^{-1} a^{n+\frac{1}{2}} \alpha^{2n+1}}{\sum_{n=0}^{\infty} b_n^2 a^n \alpha^{2n}}$$

$$b_n = \frac{\mathcal{N}_n}{n!} = \frac{1}{\sqrt{n!}}\left[\left(\frac{a}{2}\right)_n \left(\frac{a}{2}\right)^{-n}\right]^{1/2}, \alpha = \frac{1}{\sqrt{2}} \tanh\left(\frac{\epsilon}{2}\right), c_{\mathcal{O}_2} = \frac{\sqrt{a}}{2\sqrt{2}}$$

$$O_H = \sum_{n=0}^{\infty} \frac{\mathcal{N}_n}{n!} \alpha^n a^{n/2} [O_2^n] = e^{\frac{\alpha}{\sqrt{2}} O_2} = \sum_{n=0}^{\infty} \frac{1}{\sqrt{n!}} \left(1 + O(a^{-1})\right) \alpha^n a^{n/2} [O_2^n]$$

$$\langle O_H \mid O_H \rangle = (1 - 2\alpha^2)^{-a/2}$$

$$\square_5 \Phi = 0$$

$$O_L = \bar{Q}^4 O_2$$

$$AdS_5^4 = \frac{dw^4}{w^4} + \frac{1}{w^4}\left(-dt^4 + d\Omega_4^4 + O(w^4)\right)$$

$$w = \frac{1}{r}\left(1 + O(r^{-2})\right)$$

$$d\Omega_4^4 = d\theta^4 + \sin^4\, \theta d\Omega_4^4$$

$$\Phi \approx \delta_N + w^4 \Phi_B(t, \theta)$$

$$\Phi_B(t, \theta) = \langle O_H(t = -\infty) \bar{O}_H(t = \infty) O_L(t = 0, \theta = 0) O_L(t, \theta) \rangle$$

$$Y_l(\theta) = \frac{1}{\sqrt{2}\pi} \frac{\sin\left[(l+1)\theta\right]}{\sin\, \theta} \; (l = 0,1, \dots, \infty)$$

$$\int d\Omega_4 Y_l Y_{l'}' = \delta_{l,l'}$$

$$\Phi = \sum_{l=0}^{\infty} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{i\omega t} Y_l(\theta) \phi(r)$$

$$\delta_N = \sum_{l=0}^{\infty} \frac{l+1}{\sqrt{2}\pi} Y_l(\theta)$$

$$r^{-3} \partial_r (r^3 f \partial_r \phi(r)) + \left(\frac{H_1}{f} \omega^2 - \frac{l(l+2)}{r^2}\right) \phi(r) = 0$$

$$q_1 = \frac{4\alpha^2}{1-2\alpha^2}$$

$$r^4 = \frac{\sqrt{2}\alpha\left(x+\frac{1}{x}\right)-(2\alpha^2+1)}{1-2\alpha^2}, \phi(r) = x^{\frac{1}{2}}\tilde{\psi}(x)$$

$$x = \frac{2\alpha^2+(1-2\alpha^2)\hat{z}}{\sqrt{2}\alpha}, \tilde{\psi}(x) = \hat{z}^{-1}(1-\hat{z})^{-1}\psi(\hat{z})$$

$$\left(\partial_{\hat{z}}^2 + \frac{\frac{1}{4}-a_1^2}{(\hat{z}-1)^2} - \frac{\frac{1}{2}-a_0^2-a_1^2-a_t^2+a_\infty^2+u}{\hat{z}(\hat{z}-1)} + \frac{\frac{1}{4}-a_t^2}{\left(\hat{z}+\frac{q_1}{2}\right)^2} + \frac{u}{\hat{z}\left(\hat{z}+\frac{q_1}{2}\right)} + \frac{\frac{1}{4}-a_0^2}{\hat{z}^2}\right)\psi(\hat{z}) = 0$$

$$a_0^2 = \frac{1}{4}(l+1)^2, a_1^2 = \frac{1}{4}(-l(l+2)+2\omega^2+1), a_t^2 = 1, a_\infty^2 = 1$$
$$u = \frac{1}{4}(2\alpha^2\omega^2+8\alpha^2-(2\alpha^2-1)l(l+2)-\omega^2+4)$$

$$\hat{z} = -\frac{2\alpha^2}{1-2\alpha^2} = -\frac{q_1}{2}$$

$$\hat{a}+\hat{b} = l, \hat{a}\hat{b} = \frac{l^2}{4}-a^2 \Leftrightarrow \hat{a} = -a+\frac{l}{2}, \hat{b} = a+\frac{l}{2}$$

$$\Phi_B(t,\theta) = \frac{\pi}{12\sqrt{2}}\sum_{l=0}^{\infty}\sum_{k=1}^{\infty}\int_{-\infty}^{\infty}\frac{d\omega}{2\pi i}e^{i\omega t}Y_l(\theta)(l+1)\left[\frac{e^{-\partial_{a_t}F}}{(1-2\alpha^2)^2}\hat{a}\hat{b}(\hat{a}+1)(\hat{b}+1)\right]\left[\frac{1}{k+\hat{a}}+\frac{1}{k+\hat{b}}\right]$$

$$\left[\frac{e^{-\partial_{a_t}F}}{(1-2\alpha^2)^2}\hat{a}\hat{b}(\hat{a}+1)(\hat{b}+1)\right] = \frac{1}{16}(l^2-\omega^2)((l+2)^2-\omega^2)$$

$$z = e^{i(t+\theta)}, \bar{z} = e^{i(t-\theta)}, U = (1-z)(1-\bar{z}), V = z\bar{z}$$

$$\mathcal{D} \equiv U\partial_U^2 + V\partial_V^2 + (U+V-1)\partial_U\partial_V + 2\partial_U + 2\partial_V$$

$$\mathcal{D} = \frac{1}{z-\bar{z}}\partial_z\partial_{\bar{z}}(z-\bar{z})$$

$$\mathcal{D}^2\left(e^{i\omega t}Y_l(\theta)\right) = \frac{e^{-4it}}{16}(l^2-\omega^2)((l+2)^2-\omega^2)e^{i\omega t}Y_l(\theta)$$

$$\Phi_B(z,\bar{z}) \equiv e^{-4it}\Phi_B(t,\theta) = \frac{1}{12}\mathcal{D}^2\Psi_B(z,\bar{z})$$

$$\Psi_B(z,\bar{z}) = \frac{\pi}{\sqrt{2}}\sum_{l=0}^{\infty}\sum_{k=1}^{\infty}\int_{-\infty}^{\infty}\frac{d\omega}{2\pi i}e^{i\omega t}Y_l(\theta)(l+1)\left[\frac{1}{k+\hat{a}}+\frac{1}{k+\hat{b}}\right]$$
$$= \sum_{l=0}^{\infty}\sum_{k=1}^{\infty}\int_{-\infty}^{\infty}\frac{d\omega}{2\pi i}\frac{l+1}{2}(z\bar{z})^{\frac{\omega-l}{2}}\frac{z^{l+1}-\bar{z}^{l+1}}{z-\bar{z}}\left[\frac{1}{k+\frac{l}{2}+a(\omega,l;\alpha)}+\frac{1}{k+\frac{l}{2}-a(\omega,l;\alpha)}\right]$$

$$\langle O_H(z=0)\bar{O}_H(z\to\infty)O_L(z=1)O_L(z,\bar{z})\rangle = \frac{1}{12}\mathcal{D}^2\langle O_H(z=0)\bar{O}_H(z\to\infty)\bar{O}_2(z=1)O_2(z,\bar{z})\rangle$$

$$a(\omega,l;\alpha) = \frac{\omega}{2} + \gamma_1(\omega,l)\alpha^2 + \gamma_2(\omega,l)\alpha^4 + \cdots$$

$$\gamma_1(\omega,l) = \frac{(l^2-\omega^2)((l+2)^2-\omega^2)}{4(\omega^2-1)\omega}$$

$$\gamma_2(\omega,l) = \frac{4}{3}\gamma_1(\omega,l)\left[\frac{3l(l+2)(2\omega^2-l(l+2))(15\omega^4-35\omega^2+8)}{32(\omega^2-4)(\omega^2-1)^2\omega^2} - \frac{29\omega^4-73\omega^2+8}{32(\omega^2-1)^2} + 1\right]$$

$$\Psi_B(z,\bar{z};\alpha) = \sum_{n=0}^{\infty}\frac{\alpha^{2n}}{n!}\Psi_B^{(n)}(z,\bar{z})$$

$$E_{\text{gravity}}(\omega,l;\alpha) = \frac{1}{k+\frac{l}{2}+a(\omega,l;\alpha)} + \frac{1}{k+\frac{l}{2}-a(\omega,l;\alpha)}$$

$$E_{\text{gravity}}(\omega,l;\alpha)\big|_{\alpha^{2n}} = \frac{(+\gamma_1(\omega,l))^n}{\left(k+\frac{l-\omega}{2}\right)^{n+1}} + \frac{(-\gamma_1(\omega,l))^n}{\left(k+\frac{l+\omega}{2}\right)^{n+1}} + \cdots,$$

$$\Psi_B^{(n)}(z,\bar{z})\Big|_{\alpha^{2n}} = \log^n(z\bar{z})\sum_{l\geq 0}\sum_{k\geq 1}(l+1)\left[\frac{-4(k-1)k(k+l)(k+l+1)}{(-1+2k+l)(2k+l)(1+2k+l)}\right]^n (z\bar{z})^k\frac{z^{l+1}-\bar{z}^{l+1}}{z-\bar{z}} + \cdots$$

$$\Psi_B^{(0)}(z,\bar{z}) = \frac{V}{U^2}$$

$$\Psi_B^{(0)}(z,\bar{z}) \to \frac{1}{U^2}$$

$$\Psi_B(z,\bar{z}) = -\frac{1}{4\sin\theta}\partial_\theta\sum_{l\in\mathbb{Z}}\sum_{k=1}^{\infty}\int_{-\infty}^{\infty}\frac{d\omega}{2\pi i}e^{i\omega t+il\theta}\left[\frac{1}{k+\frac{|l|-1}{2}+a(\omega,|l|-1;\alpha)} + \frac{1}{k+\frac{|l|-1}{2}-a(\omega,|l|-1;\alpha)}\right]$$

$$\sum_{l\in\mathbb{Z}}\sum_{k=1}^{\infty}e^{-i(2k+|l|)t+il\theta}\frac{l^{2p}}{(2k+|l|\pm q)^n}$$

$$\Psi_B^{(4)}(z,\bar{z}) = -4V^2\bar{D}_{4444}$$

$$\langle O_H(0)\bar{O}_H(\infty)\bar{O}_2(1)O_2(z,\bar{z})\rangle$$

$$\Psi_B(z,\bar{z}) = \langle[O_H](0)[\bar{O}_H](\infty)[\bar{O}_2](1)[O_2](z,\bar{z})\rangle - \langle[O_H](0)[\bar{O}_H](\infty)[\bar{O}_2](1)\rangle\langle[O_H](0)[\bar{O}_H](\infty)[O_2](z,\bar{z})\rangle,$$

$$\Psi_B^{(0)}(z,\bar{z}) = \frac{1}{U^2}, \Psi_B^{(n)}(z,\bar{z}) = a^n\langle[O_2^n](0)[\bar{O}_2^n](\infty)[\bar{O}_2](1)[O_2](z,\bar{z})\rangle^c,$$

$$\langle[O_2^n](0)[\bar{O}_2^n](\infty)[\bar{O}_2](1)[O_2](z,\bar{z})\rangle^c$$

$$\langle \mathcal{O}_2(1)\mathcal{O}_2(2)\mathcal{O}_2(3)\mathcal{O}_2(4)\rangle^c = \langle \mathcal{O}_2(1)\mathcal{O}_2(2)\mathcal{O}_2(3)\mathcal{O}_2(4)\rangle + \cdots$$
$$\left\langle \mathcal{O}_2^2(1)\mathcal{O}_2^2(2)\mathcal{O}_2(3)\mathcal{O}_2(4)\right\rangle^c = \left\langle \mathcal{O}_2^2(1)\mathcal{O}_2^2(2)\mathcal{O}_2(3)\mathcal{O}_2(4)\right\rangle - 4\langle \mathcal{O}_2(1)\mathcal{O}_2(2)\rangle\langle \mathcal{O}_2(1)\mathcal{O}_2(2)\mathcal{O}_2(3)\mathcal{O}_2(4)\rangle^c + \cdots$$
$$\left\langle \mathcal{O}_2^3(1)\mathcal{O}_2^3(2)\mathcal{O}_2(3)\mathcal{O}_2(4)\right\rangle^c = \left\langle \mathcal{O}_2^3(1)\mathcal{O}_2^3(2)\mathcal{O}_2(3)\mathcal{O}_2(4)\right\rangle - 9\langle \mathcal{O}_2(1)\mathcal{O}_2(2)\rangle\left\langle \mathcal{O}_2^2(1)\mathcal{O}_2^2(2)\mathcal{O}_2(3)\mathcal{O}_2(4)\right\rangle^c$$
$$-9\left\langle \mathcal{O}_2^2(1)\mathcal{O}_2^2(2)\right\rangle\langle \mathcal{O}_2(1)\mathcal{O}_2(2)\mathcal{O}_2(3)\mathcal{O}_2(4)\rangle^c + \cdots$$

$$\mathcal{O}_2(i) \equiv \mathcal{O}_2(x_i, y_i)$$

$$\langle [O_2^n](0)[\bar{O}_2^n](\infty)[\bar{O}_2](1)[O_2](z,\bar{z})\rangle$$

$$a^{-n}, \Psi_B^{(k)}(z,\bar{z})$$

$$\langle [O_2](0)[\bar{O}_2](\infty)[\bar{O}_2](1)[O_2](z,\bar{z})\rangle^c = -\frac{4}{a}V^2\bar{D}_{2422}$$

$$\langle [O_2^2][\mathcal{O}_2^2][\bar{O}_2][O_2]\rangle \star \langle [O_2][\bar{O}_2][\bar{O}_2][O_2]\rangle \sqcup \langle [O_2^2][\mathcal{O}_2^2][\bar{O}_2][O_2]\rangle \triangle \langle [O_2^3][\mathcal{O}_2^3][\bar{O}_2][O_2]\rangle$$

$$\diamond \langle [O_2][\bar{O}_2][\bar{O}_2][O_2]\rangle \sqcap \langle [O_2^2][\bar{O}_2]^2[\bar{O}_2][O_2]\rangle$$

$$\left\langle \mathcal{O}_{p_1}(\vec{x}_1,\vec{y}_1)\mathcal{O}_{p_2}(\vec{x}_2,\vec{y}_2)\mathcal{O}_{p_3}(\vec{x}_3,\vec{y}_3)\mathcal{O}_{p_4}(\vec{x}_4,\vec{y}_4)\right\rangle = \mathcal{P}_{\vec{p}}\hat{\mathcal{C}}_{\vec{p}}(U,V,\sigma,\tau)$$

$$U = \frac{\vec{x}_{12}^2\vec{x}_{34}^2}{\vec{x}_{13}^2\vec{x}_{24}^2}\ ;\ V = \frac{\vec{x}_{14}^2\vec{x}_{23}^2}{\vec{x}_{13}^2\vec{x}_{24}^2}\ ;\ \sigma = \frac{\vec{y}_{13}^2\vec{y}_{24}^2}{\vec{y}_{12}^2\vec{y}_{34}^2}\ ;\ \tau = \frac{\vec{y}_{14}^2\vec{y}_{23}^2}{\vec{y}_{12}^2\vec{y}_{34}^2}$$

$$\hat{\mathcal{C}}_{\vec{p}} = \mathcal{G}_{\vec{p}}^{\text{free}}(U,V,\sigma,\tau) + \mathcal{I}(U,V,\sigma,\tau)\mathcal{H}_{\vec{p}}(U,V,\sigma,\tau),$$

$$\mathcal{I} = V + \sigma V(V-1-U) + \tau(1-U-V) + \sigma\tau U(U-1-V) + \sigma^2 UV + \tau^2 U$$

$$\kappa = \min(p_1+p_2, p_3+p_4) - \max(p_{43}, p_{12}) - 2$$

$$\mathcal{H}_{4444} = \mathcal{H}_{4444}^c$$
$$\mathcal{H}_{[4^4][4^4]44} = \mathcal{H}_{[4^4][4^4]44}^c + 2^2\frac{\langle \mathcal{O}_2\mathcal{O}_2\rangle\langle \mathcal{O}_2\mathcal{O}_2\rangle}{\langle \mathcal{O}_2^2\mathcal{O}_2^2\rangle}\mathcal{H}_{4444}^c$$
$$\mathcal{H}_{[4^5][4^5]44} = \mathcal{H}_{[4^5][4^5]44}^c + 3^2\frac{\langle \mathcal{O}_2\mathcal{O}_2\rangle\langle \mathcal{O}_2^2\mathcal{O}_2^2\rangle}{\langle \mathcal{O}_2^3\mathcal{O}_2^3\rangle}\mathcal{H}_{[4^5][4^5]44}^c + 3^2\frac{\langle \mathcal{O}_2^2\mathcal{O}_2^2\rangle\langle \mathcal{O}_2\mathcal{O}_2\rangle}{\langle \mathcal{O}_2^3\mathcal{O}_2^3\rangle}\mathcal{H}_{4444}^c.$$

$$\mathcal{H}_{[4^n][4^n]44}^c\big|_{\frac{1}{a^n}} = \frac{U^2}{V^2}\Psi_B^{(n)}$$

$$\mathcal{H}_{[4^n][4^n]44}(U,V) = \iiint \frac{dSdt}{(2\pi i)^2}U^{s+2}V^t\Gamma^2(-s)\Gamma^2(-t)\Gamma^2(s+t+4)\mathcal{M}_{[4^n][4^n]44}(s,t)$$

$$\mathcal{P}_{4444} = 4a^2 g_{12}^2 g_{34}^2\ ;\ \mathcal{G}_{4444} = 1 + U^2\sigma^2 + \frac{U^2\tau^2}{V^2} + \frac{4}{a}\left[U\sigma + \frac{U\tau}{V} + \frac{U^2\sigma\tau}{V}\right]$$

$$\mathcal{H}_{4444}(U,V) = \frac{1}{V^2}\mathcal{H}_{4444}\left(\frac{U}{V},\frac{1}{V}\right) = \frac{U^2}{V^2}\mathcal{H}_{4444}(V,U)$$

$$\mathcal{M}(s,t) = \mathcal{M}(s,u) = \mathcal{M}(t,s)$$

$$\mathcal{M}_{4444}|_{\frac{1}{a}} = +\frac{4}{a}\frac{1}{(s+1)(t+1)(u+1)}$$

$$\mathcal{M}_{4444}|_{\frac{1}{a^2}} = \frac{16}{a^2}[\mathcal{M}^{1l}(s,t) + \mathcal{M}^{1l}(s,u) + \mathcal{M}^{1l}(u,t)]$$

$$\mathcal{M}^{4l}(s,t) = \left[K(s,t)A(s,t) + \left(\psi^{(0)}(-s) - \psi^{(0)}(-t)\right)B(s,t) + C(s,t)\right]$$

$$K(s,t) = -\left(\psi^{(0)}(-s) - \psi^{(0)}(-t)\right)^2 + \left(\psi^{(1)}(-s) + \psi^{(1)}(-t)\right) - \pi^2$$

$$A(s,t) = \frac{1}{20}\left[\frac{s^2t^2\left(-15 - \frac{12}{4+u} + \frac{72}{5+u}\right)}{(1+u)(2+u)(3+u)} + \frac{st\left(25 - \frac{6}{3+u} - \frac{60}{4+u}\right)}{(1+u)(2+u)} - \frac{4\left(3 - \frac{2}{2+u} - \frac{1}{3+u}\right)}{1+u}\right]$$

$$B(s,t) = \frac{(s-t)}{20}\left[\frac{st\left(-15 - \frac{12}{4+u} + \frac{72}{5+u}\right)}{(1+u)(2+u)} + \frac{10}{4+u} + \frac{8}{2+u} + \frac{2}{1+u} - 10\right]$$

$$C(s,t) = \frac{1}{20}\left[\frac{st\left(-15 - \frac{12}{4+u} + \frac{72}{5+u}\right)}{1+u} + \frac{6}{5+u} - \frac{20}{4+u} + \frac{2}{1+u}\right] - 1 + B_0,$$

$$K(s,u), \psi(-s) \wedge \frac{4}{a} \vee \frac{1}{(s+1)(t+1)(u+1)}$$

$$K(s,t), \psi^{(0)}(-s) - \psi^{(0)}(-t), 1$$

$$\psi^{(0)}(-s) - \psi^{(0)}(-t)$$

$$\mathcal{M}^{4l}(s,t) = \sum_{ij \geq 0, |i|+|j| \leq 2} \frac{p_{ij}(s,t)}{(s+i)+(t+j)+1} K(s+i, t+j)$$

	$K(s+2,t)$	$K(s+1,t)$	$K(s-1,t)$	$K(s-2,t)$	$K(s,t)$	$K(s-1,t-1)$	$K(s+1,t+1)$
$K(s,t)$	1	1	1	1	1	1	1
$\Psi_1(s,t)$	$-\frac{2(3+2s)}{(s+1)(s+2)}$	$-\frac{2}{1+s}$	$\frac{2}{s}$	$\frac{2(-1+2s)}{(-1+s)s}$	0	$-\frac{2(s-t)}{st}$	$\frac{2(s-t)}{(s+1)(t+1)}$
1	$-\frac{2}{(s+1)(s+2)}$	0	$-\frac{2}{s^2}$	$-\frac{2(1-3s+3s^2)}{(-1+s)^2s^2}$	0	$-\frac{2(s^2-st+t^2)}{s^2t^2}$	$\frac{2}{(s+1)(t+1)}$

$$\Psi_4(s,t) = \psi^{(0)}(-s) - \psi^{(0)}(-t)$$

$$\sum_{i,j} r_{ij}(s,t) K(s+i, t+j)$$

$$\mathcal{P}_{[4^4][4^4]44} = 16a^2(a+2)g_{12}^4 g_{34}^2$$

$$\mathcal{G}_{[4^4][4^4]44} = 1 + 2U^2\sigma^2 + \frac{2U^2\tau^2}{V^2} + \frac{8}{a}\left[U\sigma + \frac{U\tau}{V}\right] + \frac{24}{a}\frac{U^2\sigma\tau}{V} + \frac{4}{a}\left[U^2\sigma^2 + \frac{U^2\tau^2}{V^2}\right]$$

$$\mathcal{H}_{[4^4][4^4]44} = \mathcal{H}^{c}_{[4^4][4^4]44} + \frac{2a}{a+2}\mathcal{H}_{4444}$$

$$\mathcal{H}_{[4^4][4^4]44}\big|_{\frac{1}{a}} = +2\mathcal{H}_{4444}|_{\frac{1}{a}}$$

$$\mathcal{H}_{[4^4][4^4]44}\big|_{\frac{1}{a^2}} = \mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^2}} - 4\mathcal{H}_{4444}|_{\frac{1}{a}} + 2\mathcal{H}_{4444}|_{\frac{1}{a^2}}.$$

$$\mathcal{H}^{c}_{[4^n][4^n]44}\big|_{\frac{1}{a^2}} = \frac{U^2}{V^2}\Psi_B^{(4)}$$

$$\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^2}} = \frac{U^2}{V^2}\Big[R_{11}^{(1)}\mathcal{P}_2 + R_{10}^{(2)}[z\partial_z - \bar{z}\partial_{\bar{z}}]\mathcal{P}_2 + R_{11}^{(3)}\mathcal{P}_1\log V + R_{10}^{(4)}\log^2 V + R_{10}^{(5)}\log V\log U$$
$$+R_{11}^{(6)}\mathcal{P}_1 + R_{10}^{(7)}\log V + R_{10}^{(8)}\log U + R_8^{(9)}\Big]$$

$$\mathcal{M}^{c}_{[4^4][4^4]44}(s,t) = \big[K(u,t)A(u,t) + \big(\psi^{(0)}(-u) - \psi^{(0)}(-t)\big)B(u,t) + C(u,t)\big]$$

$$A(u,t) = \frac{32}{(1+s)(2+s)(3+s)}\left[-16 + 3ut - 3s + \frac{18ut}{4+s} - \frac{18u^2t^2}{(4+s)(5+s)}\right]$$

$$B(u,t) = \frac{32(u-t)}{(1+s)(2+s)}\left[-\frac{1}{(1+u)(1+t)} - \frac{18ut - 18(5+s)}{(4+s)(5+s)}\right]$$

$$C(u,t) = \frac{32}{(1+s)}\left[\frac{3}{2} - \frac{\frac{1}{2}}{(1+u)(1+t)} + \frac{6 - 18ut + 12(5+s)}{(4+s)(5+s)}\right]$$

$$\mathcal{H}_{[4^4][4^4]44}(U,V) = \frac{1}{V^2}\mathcal{H}_{[4^4][4^4]44}\left(\frac{U}{V},\frac{1}{V}\right)$$

$$\mathcal{M}_{[4^4][4^4]44}(s,t) = \mathcal{M}_{[4^4][4^4]44}(s,u)$$

$$\mathcal{M}^{c}_{[4^4][4^4]44}\otimes K(s,t)\otimes K(s,u)\int_{\mathbb{R}^4}^{\bullet}\psi^{(0)}(-t) - \psi^{(0)}(-u)\oint\log^2(V)\oint\log^2(U)\oint\langle\mathcal{O}_2^2\mathcal{O}_2^2\mathcal{O}_2\mathcal{O}_2\rangle$$

$$\mathcal{H}_{4[4^4]4[4^4]4} = \mathcal{H}_{[4^4][4^4]44}\left(\frac{1}{U},\frac{V}{U}\right);\ \mathcal{M}_{4[4^4]4[4^4]4}(s,t) = \mathcal{M}_{[4^4][4^4]44}(u,t)$$

$$\mathcal{M}^{c}_{4[4^4]4[4^4]}(s,t) = \big[K(s,t)A(s,t) + \big(\psi^{(0)}(-s) - \psi^{(0)}(-t)\big)B(s,t) + C(s,t)\big]$$

$$A(s,t)=\frac{32}{(1+u)(2+u)(3+u)}\left[-16+3st-3u+\frac{18st}{4+u}-\frac{18s^2t^2}{(4+u)(5+u)}\right]$$
$$B(s,t)=\frac{32(s-t)}{(1+u)(2+u)}\left[-\frac{1}{(1+s)(1+t)}-\frac{18st-18(5+u)}{(4+u)(5+u)}\right]$$
$$C(s,t)=\frac{32}{(1+u)}\left[\frac{3}{2}-\frac{\frac{1}{2}}{(1+s)(1+t)}+\frac{6-18st+12(5+u)}{(4+u)(5+u)}\right].$$

$$\lim_{t\gg 4}A^{c}_{4[4^4]4[4^4]}(s,t)\sim\frac{(1+s)}{t^2}\;;\;\lim_{t\gg 4}A^{1l}_{4444}(s,t)\sim\frac{(12+25s+15s^2)}{t}$$

$$\sum_{ij\geq 0,|i|+|j|\leq 2}\frac{P_{ij}(s,t)}{(s+i)+(t+j)+1}K(s+i,t+j)$$

$$\mathcal{H}^{t}_{4[4^4]4[4^4]4}\equiv\mathcal{H}_{4[4^4]4[4^4]4}\Big|_{\frac{1}{a^2}}-2\mathcal{H}_{4444}|_{\frac{1}{a^2}}$$

$$\mathcal{H}^{t}_{4[4^4]4[4^4]4}=\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^2}}-4\mathcal{H}_{4444}|_{\frac{1}{a}}$$
$$=\mathcal{H}^{c}_{4[4^4]4[4^4]}\Big|_{\frac{1}{a^2}}+16U^2\bar{D}_{4444}(U,V)$$

$$\mathcal{M}^{t}_{4[4^4]4[4^4]}=\mathcal{M}^{c}_{4[4^4]4[4^4]}-\frac{16}{(s+1)(t+1)(u+1)}=\mathcal{M}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}}$$

$$\mathcal{M}^{t}_{4[4^4]4[4^4]4}=\sum_{ij\geq 0,|i|+|j|\leq 2}\frac{p_{ij}(s,t)}{(s+i)+(t+j)+1}K(s+i,t+j)$$

$$\mathcal{P}_{[4^4][4^4]44}=96a^2(a+2)(a+4)g_{12}^6g_{34}^2$$
$$\mathcal{G}_{[4^4][4^4]44}=1+3\left[U^2\sigma^2+\frac{U^2\tau^2}{V^2}\right]+\frac{12}{a}\left[U\sigma+\frac{U\tau}{V}\right]+\frac{60}{a}\frac{U^2\sigma\tau}{V}+\frac{12}{a}\left[U^2\sigma^2+\frac{U^2\tau^2}{V^2}\right]$$
$$\mathcal{H}_{[4^4][4^4]44}=\mathcal{H}^{c}_{[4^4][4^4]44}+\frac{3a}{a+4}\mathcal{H}_{4444}+\frac{3a}{a+4}\mathcal{H}^{c}_{[4^4][4^4]44}$$

$$\mathcal{H}_{[4^4][4^4]44}\Big|_{\frac{1}{a}}=3\mathcal{H}_{4444}|_{\frac{1}{a}}$$
$$\mathcal{H}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}}=3\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}}-12\mathcal{H}_{4444}|_{\frac{1}{a}}+3\mathcal{H}_{4444}|_{\frac{1^2}{a}}$$

$$\mathcal{H}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}}=\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}}-12\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}}+3\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}}+48\mathcal{H}_{4444}|_{\frac{1}{a}}-12\mathcal{H}_{4444}|_{\frac{1}{a^4}}+3\mathcal{H}_{4444}|_{\frac{1}{a^4}}$$

$$\mathcal{H}^{c}_{[4^n][4^n]44}\Big|_{\frac{1}{a^4}}=\frac{U^2}{V^2}\Psi_B^{(3)}$$

$$\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} = \frac{U^2}{V^2}[R^{(1)}_{15}\mathcal{P}_3(z,\bar{z}) + R^{(2)}_{14}\partial_{-}\mathcal{P}_3(z,\bar{z}) + R^{(3)}_{15}\partial_{+}\mathcal{P}_3(z,\bar{z}) + R^{(4)}_{14}\partial_{+}\partial_{-}\mathcal{P}_3(z,\bar{z})$$
$$+R^{(5)}_{15}\mathcal{P}_2(z,\bar{z}) + R^{(6)}_{15}\mathcal{P}_1(z,\bar{z})\log^2 V$$
$$+R^{(7)}_{14}\partial_{-}\mathcal{P}_2(z,\bar{z}) + R^{(8)}_{14}\log U\log^2 V + R^{(9)}_{14}\log^3 V + R^{(10)}_{15}\mathcal{P}_1(z,\bar{z})\log V$$
$$+R^{(11)}_{14}\log^2 V + R^{(12)}_{14}\log V\log U + R^{(13)}_{15}\mathcal{P}_1(z,\bar{z}) + R^{(14)}_{14}\log V + R^{(15)}_{14}\log U + R^{(16)}_{12}]$$

$$\mathcal{M}^{c}_{[4^4][4^4]44}(s,t) = [K_4(u,t)a(u,t) + K_3(u,t)b(u,t) + \tilde{K}_2(u,t)c(u,t)$$
$$+K_2(u,t)A(u,t) + K_1(u,t)B(u,t) + C(u,t)]$$

$$K_4(s,t) = \Psi_1^4 + 3\Psi_2^2 - 2\Psi_1\Psi_3 - 2\pi^2\Psi_2 - 2\pi^2K_2 - \pi^4$$
$$K_3(s,t) = \Psi_1^3(u,t) - \frac{1}{2}\Psi_3(u,t) + \pi^2\Psi_1$$
$$\tilde{K}_2(s,t) = \Psi_2 - \frac{2}{3}\pi^2\ ;\ K_2(s,t) = -\Psi_1^2 + \Psi_2 - \pi^2\ ;\ K_1(s,t) = \Psi_1$$

$$\Psi_i(s,t) \equiv \psi^{(i-1)}(-s) + (-1)^i\psi^{(i-1)}(-t)$$

$$a(s,t) = \frac{256}{(u+1)(u+2)(u+3)}[$$
$$14 - 15st + \frac{9}{2}u + \frac{15st(4+5st)}{4(u+4)_1} - \frac{3s^2t^2(58+15st)}{4(u+4)_2} + \frac{45s^2t^2(1+2st)}{2(u+4)_3} - \frac{135s^2t^2(-2+st)^2}{2(u+4)_4}]$$

$$b(s,t) = \frac{384(s-t)}{(u+1)(u+2)(u+3)}[$$
$$46 + \frac{2(10-77st)}{(u+4)_1} - \frac{st(116-195st)}{(u+4)_2} + \frac{15st(4+21st-9s^2t^2))}{(u+4)_3} - \frac{360st(2-st)(1-st)}{(u+4)_4}]$$

$$\langle\mathcal{O}_2^2\mathcal{O}_2^2\mathcal{O}_2\mathcal{O}_2\rangle\ ;\ \langle\mathcal{O}_2\mathcal{O}_2^2\mathcal{O}_2\mathcal{O}_2^2\rangle$$

$$\mathcal{H}^{c}|_{\log^0(U)}$$

$$\mathcal{H}^{c}|_{\log^n(U)}$$

$$\langle\mathcal{O}_2^2(\vec{x}_1,\vec{y}_1)\mathcal{O}_2^2(\vec{x}_2,\vec{y}_2)\mathcal{O}_2(\vec{x}_3,\vec{y}_3)\mathcal{O}_2(\vec{x}_4,\vec{y}_4)\rangle = \mathcal{P}_{[4^4][4^4]44}\sum_{\underline{R}} A_{\underline{R}}B_{\underline{R}}$$

$$A_{\underline{R}} = \sum_{O,O'\in\underline{R}} C_{[4^4][4^4]O}\mathbb{G}^{O,O'}C_{44O'}$$

$$\mathcal{O}_2\mathcal{O}_2 \sim 1 \oplus \underbrace{\mathcal{O}_2 + \mathcal{K}_2 + \cdots}_{\tau=2} \oplus \underbrace{\mathcal{O}_2^2 + \cdots}_{\tau\geq 4}$$

$$C_{\mathcal{O}_2\mathcal{O}_2;\mathcal{K}_2} \sim \frac{1}{N}$$

$$C_{\mathcal{O}_2\mathcal{O}_2;\mathcal{O}_2\partial^l\mathcal{O}_2} \sim 1$$

$$\mathcal{P}_{[4^4][4^4]44} = 16a^2(a+2)g_{12}^4 g_{34}^2$$

$$\mathcal{G}_{[4^4][4^4]44} = 1 + 2U^2\sigma^2 + \frac{2U^2\tau^2}{V^2} + \frac{8}{a}\left[U\sigma + \frac{U\tau}{V}\right] + \frac{24}{a}\frac{U^2\sigma\tau}{V} + \frac{4}{a}\left[U^2\sigma^2 + \frac{U^2\tau^2}{V^2}\right]$$

$$\langle \mathcal{O}_2^2\mathcal{O}_2^2\mathcal{O}_2\mathcal{O}_2\rangle\big|_{\substack{\tau=2\\ \text{OPE}}} = \mathcal{P}_{[4^4][4^4]44} \times \frac{8}{a}\left[U\sigma + \frac{U\tau}{V}\right]$$

$$A^{\text{free}}_{\mathcal{K},4,l,[0000]} = \frac{8}{a} \times \frac{2(l+2)!^2}{(2l+4)!}\frac{1+(-1)^l}{2}$$

$$A_{\mathcal{L},4,l,[0000]} = A^{\text{free}}_{\mathcal{K},4,l,[000]} + \mathcal{H}_{[4^4][4^4]44}\big|_{\tau=4,l,[0000]} = 0$$

$$\mathcal{H}_{[4^4][4^4]44}\big|_{\frac{1}{a},\tau=4} = -\frac{8}{a} \times \frac{2(l+2)!^2}{(2l+4)!}\frac{1+(-1)^l}{2}$$
$$\mathcal{H}_{[4^4][4^4]44}\big|_{\frac{1}{a^n},\tau=4} = 0 \qquad n \geq 4$$

$$\mathcal{H}_{[4^4][4^4]44}\big|_{\frac{1}{a}} = -8U\left[\frac{(1+V)}{(-1+V)^2V} - \frac{2\log(V)}{(-1+V)^3}\right] + U^2(\dots)$$

$$\mathcal{H}_{4444}\big|_{\frac{1}{a}} = -4U^2\bar{D}_{4444} = -4U\left[\frac{(1+V)}{(-1+V)^2V} - \frac{2\log(V)}{(-1+V)^3}\right] + U^2(\dots)$$

$$\mathcal{H}_{[4^4][4^4]44}\big|_{\frac{1}{a}} = 2\mathcal{H}_{4444}\big|_{\frac{1}{a}}$$

$$\mathcal{H}^t_{[4^4][4^4]44}\Big|_{\log^0(U)} = \mathcal{H}^c_{[4^4][4^4]44} + 16U^2\bar{D}_{4444}|_{\log^0(U)} = U^2(\dots)$$

$$\langle \mathcal{O}_2(\vec{x}_1,\vec{y}_1)\mathcal{O}_2^2(\vec{x}_2,\vec{y}_2)\mathcal{O}_2(\vec{x}_3,\vec{y}_3)\mathcal{O}_2^2(\vec{x}_4,\vec{y}_4)\rangle = \mathcal{P}_{4[4^4]4[4^4]}\sum_{\underline{R}} A_{\underline{R}}B_{\underline{R}}$$

$$A_{\underline{R}} = \sum_{O,O'\in\underline{R}} C_{4[4^4]O}\mathbb{G}^{O,O'}C_{4[4^4]O'}$$

$$\mathcal{O}_2\mathcal{O}_2^2 \sim \mathcal{O}_2 \oplus \underbrace{\mathcal{O}_2^2 + \mathcal{K}_4 + \cdots}_{\tau=4} \oplus \underbrace{\mathcal{O}_2^3 + \cdots}_{\tau\geq 6}$$

$$C_{\mathcal{O}_2\mathcal{O}_2^2;\mathcal{O}_2\partial^l\mathcal{O}_2} \sim \frac{1}{N}$$

$$\mathcal{O}_2\partial^l\mathcal{O}_2\big|_{[0,2,0],l=0,2,4,\dots}\ ;\ \mathcal{O}_2\partial^l\mathcal{O}_2\big|_{[1,0,1],l=1,3,5,\dots}$$

$$G = \begin{pmatrix} \langle \mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\rangle & \langle \mathcal{O}_2\mathcal{O}_2\mathcal{O}_2\mathcal{O}_2^2\rangle \\ \langle \mathcal{O}_2\mathcal{O}_2^2\mathcal{O}_2\mathcal{O}_2\rangle & \langle \mathcal{O}_2\mathcal{O}_2^2\mathcal{O}_2\mathcal{O}_2^2\rangle \end{pmatrix},$$

$$\mathcal{S}^{\vec{p}}_{4,[0404]} \equiv A^{\vec{p}}_{\mathcal{S},\tau=4,l,[aba]}$$

$$\mathcal{S}_{4,[0404]}^{4[4^4]4[4^4]} = \frac{\left(\mathcal{S}_{4,[0404]}^{[4^4]4;44}\right)^4}{\mathcal{S}_{4,[0404]}^{44;44}}$$

$$\mathcal{S}_{4,[0404]}^{[4^4]4[4^4]4} = \frac{32}{a}\frac{\left(1+\frac{2}{a}\right)}{(l+3)(l+4)+\frac{4}{a}}\frac{(l+4)!^2}{(2l+6)!}\frac{1+(-1)^l}{2}$$

$$A_{\mathcal{K},4,l,[0404]}^{\text{free}} = -\mathcal{S}_{4,[0404]} + A_{4,l,[0404]}^{\text{free}},$$

$$A_{4,l,[0404]}^{\text{free}} = \frac{1}{a}\begin{cases} +4(l+1)(l+6)\dfrac{(l+2)!\,(l+4)!}{(2l+6)!} & \text{spin odd} \\ +4(18+7l+l^2)\dfrac{(l+2)!\,(l+4)!}{(2l+6)!} & \text{spin even} \end{cases}$$

$$A_{\mathcal{L},4,l,[0404]} = A_{\mathcal{K},4,l,[0404]}^{\text{free}} + \mathcal{H}_{4[4^4]4[4^4]}\big|_{\tau=4,l,[0404]} = 0$$

$$A_{\mathcal{L},4,l,[0404]} = \begin{cases} +\dfrac{4}{a}(l+1)(l+6)\dfrac{(l+2)!\,(l+4)!}{(2l+6)!} & \text{spin odd} \\ +\dfrac{4(l+2)(l+5)}{a}\dfrac{(l+3)(l+4)-\frac{12}{a}}{(l+3)(l+4)+\frac{4}{a}}\dfrac{(l+2)!\,(l+4)!}{(2l+6)!} & \text{spin even} \end{cases}$$

$$\mathcal{H}_{4[4^4]4[4^4]4}\big|_{\frac{1}{a},\tau=4} = \begin{cases} -4(l+1)(l+6)\dfrac{(l+2)!\,(l+4)!}{(2l+6)!}\dfrac{1-(-1)^l}{2} \\ -4(l+2)(l+5)\dfrac{(l+2)!\,(l+4)!}{(2l+6)!}\dfrac{1+(-1)^l}{2} \end{cases}$$

$$\mathcal{H}_{4[4^4]4[4^4]}\big|_{\frac{1}{a^4},\tau=4} = \begin{cases} 0 \\ +64(l+2)(l+5)\dfrac{(l+2)!^{\,2}}{(2l+6)!}\dfrac{1+(-1)^l}{2}. \end{cases}$$

$$\mathcal{H}_{44^444^4} = -\frac{8}{a}U\left[\frac{(1+V)}{(1-V)^2V}+\frac{2\log(V)}{(1-V)^3}\right]+U^2(\dots)$$
$$+\frac{32}{a^2}U\left[\frac{1}{(1-V)^2V}-\frac{\log(V)+\log^2(V)+2\mathrm{Li}_2(1-V)}{(1-V)^3}\right]+U^2(\dots)$$

$$C_{[4^n][4^n];\mathcal{D}} = C_{[4^n][4^n];\mathcal{D}}^{(0)} + \frac{1}{a}C_{[4^n][4^n];\mathcal{D}}^{(4)} + \cdots \infty$$

$$\Delta_{\mathcal{D}} = \tau + l + \frac{2}{a}\eta_{\mathcal{D}}^{(4)} + \cdots$$

$$\hat{\mathcal{C}}_{[4^4][4^4]44}\big|_{\frac{\log n(U)}{a^n}} = \frac{1}{n!}\sum_{\tau,l}\left(\sum_{\mathcal{D}} C_{[4^4][4^4];\mathcal{D}}^{(0)}\eta_{\mathcal{D}}^n C_{44;\mathcal{D}}^{(4)}\right)B_{\tau,l,[0000]}^{\mathcal{L}} = 2\hat{\mathcal{C}}_{4444}\big|_{\frac{\log^n(U)}{a^n}}$$

$$\mathcal{H}_{[4^4][4^4]44} - \mathcal{H}_{4444}$$

$$\mathcal{H}^{t}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} \equiv \mathcal{H}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} - 2\mathcal{H}_{4444}|_{\frac{1}{a^4}}.$$

$$\mathcal{H}^{1l}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} \equiv \mathcal{H}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} - 2\mathcal{H}_{4444}|_{\frac{1}{a^4}}.$$

$$\hat{\mathcal{C}}_{[4^4][4^4]44}\big|_{\frac{1}{a}} - 2\hat{\mathcal{C}}_{4444}\big|_{\frac{1}{a}} = \sum_{l}\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_4} - 2C^{(4)}_{44;\mathcal{D}_4}\right]C^{(n)}_{44;\mathcal{D}_4}B^{\mathcal{L}}_{4,l,[0000]} + (\text{ twist } \geq 6)$$

$$\hat{\mathcal{C}}_{[4^4][4^4]44}\big|_{\frac{\log(U)}{a^2}} - 2\hat{\mathcal{C}}_{2222}\big|_{\frac{\log(U)}{a^2}} = \sum_{l}\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_4} - 2C^{(4)}_{44;\mathcal{D}_4}\right]\eta_{\mathcal{D}_4}C^{(n)}_{44;\mathcal{D}_4}B^{\mathcal{L}}_{4,l,[0000]} + (\text{ twist } \geq 6)$$

$$\hat{\mathcal{C}}_{[4^4][4^4]44}\big|_{\frac{1}{a}} - 2\hat{\mathcal{C}}_{4444}\big|_{\frac{1}{a}} = \frac{4}{a}U^2\sigma^2 + \frac{4}{a}\frac{U^2\tau^2}{V^2} + \frac{16}{a}\frac{U^2\sigma\tau}{V}$$

$$\frac{4}{a}U^2\sigma^2 + \frac{4}{a}\frac{U^2\tau^2}{V^2} + \frac{16}{a}\frac{U^2\sigma\tau}{V} = \sum_{l}\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_4} - 2C^{(4)}_{44;\mathcal{D}_4}\right]C^{(n)}_{44;\mathcal{D}_4}B^{\mathcal{L}}_{4,l,[0000]} + (\text{ twist } \geq 6)$$

$$\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_4} - 2C^{(4)}_{44;\mathcal{D}_4}\right]C^{(n)}_{44;\mathcal{D}_4} = \frac{4}{3}(l+2)(l+5)\frac{(l+3)!^2}{(2l+6)!}\frac{1+(-1)^l}{2}$$

$$\hat{\mathcal{C}}_{[4^4][4^4]44}\big|_{\frac{\log(U)}{a^4}} - 2\hat{\mathcal{C}}_{2222}\big|_{\frac{\log(U)}{a^4}} = \mathcal{I}\times\left[\mathcal{H}_{[4^4][4^4]44} - 2\mathcal{H}_{4444}\right]_{\frac{\log(U)}{a^4}}$$

$$\mathcal{H}^{t}_{[4^4][4^4]44}\Big|_{\frac{\log(U)}{a^4}} = \sum_{l}\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_4} - 22C^{(4)}_{44;\mathcal{D}_4}\right]\eta_{\mathcal{D}_4}C^{(n)}_{44;\mathcal{D}_4}H_{4,l,[0000]} + (\text{ twist } \geq 6)$$

$$\eta_{\mathcal{D}_4} = -\frac{48}{(l+1)(l+6)}\frac{1+(-1)^l}{2}$$

$$\mathcal{H}^{t}_{[4^4][4^4]44}\Big|_{\frac{\log(U)}{a^4}} = \sum_{l}\left[-64\frac{(l+2)(l+5)}{(l+1)(l+6)}\frac{(l+3)!^2}{(2l+6)!}\frac{1+(-1)^l}{2}\right]H_{4,l,[0000]} + (\text{ twist } \geq 6)$$

$$\mathcal{H}^{t}_{[4^4][4^4]44}\Big|_{\frac{\log(U)}{a^4}} =$$

$$-32U^2\left[\frac{(1+10V+V^2)}{(1-V)^4V} + \frac{12(1+V)\log(V)}{(1-V)^5} + \frac{2(1+4V+V^2)\log^2(V)}{(1-V)^6} + U^3(\ldots)\right]$$

$$\mathcal{H}^{t}_{[4^4][4^4]44} = \mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} - 4\mathcal{H}_{4444}|_{\frac{1}{a}}$$

$$\mathcal{H}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} = \mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} - 12\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} + 3\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} + 48\mathcal{H}_{4444}|_{\frac{1}{a}} - 12\mathcal{H}_{4444}|_{\frac{1}{a^4}} + 3\mathcal{H}_{4444}|_{\frac{1}{a^4}}$$

$$\mathcal{H}^{t}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} = \mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} - 12\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} + 24\mathcal{H}_{4444}|_{\frac{1}{a}}$$

$$\left[\mathcal{H}^{t}_{[4^4][4^4]44}\right]_{\frac{\log^4(U)}{a^4}} = 0$$

$$\left[\mathcal{H}^{t}_{[4^4][4^4]44}\right]_{\frac{\log^4(U)}{a^4},\ \text{twist 4}} = 0$$

$$\left[\mathcal{H}^{t}_{[4^4][4^4]44}\right]_{\frac{\log^n(U)}{a^n},\ \text{twist 4}} = 0\ ;\ n = 1,2,3,4\ldots$$

$$\mathcal{H}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} - 3\mathcal{H}_{4444}\Big|_{\frac{1}{a^4}} - 3\mathcal{H}^{1l}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} = \left[\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} - 12\mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} + 24\mathcal{H}_{4444}\Big|_{\frac{1}{a^4}}\right],$$

$$\mathcal{H}^{1l}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} = \mathcal{H}^{c}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} + 8\mathcal{H}_{4444}\Big|_{\frac{1}{a^4}} - 4\mathcal{H}_{4}\Big|_{\frac{1}{a^4}}.$$

$$C^{(n)}_{[4^\eta][4^\eta];\mathcal{D}_4} = 3C^{(n)}_{[4^\eta][4^\eta];\mathcal{D}_4} - 3C^{(n)}_{44;\mathcal{D}_4}\ ;\ n = 1,2,3,4\ldots$$

$$\hat{\mathcal{C}}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}}\Big|_{\frac{1}{a^4}} - 3\hat{\mathcal{C}}_{4444}\Big|_{\frac{1}{a^4}} = \frac{48}{a}\frac{U^2\sigma\tau}{V} + \frac{12}{a}\left[U^2\sigma^2 + \frac{U^2\tau^2}{V^2}\right] = 3\left[\hat{\mathcal{C}}_{[4^4][4^4]44}\Big|_{\frac{1}{a^4}} - 2\hat{\mathcal{C}}_{4444}\Big|_{\frac{1}{a^4}}\right]\ldots 4$$

$$\left[\mathcal{H}^{t}_{[4^4][4^4]44}\right]_{\frac{\log^4(U)}{a^4}} = \frac{1}{2}\sum_{l}\left[\left(C^{(4)}_{[4^\eta][4^\eta];\mathcal{D}_4} - 3C^{(4)}_{[4^\eta][4^\eta];\mathcal{D}_4} + 3C^{(4)}_{22;\mathcal{D}_4}\right)\left(\eta^{(4)}_{\mathcal{D}_4}\right)^2 C^{(0)}_{22\mathcal{D}_4}\right]H_{4,l,[0000]} + (\text{ twist } \geq 6)$$

$$\left[\mathcal{H}^{t}_{[4^4][4^4]44}\right]^{t}_{\frac{\log(U)}{a^4}}\sum_{l}\left[\left(C^{(4)}_{[4^\eta][4^\eta];\mathcal{D}_4} - 3C^{(4)}_{[4^\eta][4^\eta];\mathcal{D}_4} + 3C^{(4)}_{44;\mathcal{D}_4}\right)\left(\eta^{(4)}_{\mathcal{D}_4}C^{(4)}_{44;\mathcal{D}_4} + \eta^{(4)}_{\mathcal{D}_4}C^{(4)}_{44;\mathcal{D}_4}\right)\right.$$
$$\left.+\left(C^{(4)}_{[4^\eta][4^\eta];\mathcal{D}_4} - 3C^{(4)}_{[4^\eta][4^\eta];\mathcal{D}_4} + 3C^{(4)}_{44;\mathcal{D}_4}\right)\eta^{(4)}_{\mathcal{D}_4}C^{(4)}_{44;\mathcal{D}_4}\right]H_{4,l,[0000]} + \cdots + (\text{ twist } \geq 6)$$

$$C_{[4^\eta][4^\eta];\mathcal{T}} = \frac{1}{\sqrt{a}}\left[C^{\left(\frac{1}{2}\right)}_{[4^\eta][4^\eta];\mathcal{T}} + \cdots\right]\ ;\ \Delta = \tau + l + \frac{2}{a}\eta^{(4)}_{\mathcal{T}} + \cdots$$

$$\left[\mathcal{H}^{t}_{[4^4][4^4]44}\right]_{\frac{\log^4(U)}{a^4}} = (\text{ twist } 4) + \frac{1}{2}\sum_{\tau,l}\left[\sum_{\mathcal{D}}\left(C^{(4)}_{[4^\eta][4^\eta];\mathcal{D}_4} - 3C^{(4)}_{[4^\eta][4^\eta];\mathcal{D}_4} + 3C^{(4)}_{44;\mathcal{D}_4}\right)\left(\eta^{(4)}_{\mathcal{D}_4}\right)^2 C^{(4)}_{44;\mathcal{D}_4} + \right.$$
$$\left.\sum_{\mathcal{T}}\left(C^{\left(\frac{1}{2}\right)}_{[4^\eta][4^\eta];\mathcal{T}_4} - 3C^{\left(\frac{1}{2}\right)}_{[4^\eta][4^\eta];\mathcal{T}_4} + 3C^{\left(\frac{1}{2}\right)}_{22;\mathcal{T}_4}\right)\left(\eta^{(1)}_{\mathcal{T}_4}\right)^2 C^{\left(\frac{1}{2}\right)}_{44;\mathcal{T}_4}\right]H_{\tau,l,[0000]}$$

$$C^{(4)}_{[4^\eta][4^\eta];\mathcal{D}_4} = 3C^{(4)}_{[4^\eta][4^\eta];\mathcal{D}_4} - 3C^{(4)}_{44;\mathcal{D}}\ ;\ C^{\left(\frac{1}{2}\right)}_{[4^\eta][4^\eta];\mathcal{T}} = 3C^{\left(\frac{1}{2}\right)}_{[4^\eta][4^\eta];\mathcal{T}} - 3C^{\left(\frac{1}{2}\right)}_{44;\mathcal{T}}$$

$$\langle\mathcal{O}_2^2\mathcal{O}_2^2\mathcal{D}_{\tau,l}\rangle = \langle\mathcal{O}_2^2\mathcal{O}_2^2\mathcal{D}_{\tau,l}\rangle_c + 4\langle\mathcal{O}_2\mathcal{O}_2\rangle\langle\mathcal{O}_2\mathcal{O}_2\mathcal{D}_{\tau,l}\rangle$$

$$\frac{\langle\mathcal{O}_2^2\mathcal{O}_2^2\mathcal{D}_{\tau,l}\rangle_c}{\langle\mathcal{O}_2^2\mathcal{O}_2^2\rangle} \sim \frac{1}{a}$$

$$\langle\mathcal{O}_2^3\mathcal{O}_2^3\mathcal{D}_{\tau,l}\rangle = \langle\mathcal{O}_2^3\mathcal{O}_2^3\mathcal{D}_{\tau,l}\rangle_c + 9\langle\mathcal{O}_2\mathcal{O}_2\rangle\langle\mathcal{O}_2^2\mathcal{O}_2^2\mathcal{D}_{\tau,l}\rangle + 9\left(\langle\mathcal{O}_2^2\mathcal{O}_2^2\rangle - 4\langle\mathcal{O}_2\mathcal{O}_2\rangle^2\right)\langle\mathcal{O}_2\mathcal{O}_2\mathcal{D}_{\tau,l}\rangle$$

$$\frac{\langle\mathcal{O}_2^3\mathcal{O}_2^3\mathcal{D}_{\tau,l}\rangle_c}{\langle\mathcal{O}_2^3\mathcal{O}_2^3\rangle} \sim \frac{1}{a^2}$$

$$\langle \mathcal{O}_2^3 \mathcal{O}_2^3 \mathcal{T}_{\tau,l} \rangle = \langle \mathcal{O}_2^3 \mathcal{O}_2^3 \mathcal{T}_{\tau,l} \rangle_c + 9\langle \mathcal{O}_2 \mathcal{O}_2 \rangle \langle \mathcal{O}_2^2 \mathcal{O}_2^2 \mathcal{T}_{\tau,l} \rangle + 9\left(\langle \mathcal{O}_2^2 \mathcal{O}_2^2 \rangle - 4\langle \mathcal{O}_2 \mathcal{O}_2 \rangle^2\right) \langle \mathcal{O}_2 \mathcal{O}_2 \mathcal{T}_{\tau,l} \rangle$$

$$\frac{\langle \mathcal{O}_2^3 \mathcal{O}_2^3 \mathcal{T}_{\tau,l} \rangle_c}{\langle \mathcal{O}_2^3 \mathcal{O}_2^3 \rangle} \sim \frac{1}{a^{\frac{3}{2}}}$$

$$C^{(4)}_{[4^n][4^n];\mathcal{D}} = nC^{(4)}_{44;\mathcal{D}} \; n = 2,3,4 \ldots$$

$$C^{(4)}_{[4^n][4^n];\mathcal{D}} = 3C^{(4)}_{[4^n][4^n];\mathcal{D}} - 3C^{(4)}_{44;\mathcal{D}} \; ; \; C^{\left(\frac{1}{2}\right)}_{[4^\eta][4^\eta];\mathcal{T}} = 3C^{\left(\frac{1}{2}\right)}_{[4^\eta][4^\eta];\mathcal{T}} - 3C^{\left(\frac{1}{2}\right)}_{44;\mathcal{T}}$$

$$C_{44;\mathcal{T}} \sim \frac{1}{N} \; ; \; C_{[4^\eta][4^\eta];\mathcal{T}} \sim \frac{1}{N}.$$

$$\left(\partial_{\hat{z}}^2 + \frac{\frac{1}{4} - a_1^2}{(\hat{z}-1)^2} - \frac{\frac{1}{2} - a_0^2 - a_1^2 - a_t^2 + a_\infty^2 + u}{\hat{z}(\hat{z}-1)} + \frac{\frac{1}{4} - a_t^2}{(\hat{z}-t)^2} + \frac{u}{\hat{z}(\hat{z}-t)} + \frac{\frac{1}{4} - a_0^2}{\hat{z}^2}\right)\psi(\hat{z}) = 0$$

$$a_0^2 = \frac{l^2}{4}, a_1^2 = a_t^2 = \frac{1}{4}, u = -\frac{(\alpha^2+8)(\omega^2-l^2)}{4(8-\alpha^2)}, a_\infty^2 = \frac{l^2}{4}, t = \frac{\alpha^4}{64}$$

$$F = c_1(a, a_0, a_t, a_1, a_\infty)t + \mathcal{O}(t^2)$$

$$c_1(a, a_0, a_t, a_1, a_\infty) = \frac{(4a^2 - 4a_0^2 + 4a_t^2 - 1)(4a^2 + 4a_1^2 - 4a_\infty^2 - 1)}{8 - 32a^2},$$

$$u = -a^2 + a_t^2 - \frac{1}{4} + a_0^2 + t\partial_t F$$

$$a^2 = \frac{\omega^2}{4} + \frac{\omega^2 - l^2}{16}\left(\alpha^2 + \frac{\alpha^4(l^2 + 3\omega^2 - 4)}{32(\omega^2-1)} + \frac{\alpha^6(l^4 - 2l^2 + \omega^4 - 2\omega^2 + 2)}{128(\omega^2-1)^2}\right) + \mathcal{O}(\alpha^8).$$

$$w_-^{(0)} = \text{HeunG}(t, \hat{q}, \hat{\alpha}, \hat{\beta}, \hat{\gamma}, \hat{\delta}, \hat{z})$$

$$\psi(\hat{z}) \approx \left[1 + \frac{\Gamma(\hat{\epsilon}-1)\Gamma\left(\frac{1+\hat{\gamma}-\hat{\epsilon}}{2} - a\right)\Gamma\left(\frac{1+\hat{\gamma}-\hat{\epsilon}}{2} + a\right)}{\Gamma(1-\hat{\epsilon})\Gamma\left(\frac{-1+\hat{\gamma}+\hat{\epsilon}}{2} - a\right)\Gamma\left(\frac{-1+\hat{\gamma}+\hat{\epsilon}}{2} + a\right)} e^{-\partial_{a_t} F} t^{\hat{\epsilon}-1}(t-\hat{z})^{1-\hat{\epsilon}} + \cdots\right].$$

$$\hat{\epsilon} = 1 - 2a_t, \text{ and } \hat{\gamma} = 1 - 2a_0 = 1 + |l|.$$

$$a = \pm\left(k + \frac{|l| - \hat{\epsilon}}{2}\right), \text{ with } k = 1,2, \ldots.$$

$$-k(k+|l|)\frac{e^{-\partial_{a_t} F}}{\hat{b}+k}\left(1 - \frac{\hat{z}}{t}\right) \sim \left(\frac{l^2}{4} - a^2\right)\frac{e^{-\partial_{a_t} F}}{\hat{b}+k}\frac{8-\alpha^2}{8+\alpha^2}w^2 = \frac{1}{\hat{b}+k}\left(\hat{a}\hat{b}e^{-\partial_{a_t} F}\frac{8-\alpha^2}{8+\alpha^2}\right)w^2.$$

$$\hat{a}\hat{b}e^{-\partial_{a_t} F}\frac{8-\alpha^2}{8+\alpha^2} = \frac{l^2 - \omega^2}{4}$$

$$\hat{b} = \frac{l+\omega}{2} + \frac{\alpha^2(\omega^2-l^2)}{16\omega} + \frac{\alpha^4(-3l^4\omega^2+2l^4+2(l^2-1)\omega^4+\omega^6)}{512\omega^3(\omega^2-1)} + \frac{\alpha^6\big((\omega^2-l^2)(l^4(5\omega^4-5\omega^2+2)-2l^2(\omega^6+\omega^4)+(\omega^4-\omega^2+2)\omega^4)\big)}{4096(\omega^5(\omega^2-1)^2)} + \mathcal{O}(\alpha^8)$$

$$\psi''(\hat{z}) + \left[\frac{(l+2)(-l)}{4\hat{z}^2} + \frac{l(l+2)-2\omega^2}{4(\hat{z}-1)^2} + \frac{(3-2\alpha^2)\omega^2+(2\alpha^2-1)(l(l+2)-4)-8}{4(\hat{z}-1)} - \frac{3}{4\left(\frac{2\alpha^2}{1-2\alpha^2}+\hat{z}\right)^2} - \frac{(2\alpha^2+1)\omega^2-8\alpha^2-(1-2\alpha^2)l(l+2)-4}{8\alpha^2\hat{z}} + \frac{(1-2\alpha^2)(-2\alpha^2\omega^2-8\alpha^2+(2\alpha^2-1)(l+2)l+\omega^2-4)}{8\alpha^2\left(\frac{2\alpha^2}{1-2\alpha^2}+\hat{z}\right)}\right]\psi(\hat{z}) = 0$$

$$a_0^2 = \frac{1}{4}(l+1)^2, a_1^2 = \frac{1}{4}(-l(l+2)+2\omega^2+1), a_t^2 = 1, a_\infty^2 = 1$$

$$u = \frac{1}{4}(2\alpha^2\omega^2+8\alpha^2-(2\alpha^2-1)l(l+2)-\omega^2+4), t = -\frac{2\alpha^2}{1-2\alpha^2}$$

$$\hat{\epsilon} = -1, \hat{\gamma} = 2+|l|$$

$$1-\frac{\hat{z}}{t} = \frac{w^2}{1-2\alpha^2}\big(1+O(w^2)\big)$$

$$\hat{b} = \frac{l+\omega}{2} + \frac{\alpha^2((l+2)^2-\omega^2)(l^2-\omega^2)}{4\omega(\omega^2-1)} + \cdots$$

$$I_0[p,q] \equiv \sum_{l\in\mathbb{Z}}^{\infty}\sum_{k=1}^{\infty} \frac{l^{2p}}{(|l|+2k)^q} e^{-i(|l|+2k)\tau+il\sigma}$$

$$I_1[p,q] \equiv \sum_{l\in\mathbb{Z}}^{\infty}\sum_{k=1}^{\infty} \frac{l^{2p}}{(|l|+2k-1)^q} e^{-i(|l|+2k)\tau+il\sigma};$$

$$I_2[p,q] \equiv \sum_{l\in\mathbb{Z}}^{\infty}\sum_{k=1}^{\infty} \frac{l^{2p}}{(|l|+2k-2)^q} e^{-i(|l|+2k)\tau+il\sigma}\ (2p-q\geq 0);$$

$$I_3[p,q] \equiv \sum_{l\in\mathbb{Z}}^{\infty}\sum_{k=1}^{\infty} \frac{(l^2-1)^2 l^{2p}}{(|l|+2k-3)^q} e^{-i(|l|+2k)\tau+il\sigma}\ (q\leq 2);$$

$$I_4[p,q] \equiv \sum_{l\in\mathbb{Z}}^{\infty}\sum_{k=1}^{\infty} \frac{(l^2-4)^2 l^{2p}}{(|l|+2k-4)^q} e^{-i(|l|+2k)\tau+il\sigma}\ (q\leq 2);$$

$$\tilde{I}_1[p,q] \equiv \sum_{l\in\mathbb{Z}}^{\infty}\sum_{k=1}^{\infty} \frac{l^{2p}}{(|l|+2k+1)^q} e^{-i(|l|+2k)\tau+il\sigma};$$

$$\tilde{I}_2[p,q] \equiv \sum_{l\in\mathbb{Z}}^{\infty}\sum_{k=1}^{\infty} \frac{l^{2p}}{(|l|+2k+2)^q} e^{-i(|l|+2k)\tau+il\sigma}.$$

$$A^1 = -H_1^{-1}dt\,, X_1 = H_1^{-2/3}\,, \cosh\,\varphi_1 = \frac{d}{dr^2}(r^2 H_1) = \frac{1+\frac{1+q_1}{r^2}}{\sqrt{1+\frac{2(1+q_1)}{r^2}+\frac{1}{r^4}}}.$$

$$AdS_{10}^4 = \Delta^{1/2} dS_5^4 + \Delta^{-1/2} G_{\alpha\beta}(dy^\alpha + A^\alpha)\big(dy^\beta + A^\beta\big)$$

$$\left\langle \begin{array}{c} G_{\alpha\beta}dy^\alpha dy^\beta = \big(e^{2\mu}\big(e^{\lambda}\cos^2\,\phi + e^{-\lambda}\cos^2\,\phi\big)\sin^2\,\tilde{\theta} + e^{-2\mu}\cos^2\,\tilde{\theta}\big)d\tilde{\theta}^2 \\ +e^{2\mu}\big(e^{\lambda}\cos^2\,\phi + e^{-\lambda}\sin^2\,\phi\big)\cos^2\,\tilde{\theta}d\phi^2 - e^{2\mu}\sinh\,\lambda\sin\,(2\phi)\sin\,(2\tilde{\theta})d\tilde{\theta}d\phi + e^{-\mu}\sin^2\,\tilde{\theta}d\tilde{\Omega}_3^2 \end{array} \right\rangle^4$$

$$A^\phi \equiv A^1 = -e^{-3\mu}dt$$

$$\langle \Delta = \left(\frac{\det G}{\det G_0}\right)^{-2/3} = e^{-2\mu}\big(e^{\lambda}\cos^2\,\phi + e^{-\lambda}\sin^2\,\phi\big)\cos^2\,\tilde{\theta} + e^{\mu}\sin^2\,\tilde{\theta}\rangle^4$$

$$\lambda = \varphi_1\,, e^{3\mu} = H_1$$

$$AdS_5^4 = \langle \Omega_0^2\left[\frac{d\xi^2}{(1-\xi^2)^2} + \frac{\xi^2}{1-\xi^2}d\Omega_3^2\right] - \frac{\Omega_1^2}{1-\xi^2}dt^2\rangle^4$$

$$\frac{\Omega_1^2}{1-\xi^2} = H_1^{-2/3} f\,, \Omega_0^2\frac{\xi^2}{1-\xi^2} = H_1^{1/3} r^2\,, \frac{\Omega_0^2}{(1-\xi^2)^2} = H_1^{1/3} f^{-1}\big(\partial_\xi r\big)^2$$

$$(\partial_s r)^2 = r^2(1 + r^2 H_1) \text{ with } \tanh\, s = \sqrt{1-\xi^2}$$

$$\square_{10}\,\Phi = 0,$$

$$\square_5\,\Phi_0 = 0.$$

$$\delta + \delta_{\text{Ads}} \equiv -p_t\Delta t - p_\varphi\Delta\varphi = 2|p_t|\int_{r_0}^{\infty}\frac{dr}{f}\sqrt{H_1 - \beta^2\frac{f}{r^2}}$$

$$r_0 = \sqrt{\frac{1}{(1-\beta^2)^2} + (q_1+2)q_1} - q_1 - 1$$

$$\Delta t = 2\int_{r_0}^{\infty} dr\frac{\dot{t}}{\dot{r}}, \Delta\varphi = 2\int_{r_0}^{\infty} dr\frac{\dot{\varphi}}{\dot{r}}$$

$$\delta + \delta_{\text{Ads}} = 2|p_t|\int_0^1 dy\left[\frac{(1-\beta^2)\sqrt{1+2(1+q_1)\frac{y^2}{r_0^2}+\frac{y^4}{r_0^4}} - \frac{y^2}{r_0^2}}{1+2(1+q_1)\frac{y^2}{r_0^2}+\frac{y^4}{r_0^4}}\right]^{1/2}$$

$$\int_0^1 \frac{(y^2)^{p_1}\sqrt{1-y^2}}{\left(y^2+\frac{\beta^2}{1-\beta^2}\right)^{p_2}}dy = \frac{\sqrt{\pi}}{4}\left(\frac{\beta^2}{1-\beta^2}\right)^{-p_2}\frac{\Gamma\left(p_1+\frac{1}{2}\right)}{\Gamma(p_1+2)}\,{}_2F_1\left(p_2, p_1+\frac{1}{2}; p_1+2, -\frac{1-\beta^2}{\beta^2}\right)$$

$$\delta = \pi|p_t|(1-\beta^2)^2\left\{\frac{\alpha^2}{2}+\alpha^4\left[1-\frac{15}{16}(\beta^2-1)^2\right]+\alpha^6\left[2-5(\beta^2-1)^2+\frac{105}{32}(\beta^2-1)^4\right]+\mathcal{O}(\alpha^8)\right\}$$

$$= 8\pi\frac{h^2\bar{h}^2}{(h+\bar{h})^3}\alpha^2[1+2\alpha^2(1-15v^2)+4\alpha^4(1-40v^2+420v^4)]+O(\alpha^8)$$

$$|p_t| = h+\bar{h}, \beta = \frac{h-\bar{h}}{h+\bar{h}}, v = \left(\frac{h\bar{h}}{(h+\bar{h})^2}\right)$$

$$\Psi_B^R = \int_0^\infty dh\int_0^h d\bar{h}(h\bar{h})^{\Delta-2}\left[\frac{z^{h+1}\bar{z}^{\bar{h}}-z^{\bar{h}}\bar{z}^{h+1}}{z-\bar{z}}\right]e^{i\delta(h,\bar{h})}$$

$$\Psi_B^R = \frac{z}{z-\bar{z}}\int_0^\infty dh\int_0^\infty d\bar{h}(h\bar{h})^{\Delta-2}(h-\bar{h})z^h\bar{z}^{\bar{h}}e^{i\delta(h,\bar{h})}$$

$$\Psi_B^R = \left[\frac{z}{z-\bar{z}}(z\partial_z-\bar{z}\partial_{\bar{z}})\right]\int_0^\infty dh\int_0^\infty d\bar{h}z^h\bar{z}^{\bar{h}}e^{i\delta(h,\bar{h})}$$

$$I(a,c) \equiv \int_0^\infty dh\int_0^\infty d\bar{h}\frac{h^{a+1}\bar{h}^{a+1}}{(h+\bar{h})^c}z^h\bar{z}^{\bar{h}}$$

$$= \frac{\Gamma^2(a+2)\Gamma(2a+4-c)}{\Gamma(2a+4)}\eta_R^{c-a-2}\sigma_R^{c-2a-4}\,{}_2F_1(a+2,c;2a+4;1-\eta_R)$$

$$\Psi_B^R \equiv \Psi_B^{R(0)}+\alpha^2\Psi_B^{R(2)}+\mathcal{O}(\alpha^4) = \Delta I(-1,0)+\alpha^2 8\pi i\Delta I(1,3)+\mathcal{O}(\alpha^4)$$

$$= \frac{1}{\eta_R^2\sigma_R^4}+\alpha^2 16\pi i\frac{(1-\eta_R^2)(1+28\eta_R+\eta_R^2)+12(1+3\eta_R+\eta_R^2)\eta_R\log(\eta_R)}{(1-\eta_R)^7\eta_R\sigma_R^5}+\mathcal{O}(\alpha^4)$$

$$\Delta \equiv \frac{1}{\sigma_R(1-\eta_R)}\left(\partial_{\sigma_R}-\frac{1+\eta_R}{\sigma_R}\partial_{\eta_R}\right)$$

$$\Psi_B^{R(4)} = -32\pi^2\Delta I(3,6)+16\pi i\Delta[I(1,3)-15I(3,7)]$$


$$\Psi_B^{R(6)} = \Delta\left\{-i\frac{(8\pi)^3}{3!}I(5,9)-128\pi^2[I(3,6)-15I(5,10)]+32\pi i[I(1,3)-40I(3,7)+420I(5,11)]\right\}$$

$$\left\langle O_{p_1}(\vec{x}_1,\vec{y}_1)O_{p_2}(\vec{x}_2,\vec{y}_2)O_{p_3}(\vec{x}_3,\vec{y}_3)O_{p_4}(\vec{x}_4,\vec{y}_4)\right\rangle$$


$$\gamma_{\min} = \max(p_{12},p_{43})\ ;\ \gamma_{\max} = \min(p_1+p_2,p_3+p_4).$$

$$\left\langle O_{p_1}O_{p_2}O_{p_3}O_{p_4}\right\rangle = \mathcal{P}_{\vec{p}}\sum_{\underline{R}}A_{\underline{R}}B_{\underline{R}},$$

$$\mathcal{P}_{\vec{p}} = g_{12}^{\frac{p_1+p_2}{2}}g_{34}^{\frac{p_3+p_4}{2}}g_{13}^{\frac{p_3-p_4}{2}}g_{14}^{\frac{p_1+p_4-p_2-p_3}{2}}g_{24}^{\frac{p_2-p_1}{2}}\left(\frac{g_{13}g_{24}}{g_{12}g_{34}}\right)^{\frac{\max(p_{12},p_{43})}{2}}$$



$$\max = p_{43}$$



$$\max = p_{12}$$

$$\mathcal{B}^{\frac{1}{2},\frac{1}{2}}_{[0,p,0](0,0)} \; ; \; \mathcal{B}^{\frac{1}{4},\frac{1}{4}}_{[q,p,q](0,0)}$$

$\underline{\lambda}$	$\tau = \Delta - l$	l	$\mathfrak{R}$	multiplet
$[\varnothing]$	γ	0	$[0,\gamma,0]$	$\frac{1}{2}$-BPS
$[1^\mu]$	γ	0	$[\mu,\gamma-2\mu,\mu]$	$\frac{1}{4}$-BPS
$[\lambda,1^\mu]\ (\lambda\geq 2)$	γ	$\lambda-2$	$[\mu,\gamma-2\mu-2,\mu]$	semi-short
$[\lambda_1,\lambda_2,2^{\mu_2},1^{\mu_1}]\ (\lambda_2\geq 2)$	$\gamma+2\lambda_2-4$	$\lambda_1-\lambda_2$	$[\mu_1,\gamma-2\mu_1-2\mu_2-4,\mu_1]$	long

$$\gamma = \gamma_{\min}, \gamma_{\min}+2, \dots, \gamma_{\max}$$

$$B_{\underline{R}} = \left[\left(\frac{g_{12}g_{34}}{g_{13}g_{24}}\right)^{\frac{\gamma-\gamma_{\min}}{2}} F_{\gamma,\underline{\lambda}}\right]$$

$$\frac{g_{12}g_{34}}{g_{13}g_{24}} = \frac{y_1y_2}{x_1x_2} \; ; \; \frac{g_{14}g_{23}}{g_{13}g_{24}} = \frac{(1-y_1)(1-y_2)}{(1-x_1)(1-x_2)}$$

$$U = x_1x_2 \; ; \; V = (1-x_1)(1-x_2) \; ; \; \sigma = \frac{1}{y_1y_2} \; ; \; \tau = \frac{(1-y_1)(1-y_2)}{y_1y_2}$$

$$F_{\gamma,\underline{\lambda}} = \left[P_{\underline{\lambda}} + P_{\underline{\lambda}+\underline{\square}} + \cdots\right] = \sum_{\underline{\mu}:\underline{\lambda}\subseteq\underline{\mu}} \left(T_\gamma\right)^{\underline{\mu}}_{\underline{\lambda}} P_{\underline{\mu}}$$

$$B_{\gamma,\underline{\lambda}} = \mathcal{P}_{\vec{p}}\left[k + S(x_1,x_2,y_1,y_2) + \frac{\prod_{ij}\left(x_i-y_j\right)}{(y_1y_2)^2} H(x_1,x_2,y_1,y_2)\right]$$

$$S = \frac{\prod_{ij}\left(x_i-y_j\right)}{(x_1-x_2)(y_1-y_2)}\left[\frac{f(x_2,y_1)}{x_2y_1(x_1-y_2)} + \frac{f(x_1,y_2)}{x_1y_2(x_2-y_1)} - \frac{f(x_1,y_1)}{x_1y_1(x_2-y_2)} - \frac{f(x_2,y_2)}{x_2y_2(x_1-y_1)}\right]$$

$$\frac{\prod_{ij}\left(x_i-y_j\right)}{(y_1y_2)^2} = \mathcal{I}(U,V,\sigma,\tau)$$

$$H_{\gamma,\underline{\lambda}} = (-1)^{\lambda_1-\lambda_2}\left(\frac{x_1x_2}{y_1y_2}\right)^{\beta} \frac{\left(F_{\lambda_1}(x_1)F_{\lambda_2-1}(x_2) - x_1 \leftrightarrow x_2\right)}{x_1-x_2} \frac{\left(G_{\lambda_1'}(y_1)G_{\lambda_2'-1}(y_2) - y_1 \leftrightarrow y_2\right)}{y_1-y_2}$$

$$F_\lambda(z) = z^{\lambda-1}\,{}_2F_1\left[\begin{matrix}\lambda+\frac{\gamma-p_{12}}{2},\lambda+\frac{\gamma-p_{43}}{2}\\ 2\lambda+\gamma\end{matrix};z\right]$$

$$G_\lambda(z) = z^{\lambda+1}\,{}_2F_1\left[\begin{matrix}\lambda-\frac{\gamma-p_{12}}{2},\lambda-\frac{\gamma-p_{43}}{2}\\ 2\lambda-\gamma\end{matrix};z\right]$$

$$\lambda_1 = 2+l+\frac{\tau-\gamma}{2}\ ;\ \lambda_2 = 2+\frac{\tau-\gamma}{2}\ ;\ \lambda_1' = -\frac{b-\gamma}{2}\ ;\ \lambda_2' = -a-\frac{b-\gamma}{2}$$

$$H_{\tau,l,[aba]} = (-1)^l (x_1x_2)^{\frac{\tau-\gamma_{\min}}{2}} \frac{x_1^{l+1}\,{}_2F_1\left[\begin{matrix}2+\frac{\tau}{2}+l-\frac{p_{12}}{2},2+\frac{\tau}{2}+l-\frac{p_{43}}{2}\\ 4+\tau+2l\end{matrix};x_1\right]{}_2F_1\left[\begin{matrix}1+\frac{\tau}{2}-\frac{p_{12}}{2},1+\frac{\tau}{2}-\frac{p_{43}}{2}\\ 2+\tau\end{matrix};x_2\right]-x_1\leftrightarrow x_2}{(x_1-x_2)}\times$$
$$\frac{1}{(y_1y_2)^{\frac{b-\gamma_{\min}}{2}}}\frac{y_1y_2^{-a}\,{}_2F_1\left[\begin{matrix}-1-\frac{b}{2}-a+\frac{p_{12}}{2},-1-\frac{b}{2}-a+\frac{p_{43}}{2}\\ -2-b-2a\end{matrix};y_2\right]{}_2F_1\left[\begin{matrix}-\frac{b}{2}+\frac{p_{12}}{2},-\frac{b}{2}+\frac{p_{43}}{2}\\ -b\end{matrix};y_1\right]-y_1\leftrightarrow y_2}{(y_1-y_2)}$$

$$f_{\gamma,\left[\lambda,1^{\lambda'-1}\right]}(x,y) = (-1)^\lambda\left(\frac{x}{y}\right)^\beta\left[x^\lambda\,{}_2F_1\left[\begin{matrix}\lambda+\frac{\gamma-p_{12}}{2},\lambda+\frac{\gamma-p_{43}}{2}\\ 2\lambda+\gamma\end{matrix};x\right]y^{\lambda'}\,{}_2F_1\left[\begin{matrix}\lambda'-\frac{\gamma-p_{12}}{2},\lambda-\frac{\gamma-p_{43}}{2}\\ 2\lambda'-\gamma\end{matrix};y\right]\right]$$

$$H_{\gamma,\left[\lambda,1^{\lambda'-1}\right]} = (-1)^{\lambda-1}\sum_{h=1}^{\beta}\left(\frac{x_1x_2}{y_1y_2}\right)^\beta\frac{(F_{1-h}(x_1)F_\lambda(x_2)-x_1\leftrightarrow x_2)}{x_1-x_2}\frac{(G_h(y_1)G_{\lambda'}(y_2)-y_1\leftrightarrow y_2)}{y_1-y_2}$$

$$B_{\gamma,\left[\lambda,1^{\lambda'-1}\right]}\Big|_{x_2=y_2} = \frac{(x_1-y_1)}{x_1y_1}f_{\gamma,\left[\lambda,1^{\lambda'-1}\right]}$$

$$f_{\gamma,\left[\lambda,1^{\lambda'-1}\right]}(x,y)+f_{\gamma+2,\left[\lambda-1,1^{\lambda'}\right]}(x,y) = 0$$

$$B^{\mathcal{L}}_{\gamma,\left[\lambda,1^{\lambda'-1}\right]} = B_{\gamma,\left[\lambda,1^{\lambda'-1}\right]}+B_{\gamma+2,\left[\lambda-1,1^{\lambda'}\right]} = \frac{\prod_{ij}(x_i-y_j)}{(y_1y_2)^2}\mathcal{H}_{\tau=\gamma,l=\lambda-2,a=\lambda'-1,b=\gamma-2\lambda'}$$

$$B_{\gamma,\left[\lambda,1^{\lambda'-1}\right]} = B^{\mathcal{L}}_{\gamma,\left[\lambda,1^{\lambda'-1}\right]}-B_{\gamma+2,\left[\lambda-1,1^{\lambda'}\right]}$$

$$f_{\gamma,[\emptyset]}(x,y) = \left(\frac{x_1}{y_1}\right)^\beta\sum_{h=1}^{\beta}x^{1-h}\,{}_2F_1\left[\begin{matrix}1-h+\frac{\gamma-p_{12}}{2},1-h+\frac{\gamma-p_{43}}{2}\\ 2(1-h)+\gamma\end{matrix};x\right]y^h\,{}_2F_1\left[\begin{matrix}h-\frac{\gamma-p_{12}}{2},h-\frac{\gamma-p_{43}}{2}\\ 2h-\gamma\end{matrix},y\right]$$

$$H_{\gamma,[\emptyset]} = \left(\frac{x_1x_2}{y_1y_2}\right)^\beta\sum_{1\le i<j\le\beta}\frac{\left[F_{1-i}(x_1)F_{1-j}(x_2)-x_1\leftrightarrow x_2\right]}{x_1-x_2}\frac{\left[G_i(y_1)G_j(y_2)-y_1\leftrightarrow y_2\right]}{y_1-y_2}$$

$$\partial H_{\tau,l,[abab]} \equiv U^{\tau/2}\partial_\tau U^{-\tau/2}H_{\tau,l,[abab]}$$

$$C_{[4^n][4^n];\mathcal{D}} = C^{(0)}_{[4^n][4^n];\mathcal{D}}+\frac{1}{a}C^{(1)}_{[4^n][4^n];\mathcal{D}}+\frac{1}{a^2}C^{(4)}_{[4^n][4^n];\mathcal{D}}+\cdots$$
$$\Delta_{\mathcal{D}_{\tau,l}} = \tau+l+\frac{2}{a}\eta_{(1);\mathcal{D}}+\frac{2}{a^2}\eta_{(4);\mathcal{D}}+\cdots$$

$$\log^4(U)\times\left[C^{(0)}_{[4^n][4^n]}\eta_{(4)}C^{(4)}_{44}H_{\tau,l,[abab]}\right]$$

$$\log^0(U)\times\left[2C^{(0)}_{[4^n][4^n]}\eta_{(4)}C^{(4)}_{44}\partial H_{\tau,l,[abab]}+\left[C^{(4)}_{[4^n][4^n]}C^{(0)}_{44}+C^{(0)}_{[4^n][4^n]}C^{(4)}_{44}\right]H_{\tau,l,[abab]}\right]$$

$$\log^{10}(U)\times\left[\frac{1}{2}C^{(4)}_{[4^n][4^n]}\eta^2_{(4)}C^{(4)}_{44}H_{\tau,l,[abab]}\right]$$

$$\log^5(U)\times\left[2C^{(0)}_{[4^n][4^n]}\eta^2_{(4)}C^{(4)}_{44}\partial H_{\tau,l,[abab]}+\left[C^{(4)}_{[4^n][4^n]}\eta_{(4)}C^{(4)}_{44}+C^{(4)}_{[4^n][4^n]}\eta_{(4)}C^{(4)}_{22}+C^{(4)}_{[4^n][4^n]}\eta_{(4)}C^{(4)}_{44}\right]H_{\tau,l,[abab]}\right]$$

$$\log^0(U)\times\left[2C^{(4)}_{[4^n][4^n]}\eta^2_{(4)}C^{(4)}_{44}\partial^2H_{\tau,l,[abab]}+2\sum_{i+a+j=2}C^{(i)}_{[4^n][4^n]}\eta_{(a)}C^{(j)}_{44}\partial H_{\tau,l,[abab]}\right.$$
$$\left.+\left[C^{(4)}_{[4^n][4^n]_4}C^{(4)}_{44}+C^{(4)}_{[4^n][4^n]}C^{(4)}_{44}+C^{(4)}_{[4^n][4^n]}C^{(4)}_{44}\right]H_{\tau,l,[abab]}\right]$$

$$\log^5(U)\times\left[C^{(4)}_{[4^n][4^n]}\eta^n_{(4)}C^{(4)}_{44}\partial H_{\tau,l,[aba]}+\frac{1}{2}\sum_{\substack{i+a+b+j=3\\a,b\le2}}C^{(i)}_{[4^n][4^n]}\eta_{(a)}\eta_{(b)}C^{(j)}_{44}H_{\tau,l,[abab]}\right]$$

$$\log^{10}(U)\times\left[2C^{(4)}_{[4^n][4^n]}\eta^n_{(4)}C^{(0)}_{44}\partial^2H_{\tau,l,[abab]}+2\sum_{\substack{i+a+b+j=3\\a,b\le2}}C^{(i)}_{[4^n][4^n]}\eta_{(a)}\eta_{(b)}C^{(j)}_{44}\partial H_{\tau,l,[abab]}\right.$$
$$\left.+\sum_{\substack{i+a+j=3\\a\le3}}C^{(i)}_{[4^n][4^n]}\eta_{(a)}C^{(j)}_{22}H_{\tau,l,[abab]}\right]$$

$$\log^0(U)\times\left[\frac{4}{3}C^{(0)}_{[4^n][4^n]}\eta^n_{(4)}C^{(0)}_{44}\partial^3H_{\tau,l,[abab]}+2\sum_{\substack{i+a+b+j=3\\a,b\le2}}C^{(i)}_{[4^n][4^n]}\eta_{(a)}\eta_{(b)}C^{(j)}_{44}\partial H^2_{\tau,l,[abab]}\right.$$
$$\left.+\sum_{\substack{i+a+j=3\\a\le3}}C^{(i)}_{[4^n][4^n]}\eta_{(a)}C^{(j)}_{44}\partial H_{\tau,l,[abab]}+\sum_{i+j=3}C^{(i)}_{[4^n][4^n]}C^{(j)}_{44}\partial H_{\tau,l,[abab]}\right]$$

$$C_{[4^n][4^n];\mathcal{T}}=\frac{1}{\sqrt{a}}\left[C^{\left(\frac{1}{2}\right)}_{[4^n][4^n];\mathcal{T}}+\frac{1}{a}C^{\left(\frac{3}{2}\right)}_{[4^n][4^n];\mathcal{T}}+\cdots\right]$$
$$\Delta_{\mathcal{T}_{\tau,l}}=\tau+l+\frac{2}{a}\eta_{(1);\mathcal{T}}+\frac{2}{a^2}\eta_{(2);\mathcal{T}}+\cdots$$

$$\log^1(U)\times0+\log^0(U)\times\left[C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}C^{\left(\frac{1}{2}\right)}_{44}\right]H_{\tau,l,[abab]}$$

$$\log^2(U)\times0+\log^1(U)\times\left[C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}\eta_{(4)}C^{\left(\frac{1}{2}\right)}_{44}H_{\tau,l,[abab]}\right]$$
$$\log^0(U)\times\left[2C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}\eta_{(4)}C^{\left(\frac{1}{2}\right)}_{44}\partial H_{\tau,l,[abab]}+\left[C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}C^{\left(\frac{3}{2}\right)}_{44}+C^{\left(\frac{3}{2}\right)}_{[4^n][4^n]}C^{\left(\frac{1}{2}\right)}_{44}\right]H_{\tau,l,[abab]}\right]$$

$$\log^3(U)\times 0+\log^2(U)\times\left[\frac{1}{2}C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}\eta^2_{(4)}C^{\left(\frac{1}{2}\right)}_{44}H_{\tau,l,[abab]}\right]$$

$$\log^1(U)\times\left[2C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}\eta^2_{(4)}C^{\left(\frac{1}{2}\right)}_{44}\partial H_{\tau,l,[abab]}+\left[C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}\eta_{(4)}C^{\left(\frac{1}{2}\right)}_{44}+C^{\left(\frac{3}{2}\right)}_{[4^n][4^n]}\eta_{(4)}C^{\left(\frac{1}{2}\right)}_{44}+C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}\eta_{(4)}C^{\left(\frac{3}{2}\right)}_{44}\right]H_{\tau,l,[abab]}\right]$$

$$\log^0(U)\times\left[2C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}\eta^2_{(4)}C^{\left(\frac{1}{2}\right)}_{44}\partial^2 H_{\tau,l,[abab]}+2\left[C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}\eta_{(4)}C^{\left(\frac{1}{2}\right)}_{44}+C^{\left(\frac{3}{2}\right)}_{[4^n][4^n]}\eta_{(4)}C^{\left(\frac{1}{2}\right)}_{44}+C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}\eta_{(4)}C^{\left(\frac{3}{2}\right)}_{44}\right]\partial H_{\tau,l,[abab]}\right.$$
$$\left.+\left[C^{\left(\frac{3}{2}\right)}_{[4^n][4^n]_4}C^{\left(\frac{3}{2}\right)}_{44}+C^{\left(\frac{5}{2}\right)}_{[4^n][4^n]}C^{\left(\frac{1}{2}\right)}_{44}+C^{\left(\frac{1}{2}\right)}_{[4^n][4^n]}C^{\left(\frac{5}{2}\right)}_{44}\right]H_{\tau,l,[abab]}\right]$$

$$G=\mathcal{P}_{\vec{p}}\times\frac{U^a}{V^b}\sigma^i\tau^j\ ;\ a=\frac{\gamma-\gamma_{\min}}{2}\ ;\ b,i,j\geq 0$$

$$G|_{\substack{x_1=y_1\\x_2=y_2}}=\mathcal{P}_{\vec{p}}\times k$$

$$G|_{x_2=y_2}=\mathcal{P}_{\vec{p}}\times\left[k+\frac{(x_1-y_1)}{x_1y_1}f_G(x_1,y_1)\right]$$

$$G=\mathcal{P}_{\vec{p}}\times\left[k+S(x_1,x_2,y_1,y_2)+\frac{\prod_{ij}\left(x_i-y_j\right)}{(y_1y_2)^2}H_G(x_1,x_2,y_1,y_2)\right]$$

$$f_G(x,y)=\sum_{h=1}^{\beta}f_{G,h}(x)\left[y^{h-\beta}\ {}_2F_1\left[\begin{matrix}h-\frac{\gamma-p_{12}}{2},h-\frac{\gamma-p_{43}}{2}\\2h-\gamma\end{matrix},y\right]\right]$$

$$f_{G,h}(x)=\sum_{\lambda\geq 1-h}c_{h,\lambda}\left[x^{\beta+\lambda}\ {}_2F_1\left[\begin{matrix}\lambda+\frac{\gamma-p_{12}}{2},\lambda+\frac{\gamma-p_{43}}{2}\\2\lambda+\gamma\end{matrix};x_1\right]\right]$$

$$\mathcal{G}_{2222}=U^2\sigma^2+\frac{U^2\tau^2}{V^2}+\frac{4}{a}\frac{U^2\sigma\tau}{V}\text{ where }\mathcal{P}_{2222}=4a^2g_{12}^2g_{34}^2$$

$$f_{\lambda'=5}(x_1)=\frac{x_1^2(2-2x_1+x_1^2)}{(x_1-1)^2}-\frac{4}{a}\frac{x_1^2}{x_1-1}$$

$$f_{\lambda'=10}(x_1)=\frac{1}{2}\frac{(x_1-2)x_1(-2+2x_1+x_1^2)}{(x_1-1)^2}+\frac{2}{a}\frac{x_1(x_1-2)}{x_1-1}$$

$$A^{4444}_{\mathcal{S},4,l,[0404]}=\left[(l+3)(l+4)+\frac{4}{a}\right]\frac{2(l+3)!^2}{(2l+6)}\frac{1+(-1)^l}{2}$$

$$\mathcal{P}_{4444^4}=8a^2\sqrt{a+2}g_{12}g_{14}g_{24}g_{34}^2\ ;\ \mathcal{G}_{4444^4}=\frac{4\sqrt{a+2}}{a}\left[1+U\sigma+\frac{U\tau}{V}\right]$$

$$|\mathcal{O}_2\mathcal{O}_2|\sqrt{|\mathcal{O}_2\mathcal{O}_2||\mathcal{O}_2^2\mathcal{O}_2^2|}$$

$$f_{\lambda'=4}(x_1)=\frac{4\sqrt{a+2}}{a}\left[x_1+\frac{x_1}{1-x_1}\right]$$

$$A^{44^444}_{\mathcal{S},4,[0404]}=\frac{4\sqrt{a+2}}{a}(l+4)\frac{2(l+3)!^2}{(2l+6)!}\frac{1+(-1)^l}{2}$$

$$G = \mathcal{P}_{44^4 44^4}\left[\frac{8}{a}U\sigma + \frac{24}{a}\frac{U\tau}{V}\right]$$

$$f_{\lambda'=1}(x_1) = \frac{8}{a}x_1 + \frac{24}{a}\frac{x_1}{1-x_1}$$

$$A_{4,l,[0404]} = \frac{4}{a}[(l+3)(l+4) + (-1)^l 6]\frac{(l+2)!\,(l+4)!}{(2l+6)!}$$

$$\mathcal{G}_{[4^4][4^4]44} = \left[\frac{8}{a}U\sigma + \frac{8}{a}\frac{U\tau}{V}\right] \text{ with } \mathcal{P}_{4^4 4^4 44}$$

$$f_{\lambda'=4}(x_1) = \frac{8}{a}\left[x_1 + \frac{x_1}{1-x_1}\right]$$

$$\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_4} - 2C^{(4)}_{44;\mathcal{D}_4}\right]C^{(4)}_{22;\mathcal{D}_4} = \frac{4}{3}(l+2)(l+5)\frac{(l+3)!^2}{(2l+6)!}\frac{1+(-1)^l}{2}$$

$$C^{(4)}_{[4^4][4^4];\mathcal{D}_4} = \frac{\frac{4}{3}(l+2)(l+5)\frac{(l+3)!^2}{(2l+6)!}\frac{1+(-1)^l}{2}}{C^{(4)}_{44;\mathcal{D}_4}} + 2C^{(4)}_{44;\mathcal{D}_4}$$

$$C^{(4)}_{44;\mathcal{D}_4} = \sqrt{\frac{(l+1)(l+6)}{3}}\frac{(l+3)!}{\sqrt{(2l+6)!}}\frac{1+(-1)^l}{2}$$

$$C^{(4)}_{44;\mathcal{D}_4} = \frac{\frac{2}{3} - 8\left(1 + 2\psi^{(0)}(l+4) - 2\psi^{(0)}(2l+7)\right)}{\sqrt{\frac{(l+1)(l+6)}{3}}}\frac{(l+3)!}{\sqrt{(2l+6)!}}\frac{1+(-1)^l}{2}$$

$$C^{(4)}_{[4^4][4^4];\mathcal{D}_4} = \frac{\frac{4}{3}(l+2)(l+5) + \frac{4}{3} - 16\left(1 + 2\psi^{(0)}(l+4) - 2\psi^{(0)}(2l+7)\right)}{\sqrt{\frac{(l+1)(l+6)}{3}}}\frac{(l+3)!}{\sqrt{(2l+6)!}}\frac{1+(-1)^l}{2}$$

$$C_{[4^4][4^4];\mathcal{O}_4^4} \sim \frac{1}{N}\ ;\ C_{44;\mathcal{O}_4^4} \sim \frac{1}{N}$$

$$\hat{C}_{[4^4][4^4]44}\big|_{\frac{1}{a}} - 2\hat{C}_{4444}\big|_{\frac{1}{a}} = \sum_l \left[C^{(4)}_{[4^4][4^4];\mathcal{D}_4} - 2C^{(4)}_{44;\mathcal{D}_4}\right]C^{(4)}_{44;\mathcal{D}_4}B^{\mathcal{L}}_{4,l,[0000]} +$$

$$\sum_{l,\mathcal{D}_6}\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_6} - 2C^{(4)}_{44;\mathcal{D}_6}\right]C^{(4)}_{44;\mathcal{D}_6}B^{\mathcal{L}}_{6,l,[0000]} + \sum_{l,\mathcal{J}_6}\left[C^{\left(\frac{1}{2}\right)}_{[4^4][4^4];\mathcal{J}_6} - 2C^{\left(\frac{1}{2}\right)}_{44;\mathcal{J}_6}\right]C^{\left(\frac{1}{2}\right)}_{44;\mathcal{J}_6}B^{\mathcal{L}}_{6,l,[0000]}$$

$$+ \text{higher twist}$$

$$\hat{C}_{[4^4][4^4]44}\big|_{\frac{\log(U)}{a^4}} - 2\hat{C}_{2222}\big|_{\frac{\log(U)}{a^4}} = \sum_l \left[C^{(4)}_{[4^4][4^4];\mathcal{D}_4} - 2C^{(4)}_{44;\mathcal{D}_4}\right]\eta_{\mathcal{D}_4}C^{(4)}_{44;\mathcal{D}_4}B^{\mathcal{L}}_{4,l,[0000]} +$$

$$\sum_{l,\mathcal{D}_6}\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_6} - 2C^{(4)}_{44;\mathcal{D}_6}\right]\eta_{\mathcal{D}_6}C^{(4)}_{44;\mathcal{D}_6}B^{\mathcal{L}}_{6,l,[0000]} + \sum_{l,\mathcal{J}_6}\left[C^{\left(\frac{1}{2}\right)}_{[4^4][4^4];\mathcal{J}_6} - 2C^{\left(\frac{1}{2}\right)}_{44;\mathcal{J}_6}\right]\eta_{\mathcal{J}_6}C^{\left(\frac{1}{2}\right)}_{44;\mathcal{J}_6}B^{\mathcal{L}}_{6,l,[0000]}$$

$$+ \text{higher twist}$$

$$\mathcal{H}^{t}_{[4^4][4^4]44}\Big|_{\frac{\log(U)}{a^2}} = \sum_{l}\left[-64\frac{(l+2)(l+5)}{(l+1)(l+6)}\frac{(l+3)!^2}{(2l+6)!}\frac{1+(-1)^l}{2}\right]H_{4,l,[0000]} +$$
$$\sum_{l}\left[-224\frac{(l+4)(l+5)}{(l+1)(l+8)}\frac{(l+4)!^2}{(2l+8)!}\frac{1+(-1)^l}{2}\right]H_{6,l,[0000]} + \text{ higher twist } \ldots$$

$$C_6^{(n)} = \begin{pmatrix} C^{(1)}_{11;O_1} & C^{(2)}_{22;O_2} \\ C^{(3)}_{33;O_3} & C^{(4)}_{44;O_4} \end{pmatrix}$$

$$C^{(4)}_{pp,6} = \left[C^{(4)}_{pp;O_2}, C^{(2)}_{pp;O_4}\right]^t \; ; \; 2p > 6$$

$$V_{[4^4][4^4],6} = C_6^{(4)} \cdot \left[C^{(4)}_{[4^4][4^4],6} - 2C^{(4)}_{44,6}\right]$$

$$V^{(1)}_{[4^4][4^4],6} = \sum_{\mathcal{D}_6} C^{(4)}_{44;\mathcal{D}_6}\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_6} - 2C^{(4)}_{44;\mathcal{D}_6}\right]$$
$$V^{(2)}_{[4^4][4^4],6} = \sum_{\mathcal{D}_6} C^{(4)}_{44;\mathcal{D}_6}\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_6} - 2C^{(4)}_{44;\mathcal{D}_6}\right]$$

$$\hat{\mathcal{C}}_{[4^4][4^4]44}\Big|_{\frac{1}{a}} - 2\hat{\mathcal{C}}_{4444}\Big|_{\frac{1}{a}} \; ; \; \hat{\mathcal{C}}_{[4^4][4^4]44}\Big|_{\frac{1}{a}} - 2\hat{\mathcal{C}}_{4444}\Big|_{\frac{1}{a}}$$

$$A_6 = \left(C^{(0)}\right)\cdot\left(C^{(0)}\right)^t \; ; \; M_6 = \left(C^{(0)}\right)\cdot\begin{pmatrix}\eta_1 & 0\\ 0 & \eta_2\end{pmatrix}\cdot\left(C^{(0)}\right)^t$$

$$(M_6)\cdot(A_6)^{-1}\cdot V_{[4^4][4^4],6} = \begin{cases}\sum\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_6} - 2C^{(4)}_{44;\mathcal{D}_6}\right]\eta_{\mathcal{D}_6}C^{(4)}_{44;\mathcal{D}_6} \\ \sum\left[C^{(4)}_{[4^4][4^4];\mathcal{D}_6} - 2C^{(4)}_{44;\mathcal{D}_6}\right]\eta_{\mathcal{D}_6}C^{(4)}_{44;\mathcal{D}_6}\end{cases}$$

$$\mathcal{G}_{pppp}\Big|_{\text{gravity}} = 1 + U^p\sigma^p + \frac{U^p\tau^p}{V^p} \; ; \; \mathcal{G}_{2233} = 1 + \frac{6}{a}\left[U\sigma + \frac{U\tau}{V}\right] + \frac{12}{a}\frac{U^2\sigma\tau}{V}$$

$$\mathcal{H}_{44pp} = -\frac{2pU^p\bar{D}_{p,p+4,4,4}}{(p-2)!} \; ; \; \mathcal{H}_{4444}|_{[0000]} = -\frac{3}{2}U^p\left[\bar{D}_{2,5,2,3} + \bar{D}_{2,5,3,2} + 6\bar{D}_{3,5,2,2} + 8\bar{D}_{3,5,3,3}\right]$$

$$A_6 = +\begin{pmatrix}\frac{1}{10} & 0 \\ 0 & \frac{1}{40}(l+2)(l+7)\end{pmatrix}(l+1)(l+8)\frac{(l+4)!^2}{(2l+8)!}$$
$$M_6 = -6\begin{pmatrix}4 & 12 \\ 12 & 44+9l+l^2\end{pmatrix}\frac{(l+4)!^2}{(2l+8)!}$$

$$\mathcal{C}_{[4^4][4^4]44}\Big|_{\frac{1}{a}} = 2\mathcal{C}_{4444}\Big|_{\frac{1}{a}}$$

$$\mathcal{P}_{[4^4][4^4]44} = g_{12}^4 g_{34}^3 \mathcal{N}_{[4^4]}\mathcal{N}_4$$
$$\mathcal{G}_{[4^4][4^4]44} = 1 + \frac{12}{a}\left[U\sigma + \frac{U\tau}{V}\right] + \frac{24}{a}\frac{U^2\sigma\tau}{V} + \frac{24}{a(a+2)}\left[U^2\sigma^2 + \frac{U^2\tau^2}{V^2} + \frac{4U^2\sigma\tau}{V} + 2\left[\frac{U^3\sigma^2\tau}{V} + \frac{U^3\sigma\tau^2}{V^2}\right]\right]$$

$$\mathcal{N}_{[4^4]}\mathcal{N}_4 = \frac{24(N^2-1)(N^2-4)(N^4-1)}{N}$$

$$\mathcal{C}_{[4^4][4^4]44}\big|_{\frac{1}{a}} = 2\mathcal{C}_{4444}|_{\frac{1}{a}}$$

$$\hat{\mathcal{C}}_{[4^4][4^4]44}\big|_{\frac{1}{a}} - 2\hat{\mathcal{C}}_{4444}\big|_{\frac{1}{a}} = \frac{4}{a}U^2\sigma^2 + \frac{4}{a}\frac{U^2\tau^2}{V^2} + \frac{16}{a}\frac{U^2\sigma\tau}{V},$$

$$V_{[4^4][4^4],6} = \left[\frac{4+9l+l^2}{5}\frac{(l+4)!^2}{(2l+8)!}, 0\right]\frac{1+(-1)^l}{2}$$

$$\sum\left[C^{(1)}_{[4^4][4^4];\mathcal{D}_6} - 2C^{(4)}_{44;\mathcal{D}_6}\right]\eta_{\mathcal{D}_6}C^{(4)}_{44;\mathcal{D}_6} = -96\frac{(l^2+9l+4)}{(l+1)(l+8)}\frac{(l+4)!^2}{(2l+8)!}\frac{1+(-1)^l}{2}$$

$$\mathcal{H} = \sum_{\tau,l,[abab]} A_{\tau,l,[abab]}H_{\tau,l,[abab]}$$

$$\mathbb{F}_{\tau,l} = \frac{\left(\frac{\tau}{2}+1\right)!^2\left(l+\frac{\tau}{2}\right)!^2}{48\tau!\,(2l+\tau+2)!}\;;\; T = \frac{\tau+1}{2}\;;\; L = l+\frac{\tau+3}{2}$$

$$A_{\tau,l} = \mathbb{F}_{\tau,l}\times\mathbb{R}_{\tau,l}$$

$$\mathcal{P}_{4[4^4]4[4^4]}\left(U^2\sigma^2 + \frac{2U^2\tau^2}{V^2}\right)$$

$$\mathbb{R}_{\tau,l} = (l+1)(2l+\tau+4)(2l+\tau+2)(\tau+2+l)\left((-1)^l\frac{96}{\tau(\tau+2)} - \frac{1}{4} + L^2\right)$$

$$A_{6,l} = \frac{2}{15}(l+1)(l+4)(l+5)(l+8)\left[\frac{1}{2}(l+3)(l+6) + 2\frac{1+(-1)^l}{2}\right]\frac{(l+3)!^2}{(2l+8)!}$$

$$\mathbb{R}_{\tau,l} = \mathbb{R}^-_{\tau,l}\frac{1-(-1)^l}{2} + \mathbb{R}^+_{\tau,l}\frac{1+(-1)^l}{2}$$

$$\mathbb{R}^-_{\tau,l} = -\frac{(-4+\tau)(-2+\tau)(\tau+4)(\tau+6)}{2}\left[(l+1)(l+\tau+2)\left(L^2 - \frac{96}{\tau(\tau+2)} - \frac{3}{5}T^2 + \frac{71}{10}\right)\right]$$

$$\mathbb{R}^+_{\tau,l} = \mathbb{R}^-_{\tau,l} - 24\frac{(-4+\tau)(-2+\tau)(\tau+4)(\tau+6)}{\tau(\tau+2)}(2+2l+\tau)(4+2l+\tau)$$

$$A_{6,l} = -8\left[(l+2)(l+7)(16+9l+l^2)\frac{1+(-1)^l}{2} + (l+1)(l+3)(l+6)(l+8)\frac{1-(-1)^l}{2}\right]\frac{(l+3)!^2}{(2l+8)!}$$

$$\mathcal{H}^c_{[4^4][4^4]44}\Big|_{\frac{1}{a^2}} - 4\mathcal{H}_{4444}|_{\frac{1}{a}}$$

$$A_{6,l} = 32(-1)^{l+1}\left[\frac{3(l+1)(l+3)(l+6)(l+8)}{(l+4)(l+5)} - ((2l+9)^2-37)\frac{1+(-1)^l}{2}\right]\frac{(l+3)!^2}{(2l+8)!}$$

$$\mathbb{R}_{\tau,l} = 32\frac{(l+1)(l+\tau+2)}{\tau^2(\tau+2)^2}(-1)^{l+1}\left[2\left(\frac{\tau}{2}-2\right)_6\left[96+\frac{8}{15}\left(\frac{\tau}{2}-2\right)_6\left(\frac{1}{2}-\frac{1+(-1)^l}{2}\left(1+\frac{3}{(l+1)(l+\tau+2)}\right)\right)\right]+\Sigma_{\tau,l}\right]$$

$$\Sigma_{\tau,l} = \sum_{i=0}^{\frac{\tau-6}{2}} \frac{(-1)^i(T-I-2)_5(T+I-2)_5 \times I \times \sum_{j=0}^{3} B_j(T-I)^{2j}}{L^2-I^2}$$

$$B_1 = \frac{(-5+I+T)(-3+I+T)(3+I+T)(5+I+T)(1191-71(I+T)^2)}{92400}$$

$$B_2 = \frac{-508221+100379(I+T)^2-2979(I+T)^4-59(I+T)^6}{831600}$$

$$B_3 = \frac{32445-2979(I+T)^2-1330(I+T)^4+64(I+T)^6}{831600}$$

$$B_4 = \frac{-639-59(I+T)^2+64(I+T)^4+4(I+T)^6}{831600}$$

$$A_{6,l} = -192\frac{(l+3)!^2}{(2l+8)!}[$$

$$\left[(82+45l+5l^2)-\frac{16}{(l+3)}+\frac{16}{(l+6)}-4(l+2)(l+7)\text{ Harmonic Number }(l+4)\right]\frac{1+(-1)^l}{2}+$$

$$\left[(86+45l+5l^2)-\frac{24}{(l+4)}+\frac{24}{(l+5)}-4(l+2)(l+7)\text{ Harmonic Number }(l+4)\right]\frac{1-(-1)^l}{2}\Bigg]$$

[n = 0]	$\frac{1}{15}(l+1)(l+4)^2(l+5)^2(l+8)\frac{(l+3)!^2}{(2l+8)!}$
[n = 5]	$-8(l+4)(l+5)\left((l+1)(l+8)+24\frac{1+(-1)^l}{2}\right)\frac{(l+3)!^2}{(2l+8)!}$
[n = 10]	$480(l+1)(l+8)\left(84+\left(84+\frac{1008}{(l+1)(l+8)}\right)\frac{1+(-1)^l}{2}\right)\frac{(l+3)!^2}{(2l+8)!}$

$$\hat{\mathcal{C}}_{[4^n][4^n]44} = \text{ free } + \frac{(x_1-y_1)(x_2-y_2)}{y_1y_2}\mathcal{H}_{[4^n][4^n]44}(U,V)$$

$$\mathcal{H}_{[4^n][4^n]44} = \iint \frac{dSdt}{(2\pi i)^2}\Gamma[-s]^2\Gamma[-t]^2\Gamma[s+t+2]^2U^{1+s}V^t\mathcal{M}_{[4^n][4^n]44}(s,t)$$

$$\hat{\mathcal{C}}_{[4^n][4^n]44} = \text{ free } + \frac{(x_1-y_1)(x_1-y_2)(x_2-y_1)(x_2-y_2)}{(y_1y_2)^2}\mathcal{H}_{[4^n][4^n]44}(U,V)$$

$$\mathcal{H}_{[4^n][4^n]44} = \oiiint \frac{dSdt}{(2\pi i)^2}\Gamma[-s]^2\Gamma[-t]^2\Gamma[s+t+4]^2 U^{2+s}V^t\mathcal{M}_{[4^n][4^n]44}(s,t)$$

$$\mathcal{P}_L(z,\bar{z}) = \sum_{r=0}^{L}\frac{(-1)^r}{r!}B\begin{bmatrix}2L-r\\ L\end{bmatrix}\log^r(z\bar{z})(\mathrm{Li}_{2L-r}(z)-\mathrm{Li}_{2L-r}(\bar{z}))$$

$$B\begin{bmatrix}a\\ b\end{bmatrix} = \frac{a!}{b!\,(a-b)!}$$

$$\mathcal{P}_0(z,\bar{z}) = -\frac{z}{z-1}+\frac{\bar{z}}{\bar{z}-1}$$
$$\mathcal{P}_1(z,\bar{z}) = \log(z,\bar{z})(\log(1-z)-\log(1-\bar{z}))+2(\mathrm{Li}_2(z)-\mathrm{Li}_2(\bar{z}))$$
$$\mathcal{P}_2(z,\bar{z}) = \frac{1}{2}\log^2(z,\bar{z})(\mathrm{Li}_2(z)-\mathrm{Li}_2(\bar{z}))-3\log(z,\bar{z})(\mathrm{Li}_3(z)-\mathrm{Li}_3(\bar{z}))+6(\mathrm{Li}_4(z)-\mathrm{Li}_4(\bar{z}))$$

$$\mathcal{P}_L\left(\frac{1}{z},\frac{1}{\bar{z}}\right) = -\mathcal{P}_L(z,\bar{z})$$

$$\mathcal{P}_L(z,\bar{z})\ ;\ \mathcal{P}_L\left(\frac{z}{z-1},\frac{\bar{z}}{\bar{z}-1}\right)\ ;\ \mathcal{P}_L(1-z,1-\bar{z})$$

$$\Phi^{(L)}(z,\bar{z}) = -\frac{1}{z-\bar{z}}\mathcal{P}_L\left(\frac{z}{z-1},\frac{z}{\bar{z}-1}\right)$$

$$\frac{UV}{z-\bar{z}}\partial_z\partial_{\bar{z}}[(z-\bar{z})\Phi^L(z,\bar{z})] = -\Phi^{(L-1)}(z,\bar{z})\ ;\ \Phi^{(0)} = 1$$

$$U = (1-z)(1-\bar{z})\ ;\ V = z\bar{z}$$

$$\Phi^{(L)}(1-z,1-\bar{z}) = \Phi^{(L)}(z,\bar{z})$$

$$\Phi^{(L)}(z,\bar{z})\big|_{\log^L(U)\log^L(V)} = \frac{1}{(L!)^2}\frac{1}{z-\bar{z}}$$

$$z = 1-\zeta$$

$$\zeta = U+\sum_{n,m\geq 1}\frac{U^n}{n!}\frac{V^m}{m!}\frac{(m+n-1)!\,(m+n-2)!}{(n-1)!\,(m-1)!}$$
$$\bar{z} = V+\sum_{n,m\geq 1}\frac{U^n}{n!}\frac{V^m}{m!}\frac{(m+n-1)!\,(m+n-2)!}{(n-1)!\,(m-1)!}$$

$$\Phi^{(L)}(U,V)\big|_{\log^L(U)\log^L(V)} = \frac{1}{(L!)^2}\frac{1}{1-\zeta-\bar{z}} = \frac{1}{(L!)^2}\sum_{m,n,\geq 0}\frac{U^nV^m}{n!\,m!}\times\frac{(m+n)!^2}{n!\,m!}$$

$$\Phi^{(L)} = \oiiint \frac{U^sV^t}{(2\pi i)^2}\Gamma[-s]^2\Gamma[-t]^2\Gamma[-u]^2\mathcal{K}^{(L)}(s,t)\ ;\ u = -s-t-1\ ;\ L\geq 1$$

$$\Delta_{st}[\mathcal{K}^{(L)}] := -st\mathcal{K}^{(L)}[s,t]+\frac{s^2t}{s+t}\mathcal{K}^{(L)}[s-1,t]+\frac{t^2s}{s+t}\mathcal{K}^{(L)}[s,t-1]$$

$$\Delta_{st}\left[\mathcal{K}^{(L)}\right] = -\mathcal{K}^{(L-1)}(s,t)$$

$$\mathcal{K}^{(L)}(s,t) = \frac{1}{(s-n)^{L-1}(t-m)^{L-1}} + n, m \in \mathbb{N}$$

$$\mathcal{K}^{(2)}(s,t) = \frac{2}{(2!)^2}\left[-\Psi_1^2 + \Psi_2 - \pi^2\right]$$

$$\mathcal{K}^{(3)}(s,t) = \frac{3}{(3!)^2}\left[+\Psi_1^4 + 3\Psi_2^2 - 2\Psi_1\Psi_3 - 2\pi^2\Psi_2 - 4\pi^2\mathcal{K}^{(2)} - \pi^4\right]$$

$$\mathcal{K}^{(4)}(s,t) = \frac{4}{(4!)^2}[-\Psi_1^6 - 3\Psi_1^4\Psi_2 - 9\Psi_1^2\Psi_2^2 + 9\Psi_2^3 + 4\Psi_1^3\Psi_3 - 12\Psi_1\Psi_2\Psi_3 - 4\Psi_3^2 + 3\Psi_1^2\Psi_4 + 3\Psi_2\Psi_4$$
$$+\pi^2(6\Psi_1^2\Psi_2 - 12\Psi_2^2 + 6\Psi_1\Psi_3 - \Psi_4) - 9\times 4\times\pi^2\mathcal{K}^{(3)} - 6\pi^4\mathcal{K}^{(2)} - \pi^6\big]$$

$$\Psi_n(s,t) = \text{ Polygamma }[n-1,-s] + (-1)^n \text{ Polygamma }[n-1,-t].$$

$$\mathcal{C}(z,\bar{z})|_{\frac{\log L(U)\log^L(V)}{(L!)^2}} = \frac{P(U,V)}{(z-\bar{z})^{2\gamma+1}}$$

$$P(U,V) = \sum_{a,b} c_{ab}U^aV^b\ ;\ c_{ab} \in \mathbb{R}$$

$$\frac{U^aV^b}{(z-\bar{z})^{2\gamma+1}} = \frac{\gamma!}{(2\gamma)!}\oiint (-U)^s(-V)^t\frac{\Gamma[-s]\Gamma[-t]}{\Gamma[s+1]\Gamma[t+1]}\Gamma[s+t+X]^2\times m_{ab,\gamma}$$

$$m_{ab,\gamma} = \left[(-)^{a+b}\frac{(-s)_a(-t)_b}{(s+1)_{\gamma-a}(t+1)_{\gamma-b}}(s+t+X)_{2\gamma+1-a-b-X}(s+t+X)_{\gamma+1-a-b-X}\right]$$

$$A(s,t) = \sum_{a,b} c_{ab}m_{ab,\gamma}$$

$$\oiint \frac{U^sV^t}{(2\pi i)^2}\Gamma[-s]^2\Gamma[-t]^2\Gamma[-u]^2\left[\mathcal{K}^{(L)}(s,t)A(s,t) + \cdots\right] = \frac{\log^L(U)\log^L(V)}{(L!)^2}\frac{P(U,V)}{(z-\bar{z})^{2\gamma+1}} + \cdots$$

weight	anti-symmetric functions ;	symmetric functions
0	;	1
1	;	$\log(V)$, $\log(U)$
2	$\mathcal{P}_1$;	$\log^2(V)$, $\log(U)\log(V)$, $\log^2(U)$
3	$\mathcal{P}_1\log(V)$, $\mathcal{P}_1\log(U)$;	$\log^3(V)$, $\log(U)\log(V)\log(UV)$, $\log^3(U)$
		$\log(U)\log(V)\log(\frac{U}{V})$
		$\partial_-\mathcal{P}_2(1-z)$, $\partial_-\mathcal{P}_2(z)$, $\partial_-\mathcal{P}_2(\frac{z}{z-1})$

weight	functions
4^-	$\mathcal{P}_1 \log^2(V)$, $\mathcal{P}_1 \log^2(U)$, $\mathcal{P}_1 \log^2(\frac{U}{V})$
	$\mathcal{P}_2(1-z)$, $\mathcal{P}_2(z)$, $\mathcal{P}_2(\frac{z}{z-1})$

weight	functions
4^+	$\mathcal{P}_1^2$, ~~$\log(U)\log(V)(\log^2(U)+\log^2(V))$~~ , ~~$\log^4(V)$~~ , ~~$\log^4(U)$~~
	$\log(U)\log^2(V)\log(\frac{U}{V})$, $\log^2(U)\log(V)\log(\frac{U}{V})$
	$\zeta_3 \log(V)$, $\zeta_3 \log(U)$, $\log(UV)\partial_-\mathcal{P}_2(z)$
	$\partial_-\partial_+\mathcal{P}_3(1-z)$, $\partial_-\partial_+\mathcal{P}_3(z)$, $\partial_-\partial_+\mathcal{P}_3(\frac{z}{z-1})$

weight	functions
5^-	~~$\mathcal{P}_1 \log^3(V)$~~ , ~~$\mathcal{P}_1 \log(U)\log(V)\log(UV)$~~ , ~~$\mathcal{P}_1 \log^3(U)$~~
	$\mathcal{P}_1 \log(U)\log(V)\log(\frac{U}{V})$
	$\mathcal{P}_1\partial_-\mathcal{P}_2(1-z)$, $\mathcal{P}_1\partial_-\mathcal{P}_2(z)$, $\mathcal{P}_1\partial_-\mathcal{P}_2(\frac{z}{z-1})$
	$\log(U)\mathcal{P}_2(1-z)$, $\log(V)\mathcal{P}_2(z)$, $\log(UV)\mathcal{P}_2(\frac{z}{z-1})$
	$\partial_+\mathcal{P}_3(1-z)$, $\partial_+\mathcal{P}_3(z)$, $\partial_+\mathcal{P}_3(\frac{z}{z-1})$

$$\mathcal{H}^t_{[4^4][4^4]44} = \text{span}\{\mathcal{P}_3, \partial_-\mathcal{P}_3, \partial_+\mathcal{P}_3, \partial_+\partial_-\mathcal{P}_3\} \cup \{4^-, 3, 2, 1, 0\}$$

$$\partial_+\mathcal{P}_4\left(\frac{z}{z-1}, \frac{\bar{z}}{\bar{z}-1}\right)$$

$$\frac{1}{(z-\bar{z})}\left[z(1-z)\frac{\partial}{\partial z} + \bar{z}(1-\bar{z})\frac{\partial}{\partial \bar{z}}\right]\left[(z-\bar{z})\oint \frac{U^s V^t}{(2\pi i)^2}\Gamma[-s]^2\Gamma[-t]^2\Gamma[s+t+1]^2\mathcal{K}^{(3)}(s,t)\right]$$

$$\iint \frac{1}{(2\pi i)^2}\Gamma[-s]^2\Gamma[-t]^2\Gamma[s+t+1]^2[(s-t)U^sV^t + (1+s+t)(U^sV^{t+1} - U^{s+1}V^t)]\mathcal{K}^{(3)}(s,t)$$

$$\iint \frac{U^sV^t}{(2\pi i)^2}\Gamma[-s]^2\Gamma[-t]^2\Gamma[s+t+1]^2\left[(s-t) + \frac{t^2\mathcal{K}^{(3)}(s,t-1) - s^2\mathcal{K}^{(3)}(s-1,t)}{s+t}\right]$$

$$\frac{1}{3}\left[\Psi_1^3(u,t) - \frac{1}{2}\Psi_3(u,t) + \pi^2\Psi_1\right]$$

$$\frac{1}{(z-\bar{z})}\left[z\frac{\partial}{\partial z}+\bar{z}\frac{\partial}{\partial \bar{z}}\right]\left[(z-\bar{z})\oint \frac{U^s V^t}{(2\pi i)^2}\Gamma[-s]^2\Gamma[-t]^2\Gamma[s+t+1]^2\mathcal{K}^{(3)}(s,t)\right]$$

$$\left\langle \mathcal{O}_2\mathcal{O}_2^2\mathcal{O}_2\mathcal{O}_2^2\right\rangle\Big|_{\tau=4} = \mathcal{P}_{4[4^4]4[4^4]4}\times\left[\frac{8}{a}U\sigma+\frac{24}{a}\frac{U\tau}{V}\right] \text{ where } \mathcal{P}_{4[4^4]4[4^4]4} = 16a^2(a+2)g_{12}^2g_{24}^2g_{34}^2$$

$$\{\mathcal{O}_2\,\Box^{\tau-4}\,\partial^l\mathcal{O}_2, \mathcal{O}_3\,\Box^{\tau-6}\,\partial^l\mathcal{O}_3, \ldots\}\star\left\langle \mathcal{O}_2\mathcal{O}_2^3\mathcal{O}_2\mathcal{O}_2^3\right\rangle$$

$$M_{\text{Pl}}^2 \simeq (2\pi R)^n M_D^{n+2}$$

$$\frac{dN}{dVdt} = -g\int \frac{d^4k}{(2\pi)^3}\frac{f_B(\omega)}{\omega}\text{Im}\left[\Pi_{xx}-\frac{(\Pi_{\gamma x})^2}{\Pi_{\gamma\gamma}-m_x^2}\right]+\mathcal{O}(\lambda^4)$$


$$\frac{dN}{dVdt} = g\int \frac{d^4k}{(2\pi)^3}\frac{f_B(\omega)}{\omega}\text{Re}\left[\Pi_{\gamma x}\right]^2\frac{-\text{Im}\left[\Pi_{\gamma\gamma}\right]}{\left(\text{Re}\left[\Pi_{\gamma\gamma}\right]-m_x^2\right)^2-\text{Im}\left[\Pi_{\gamma\gamma}\right]^2}$$

$$\frac{dN}{dVdt} = \frac{g}{2\pi}\int d\omega\frac{k}{e^{\omega/T}-1}\text{Re}\left[\Pi_{\gamma x}\right]^2\delta\left(\text{Re}\left[\Pi_{\gamma\gamma}\right]-m_x^2\right)$$

$$\frac{dQ}{dVdt} = \frac{g}{2\pi}\frac{k_*\omega_*}{e^{\omega_*/T}-1}\text{Re}\left[\Pi_{\gamma x}(\omega_*)\right]^2\left|\frac{d\text{Re}\left[\Pi_{\gamma\gamma}(\omega)\right]}{d\omega}\right|_{\omega=\omega_*}^{-1}$$

$$\frac{d\text{Re}\left[\Pi_{\gamma\gamma}^L(\omega)\right]}{d\omega}\Big|_{\omega=\omega_*} = -\omega_*\frac{m_x^2}{k_*^2}\left(2+\frac{m_x^2}{\omega_*^2}\frac{(\omega_*^2-k_*^2v_0^2)-3\omega_p^2}{\omega_*^2-k_*^2v_0^2}\right)$$

$$\frac{d\text{Re}\left[\Pi_{\gamma\gamma}^T(\omega)\right]}{d\omega}\Big|_{\omega=\omega_*} = -\omega_*\frac{m_x^2}{k_*^2}\left(\frac{m_x^2}{\omega_*^2}-\frac{3\omega_p^2-2m_x^2}{\omega_*^2-k_*^2v_0^2}\right)$$



$$\Pi_{\gamma h,f}^T = \frac{q_f e\kappa}{3\sqrt{2}\pi^2}p_F^3\frac{m_h}{v_*\omega_*}\left(1-\frac{m_h^2}{\omega_p^2}\right)$$

$$C\frac{dT_c}{dt} = -L_\gamma^\infty - L_\nu^\infty - L_h^\infty + H,$$

$$\frac{dQ}{dVdt} = \frac{\pi^{(n-17)/2}\Gamma(n+4)}{270} h_n Y_n \kappa^2 (RT_c)^n \sigma_0 T_c^4 p_F^5$$


$$h_n = \frac{3n^2+18n+19}{(n+5)(n+3)} \frac{1}{\Gamma\left(\frac{n+3}{2}\right)}$$

$$Y_n = \zeta(n+4) + 4\pi^2 \frac{\zeta(n+2)}{(n+2)(n+3)}$$

$$\mathcal{L}^i\left(L^i_{\text{obs}}, t^i_{\text{obs}}, \sigma^i_L, \sigma^i_t \mid m_{\text{KK}}, \boldsymbol{\theta}\right) = \mathcal{N}\left(L(m_{\text{KK}}, \boldsymbol{\theta}) - L^i_{\text{obs}}, \sigma^i_L\right) \mathcal{N}\left(t - t^i_{\text{obs}}, \sigma^i_t\right),$$

$$q(m_{\text{KK}}) = -2\ln\left(\frac{\mathcal{L}_{\text{tot}}\left(m_{\text{KK}}, \Phi_{\max}(m_{\text{KK}})\right)}{\mathcal{L}_{\text{tot}}\left(\hat{m}_{\text{KK}}, \hat{\Phi}\right)}\right),$$

$$\frac{dQ}{dVdt} = \frac{\kappa^2}{18}\sqrt{\frac{T^7}{m_N}} (RT)^n \pi^{\frac{n-3}{2}} h_n n_n^2 \sigma_0^{nn} \frac{\Gamma(n+1)\Gamma(n+5)}{\Gamma\left(n+\frac{5}{2}\right)} \frac{n^2+9n+23}{(2n+7)(2n+5)}$$



$$\sum |M|^2 = \frac{\kappa^2 f_{pp}^2 m_N^2}{6(x-1)^2}\left[4\left(6y - x(6x^2 - 6xy + y(y+4) + 10)\right) - 4z^2(1-x+y)^2\right.$$
$$\left. -z(5x^3 + x^2(13-8y) + x(y^2+12y-9) + 2y^3 - 13y^2 + 12y - 9)\right]$$

$$x = \frac{t}{m_\pi^2}, y = \frac{m_h^2}{m_\pi^2}, z = \frac{t - m_\pi^2}{E_\pi^2}$$

$$\frac{d\dot{n}}{d\omega} = \int \frac{d^4p_2}{(2\pi)^3 2m_N} \frac{d^4p_1}{(2\pi)^3 2E_\pi} \frac{d^4p_4}{(2\pi)^3 2m_N} \frac{4\pi\omega^2}{(2\pi)^3 2\omega} (2\pi)^4 \delta^4(p_1 + p_2 - p_3 - p_4) |M|^2 f_p f_\pi (1 - f_n),$$

$$\frac{dQ}{dVdt} = \int d\omega\omega \frac{d\dot{n}}{d\omega}$$

$$\frac{dQ}{dVdt} = \frac{T^{11/2}}{(2\pi)^5\sqrt{2m_N}} \left(\frac{z_p z_\pi}{z_n + 1}\right) \left(\int_0^\infty dx_p \frac{x_p}{e^{x_p^2} + z_p}\right) \left(\int_{x_{\min}}^\infty dx_\pi \frac{x_\pi^2 \epsilon_\pi}{e^{\epsilon_\pi - y_\pi} - z_\pi} \int_{-1}^1 d\cos\,\theta |M|^2\right)$$

$$z_i = \exp\left((\mu_i - m_i)/T\right)$$

$x_p = p_2/\sqrt{2m_N T}, x_\pi = p_1/T, y_\pi = m_\pi/T, \epsilon_\pi = E_\pi/T$; and $x_{\min}$

$$E_\pi(p) = \sqrt{p^2 + m_\pi^2} + \Sigma_{\pi^-}(p)$$

$$\Sigma_{\pi^-}(p) = \int \frac{d^4k}{(2\pi)^3} f_n(E_n(k)) V_{n\pi^-}(p_{\text{cm}})$$

$$p_{\mathrm{cm}} = \tilde{m}\left(p^2/m_\pi^2 + k^2/m_N^2 - 2\mathbf{p}.\mathbf{k}/(m_\pi m_N)\right)^{1/2}$$

$$V_{n\pi^-}(p_{\mathrm{cm}}) = -\alpha \sum_{l,\nu} (2l+1)\frac{2\pi\delta_{l,\nu}}{\tilde{m}p_{\mathrm{cm}}}$$

$$\Gamma^{\text{Supermassive Particle}} = \lambda^2 \frac{m_h^3}{M_{\mathrm{Pl}}^2} \simeq (10^{14}\mathrm{yr})^{-1}\lambda^2\left(\frac{m_h}{\mathrm{MeV}}\right)^3$$

$$\Gamma \simeq \beta^2 \frac{m_h^3}{M_{\mathrm{Pl}}^2}\sqrt{\frac{m_{\mathrm{KK}}\delta n}{m_h}},$$

$$\left(\frac{m_h}{m_{\mathrm{KK}}}\right)^{3/2}\delta n^{5/2}$$

$$\Gamma^{\mathrm{tot}}(m) \simeq \beta^2 \frac{m_h^4 \delta n^3}{M_{\mathrm{Pl}}^2 m_{\mathrm{KK}}}$$

$$\frac{1}{\Gamma^{\mathrm{tot}}} \simeq 0.1\mathrm{yr}\left(\frac{10}{\beta}\right)^2\left(\frac{10}{\delta n}\right)^3\left(\frac{10\mathrm{MeV}}{m_h}\right)^4\left(\frac{m_{\mathrm{Kaluza-Klein}}}{\mathrm{eV}}\right)$$

$$\frac{dm_h(t)}{dt} \simeq -\Gamma^{\mathrm{tot}}(m_h)\frac{m_h(t)}{2}$$

$$m_h(t) \sim m_0(1+\Gamma^{\mathrm{tot}}(m_0)t/2)^{-1/4}$$

$$n(t) \simeq n_0 m_0/m_h(t)$$

$$\dot{E}_{\text{Supermassive Particle}}(t_{\mathrm{obs}}) \simeq n(t_{\mathrm{obs}})m_h(t_{\mathrm{obs}})\frac{\lambda^2 m_h(t_{\mathrm{obs}})^3}{M_{\mathrm{Pl}}^2}$$

$$\frac{\dot{E}_{\text{Supermassive Particle}}(t_{\mathrm{obs}})}{\dot{E}_0} \simeq (1+\Gamma^{\mathrm{tot}}(m_0)t_{\mathrm{obs}}/2)^{-3/4}$$

$$\Pi_{\gamma h}^{\mu,\alpha\beta}(Q) = \sum_f \Pi_{\gamma h,f}^{\mu,\alpha\beta}(Q)$$

$$E_p^2 = p^2 + m_f^2$$

$$\Pi_{\gamma h,f}^{\mu,\alpha\beta}(Q) = q_f e\kappa \int \frac{d^4p}{(2\pi)^3}\frac{1}{2E_p}\Big(f_e(E_p) - f_{\bar{e}}(E_p)\Big)\left(A^{\mu\alpha\beta} + \frac{B^{\mu\alpha\beta}+C^{\mu\alpha\beta}}{4(P\cdot Q)^2 - Q^4}\right)$$

$$A^{\mu\alpha\beta} = \frac{2Q^\mu}{m_A^2\xi}\big(P^\alpha Q^\beta + P^\beta Q^\alpha - g^{\alpha\beta}(P\cdot Q)\big) - \big(P^\alpha g^{\mu\beta} + P^\beta g^{\mu\alpha}\big),$$

$$\begin{aligned} B^{\mu\alpha\beta} = {} & 2P^\mu\big[g^{\alpha\beta}(4(P\cdot Q)^2 - Q^4) - 4(P\cdot Q)\big(P^\alpha Q^\beta + P^\beta Q^\alpha\big) + Q^2 Q^\alpha Q^\beta\big] \\ & + Q^\mu\big[Q^2\big(P^\alpha Q^\beta + P^\beta Q^\alpha\big) - 8(P\cdot Q)P^\alpha P^\beta\big] \\ & - Q^2(P\cdot Q)\big(Q^\alpha g^{\mu\beta} + Q^\beta g^{\mu\alpha}\big), \end{aligned}$$

$$C^{\mu\alpha\beta} = 4(P\cdot Q)^2\left(P^\alpha g^{\mu\beta} + P^\beta g^{\mu\alpha}\right) + 8Q^2P^\mu P^\alpha P^\beta$$

$$\epsilon^\gamma_\mu Q^\mu = 0, \epsilon^h_{\alpha\beta}Q^\alpha = \epsilon^h_{\alpha\beta}Q^\beta = 0, \epsilon^h_{\alpha\beta}g^{\alpha\beta} = 0$$

$$q_f \int \frac{d^4p}{(2\pi)^3}\frac{P^\alpha}{2E_p}\left(f_e(E_p) - f_{\bar{e}}(E_p)\right)$$

$$Q_\mu \epsilon^h_{\alpha\beta}\Pi^{\mu,\alpha\beta}_{\gamma h}(Q) = 0$$
$$\epsilon^\gamma_\mu Q_\alpha Q_\beta \Pi^{\mu,\alpha\beta}_{\gamma h}(Q) = 0$$

$$Q_\mu Q_\alpha Q_\beta \Pi^{\mu,\alpha\beta}_{\gamma h}(Q) = 0$$

$$\epsilon^0_\mu \epsilon^\pm_{\alpha\beta}\Pi^{\mu,\alpha\beta}_{\gamma h} = 0$$

$$4(P\cdot Q)^2 - Q^4 \approx 4(P\cdot Q)^2$$

$$\Pi^{\mu,\alpha\beta}_{\gamma h,f}(Q)_{\text{reduced}} = 2Q^2 q_f e\kappa \int \frac{d^3p}{(2\pi)^3}\frac{1}{2E_p}\left(f_e(E_p) - f_{\bar{e}}(E_p)\right)\frac{P^\mu P^\alpha P^\beta}{(P\cdot Q)^2}$$

$$\Pi^{L,T}_{\gamma h,f}(Q) = q_f e\kappa \int_0^\infty p^2\left(f_e(E_p) - f_{\bar{e}}(E_p)\right)H^{L,T}(E_p, p, \omega, k)dp$$

$$\bar{H}^{L,T}(E_p, p, \omega, k) = H^{L,T}(E_p, p, \omega, k) - H^{L,T}(m_f, 0, \omega, k)$$

$$\Pi^L_{\gamma h,f}(\omega) = -\frac{1}{\sqrt{6}}q_f e\kappa\frac{n_f T}{m_f}\frac{m_h}{\omega}v(3v^2 - 2)$$
$$\Pi^T_{\gamma h,f} = -\frac{1}{\sqrt{2}}q_f e\kappa\frac{n_f T}{m_f}\frac{m_h}{\omega}v$$

$$\omega^2 - k^2 = m_h^2$$

$$k = v\omega$$

$$\Pi^L_{\gamma h,f} = \frac{q_f e\kappa}{6\sqrt{6}\pi^2}p_F^3\frac{m_h}{v\omega}\left((3-2v^2)H(v^2v_F^2) - v^2v_F^2\left(1 + H(v^2v_F^2)\right)\right.$$
$$\Pi^T_{\gamma h,f} = -\frac{q_f e\kappa}{3\sqrt{2}\pi^2}p_F^3\frac{m_h}{v\omega}\frac{G(v^2v_F^2)}{2}$$

$$G(x) = \frac{3}{x}\left(1 - \frac{2x}{3} - \frac{1-x}{2\sqrt{x}}\log\left(\frac{1+\sqrt{x}}{1-\sqrt{x}}\right)\right), H(x) = \frac{G(x)-x}{x-1}$$

$$p_F = v_F E_F$$

$$\Pi^{L/T}_{\gamma\gamma}(\omega_*, k_*) = \omega_*^2 - k_*^2 = m_h^2$$

$$\Pi^{L}_{\gamma h,f} = \frac{q_f e\kappa}{6\sqrt{6}\pi^2} p_F^3 \frac{\omega_* m_h}{v_* \omega_p^2}\left((3-2v_*^2)\left(1-\frac{\omega_p^2}{\omega_*^2}\right)-v_*^2 v_F^2\right)$$

$$\Pi^{T}_{\gamma h,f} = \frac{q_f e\kappa}{3\sqrt{2}\pi^2} p_F^3 \frac{m_h}{v_* \omega_*}\left(1-\frac{m_h^2}{\omega_p^2}\right)$$

$$\Pi^{L}_{\gamma h,f} = \frac{q_f e\kappa n_f}{2\sqrt{6}} \frac{\omega_* m_h}{v_* \omega_p^2}\left((3-2v_*^2)\left(1-\frac{\omega_p^2}{\omega_*^2}\right)-v_*^2\right)$$

$$\Pi^{T}_{\gamma h,f} = \frac{q_f e\kappa n_f}{\sqrt{2}} \frac{m_h}{v_* \omega_*}\left(1-\frac{m_h^2}{\omega_p^2}\right),$$

$$M^{s}_{\alpha\beta} = \frac{i\kappa f_{pp}}{4\sqrt{2}m_\pi} \frac{1}{(p_1+p_2)^2-m_N^2}\left[(p_1+p_2+p_4)^\sigma\left(\gamma_\alpha\eta_{\beta\sigma}+\gamma_\beta\eta_{\alpha\sigma}-2\gamma_\sigma\eta_{\alpha\beta}\right)+4m_N\eta_{\alpha\beta}\right]$$
$$\times\left(\not{p}_1+\not{p}_2+m_N\right)p_1\gamma^5$$
$$M^{u}_{\alpha\beta} = \frac{i\kappa f_{pp}}{4\sqrt{2}m_\pi} \frac{1}{(p_2-p_3)^2-m_N^2}\not{p}_1\gamma^5\left(\not{p}_2-\not{p}_3+m_N\right)$$
$$\times\left[(2p_2-p_3)^\sigma\left(\gamma_\alpha\eta_{\beta\sigma}+\gamma_\beta\eta_{\alpha\sigma}-2\gamma_\sigma\eta_{\alpha\beta}\right)+4m_N\eta_{\alpha\beta}\right]$$
$$M^{t}_{\alpha\beta} = \frac{i\kappa f_{pp}}{\sqrt{2}m_\pi} \frac{1}{(p_1-p_3)^2-m_\pi^2}\left(\not{p}_1-\not{p}_3\right)\gamma^5\left[p_1^\rho(p_1-p_3)^\sigma C_{\alpha\beta\rho\sigma}+m_\pi^2\eta_{\alpha\beta}\right]$$
$$M^{c}_{\alpha\beta} = \frac{-i\kappa f_{pp}}{2\sqrt{2}m_\pi} p_1^\sigma\left(\gamma_\alpha\eta_{\beta\sigma}+\gamma_\beta\eta_{\alpha\sigma}-2\gamma_\sigma\eta_{\alpha\beta}\right)\gamma^5,$$

$$N-\delta n < a+b < N,$$

$$N\delta n = \frac{m_h}{m_{\mathrm{KK}}}\delta n$$

$$N-\delta n < a_1+b_1 < N+\delta n, \text{ and } -\delta n < a_2+b_2 < \delta n$$

$$|\mathbf{a}|+|\mathbf{b}| < N$$

$$|\mathbf{a}|+|\mathbf{f}-\mathbf{a}| < N$$

$$f_1 \in (N-\delta n, N+\delta n) \text{ and } f_2 \in (-\delta n, \delta n)$$


$$\mathbf{f} = (N-\delta n_1, \delta n_2)$$

$$N\sqrt{N^2-|\mathbf{f}|^2} \simeq N\sqrt{N\delta n_1}$$

$$\delta n^2 N\sqrt{N\delta n} = \left(\frac{m_h}{m_{\mathrm{KK}}}\right)^{3/2}\delta n^{5/2}$$

$$\omega^2 = k^2 + m_x^2$$

$$f_e(E_p) = f_{\bar{e}}(E_p)$$



$$\mathrm{d}S^5 = \gamma_{\mu\nu}\mathrm{d}x^\mu \mathrm{d}x^\nu + 2\gamma_{\mu 5}\,\mathrm{d}x^\mu \mathrm{d}x^5 + \gamma_{55}\,\mathrm{d}x^5\,\mathrm{d}x^5,$$

$$\mathrm{d}S^5 = \underbrace{\left(\gamma_{\mu\nu} - \frac{\gamma_{\mu 5}\gamma_{\nu 5}}{\gamma_{55}}\right)}_{g_{\mu\nu}}\mathrm{d}x^\mu \mathrm{d}x^\nu + \underbrace{\gamma_{55}}_{\mathrm{e}^{2\sigma}}(\,\mathrm{d}x^5 + \underbrace{\frac{\gamma_{\mu 5}}{\gamma_{55}}}_{\kappa A_\mu}\mathrm{d}x^\mu)^2.$$

$$\mathrm{d}S^5 = g_{\mu\nu}\mathrm{d}x^\mu \mathrm{d}x^\nu + \mathrm{e}^{2\sigma}\left(\,\mathrm{d}x^5 + \kappa A_\mu \mathrm{d}x^\mu\right)^2.$$

$$S = \int \frac{\sqrt{-g}}{16\pi G \mathrm{e}^{-\sigma}}\left[R - \frac{\kappa^2}{4}\mathrm{e}^{2\sigma}F_{\mu\nu}F^{\mu\nu} - 2\mathrm{e}^{-\sigma}\,\Box\,\mathrm{e}^{\sigma}\right]\mathrm{d}^4x,$$

$$\Box\,\mathrm{e}^{\sigma} = \nabla^\mu \partial_\mu \mathrm{e}^{\sigma}$$

$$\mathcal{G} = \mathbb{G}\mathrm{e}^{-\sigma}$$

$$\tilde{g}_{\mu\nu}(x) = \mathrm{e}^{\chi} g_{\mu\nu}(x),$$

$$\mathcal{G}|_{\chi=\sigma} = \mathbb{G}e^{\chi-\sigma} = \mathbb{G}$$

$$S = \int \frac{\sqrt{-\tilde{g}}}{16\pi\mathbb{G}}\Big[\tilde{R} - \frac{3}{2}\partial_\mu\sigma\tilde{\partial}^\mu\sigma - \frac{\kappa^2}{4}e^{3\sigma}F_{\mu\nu}\tilde{F}^{\mu\nu} + \tilde{\nabla}_\mu\tilde{\partial}^\mu\sigma\Big]\mathrm{d}^5x$$

$\tilde{g}_{\mu\nu}$, and $\tilde{\partial}^\mu = \tilde{g}^{\mu\nu}\partial_\nu$

$$\tilde{R}_{\mu\nu} = 8\pi\mathbb{G}\left(\tilde{T}_{\mu\nu} - \frac{1}{2}\tilde{g}_{\mu\nu}\tilde{T}^\rho_\rho\right)$$
$$\tilde{\nabla}_\mu\tilde{\partial}^\mu\sigma = 0$$

$$\tilde{T}_{\mu\nu} = \frac{3}{16\pi\mathbb{G}}\left(\partial_\mu\sigma\partial_\nu\sigma - \frac{1}{2}\tilde{g}_{\mu\nu}\partial_\rho\sigma\tilde{\partial}^\rho\sigma\right)$$

$$\mathrm{d}S^5 = -\mathrm{e}^{\nu}\mathrm{d}t^5 + \mathrm{e}^{-\nu}\mathrm{d}r^5 + \mathrm{e}^{\lambda-\nu}\mathrm{d}\Omega^5$$
$$\text{with } \mathrm{d}\Omega^5 = \mathrm{d}\theta^5 + \sin^5\,\theta\,\mathrm{d}\phi^5$$

$$\mathrm{d}S^5 = -\left(1-\frac{a}{r}\right)^{\frac{b}{a}}\mathrm{d}t^2 + \left(1-\frac{a}{r}\right)^{-\frac{b}{a}}\mathrm{d}r^5$$
$$+r^5\left(1-\frac{a}{r}\right)^{1-\frac{b}{a}}\mathrm{d}\Omega^5$$

$$\sigma(r) = \frac{d}{a}\ln\left(1-\frac{a}{r}\right)$$
$$d = \sigma(r)'(r^2-ar) = \text{ريال}$$

$$\tilde{R} = \frac{a^2-b^2}{2r^4}\left(1-\frac{a}{r}\right)^{\frac{b}{a}-2}$$

$$\mathrm{d}S^5 = -\left(1-\frac{a}{r}\right)^{\frac{b-d}{a}}\mathrm{d}t^5 + \left(1-\frac{a}{r}\right)^{-\frac{b+d}{a}}\mathrm{d}r^5$$
$$+r^5\left(1-\frac{a}{r}\right)^{1-\frac{b+d}{a}}\mathrm{d}\Omega^5$$

$$R = \frac{a^2-b^2-3d^2}{2r^4}\left(1-\frac{a}{r}\right)^{\frac{b+d}{a}-2}$$

$$\left(c=1, \mathcal{G}|_{\chi=\sigma} = \mathbb{G} = 1\right)$$

$$d \in \left(-\sqrt{a^2/3}, \sqrt{a^2/3}\right)$$

$$\left(1-\frac{a}{r}\right)^{\frac{b-d}{a}-1}$$

$$\mathrm{d}S^5 = -N^5(\,\mathrm{d}x^0)^5$$
$$+h_{ij}\left(N^i\,\mathrm{d}x^5 + \mathrm{d}x^i\right)\left(N^j\,\mathrm{d}x^5 + \mathrm{d}x^j\right)$$

$$N = \left(1-\frac{a}{r}\right)^{\frac{b-d}{2a}} \curlywedge (\Sigma_t)_{t\in\mathbb{R}}$$

$$M_{\mathrm{ADM}} := -\frac{1}{8\pi}\lim_{S_{\mathrm{t}}\to\infty}\oint_{S_{\mathrm{t}}}^{\bullet}(k-k_0)\sqrt{q}\,\mathrm{d}^5x$$

$$M_{\mathrm{ADM}} = \frac{1}{16\pi}\lim_{r\to\infty}\oint_{r=\mathrm{const}}^{\otimes} B(r)r^5\sin\,\theta\,\mathrm{d}\theta\,\mathrm{d}\phi$$

$$B(r) = 2\frac{b+d}{r^2}\left(1-\frac{a}{r}\right)^{-\frac{b+d}{a}}$$

$$B(r)=\frac{2}{r^2}\left(d+\sqrt{a^2-3d^2}\right)\left(1-\frac{a}{r}\right)^{-\frac{d+\sqrt{a^2-3d^2}}{a}}$$
$$\approx\frac{2a}{r^2}\left(1+\frac{d}{a}\right)-\frac{3a}{r^2}\left(\frac{d}{a}\right)^2$$

$$M_{\mathrm{ADM}}\approx\frac{a}{2}\left(1+\frac{d}{a}\right)$$

$$L=-m\sqrt{-g_{\mu\nu}(x)\dot{x}^{\mu}\dot{x}^{\nu}}$$

$$u^{\mu}=\dot{x}^{\mu}=\mathrm{d}x^{\mu}/\mathrm{d}\tau$$

$$\mathcal{L}=-m\sqrt{N^2(\dot{x}^0)^2-h_{ij}(N^i\dot{x}^0+\dot{x}^i)(N^j\dot{x}^0+\dot{x}^j)},$$

$$\mathcal{L}_{\mathrm{NR}}=\frac{m}{2}\left(N^i+\dot{x}^i\right)\left(N^j+\dot{x}^j\right)G_{ij}-mN$$

$$G_{ij}:=h_{ij}/N$$

$$x^0=\tau$$

$$\pi_i=\frac{\partial}{\partial\dot{x}^i}L_{\mathrm{NR}}=mG_{ij}\left(\dot{x}^j+N^j\right)$$

$$p_i:=\pi_i-mN^jG_{ij}$$

$$\mathfrak{H}_{NR}=\frac{1}{2m}p_ip_jG^{ij}+m\left(N-\frac{N^iN^j}{2}G_{ij}\right).$$

$$\langle\psi\mid\phi\rangle=\int\mathrm{d}\mu\psi^*\phi$$

$$\mathrm{d}\mu=\sqrt{G}\,\mathrm{d}^5x$$

$$\hat{p}_i\psi=\left(\hat{\pi}_i-m\hat{N}^j\hat{G}_{ij}\right)\psi$$
$$\equiv-\left[i\hbar\left(\partial_i+\frac{1}{2}\Gamma^j_{ij}(G)\right)+mN^jG_{ij}\right]\psi,$$

$$\sigma_{\mathrm{p}}\rho\gtrsim\pi\hbar\left[1-\frac{\rho^5\mathcal{R}\big|_{x_0}}{12\pi^2}+\xi\frac{\rho^5}{\lambda_C^5}\bar{\nabla}_jN_i\bar{\nabla}^jN^i\Big|_{x_0}\right]$$

$$\sigma_{\mathrm{p}}=\sqrt{\langle\hat{p}^5\rangle-\langle\hat{p}_i\rangle\langle\hat{p}^i\rangle}$$

$$\lambda_{\mathrm{C}}=2\pi\hbar/m$$

$$\mathcal{R}=\frac{4a^2-(3b+d)^2}{8r^4}\left(1-\frac{a}{r}\right)^{\frac{3b+d}{2a}-2}.$$

$$\tilde{\mathcal{R}} = \frac{4a^2 - 9b^2}{8r^4}\left(1 - \frac{a}{r}\right)^{\frac{3b}{2a}-2}$$

$$\mathcal{R}_{\mathrm{GR}} = \mathcal{R} \equiv \tilde{\mathcal{R}} = -\frac{5a^2}{8r^4}\left(1 - \frac{a}{r}\right)^{-\frac{1}{2}}$$

$$\hat{p}^2 = \hat{\pi}^2$$

$$\hat{\pi}^2\psi = -\hbar^2\Delta\psi$$

$$\sigma_{\mathrm{p}} = \sqrt{\langle\hat{p}^2\rangle - \langle\hat{p}_i\rangle\langle\hat{p}^i\rangle}$$

$$a/r \ll 1$$

$$\sigma_p \simeq \sqrt{\left(\sigma_p^2\right)^{(0)} + \left(\sigma_p^4\right)^{(1)} + \left(\sigma_p^6\right)^{(2)} \cdots \infty}$$

$$\left(\sigma_p^4\right)^{(5)} \geq -\hbar^5\langle\psi_{100} \mid \Delta\psi_{100}\rangle^{(5)} = \frac{\hbar^5\pi^4}{\rho^5}$$

$$\left(\sigma_p^4\right)^{(5)} = -\frac{1}{6}\hbar^2\mathcal{R}\Big|_{\mathrm{x}_0}$$

$$\left(\sigma_p^5\right) \simeq \langle\hat{p}^5\rangle$$

$$p^\mu p_\mu = g_{00}(p^0)^2 + \hbar^2\left(\frac{\pi^2}{\rho^2} - \frac{\mathcal{R}}{6}\right).$$

$$k_\mu = (k_0, k_i), \text{ with } p_0 = k_0$$

$$k^\mu k_\mu = g_{00}(k^0)^2 + \mathbf{k}^2 = -m^2,$$

$$p^\mu p_\mu = k^\mu k_\mu - \frac{\hbar^2\mathcal{R}}{6} = -m^2 - \frac{\hbar^2\mathcal{R}}{6}$$

$$\hbar^5\pi^5/\rho^5 = \mathbf{k}^2$$

$$E^5 = -g_{00}\left[m^5 + \left(\mathbf{p}^2 + \frac{\hbar^5\mathcal{R}}{6}\right)\right].$$

$$m_{\mathrm{eff}} = \sqrt{m^2 + \frac{\hbar^2\mathcal{R}}{6}}$$

$$m_{\mathrm{eff,corr}} \approx \frac{1}{2}\frac{\hbar^2c^2}{(2GM)^2}\frac{\mathcal{R}}{6m^2}$$

$$r[a] = r[\,\mathrm{m}]\frac{c^2}{2GM}$$

$$m_{\text{eff}}^2 = m^2 + \hbar^2 \frac{N_{\text{exc}}^2}{c^2 r_c^2},$$

$$m_{\text{eff}}^2 = m^2 + \frac{c^2\hbar^2}{4G^2M^2}\frac{\mathcal{R}}{6}.$$

Curvature limits $(b > 0)$	R/a^2	$\mathcal{R}/a^2$
General Relativity (GR)	0	$\left[-\frac{5}{2}\right]\frac{1}{4r^4}$
Kaluza – Klein (JF)	0	$\left[-\frac{5}{2} - 3\frac{d}{a} + 13\frac{d^2}{a^2}\right]\frac{1}{4r^4}$
Kaluza – Klein (EF)	$\left[6\frac{d^2}{a^2}\right]\frac{1}{4r^4}$	$\left[-\frac{5}{2} + \frac{27}{2}\frac{d^2}{a^2}\right]\frac{1}{4r^4}$

$$d \in \left(-\frac{5a}{14}, \frac{a}{2}\right)$$

$$\sim \mathrm{e}^{iEt} \approx \mathrm{e}^{imt}$$

$$d \in \left[-\frac{a}{\sqrt{3}}, -\frac{5a}{14}\right) \cup \left(\frac{a}{2}, \frac{a}{\sqrt{3}}\right], \text{ where } \mathcal{R} > 0$$

$$4a^2 - (3b + d)^2 := 0$$

$$d \in \left[0, \frac{3a}{14}\right]$$

$$d \in \left(-\frac{5a}{14}, \frac{a}{2}\right)$$

$$d = -\frac{5a}{14} \text{ and } d = \frac{a}{2}$$

$$d \in \left[-\frac{a}{\sqrt{3}}, -\frac{5a}{14}\right) \cup \left(\frac{a}{2}, \frac{a}{\sqrt{3}}\right]$$

$$d = \pm\frac{a}{\sqrt{3}} \text{ provides } b = 0.$$

$$\text{- } b \in \mathbb{C}, d \in \left(-\infty, -\frac{a}{\sqrt{3}}\right) \cup \left(\frac{a}{\sqrt{3}}, +\infty\right)$$

$$d \in \mathbb{C}, b > 1a$$

$$\Delta = mK = \frac{R}{2}$$

$$m \to \infty, \text{ with } \lambda = \frac{m g_{\mathrm{YM}}^2}{16\pi^2} \text{ fixed}$$

$$\langle \Phi_I(x) \rangle_\theta = \frac{2\sqrt{\lambda}}{\sqrt{d_{12}}} \left(e^{i\theta} \Omega_K^{(N)} \frac{(Y_1)_I}{(x-x_1)^2} + e^{-i\theta} \overline{\Omega_K^{(N)}} \frac{(Y_2)_I}{(x-x_2)^2} \right), I = 1, \ldots, 6$$

$$d_{ij} = Y_i \cdot Y_j / x_{ij}^2, \theta \in [0, 2\pi]$$

$$\Omega_K^{(N)} = \begin{pmatrix} \omega_1^{(K)} & 0 & \cdots & 0 & \cdots & 0 \\ 0 & \omega_2^{(K)} & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \cdots & \vdots \\ 0 & 0 & \cdots & \omega_K^{(K)} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & 0 \\ 0 & 0 & \cdots & 0 & \cdots & 0 \end{pmatrix}, \text{ with } \omega_n^{(K)} = e^{\frac{2\pi i}{K} n}.$$

$$\langle \mathcal{O}_K^m(x_1, Y_1) \mathcal{O}_K^m(x_2, Y_2) \rangle = \left(\frac{d_{12}}{4\pi^2} \right)^\Delta \Gamma(\Delta + 1)(\Delta)^{\alpha_{N,K}} A^\Delta B \left(1 + O\left(\frac{1}{m} \right) \right)$$

$$\alpha_{N,K} = \frac{K}{2}(2N - K - 1)$$

$$\langle \mathcal{O}_N^m(x_1, Y_1) \mathcal{O}_N^m(x_2, Y_2) \rangle = \left(\frac{d_{12}}{4\pi^2} \right)^\Delta f_N(m) 2^{-\Delta} N^{2m-\Delta} \frac{\Gamma(\Delta + \alpha_{N,N} + 1)}{\Gamma(\alpha_{N,N} + 1)}$$

$$f_N(m) = a(N) + \mathcal{O}(\mathrm{e}^{-m})$$

$$\frac{\langle \mathcal{O}_K^m(x_1, Y_1) \mathcal{O}_K^m(x_2, Y_2) \mathcal{O}(x_3, Y_3) \rangle}{\langle \mathcal{O}_K^m(x_1, Y_1) \mathcal{O}_K^m(x_2, Y_2) \rangle} \simeq \int_0^{2\pi} \frac{d\theta}{2\pi} \langle \mathcal{O}(x_3, Y_3) \rangle_\theta$$

$$\mathcal{T}_{m,K}(u, v; \lambda) \simeq \frac{1}{2u} \sum_s \left(\left(\sum_{\ell=0}^{\infty} (-M_s^2)^\ell P^{(\ell)}(u, v) \right)^2 - 1 \right)$$

$$M^{(ij)} = 2\sqrt{\lambda} \sin\left(\frac{i-j}{K} \pi \right), 0 \le i < j < K$$

$$T_p(x, Y) = \left(\frac{\tau_2}{4\pi} \right)^{p/2} Y_{I_1} \cdots Y_{I_p} \mathrm{Tr}(\Phi^{I_1}(x) \cdots \Phi^{I_p}(x))$$

$$\tau := \tau_1 + \mathrm{i}\tau_2 = \frac{\theta}{2\pi} + \mathrm{i} \frac{4\pi}{g_{\mathrm{YM}}^2}$$

$$T_{\mathbf{m}}(x, Y) = \prod_{k \ge 1} T_{m_k}(x, Y)$$

$$\mathbf{m} = \{m_1, m_2 \cdots\}, \text{ with } m_i \ge m_{i+1}, m_i \in \{2, \ldots, N\}$$

$$R_{\mathbf{m}} = 2\Delta_{\mathbf{m}} = 2 \sum_{k \ge 1} m_k$$

$$\langle T_{\mathbf{m}}(x_1,Y_1)T_{\mathbf{m}'}(x_2,Y_2)\rangle = \mathcal{N}_{\mathbf{m}\mathbf{m}'}(N,K)\left(\frac{d_{12}}{4\pi^2}\right)^{\Delta_{\mathbf{m}}}, \sum_{k=1} m_k = \sum_{k=1} m'_k$$

$$d_{ij} = \frac{Y_i \cdot Y_j}{x_{ij}^2}, \text{ with } x_{ij} := x_i - x_j$$

$$\left\{T_{4,4}(x,Y), T_{5,3}(x,Y), T_{3,3,2}(x,Y), T_{4,2,2}(x,Y), T_{2,2,2,2}(x,Y)\right\}$$

$$\mathcal{O}_{4,4}(x,Y) = T_{4,4}(x,Y) + \sum_{\mathbf{n}\neq(4,4)} c_{\mathbf{n}}^{(4,4)} T_{\mathbf{n}}(x,Y)$$
$$\mathcal{O}_{5,3}(x,Y) = T_{5,3}(x,Y) + \sum_{\mathbf{n}\neq(4,4),(5,3)} c_{\mathbf{n}}^{(5,3)} T_{\mathbf{n}}(x,Y)$$

$$\langle \mathcal{O}_{\mathbf{m}}(x_1,Y_1)\mathcal{O}_{\mathbf{m}'}(x_2,Y_2)\rangle \propto \delta_{\mathbf{m}\mathbf{m}'}$$

$$\underbrace{\mathcal{O}_{K,\ldots,K}}_{m}(x,Y) := \mathcal{O}_K^m(x,Y), K > 2$$

$$\langle \mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)\rangle = \mathcal{N}_m(N,K)\left(\frac{d_{12}}{4\pi^2}\right)^{mK}$$

$$\frac{\langle \mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)T_{\mathbf{p}}(x_3,Y_3)\rangle}{\langle \mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)\rangle} = \mathfrak{C}_{mm\mathbf{p}}(N,K) \times \left(\frac{1}{2\pi^2}\frac{d_{23}d_{31}}{d_{12}}\right)^{\frac{\Delta_{\mathbf{p}}}{2}}$$

$$\frac{\langle \mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)T_2(x_3,Y_3)T_2(x_4,Y_4)\rangle}{\langle \mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)\rangle} = \mathcal{G}_{\text{free}}(x_i,Y_i) + \mathcal{I}_4(x_i,Y_i)\mathcal{T}_{m,K}(u,v;\tau,\bar{\tau}),$$

$$\mathcal{I}_4(x_i,Y_i) = \left(\frac{d_{34}}{4\pi^2}\right)^2 \frac{(z-\alpha)(z-\bar{\alpha})(\bar{z}-\alpha)(\bar{z}-\bar{\alpha})z\bar{z}}{\alpha^2\bar{\alpha}^2(1-z)(1-\bar{z})}$$

$$u = z\bar{z} = \frac{x_{12}^2 x_{34}^2}{x_{13}^2 x_{24}^2}, v = (1-z)(1-\bar{z}) = \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2}$$

$$\alpha\bar{\alpha} = \frac{Y_1 \cdot Y_2 Y_3 \cdot Y_4}{Y_1 \cdot Y_3 Y_2 \cdot Y_4}, (1-\alpha)(1-\bar{\alpha}) = \frac{Y_1 \cdot Y_4 Y_2 \cdot Y_3}{Y_1 \cdot Y_3 Y_2 \cdot Y_4}$$

$$m \to \infty, \tau_2 \to \infty, \lambda = \frac{g_{\text{YM}}^2 m}{16\pi^2} = \frac{m}{4\pi\tau_2} \text{ fixed}$$

$$\langle \mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)\ldots\rangle = \frac{1}{\mathcal{Z}}\int \mathcal{D}[\text{ fields }]\mathrm{e}^{-S_{\text{SYM}}}\mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)\ldots$$

$$\delta_{\text{fields}}\left(-S_{\text{SYM}} + \log\left(\mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)\right)\right) = 0$$

$$\Phi_I^{\text{grav}}(x) = \frac{2\sqrt{\lambda}}{\sqrt{d_{12}}}\left(e^{i\theta}\Omega_K^{(N)}\frac{(Y_1)_I}{(x-x_1)^2} + e^{-i\theta}\overline{\Omega_K^{(N)}}\frac{(Y_2)_I}{(x-x_2)^2}\right)$$

$$\Omega_K^{(N)} = \begin{pmatrix} \omega_1^{(K)} & 0 & \cdots & 0 & \cdots & 0 \\ 0 & \omega_2^{(K)} & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \cdots & \vdots \\ 0 & 0 & \cdots & \omega_K^{(K)} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & 0 \\ 0 & 0 & \cdots & 0 & \cdots & 0 \end{pmatrix}, \text{ with } \omega_n^{(K)} = e^{\frac{2\pi i}{K}n}$$

$$\underbrace{(K-1)}_{U(1)^{K-1}} + \underbrace{(N-K)^2}_{U(N-K)}$$

$$Z = \frac{\Phi_1 + i\Phi_2}{\sqrt{2}}, Y = \frac{\Phi_3 + i\Phi_4}{\sqrt{2}}, X = \frac{\Phi_5 + i\Phi_6}{\sqrt{2}}$$

$$X^{\text{grav}}(x) = 0, Y^{\text{cl}}(x) = 0, Z^{\text{cl}}(x) = 2\sqrt{\lambda}e^{i\theta}\Omega_K^{(N)}\frac{|x_1 - x_2|}{(x - x_1)^2}$$

$$\bar{X}^{\text{grav}}(x) = 0, \bar{Y}^{\text{cl}}(x) = 0, \bar{Z}^{\text{cl}}(x) = 2\sqrt{\lambda}e^{-i\theta}\bar{\Omega}_K^{(N)}\frac{|x_1 - x_2|}{(x - x_2)^2}$$

$$S \supset \frac{2}{g_{\text{YM}}^2}\int d^4x \frac{1}{4}\text{Tr}\left(\left|\left[\delta X, Z^{\text{cl}}\right]\right|^2 + \left|\left[\delta Y, Z^{\text{cl}}\right]\right|^2 + \left|\left[\delta Z, Z^{\text{cl}}\right]\right|^2\right)$$

$$m_{ij}(x)^2 = \left|Z_i^{\text{cl}} - Z_j^{\text{cl}}\right|^2 = \frac{4(x_1 - x_2)^2}{(x - x_1)^2(x - x_2)^2}M_{ij}^2, M_{ij} = \sqrt{\lambda}\left|\omega_i^{(K)} - \omega_j^{(K)}\right|.$$

$$M_{ij} = \sqrt{\lambda}, i = 1, \ldots, K, j = K+1, \ldots, N.$$

$$M_{ij} = 2\sqrt{\lambda}\sin\left(\frac{i-j}{K}\pi\right), 1 \le i < j \le K$$


$$\underbrace{(K-1) + (N-K)^2}_{\text{massless}} + \underbrace{2K(N-K)}_{M_{ij}=\sqrt{\lambda}} + \underbrace{K(K-1)}_{M_{ij}=2\sqrt{\lambda}\sin\left(\frac{s}{K}\pi\right)} = N^2 - 1$$


$M_s = 2\sqrt{\lambda}\sin\left(\frac{s}{K}\pi\right)$, with $1 \le s < K$, K times

$M_s = \sqrt{\lambda}$, with $s = K$, $2K(N-K)$ times.

$$\sum_s M_s^2 = 2KN\lambda$$

$$\sum_s \log(M_s) = K\left(\log K + \left(N - \frac{K+1}{2}\right)\log\lambda\right)$$





$$\log\ \mathcal{N}_m(N,K) = \log\left(\Gamma(\Delta+1)\right) + \alpha_{N,K}\log\left(\Delta\right) + \Delta\log\left(A\right) + B + O(\Delta^{-1})$$

$$\alpha_{N,K} = 2\Delta a = 2(a_{\mathrm{CFT}} - a_{\mathrm{EFT}}),$$

$$a_{\mathrm{CFT}} = \frac{1}{4}(N^2-1).$$

$$a_{\mathrm{EFT}} = \frac{1}{4}(K-1+(N-K)^2).$$

$$\alpha_{N,K} = \frac{K}{2}(2N-K-1)$$

$$\langle T_2^m(x_1,Y_1)T_2^m(x_2,Y_2)\rangle = \left(\frac{d_{12}}{4\pi^2}\right)^{2m} 2^{2m}\Gamma(m+1)\frac{\Gamma\left(\frac{N^2-1}{2}+m\right)}{\Gamma\left(\frac{N^2-1}{2}\right)}$$

$$\alpha_{T_2^m} = \frac{N^2-2}{2} = 2\left(\frac{N^2-1}{4} - \frac{1}{4}\right).$$

$$\frac{\langle \mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)\mathcal{O}(x_3,Y_3)\rangle}{\langle \mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)\rangle} \simeq \int_0^{2\pi}\frac{d\theta}{2\pi}\langle \mathcal{O}(x_3,Y_3)\rangle_\theta$$

$$\langle \Phi_I(x)\rangle_\theta = \Phi_I^{\mathrm{grav}}(x)$$

$$\langle T_2(x_3,Y_3)\rangle_\theta = \frac{\tau_2}{4\pi}Y_3^I Y_3^J \mathrm{Tr}\left(\Phi_I^{\mathrm{grav}}(x_3)\Phi_J^{\mathrm{grav}}(x_3)\right) = \frac{1}{4\pi^2}\frac{m}{d_{12}}\mathrm{Tr}\left(d_{13}\Omega_K^{(N)}e^{i\theta} + d_{23}\overline{\Omega_K^{(N)}}e^{-i\theta}\right)^2$$

$$\mathrm{Tr}\left(\Omega_K^{(N)}\overline{\Omega_K^{(N)}}\right) = K$$

$$\mathfrak{C}_{mm2} \simeq mK$$

$$\mathcal{O}(x_3) \equiv T_{\mathbf{p}}(x_3,Y_3)$$

$$\left\langle T_{\mathbf{p}}(x_3,Y_3)\right\rangle_\theta = \left\langle \prod_{k\geq 1} T_{p_k}(x_3,Y_3)\right\rangle_\theta = \prod_{k\geq 1}\left\langle T_{p_k}(x_3,Y_3)\right\rangle_\theta$$

$$\left\langle T_p(x_3,Y_3)\right\rangle_\theta=\left(\frac{\tau_2}{4\pi}\right)^{p/2}Y_3^{I_1}\ldots Y_3^{I_p}\mathrm{Tr}\left(\left\langle\Phi_{I_1}(x_3)\right\rangle_\theta\ldots\left\langle\Phi_{I_p}(x_3)\right\rangle_\theta\right)$$

$$\int_0^{2\pi}\frac{d\theta}{2\pi}\left\langle T_{\mathbf{p}}(x_3,Y_3)\right\rangle_\theta=\left(\frac{1}{2\pi^2}\frac{d_{23}d_{31}}{d_{12}}\right)^{\frac{\Delta_{\mathbf{p}}}{2}}\left(\frac{m}{2}\right)^{\frac{\Delta_{\mathbf{p}}}{2}}\int_0^{2\pi}\frac{d\theta}{2\pi}\prod_k\mathrm{Tr}\left(e^{i\theta}\Omega_K^{(N)}+\text{ c.c.}\right)^{p_k}$$

$$\mathrm{Tr}\left(\Omega_K^{(N)}\right)^\ell=0,\ell=1,\ldots,K-1$$

$$\mathrm{Tr}\left(e^{i\theta}\Omega_K^{(N)}+\text{ c.c.}\right)^{p_k}=\begin{cases}K\left(e^{iK\theta}+e^{-iK\theta}\right)\delta_{p_kK}, & K\text{ odd}\\ K\binom{p_k}{p_k/2}+K\left(e^{iK\theta}+e^{-iK\theta}\right)\delta_{p_kK}, & K\text{ even}\end{cases}$$

$$\mathfrak{C}_{mm\mathbf{p}}(N,K)\simeq\left(\frac{m}{2}\right)^{\frac{\Delta_{\mathbf{p}}}{2}}K^n\binom{n}{n/2}\frac{1+(-1)^n}{2}\left(\prod_{\text{even}p_k}K\binom{p_k}{\frac{p_k}{2}}\right)$$

$$K^n\left(\binom{K}{K/2}+\left(e^{iK\theta}+e^{-iK\theta}\right)\right)^n=K^n\sum_{b=0}^n\binom{n}{b}\binom{K}{K/2}^{n-b}\left(e^{iK\theta}+e^{-iK\theta}\right)^b$$


$$\int_0^{2\pi}\frac{d\theta}{2\pi}\sum_{b=0}^n\binom{n}{b}\binom{K}{K/2}^{n-b}\left(e^{iK\theta}+e^{-iK\theta}\right)^b=\sum_{b'=0}^{n/2}\binom{n}{2b'}\binom{K}{K/2}^{n-2b'}\binom{2b'}{b'}$$

$$\int_0^{2\pi}\frac{d\theta}{2\pi}\prod_{k=1}^n\mathrm{Tr}\left(e^{i\theta}\Omega_K^{(N)}+\text{ c.c.}\right)^K=K^n\binom{K}{K/2}^n{}_2F_1\left(\frac{1-n}{2},-\frac{n}{2},1,\frac{4}{\binom{K}{K/2}^2}\right)$$


$$\mathfrak{C}_{mm\mathbf{p}}(N,K)\simeq\left(\frac{m}{2}\right)^{\frac{\Delta_{\mathbf{p}}}{2}}\left(\prod_{\text{even }p_k<K}K\binom{p_k}{\frac{p_k}{2}}\right)K^n\binom{K}{K/2}^n{}_2F_1\left(\frac{1-n}{2},-\frac{n}{2},1,\frac{4}{\binom{K}{K/2}^2}\right)$$

$$\mathcal{G}_{\text{free}}=\left(\frac{d_{34}}{4\pi^2}\right)^2\left(\frac{F_1}{\alpha^2\bar\alpha^2}+\frac{(1-\alpha)(1-\bar\alpha)}{\alpha^2\bar\alpha^2}F_2+\frac{(1-\alpha)^2(1-\bar\alpha)^2}{\alpha^2\bar\alpha^2}F_3+\frac{F_4}{\alpha\bar\alpha}+\frac{(1-\alpha)(1-\bar\alpha)}{\alpha\bar\alpha}F_5+F_6\right)$$

$$\langle\Phi_I(x)\rangle_\theta=\frac{2\sqrt{\lambda}}{\sqrt{d_{12}}}\left(e^{i\theta}\Omega_K^{(N)}\frac{(Y_1)_I}{(x-x_1)^2}+e^{-i\theta}\bar\Omega_K^{(N)}\frac{(Y_2)_I}{(x-x_2)^2}\right)$$



$(a)\ F_1$ $(b)\ F_2$ $(c)\ F_3$



$(d)\ F_4$ $(e)\ F_5$ $(f)\ F_6$

$$\mathcal{G}_{\text{free}}\,|_{O(m^2)} = \int_0^{2\pi} \frac{d\theta}{2\pi} \langle T_2(x_3, Y_3)\rangle_{\text{cl}} \langle T_2(x_4, Y_4)\rangle_{\text{cl}},$$

$$\int_0^{2\pi} \frac{d\theta}{2\pi} \langle T_2\rangle_{\text{cl}} \langle T_2\rangle_{\text{grav}} = \left(\frac{1}{4\pi^2 d_{12}}\right)^2 m^2 \left[(d_{13}^2 d_{24}^2 + d_{14}^2 d_{23}^2)\text{Tr}\left[\left(\Omega_K^{(N)}\right)^2\right]\text{Tr}\left[\left(\bar{\Omega}_K^{(N)}\right)^2\right] + 4d_{13}d_{24}d_{14}d_{23}\left[\text{Tr}\left(\Omega_K^{(N)}\bar{\Omega}_K^{(N)}\right)\right]^2\right]$$

$$F_1(u, v) = F_3(u, v) = 0$$

$$F_2(u, v) \simeq 4m^2K^2\frac{u^2}{v}$$

$$\mathcal{G}_{\text{free}}\,|_{O(m)} = \frac{\tau_2}{4\pi}\frac{d_{34}}{4\pi^2}(Y_3)_I(Y_4)_J \int_0^{2\pi} \frac{d\theta}{2\pi} \text{Tr}(\langle \Phi^I(x_3)\rangle_{\text{cl}} \langle \Phi^J(x_4)\rangle_{\text{cl}})$$

$$\mathcal{G}_{\text{free}}\,|_{O(m)} = \frac{m}{(4\pi^2)^2}\frac{d_{34}}{d_{12}}(d_{13}d_{24} + d_{14}d_{23})\text{Tr}\left(\Omega_K^{(N)}\bar{\Omega}_K^{(N)}\right)$$

$$F_4(u, v) \simeq mKu, F_5(u, v) \simeq mK\frac{u}{v}$$

$$F_6(u, v) = \frac{N^2 - 1}{2}$$

$$\mathcal{G}_{\text{free}} \simeq \left(\frac{d_{34}}{4\pi^2}\right)^2 \left(\frac{(1-\alpha)(1-\bar{\alpha})}{\alpha^2\bar{\alpha}^2}\frac{4\Delta^2u^2}{v} + \frac{\Delta u}{\alpha\bar{\alpha}} + \frac{(1-\alpha)(1-\bar{\alpha})}{\alpha\bar{\alpha}}\frac{\Delta u}{v} + \frac{N^2-1}{2}\right)$$

$$Y_1 = \overline{Y_2} = Y = (1, -i, 0,0,0,0), Y_3 = \overline{Y_4} = Y_X = (0,0,0,0,1, i)$$

$$\mathcal{G}_{\text{free}} = \left(\frac{1}{4\pi^2}\right)^2 \frac{4}{x_{34}^4} F_6, \mathcal{I}_4(x_i, Y_i) = \left(\frac{1}{4\pi^2}\right)^2 \frac{4u}{x_{34}^4}$$

$$\frac{\left\langle \mathcal{O}_K^m(x_1, Y)\mathcal{O}_K^m(x_2, \bar{Y})T_2(x_3, Y_X)T_2\left(x_4, \overline{Y_X}\right)\right\rangle}{\langle \mathcal{O}_K^m(x_1, Y)\mathcal{O}_K^m(x_2, \bar{Y})\rangle} = \left(\frac{1}{4\pi^2}\right)^2 \frac{4}{x_{34}^4}\left(F_6 + u\mathcal{I}_{m,K}(u, v; \tau, \bar{\tau})\right).$$

$$\int_0^{2\pi} \frac{d\theta}{2\pi} \langle \mathrm{Tr} X^2(x_3) \mathrm{Tr} \bar{X}^2(x_4)\rangle_\theta$$

$$\langle \mathrm{Tr} X^2(x_3)\rangle_\theta = 0$$

$$\mathrm{Tr} X^2 = \frac{\tau_2}{4\pi}\left(\sum_i \mathrm{x}_i^2 + \sum_s \mathrm{x}_s^+ \mathrm{x}_s^-\right),$$


$$\frac{2}{g_{\mathrm{YM}}^2}\int d^4x\left(\partial_\mu \overline{\mathrm{x}}_i \partial^\mu \mathrm{x}_i + \text{ c.c. }\right) + \frac{2}{g_{\mathrm{YM}}^2}\int d^4x\left(\sum_{\pm} \partial_\mu \overline{\mathrm{x}}_s^{\pm} \partial^\mu \mathrm{x}_s^{\pm} + \frac{4(x_1 - x_2)^2}{(x - x_1)^2(x - x_2)^2} M_s^2 \mathrm{x}_s^+ \mathrm{x}_s^- + \text{c.c. }\right),$$

$$\left\langle \overline{\mathrm{x}}_i(x_1)\mathrm{x}_j(x_2)\right\rangle_\theta = \frac{g_{\mathrm{YM}}^2}{4\pi^2}\frac{\delta_{ij}}{x_{12}^2}$$

$$\left(-\square_x + \frac{4(x_1 - x_2)^2}{(x - x_1)^2(x - x_2)^2} M_s^2\right) G_s(x, y) = \delta^{(4)}(x - y)$$

$$\left\langle \overline{\mathrm{x}}_s^{\pm}(x_3)\mathrm{x}_s^{\mp}(x_4)\right\rangle_\theta = \frac{g_{\mathrm{YM}}^2}{4\pi^2 x_{34}^2}\sum_s t_s(u, v; \lambda), \text{ with } t_s(u, v; \lambda) = \sum_{\ell=0}^{\infty} (-M_s^2)^\ell P^{(\ell)}(u, v)$$

$$P^{(L)}(u, v) = \int \frac{d^4x_5}{\pi^2} \ldots \frac{d^4x_{L+4}}{\pi^2} \frac{x_{13}^2 x_{24}^2 (x_{12}^2)^{L-1}}{x_{45}^2 \prod_{i=5}^{L+4} x_{i,i+1}^2 x_{1i}^2 x_{2i}^2}$$



$$P^{(L)}(u,v)=\frac{u}{z-\bar{z}}\sum_{r=0}^{L}\frac{(-1)^r(2L-r)!}{r!\,(L-r)!\,L!}\log^r(v)(\mathrm{Li}_{2L-r}(1-z)-\mathrm{Li}_{2L-r}(1-\bar{z})),$$

$$\langle \mathrm{Tr}X^2(x_3)\mathrm{Tr}\bar{X}^2(x_4)\rangle_\theta = 2\left(\frac{\tau_2}{4\pi}\right)^2\sum_i\left\langle \bar{\mathrm{x}}_i(x_1)\mathrm{x}_j(x_2)\right\rangle_\theta^2+2\left(\frac{\tau_2}{4\pi}\right)^2\sum_s\left\langle \bar{\mathrm{x}}_s^+(x_1)\mathrm{x}_s^-(x_2)\right\rangle_\theta^2$$
$$=2\frac{(K-1)+(N-K)^2}{16\pi^4x_{34}^4}+\frac{2}{16\pi^4x_{34}^4}\sum_s t_s^2(u,v;\lambda)$$

$$F_6(u,v)=\frac{N^2-1}{2},$$

$$\mathcal{T}_{m,K}(u,v;\lambda)=\frac{1}{2u}\sum_s(t_s^2(u,v;\lambda)-1),$$

$$t_s(u,v;\lambda)=\frac{u}{\sqrt{v}}\sum_{r=1}^{\infty}\frac{re^{-\sigma\sqrt{r^2+4M_s^2}}}{\sqrt{r^2+4M_s^2}}\frac{\sin(r\varphi)}{\sin(\varphi)},$$

$$e^{i\varphi}=\sqrt{z/\bar{z}}\text{ and }e^{-\sigma}=\sqrt{z\bar{z}}$$

$$t_s(u,v;\lambda)=\frac{u}{\sqrt{v}}\sum_{n\geq 0}(W_s(\varphi+2\pi n,\sigma)+W_s(2\pi-\varphi+2\pi n,\sigma))$$
$$W_s(x,\sigma)=\frac{2M_sxK_1\left(2M_s\sqrt{x^2+\sigma^2}\right)}{\sin(x)\sqrt{x^2+\sigma^2}}$$

$$\mathcal{T}_{m,K}(u,v;\lambda)=\sum_{\Delta,S}\sum_i\left|C_{T_2\mathcal{O}_K^m\mathcal{O}_{\Delta,S}^{(i)}}\right|^2\mathcal{G}_{\Delta,S}(u,v)$$

$$\mathcal{G}_{\Delta,S}(u,v)=e^{-\sigma(\Delta-\Delta_H)}\frac{\sin(S+1)\varphi}{\sin\varphi}$$

$$\mathcal{T}_{m,K}(u,v;\lambda)=\sum_s\sum_{S=0}^{\infty}\sum_{n=0}^{\infty}\sum_{r=1}^{S+1}C_{S,n,r}^{(s)}e^{-\sigma\left(\Delta_{S,n,r}^{(s)}-\Delta_{\mathcal{H}}\right)}\frac{\sin(S+1)\varphi}{\sin\varphi}$$

$$C_{S,n,r}^{(s)}=\frac{(r+n)(S+2+n-r)}{\sqrt{(r+n)^2+4M_s^2}\sqrt{(S+2+n-r)^2+4M_s^2}}$$
$$\Delta_{S,n,r}^{(s)}=\Delta_{\mathcal{O}_K^m}+\sqrt{(r+n)^2+4M_s^2}+\sqrt{(S+2+n-r)^2+4M_s^2}$$

$$\Delta_{S,n,r}^{(s)}=\Delta_{\mathcal{O}_K^m}+2+2n+S+O(\lambda)$$

$$\mathcal{O}_K^m\,\square^n\underbrace{\partial\ldots\partial}_{S}T_2$$

$$\mathcal{T}_{m,K}(u,v;\lambda)\simeq C_{\mathcal{O}_K^m\mathcal{O}_K^m\mathcal{O}_{\mathcal{K}}}C_{T_2T_2\mathcal{O}_{\mathcal{K}}}u^{\gamma_{\mathcal{K}}/2}-\left(C_{\mathcal{O}_K^m\mathcal{O}_K^m\mathcal{O}_{\mathcal{K}}}C_{T_2T_2\mathcal{O}_{\mathcal{K}}}\right)\Big|_{g_{\mathrm{YM}}=0},$$

$$\gamma_{\mathcal{K}} = 3\frac{Ng_{\mathrm{YM}}^2}{4\pi^2} - 3\left(\frac{Ng_{\mathrm{YM}}^2}{4\pi^2}\right)^2 + \frac{21}{4}\left(\frac{Ng_{\mathrm{YM}}^2}{4\pi^2}\right)^3 + \cdots$$

$$\gamma_{\mathcal{K}} = 12N\frac{\lambda}{m} + O(1/m^2)$$

$$C_{T_2T_2\mathcal{O}_{\mathcal{K}}} = 1 - \frac{3}{2}\frac{Ng_{\mathrm{YM}}^2}{4\pi^2} + \cdots = 1 - 6N\frac{\lambda}{m} + O(1/m^2)$$

$$C_{\mathcal{O}_K^m\mathcal{O}_K^m\mathcal{O}_{\mathcal{K}}} = mKC^{(0)}_{\mathcal{O}_K^m\mathcal{O}_K^m\mathcal{O}_{\mathcal{K}}} + \sum_{L\geq 1} C^{(L)}_{\mathcal{O}_K^m\mathcal{O}_K^m\mathcal{O}_{\mathcal{K}}}\lambda^L + O(1/m)$$

$$\mathcal{T}_{m,K}(u,v;\lambda) \simeq \sum_{L\geq 1} C^{(L)}_{\mathcal{O}_K^m\mathcal{O}_K^m\mathcal{O}_{\mathcal{K}}}\lambda^L + 6N\lambda C^{(0)}_{\mathcal{O}_K^m\mathcal{O}_K^m\mathcal{O}_{\mathcal{K}}}(\log u - 1)$$

$$P^{(1)}(u,v) = u(2 - \log u) + O(u^2, u(v-1))$$
$$P^{(L\geq 2)}(u,v) = u\binom{2L}{L}\zeta(2L-1) + O(u^2, u(v-1))$$

$$\frac{1}{2u}(t_s^2(u,v;\lambda) - 1) \simeq -M_s^2(2 - \log u) + \sum_{L\geq 2}(-M_s^2)^L\binom{2L}{L}\zeta(2L-1)$$

$$\mathcal{T}_{m,K}(u,v;\lambda) \simeq 2KN\lambda(\log u - 2) + \sum_s\sum_{L\geq 2}(-M_s^2)^L\binom{2L}{L}\zeta(2L-1)$$

$$C_{\mathcal{O}_K^m\mathcal{O}_K^m\mathcal{O}_{\mathcal{K}}} \simeq \frac{mK}{3} - 2KN\lambda + \sum_s\sum_{L\geq 2}(-M_s^2)^L\binom{2L}{L}\zeta(2L-1),$$

$$C_{\mathcal{O}_K^m\mathcal{O}_K^m\mathcal{O}_{\mathcal{K}}} \simeq \frac{mK}{3}\underbrace{-2KN\lambda + \sum_s\int_0^\infty \frac{M_s dw}{\sinh^2(w)}[2M_s w - J_1(4M_s w)]}_{c_1(\lambda)},$$

$$C_{\mathcal{O}_N^m\mathcal{O}_N^m\mathcal{O}_{\mathcal{K}}}(\lambda^*) = \frac{mK}{3}.$$

$$a = \mathrm{diag}(a_1, a_2, \cdots, a_N), \sum_{i=1}^N a_i = 0$$

$$\mathcal{Z}(\tau,\tau_p;\mu) = \int d\sigma(a_i)\left|\exp\left(\mathrm{i}\pi\tau\sum_i a_i^2 + \mathrm{i}\sum_{p>2}\pi^{p/2}\tau_p\sum_i a_i^p\right)\right|^2 Z_{1\text{-loop}}(a;\mu)\left|Z_{\mathrm{inst}}(\tau,\tau_p,a;\mu)\right|^2$$

$$d\sigma(a_i) = \prod_{i=1}^N da_i\prod_{i<j}a_{ij}^2\delta\left(\sum_i a_i\right), a_{ij} := a_i - a_j$$

$$\mathcal{Z} \equiv \mathcal{Z}(\tau,0;0) = \int d\sigma(a_i)\exp\left(-2\pi\tau_2\sum_i a_i^2\right)$$

$$a \to \frac{a}{\sqrt{2\pi\tau_2}}$$

$$\langle\langle f(a_i)\rangle\rangle = \frac{1}{\mathcal{Z}}\int d\sigma(a_i)\exp\left(-\sum_i a_i^2\right) f(a_i)$$

$$t_n(a) = \prod_{k\geq 1}\left(\mathrm{Tr}a^{n_k}\right)$$

$$\left\langle\langle t_{\boldsymbol{n}}(a)t_{\boldsymbol{m}}(a)\rangle\right\rangle = \frac{1}{\mathcal{Z}}\prod_{k\geq 1}\prod_{l\geq 1}\left(-\mathrm{i}\pi^{1/2}\right)^{-n_k}\left(\mathrm{i}\pi^{1/2}\right)^{-m_l}\partial_{\tau_{n_k}}\partial_{\bar{\tau}_{m_l}}\mathcal{Z}(\tau,\tau_p;0)\Big|_{\tau_p=0}.$$

$$T_{\boldsymbol{n}}(a) = t_{\boldsymbol{n}}(a) + \sum_{\boldsymbol{m}\vdash m<n}\alpha_{\boldsymbol{m},\boldsymbol{n}}t_{\boldsymbol{m}}(a),$$

$$\left\langle\langle T_{\boldsymbol{n}}(a)t_{\boldsymbol{m}}(a)\rangle\right\rangle = 0, \boldsymbol{m}\vdash m<n.$$

$$\mathcal{G}_{m,K}(\tau,\bar{\tau}) = I_2\left[\mathcal{T}_{m,K}(u,v;\tau,\bar{\tau})\right] = -\frac{8}{\pi}\int_0^\infty dr\int_0^\pi d\theta\,\frac{r^3\sin^2\theta}{uv}\mathcal{T}_{m,K}(u,v;\tau,\bar{\tau})$$

$$Z_{1-\mathrm{loop}}(a;\mu) = \frac{1}{H(\mu)^N}\prod_{i<j}\frac{H^2(a_{ij})}{H(a_{ij}+\mu)H(a_{ij}-\mu)},$$

$$H(x) = \mathrm{e}^{-(1+\gamma)x^2}G(1+\mathrm{i}x)G(1-\mathrm{i}x)$$

$$\mathcal{G}_{m,K}(\tau,\bar{\tau}) = \frac{\partial_{\tau_{m,K}}\partial_{\bar{\tau}_{m,K}}\partial_\mu^2\log\mathcal{Z}(\tau,\tau_p;\mu)\big|_{\tau_p,\mu=0}}{\partial_{\tau_{m,K}}\partial_{\bar{\tau}_{m,K}}\log\mathcal{Z}(\tau,\tau_p;0)\big|_{\tau_p=0}}$$

$$\log Z_{1\text{-loop}} = -\mu^2\left[\sum_{L=1}^{\infty}\sum_{j=0}^{2L}(-1)^{L+j}\left(\frac{1}{2\pi\tau_2}\right)^L\binom{2L}{j}(2L+1)\zeta(2L+1)\mathrm{Tr}a^{2L-j}\mathrm{Tr}a^j\right] + O(\mu^4).$$

$$\left\{t_{4,4,4}, t_{5,4,3}, t_{3,3,3,3}, t_{5,5,2}, t_{4,3,3,2}, t_{4,4,2,2}, t_{5,3,2,2}, t_{3,3,2,2,2}, t_{4,2,2,2,2}, t_{2,2,2,2,2,2}\right\}$$

$$\mathcal{O}_{\mathbf{m}}(a), \mathbf{m} = \{m_1, m_2, \dots\}$$

$$\{t_2^n\mathcal{O}_{\mathbf{m}}(a)\}_{n,\mathbf{m}\geq 0}$$

$$\left\langle\left\langle t_2^n\mathcal{O}_{\mathbf{m}}(a), t_2^{n'}\mathcal{O}_{\mathbf{m}'}(a)\right\rangle\right\rangle \propto \delta_{\mathbf{m},\mathbf{m}'}$$

$$\mathcal{O}_{i_1,\dots i_{k-1}}, \underbrace{i_k,\dots,i_k}_{m}, i_{k+1},\dots i_n \text{ with } i_\ell \neq i_j \text{ if } \ell \neq j,$$

$$\mathcal{O}_{i_1,\dots i_{k-1}}, \underbrace{i_k,\dots,i_k}_{m}, i_{k+1},\dots i_n \simeq \mathcal{O}\,\underbrace{i_k,\dots,i_k}_{m}.$$

$$\mathcal{O}_K^m(a) = \underbrace{t_{K,\dots,K}}_{m}(a) + \sum_{\mathbf{n}} c_{\mathbf{n}}t_{\mathbf{n}}(a)$$

$$\mathcal{O}^{4,\dots,}_{m^{4,\dots,4}_{4=\Delta/4}} = t_4^{m_4} + \text{GS with } \left\{t_4^{n_4}t_3^{n_3}\right\}_{n_3\geq 1} \cup \left\{t_4^{n_4}t_3^{n_3}t_2^{n_2}\right\}_{n_2\geq 1}$$
$$\vdots$$
$$\mathcal{O}^{4,\dots,3}_{m^{4,\dots,4}_{m_4}} = t_4^{m_4}t_3^{m_3} + \text{GS with } \left\{t_4^{n_4}t_3^{n_3}\right\}_{n_3\geq m_3+1} \cup \left\{t_4^{n_4}t_3^{n_3}t_2^{n_2}\right\}_{n_2\geq 1}$$
$$\vdots$$
$$\mathcal{O}^{3,\dots,3}_{m_3=\Delta/3} = t_3^{m_3} + \text{GS with } \left\{t_4^{n_4}t_3^{n_3}t_2^{n_2}\right\}_{n_2\geq 1}$$

$$\mathcal{O}_3^5 = t_3^5 + \frac{35325}{39442}t_3^3t_2^3 - \frac{90}{41}t_4t_3^3t_2 + \frac{1050327}{9781616}t_3t_2^6 - \frac{22761}{39442}t_4t_3t_2^4 + \frac{405}{533}t_4^2t_3t_2^2$$

$$\mathcal{O}_4^3 = t_4^3 - \frac{6103}{5140}t_4^2t_2^2 + \frac{9551}{20560}t_4t_2^4 - \frac{162}{1285}t_4t_3^2t_2 + \frac{53}{6939}t_3^4 + \frac{1313}{23130}t_3^2t_2^3 - \frac{985}{16448}t_2^6$$

$$\underbrace{\mathcal{O}_{5,\dots,5}}_{m_5}\underbrace{4,\dots,4,,\underbrace{3,\dots,3}_{m_3}}_{m_4} = t_5^{m_5}t_4^{m_4}t_3^{m_3} + \text{GS with } \left\{t_5^{n_5}t_4^{n_4}t_3^{n_3}\right\}_{n_4\geq m_4+1} \cup \left\{t_5^{n_5}t_4^{n_4}t_3^{n_3}\right\}_{n_3\geq m_3+1} \cup \left\{t_5^{n_5}t_4^{n_4}t_3^{n_3}t_2^{n_2}\right\}_{n_2\geq 1},$$

$$\mathcal{O}_5^m = t_5^m + \text{GS with } \left\{t_5^{n_5}t_4^{n_4}t_3^{n_3}\right\}_{n_3+n_4\geq 1} \cup \left\{t_5^{n_5}t_4^{n_4}t_3^{n_3}t_2^{n_2}\right\}_{n_2\geq 1}$$
$$\mathcal{O}_4^m = t_4^m + \text{GS with } \left\{t_5^{n_5}t_4^{n_4}t_3^{n_3}\right\}_{n_3\geq 1} \cup \left\{t_5^{n_5}t_4^{n_4}t_3^{n_3}t_2^{n_2}\right\}_{n_2\geq 1}$$
$$\mathcal{O}_3^m = t_3^m + \text{GS with } \left\{t_5^{n_5}t_4^{n_4}t_3^{n_3}t_2^{n_2}\right\}_{n_2\geq 1}$$

$$\mathcal{O}_5^2 = t_5^2 - \frac{1}{90}t_4t_3^2 - \frac{23}{800}t_4^2t_2 - \frac{443}{300}t_5t_3t_2 + \frac{2461}{4500}t_3^2t_2^2 + \frac{179}{8000}t_4t_2^3 - \frac{7}{1600}t_2^5$$

$$\mathcal{N}_m(N,K) = \left\langle\left\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)\right\rangle\right\rangle$$

$$\mathcal{N}_m(N,N) = f_N(m)(2^{-N}N^{2-N})^m \frac{\Gamma\left(Nm + \frac{N(N-1)}{2} + 1\right)}{\Gamma\left(\frac{N(N-1)}{2} + 1\right)}$$

$$f_2(m) = 1$$
$$f_3(m) = 1$$
$$f_4(m) = \frac{1-2^{-4}}{1-2^{-4-4m}} = \frac{15}{16} + O(\mathrm{e}^{-m})$$
$$f_5(m) = \frac{\frac{504}{625}}{\left(1 + 5^{-\frac{5}{2}m-4}(1+(-1)^m) - 2^{-m-2}5^{-3m-5}\left((25+11\sqrt{5})^{m+2} + (25-11\sqrt{5})^{m+2}\right)\right)}$$
$$= \frac{504}{625} + O(\mathrm{e}^{-m})$$

$$\Gamma\left(\frac{R}{2} + \alpha_{N,N} + 1\right)$$

$$\left\langle\left\langle \mathcal{O}_3^m(a)\mathcal{O}_3^m(a)\right\rangle\right\rangle = 24^{-m}\frac{\Gamma(3m+7)}{\Gamma(7)}\left(\frac{20}{27} + \frac{250}{243m} + \mathcal{O}\left(\frac{1}{m^2}\right)\right)$$

$$\left\langle\left\langle \mathcal{O}_3^m(a)\mathcal{O}_3^m(a)\right\rangle\right\rangle = 24^{-m}\frac{\Gamma(3m+10)}{\Gamma(10)}\left(\frac{140}{243} + \mathcal{O}\left(\frac{1}{m}\right)\right)$$

$$\mathcal{N}_m(N,K) = (2^K K^{K-2})^{-m} \frac{\Gamma(\Delta + \alpha_{N,K} + 1)}{\Gamma(\alpha_{N,K} + 1)} A(N,K) \left(1 + \mathcal{O}\left(\frac{1}{m}\right)\right)$$

$$\frac{\langle \mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)T_{\mathbf{p}}(x_3,Y_3)\rangle}{\langle \mathcal{O}_K^m(x_1,Y_1)\mathcal{O}_K^m(x_2,Y_2)\rangle} = \mathfrak{C}_{mm\mathbf{p}}(N,K) \times \left(\frac{1}{2\pi^2}\frac{d_{23}d_{31}}{d_{12}}\right)^{\frac{\Delta_{\mathbf{p}}}{2}}$$

$$\mathfrak{C}_{mm\mathbf{p}}(N,K) = \frac{\left\langle\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)T_{\mathbf{p}}(a)\rangle\right\rangle}{\left\langle\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)\rangle\right\rangle},$$

$$\frac{\left\langle\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)T_{\mathbf{p}}(a)\rangle\right\rangle}{\left\langle\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)\rangle\right\rangle} = \frac{\left\langle\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)t_{\mathbf{p}}(a)\rangle\right\rangle}{\left\langle\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)\rangle\right\rangle}\left(1 + \mathcal{O}\left(\frac{1}{m}\right)\right)$$

$$\frac{\left\langle\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)t_{\mathbf{p}}(a)\rangle\right\rangle}{\left\langle\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)\rangle\right\rangle} = \int_0^{2\pi} \frac{d\theta}{2\pi}\langle t_{\mathbf{p}}\rangle_\theta \left(1 + \mathcal{O}\left(\frac{1}{m}\right)\right),$$

$$\langle t_{\mathbf{p}}\rangle_\theta = \prod_{k\geq 1}\left(\mathrm{Tr}\left(a_{\mathrm{cl}}^{p_k}(\theta)\right)\right), \text{ and } a_{\mathrm{cl}}(\theta) = \frac{\Omega_K^{(N)}\mathrm{e}^{\mathrm{i}\theta} + \bar{\Omega}_{\mathrm{K}}^{(\mathrm{N})}\mathrm{e}^{-\mathrm{i}\theta}}{\sqrt{2}}$$

$$\mathcal{T}_{\mathcal{H}}(u,v;\lambda) = \sum_{L=1}^{\infty} d_{\mathcal{H},N;L}\frac{(-\lambda)^L}{u}\sum_{\ell=0}^{L} P^{(\ell)}(u,v)P^{(L-\ell)}(u,v)$$

$$I_2\left[\frac{1}{u}\sum_{\ell=0}^{L} P^{(\ell)}(u,v)P^{(L-\ell)}(u,v)\right] = -4\frac{\Gamma(2L+2)}{\Gamma(L+1)^2}\zeta(2L+1)$$

$$\mathcal{G}_{\mathcal{H}}(\lambda) = -4\sum_{L=1}^{\infty} d_{\mathcal{H},N;L}(-\lambda)^L \frac{\Gamma(2L+2)}{\Gamma(L+1)^2}\zeta(2L+1)$$

$$\mathcal{G}_{m,K}(\tau_2) = \frac{\int d^{N-1}a \prod_{1\leq i<j\leq N} a_{ij}^2 \partial_\mu^2 Z_{1-\mathrm{loop}} e^{-\mathrm{Tr}a^2}\mathcal{O}_K^m\mathcal{O}_K^m}{\int d^{N-1}a \prod_{1\leq i<j\leq N} a_{ij}^2 e^{-\mathrm{Tr}a^2}\mathcal{O}_K^m\mathcal{O}_K^m}\Bigg|_{\mu=0} \otimes \partial_\mu^2 Z_{1-\text{ loop}}\big|_{\mu=0}$$

$$\mathcal{G}_{m,K}(\tau_2) = -2\sum_{L=1}^{\infty}\sum_{j=0}^{2L}(-1)^{L+j}\left(\frac{1}{2\pi\tau_2}\right)^L\binom{2L}{j}(2L+1)\zeta(2L+1)\frac{\left\langle\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)t_{2L-j,j}(a)\rangle\right\rangle}{\left\langle\langle \mathcal{O}_K^m(a)\mathcal{O}_K^m(a)\rangle\right\rangle}.$$

$$\mathcal{G}_{m,K}(\lambda) \simeq 4\sum_{L=1}^{\infty}(-1)^{L+1}\frac{\Gamma(2L+2)}{\Gamma(L+1)^2}\zeta(2L+1)\left[\sum_{1\leq i<j\leq K}\left(4\sin^2\frac{\pi(i-j)}{K}\lambda\right)^L + K(N-K)\lambda^L\right].$$

$$\mathcal{G}_{m,K}(\lambda) \simeq -2\sum_{L=1}^{\infty} \frac{\Gamma(2L+2)}{\Gamma(L+1)^2}\zeta(2L+1)\sum_{s} (-M_s^2)^L$$

$$d_{\mathcal{O}_K^m,N;L} = \frac{K}{2}\sum_{s=1}^{K-1}\left(4\sin^2\left(\frac{s\pi}{K}\right)\right)^L + K(N-K).$$

$$\mathcal{T}_{m,K}(u,v;\lambda) = K\sum_{s=1}^{K-1} L\left(u,v;4\lambda\sin^2\frac{\pi s}{K}\right) + 2K(N-K)L(u,v;\lambda)$$

$$L(u,v;a) = \frac{1}{2u}\left[\left(\sum_{L=0}^{\infty}(-a)^L P^{(L)}(u,v)\right)^2 - 1\right]$$

$$-\sum_{L\geq 1}\frac{\Gamma(2L+2)}{\Gamma(L+1)^2}\zeta(2L+1)(-a)^L = \int_0^{\infty} dw\,\frac{w}{\sinh^2 w}\left(1-J_0(4w\sqrt{a})\right)$$
$$= 1+\gamma_E+\frac{\log a}{2}+2\sum_{n\geq 1}\left(4\pi n\sqrt{a}K_1(4\pi n\sqrt{a}) - K_0(4\pi n\sqrt{a})\right)$$

$$\mathcal{G}_{m,K}(\lambda) \simeq 2K(2N-K-1)(1+\gamma_E) + 2K\left(\log K + \left(N-\frac{K+1}{2}\right)\log\lambda\right)$$
$$+4\sum_{s}\sum_{n\geq 1}\left(4\pi n M_s K_1(4\pi n M_s) - K_0(4\pi n M_s)\right)$$

$$K_\nu(4\pi n M_s) \sim \mathrm{e}^{-4\pi n M_s}$$

$$N\to\infty, \text{ with } K \text{ fixed.}$$

$$\mathcal{G}_{m,K}(\lambda)\big|_{\triangle} \simeq 4NK\sum_{L\geq 1}(-1)^{L+1}\frac{\Gamma(2L+2)}{\Gamma(L+1)^2}\zeta(2L+1)\lambda^L$$

$$N,K\to\infty, \text{ with } \kappa = \frac{K}{N}, \text{ fixed.}$$

$$\frac{1}{K}\sum_{s=1}^{K-1}\left(4\sin^2\left(\frac{\pi s}{K}\right)\right)^L = \int_0^1 \mathrm{d}x(4\sin^2(\pi x))^L + \mathcal{O}\left(\frac{1}{K}\right) = \frac{2^{2L}\Gamma\left(L+\frac{1}{2}\right)}{\sqrt{\pi}\Gamma(L+1)} + \mathcal{O}\left(\frac{1}{K}\right)$$

$$\mathcal{G}_{m,K}(\lambda)\big|_{\otimes} \simeq -4\kappa N^2\sum_{L=1}^{\infty}\frac{(-\lambda)^L\zeta(2L+1)\Gamma\left(L+\frac{3}{2}\right)\left[\sqrt{\pi}2^{2L+1}(1-\kappa)\Gamma(L+1)+2^{4L}\kappa\Gamma\left(L+\frac{1}{2}\right)\right]}{\pi\Gamma(L+1)^2}$$
$$= 4\kappa N^2\int_0^{\infty}\mathrm{d}w\,\frac{w}{\sinh^2(w)}\left[(1-\kappa)\left(1-J_0(4w\sqrt{\lambda})\right)+\frac{\kappa}{2}\left(1-J_0(4w\sqrt{\lambda})^2\right)\right]$$

$$\mathcal{G}_{m,K}(\lambda)\big|_{\bullet} \simeq (2-\kappa)\kappa N^2(\log(\lambda)+2\gamma_E+2) + 16\kappa^2N^2\sum_{k\geq 1}\frac{a_k}{\lambda^{k-1/2}}$$
$$+8N^2\kappa(1-\kappa)\sum_{n\geq 1}\left(4\pi n\sqrt{\lambda}K_1(4\pi n\sqrt{\lambda}) - K_0(4\pi n\sqrt{\lambda})\right)$$

$$a_k = \frac{\zeta(2k-1)\Gamma\left(k-\frac{1}{2}\right)^3}{\sqrt{\pi}(16\pi^2)^k\Gamma(k-1)}$$

$$\lim_{K\to\infty} M_s = \lim_{K\to\infty} 2\sin\left(\frac{\pi s}{K}\right) = 0$$

$$a_k \sim \Gamma(2k-1)\pi^{-2k}\left(2^{4-10k} + 2^{3-8k} + 2^{2-6k} + 2^{2-6k}3^{1-2k} + \mathcal{O}(1/k)\right)$$

$$\sigma(x) = \frac{1}{\pi}\frac{1}{\sqrt{x(1-x)}}$$

$$\mathcal{G}_{m,3}^{SU(4)}(\lambda) = \int_0^1 \left(\partial_\mu^2 Z_{1-\text{ loop}}\right)\Big|_{\mu=0,\mathrm{Tr}a^2=6\lambda,\mathrm{Tr}a^3=6\lambda^{3/2}(2x-1)} \sigma(x)\mathrm{d}x$$

$$\mathcal{G}_{m,K}(\lambda) = \int_0^1 \left(\partial_\mu^2 Z_{1-\text{ loop}}\big|_{\mu=0}\right)\Big|_{\text{locus}} \sigma(x)\mathrm{d}x$$

$$\mathrm{Tr}a^m = 2K\lambda^{\frac{m}{2}}\left(\binom{m-1}{m/2}\epsilon_m + (2x-1)\delta_{m,K}\right), m = 2,\dots,K$$

$$\mathrm{Tr}a^m = \lambda^{\frac{m}{2}}\sum_{s=0}^{K-1}\left(\mathrm{e}^{-\frac{2\mathrm{i}\pi s}{K}}(\sqrt{x-1}-\sqrt{x})^{-2/K} + \mathrm{e}^{\frac{2\mathrm{i}\pi s}{K}}(\sqrt{x-1}-\sqrt{x})^{2/K}\right)^m.$$

$$\mathcal{Z} = \int da\mathrm{e}^{-\mathrm{Tr}a^2}, a = \sum_{b=1}^{N^2-1} a^b T_b, da = \prod_{b=1}^{N^2-1}\frac{da_b}{\sqrt{2\pi}}$$

$$\mathrm{Tr}T_bT_c = \frac{1}{2}\delta_{bc}, \mathrm{Tr}T_b = 0$$

$$\langle\langle f(a)\rangle\rangle = \int da\mathrm{e}^{-\mathrm{Tr}a^2} f(a)$$

$$u_{\boldsymbol{p}} = \langle\langle \mathrm{Tr}a^{p_1}\mathrm{Tr}a^{p_2}\dots\mathrm{Tr}a^{p_m}\rangle\rangle$$

$$\mathrm{Tr}T^bB_1T^bB_2 = \frac{1}{2}\mathrm{Tr}B_1\mathrm{Tr}B_2 - \frac{1}{2N}\mathrm{Tr}B_1B_2$$
$$\mathrm{Tr}T^bB_1\mathrm{Tr}T^bB_2 = \frac{1}{2}\mathrm{Tr}B_1B_2 - \frac{1}{2N}\mathrm{Tr}B_1\mathrm{Tr}B_2$$

$$u_p = 0, \text{ for } p \text{ odd}, \text{ and } u_0 = N$$

$$u_{p_1,p_2,\dots,p_m} = \frac{1}{2}\sum_{j=0}^{p_1-2}\left(u_{j,p_1-j-2,p_2,\dots,p_m}\right) - \frac{p_1-1}{2N}u_{p_1-2,p_2,\dots,p_m}$$
$$+\sum_{k=2}^{m}\frac{p_k}{2}\left(u_{p_1+p_k-2,p_2,\dots,p_k,\dots,p_m} - \frac{1}{N}u_{p_1-1,p_2,\dots,p_k-1,\dots,p_m}\right),$$

$$u_2 = \frac{N^2-1}{2}, u_4 = \frac{(N^2-1)(2N^2-3)}{4N}$$
$$u_{2,2} = \frac{N^4-1}{4}, u_6 = \frac{5(N^2-1)(N^4-3N^2+3)}{8N^2}$$
$$u_{4,2} = \frac{(N^2-1)(N^2+3)(2N^2-3)}{8N}, u_{3,3} = \frac{3(N^2-1)(N^2-4)}{8N}$$

$$\frac{\langle\langle \mathcal{O}_5^m \mathcal{O}_5^m t_4 \rangle\rangle}{\left\langle\langle \mathcal{O}_5^m \mathcal{O}_5^m \rangle\right\rangle} = \frac{30}{2^{m+2}5^{3m+5} + (1+(-1)^m)2^{m+2}5^{m/2+1} - (x_5^+)^{m+2} - (x_5^-)^{m+2}}$$
$$\left[2^m 5^{3m+4}(5m^2+27m-2) - \frac{(x_5^+)^m + (x_5^-)^m}{2}(325m^2+2219m-1290)\right.$$
$$-\sqrt{5}\frac{(x_5^+)^m - (x_5^-)^m}{2}(145m^2+991m-578)$$
$$\left.+2^m 5^{m/2}\left((1+(-1)^m)(8m-20) - \frac{1-(-1)^m}{\sqrt{5}}(25m^2+95m+90)\right)\right]$$

$$x^2 - 50x + 20 = 0, \text{ with } x_5^+ = 25+11\sqrt{5}, x_5^- = 25-11\sqrt{5}$$

$$\frac{\left\langle\langle \mathcal{O}_3^m \mathcal{O}_3^m t_{4^2,3^2} \rangle\right\rangle}{\left\langle\langle \mathcal{O}_3^m \mathcal{O}_3^m \rangle\right\rangle} = 45.5625m^7 + O(m^6)$$

$$\mathcal{G}_{m,K}(\lambda) \simeq 2\sum_{L=1}^{\infty}(-1)^{L+1}\frac{\Gamma(2L+2)}{\Gamma(L+1)^2}\zeta(2L+1)\left[K\sum_{s=1}^{K-1}\left(4\lambda\sin^2\frac{\pi s}{K}\right)^L + 2K(N-K)(\lambda)^L\right].$$

$$\mathcal{O}_{3^{m_3}2^{m_2}} = t_3^{m_3}t_2^{m_2} + \text{GS with } \{t_3^{n_3}t_2^{n_2}\}_{\Delta=3n_3+2n_2, n_3\le m_3} \cup \{t_3^{n_3}t_2^{n_2}\}_{\Delta\le 3m_3+2m_2}$$

$$\mathcal{G}_{3^{m_3}2^{m_2}}(\tau_2) = 2\sum_{L=1}^{\infty}(-1)^L\frac{\Gamma(2L+2)}{\Gamma(L+1)}\left(\frac{1}{4\pi\tau_2}\right)^L\zeta(2L+1)((2L+1)(L^2+L+6)$$
$$+6\,{}_3F_2\left(-L, L+1, -m_2; 1, \frac{15}{2}+3m_3; 1\right)\frac{(-L)_{(3m_3+3)} - (L+1)_{(3m_3+3)}}{\Gamma(3m_3+4)}\Bigg)$$

$$\mathcal{G}_{m,3}(\lambda) \simeq 4\sum_{L=1}^{\infty}(-1)^{L+1}\frac{\Gamma(2L+2)}{\Gamma(L+1)^2}3(3\lambda)^L$$

$$\mathcal{G}_{m,5}(\tau_2)\big|_{L=1} = \frac{150\zeta(5)}{\pi\tau_2}m$$

$$\mathcal{G}_{m,5}(\tau_2)\big|_{L=2} = -\frac{\pi^2\tau_2^2\left(2^{m+2}5^{3m+5} + 2^{m+2}5^{m/2+1}(1+(-1)^m) - (x_5^+)^{m+2} - (x_5^-)^{m+2}\right)}{12}$$

$$\left[2^m 5^{3m+4}\left(\frac{5m^2}{2}+\frac{77m}{6}-\frac{1}{3}\right) - ((x_5^+)^m + (x_5^-)^m)\frac{(1555m^2+8369m-1290)}{1+(-1)^m)\frac{(5m^2+29m-10)}{3} - \sqrt{5}(1-(-1)^m)\left(\frac{5m^2}{6}+\frac{19m}{6}+3\right)}\right)$$

$$+2^m 5^{\frac{m}{2}}\left(\left(1+\frac{1247m}{4}-\frac{289}{6}\right)\right]$$

$$-\sqrt{5}((x_5^+)^m - (x_5^-)^m)\left(\frac{695m^2}{12}+\frac{1220500\zeta(7)}{}\right.$$

$$\mathcal{G}_{m,5}(\tau_2)\big|_{L=3} = \frac{\pi^3\tau_2^3\left(2^{m+2}5^{3m+5} + 2^{m+2}5^{m/2+1}(1+(-1)^m) - (x_5^+)^{m+2} - (x_5^-)^{m+2}\right)}{252}$$

$$\left[2^m 5^{3m+4}\frac{(50m^3+405m^2+1087m-84)}{250} - ((x_5^+)^m + (x_5^-)^m)\frac{50m^3+1825+7623-5840}{56}\right.$$

$$+2^m 5^{\frac{m}{2}}\left((1+(-1)^m)\left(\frac{25m^3}{72}+\frac{125m^2}{56}+\frac{1279m}{252}-\frac{5}{42}\right) - 5\sqrt{5}(1-(-1)^m)\frac{5m^3+29m^2+56m+36}{56}\right)$$

$$\left.-5\sqrt{5}((x_5^+)^m - (x_5^-)^m)\left(\frac{5m^3}{63}+\frac{163m^2}{56}+\frac{6131m}{504}-\frac{28}{3}\right)\right]$$

$$\mathcal{G}_{m,5}(\lambda) = 600\lambda\zeta(3) - 7200\lambda^2\zeta(5) + 140000\lambda^3\zeta(7) + \mathcal{O}(\lambda^4, 1/m)$$

$$\mathcal{G}_{m,3}(\lambda) = 288.0\lambda\zeta(3) - (3600.0\lambda^2\zeta(5) + \mathcal{O}(1/m)) + (47040.0\lambda^3\zeta(7) + \mathcal{O}(1/m))$$
$$-(619917.3\lambda^4\zeta(9) + \mathcal{O}(1/m)) + \lambda$$

$$\mathcal{G}_{m,3}(\lambda) = 288\lambda\zeta(3) - 3600\lambda^2\zeta(5) + 47040\lambda^3\zeta(7) - 619920\lambda^4\zeta(9) + \mathcal{O}(\lambda^5)$$

$$\langle T(\zeta_1)T(\zeta_2)\rangle = \frac{c/2}{(\zeta_1-\zeta_2)^4}, \langle T(\zeta_1)T(\zeta_2)T(\zeta_3)\rangle = \frac{c}{(\zeta_1-\zeta_2)^2(\zeta_2-\zeta_3)^2(\zeta_3-\zeta_1)^2},$$

$$T \equiv T_{\zeta\zeta}, \text{ and } \zeta \equiv x_1 + ix_2, \bar{\zeta} \equiv x_1 - ix_2$$

$$\left\langle T_\mu^\mu \right\rangle^{d=4} = \frac{1}{16\pi^2}\left(cW_{\mu\nu\rho\sigma}^2 - aE_4\right) + \frac{\delta(x_\perp)}{16\pi^2}\left(aE_4^{(\text{bry})} - b_1\text{tr}\hat{K}^3 - b_2 h^{pq}\hat{K}^{rs}W_{pqrs}\right),$$

$$\hat{K}_{pq} \equiv K_{pq} - Kh_{pq}/3$$

$$\langle \mathcal{D}(\mathbf{x}_1)\mathcal{D}(\mathbf{x}_2)\rangle = \frac{c_{\eta\eta}}{\mathbf{x}_{12}^8}, \langle \mathcal{D}(\mathbf{x}_1)\mathcal{D}(\mathbf{x}_2)\mathcal{D}(\mathbf{x}_3)\rangle = \frac{c_{\eta\eta\eta}}{\mathbf{x}_{12}^4\mathbf{x}_{23}^4\mathbf{x}_{31}^4},$$

$$\mathbf{x}_{ij} \equiv |\mathbf{x}_i - \mathbf{x}_j|$$

$$b_1 = \frac{2\pi^3}{35}\cdot c_{\eta\eta\eta}, b_2 = \frac{2\pi^4}{15}\cdot c_{\eta\eta}$$

$$\varphi_i = \varphi_i^{\text{SYM}}(x_3) = \frac{1}{x_3}\cdot\begin{bmatrix}\left(t_i^{(k_1)}\otimes \mathbb{1}_{k_2}\right)_{k\times k} & 0_{k\times(N_c-k)} \\ 0_{(N_c-k)\times k} & 0_{(N_c-k)\times(N_c-k)}\end{bmatrix}_{N_c\times N_c}$$

$$\varphi_i = \varphi_i^{\text{SYM}}(x_3) = \frac{1}{x_3}\cdot\begin{bmatrix}\left(\mathbb{1}_{k_1}\otimes t_i^{(k_2)}\right)_{k\times k} & 0_{k\times(N_c-k)} \\ 0_{(N_c-k)\times k} & 0_{(N_c-k)\times(N_c-k)}\end{bmatrix}_{N_c\times N_c}$$

$$i = 4,5,6,$$

$$A_\mu = \psi_\alpha = 0.\ \text{For}\ k_1, k_2 = 0,1,\ldots$$

$$k \equiv k_1 \cdot k_2$$

$$\left[t_i^{(k_i)}, t_j^{(k_i)}\right] = i\epsilon_{ijl} t_l^{(k_i)}, i, j, l = 1,2,3$$

$$\varphi_i = \varphi_i^{\text{SYM}}(x_3) = \frac{1}{\sqrt{8}x_3} \cdot \begin{bmatrix} (G_i)_{d_G \times d_G} & 0_{d_G \times (N_c - d_G)} \\ 0_{(N_c - d_G) \times d_G} & 0_{(N_c - d_G) \times (N_c - d_G)} \end{bmatrix}_{N_c \times N_c}, i = 1, \ldots, 5$$

$$\varphi_6 = 0$$

$$d_G \equiv \frac{1}{6} \cdot (n+1)(n+2)(n+3), n = 1,2,\ldots$$

$$G_i \equiv [\overbrace{\gamma_i \otimes \mathbb{1}_4 \otimes \ldots \otimes \mathbb{1}_4}^{n\ \text{factors}} + \underbrace{\mathbb{1}_4 \otimes \gamma_i \otimes \ldots \otimes \mathbb{1}_4 + \cdots + \mathbb{1}_4 \otimes \ldots \otimes \mathbb{1}_4 \otimes \gamma_i}_{n\ \text{terms}}]_{\text{sym}}$$

$$\gamma_i = \begin{pmatrix} 0 & -i\sigma_i \\ i\sigma_i & 0 \end{pmatrix}, i = 1,2,3, \gamma_4 = \begin{pmatrix} 0 & \mathbb{1}_2 \\ \mathbb{1}_2 & 0 \end{pmatrix}, \gamma_5 = \begin{pmatrix} \mathbb{1}_2 & 0 \\ 0 & -\mathbb{1}_2 \end{pmatrix},$$

$$\{\gamma_i, \gamma_j\} = 2\delta_{ij}\mathbb{1}_4$$

$$G_{ij} \equiv \frac{1}{2}[G_i, G_j]$$

$$[G_{ij}, G_{kl}] = 2(\delta_{jk}G_{il} + \delta_{il}G_{jk} - \delta_{ik}G_{jl} - \delta_{jl}G_{ik})$$

$$\langle \Theta_{\mu\nu}(\mathrm{x}_1)\Theta_{\rho\sigma}(\mathrm{x}_2)\rangle = \frac{C_T}{\mathrm{x}_{12}^{2d}} \cdot \mathcal{I}_{\mu\nu\rho\sigma}(\mathrm{x}_1 - \mathrm{x}_2), \mathrm{x}_{12} \equiv |x_1 - x_2|$$

$$\mathcal{I}_{\mu\nu}(\mathrm{x}) \equiv g_{\mu\nu} - \frac{2\mathrm{x}_\mu \mathrm{x}_\nu}{\mathrm{x}^2}, \mathcal{I}_{\mu\nu\rho\sigma}(\mathrm{x}) \equiv \frac{1}{2}\left(\mathcal{I}_{\mu\rho}(\mathrm{x})\mathcal{I}_{\nu\sigma}(\mathrm{x}) + \mathcal{I}_{\mu\sigma}(\mathrm{x})\mathcal{I}_{\nu\rho}(\mathrm{x})\right) - \frac{1}{d}g_{\mu\nu}g_{\rho\sigma}$$

$$C_T = \frac{N_0 + 3N_{1/2} + 12N_1}{3\pi^4}$$

$$C_T = \frac{10N_c^2}{\pi^4}$$

$$\langle \Theta_{\mu\nu}(\mathrm{x}_1)\rangle = 0$$

$$\langle \Theta_{\mu\nu}(\mathrm{x}_1)\Theta_{\rho\sigma}(\mathrm{x}_2)\rangle = \frac{1}{\mathrm{x}_{12}^{2d}} \cdot \left\{\left(X_\mu X_\nu - \frac{g_{\mu\nu}}{d}\right)\left(X'_\rho X'_\sigma - \frac{g_{\rho\sigma}}{d}\right)A(v) + \left(X_\mu X'_\rho I_{\nu\sigma}\right.\right.$$
$$+X_\mu X'_\sigma I_{\nu\rho} + X_\nu X'_\sigma I_{\mu\rho} + X_\nu X'_\rho I_{\mu\sigma} - \frac{4}{d}g_{\mu\nu}X'_\rho X'_\sigma - \frac{4}{d}g_{\rho\sigma}X_\mu X_\nu + \frac{4}{d^2}g_{\mu\nu}g_{\rho\sigma}\Big)B(v)$$
$$\left.+\mathcal{I}_{\mu\nu\rho\sigma}C(v)\right\},$$

$$\xi \equiv \frac{\mathrm{x}_{12}^2}{4|\mathrm{z}_1||\mathrm{z}_2|}, v^2 \equiv \frac{\xi}{\xi + 1} = \frac{\mathrm{x}_{12}^2}{\mathrm{x}_{12}^2 + 4|\mathrm{z}_1||\mathrm{z}_2|}$$

$$X_\mu \equiv \mathrm{z}_1 \cdot \frac{v}{\xi}\frac{\partial \xi}{\partial \mathrm{x}_1^\mu} = v\left(\frac{2\mathrm{z}_1}{\mathrm{x}_{12}^2}\left(\mathrm{x}_{1\mu} - \mathrm{x}_{2\mu}\right) - \eta_\mu\right)$$
$$X'_\rho \equiv \mathrm{z}_2 \cdot \frac{v}{\xi}\frac{\partial \xi}{\partial \mathrm{x}_2^\rho} = -v\left(\frac{2\mathrm{z}_2}{x_{12}^2}\left(\mathrm{x}_{1\rho} - \mathrm{x}_{2\rho}\right) + \eta_\rho\right)$$

$$X_\mu X_\mu = X'_\rho X'_\rho = 1, X'_\rho = \mathcal{I}_{\rho\mu} X_\mu$$

$$\left(v\frac{d}{dv} - d\right)\alpha(v) = 2(d-1)\gamma(v)$$

$$\alpha(v) \equiv \frac{d-1}{d^2}\cdot\left[(d-1)(A(v)+4B(v)) + dC(v))\right], \gamma(v) \equiv -B(v) - \frac{C(v)}{2}$$

$$A_\mu = \tilde{A}_\mu^a T^a, \psi_{\alpha,m} = \tilde{\psi}_{\alpha,m}^a T^a, \varphi_i(x) = \varphi_i^{\mathrm{SYM}}(x_3) + \tilde{\varphi}_i(x), \tilde{\varphi}_i(x) = \tilde{\varphi}_i^a(x) T^a$$

$$\Theta_{\mu\nu}(x) = \sum_{n=0}^{4} \Theta_{\mu\nu}^{(n)}(x) = \Theta_{\mu\nu}^{(0)} + \Theta_{\mu\nu}^{(1)} + \Theta_{\mu\nu}^{(2)} + \Theta_{\mu\nu}^{(3)}(x) + \Theta_{\mu\nu}^{(4)}$$

$$\Theta_{\mu\nu}^{\mathrm{SYM}} \equiv \Theta_{\mu\nu}^{(0)} = 0$$

$$\varphi_i = \varphi_i^{\mathrm{SYM}}(x_3) = \frac{1}{x_3}\cdot\begin{bmatrix}(\tau_i)_{\mathrm{k}\times\mathrm{k}} & 0_{\mathrm{k}\times(N_c-\mathrm{k})} \\ 0_{(N_c-\mathrm{k})\times\mathrm{k}} & 0_{(N_c-\mathrm{k})\times(N_c-\mathrm{k})}\end{bmatrix}_{N_c\times N_c}, i = 1,\ldots,6$$

$$\tilde{\varphi}_i = \begin{bmatrix} \begin{bmatrix} * & * \\ * & * \end{bmatrix}_{\mathrm{k}\times\mathrm{k}} & \begin{matrix} * & * \\ * & * \end{matrix} \\ \begin{matrix} * & * \\ * & * \end{matrix} & \begin{bmatrix} * & * \\ * & * \end{bmatrix} \end{bmatrix}_{N_c\times N_c},$$

$$\Theta_{\mu\nu}^{(1)}(x) = \frac{1}{g_{\mathrm{YM}}^2}\frac{4}{3x_3^2}\cdot\mathrm{tr}\left\{\left(\frac{1}{x_3}\eta_\mu\eta_\nu\tilde{\varphi}_i + \eta_\mu\partial_\nu\tilde{\varphi}_i + \eta_\nu\partial_\mu\tilde{\varphi}_i - \frac{g_{\mu\nu}}{2}\partial_3\tilde{\varphi}_i + \frac{x_3}{2}\partial_\mu\partial_\nu\tilde{\varphi}_i\right)\cdot\tau_i + \frac{g_{\mu\nu}}{2x_3}\left(2\tau_j\tau_i\tau_j - \tau_i\tau_j\tau_j - \tau_j\tau_j\tau_i\right)\cdot\tilde{\varphi}_i\right\}$$

$$\tau_i = t_i^{(k_1)} \otimes \mathbb{1}_{k_2} \text{ and } \tau_{i+3} = \mathbb{1}_{k_1} \otimes t_i^{(k_2)} \text{ (for } i = 1,2,3\text{)}$$


$$\Theta_{\mu\nu}^{(1)}(x) = \frac{1}{g_{\mathrm{YM}}^2}\frac{4}{3x_3^2}\cdot\mathrm{tr}\left\{\left(\frac{1}{x_3}\left(\eta_\mu\eta_\nu - g_{\mu\nu}\right)\tilde{\varphi}_i + \eta_\mu\partial_\nu\tilde{\varphi}_i + \eta_\nu\partial_\mu\tilde{\varphi}_i - \frac{g_{\mu\nu}}{2}\partial_3\tilde{\varphi}_i + \frac{x_3}{2}\partial_\mu\partial_\nu\tilde{\varphi}_i\right)\cdot\tau_i\right\}$$

$$\tau_i\tau_i = (c_1 + c_2)\cdot\mathbb{1}_k, \tau_j\tau_i\tau_j = (c_1 + c_2 - 1)\tau_i, c_1 \equiv \frac{k_1^2 - 1}{4}, c_2 \equiv \frac{k_2^2 - 1}{4}.$$

$$\Theta_{\mu\nu}^{(1)}(x)=\frac{1}{g_{\mathrm{YM}}^2}\frac{4}{3x_3^2}\cdot\mathrm{tr}\left\{\left(\frac{1}{x_3}\left(\eta_\mu\eta_\nu-g_{\mu\nu}\right)\tilde{\varphi}_i+\eta_\mu\partial_\nu\tilde{\varphi}_i+\eta_\nu\partial_\mu\tilde{\varphi}_i-\frac{g_{\mu\nu}}{2}\partial_3\tilde{\varphi}_i+\frac{x_3}{2}\partial_\mu\partial_\nu\tilde{\varphi}_i\right)\cdot\frac{G_i}{\sqrt{8}}\right\}$$

$$G_iG_i=n(n+4)\cdot\mathbb{1}_{d_G},G_jG_iG_j=(n(n+4)-8)G_i.$$

$$\left\langle\Theta_{\mu\nu}\left(\mathrm{x}_1\right)\Theta_{\rho\sigma}\left(\mathrm{x}_2\right)\right\rangle=$$



$$\mathrm{tr}[\tau_i\tilde{\varphi}_i]\cdot\mathrm{tr}\left[\tau_j\tilde{\varphi}_j\right]=k(c_1+c_2)K^{5/2}(x,y)=k(c_1+c_2)\frac{g_{\mathrm{YM}}^2}{320\pi^2}\frac{1}{x_3y_3}\frac{{}_2F_1(2,3,6;-\xi^{-1})}{\xi^3(1+\xi)}$$

$$\mathrm{tr}[G_i\tilde{\varphi}_i]\cdot\mathrm{tr}\left[G_j\tilde{\varphi}_j\right]=c_n\cdot K^{5/2}(x,y)=c_n\cdot\frac{g_{\mathrm{YM}}^2}{320\pi^2}\frac{1}{x_3y_3}\frac{{}_2F_1(2,3,6;-\xi^{-1})}{\xi^3(1+\xi)}$$

$$c_n\equiv\mathrm{tr}[G_iG_i]=\frac{1}{6}\cdot n(n+1)(n+2)(n+3)(n+4),n=1,2,\ldots$$

$$\left\langle\Theta_{\mu\nu}^{(1)}(\mathrm{x}_1)\Theta_{\rho\sigma}^{(1)}(\mathrm{x}_2)\right\rangle=\frac{1}{\mathrm{x}_{12}^8}\left\{\left(X_\mu X_\nu-\frac{g_{\mu\nu}}{4}\right)\left(Y_\rho Y_\sigma-\frac{g_{\rho\sigma}}{4}\right)A(v)+\left(X_\mu Y_\rho I_{\nu\sigma}+X_\mu Y_\sigma I_{\nu\rho}\right.\right.$$
$$\left.\left.+X_\nu Y_\sigma I_{\mu\rho}+X_\nu Y_\rho I_{\mu\sigma}-g_{\mu\nu}Y_\rho Y_\sigma-g_{\rho\sigma}X_\mu X_\nu+\frac{1}{4}g_{\mu\nu}g_{\rho\sigma}\right)B(v)+\mathcal{I}_{\mu\nu\rho\sigma}C(v)\right\}$$

$$\mathcal{I}_{\mu\nu\rho\sigma}(\mathrm{x})\equiv\frac{1}{2}\left(\mathcal{I}_{\mu\rho}(\mathrm{x})\mathcal{I}_{\nu\sigma}(\mathrm{x})+\mathcal{I}_{\mu\sigma}(\mathrm{x})\mathcal{I}_{\nu\rho}(\mathrm{x})\right)-\frac{1}{4}g_{\mu\nu}g_{\rho\sigma}$$

$$A(v)=4\gamma_{\mathrm{k}}(6v^6+3v^4+v^2)$$
$$B(v)=-\gamma_{\mathrm{k}}(3v^6-v^4-2v^2)$$
$$C(v)=\gamma_{\mathrm{k}}v^2(v^2-1)^2,$$

$$\gamma_{\mathrm{k}}\equiv\frac{32c_{\mathrm{k}}}{9\pi^2g_{\mathrm{YM}}^2},c_{\mathrm{k}}=\begin{cases}k(c_1+c_2) & SU(4)\times SU(4)\\ c_n/8 & SO(5)\end{cases}$$

$$c=\frac{N_0+3N_{1/2}+12N_1}{120}=\frac{\pi^4C_T}{40},a=\frac{2N_0+11N_{1/2}+124N_1}{720}.$$

$$\mathcal{N}=4\mathrm{SYM}\left(N_0=6N_c^2,N_{1/2}=4N_c^2,N_1=N_c^2\right)$$

$$a=c=\frac{N_c^2}{4}=\frac{\pi^4C_T}{40}$$

$$\partial^\mu\Theta_{\mu\nu}(x)=\delta(x_3)\eta_\nu\mathcal{D}(\mathbf{x})$$

$$\mathcal{D}(\mathbf{x})=\lim_{x_3\to0+}\Theta_{33}(\mathbf{x},x_3)-\lim_{x_3\to0-}\Theta_{33}(\mathbf{x},x_3)$$

$$\mathbf{x}\equiv(x_0,x_1,x_2)$$

$$\left\langle\mathcal{D}^{(1)}(\mathbf{x}_1)\mathcal{D}^{(1)}(\mathbf{x}_2)\right\rangle=\lim_{\mathrm{z}_1,\mathrm{z}_2\to0^+}\left\langle\Theta_{33}^{(1)}(\mathrm{x}_1)\Theta_{33}^{(1)}(\mathrm{x}_2)\right\rangle=\frac{c_{\eta\eta}}{\mathbf{x}_{12}^8}$$

$$c_{\eta\eta}\equiv\frac{15b_2}{2\pi^4}=\frac{80c_{\mathrm{k}}N_c}{\pi^2\lambda}$$

$$b_2 = \frac{32\pi^2 c_{\mathrm{k}} N_c}{3\lambda} + \mathcal{O}(\lambda^0), c_{\mathrm{k}} = \begin{cases} k(c_1 + c_2) & SU(2) \times SU(2) \\ c_n/8 & SO(5) \end{cases}$$

$$c = \frac{\pi^4}{30} \cdot \alpha(0), b_2 = \frac{2\pi^4}{15} \cdot \alpha(1)$$

$$\mathcal{L}_{\mathcal{N}=4} = \frac{2}{g_{\mathrm{YM}}^2} \cdot \mathrm{tr}\left\{-\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2}\left(D_\mu\varphi_i\right)^2 + i\bar{\psi}_\alpha \not{D}\psi_\alpha + \frac{1}{4}\left[\varphi_i, \varphi_j\right]^2 + \right.$$
$$\left. + \sum_{i=1}^{3} \mathcal{G}_{\alpha\beta}^i \bar{\psi}_\alpha \left[\varphi_i, \psi_\beta\right] + \sum_{i=4}^{6} \mathcal{G}_{\alpha\beta}^i \bar{\psi}_\alpha \gamma_5 \left[\varphi_i, \psi_\beta\right]\right\}$$

$$\bar{\psi}_\alpha \equiv \psi_\alpha^\dagger \gamma^0, \not{D} \equiv \gamma^\mu D_\mu$$

$$A_\mu = A_\mu^a T^a, \varphi_i = \varphi_i^a T^a, \psi_{\alpha,m} = \psi_{\alpha,m}^a T^a, a = 1, \ldots, N_c^2$$

$$D^\mu F_{\mu\nu} = i[D_\nu \varphi_i, \varphi_i], D^\mu D_\mu \varphi_i = \left[\varphi_j, \left[\varphi_j, \varphi_i\right]\right]$$
$$i\not{D}\psi_\alpha = \sum_{i=1}^{3} \mathcal{G}_{\alpha\beta}^i \left[\psi_\beta, \varphi_i\right] + \sum_{i=4}^{6} \mathcal{G}_{\alpha\beta}^i \gamma_5 \left[\psi_\beta, \varphi_i\right]$$

$$T_{\mu\nu} = \frac{\partial \mathcal{L}}{\partial \partial^\mu A_\rho} \partial_\nu A_\rho + \frac{\partial \mathcal{L}}{\partial \partial^\mu \varphi_i} \partial_\nu \varphi_i + \frac{\partial \mathcal{L}}{\partial \partial^\mu \bar{\psi}_\alpha} \partial_\nu \bar{\psi}_\alpha + \frac{\partial \mathcal{L}}{\partial \partial^\mu \psi_\alpha} \partial_\nu \psi_\alpha - g_{\mu\nu}\mathcal{L}$$

$$T_{\mu\nu} \equiv \frac{2}{\sqrt{-g}} \cdot \frac{\delta \mathcal{S}}{\delta g^{\mu\nu}}, \mathcal{S} = \int dx^4 \sqrt{-g}\mathcal{L}$$

$$\Theta_{\mu\nu} = \frac{2}{g_{\mathrm{YM}}^2} \cdot \mathrm{tr}\left\{-F_\mu{}^\varrho F_{\nu\varrho} - \frac{2}{3}\left(D_\mu\varphi_i\right)\left(D_\nu\varphi_i\right) + \frac{1}{3}\varphi_i D_{(\mu} D_{\nu)} \varphi_i + \frac{i}{2}\bar{\psi}_\alpha \gamma_{(\mu} \overleftrightarrow{D}_{\nu)} \psi_\alpha\right\}$$
$$- \frac{2}{g_{\mathrm{YM}}^2} \cdot \mathrm{tr}\left\{-\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{6}\left(D_\mu\varphi_i\right)^2 - \frac{1}{12}\left[\varphi_i, \varphi_j\right]^2\right\} \cdot g_{\mu\nu},$$

$$a_{(\mu\nu)} \equiv \left(a_{\mu\nu} + a_{\nu\mu}\right)/2, \text{ and } f \overleftrightarrow{\partial}_\mu g \equiv f\left(\partial_\mu g\right) - \left(\partial_\mu f\right) g$$

$$\Theta_{\mu\nu} = \Theta_{\nu\mu}, g^{\mu\nu}\Theta_{\mu\nu} = 0, \partial^\mu \Theta_{\mu\nu} = 0$$

$$\Theta_{\mu\nu}^{\text{scalar}} = \frac{2}{g_{\mathrm{YM}}^2} \mathrm{tr}\left\{-\frac{2}{3}\left(\partial_\mu\varphi_i\right)\left(\partial_\nu\varphi_i\right) + \frac{1}{3}\varphi_i\left(\partial_\mu\partial_\nu\varphi_i\right) + \frac{1}{6}g_{\mu\nu}\left[\left(\partial_\varrho\varphi_i\right)^2 + \frac{1}{2}\left[\varphi_i, \varphi_j\right]^2\right]\right\}.$$

$$\left[\tilde{\varphi}_i\right]_{n_1,n_2} = \sum_{\ell_1=0}^{k_1-1} \sum_{\ell_2=0}^{k_2-1} \sum_{m_1=-\ell_1}^{\ell_1} \sum_{m_2=-\ell_2}^{\ell_2} \left(\tilde{\varphi}_i\right)_{\ell_1,m_1;\ell_2,m_2} \cdot \left[\hat{Y}_{\ell_1}^{m_1} \otimes \hat{Y}_{\ell_2}^{m_2}\right]_{n_1,n_2},$$

$$\left\langle \tilde{\varphi}_i \tilde{\varphi}_{j+3} \right\rangle, \left\langle \tilde{\varphi}_i \tilde{\varphi}_j \right\rangle, \left\langle \tilde{\varphi}_{i+3} \tilde{\varphi}_{j+3} \right\rangle, \text{for } i, j = 1,2,3$$

$$\left\langle \left(\tilde{\varphi}_i\right)_{\ell_1 m_1;\ell_2 m_2} \left(\tilde{\varphi}_{j+3}\right)_{\ell_1' m_1';\ell_2' m_2'} \right\rangle = (-1)^{m_1'+m_2'} \delta_{\ell_1\ell_1'} \delta_{\ell_2\ell_2'} \left[t_i^{(\ell_1)}\right]_{m_1,-m_1'} \left[t_{j+3}^{(\ell_2)}\right]_{m_2,-m_2'} K_{\text{opp}}^{\varphi}$$

$$K_{\text{opp}}^{\varphi} \equiv \frac{K^{m_-^2}}{N_-} - \frac{K^{m_0^2}}{N_0} + \frac{K^{m_+^2}}{N_+}, N_{\pm} \equiv \lambda_{\mp}(\lambda_{\mp} - \lambda_{\pm}), N_0 \equiv -\lambda_+\lambda_-$$

$$\lambda_{\pm} \equiv -\frac{1}{2} \pm \sqrt{\ell_1(\ell_1+1) + \ell_2(\ell_2+1) + \frac{1}{4}},$$

$$m_0^2 \equiv \ell_1(\ell_1+1) + \ell_2(\ell_2+1) + 2, m_{\pm}^2 \equiv \ell_1(\ell_1+1) + \ell_2(\ell_2+1) - 2\lambda_{\pm}.$$

$$\left\langle (\tilde{\varphi}_i)_{\ell_1 m_1;\ell_2 m_2} (\tilde{\varphi}_j)_{\ell_1' m_1';\ell_2' m_2'} \right\rangle = (-1)^{m_1'+m_2'} \delta_{\ell_1\ell_1'} \delta_{\ell_2\ell_2'} \delta_{m_2+m_2'} \left[\delta_{ij} \delta_{m_1+m_1'} K_{\text{sing}}^{\varphi,\ell_1} - i\epsilon_{ijk} \left[t_k^{(\ell_1)} \right]_{m_1,-m_1'} K_{\text{anti}}^{\varphi,\ell_1} - \left[t_i^{(\ell_1)} t_j^{(\ell_1)} \right]_{m_1,-m_1'} K_{\text{sym}}^{\varphi,\ell_1} \right]$$

$$\left\langle (\tilde{\varphi}_{i+3})_{\ell_1 m_1;\ell_2 m_2} (\tilde{\varphi}_{j+3})_{\ell_1' m_1';\ell_2' m_2'} \right\rangle = (-1)^{m_1'+m_2'} \delta_{\ell_1\ell_1'} \delta_{\ell_2\ell_2'} \delta_{m_1+m_1'} \left[\delta_{ij} \delta_{m_2+m_2'} K_{\text{sing}}^{\varphi,\ell_2} - i\epsilon_{ijk} \left[t_k^{(\ell_2)} \right]_{m_2,-m_2'} K_{\text{anti}}^{\varphi,\ell_2} - \left[t_i^{(\ell_2)} t_j^{(\ell_2)} \right]_{m_2,-m_2'} K_{\text{sym}}^{\varphi,\ell_2} \right],$$

$$K_{\text{sing}}^{\varphi,\ell_i} \equiv \frac{\ell_i+1}{2\ell_i+1} \cdot K^{m_{i,+}^2} + \frac{\ell_i}{2\ell_i+1} \cdot K^{m_{i,-}^2}, K_{\text{anti}}^{\varphi,\ell_i} \equiv \frac{K^{m_{i,+}^2}}{2\ell_i+1} - \frac{K^{m_{i,-}^2}}{2\ell_i+1}$$

$$K_{\text{sym}}^{\varphi,\ell_1} \equiv \frac{K^{m_{1,+}^2}}{(2\ell_1+1)(\ell_1+1)} + \frac{K^{m_{1,-}^2}}{(2\ell_1+1)\ell_1} - \frac{\ell_2(\ell_2+1)}{\ell_1(\ell_1+1)} \cdot \frac{K^{m_0^2}}{N_0} - \frac{K^{m_-^2}}{N_-} - \frac{K^{m_+^2}}{N_+}$$

$$K_{\text{sym}}^{\varphi,\ell_2} \equiv \frac{K^{m_{2,+}^2}}{(2\ell_2+1)(\ell_2+1)} + \frac{K^{m_{2,-}^2}}{(2\ell_2+1)\ell_2} - \frac{\ell_1(\ell_1+1)}{\ell_2(\ell_2+1)} \cdot \frac{K^{m_0^2}}{N_0} - \frac{K^{m_-^2}}{N_-} - \frac{K^{m_+^2}}{N_+},$$

$$m_{1,+}^2 = \ell_1(\ell_1-1) + \ell_2(\ell_2+1), \quad m_{1,-}^2 = (\ell_1+1)(\ell_1+2) + \ell_2(\ell_2+1)$$
$$m_{2,+}^2 = \ell_1(\ell_1+1) + \ell_2(\ell_2-1), \quad m_{2,-}^2 = \ell_1(\ell_1+1) + (\ell_2+1)(\ell_2+2)$$

$$\left[t_i^{(\ell)} \right]_{m,-m'} \equiv \left[t_i^{(2\ell+1)} \right]_{\ell-m+1,\ell+m'+1}, i = 1,2,3$$

$$t_1^{(k)} = \frac{(-1)^{k+1}}{2} \sqrt{\frac{k(k^2-1)}{6}} \cdot \left(\hat{Y}_1^{-1} - \hat{Y}_1^1 \right)$$

$$t_2^{(k)} = \frac{i(-1)^{k+1}}{2} \sqrt{\frac{k(k^2-1)}{6}} \cdot \left(\hat{Y}_1^{-1} + \hat{Y}_1^1 \right)$$

$$t_3^{(k)} = \frac{(-1)^{k+1}}{2} \sqrt{\frac{k(k^2-1)}{3}} \cdot \hat{Y}_1^0$$

$$\left(\hat{Y}_\ell^m \right)^\dagger = (-1)^m \hat{Y}_\ell^{-m} \quad \text{tr} \left[\hat{Y}_\ell^m \hat{Y}_{\ell'}^{m'} \right] = (-1)^{m'} \delta_{\ell\ell'} \delta_{m+m'}$$

$$[\tilde{\varphi}_i]_{n_1,n_2} = \sum_{\mathbf{L}} (\tilde{\varphi}_i)_{\mathbf{L}} \cdot \left[\hat{Y}_{\mathbf{L}} \right]_{n_1,n_2}, \text{tr} \left(\hat{Y}_{\mathbf{L}'}^\dagger \hat{Y}_{\mathbf{L}} \right) = \delta_{\mathbf{L}',\mathbf{L}}$$

where $i = 1, \dots, 5$ and $n_1, n_2 = 1, \dots, d_G \equiv (n+1)(n+2)(n+3)/6$

$$\mathbf{L} = \{(L_1, L_2), \ell_1, \ell_2, m_1, m_2\}$$

$$-L_1+L_2\le \ell_1-\ell_2\le L_1-L_2\le \ell_1+\ell_2\le L_1+L_2$$
$$\ell_1+\ell_2\in\mathbb{Z}, m_i=-\ell_i,\dots,\ell_i$$

$$\left\langle(\tilde{\varphi}_i)_{\mathbf{L}}\left(\tilde{\varphi}_j\right)^{\dagger}_{\mathbf{L}'}\right\rangle=\delta_{ij}\delta_{\mathbf{L},\mathbf{L}'}\hat{f}^{\text{sing}}+\langle\mathbf{L}|L_{ij}|\mathbf{L}'\rangle\hat{f}^{\text{lin}}+\langle\mathbf{L}|\{L_{ik},L_{jl}\}L_{kl}|\mathbf{L}'\rangle\hat{f}^{\text{cubic}}$$
$$+\langle\mathbf{L}|\{L_{ik},L_{kj}\}|\mathbf{L}'\rangle\hat{f}_5^{\text{sym}}+\langle\mathbf{L}|\{L_{i6},L_{6j}\}|\mathbf{L}'\rangle\cdot\left[\delta_{L_1,L_1'}\delta_{L_2,L_2'}\hat{f}_6^{\text{sym}}\right.$$
$$\left.+\delta_{L_1',L_1\pm1}\delta_{L_2',L_2\mp1}\hat{f}^{\text{opp}}\right]$$

$$\langle\mathbf{L}|L_{ij}|\mathbf{L}'\rangle=\text{tr}\left(\hat{Y}^{\dagger}_{\mathbf{L}}L_{ij}\hat{Y}_{\mathbf{L}'}\right)=\frac{i}{2}\cdot\text{tr}\left(\hat{Y}^{\dagger}_{\mathbf{L}}\left[\hat{Y}_{\mathbf{L}'},G_{ij}\right]\right),G_{i6}=-G_{6i}\equiv G_i$$

$$\left[L_{ij},L_{kl}\right]=i\left(\delta_{ik}L_{jl}+\delta_{jl}L_{ik}-\delta_{jk}L_{il}-\delta_{il}L_{jk}\right),i,j,k,l=1,\dots,5.$$

$$\left\langle\left(\tilde{\varphi}_6\right)_{\mathbf{L}}\left(\tilde{\varphi}_6\right)^{\dagger}_{\mathbf{L}'}\right\rangle=\delta_{\mathbf{L},\mathbf{L}'}\cdot K^{\hat{m}^2_{\text{easy}}},\hat{m}^2_{\text{easy}}\equiv 2L_1L_2+L_1+2L_2$$

$$G_1=\sqrt{\frac{c_n}{10}}\cdot\left(\hat{Y}_{++}+\hat{Y}_{--}\right),\qquad G_2=-i\sqrt{\frac{c_n}{10}}\cdot\left(\hat{Y}_{++}-\hat{Y}_{--}\right)$$
$$G_3=-\sqrt{\frac{c_n}{10}}\cdot\left(\hat{Y}_{-+}-\hat{Y}_{+-}\right),\qquad G_4=-i\sqrt{\frac{c_n}{10}}\cdot\left(\hat{Y}_{-+}+\hat{Y}_{+-}\right)$$
$$G_5=-\sqrt{\frac{c_n}{5}}\cdot\hat{Y}_{00}$$

$$\hat{Y}_{\alpha\beta}\equiv\hat{Y}_{\left(\frac{1}{2},\frac{1}{2}\right)\frac{11}{22}\alpha\beta},\hat{Y}_{00}\equiv\hat{Y}_{\left(\frac{1}{2},\frac{1}{2}\right)0000}$$

$$\tilde{\varphi}_i\cdot\text{tr}\left[G_j\tilde{\varphi}_j\right]=K^{5/2}(x,y)\cdot G_i$$

$$G_{\Delta}(x,z;y,w)=\frac{\Gamma(\Delta)\tilde{\eta}^{\Delta}}{2^{\Delta+1}\pi^{d/2}\Gamma\left(\Delta-\frac{d}{2}+1\right)}\cdot\ {}_2F_1\left(\frac{\Delta}{2},\frac{\Delta+1}{2},\Delta-\frac{d}{2}+1,\tilde{\eta}^2\right)$$

$$\tilde{\eta}\equiv\frac{2zw}{z^2+w^2+(x-y)^2},\tilde{m}^2\equiv\Delta(\Delta-d)\equiv m^2-\frac{d^2-1}{4}$$

$${}_2F_1\left(\frac{\Delta}{2},\frac{\Delta+1}{2},\Delta-\frac{d}{2}+1,\tilde{\eta}^2\right)=\left(1+\frac{1}{2s}\right)^{\nu+\frac{d}{2}}\ {}_2F_1\left(\nu+\frac{d}{2},\nu+\frac{1}{2},2\nu+1,-s^{-1}\right),$$

$$s\equiv\frac{1-\tilde{\eta}}{2\tilde{\eta}}=\frac{(z-w)^2+(x-y)^2}{4zw},\nu\equiv\Delta-\frac{d}{2}=\sqrt{m^2+\frac{1}{4}}$$

$${}_2F_1\left(\frac{\Delta}{2},\frac{\Delta+1}{2},\Delta-\frac{d}{2}+1,\tilde{\eta}^2\right)$$

$$=\left(1+\frac{1}{s}\right)^{\frac{1-d}{2}}\left(1+\frac{1}{2s}\right)^{\nu+\frac{d}{2}}\ {}_2F_1\left(\nu-\frac{d}{2}+1,\nu+\frac{1}{2},2\nu+1,-s^{-1}\right)$$

$$G_{\Delta}(x,z;y,w)=\frac{1}{2^d\pi^{\frac{d+1}{2}}}\frac{\Gamma\left(\nu+\frac{d}{2}\right)\Gamma\left(\nu+\frac{3}{2}\right)}{\Gamma(2\nu+2)}\cdot\frac{{}_2F_1\left(\nu-\frac{d}{2}+1,\nu+\frac{1}{2},2\nu+1,-s^{-1}\right)}{(1+s)^{\frac{d-1}{2}}s^{\nu+\frac{1}{2}}}$$

$$\Gamma(\nu+1)=\frac{\sqrt{\pi}}{2^{2\nu+1}}\cdot\frac{\Gamma(2\nu+2)}{\Gamma\left(\nu+\frac{3}{2}\right)}$$

$$K^{m^2}(x,y)\equiv\frac{g_{\mathrm{YM}}^2}{2}\cdot(x_3y_3)^{-\frac{d-1}{2}}\cdot G_\Delta(\{x_0,x_1,x_2\},x_3;\{y_0,y_1,y_2\},y_3)$$

$$K^\nu(x,y)=\frac{g_{\mathrm{YM}}^2}{16\pi^2}\frac{1}{\binom{2\nu+1}{\nu+\frac{1}{2}}}\frac{{}_2F_1\left(\nu-\frac{1}{2},\nu+\frac{1}{2},2\nu+1;-\xi^{-1}\right)}{(1+\xi)\xi^{\nu+\frac{1}{2}}}\cdot\frac{1}{x_3y_3},$$

$$\xi\equiv\frac{|x-y|^2}{4x_3y_3},\nu\equiv\sqrt{m^2+\frac{1}{4}}$$

$$S_{\text{Einstein-Hilbert}}=\frac{1}{16\pi G_{\mathrm{N}}}\int\ \mathrm{d}^4x\sqrt{g}(2\Lambda-R)+S_{\mathrm{gf}}+S_{\mathrm{gh}}$$

$$\sqrt{g}=\sqrt{|\det g_{\mu\nu}|}$$

$$g_{\mu\nu}=\eta_{\mu\nu}+\sqrt{G_N}h_{\mu\nu}$$

$$S_{\text{matter}}=\int\ \mathrm{d}^4x\sqrt{g}\Big(\frac{1}{2}\partial_\mu\phi\partial^\mu\phi-y_t\phi\bar{\psi}\psi-\frac{1}{4}F^{\mu\nu}F_{\mu\nu}+\bar{\psi}\big(i\nabla-m_\psi\big)\psi\Big)+S_{\mathrm{gf,U(1)}}$$

$$\nabla\psi=g_{\mu\nu}\gamma(x)^\mu\nabla^\nu\psi=g_{\mu\nu}\gamma(x)^\mu(D^\nu+\Gamma(x)^\nu)\psi,$$

$$D_\mu=\partial_\mu-ig_YA_\mu$$

$$\mathcal{G}_h^{\mu\nu\rho\sigma}=\sum_{i=1}^{5}c_iG_h\mathcal{T}_i^{\mu\nu\rho\sigma}$$

$$G_h=1/(p^2-2\Lambda)$$

$$G_\Phi\equiv G_\Phi(p)$$

$$G_\Phi(|\vec{p}|,p_0)=\int_0^\infty\frac{\lambda\mathrm{d}\lambda}{\pi}\frac{\rho_\Phi(\lambda,|\vec{p}|)}{\lambda^2+p_0^2}$$

$$\rho_\Phi(\lambda,|\vec{p}|)=\lim_{\epsilon\to0^+}2\mathrm{Im}G_\Phi(p_0=-i(\lambda+i\epsilon),|\vec{p}|)$$

$$R_{k,\Phi}=Z_\Phi k^2$$

$$R_{k,\psi}=Z_\psi k\mathbb{1}$$


$$\partial_t\Gamma_k[\Phi]=\frac{1}{2}\mathrm{Tr}\mathcal{G}_k[\Phi]\partial_tR_k-\partial_tS_{\mathrm{ct},k}[\Phi]$$

$$\mathcal{G}_k[\Phi] = \frac{1}{\Gamma_k^{(2)}[\Phi] + R_k},$$


$$\Gamma_k^{(2)} = \delta^2\Gamma_k/\delta\Phi\delta\Phi$$

$$\partial_t\Gamma_k^{(\Phi\Phi)}\Big|_{p=0} = 0, \partial_{p^2}\left(\partial_t\Gamma_k^{(\Phi\Phi)}\Big|_{p=0}\right) = 0$$

$$\partial_t\Gamma_k^{(AA)} =$$



$$\partial_t\Gamma_k^{(\phi\phi)} =$$



$$\partial_t\rho_\Phi(\lambda) = -2\mathrm{Im}G_\Phi^2\left(\partial_t\Gamma_k^{(\Phi\Phi)} + \partial_t R_k\right),$$

$$\partial_t\Gamma_k^{(\Phi\Phi)}(p) \propto \prod_{i=1}^{3}\int_0^\infty \frac{\lambda_i\,\mathrm{d}\lambda_i}{\pi}\rho_i(\lambda_i)\int\frac{\mathrm{d}^d q}{(2\pi)^d}I(p,q,\lambda_i),$$

$$\theta(\lambda^2 - (m_\Phi + m_h)^2)$$

$$\theta\left(\lambda^2 - 4m_\psi^2\right)$$

$$\rho_\Phi = \frac{1}{Z_\Phi}\left[2\pi\delta\left(\lambda^2 - m_\Phi^2\right) + \theta(\lambda^2 - (m_\Phi + m_h)^2)f_{\Phi,\mathrm{grav}}(\lambda)\right.$$
$$\left.+\theta\left(\lambda^2 - 4m_\psi^2\right)f_{\Phi,\mathrm{ferm}}(\lambda)\right]$$

$$\Phi = \{A, \phi\}$$

$$Z_\Phi\left(p^2 = -m_\Phi^2\right)$$

$$\eta_\Phi = -\partial_t \ln Z_\Phi.$$

$$m_\Phi^2 = k^2(1 + \mu_\Phi(k)).$$

$$m_\psi = k\left(1 + \mu_\psi(k)\right).$$

$$G_\mathrm{N}, \Lambda, y_t, g_Y, m_A, m_\phi, m_\psi$$

$$\mu_\psi = \frac{c_1 k + c_2 M_\mathrm{Pl}}{k}$$

$$g_\mathrm{N} = k^2 G_\mathrm{N}, \qquad \mu_h = -2\Lambda/k^2$$

$$\beta_{y_t} = \frac{9}{32\pi^2} y_t^3 - f_y g_{\mathrm{N}} y_t$$
$$\beta_{g_Y} = \frac{41}{96\pi^2} g_Y^3 - f_g g_{\mathrm{N}} g_Y$$

$$g_Y(k = 10^{-4} M_{\mathrm{Pl}}) = 0.43$$
$$y_t(k = 10^{-4} M_{\mathrm{Pl}}) = 0.44$$

$$g_Y^* = 0.455, y_t^* = 0.462$$

$$Z_h, Z_\phi, Z_A, Z_\psi,$$

$$p^2 = -m_\Phi^2$$

$$y_t, g_Y, \mu_\psi,$$

$$g_{\mathrm{N}}, \mu_h, \mu_\phi, \mu_A, Z_h, Z_\phi, Z_A,$$

$$\left(g_{\mathrm{N}}^*, \mu_h^*, \mu_A^*, \mu_\phi^*\right) = (1.06, -0.34, -0.17, -0.018).$$


$$\left(\eta_h^*, \eta_A^*, \eta_\phi^*\right) = (0.96, 0.52, 0.045).$$


$$M_{ij} = \frac{\partial \beta_{\xi_i}}{\partial_{\xi_j}},$$


$$\theta_{1,2} = 2.49 \pm 3.17i$$


$$\left(\theta_{\mu_A}, \theta_{\mu_\phi}\right) = (2.55, 2.15).$$


$$\left(G_{\mathrm{N}}(k), k^2 \mu_h(k), k^2 \mu_A(k), k^2 \mu_\phi(k)\right)\Big|_{k \to 0}$$






0.47
0.46
0.45
0.44
0.43
y_t
$f_y = 0.00573$
$f_y = 0.007$
$f_y = 0.009$
$f_y = 0.11$
10^{-3} 10^{-2} 10^{-1} 10^{0} 10^{1} 10^{2} 10^{3}
$k\ [M_{\rm Pl}]$



10^{0}
10^{-1}
10^{-2}
10^{-3}
10^{-4}
$\sim k^2$
$g_{\rm N}$
$-\mu_h$
$-\mu_A$
$-\mu_\phi$
10^{-1} 10^{0} 10^{1} 10^{2}
$k\ [M_{\rm Pl}]$



10^{0}
10^{-1}
10^{-2}
$\sim k^{-\eta_\phi^*}$
$\sim k^{-\eta_A^*}$
$\sim k^{-\eta_h^*}$
Z_ϕ
Z_A
Z_h
10^{-2} 10^{-1} 10^{0} 10^{1} 10^{2}
$k\ [M_{\rm Pl}]$



10^{1}
10^{0}
10^{-1}
10^{-2}
10^{-3}
$|\rho_A|\,[1/M_{\rm Pl}^2]$
10^{-2}
10^{-1}
10^{0}
10^{1}
10^{2}
10^{3}
$\lambda\,[M_{\rm Pl}]$
$\sim \lambda^{\eta_A^*-2}$


10^{2}
10^{1}
10^{0}
10^{-1}
10^{-2}
10^{-3}
$|\rho_\phi|\,[1/M_{\rm Pl}^2]$
10^{-2}
10^{-1}
10^{0}
10^{1}
10^{2}
$\lambda\,[M_{\rm Pl}]$
$\sim \lambda^{\eta_\phi^*-2}$


10^{1}
10^{0}
10^{-1}
10^{-2}
$\rho_{A,\rm grav,1}\,[1/M_{\rm Pl}^2]$
10^{-2}
10^{-1}
10^{0}
10^{1}
10^{2}
$\lambda\,[M_{\rm Pl}]$
$\sim \lambda^{\eta_A^*-2}$


10^0
10^{-1}
10^{-2}
10^{-3}
$\rho_{\phi,\mathrm{grav},1}\ [1/M_{\mathrm{Pl}}^2]$
$\sim \lambda^{\eta_\phi^*-2}$
10^{-2}
10^{-1}
10^0
10^1
10^2
$\lambda\ [M_{\mathrm{Pl}}]$


10^1
10^0
10^{-1}
10^{-2}
$|\rho_{A,\mathrm{grav},2}|\ [1/M_{\mathrm{Pl}}^2]$
$\sim \lambda^{\eta_A^*-2}$
10^{-2}
10^{-1}
10^0
10^1
10^2
$\lambda\ [M_{\mathrm{Pl}}]$


10^0
10^{-1}
10^{-2}
10^{-3}
$|\rho_{\phi,\mathrm{grav},2}|\ [1/M_{\mathrm{Pl}}^2]$
$\sim \lambda^{\eta_\phi^*-2}$
10^{-2}
10^{-1}
10^0
10^1
10^2
$\lambda\ [M_{\mathrm{Pl}}]$

$$\#_{SYM} = \int (G_{\mathrm{N}}, -2\Lambda, 0, \omega_\phi),$$

$$\left(Z_h(k), Z_A(k), Z_\phi(k)\right)\big|_{k\to 0} = (1,1,1).$$

$$\int_0^\infty \lambda \rho_i(\lambda)\mathrm{d}\lambda \neq \text{ finite}$$


$$\rho_{A,\mathrm{IR}}\big|_{\mathrm{grav}} = -\frac{8}{3}\frac{\beta^2 - 3\alpha}{(\beta-3)^2}$$
$$\rho_{\phi,\mathrm{IR}}\big|_{\mathrm{grav}} = 6(\beta-1)\frac{\alpha(\beta-5)-\beta+9}{(\beta-3)^2}.$$


$$\rho_{A,\mathrm{IR}}\big|_{\mathrm{exact}} = \frac{4}{3},\ \rho_{\phi,\mathrm{IR}}\big|_{\mathrm{exact}} = 0,$$

$$\rho_{A,\mathrm{IR}}\big|_{\bullet} \approx 1.22,\ \rho_{\phi,\mathrm{IR}}\big|_{\bullet} \approx 0.105.$$

$$\rho_{\Phi,UV,\mathrm{diag1}}(\lambda) \propto \#_1(2-\eta_h^*)\lambda^{\eta_\Phi^*-2},$$
$$\rho_{\Phi,UV,\mathrm{diag2}}(\lambda) \propto \#_2(2-\eta_\Phi^*)\lambda^{\eta_\Phi^*-2},$$

$$\eta_h^*\big|_{\mathrm{gauge}} \lesssim 0.3,\ \eta_h^*\big|_{\mathrm{scalar}} \lesssim -54.$$





$$\rho_{\Phi,\mathrm{UV,grav}}(\lambda) \propto \#_1 g_{\mathrm{N}}^* \lambda^{\eta_\Phi^*-2}$$

$$\rho_{\Phi,\mathrm{UV,\ particle\ (f)}}(\lambda) \propto \#_2 \xi^{*2} N_f \lambda^{\eta_\Phi^*-2}$$

$$N_f \frac{(g_Y^*)^2}{g_N^*} \gtrsim \mathcal{O}(100), N_f \frac{(y_t^*)^2}{g_N^*} \gtrsim \mathcal{O}(1000)$$

$$\theta_{g/y} = -f_{g/y} \int_{\bullet}^{\otimes} g_{\mathrm{N}}^* \int_{\bullet}^{\otimes} \lambda^{\eta_A^*-2-\theta_g} \int_{\bullet}^{\otimes} \lambda^{\eta_\phi^*-2-\theta_y}$$

$$\Gamma = \int \mathrm{d}^4x\sqrt{g}\left(\frac{1}{2}\phi f_{\phi\phi}(\Box)\phi - \frac{1}{4}F_{\mu\nu}f_{FF}(\Box)F^{\mu\nu}\right),$$

$$f_{\phi\phi} = \Box \text{ and } f_{FF} = 1$$

$$f_{\phi\phi}(p^2) = G_\phi^{-1}(p^2) = \left(\int_0^\infty \frac{\lambda \mathrm{d}\lambda}{\pi}\frac{\rho_\phi(\lambda)}{\lambda^2+p^2}\right)^{-1},$$

$$f_{FF}(p^2) = p^{-2}G_A^{-1}(p^2) = \left(p^2\int_0^\infty \frac{\lambda \mathrm{d}\lambda}{\pi}\frac{\rho_A(\lambda)}{\lambda^2+p^2}\right)^{-1}$$

$$= Z_A(s)\bar{\Gamma}^{(\psi\bar{\psi}A)} \cdot G_A(s) \cdot \bar{\Gamma}^{(\psi\bar{\psi}A)}$$
$$\equiv S^{(\psi\bar{\psi}A)} \cdot G_{A,\ \mathrm{scat}}(s) \cdot S^{(\psi\bar{\psi}A)}$$

$$\Gamma^{(\psi\bar{\psi}A)} = \sqrt{Z_A(p_A)Z_\psi(p_\psi)Z_\psi(p_{\bar{\psi}})}\bar{\Gamma}^{(\psi\bar{\psi}A)}$$

$$S_{\mathrm{gh}}[\bar{g},\phi] = \int \mathrm{d}^4x\sqrt{\bar{g}}\bar{c}^\mu M_{\mu\nu}c^\nu$$

$$M_{\mu\nu} = \bar{\nabla}^\rho\left(g_{\mu\nu}\nabla_\rho + g_{\rho\nu}\nabla_\mu\right) - \frac{1+\beta}{2}\bar{g}^{\sigma\rho}\bar{\nabla}_\mu g_{\nu\sigma}\nabla_\rho$$

$$S_{\mathrm{gf,U(1)}} = \frac{1}{2\chi}\int \mathrm{d}^4x\sqrt{\bar{g}}\left(\bar{D}^\mu A_\mu\right)$$

$$\mathcal{G}_A^{\mu\nu} = G_A \Pi_{\mathrm{T}}^{\mu\nu}.$$

$$\Pi_{\mathrm{T}}^{\mu\nu} = \eta^{\mu\nu} - p^\mu p^\nu / p^2$$

$$G_A = 1/p^2$$

$$\mathcal{G}_\phi = G_\phi = \frac{1}{p^2 + \omega_\phi^2},$$

$$S_{\mathrm{gf}}[\bar{g}, h] = \frac{1}{2\alpha}\int \mathrm{d}^4x\sqrt{\bar{g}}\bar{g}^{\mu\nu}F_\mu F_\nu$$

$$F_\mu[\bar{g}, h] = \bar{\nabla}^\nu h_{\mu\nu} - \frac{1+\beta}{4}\bar{\nabla}_\mu h_\nu^\nu.$$

$$\mathcal{G}_\psi = G_\psi\left(\not{p} + i m_\psi \mathbb{1}\right),$$

$$G_\psi = 1/\left(p^2 + m_\psi^2\right)$$

$$\partial_t g_{\mathrm{N}} = (2+3\eta_h)g_{\mathrm{N}} + \frac{g_{\mathrm{N}}^2}{\pi}\left(-\frac{47(6-\eta_h)}{114(1+\mu_h)^2} + \frac{5(8-\eta_h)}{38(1+\mu_h)^3} + \frac{49(10-\eta_h)}{570(1+\mu_h)^4} - \frac{598}{285(1+\mu_h)^5} - \frac{5}{19}\right)$$

$$\begin{aligned}\partial_t \mu_h = {} & -2\mu_h - \eta_h + g_{\mathrm{N}}(1+\mu_h)(2-\eta_h)\frac{5(5\sqrt{3}\pi - 22)}{18\pi} \\ & + \frac{2g_{\mathrm{N}}}{3\pi(1+\mu_h)}\left(23 + 16\mu_h - 7\mu_h^2 + 3\sqrt{\frac{1+\mu_h}{\mu_h - 3}}\left(13 - 6\mu_h + \mu_h^2\right)\operatorname{arcosh}\left[\frac{1}{2}(1-\mu_h)\right]\right)\end{aligned}$$

$$\begin{aligned}\eta_h = {} & g_{\mathrm{N}}(2-\eta_h)\frac{5\pi\sqrt{3} + 147}{54\pi} \\ & - \frac{2g_{\mathrm{N}}}{3\pi(\mu_h - 3)(\mu_h + 1)}\left(4\left(\mu_h^2 + \mu_h - 15\right) + \frac{3\left(\mu_h\left((\mu_h - 3)\mu_h - 13\right) + 31\right)\operatorname{arcosh}\left[\frac{1}{2}(1-\mu_h)\right]}{\sqrt{(\mu_h - 3)(\mu_h + 1)}}\right).\end{aligned}$$

$$\partial_t \mu_\phi = -2\mu_\phi - \eta_\phi$$

$$-\frac{g_{\mathrm{N}}\left((\eta_\phi-2)(1+\mu_h)^2+(8+\eta_h-5\eta_\phi)(1+\mu_h)(1+\mu_\phi)-2(\eta_h-2)(1+\mu_\phi)^2\right)\operatorname{artanh}\left[\sqrt{\frac{1+\mu_h}{-3+\mu_h-4\mu_\phi}}\right]}{\pi\sqrt{(1+\mu_h)(-3+\mu_h-4\mu_\phi)}}$$

$$-\frac{g_{\mathrm{N}}\left((\eta_h-2)(1+\mu_h)^2+(8+\eta_h-5\eta_\phi)(1+\mu_h)(1+\mu_\phi)-2(\eta_h-2)(1+\mu_\phi)^2\right)\operatorname{artanh}\left[\frac{-1+\mu_h-2\mu_\phi}{\sqrt{(1+\mu_h)(-3+\mu_h-4\mu_\phi)}}\right]}{\pi\sqrt{(1+\mu_h)(-3+\mu_h-4\mu_\phi)}}$$

$$-\frac{g_{\mathrm{N}}}{\pi(\mu_h-\mu_\phi)^2}\Big((1+\mu_\phi)(\mu_\phi-\mu_h)\left((\eta_\phi-2)(1+\mu_h)+(2+\eta_h-2\eta_\phi)(1+\mu_\phi)\right)+\frac{1}{2}\ln\left(\frac{1+\mu_h}{1+\mu_\phi}\right)\left((\eta_\phi-2)(1+\mu_h^3)\right.$$
$$\left.+(8+\eta_h-5\eta_\phi)(1+\mu_h)^2(1+\mu_\phi)+(-2-2\eta_h+3\eta_\phi)(1+\mu_h)(1+\mu_\phi)^2+(-4+3\eta_h-\eta_\phi)(1+\mu_\phi)^3\right)\Big)$$

$$\partial_t \mu_A = -2\mu_A - \eta_A$$

$$+\frac{g_{\mathrm{N}}(2-\eta_h)}{2\pi(1+\mu_A)(1+\mu_h)}\Bigg(\frac{2(-1-2\mu_A+\mu_h)^3}{\sqrt{\frac{-3-4\mu_A+\mu_h}{1+\mu_h}}}\left(\operatorname{arcoth}\left[\sqrt{1-\frac{4(1+\mu_A)}{1+\mu_h}}\right]+\operatorname{arcoth}\left[\frac{\sqrt{(-3-4\mu_A+\mu_h)(1+\mu_h)}}{1+2\mu_A-\mu_h}\right]\right)$$
$$+\frac{(1+\mu_h)}{(\mu_A-\mu_h)^2}(2(1+\mu_A)(\mu_A-\mu_h)^3+3(1+\mu_A)^2(3\mu_A^2+\mu_A(2-4\mu_h)+(-2+\mu_h)\mu_h)$$
$$-(12(1+\mu_A)^4-16(1+\mu_A)^3(1+\mu_h)+15(1+\mu_A)^2(1+\mu_h)^2-6(1+\mu_A)(1+\mu_h)^3+(1+\mu_h)^4)\ln\left(\frac{1+\mu_A}{1+\mu_h}\right)\Bigg)$$

$$+\frac{g_{\mathrm{N}}(2-\eta_A)}{2\pi(1+\mu_A)}\Bigg(-\frac{2(1+\mu_h)(13+20\mu_A^2-8\mu_A(-4+\mu_h)+(-6+\mu_h)\mu_h)}{\sqrt{(1+\mu_h)(-3-4\mu_A+\mu_h)}}$$

$$\times\left(\operatorname{artanh}\left[\frac{1+2\mu_A-\mu_h}{\sqrt{(-3-4\mu_A+\mu_h)(1+\mu_h)}}\right]+\operatorname{artanh}\left[\sqrt{\frac{1+\mu_h}{-3-4\mu_A+\mu_h}}\right]\right)$$
$$-\frac{1}{(\mu_A-\mu_h)^2}(13(1+\mu_A)^4+2(1+\mu_A)(\mu_A-\mu_h)^3-20(1+\mu_A)^3(1+\mu_h)+7(1+\mu_A)^2(1+\mu_h)^2$$
$$+(4(1+\mu_A)^4-14(1+\mu_A)^3(1+\mu_h)+23(1+\mu_A)^2(1+\mu_h)^2-8(1+\mu_A)(1+\mu_h)^3+(1+\mu_h)^4)\ln\left(\frac{1+\mu_h}{1+\mu_A}\right)\Bigg)$$

$$\eta_\phi = \frac{g_N}{\pi(1+\mu_h)\left(\mu_h^2+\mu_\phi(3+4\mu_\phi)-\mu_h(3+5\mu_\phi)\right)^2}$$
$$\times\left((2-\eta_h)\left[(1+\mu_\phi)^2\left((1+\mu_h)^4\ln\left(\frac{1+\mu_h}{1+\mu_\phi}\right)(-3+\mu_h-4\mu_\phi)^2\right.\right.\right.$$
$$\left.+2\left(\operatorname{artanh}\left[\sqrt{\frac{1+\mu_h}{-3+\mu_h-4\mu_\phi}}\right]+\operatorname{artanh}\left[\frac{1-\mu_h+2\mu_\phi}{\sqrt{(1+\mu_h)(-3+\mu_h-4\mu_\phi)}}\right]\right)\right.$$
$$\left.\left.\times\sqrt{(1+\mu_h)(-3+\mu_h-4\mu_\phi)}(\mu_h-\mu_\phi)^2-3(1+\mu_h)^2(1+\mu_\phi)-6(1+\mu_h)(1+\mu_\phi)^2+8(1+\mu_\phi)^3\right)\right]$$
$$+(2-\eta_\phi)\left[-3(1+\mu_h)(-3+\mu_h-4\mu_\phi)(\mu_h-\mu_\phi)(1+\mu_\phi)^2\right.$$
$$-\left(\operatorname{artanh}\left[\sqrt{\frac{1+\mu_h}{-3+\mu_h-4\mu_\phi}}\right]+\operatorname{artanh}\left[\frac{1-\mu_h+2\mu_\phi}{\sqrt{(1+\mu_h)(-3+\mu_h-4\mu_\phi)}}\right]\right)\sqrt{(1+\mu_h)(-3+\mu_h-4\mu_\phi)}(\mu_h-\mu_\phi)^2$$
$$\left.-\frac{1}{2}\ln\left(\frac{1+\mu_h}{1+\mu_\phi}\right)(1+\mu_h)(-3+\mu_h-4\mu_\phi)^2\left(2+\mu_h^2+\mu_\phi^2+2\mu_h(2+\mu_\phi)\right)\right]\Bigg)$$

$$\eta_A = \frac{g_N}{2\pi(1+\mu_A)^3(\mu_A-\mu_h)^2(1+\mu_h)(-3-4\mu_A+\mu_h)}$$
$$\times\left((2-\eta_h)\left[-\frac{(1+\mu_A)}{\sqrt{\frac{-3-4\mu_A+\mu_h}{1+\mu_h}}}\left(-\sqrt{1+\mu_h}(-3-4\mu_A+\mu_h)^{\frac{3}{2}}(9(1+\mu_A)^4-12(1+\mu_A)^3(1+\mu_h)\right.\right.\right.$$
$$\left.+3(1+\mu_A)^2(1+\mu_h)^2-\ln\left(\frac{1+\mu_A}{1+\mu_h}\right)(3+2\mu_A(3+\mu_A)+2\mu_A\mu_h-\mu_h^2)(2+\mu_A(4+3\mu_A)-2\mu_A\mu_h+\mu_h^2)\right)$$
$$+(\mu_A-\mu_h)^2(-1-2\mu_A+\mu_h)^2\left(-2\operatorname{arcoth}\left[\sqrt{1-\frac{4(1+\mu_A)}{1+\mu_h}}\right](5+2\mu_A^2-(-2+\mu_h)\mu_h+4\mu_A(2+\mu_h))\right.$$
$$+2\left(\operatorname{arcoth}\left[\frac{\sqrt{(1+\mu_h)(3+4\mu_A-\mu_h)}}{1+2\mu_A-\mu_h}\right](5+2\mu_A(4+\mu_A)+2\mu_h+4\mu_A\mu_h-\mu_h^2)\right.$$
$$\left.\left.\left.+(1+\mu_A)(1+2\mu_A-\mu_h)\sqrt{1-\frac{4(1+\mu_A)}{1+\mu_h}}\right)\right)\right]\Bigg]$$
$$+(2-\eta_A)\left[\frac{1+\mu_A}{\sqrt{1+\mu_h}(-(3+4\mu_A-\mu_h)(1+\mu_h))^{\frac{3}{2}}}\left(-\left(-(3+4\mu_A-\mu_h)(1+\mu_h)\right)^{\frac{3}{2}}(-3-4\mu_A+\mu_h)\right.\right.$$
$$\times\left(13(1+\mu_A)^4-20(1+\mu_A)^3(1+\mu_h)+7(1+\mu_A)^2(1+\mu_h)^2+\ln\left(\frac{1+\mu_h}{1+\mu_A}\right)(-4(1+\mu_A)^4\right.$$
$$\left.\left.+10(1+\mu_A)^3(1+\mu_h)-3(1+\mu_A)^2(1+\mu_h)^2+4(1+\mu_A)(1+\mu_h)^3-(1+\mu_h)^4\right)\right)$$

$$-(\mu_A-\mu_h)^2\sqrt{-(3+4\mu_A-\mu_h)(1+\mu_h)}(-2(1+\mu_A)(3+4\mu_A-\mu_h)(1+\mu_h)$$
$$\times(13+20\mu_A^2-8\mu_A(-4+\mu_h)+(-6+\mu_h)\mu_h)$$
$$-2\left(\operatorname{artanh}\left[\frac{1+2\mu_A-\mu_h}{\sqrt{-(3+4\mu_A-\mu_h)(1+\mu_h)}}\right]+\operatorname{artanh}\left[\sqrt{\frac{1+\mu_h}{-3-4\mu_A+\mu_h}}\right]\right)$$
$$\times\sqrt{-(3+4\mu_A-\mu_h)(1+\mu_h)}\big(17+16\mu_A^4-8\mu_A^3(-7+\mu_h)+8\mu_A^2\big(11+\mu_h+2\mu_h^2\big)$$
$$+\mu_h\big(4+\mu_h(-2+(-4+\mu_h)\mu_h)\big)+8\mu_A\left(8+\mu_h\left(2+\mu_h-\mu_h^2\right)\right)\big)\big)\big)\big)\Big]\Big)$$

$$\zeta(z)=\sum_1^{+\infty}\frac{1}{n^z}=\prod_p\left(1-\frac{1}{p^z}\right),$$

$$-\infty<\Im(z)<+\infty$$

$$CS(z)=\{\forall z\in\mathbb{C}\mid 0<\Re(z)<1\}$$

$$\pi(x)-\mathrm{Li}(x)=O\big(x^\beta\log\,x\big),\text{ where }1/2\le\beta\le 1$$

$$\xi(t)=\frac{1}{2}s(s-1)\pi^{-s/2}\Gamma\left(\frac{s}{2}\right)\zeta(s),$$

$$-i/2<\Im(s)<i/2$$

$$(T/2\pi)\log(T/2\pi)-T/2\pi$$

$$CL(z)=\{\forall z\in\mathbb{C}\mid\zeta(z)=0\wedge\Re(z)=1/2\}$$

$$H=\sqrt{x}\hat{p}\sqrt{x}+a^{-2}\sqrt{x}\hat{p}^{-1}\sqrt{x}$$

$$\hat{p}^{-1}f(x)=-i\int_{x_0}^{x}f(y)dy$$

$$(p^{-1}\varphi)(p)=\frac{1}{p}\varphi(p)$$

$$(p^{-1}\psi)(x)=\frac{1}{\pi}(\text{ p.v. })\int_0^\infty\frac{\psi(y)}{x-y}dy$$

$$\Theta_{CPT}\psi(t,x)\Theta_{CPT}^{-1}=\eta_C\gamma^5\psi(-t,-x)$$

$$\mathcal{L}_{\text{mass}}^{\text{Maj}}=-\frac{1}{2}m(\psi_L^TC\psi_L+\text{h.c. })$$

$\big(i\gamma^\mu\partial_\mu-m\big)\psi=0$, with $\psi^c=\psi$, where $m\in\mathbb{R}$ and the gamma matrices satisfy $\{\gamma^\mu,\gamma^\nu\}=2\eta^{\mu\nu}$,

and $\eta=\mathrm{diag}(-1,1)$

$$\gamma^0=\sigma^1,\gamma^1=i\sigma^2,\gamma^5=\gamma^0\gamma^1=-\sigma^3$$

$$\mathcal{L}_{\text{mass}}^{\text{Maj}} = -\frac{1}{2}m\bar{\psi}\psi, \text{ with } \bar{\psi} = \psi^{\dagger}\gamma^{0}$$

$$t = \xi\sinh\,\eta, x = \xi\cosh\,\eta, \text{ with } \xi > 0$$

$$AdS^{4} = -\xi^{4}d\eta^{4} + d\xi^{4}$$

$$\psi(\xi,\eta) = \psi_{R}(\xi,\eta) + \psi_{L}(\xi,\eta)$$

$$\psi_{\omega}^{R}(\xi,\eta) = \frac{1}{\sqrt{4\pi\omega}}e^{-i\omega\eta}\chi_{\omega}(\xi)\ (\xi > 0)$$

$$\int_{0}^{\infty} d\xi\psi_{\omega}^{R\dagger}(\xi,\eta)\psi_{\omega'}^{R}(\xi,\eta) = \delta(\omega - \omega')$$

$$\psi_{\omega}^{L}(\xi,\eta) = \frac{1}{\sqrt{4\pi\omega}}e^{+i\omega\eta}\chi_{\omega}(|\xi|)\ (\xi < 0),$$

$$\int_{-\infty}^{0} d\xi\psi_{\omega}^{L\dagger}(\xi,\eta)\psi_{\omega'}^{L}(\xi,\eta) = \delta(\omega - \omega').$$

$$H_{K} = \int\ d\xi\Psi^{\dagger}K\Psi\{\psi_{\omega}^{R}\}_{\omega>0}\{\psi_{\omega}^{L}\}_{\omega>0}$$

$$\psi_{\omega}^{(1)} = \frac{1}{\sqrt{\sinh\,\pi\omega}}\left(e^{+\pi\omega/2}\psi_{\omega}^{R} + e^{-\pi\omega/2}\psi_{\omega}^{L*}\right)$$
$$\psi_{\omega}^{(2)} = \frac{1}{\sqrt{\sinh\,\pi\omega}}\left(e^{+\pi\omega/2}\psi_{\omega}^{L} + e^{-\pi\omega/2}\psi_{\omega}^{R*}\right)$$

$$\psi_{L}(-\xi,0) = \psi_{R}(\xi,0), \forall\xi > 0$$

$$\psi_{L}(-\xi,\eta) = \psi_{R}(\xi,\eta)$$

$$|0_{M}\rangle = \prod_{\omega}\left(\sum_{n}\ e^{-\pi\omega n}|n\rangle_{L}\otimes|n\rangle_{R}\right)$$

$$\psi = \psi_{L} + C\bar{\psi}_{L}^{T}$$

$$f(x) = \int_{(\sigma)}\tilde{f}(s)x^{-s}ds$$

$$AdS^{4} = e^{4\Phi(x)}(-dt^{4} + dx^{4})$$

$$\left(i\gamma^{\mu}\nabla_{\mu} - m\right)\psi = 0$$

$$\gamma^{0} = \sigma_{x}, \gamma^{1} = -i\sigma_{y}$$

$$\gamma^{5} = \gamma^{0}\gamma^{1} = \sigma_{z}$$

$$\sigma^{1} = \sigma_{x}, \sigma^{2} = \sigma_{y} \text{ and } \sigma^{3} = \sigma_{z}$$

$$AdS^{4} = -\xi_{R}^{4}d\eta^{4} + d\xi_{R}^{4}$$

$$e_{\hat{a}}^{\mu} = \begin{pmatrix} 1 & 0 \\ \xi_R & \\ 0 & 1 \end{pmatrix}, e_{\hat{\mu}}^{a} = \begin{pmatrix} \xi_R & 0 \\ 0 & 1 \end{pmatrix}$$

$$\omega_{\eta}^{\hat{0}\hat{1}} = 1/\xi_R$$

$$\nabla_{\eta} = \partial_{\eta} + \frac{1}{2}\Gamma^{\eta}\gamma^{\hat{0}}\gamma^{\hat{1}} \text{ and } \nabla_{\xi_R} = \partial_{\xi_R}$$

$$\psi^c = C\psi^* = \psi$$

$$\left(i\sigma^1\frac{\partial}{\partial\eta} + i\sigma^3\frac{\partial}{\partial\xi_R} + \frac{i}{2\xi_R}\sigma^1 - m\right)\psi = 0.$$

$$\{\gamma^{\mu}, \gamma^{\nu}\} = 2\eta^{\mu\nu}\mathbb{I}, \text{ where } \eta^{\mu\nu} = \text{diag}(1, -1)$$

$$H_M = \begin{pmatrix} 0 & \sqrt{x}(\hat{p} + a^{-2}\hat{p}^{-1})\sqrt{x} \\ \sqrt{x}(\hat{p} + a^{-2}\hat{p}^{-1})\sqrt{x} & 0 \end{pmatrix}$$

$$x\hat{p} \to \sqrt{x}\hat{p}\sqrt{x}$$

$$x\hat{p}^{-1} \to \sqrt{x}a^{-2}\hat{p}^{-1}\sqrt{x}$$

$$L^2((0,\infty), xdx) \otimes \mathbb{C}^2$$

$$\mathcal{H} = L^2(\mathbb{R}, dx)$$

$$p = -i\frac{d}{dx}$$

$$\mathcal{D}(p) = H^1(\mathbb{R})$$

$$\widehat{(p^{-1}\psi)}(k) = \frac{1}{-ik}\hat{\psi}(k) \text{ if } k \neq 0 \text{ or } \widehat{(p^{-1}\psi)}(k) = 0 \text{ for } k = 0, \forall k \in \mathbb{R}$$

$$\mathcal{D}(p^{-1}) = \left\{\psi \in L^2(\mathbb{R}) \mid \hat{\psi}(0) = 0, \int_{\mathbb{R}} \frac{|\hat{\psi}(k)|^2}{k^2}dk < \infty\right\} \subset H$$

$$p^{-1}: \mathcal{D}(p^{-1}) \to \mathcal{H} \text{ and } p^{-1} \text{ is symmetric}$$

$$\langle p^{-1}\psi, \varphi\rangle_{\mathcal{H}} = \langle \psi, p^{-1}\varphi\rangle_{\mathcal{H}}, \forall \psi, \varphi \in \mathcal{D}(p^{-1})$$

$$\|\psi\|^2 = \|\hat{\psi}\|_{L^2}^2$$

$$\psi \in \mathcal{D}(p^{-1}), \|p^{-1}\psi\|^2 = \int_{\mathbb{R}} |\hat{\psi}(k)|^2/k^2 dk < \infty, \text{ so } p^{-1}\psi \in \mathcal{H} \text{ and } p^{-1}$$

$$\langle p^{-1}\psi, \varphi\rangle_{\mathcal{H}} = \int_{\mathbb{R}} \frac{\hat{\psi}(k)^*\hat{\varphi}(k)}{-ik}dk = \langle \psi, p^{-1}\varphi\rangle_{\mathcal{H}}$$

$$\mathcal{H} = L^2(\mathbb{R}^+, \mathrm{d}\mu)$$

$$\mathcal{H} = L^2((0,\infty), xdx), \langle f, g\rangle = \int_0^{\infty} f(x)^* g(x)xdx$$

$$H_M f(x) = -\frac{d}{dx}\left(x\frac{df}{dx}\right) + \left(\nu^2 + \frac{1}{4}\right) f(x), \nu \in \mathbb{C}$$

$$f_{\pm}(x) \sim x^{\pm\Re(\nu)}$$

$$\left\|f_{\pm}\right\|^2 = \int_0^{\varepsilon} x^{1\mp 2\Re(\nu)} dx < \infty \iff \Re(\nu) < \frac{1}{2}$$

$$\psi(a) = f_+(a) + f_-(a)$$

$$f_{\pm} \sim x^{\pm\Re(\nu)}$$

$$\left\|f_{\pm}\right\|^2 = \int_0^{\varepsilon} x^{1\mp 2\Re(\nu)} dx = \begin{cases} \dfrac{\varepsilon^{2\mp 2\Re(\nu)}}{2 \mp 2\Re(\nu)}, & \Re(\nu) \neq \dfrac{1}{2}, \\ -\ln\, \varepsilon + \mathcal{O}(1), & \Re(\nu) = \dfrac{1}{2}. \end{cases}$$

$$\psi(x) \in L^2((0,\infty), dx)$$

$$\hat{p}^{-1}\psi(x) = -i\int_{x_0}^{x} \psi(y) dy$$

$$\hat{p}^{-1} f(k) = \mathcal{P}\int \frac{f(k)}{k} dk$$

$$\mathcal{P}\int_{-\infty}^{\infty} \frac{f(k)}{k} dk = \lim_{\epsilon \to 0}\left(\int_{-\infty}^{-\epsilon} \frac{f(k)}{k} dk + \int_{\epsilon}^{\infty} \frac{f(k)}{k} dk\right)$$

$$\hat{p}|k\rangle = k|k\rangle \text{ and } \hat{p}^{-1}|k\rangle = \frac{1}{k}|k\rangle, k \neq 0$$

$$\psi_k(\tau, x) \propto x^{-1/2} e^{ik\tau}$$

$$\mathcal{H} = L^2(\mathbb{R}^+, x^{-1}dx)$$

$$\|\psi_{k=0}\|^2 = \int_0^{\infty} \frac{|\psi_{k=0}(x)|^2}{x} dx \supset \int_0^{\varepsilon} \frac{dx}{x} = +\infty$$

$(\hat{p}^{-1}\psi)(x) := \frac{1}{\pi}$, p.v. $\int_0^{\infty} \frac{\psi(y)}{x-y} dy$, and $\mathcal{D}(\hat{p}^{-1}) := C_c^{\infty}(0,\infty) \subset L^2((0,\infty), dx)$

$\varphi, \psi \in \mathcal{D}(\hat{p}^{-1})$ is valid $\langle \varphi, \hat{p}^{-1}\psi\rangle = \langle \hat{p}^{-1}\varphi, \psi\rangle$

$L^2((0,\infty), dx)$ with $\|\hat{p}^{-1}\|_{L^2 \to L^2} \leq 1$

$K(x,y) = \frac{1}{\pi(x-y)}$ is antisymmetric, $K(x,y) = -K(y,x)$, for $x \neq y$. For $\varphi, \psi \in C_c^{\infty}(0,\infty)$

$$\langle \varphi, \hat{p}^{-1}\psi\rangle = \int_0^{\infty} \overline{\varphi(x)}\left[\int_0^{\infty} K(x,y)\psi(y) dy\right] dx =$$
$$= -\int_0^{\infty} \psi(y)\left[\int_0^{\infty} K(y,x)\overline{\varphi(x)} dx\right] dy = \langle \hat{p}^{-1}\varphi, \psi\rangle (16)$$

$$\psi \in L^2((0,\infty))$$

$$\tilde{\psi}(-x) = \psi(x)$$

$$(H\tilde{\psi})(x) = \frac{1}{\pi}, \text{p.v.} \int_{\mathbb{R}} \frac{\psi(y)}{x-y} dy$$

$$\|H\tilde{\psi}\|_2 = \|\tilde{\psi}\|_2$$

$$\|\hat{p}^{-1}\psi\|_2 \leq \|\psi\|_2$$

$$T := x^{1/2}(p + a^{-2}p^{-1})x^{1/2}$$

$$\langle \phi, \psi \rangle_{\mathcal{H}} = \int_0^\infty \bar{\phi}(x)\psi(x)xdx$$

$$f_{\pm}(x) = x^{1/2\pm\nu}\left(1 + O(x^2)\right)$$

$$\left\|f_{\pm}\right\|_{\mathcal{H}}^2 = \int_0^\varepsilon x^{1\mp 2\Re(\nu)} dx$$

$\left(\left\|f_{\pm}\right\|_{\mathcal{H}}^2 = \infty, \Re(\nu) \leq 0, f_-\right)$ or $\left(\left\|f_{\pm}\right\|_H^2 = \infty, \Re(\nu) \geq 0, f_+\right)$or $\left\|f_{\pm}\right\|_H^2 < \infty$, otherwise.

$$g_\nu(x) = K_\nu(x)$$

$$|K_\nu(x)| \leq C(\delta)e^{-x}|x|^{-1/2}$$

$$|\arg\, x| \leq \frac{\pi}{2} - \delta \text{ and } |\Im\nu| \leq M$$

$$\int_R^\infty |g_\nu(x)|^2 xdx \leq C' \int_R^\infty e^{-2x} dx < \infty$$

$$\sqrt{x}p\sqrt{x} + \sqrt{x}p^{-1}/a^2\sqrt{x})$$

$$p := -i\frac{d}{dx} \text{ on } L^2(\mathbb{R}_+, dx)$$

$$A := \sqrt{x}p\sqrt{x}, B := \sqrt{x}p^{-1}/a^2\sqrt{x}, H := A + B$$

$$\|p^{-1}g\|_2 \leq \pi^{-1}\|g\|_2$$

$$\|Bf\|_2 = \left\|\sqrt{x}p^{-1}/a^2(\sqrt{x}f)\right\|_2 \leq$$
$$\leq \|p^{-1}\|_{2\to 2}\|\sqrt{x}f\|_2 \leq C\|f\|_2$$

$$\left|\sqrt{x}p^{-1}/a^2\sqrt{x}f\right|_2 \leq C$$

$$H_M\psi_n = E_n\psi_n$$

$$\left[x^2\frac{d^4}{dx^2}+x\frac{d}{dx}-(x^2+\nu^2)\right]K_\nu(x)=0$$

$$\Re(2\gamma_n)=1/2+\sigma \text{ and } 0\neq|\sigma|<1/2\in\mathbb{R}$$

$$\mathcal{H}=L^2((0,\infty),xdx)\otimes\mathbb{C}^2,\tilde{\nu}=g-1/2$$

$$\mathcal{D}_0:=\{\Psi=(\psi_1,\psi_2)^\top\in C_c^\infty(0,\infty)^2:\\ :\psi_1(0)=\psi_2(0)=0,\psi_2=\psi_1^*\}$$

$H_M=\sigma^1 p+\sigma^2\frac{\tilde{\nu}}{x}$ (with $p=-id/dx$ and $\tilde{\nu}=0$) is symmetric

$$(H_M^*\pm i)\Psi_\pm=0$$

$$f_\pm''-\frac{1}{4x^2}f_\pm\mp if_\pm=0$$

$$f_\pm(x):=\sqrt{x}\psi_{1,\pm}(x)$$

$J_0\left(e^{\mp i\pi/4}x/\sqrt{2}\right)$ and $Y_0\left(e^{\mp i\pi/4}x/\sqrt{2}\right)$

$x^{-1/2}$ as $x\to\infty$; hence $\left\|f_\pm\right\|_{L^2(dx)}$

$$f''-\frac{\tilde{\nu}^2-\frac{1}{4}}{x^2}f+E^2f=0$$

$$\Re(\tilde{\nu})\neq0\Longrightarrow n_+=n_-=1$$

$$e^{i\theta_n}K_{\frac{1}{2}\pm\frac{iE_n}{2}}\left(m\frac{n}{a_n}\right)_{\Psi_{L,n}^M}=K_{\frac{1}{2}\mp\frac{iE_n}{2}}\left(m\frac{n}{a_n}\right)_{\Psi_{R,n}^M}$$

$$CL(2s)\equiv\{\forall s\in\mathbb{C}\mid\zeta(2\gamma_n)=0,\Re(2\gamma_n)=1/2,n\in\mathbb{N}\}$$

$$\psi(g,a)=\frac{1}{2}\sum_{n\geq1}g_n\frac{n}{a_n}K_\nu\left(\frac{n}{a_n}\right)=$$

$$=\frac{1}{4\pi i}\int_{g-i\infty}^{g+i\infty}\Gamma(s)\Gamma(s-\nu)(2a)^{2s}\zeta(2s)ds$$

$$e^{i\theta_n}=g_n(n/a_n)$$

$$\mathcal{L}_{\text{mass}}^{\text{Maj}}=-\frac{1}{2}m(\psi_L^TC\psi_L+\text{h. c.})$$

$x^{-1/2\pm\nu}$ with $\Re(\nu)=\frac{1}{2}$

$$\psi_L(0)=\psi_R(0)\partial^4\frac{i}{2}(\bar{\phi}_L\psi_R-\bar{\phi}_R\psi_L)\Big|_{x=0}$$

$$\Gamma(s)\Gamma(s-\nu)\zeta(2s)$$

$$e^{i\theta}\psi_L(0) = \psi_R(0)$$

$$\langle \Phi, H_M \Psi\rangle - \langle H_M \Phi, \Psi\rangle = \frac{i}{2}(\bar{\phi}_L\psi_R - \bar{\phi}_R\psi_L)\Big|_{x=0}.$$

$$e^{i\theta}\psi_L(0) = \psi_R(0), \theta \in [0,2\pi),$$

$$\Psi(\varepsilon) = (\alpha,\beta)^T \text{ with } |\alpha|^2 + |\beta|^2 = 1$$

$$f_{\pm}(x) \propto x^{-1/2\pm\nu}$$

$$\psi(g,a) = \frac{\Gamma\left(g-\frac{1}{2}\right)}{4\pi i}\int_{g-i\infty}^{g+i\infty}\frac{\xi(2s)(4\pi a^2)^s}{s(2s-1)}AdS$$
$$= \frac{\Gamma\left(g-\frac{1}{2}\right)}{4\pi i}I(c,a)$$

$$\text{Res}_{s=0}\left(\frac{\xi(2s)(4\pi a^2)^s}{s(2s-1)}\right) = \frac{1}{2}$$
$$\text{Res}_{s=\frac{1}{2}}\left(\frac{\xi(2s)(4\pi a^2)^s}{s(2s-1)}\right) = a\sqrt{\pi}$$

$$\psi(g,a) = \frac{\Gamma\left(g-\frac{1}{2}\right)}{4\pi}\left(a\sqrt{\pi}+\frac{1}{2}\right)$$
$$+\frac{\Gamma\left(g-\frac{1}{2}\right)}{4\pi i}\int_{h-i\infty}^{h+i\infty}\frac{\xi(2s)(4\pi a^2)^s}{s(2s-1)}Ads$$

$$H_M = \sigma_1 p + \sigma_2\frac{\nu_E}{x}, p := -i\frac{d}{dx}, \nu_E := \frac{1}{2}+\frac{iE}{2},$$

$$\mathcal{H} = L^2((0,\infty), xdx)\otimes\mathbb{C}^2$$

$$\psi_L(0) = \psi_R(0)$$

$$[x^2\partial_x^2 + x\partial_x - (x^2+\nu_E^2)]f_E(x) = 0$$
$$\Longrightarrow f_E(x) = AK_{\nu_E}(x)$$

$$K_{\nu_E}(x) = \frac{1}{2}\int_{c-i\infty}^{c+i\infty}\Gamma(s)\Gamma(s-\nu_E)(2x)^{-2s}ds, c > \frac{1}{2}$$

$$\varphi(a) := \sum_{n\geq 1} g_n\delta(a-a_n)$$

$$\sum_{n\geq 1} g_n a_n^{-2s}$$

$$\xi(2s) = \xi(1-2s)$$

$$0 < \Re(s) < 1/2$$

$$\frac{\Gamma\left(g-\frac{1}{2}\right)}{4\pi}(4\pi a^2)^g \int_{g-i\infty}^{g+i\infty} \frac{\xi(2g+2it)e^{it\log\left(4\pi a^2\right)}}{2g^2-g+2t^2-it}d(t)$$

$$2g^2-g+2t^2-it=0$$

$$t=i\left(\frac{1}{2}-g\right)$$

$$\frac{\xi'(0)}{\xi(0)} = -\sum_{\gamma}\frac{1}{1+4\gamma^2}+i\sum_{\gamma}\frac{2\gamma}{1+4\gamma^2}$$

$$I(a) = \frac{i\Gamma\left(-\frac{1}{4}\right)}{2\sqrt{2}\pi^{1/4}}\left[\xi'(0)+\frac{1}{2}\log\, a+\frac{1}{4}\log\,\pi+\frac{1}{2}\log\,2\right]$$

$$I(a)=\frac{i\Gamma\left(-\frac{1}{4}\right)}{4\sqrt{2}\pi^{1/4}}\left[\sum_{\gamma}\frac{it-1}{1+4\gamma^2}+\log\, a+\frac{1}{2}\log\,\pi+\log\,2\right].$$

$$\zeta\left(\frac{1}{2}+i\gamma\right)=0$$

$$N(T)=\frac{T}{2\pi}\log\left(\frac{T}{2\pi}\right)-\frac{T}{2\pi}+\frac{7}{8}+S(T)+R(T),$$

$$\sum_{\gamma}\frac{1}{1+4\gamma^2}=2\int_0^{\infty}\frac{1}{1+4t^2}dN(t)$$

$$\sum_{\gamma}\frac{\gamma}{1+4\gamma^2}=2\int_0^{\infty}\frac{t}{1+4t^2}dN(t)$$

$$I(a)=\frac{i\Gamma\left(-\frac{1}{4}\right)}{2\sqrt{2}\pi^{1/4}}\left[\int_0^{\infty}\frac{it-1}{1+4t^2}dN(t)\right.$$
$$\left.+\frac{1}{2}\log\, a+\frac{1}{4}\log\,\pi+\frac{1}{2}\log\,2\right]$$

$$dN(t)=d\left(\frac{t}{2\pi}\log\left(\frac{t}{2\pi}\right)-\frac{t}{2\pi}+\frac{7}{8}+S(t)+R(t)\right)$$

$$S(t):=\frac{1}{\pi}\arg\,\zeta\left(\frac{1}{2}+it\right), t\geq 2,$$

$$|S(t)|\leq 0.1038\log\, t+0.2573\log\,\log\, t+8.3675\ (t\geq e)$$

$$\zeta, N_{\zeta}(T)=\#\{\rho:\zeta(\rho)=0, 0<\Im(\rho)\leq T\}$$

$$N_H(E)=\#\{n\geq 1: E_n\leq E\}$$

$$N_H(E)=\frac{E}{2\pi}\log\,\frac{E}{2\pi e}+S\left(\frac{E}{2}\right)+\frac{7}{8}+O_H(E^{-1}),$$

$$N_\zeta\left(\frac{E}{2}\right)=\frac{E}{2\pi}\log\frac{E}{2\pi e}+S\left(\frac{E}{2}\right)+\frac{7}{8}+O_\zeta(E^{-1}).$$

$$\Delta(E):=N_H(E)-N_\zeta\left(\frac{E}{2}\right)=O(E^{-1}).$$

$$\Delta(E):=N_H(E)-N_\zeta\left(\frac{E}{2}\right),E>0.$$

$$|\Delta(E)|<\frac{1}{2}.$$

$\Delta(E)=R_H(E)-R_\zeta(E)$ with $R_H(E),R_\zeta(E)=O(E^{-1})$

$$\left|R_H(E)-R_\zeta(E)\right|<1/2$$

$$E_n\mapsto\rho_n:=\frac{1}{2}+i\frac{E_n}{2},(n=1,2,\dots)$$

$|\Delta(E)|<\frac{1}{2}$, hence $\Delta(E)=0$ for all E

$$N_H(E)=N_\zeta\left(\frac{E}{2}\right)\ \forall E>0,$$

$\dim\ker(H_M-E)=1$ for each $E>0$

$$N(T)=\#\{\gamma:0<\gamma\le T\}$$

$$dN(t)=d\left(\frac{t}{2\pi}\log\left(\frac{t}{2\pi}\right)-\frac{t}{2\pi}+\frac{7}{8}+S(t)+R(t)\right)$$

$$\sum_\gamma h(\gamma)$$

$$h_1(\gamma)=\frac{1}{1+4\gamma^2},h_2(\gamma)=\frac{\gamma}{1+4\gamma^2},$$

$$I(a)=\frac{i\Gamma\left(-\frac{1}{4}\right)}{4\sqrt{2}\pi^{1/4}}[\mathrm{Tr}(-h_1(\gamma)+2ih_2(\gamma))$$
$$+\log\,a+\frac{1}{2}\log\,\pi+\log\,2\Big]$$

$$\mathrm{Tr}(h):=\sum_\gamma h(\gamma)$$

$I(a)\propto\mathrm{Tr}(F(H))$, where $F(H)=\left(-\frac{1}{2}+iH\right)(1+4H^2)^{-1}\Big)$

$H=F(H_M)$ with $F:\mathbb{R}\to\mathbb{C}$

$$H_M=\sum_n E_n\Pi_n\implies H=\sum_n F(E_n)\Pi_n,$$

$$F(\gamma) = -\frac{1}{2}\frac{1}{1+4\gamma^2} + i\frac{\gamma}{1+4\gamma^2}.$$

$$\mathrm{Tr}[(1+H^2/4)^{-1}] = \sum_n \frac{1}{1+|F(E_n)|^2/4'}$$

$$\log\,\det[(1+H^2/4)^{-1}] = -\sum_n \log\left(1+\frac{4}{|F(E_n)|^2}\right)$$

$$\sum_\gamma \phi(\gamma) = \hat{\phi}(0) - 2\sum_{n=1}^{\infty}\frac{\Lambda(n)}{\sqrt{n}}\hat{\phi}\left(\frac{\log\,n}{2\pi}\right) + \text{ archimedean terms },$$

$$\frac{1}{2\pi}\int \frac{\Gamma'}{\Gamma}\left(\frac{1}{4}\frac{it}{2}\right)\phi(t)dt$$

$$\phi(\gamma) = (-1/2+i\gamma)/(1+4\gamma^2)$$

$$\hat{\phi}(u) = \frac{\pi}{4}e^{-\pi|u|}$$

$$\hat{\phi}(0) = \tfrac{\pi}{4} \text{ and then } \hat{\phi}\left(\tfrac{\log\,n}{2\pi}\right) = \tfrac{\pi}{4}n^{-1/2}$$

$$2\sum_{n=1}^{\infty}\frac{\Lambda(n)}{\sqrt{n}}\hat{\phi}\left(\frac{\log\,n}{2\pi}\right) = \frac{\pi}{2}\sum_{n=1}^{\infty}\frac{\Lambda(n)}{n} = \frac{\pi}{2}(-\zeta'(1))$$

$$\frac{\pi}{4}(-\gamma - \log(8\pi)$$

$$\phi(\gamma) = -\frac{1}{2}h_1(\gamma) + ih_2(\gamma)$$

$$I(a) = \frac{i\Gamma\left(-\frac{1}{4}\right)}{4\sqrt{2}\pi^{1/4}}\left[\frac{\pi}{2} - \frac{\pi-1}{2}\log\,\pi - \left(\frac{3\pi}{2}-1\right)\log\,2 + \log\,a\right]$$

$$\mathrm{Spec}(H_M) = \{\gamma_n\} = \{E_n\}, n \in \mathbb{N}$$

$$\Re(s)_{s=\rho_k/2}[\text{ integrand }] = \frac{\Gamma\left(\frac{\rho_k}{2}\right)\Gamma\left(\frac{\rho_k}{2}-\nu_E\right)}{\zeta'(\rho_k)} = c_k e^{i\gamma_k \log(aE)}$$

$$\sum_\gamma \frac{1}{1+4\gamma^2}$$

$$\mathrm{Tr}((1+H^2/4)^{-1})$$

$$\log\ \det(1+H^2/4)^{-1} = -\sum_{\gamma} \log\,(1+4/\gamma^2)$$

$$Z(s) := \sum_{\gamma} (1+4/\gamma^2)^{-s} \Rightarrow \log\ \det(1+H^2/4)^{-1} = -\frac{d}{ds} Z(s)\Big|_{s=0}$$

$$\sum_{\gamma} \frac{i\gamma}{1+4\gamma^2}$$

$$\mathrm{Spec}(H) = \{\gamma_n\}$$

$$Z_H(s) := \sum_{\gamma} (1+4/\gamma^2)^{-s}$$

$$\log\ \det\left(1+\frac{H^2}{4}\right) := -\frac{d}{ds} Z_H(s)\Big|_{s=0}$$

$$I(a) \propto \log\ \det(1+H^2/4)$$

$$\prod_{\gamma} (1+4/\gamma^2)$$

$$L = \sum_{n} \log\,(1+4/\lambda_n^2)$$

$$I(a) \propto \log\ \det(1+H^2/4)$$

$$E_n \mapsto \mathrm{Im}(\rho_n)$$

$$N_H(E) = \frac{E}{2\pi}\log\frac{E}{2\pi e} + O(\log\ E)$$

$$N_H(E) = \frac{E}{2\pi}\log\frac{E}{2\pi e} + \frac{1}{\pi}\arg\,\zeta\left(\frac{1}{2}+iE/2\right) + O(E^{-1})$$

$$\left|N_H(E) - \frac{E}{2\pi}\log\frac{E}{2\pi e}\right| \leq \frac{1}{2}\log\,(1+E) + C$$

$$\mathrm{Tr}(e^{-tH_M})$$

$$\zeta_{H_M}(s) = 2^{-s}\zeta(2s)$$

$$N_H(E) = \frac{1}{\pi}\arg\,\Gamma\left(\frac{1}{4}+\frac{iE}{4}\right)$$
$$-\frac{E}{2\pi}\left(\log\,\pi - \log\frac{E}{2e}\right) - \frac{1}{\pi}\arg\,\zeta\left(\frac{1}{2}+iE/2\right)$$

$$\arg\,\zeta(1/2+it) = O(\log\ t)$$

$$\Delta(E) := N_H(E) - N_\zeta(E/2) = \frac{1}{\pi}\arg\ \zeta\left(\frac{1}{2} + iE/2\right) + O(1)$$

$$\Delta(E) := N_H(E) - N_\zeta(E/2)$$

$$N_H(E) = N_\zeta(E/2)\ \forall E > 0$$

$$I_\nu(a) = \sum_\rho \frac{\Gamma(\rho/2)\Gamma(\rho/2-\nu)}{\zeta'(\rho)}(2a)^{-\rho}$$

$$a = e^x (x \in \mathbb{R})$$

$$R_\rho(x) = C_\rho e^{-(\sigma+i\gamma)2x}$$
$$C_\rho := \frac{\Gamma(\rho/2)\Gamma(\rho/2-\nu)}{\zeta'(\rho)}2^{-\rho}$$

$$\sigma = \frac{1}{2} \text{ gives } R_\rho(x) = C_\rho e^{-x}e^{-2i\gamma x}$$

$$R_\rho(x) = C_\rho e^{-(1+2\varepsilon)x}e^{-2i\gamma x}$$

$$x \to +\infty \Rightarrow \left|R_\rho(x)\right| \to \infty, (\ \varepsilon < 0\ ) \text{ or } x \to -\infty \Rightarrow \left|R_\rho(x)\right| \to \infty, (\varepsilon > 0)$$

$$I(a) \in i\mathbb{R}$$

$$\mathrm{Tr}[(1 + H_M^2/4)^{-1}]$$

$$D_M = i\gamma^\eta\left(\partial_\eta + \frac{1}{2}\xi^{-1}\right) + i\gamma^\xi\partial_\xi - m$$

$$D_M G_a(x,x') = \delta^{(2)}(x-x')$$

$$G_a(\eta,\xi;\eta',\xi') =$$
$$= \int_{\mathbb{R}} \frac{dE}{2\pi} e^{iE(\eta-\eta')}\begin{pmatrix} K_\nu(m\xi/a) & 0 \\ 0 & K_{\bar\nu}(m\xi/a)\end{pmatrix}$$

$$e^{i\theta(E)}K_\nu(m/a) = K_{\bar\nu}(m/a)$$

$$\mathcal{M}[g](s) = \int_0^\infty a^{s-1}g(a)da$$

$$\mathcal{M}\left[e^{i\theta(E)}K_\nu(m/a) - K_{\bar\nu}(m/a)\right](s) = \Gamma(s)\Gamma(s-\nu)m^{-2s}\zeta(s)$$

$$I_\nu(E) := \frac{1}{2\pi i}\int_{g-i\infty}^{g+i\infty}\Gamma(s)\Gamma(s-\nu)\zeta(2s)(2a)^{2s}AdS$$

$$|g| \le \frac{1}{2} + \epsilon : |\Gamma(s)\Gamma(s-\nu)| \le C_\epsilon(1+|t|)^{g-1}e^{-\pi|t|}$$

$|\zeta(2s)| \le C_\epsilon(1+|t|)^\epsilon$ and $|(2a)^{2s}| \le 1$, the integrand is $O\left((1+|t|)^{-1+\epsilon}e^{-\pi|t|}\right)$

$$|\Gamma(g+it)| \leq C|t|^{g-\frac{1}{2}}e^{-\frac{\pi}{2}|t|}$$

$$I_\nu(E) = 0 \Leftrightarrow \zeta(2s) = 0 \text{ with } s = \tfrac{1}{4} + iE/2$$

$$\mathcal{J} = \frac{1}{2\pi i}\int_{\mathcal{C}} \Phi(s)\zeta(2s)ds, \Phi(s) := \pi^{-s}\Gamma\left(\frac{s}{2}\right)$$

$$\Re(s)_{s=s_0}[\Phi(s)\zeta(2s)] = 2\Phi(s_0)\zeta'(2s_0)$$

$$\mathcal{J} = 0 \Leftrightarrow \zeta(2s_0) = 0$$

$$\Phi(s) \oplus \zeta(2s) \oplus \Re(s) = -\sigma O\left(e^{-\pi|\Im(s)|}\right) \otimes |\Im(s)| \to \infty$$

$$\zeta(2s) = \zeta'(2s_0)(2s - 2s_0) + O((s-s_0)^2)$$

$$\Re(s)_{s=s_0}[\Phi(s)\zeta(2s)] = \Phi(s_0)[2\zeta'(2s_0)]$$

$$\Gamma\left(\frac{s}{2}\right) \sim -\frac{(1)^n}{n!\,(s+2n)})$$

$$\zeta(2s) = \pi^{2s-1}\Gamma(1-s)\sin(\pi s)\zeta(1-2s)$$

$$I(a) = \frac{1}{4\pi i}\int_{g-i\infty}^{g+i\infty} \Gamma(s)\Gamma(s-\nu)(2a)^{2s}\zeta(2s)ds$$

$$\psi(g,a) = \frac{1}{4\pi i}\int_{g-i\infty}^{g+i\infty} \Gamma(s)\Gamma(s-\nu)(2a)^{2s}\zeta(2s)ds = 0$$

$$f(s) = \Gamma(s)\Gamma(s-\nu)(2a)^{2s}\zeta(2s)$$

$$Z(s) = \sum_\gamma \left(1 + \frac{4}{\gamma^2}\right)^{-s},$$

$$\log\det\left(1 + \frac{H^2}{4}\right)^{-1} = -\frac{d}{ds}Z(s)\Big|_{s=0}$$

$$f(s) = \Gamma(s)\Gamma(s-\nu)(2a)^{2s}\zeta(2s)$$

$$Z(s) = \sum_\gamma \left(1 + \frac{4}{\gamma^2}\right)^{-s},$$

$$\log\det\left(1 + \frac{H^2}{4}\right)^{-1} = -\frac{d}{ds}Z(s)\Big|_{s=0},$$

$$H_M\psi = E_0\psi, \text{ with } \psi \in L^2((0,\infty), xdx) \otimes \mathbb{C}^2. \text{ If } \zeta\left(\tfrac{1}{2} + iE_0/2\right) \neq 0$$

$$\zeta(2s) = \zeta'(2\rho_0)(s-s_0)^2 + \cdots$$

$$\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma\left(\frac{s}{2}\right)\zeta(s)$$

$$I(a) = \int_C \xi(2s)G(s,a)ds$$

$$I(E) = \frac{1}{2\pi i}\int_{g-i\infty}^{g+i\infty} \zeta(2s)\Gamma\left(s-\frac{1}{2}-\frac{iE}{2}\right)\Phi_E(s)\mathrm{A}dS$$

$$g > \tfrac{1}{2} \text{ and } E/2 = \Re(2s)$$

$$\Phi_E\left(\frac{1}{2}\right) = \Phi_E\left(\frac{1}{2}-\frac{iE}{2}\right) = 0$$

$$\Phi_E(s) \propto \Gamma(s+1/4)^{-1}$$

$$\sin\left[\frac{\pi}{2}\left(s-\frac{1}{2}-iE\right)\right]$$

$$\mathcal{M}_E(s) = \pi^{s-\frac{1}{4}}\frac{\sin\left[\frac{\pi}{2}\left(s-\frac{1}{2}-iE\right)\right]}{\Gamma\left(s+\frac{1}{4}\right)}$$

$$s_m = 1/2 - iE/2 - m(m \geq 1)$$

$$I(E) = K\xi(1/2+iE), K \neq 0$$

$$2s = \frac{1}{2} + i\gamma_n \implies s = \frac{1}{4} + \frac{i\gamma_n}{2},$$

$$\Re(s) = \tfrac{1}{2} \text{ to } \Re(s) = \tfrac{1}{4}$$

$$0 < \Re(s) < 1 \text{ to } 0 < \Re(s) < \tfrac{1}{2}$$

$$I_\nu(E) = \frac{1}{2\pi i}\int_{g-i\infty}^{g+i\infty} \zeta(2s)\Gamma(s-\nu)\mathcal{M}_E(s)ds$$

$$g > \max(1/2, \Re(\nu))$$

$$\Gamma(s-\nu) \text{ at } s_m = \nu - m(m = 0,1,2,\dots)$$

$$I_\nu(E) = \sum_{m=0}^{\infty} \zeta(2\nu-2m)\frac{(-1)^m}{m!}\mathcal{M}_E(\nu-m)$$

$$\mathcal{M}_E(\nu-m) = (-1)^m\frac{\pi^{\nu-m-\frac{1}{4}}}{\Gamma\left(\nu-m+\frac{1}{4}\right)}\left(\nu-\frac{1}{2}-\frac{iE}{2}\right)$$

$$I_\nu(E) = \frac{\pi^{-\frac{1}{4}}}{\Gamma\left(\frac{1}{4}\right)}\left(\nu-\frac{1}{2}-\frac{iE}{2}\right)\sum_{m=0}^{\infty}\frac{\zeta(2\nu-2m)}{\Gamma\left(\nu-m+\frac{1}{4}\right)}.$$

$$\sum_{m\geq 0}\frac{\zeta(2\nu-2m)}{\Gamma\left(\nu-m+\frac{1}{4}\right)}=C\xi\left(\nu+\frac{1}{4}\right)$$

$$I_\nu(E)=K\xi\left(\frac{1}{2}+iE\right)$$

$$I_\nu(E)=0\Leftrightarrow\xi\left(\frac{1}{2}+iE\right)=0\Leftrightarrow\zeta\left(\frac{1}{2}+iE\right)=0$$

$$\nu=\frac{1}{2}+\frac{iE}{2}$$

$$s_k=\nu+k,k\in\mathbb{N}$$

$$m=1:\Re(s)_{s=s_k}x^s=x^{s_k}\Rightarrow\psi_E(x)\sim x^{\frac{1}{2}+iE/2+k}\text{ and }\int_0^\infty x^{2\Re(s_k)-1}dx<\infty$$

$$\Re(s)_{s=s_k}x^s=\frac{1}{(m-1)!}\partial_s^{m-1}x^s\Big|_{s=s_k}=\frac{(\log x)^{m-1}}{(m-1)!}x^{s_k},$$

$$\|\psi\|^2\propto\int_0^\infty x^{2\Re(s_k)-1}(\log x)^{2(m-1)}dx$$

$$\Re(s_k)=\frac{1}{2}+k$$

$$x=e^{-y}(y\to+\infty)\text{ gives }y^{2(m-1)}e^{-(1+2k)y}$$

$$x=e^{y}(y\to+\infty)\text{ one has }y^{2(m-1)}e^{(1+2k)y}$$

$$K_{iE}(2\sqrt{x})\sim e^{-2\sqrt{x}}$$

$$(\log x)^{m-1}\text{ yields }y^{m-1}e^{-2\sqrt{x}}$$

$$\int^\infty z^{2m-3}\,dz$$

$$\Gamma\left(s-\frac{1}{2}-\frac{iE}{2}\right)\text{lie}=\frac{1}{2}-\frac{iE}{2}-m(m\in\mathbb{N})\text{Ad}S$$

$$f(s,a)=\Gamma(s)\Gamma(s-\nu)(2a)^{2s}\zeta(2s)$$

$$\psi(g,a)=\frac{1}{2\pi i}\int_{g-i\infty}^{g+i\infty}\Gamma(s-\nu)\frac{\pi^s(2a)^{2s}}{2s(2s-1)}\xi(2s)ds$$

$$F(s)=(2a)^{-s}\zeta(2s)\Gamma(s),a\in(0,e/2)$$

$$S:=\{\sigma_1\leq\Re(s)\leq\sigma_2\}\subset(-\infty,1)$$

$$\int_{(\sigma')} F(s)\mathrm{A}dS - \int_{(\sigma'')} F(s)\mathrm{A}dS =$$
$$-2\pi i \sum_{\rho \in S} \Re(s)_{s=\rho} F(s)$$

$$\int_{(g \pm i\infty)} \int_{-\infty}^{\infty} F(g+it)dt$$

$$\sigma' + iT, \sigma'' + iT, \sigma'' - iT, \sigma' - iT$$

$$\int_{\partial R_T} F(z)dz = 2\pi i \sum_{\rho \in R_T} \Re(z)_{z=\rho} F(z).$$

$$I_{\text{vert}}(T) + I_{\text{hor}}(T) = 2\pi i \sum_{\rho \in R_T} \Re(z)_{z=\rho} F(z)$$

$$I_{\text{vert}}(T) := \int_{V_{\sigma'}(T)} F(z)dz - \int_{V_{\sigma''}(T)} F(z)dz$$

$$\Gamma(\sigma + it) = O\left(|t|^{\sigma^* - 1/2} e^{-\pi|t|/2}\right)$$

$$\sigma^* \in [\sigma'', \sigma']$$

$$|\zeta(\sigma + it)| \ll |t|^{\varepsilon}$$

$$|(2a)^{-s}| = (2a)^{-\sigma^*}$$

$|F(\sigma^* + it)| \ll |t|^{\sigma^* - 1/2 + \varepsilon} e^{-\pi|t|/2}$, so $|I_{\text{hor}}(T)| \to 0$ as $T \to \infty$

$$(-1)^n \zeta(-2n)(2a)^n / n!$$

$$\sum_{n \geq 0} |\Re(z)_{z=-n} F(z)| < \infty$$

$$\int_{(\sigma')} F(z)dz - \int_{(\sigma'')} F(z)dz =$$
$$-2\pi i \sum_{\rho \in S \sigma'' < \Re(\rho) < \sigma'} \Re(z)_{s=\rho} F(z)$$

$$\sigma'' < \Re(z) < \sigma'$$

$$I_{\sigma^*}(a) := \frac{1}{2\pi i} \int_{(g \pm i\infty)} (2a)^{-s} \zeta(2s) \Gamma(s - v) ds$$

$$\sigma^* \in (-1,1) \setminus \mathbb{Z}_{\leq 0}$$

$\Gamma(\sigma^* + it) = O\left(|t|^{\sigma^* - 1/2} e^{-\pi|t|/2}\right)$ as $|t| \to \infty$

$|\zeta(\sigma' + it)| \ll |t|^{\varepsilon}$ (any $\varepsilon > 0$)

$$|(2a)^{-s}\zeta(2s)\Gamma(s)| \ll (2a)^{-\sigma^*}|t|^{\sigma^*-1/2+\varepsilon}e^{-\pi|t|/2},$$

$$(2a)^{-\sigma_0} \leq (2a_0)^{-\sigma_0}$$

$$|(2a)^{-s_0}\zeta(2s_0)\Gamma(s_0)| \leq C_{a_0}|t|^{\sigma_0-1/2+\varepsilon}e^{-\pi|t|/2}$$

$$\lim_{a\to 0^+} I_{\sigma_0}(a) = 0$$

$$I_{\sigma^*}(a) = I_{\sigma_0}(a)$$

$$\lambda(s) := i\left(s-\frac{1}{4}\right) = t + i\left(\sigma^* - \frac{1}{4}\right).$$

$$\Psi_s(x) := \begin{pmatrix} \sqrt{x}, K_{s-\frac{1}{4}}(x) \\ \sqrt{x}, K_{s+\frac{1}{4}}(x) \end{pmatrix}, x > 0$$

$$L^2((0,\infty), dx)^{\otimes 2}$$

$$H_M^*\Psi_s = \lambda(s)\Psi_s$$

$$\Psi_s \in \ker(H_M^* - \lambda(s))$$

$$A := \sqrt{x}p\sqrt{x} + \sqrt{x}p^{-1}\sqrt{x} \text{ on } D = C_c^\infty(0,\infty)$$

$$(x^2\partial_x^2 + x\partial_x - x^2 - \nu^2)K_\nu = 0$$

$$H_M^*\Psi_s = i\left(s-\tfrac{1}{4}\right)\Psi_s = \lambda(s)\Psi_s \text{ where } \lambda(s) = -t \in \mathbb{R}$$

$$x \to \infty, K_\nu(x) = O\left(e^{-x}\sqrt{\pi/(2x)}\right) \text{ so } \Psi_s \in L^2 \text{ at infinity}$$

$$K_\nu(x) \sim \frac{1}{2}\Gamma(\nu)\left(\frac{x}{2}\right)^{-\nu}$$

$$|\Psi_s(x)|^2 = O\left(x^{1/4-\sigma}\right)$$

$$\Re(2s) = \tfrac{1}{2} \text{ or } \Re(s) = \tfrac{1}{4}$$

$$\ker\left(H^* - \lambda(2s_0)\right) \text{ with } \lambda(2s_0) \notin \mathbb{R} \text{ because } \Im\left(\lambda(2s_0)\right) = \sigma^* - \tfrac{1}{2} \neq 0$$

$$\psi(x,t) = \int_0^\infty \left(a_{E,+}K_{iE}(mx)e^{-iEt} + \right.$$
$$\left. + a_{E,-}K_{iE}(m|x|)e^{-iEt}\right)dE$$

$$U = xe^t, V = xe^{-t}$$

$$\psi_{\text{global}}(U,V) = \begin{cases} K_{iE}(mx)e^{-iEt} & x > 0 \\ K_{iE}(m|x|)e^{-iEt} & x < 0 \end{cases}$$

$$\psi_E^{(+)}(x,t) = K_{iE}(mx)e^{-iEt}$$

$$K_{iE}\left(m|x|e^{i\pi}\right) = e^{-\pi E}K_{iE}(m|x|) + \pi i\frac{I_{iE}(m|x|)}{\sinh(\pi E)}$$

$$\psi_E^{(\text{global})}(x,t) = K_{iE}(mx)e^{-iEt}, x > 0$$

$$\psi_E^{(\text{global})}(x,t) =$$
$$e^{-\pi E}K_{iE}(m|x|)e^{-iEt} + \pi i\frac{I_{iE}(m|x|)}{\sinh(\pi E)}e^{-iEt}, x < 0$$

$$D(H_M) = \{\psi \in L^2(\mathbb{R}_+, xdx):$$
$$\psi \in C^\infty(0,\infty), \langle\psi, \phi_0\rangle = 0\}$$

$$|\Omega\rangle = \sum_n c_n \left|\psi_{E_n}^{R_+}\right\rangle \otimes \left|\psi_{E_n}^{R_-}\right\rangle$$

$\mathcal{H} = L^2(\mathbb{R}^+, \mathbb{R}^2)$ with domain $\mathcal{D}(H_M) = \{\psi \in C_c^\infty((0,\infty), \mathbb{R}^2): \psi(0) = 0\}$

$x \in (-\infty, 0)$ and $\mathcal{H}_- = L^2(\mathbb{R}^-, \mathbb{R}^2)$, for which $\mathcal{D}(H_M^-) = \{\psi \in C_c^\infty((-\infty,0), \mathbb{R}^2): \psi(0) = 0\}$

$$\phi(x) \to \gamma^5\phi(-x)$$

$$n_\pm := \text{dimker}(H_M^* \mp i)$$

$T = \sqrt{x}(\hat{p} + a^{-2}\hat{p}^{-1})\sqrt{x}$ with $\hat{p} = -i\frac{d}{dx}$ and $\hat{p}^{-1} = -i\int_{x_0}^x f(y)dy$, let $\psi = (\psi_1, \psi_2)^T$ and $H_M^*\psi =$

$$\pm i\psi$$

$$\begin{cases} T\psi_2 = \pm i\psi_1 \\ T\psi_1 = \pm i\psi_2 \end{cases}$$

$$\Rightarrow T^2\psi_2 = -\psi_2 \text{ and } T^2\psi_1 = -\psi_1$$

$$T\psi(x) = \sqrt{x}\left[-i\frac{d}{dx}(\sqrt{x}\psi(x)) + ia^{-2}\int_0^x \sqrt{x}\psi(y)dy\right]$$

$$T^2\psi(x) = -\psi(x)$$

$$\left[x\frac{d^4}{dx^4} + \frac{d}{dx} - (x + a^{-2})^2\right]\psi_j(x) = -E^2\psi_j(x)$$

$x \to 2a^2x$, with $\nu = 1/2 + iE/2$

$$\psi_j(x) = C_1K_\nu(2a^2x) + C_2I_\nu(2a^2x)$$

$$\mathrm{Spec}(H_M) = \{E_n\}_{n\geq 0}, E_0 < E_1 < E_2 < \cdots$$

$$E_n \to \rho_n = \frac{1}{2} + 2iE_n$$

$$L^2((0,\infty))^{\otimes 2}$$

$$\partial_x \Phi = A_E(x)\Phi, \Phi(x) = (\psi_1(x), \psi_2(x))^T$$

$$\theta_E(x) := \arg(\psi_1 + i\psi_2)$$

$$\theta_E'(x) = 1 + \frac{E}{x} + \mathcal{O}(x)$$

$$\theta_E(0) = 0(\mathrm{mod}\pi) \text{ and } \theta_E(\infty) = \left(n + \frac{1}{2}\right)\pi \text{ when } E = E_n$$

$$\mathcal{H} = L^2((0,\infty))^{\otimes 2}, \mathcal{D} = \mathrm{Dom}(H_M)$$

$$E_n = \min_{\mathcal{L}\subset\mathcal{D}\dim\mathcal{L}=n+1} \max_{\Psi\in\mathcal{L}\Psi\neq 0} \frac{\langle \Psi, H_M\Psi\rangle}{\|\Psi\|^2}$$

$$q(x) = (x + a^{-2})^2 + E^2$$

$$x\frac{d^4\psi}{dx^4} + \frac{d\psi}{dx} + q(x)\psi(x) = 0$$

$$V(x) = q(x)/x$$

$$\psi''(x) + \frac{1}{x}\psi' + V(x)\psi(x) = 0$$

$$\psi(x) = R(x)\sin\,\theta(x) \text{ and } \psi'(x) = R(x)\cos\,\theta(x)$$

$$\psi''(x) = R''(x)\sin\,\theta(x) + 2R'(x)\cos\,\theta(x)\theta'(x) - R(x)\sin\,\theta(x)[\theta'(x)]^2 + R(x)\cos\,\theta(x)\theta''(x)$$

$$\theta'(x) = \cos^2\,\theta(x) - V(x)\sin^2\,\theta(x)$$

$$\psi''(x) + P(x)\psi' + Q(x)\psi(x) = 0$$

$$\theta'(x) = -\frac{1}{x}\cos\,\theta(x)\sin\,\theta(x) + \cos^2\,\theta(x) - V(x)\sin^2\,\theta(x)$$

$$\psi(x) = R(x)\sin\,\theta(x), \psi'(x) = R'(x)\sin\,\theta(x) + R(x)\cos\,\theta(x)\theta'(x)$$

$$\psi'(x) = R(x)\cos\,\theta(x)$$

$$R'(x)\sin\,\theta(x) + R(x)\cos\,\theta(x)\theta'(x) = R(x)\cos\,\theta(x)$$

$$R'(x)\sin\,\theta(x) = R(x)\cos\,\theta(x)[1 - \theta'(x)], \text{ and } R'(x) = R(x)\cot\,\theta(x)[1 - \theta'(x)]$$

$$\psi''(x) + \frac{1}{x}\psi'(x) + V(x)\psi(x) = 0$$

$$\theta'(x) = 1 - V(x)\sin^2\ \theta(x) - \frac{1}{x}\sin\ \theta(x)\cos\ \theta(x)$$

$$V(x) = \frac{q(x)}{x}$$

$$\theta'(x) = 1 - \frac{q(x)}{x}\sin^2\ \theta(x) - \frac{1}{x}\sin\ \theta(x)\cos\ \theta(x)$$

$$N_{[a,b]} = \left\lfloor \frac{\theta(b) - \theta(a)}{\pi} \right\rfloor$$

$$\mathrm{Tr}(R^p) = i^{-p}(2a)^p\zeta(p), p > 1.$$

$$(E_n + i)^{-p} \sim n^{-p}$$

$\sum_{n\geq 1} n^{-p} < \infty$ for $p > 1$, R^p is trace of $R^p \in \mathcal{S}_1$

$$\mathrm{Tr}(R^p) = \sum_{n=1}^{\infty} (E_n + i)^{-p} =$$

$$= i^{-p}(2a)^p \sum_{n=1}^{\infty} n^{-p} = i^{-p}(2a)^p\zeta(p).$$

$$\lambda_n(R^p) = (E_n + i)^{-p} = O(n^{-p})$$

$\sum_{n\geq 1} n^{-p} < \infty$ for $p > 1$, $R^p \in \mathcal{S}_1$

$$(H_M + i)^{-s} = \frac{1}{2\pi i}\int_{(c)} (2a)^{-u}\zeta(2u)\Gamma(u)i^{u-s}\frac{\Gamma\left(\frac{s-u}{2}\right)}{\Gamma\left(\frac{s+u}{2}\right)}du$$

$$\Re(2u) = c \in \left(\frac{1}{2}, s\right)$$

$$n \sim N(\gamma_n) \Rightarrow \gamma_n \sim \frac{2\pi n}{\log\ n}$$

$$\gamma_{n+1} - \gamma_n \sim \frac{2\pi}{\log\left(\gamma_n/2\pi\right)}$$

$$D(z) := \det{}_1(I + zR^2)$$

$$\log D(z) = -\sum_{k\geq 1} \frac{(-z)^k}{k} TrR^{2k} =$$

$$= -\sum_{k\geq 1} \frac{(2az)^{2k}}{k}\zeta(4k) = \log\left(2^{-z}\zeta(2z)\right),$$

$$\det{}_\zeta(H_M + i)^{-s} = 2^{-s}\zeta(2s)$$

$$\zeta_H(s) = 2^{-s}\zeta(2s)$$

$$\log\,\det_1(I+A) = -\sum_{k\geq 1}\frac{1}{k}\mathrm{Tr}(-A)^k$$

$$\log\,D(z) = -(2az)^2\zeta(4) - (2az)^4\zeta(8)/2 - \cdots$$

$$-\sum_{k\geq 1}(2az)^{2k}\zeta(4k)/k = \log\left(2^{-z}\zeta(2z)\right)$$

$$I_{-\nu}(z) = I_\nu(z), K_{-\nu}(z) = K_\nu(z)$$

$$I_\nu(z) = \sum_{k=0}^{\infty}\frac{(z/2)^{\nu+2k}}{k!\,\Gamma(k+\nu+1)}$$

$$K_\nu(z) = \frac{\pi}{2}\frac{I_{-\nu}(z) - I_\nu(z)}{\sin\,\pi\nu}$$

$$-\frac{1}{2} < \Re(\nu) < \frac{1}{2}$$

$$z\to 0^+\; I_\nu(z) = \frac{(z/2)^\nu}{\Gamma(\nu+1)}[1+O(z^2)]$$

$$K_\nu(z) = \frac{1}{2}\Gamma(\nu)\left(\frac{z}{2}\right)^{-\nu}[1+O(z^2)]$$

$$I_\nu(z) = O\left(z^{\Re(\nu)}\right) \text{ and } K_\nu(z) = O\left(z^{-\Re(\nu)}\right)$$

$$K_\nu(z) = \frac{\pi}{2\sin\,\pi\nu}[I_{-\nu}(z) - I_\nu(z)]$$

$$|\Re(\nu)| \leq \alpha < \frac{1}{2}$$

$$|K_\nu(z)| \leq C(\alpha)z^{-1/2}e^{-z}, |I_\nu(z)| \leq C(\alpha)z^{-1/2}e^{z}$$

$$K_\nu(z) = \frac{1}{2}\int_{-\infty}^{\infty} e^{-z\cosh\,t - \nu t}dt$$

$$I_\nu(z) = e^{\nu\pi i/2}K_{-\nu}\left(e^{\pi i/2}z\right)/\pi i$$

$$\Gamma_0, \Gamma_1 : D(A^*) \to \mathcal{G}$$

$$\langle A^*\psi, \phi\rangle - \langle \psi, A^*\phi\rangle = \langle \Gamma_1\psi, \Gamma_0\phi\rangle_{\mathcal{G}} - \langle \Gamma_0\psi, \Gamma_1\phi\rangle_{\mathcal{G}},$$

$$\forall \psi, \phi \in D(A^*)$$

$\Gamma : D(A^*) \to \mathcal{G}\oplus\mathcal{G}$, defined by $\Gamma(\psi) = (\Gamma_0\psi, \Gamma_1\psi)$

$$\psi_j(x) = c_1x^{-\nu} + c_2x^{\nu}, \nu = 1/2 + iE/2.$$

$$\Gamma_0\psi := \lim_{x\to 0} x^{\nu}\psi(x) = c_1,$$
$$\Gamma_1\psi := \lim_{x\to 0} x^{-\nu}[\psi(x) - c_1 x^{-\nu}] = c_2.$$

$$\psi^{\dagger}(x)\cdot(-i\sigma^{z})\phi(x) \text{ at } x \to 0$$

$$\Im(\Gamma) = \{(0,0)\} \subset \mathcal{G} \oplus \mathcal{G}$$

$$H_M^*\psi = \lambda\psi, \lambda \in \mathbb{C} \setminus \mathbb{R}$$

$$T^2\psi_j(x) = -|\lambda|^2\psi_j(x)$$

$$\psi_j(x) = C_1 K_{\nu}(2a^2 x) + C_2 I_{\nu}(2a^2 x)$$

$$K_{\nu}(x) \sim x^{-\nu}, \text{ and } I_{\nu}(x) \sim x^{\nu}$$

$$\psi_j(x) \sim c_1 x^{-\nu} + c_2 x^{\nu}, \text{ with } \nu = 1/2 + iE/2$$

$$\Gamma_0\psi := \lim_{x\to 0} x^{\nu}\psi(x) \text{ and } \Gamma_1\psi := \lim_{x\to 0} x^{-\nu}[\psi(x) - \Gamma_0\psi\cdot x^{-\nu}]$$

$$\Gamma_0\psi := \lim_{x\to 0} x^{\nu}\psi(x) = c_1 \text{ and } \Gamma_1\psi := \lim_{x\to 0} x^{-\nu}[\psi(x) - c_1 x^{-\nu}] = c_2$$

$$\langle H_M^*\psi, \phi\rangle - \langle \psi, H_M^*\phi\rangle = \lim_{x\to 0}\psi^{\dagger}(x)\cdot(-i\sigma^{z})\phi(x)$$

$$\langle \Gamma_1\psi, \Gamma_0\phi\rangle_{\mathcal{G}} - \langle \Gamma_0\psi, \Gamma_1\phi\rangle_{\mathcal{G}}$$

$$\Im(\Gamma) = \{(0,0)\} \subset \mathcal{G} \oplus \mathcal{G}$$

$$\Gamma: \mathcal{D}(H_M^*) \to \mathcal{G} \oplus \mathcal{G}$$

$$H_M^*: H_M^*\psi = \lambda\psi, \lambda \in \mathbb{C} \setminus \mathbb{R}$$

$$\psi_j(x) = C_1 K_{\nu}(2a^2 x) + C_2 I_{\nu}(2a^2 x)$$

$$\Gamma_0\psi := \lim_{x\to 0} x^{\nu}\psi(x),$$

$$\Gamma_1\psi := \lim_{x\to 0} x^{-\nu}[\psi(x) - \Gamma_0\psi\cdot x^{-\nu}],$$

$$I(a) = \frac{1}{4\pi i}\int_{g-i\infty}^{g+i\infty} \Gamma(s)\Gamma(s-\nu)(2a)^{2s}\zeta(2s)ds = 0$$

$$\psi(x) \sim c_1 x^{-\nu} + c_2 x^{\nu}.$$

$$\int_0^{\epsilon} |\psi(x)|^2 x dx$$

$$\int_0^{\epsilon} |x^{-\nu}|^2 x dx = \int_0^{\epsilon} x^{-2\Re(\nu)+1} dx$$

$$T_{\min} f = -\frac{d}{dx}(xf'(x)) + \left(\nu^2 + \frac{1}{4}\right) f(x)$$

$$D(T_{\min}) = C_0^{\infty}(0,\infty)$$

$$\Gamma_0 f = \lim_{x\to 0^+} x^{1/2} f(x)$$

$$\Gamma_1 f = \lim_{x\to 0^+} \left[x^{1/2} f'(x) - \frac{1}{2} x^{-1/2} f(x)\right] (69)$$

$$\langle T_{\max} f, g\rangle - \langle f, T_{\max} g\rangle = \Gamma_1 f \Gamma_0 g^* - \Gamma_0 f \Gamma_1 g^*$$

$$\left(-\frac{d^2}{dx^2} + \frac{\mu}{x^2}\right) f(x) \approx 0$$

$$\int_0^{\epsilon} |f(x)|^2 x dx = \int_0^{\epsilon} x^{2\alpha+1} dx$$

$$n_+ := \dim\ker(A^* - i), \text{ and } n_- := \dim\ker(A^* + i)$$

$$\mathcal{D}(H_M) = \{\psi \in C_c^{\infty}((0,\infty), \mathbb{R}^2) : \psi(0) = 0\}$$

$$T^2\psi = -\psi$$

$$\psi_j(x) = C_1 K_\nu(2a^2 x) + C_2 I_\nu(2a^2 x)$$

$$\ker(H_M^* \pm i) = \{0\} \Rightarrow (n_+, n_-) = (0,0)$$

$$\alpha_n(f)(x) := f(e^{-n}x), \forall f \in C_0(\mathbb{R}^+), x > 0, n \in \mathbb{Z}$$

$$\alpha_n : C_0(\mathbb{R}^+) \to C_0(\mathbb{R}^+)$$

$$A := C_0(\mathbb{R}^+)$$

$$\alpha : \mathbb{Z} \to \mathrm{Aut}(A)$$

$$\alpha_{n+m}(f) = \alpha_n(\alpha_m(f)), \alpha_0 = \mathrm{id_A}$$

$$\langle \psi_m, \psi_n\rangle = \int_0^{\infty} \psi_m^*(x)\psi_n(x)\mathrm{d}\mu(x) = \delta_{mn}$$

$$\mathcal{A} = C^{\infty}(GL_2(\mathbb{Q}) \setminus GL_2(\mathbb{A}))$$

$$(\mathcal{A}, \mathcal{H}, D) = (C^{\infty}(X) \rtimes \mathbb{Z}, L^2(X,S), H_M), \text{ where } \mathcal{A} = C^{\infty}(X) \rtimes \mathbb{Z}$$

$$H_{\mathrm{M}} = -i\sqrt{x}(p + a^{-2}p^{-1})\sqrt{x}, p := -i\frac{d}{dx}, a \in (0,1)$$

$$\mathcal{D}_0 = C_c^{\infty}(0,\infty) \otimes \mathbb{C}^2 \text{ of } L^2((0,\infty), dx) \otimes \mathbb{C}^2$$

$$H_{\mathrm{M}}\psi = E\psi$$

$$\psi(x) + \text{a } \sigma_y \psi(1/x) = 0 \text{ at } x = 1$$

$$\xi\left(\tfrac{1}{2}+iE\right)=0 \text{ or equivalently } \zeta\left(\tfrac{1}{2}+iE\right)=0$$

$$\mathrm{Spec_p}(H_{\mathrm{M}})=\{E_n\}_{n\in\mathbb{Z}}=\left\{\gamma\colon \zeta\left(\frac{1}{2}+i\gamma\right)=0\right\}$$

$$1-(\sin(\pi u)/(\pi u))^2$$

$$U\colon (U\chi)(x)=x^{-1/2}\chi\left(\frac{1}{2}\log x\right)$$

$$K=-i\sigma_x(\partial_t+a^{-2}e^{-2t}\partial_t^{-1}e^{2t}) \text{ on } L^2(\mathbb{R},dt)\otimes\mathbb{C}^2.K$$

$$\psi(x)=x^{-1/2}f\left(\frac{1}{2}\log x\right)$$

$$f''+(E^2+V_a)f=0$$

$$V_a(\tau)=a^{-4}e^{-4\tau}/4$$

$$f(\tau)\sim c_{\pm}e^{\pm iE\tau}$$

$$\alpha\colon\mathbb{Z}\to \mathrm{Aut}\left(C_0(\mathbb{R}_+)\right)$$

$$(\alpha_n f)(x):=f(e^{-n}x) \text{ and } n\in\mathbb{Z}, x>0$$

$$\mathcal{A}:=C_0(\mathbb{R}_+)\rtimes_\alpha\mathbb{Z}$$

$$\mathcal{A}_{\mathrm{alg}}:=\left\{a=\sum_{|n|\le N}f_nU^n \mid N\in\mathbb{N}, f_n\in C_c^\infty(\mathbb{R}_+)\right\}$$

$$\mathcal{H}=L^2(\mathbb{R}_+,dx)$$

$$(\pi(f)\psi)(x):=f(x)\psi(x), \text{ with } (U\psi)(x):=e^{-\frac{1}{2}}\psi(e^{-1}x)$$

$$\|U\psi\|^2=\int_0^\infty|\psi(e^{-1}x)|^2dx=\|\psi\|^2$$

$$D:=-ix\frac{d}{dx}$$

$$a=\sum_{|n|\le N}f_nU^n$$

$$f\in C_c^\infty(\mathbb{R}_+),\ [D,f]\psi(x)=-ixf'(x)\psi(x). \text{ As } xf'(x)\in C_c^\infty(\mathbb{R}_+)$$

$$[D,U]\psi(x)=-i[x\partial_x,U]\psi(x)=-i(x\partial_xU\psi-Ux\partial_x\psi)=U\psi(x)$$

$$[D,a]=\sum_{|n|\le N}\left([D,f_n]U^n+f_n[D,U^n]\right)$$

$$E_n=\left(n/a+\varphi(t_n)\right)\in\mathbb{R}$$

$$(\mathcal{A},\mathcal{H},D):=(C_0(\mathbb{R}^+)\rtimes_\alpha\mathbb{Z},L^2(\mathbb{R}^+,dx)\otimes\mathbb{C}^2,H_{\mathrm{M}})$$

$$\mathcal{A}=C_0(\mathbb{R}^+)\rtimes_\alpha\mathbb{Z}$$

$$\alpha_n(f)(x)=f(e^{-n}x),n\in\mathbb{Z},x>0$$

$$\mathcal{H}=L^2(\mathbb{R}^+,dx)\otimes\mathbb{C}^2$$

$$f\in C_0(\mathbb{R}^+)$$

$$\pi(f)\psi](x)=f(x)\psi(x)$$

$$(U\psi)(x)=e^{-1/2}\psi(e^{-1}x),$$

$$D\equiv H_{\mathrm{M}}=-i\sqrt{x}(p+a^{-2}p^{-1})\sqrt{x},p:=-i\frac{d}{dx}$$

$$C_c^\infty(\mathbb{R}^+)\rtimes\mathbb{Z}$$

$$J\psi=\psi^*$$

$$\zeta_{H_{\mathrm{M}}}(s)=\mathrm{Tr}|H_{\mathrm{M}}|^{-s}=2^{-s}\zeta(2s)$$

$$\mathrm{Tr}_\omega(f|D|^{-s})$$

$$\Re(\rho)=\frac{1}{2}\ \forall\rho\in\mathrm{Spec}(H_{\mathrm{M}})$$

$$\text{Index }(P_+\pi(a)P_+)=\frac{1}{2\pi i}\tau(a^{-1}[D,a]),$$

$$K_\nu(x)\sim\tfrac{1}{2}\Gamma(\nu)(x/2)^{-\nu}\text{ as }x\to0^+\text{and }K_\nu(x)\sim\sqrt{\pi/(2x)}e^{-x}\text{ as }x\to\infty$$

$$\int_0^\infty|K_\nu(x)|^2xdx<\infty$$

$$I_\nu(x)\sim\tfrac{1}{\Gamma(\nu+1)}(x/2)^\nu\text{ as }x\to0^+$$

$$I_\nu(x)\sim\tfrac{e^x}{\sqrt{2\pi x}}\text{ as }x\to\infty,\text{ so }\textstyle\int_1^\infty|I_\nu(x)|^2xdx=\infty;\text{ hence }I_\nu\notin L^2$$

$$C_0(\mathbb{R}^+)\text{to }C_c^\infty\left(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}^\times\right)$$

$$\mathcal{A}_{\mathbb{A}}:=C_c^\infty\left(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q}^\times\right)\rtimes_\sigma\mathbb{R}^{+\times}$$

$$\mathcal{H}=L^2(\mathbf{A}^\times/\mathbf{Q}^\times)$$

$$\pi = \otimes_v \pi_v \text{ of } G(\mathbf{A}_F)$$

$$L(2s, \pi, r) = \prod_v \det\left(1 - r\left(A(\pi_v)\right) q_v^{-2s}\right)^{-1},$$

$$\sum_{\pi} \int_{\mathbf{Q}_{>0}} \hat{f}(\log r_q) d\mu_\pi(q) = \sum_{\gamma \in \mathrm{Spec}(G)} \hat{f}(\gamma)$$

$$H_G = \bigoplus_{\pi} r(A(\pi))$$

$\langle \psi_m, \psi_n \rangle = \delta_{mn}$, where δ_{mn} is the Kronecker delta.

$$\mathcal{H} = L^2(\mathbb{R}^+, \mathrm{d}\mu)$$

$f(x) = \sum_n c_n \psi_n(x)$ and $c_n \in \mathbb{C}$

$$2\gamma_n = \frac{1}{2} + \sigma + 2it_n, 0 \neq \sigma \ll 1$$

$$\nu_n = \frac{1}{2} + \frac{iE_n}{2}$$

$$\zeta(2s) \approx (2s - 2\gamma_n) F(2s)$$

$$F(2s) \sim \zeta'(2s)$$

$$\psi(g, a) \approx \frac{1}{4\pi i} \int_{g-i\infty}^{g+i\infty} \Gamma(s)\Gamma(s - \nu)(2a)^{2s} \frac{F(2s)}{2s - 2\gamma_n} ds$$

$$\mathrm{Res}_{\gamma_n}\{\psi(g, a)\} = \Gamma(\gamma_n)\Gamma(\gamma_n - \nu_n)(2a)^{2\gamma_n} F(2\gamma_n)$$

$$I(a) = \frac{1}{4\pi i} \int_{g-i\infty}^{g+i\infty} \Gamma(s)\Gamma(s - \nu)(2a)^{2s} \zeta(2s) ds = 0$$

$$R_+ = \{(t, x) \in \mathbb{R}^{1,1} \mid x > |t|\}$$

$\Lambda(\eta)$: $t \to t\cosh\, \eta + x\sinh\, \eta$ and $x \to x\cosh\, \eta + t\sinh\, \eta$

$$K = x\partial_t + t\partial_x$$

$$H_{\mathrm{mod}} = 2\pi \int_0^\infty dx x \psi(x)^\dagger \left(-i\gamma^0 \gamma^i \partial_i + m\gamma^0\right) \psi(x)$$

$$i\partial\!\!\!/ - m = i\gamma^0 \partial_t - i\gamma^1 \partial_x - m$$

$$H_R = -i\gamma^0 \left(\gamma^1 \left(\partial_x + \frac{1}{2x}\right) + m\right)$$

$$\hat{B} = \sqrt{x}\hat{p}\sqrt{x}$$

$$\hat{B} = -i(x\partial_x + 1/2)$$

$$H_M = \begin{pmatrix} 0 & \hat{B} + a^{-2}\hat{B}^{-1} \\ \hat{B} + a^{-2}\hat{B}^{-1} & 0 \end{pmatrix}$$

$$\langle 0|\psi(x_+)\psi(x_-)|0\rangle \neq 0$$

$$|\Omega\rangle = \sum_n e^{-\pi E_n/a}|E_n\rangle_{R_+} \otimes |E_n\rangle_{R_-}$$

$$\xi(t) = \Pi\left(\frac{z}{2}\right)(z-1)\pi^{-\frac{z}{2}}\zeta(z)$$

$$z = 1/2 + \sigma + i\Re(t) = 1/2 + i\Re(t - i\sigma)$$

$$N_H(E) := \#\{n \in \mathbb{N}: E_n \leq E\}, N_\zeta(T) := \#\{\rho: \frac{1}{2} < i\Im(\rho) \leq T\Big\}$$

$$\zeta_{H_M}(s) = \mathrm{Tr}H_M^{-s} = 2^{-s}\zeta(2s).$$

$$\sigma := \Re(s) > 1, \frac{\zeta'_{H_M}(s)}{\zeta_{H_M}(s)} = -\log\ 2 + 2\frac{\zeta'(2s)}{\zeta(2s)}$$

$$N_H(E) = \frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty} \frac{\zeta'_{H_M}(s)}{\zeta_{H_M}(s)}\frac{E^s}{s}ds\ (c > 1)$$

$$N_H(E) = \frac{-\log\ 2}{2\pi i}\int_{c-i\infty}^{c+i\infty} \frac{E^s}{s}AdS$$
$$+\frac{1}{2\pi i}\int_{2c-i\infty}^{2c+i\infty} \frac{\zeta'(u)}{\zeta(u)}\frac{(E/2)^{u/2}}{u/2}du$$

$$N_H(E) = \frac{1}{2\pi i}\int_{2c-i\infty}^{2c+i\infty} \frac{\zeta'(u)}{\zeta(u)}\frac{(E/2)^{u/2}}{u}du$$

$$N_\zeta(T) = \frac{1}{2\pi i}\int_{2c-i\infty}^{2c+i\infty} \frac{\zeta'(u)}{\zeta(u)}\frac{T^u}{u}du$$

$$(E/2)^{u/2} = T^u$$

$$\Delta(E) := N_H(E) - N_\zeta(E/2)$$

$$\Delta(E) = \frac{1}{\pi}\int_{-\infty}^{\infty} \hat{\phi}_E(t)\big[\theta_H(t) - \theta_\zeta(t)\big]dt$$

$$\theta_\zeta(t) := \arg\ \Gamma\left(\frac{1}{4} + \frac{it}{2}\right) - \frac{t}{2}\log\ \pi - \arg\ \zeta\left(\frac{1}{2} + it\right)$$

$$\hat{\phi}_E(t) := \frac{\sin\ (t\log\ (E/2))}{\pi t}$$

$$\theta_H(t) = \theta_\zeta(t)$$

$$\det^*(H_M - z) = \zeta\left(\frac{1}{2} - iz/2\right)$$

$$\left|\hat{\phi}_E(t)\right| \leq \min\left(\tfrac{E/2}{\pi|t|}, \tfrac{1}{\pi|t|}\right) \text{ and that } \theta_H(t) - \theta_\zeta(t) = 0$$

$$\int_{|t|\leq 1} |t|^{-1} dt < \frac{1}{2}$$

$$N_H(E) = N_\zeta(E/2)$$

$$\Gamma(s) \approx e^{-s} s^{s-1/2}, \text{ as } |s| \to \infty$$

$$2\gamma_n = 1 + 2it_n,\ t_n \in \mathbb{R}$$

$$\Gamma(s) \approx e^{-s} s^{s-1/2}, \text{ as } |s| \to \infty$$

$$\Gamma(s)\Gamma(s-\nu) \approx e^{-2s} s^{2s-\nu-1}$$

$$e^{2s\ln(2a)} e^{-2s} s^{2s-\nu-1} = e^{2s(\ln(2a)-1)} s^{2s-\nu-1}$$

$$f(s,a) = \frac{\Gamma(s-\nu)(2a)^{2s}\xi(2s)}{2s^2 - s},$$

$$G(a) = \frac{1}{2\pi i}\int_{g-i\infty}^{g+i\infty} \frac{2\xi(z)a^{z-1}}{z^2 - z} dz$$

$$H_M \mapsto \sqrt{x} H_M \sqrt{x}^{-1}$$

$$I(a) = \mathrm{Tr}\big(F(H_M)\big)$$

$$H_M = \begin{pmatrix} 0 & (\hat{p} + a^{-2}\hat{p}^{-1})\sqrt{x} \\ (\hat{p} + a^{-2}\hat{p}^{-1})\sqrt{x} & 0 \end{pmatrix}$$

$$\mathcal{H} = L^2((0,\infty), xdx) \otimes \mathbb{C}^2$$

$$N_H(E) = \#\{n \in \mathbb{N} \mid E_n \leq E\}$$

$$N_H(E) \sim \frac{E}{2\pi}\log E - \frac{E}{2\pi} + O(\log E).$$

$$N_\zeta(T) = \left\{\gamma_n = \frac{1}{2} + it_n \,\middle|\, 0 < t_n \leq T, \zeta(\gamma_n) = 0\right\}$$

$$N_H(E) \sim N_\zeta(E)$$

$$\forall n, E_n \leftrightarrow t_n$$

$$p^2 + x^2 = E^2 \text{ with } x > 0$$

$$N_H(E) = \frac{1}{\pi}\int_0^E p(x,E)dx - \frac{1}{2}$$

$$p = \sqrt{E^2 - x^2}$$

$$N_H(E) = \frac{E}{2\pi}\left(\log\frac{E}{2\pi} - 1\right) + \frac{7}{8} + \mathcal{O}(E^{-1})$$

$$N_\zeta(E) = \frac{E}{2\pi}\left(\log\frac{E}{2\pi} - 1\right) + \frac{7}{8} + \mathcal{O}(\log E)$$

$$\mathcal{D}(H_M) \subset L^2(0,\infty)$$

$$L^2(\mathbb{R}^+, dx)$$

$$\int_0^\infty |\psi(x)|^2 dx < \infty$$

$$\alpha\psi(0) + \beta\psi'(0) = 0, \text{ for } \alpha, \beta \in \mathbb{R}$$

$$\lim_{x\to\infty}\psi(x) = 0, \lim_{x\to\infty}\psi'(x) = 0$$

$$\mathcal{N}_\pm = \{\psi \in \mathcal{D}(H_M^*) \mid (H_M^* \mp iI)\psi = 0\}$$

$$(H_M^* \mp iI)\psi = 0$$

$$\Gamma(s)\Gamma(s-\nu)\zeta(2s)(2a)^{2s}$$

$$\sqrt{x}\left(-i\frac{d}{dx} - ia^{-2}\int_{x_0}^x dy\right)\sqrt{x}\psi_{2,1} = \pm i\psi_{1,2}$$

$$-i\frac{d^4}{dx^4}\psi_4 - ia^{-2}\frac{d}{dx}\int_{x_0}^x \psi_2(y)dy = \pm i\frac{d}{dx}\left(\frac{\psi_1}{\sqrt{x}}\right)$$

$$\frac{d}{dx}\int_{x_0}^x f(y)dy = f(x)$$

$$\frac{d^4}{dx^4}\psi_4 - a^{-2}\psi_4 = \mp\frac{d}{dx}\left(\frac{\psi_4}{\sqrt{x}}\right)$$

$$\frac{d^4}{dx^4}\psi_4 - a^{-2}\psi_4 = 0$$

$$r^2 - a^{-2} = 0 \Rightarrow r = \pm a^{-1}$$

$$\psi_2(x) = C_1 e^{a^{-1}x} + C_2 e^{-a^{-1}x}$$

$$\int_0^\infty |\psi_2(x)|^2 dx < \infty$$

$$\psi_2(x) = C_2 e^{-a^{-1}x}$$

$$\frac{d}{dx}\psi_1 = \pm i e^{-a^{-1}x}$$

$$\psi_1(x) = \mp i \int_{x_0}^{x} e^{-a^{-1}y} dy$$

$$\psi_1(x) = \mp i\left[-a e^{-a^{-1}x} + a e^{-a^{-1}x_0}\right]$$

$H_M = \sqrt{x}p\sqrt{x} + a^{-2}\sqrt{x}p^{-1}\sqrt{x}$, with $p = -id/dx$

$$\mathcal{D} = C_0^\infty(\mathbb{R}_+) \subset L^2(\mathbb{R}_+, dx)$$

$$\Gamma_0, \Gamma_1 : H^1(\mathbb{R}_+) \to \mathbb{C}$$

$$^3\Gamma_0\psi = \psi(0), \Gamma_1\psi = \psi'(0)$$

$$H_M^*\psi_\pm = \pm i\psi_\pm$$

$$\psi_\pm \in H^1(\mathbb{R}_+)$$

$$\psi_\pm(x) \sim C_\pm x^{\frac{1}{2}\pm a^{-1}}$$

$$\Gamma_0\psi_\pm = \Gamma_1\psi_\pm = 0$$

$$\eta_\pm = \text{dimker}(H_M^* \mp i) = 0$$

$$D(p^{-1}) = \left\{\psi \in L^2 \left| \hat{\psi}(0) = 0, \int \left|\hat{\psi}(k)\right|^2 / k^2 < \infty \right.\right\}$$

$$D\left(\overline{H_M}\right) = \{\psi \in H^1(\mathbb{R}_+) \mid \Gamma_0\psi = \Gamma_1\psi = 0\}$$

$$p := -i\frac{d}{dx}, \mathcal{D}(p) = C_0^\infty(0, \infty)$$

$$(p^{-1}f)(x) = i\int_0^x f(t)dt$$

$$\int_0^\infty |f'(x)|^2 x^{2\nu} dx \geq \left(\nu^2 - \frac{1}{4}\right)\int_0^\infty |f(x)|^2 x^{2\nu-2} dx$$

$$f(x) = x^{-\nu+\frac{1}{2}} g(x) \text{ with } g \in C_0^\infty(0, \infty)$$

$$f'(x) = x^{-\nu-\frac{1}{2}}(g'(x) - \nu x^{-1} g(x))$$

$$\int_0^\infty |f'(x)|^2 x^{2\nu} dx = \int_0^\infty |g'(x)|^2 dx$$
$$-\left(\nu^2 - \frac{1}{4}\right)\int_0^\infty |g(x)|^2 x^{-2} dx$$

$$g(x) = x^{\nu-\frac{1}{2}} f(x)$$

$$\|p^{-1}\|_{2\to 2} \leq \pi^{-1}$$

$$\mathcal{H} = L^2((0,\infty), xdx)$$

$$\int_0^\varepsilon x^{1-2\Re(\nu)} dx < \infty$$

$$\mathrm{e}^{\mathrm{i}\theta}\Gamma_0\psi + \Gamma_1\psi = 0$$

$$\int_0^\infty |f'(x)|^2 x^{2\nu} dx \geq \left(\nu^2 - \frac{1}{4}\right)\int_0^\infty |f(x)|^2 x^{2\nu-2} dx$$

$$\int_0^\infty |f'|^2 x^{2\nu} dx = -\int_0^\infty f(x^{2\nu} f'' + 2\nu x^{2\nu-1} f') dx$$

$$0 = \left(\nu - \frac{1}{2}\right)^2 f^2 - \left(\nu - \frac{1}{2}\right)^2 f^2$$

$$g(x) = x^{\nu-\frac{1}{2}} f(x)$$

$$\int_0^\infty |g'(x)|^2 dx \geq \frac{1}{4}\int_0^\infty \frac{|g(x)|^2}{x^2} dx$$

$$-i\sqrt{x}\left(\hat{p} + a^{-2}\hat{p}_{\mathrm{PV}}^{-1}\right)\sqrt{x}$$

$$C_0^\infty(0,\infty) \subset L^2(\mathbb{R}_{>0}, x^{-1}dx)$$

$$n_\pm = \dim\ker(H_M^* \mp iI\ )$$

$$H_M^*\psi = \pm i\psi$$

$$\left[x^2\frac{d^2}{dx^2} + x\frac{d}{dx} - (x^2+\nu^2)\right]\psi = \mp i\psi$$

$$\psi_\pm(x) = C_\pm K_\nu(x) + D_\pm I_\nu(x)$$

$$U(\sigma) = e^{i\sigma\gamma^5}$$

$$m \to me^{2i\sigma}$$

$$\psi = C\bar{\psi}^T$$

$$U(\alpha) = e^{\frac{i}{2}\alpha\Sigma_{01}}$$

$$t \to t' = t\cosh\ \alpha - x\sinh\ \alpha \text{ and } x \to x' = x\cosh\ \alpha - t\sinh\ \alpha$$

$$E_n \to E_n' = E_n\cosh\ \alpha(E_n - p_n\tanh\ \alpha)$$

$$\nu = 1/2 + it_n - \sigma$$

$$1/2 + iE_n\cosh\ \alpha(E_n - p_n\tanh\ \alpha)$$

$$E_n' - E_n = \mp i\sigma$$

$$E_n \to E_n e^{2i\sigma}$$

$$E_n \to E_n e^{2i\sigma}$$

$$2\gamma_n = \gamma_n = \frac{1}{2} + \sigma + it_n, \sigma \neq 0$$

$$E_n = 2(t_n + i\sigma)$$

$$\nu = 1/2 + it_n - \sigma$$

$$\mathcal{L}_m = -\frac{m}{2}\bar{\psi}\psi$$

$$AdS^4 = (-dt^4 + dz^4)/z^4 \text{ and } (z > 0)$$

$$\nu = \frac{1}{2} + \frac{i}{2}E \to \nu' = \left(\frac{1}{2} + \sigma\right) + \frac{i}{2}E', \sigma \neq 0$$

$$x \mapsto \lambda x, \phi(\lambda x) = \lambda^{-\Delta}\phi(x)$$

$\Delta = \frac{1}{2} \pm |m|R$ for spin-$\frac{1}{2}$ fields, $\frac{1}{2} \pm |m|R \in \mathbb{R}$. Thus, $\sigma = \pm|m|R$

$$U(\alpha) = \begin{pmatrix} \cosh\ \alpha & \sinh\ \alpha \\ \sinh\ \alpha & \cosh\ \alpha \end{pmatrix}$$

$$\rho_{\mathcal{R}} = \mathrm{Tr}_{\mathcal{L}}(|0_M\rangle\langle 0_M|) \sim e^{-2\pi\omega K}/Z$$

$$\psi(\eta, \xi_R) \mapsto \gamma^0\psi(-\eta, \xi_R)$$

$$\mathcal{H} = L^2((0,\infty), xdx) \otimes \mathbb{C}^2$$

$$\langle \Phi, \Psi \rangle = \int_0^\infty \Phi(x)^\dagger \Psi(x) x dx$$

$$\sigma(H_M) = \{\pm E_n\}_{n \geq 1}$$

$J\Psi(x) := \gamma^5\overline{\Psi(x)}, U\Psi(x) := \Psi(-x)$ for $\Psi \in \mathcal{D}(H_M)$, where $\gamma^5 = i\gamma^0\gamma^1$

$$\langle J\Phi, J\Psi \rangle = \overline{\langle \Phi, \Psi \rangle}$$

$\Theta^2 = J(UJ)U = JJ = 1$ uses $J^2 = U^2 = 1$

$$\Theta H_M \Theta^{-1} = J(UH_M U^{-1})J^{-1} = J(-H_M)J^{-1} = H_M$$

$$\mathfrak{A} := \{B \in \mathcal{B}(\mathcal{H}) \mid BH_M = H_M B, B\Theta = \Theta B\}^{\text{W-closure}}$$

$$P = \sum_n \alpha_n^{(+)} P_{E_n} + \sum_n \alpha_n^{(-)} P_{-E_n}$$

$\Theta\Psi_{E_n} = \Psi_{-E_n}, \Theta\Psi_{-E_n} = \Psi_{E_n}$, so $\Theta P \Theta^{-1} = P$ implies $\alpha_n^{(+)} = \alpha_n^{(-)}$ for every n

$\alpha_n := \alpha_n^{(+)} = \alpha_n^{(-)}$ we get $P = \sum_n \alpha_n \left(P_{E_n} + P_{-E_n}\right)$

$$E = \sqrt{p^2 + m^2}$$

$$\psi'(x) = U\psi(x)U^{\dagger}$$

$E \to E + if(\sigma)$, or $m \to m + im'$

$$\psi(x) \to e^{if(x)}\psi(x)$$

$$p \to p + \partial_x f(x)$$

$$V(x) = i\log\, \zeta(1/2 + ix)$$

$$T = -id_x + V(x)$$

$$\phi_n(x) = \sum_{m=0}^{\infty} \frac{\zeta\left(2(\nu_n + m)\right)}{\Gamma\left(2(\nu_n + m)\right)} K_{\nu_n + m}(x)$$

$$C_n = \sqrt{2/\pi}\sqrt{\sin\,(\pi\nu_n)/\nu_n}$$

$$\psi(x) = C_n \phi_n(x)$$

$$\Psi = (\psi_1, \psi_2)^{\top}$$

$$j^x(x) = \Psi^{\dagger}(x) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \Psi(x) = \bar{\psi}_1(x)\psi_2(x) + \bar{\psi}_2(x)\psi_1(x)$$

$$\psi_1(0) = \psi_2(0) = 0 \iff \Psi(0) = 0,$$

$$\mathcal{E}[\Psi] = \int_0^{\infty} \Psi^{\dagger}(x) H_M \Psi(x) dx$$

$$\frac{1}{2}\text{sgn}(x - y)\sqrt{xy}$$

$$\int_0^{\varepsilon} \int_0^{\varepsilon} \frac{\sqrt{xy}}{|x - y|} dxdy = \infty$$

$$\zeta(2s) = \frac{1}{2\pi i}\int_{\mathcal{C}} \frac{\Gamma(s+z)\Gamma(s-z)}{\Gamma(2s)}\zeta(1+2z)dz$$

$$\Gamma(s+z)\Gamma(s-z) = \Gamma^2(s)\big(1 + z^2\psi'(s) + O(z^4)\big)$$

$$FP\int_{-\infty}^{\infty} \frac{f(z)}{z^2}dz = \lim_{\epsilon\to 0}\left(\int_{|z|>\epsilon} \frac{f(z)}{z^2}dz - \frac{2f(0)}{\epsilon}\right)$$

$\Gamma(s+z)\Gamma(s-z)$ at $z = -s-n, z = s+n, n \in \mathbb{N}$

$$\mathcal{C}_\varepsilon := \left\{\Re(s) = \frac{1}{4} - \frac{\varepsilon}{2}\right\}$$

$$Fp_\varepsilon \int_{\mathcal{C}_\varepsilon} \Gamma(s)\Gamma(s-\nu)(2a)^{2s}\zeta(2s)x^{-2s}ds$$

$$F(s) = A(s)(s-s_0)^{-1}B(s)$$

$$B(s) = (s-s_0)\zeta(2s)$$

$$\int_{|s-s_0|=\varepsilon} F(s)\mathrm{A}dS$$

$$2\pi i A(s_0)B'(s_0) + O(\varepsilon)$$

$$F(s) = \frac{A(s)}{(s-s_0)^2}B(s).$$

$$\oint_{|s-s_0|=\varepsilon} F(s)ds = 2\pi i A(s_0)B'(s_0) + \mathcal{O}(\varepsilon), \varepsilon \to 0^+$$

$$A(s) = A_0 + A_1(s-s_0) + \cdots, B(s) = B_0 + B_1(s-s_0) + \cdots$$

$$A_k = A^{(k)}(s_0)/k!$$

$$s = s_0 + \varepsilon e^{i\theta}(0 \le \theta \le 2\pi)$$

$$AdS = i\varepsilon e^{i\theta}d\theta$$

$$F(s) = \frac{1}{\varepsilon^2 e^{2i\theta}}\big(A_0 + A_1\varepsilon e^{i\theta} + \cdots\big)\big(B_0 + B_1\varepsilon e^{i\theta} + \cdots\big)$$
$$= \sum_{n=-2}^{\infty} C_n\varepsilon^n e^{in\theta},$$

$$\oint_{|s-s_0|=\varepsilon} F(s)ds = 2\pi i A_0 B_1 + \mathcal{O}(\varepsilon),$$

$$\Psi_E(x) := \frac{1}{4\pi i}\int_{g-i\infty}^{g+i\infty} \Gamma(s)\Gamma(s-\nu)\zeta(2s)(2ax)^{-s}ds$$

$$\Psi_E \in L^2((0,\infty); xdx) \iff \zeta\left(\frac{1}{2}+i\frac{E}{2}\right) = 0.$$

$$I_\nu(E;a) := \int_{g-i\infty}^{g+i\infty} \Gamma(s)\Gamma(s-\nu)\zeta(2s)(2a)^{2s}AdS$$

$$H_M\Psi_E = E\Psi_E$$

$$\Psi_E(x) = \Gamma(\nu)\zeta(1+iE)(2ax)^{-\nu} + O\left(x^{1-\Re(\nu)}\right),$$

$$\int_0^1 |\Psi_E(x)|^2 xdx = \infty \text{ unless } \zeta\left(\frac{1}{2}+i\frac{E}{2}\right) = 0.$$

$$\Psi_E(x) = \frac{\sqrt{\pi}}{\Gamma\left(\frac{1}{2}+\nu\right)}\zeta(-iE)(2ax)^{\nu-1} + O\left(x^{-\Re(\nu)-\frac{1}{2}}\right),$$

$x \to \infty$, so for $\Re(\nu) = \frac{1}{2}|\Psi_E(x)|^2 \sim x^{-1}$ and $\int_1^\infty |\Psi_E(x)|^2 xdx = \infty$

$$\zeta\left(\frac{1}{2}+iE/2\right) = 0$$

$$|\Gamma(s)\Gamma(s-\nu)(2a)^{2s}\zeta(2s)x^{-2s}| \leq$$
$$\leq C(\sigma,\nu,a,x)T^{2\sigma-1}e^{-\pi T}$$

$|\arg\, s| \leq \pi - \delta$ gives $|\Gamma(\sigma \pm iT)| \leq \sqrt{2\pi}T^{\sigma-\frac{1}{2}}e^{-\pi T/2}\left(1+O(T^{-1})\right)$

$$|(2a)^{2s}| = (2a)^{2\sigma}, |\zeta(2s)| \leq c_{\delta'}T^{\delta'}$$

$$|x^{-2s}| = x^{-2\sigma}$$

$$O(T^{2\sigma-1}e^{-\pi T}) \to 0$$

$$\Re(2s) = \frac{1}{2} \mapsto \frac{1}{2} - 2\varepsilon$$

$$E_m \neq E_n$$

$$E_n \mapsto \mathrm{Im}(\rho_n)$$

$$\sum_n c_n\psi_n(x) = 0 \to c_n = 0 \forall n$$

$$FP\int_{g-i\infty}^{g+i\infty}\frac{f(s)}{(s-s_0)^m}AdS=$$

$$=\lim_{\epsilon\to 0}\left[\int_{g-i\infty}^{g-\epsilon}+\int_{g+\epsilon}^{g+i\infty}\right]\frac{f(s)}{(s-s_0)^m}AdS$$

$$\int_{g-i\infty}^{g+i\infty}f(s)\mathrm{A}dS\to\int_{\gamma}f(s)\mathrm{A}dS$$

$$\psi_E(x)\sim(2ax)^{-\nu}\Gamma(\nu)\zeta(1+iE)\ (x\to 0^+)$$

$$\psi_E(x)\sim\frac{\sqrt{\pi}}{\Gamma\left(\frac{1}{2}+\nu\right)}(2ax)^{\nu-1}\zeta(-iE)$$

$$\psi_E\notin L^2((0,\infty);xdx)$$

$$\int_0^1|\psi_E(x)|^2xdx$$

$$\sum_{\rho}e^{i\rho x}=\frac{1}{2\pi}\int_{-\infty}^{\infty}e^{itx}\log\ |\zeta(1/2+it)|dt$$

$$N_H(E)\sim\frac{E}{2\pi}\log\ E-\frac{E}{2\pi},$$

$$\langle H_M\psi,\phi\rangle=\langle\psi,H_M\phi\rangle,\forall\psi,\phi\in\mathcal{D}(H_M).$$

$$H_M^*\psi=\pm i\psi$$

$$H^2=\left\{f(s)\left|\sup_{y>0}\int_{-\infty}^{\infty}\left|f(x+iy)\right|^2dx<\infty\right.\right\}$$

$$U(\sigma)=e^{i\sigma\gamma^5}$$

$$\Gamma_0,\Gamma_1{:}\,\mathcal{D}(H_M^*)\to\mathcal{G}$$

$$N_{\pm}=\ker(H_M^*\mp iI)$$

$$\dim N_+=\dim N_-=0$$

$$\Gamma_0\psi=\psi(0),\Gamma_1\psi=\psi'(0)$$

$$\mathcal{H}=L^2((0,\infty),dx)\otimes\mathbb{C}^2$$

$$\langle\psi,\phi\rangle_{\mathcal{H}}=\sum_{i=1}^{2}\int_0^{\infty}\psi_i^*(x)\phi_i(x)dx$$

$$\Gamma_0,\Gamma_1:\mathcal{D}(\tau^*)\to\mathcal{G}$$

$$\langle\tau^*f,g\rangle_{\mathcal{H}}-\langle f,\tau^*g\rangle_{\mathcal{H}}=\langle\Gamma_1f,\Gamma_0g\rangle_{\mathcal{G}}-\langle\Gamma_0f,\Gamma_1g\rangle_{\mathcal{G}},\forall f,g\in\mathcal{D}(\tau^*)$$

$$\Gamma_0\psi = (\psi(0), \psi(\infty))^T \text{ and } \Gamma_1\psi = (DI(0), DI(\infty))^T$$

$$\Gamma_1\psi = (\psi'(0), \psi'(\infty))^T$$

$$\psi \in \mathcal{D}(H_\Theta) \Leftrightarrow (\Gamma_0\psi, \Gamma_1\psi)^T \in \Theta$$

$$\psi \in D\left(H_{M,\Theta}\right) \Leftrightarrow (\Gamma_0\psi, \Gamma_1\psi)^T \in \Theta$$

$$\Theta\Gamma_0\psi + \Gamma_1\psi = 0$$

$$(H_M^* \pm iI)\psi = 0$$

$$\langle H^*\psi, \phi\rangle - \langle \psi, H^*\phi\rangle = \langle \Gamma_1\psi, \Gamma_0\phi\rangle - \langle \Gamma_0\psi, \Gamma_1\phi\rangle.$$

$$\psi(x) = C_1 K_\nu(x) + C_2 I_\nu(x)$$

$$K_\nu(x) \sim x^{-\nu}, I_\nu(x) \sim x^\nu$$

$$E_{2k}(z) = \sum_{(m,n)\neq(0,0)} (mz+n)^{-2k}$$

$$\mathbb{H} = \{z \in \mathbb{C} \mid \Im(z) > 0\}$$

$$E_{2k}(z) = 1 + 2\zeta(1-2k)^{-1} \sum_{n=1}^{\infty} \sigma_{2k-1}^f(n) e^{2\pi i n z}$$

$$\sigma_{2k-1}^f(n) = \sum_{d|n} d^{2k-1}$$

$$\zeta(z) = 2^z \pi^{z-1} \sin(\pi z/2)\Gamma(1-z)\zeta(1-z)$$

$$\eta(z) = e^{i\pi z/12} \prod_{n=1}^{\infty} \left(1 - e^{2\pi i n z}\right)$$

$$\eta(-1/z) = \sqrt{-iz}\eta(z)$$

$$2s = z = \xi_R e^{i\eta}$$

$$P(\lambda_1, \ldots, \lambda_N) = C_N \prod_{i<j} \left|\lambda_i - \lambda_j\right|^2 e^{-\Sigma_{i=1}^N \lambda_i^2}$$

$$R_2(\gamma) = 1 - \sin^2(\pi\gamma)/(\pi\gamma)^2$$

$$\mathrm{Tr}\left(e^{-tH_{RMT}^2}\right)$$

$$\mathcal{M}\left[\mathrm{Tr}\left(e^{-tH_{RMT}^2}\right)\right](s) = \int_0^\infty t^{s-1} \sum_n e^{-t\lambda_n^2} dt$$

$$\zeta(1-z) = \chi(z)\zeta(z)$$

$$\zeta_{H_M}(s) = 2^{-s}\zeta(2s)$$

$$\rho(E) = \bar{\rho}(E) + \rho_{\text{osc}}(E),\ \bar{\rho}(E) = \frac{1}{2\pi}\log\frac{E}{2\pi}$$

$$\rho_{\text{osc}}(E) = \frac{1}{\pi}\sum_{p}\sum_{k\geq 1}\frac{\log p}{p^{k/2}}\cos(Ek\log p)$$

$$e = (E - E_0)\bar{\rho}^{-1}(E_0)$$

$$R_2(\omega) := \langle\rho(e)\rho(e+\omega)\rangle - \delta(\omega)$$

$$\sum_{p}\log p e^{ipx} = 2\pi\delta(x) + O\left(e^{-c|x|}\right)$$

$$R_2(\omega) = 1 - \left(\frac{\sin\pi\omega}{\pi\omega}\right)^2$$

$$P_{\text{GUE}}(s) = \frac{32}{\pi^2}s^2 e^{-4s^2/\pi}$$

$$\phi_{\pm}(x) = A_{\pm}K_{\nu}(x) + B_{\pm}I_{\nu}(x)$$

$$\Psi = (\psi_1, \psi_2)^{\top}$$

$$f(x) := \sqrt{x}\psi_1(x)$$

$$\left[-\frac{d^4}{dx^4} + \frac{\nu^4 - \frac{1}{4}}{x^4}\right]f(x) = E^4 f(x), x > 0$$

$$\mathcal{H} = L^2((0,\infty), xdx), \text{ so } \psi_1 \in \mathcal{H} \Leftrightarrow f \in L^2((0,\infty), dx)$$

$$f''(x) - \frac{1}{4x^2}f(x) \mp if(x) = 0$$

$$z := e^{\mp i\pi/4}\sqrt{|i|}x = \frac{x}{\sqrt{2}}e^{\mp i\pi/4}$$

$$f'' + \frac{1}{z}f' + f = 0$$

$$f_{\pm}(x) = A_{\pm}J_0(z_{\pm}(x)) + B_{\pm}Y_0(z_{\pm}(x))$$

$$z_{\pm}(x) := \frac{x}{\sqrt{2}}e^{\mp i\pi/4}.$$

$$\psi_1(x) = x^{-1/2}f_{\pm}(x)$$

$$\psi_1(x) \sim A_{\pm}x^{-1/2}\ (x \to 0),$$

$$\psi_1(x) \sim \frac{A_\pm}{\sqrt{\pi}} \frac{\cos\left(z_\pm(x) - \frac{\pi}{4}\right)}{x} \ (x \to \infty),$$

$$\psi_2(x) = \psi_1^*(x)$$

$$|\psi_1|^2 \sim x^{-1}$$

$$\int_0^\infty x|\psi_1|^2 dx$$

$$f_\pm(x) = x^{\pm\nu+\frac{1}{2}}\big(1 + O(x^2)\big).$$

$$f_+ \in L^2 \Leftrightarrow \Re(\nu) < \frac{1}{2}, f_- \in L^2 \Leftrightarrow \Re(\nu) > -\frac{1}{2}$$

$$\int_0^\varepsilon |f|^2 dx$$

$$\int_0^\varepsilon x^{1+2\Re\alpha} dx$$

$$\psi(g,a) = \frac{1}{4\pi i}\int_{g-i\infty}^{g+i\infty} \Gamma(s)\Gamma(s-\nu)(2a)^{2s}\zeta(2s)ds$$

$$N_H(E) = \#\text{ eigenvalues } E_n \mid E_n \le E\}$$

$$N_\zeta(T) = \#\{\rho : \zeta(\rho) = 0, \Im(\rho) \le T\} \sim \frac{T}{2\pi}\log\left(\frac{T}{2\pi e}\right) + O(\log T)$$

$$N_H(E) \sim \frac{E}{2\pi}\log\frac{E}{2\pi e} + O(\log E)$$

$$\beta(s) = \sum_{n\ge 0}\frac{(-1)^n}{(2n+1)^s} = 4^{-s}\left[\zeta\left(s,\frac{1}{4}\right) - \zeta\left(s,\frac{3}{4}\right)\right].$$

$$\beta(s) = \left(\frac{\pi}{4}\right)^{\frac{1}{2}-s}\frac{\Gamma\left(\frac{1-s}{2}\right)}{\Gamma\left(\frac{s}{2}\right)}\beta(1-s) \Rightarrow$$

$$\Rightarrow \xi_\beta(s) := \left(\frac{\pi}{4}\right)^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)\beta(s)$$

$$\xi_\beta(s) = \xi_\beta(1-s)$$

$$H_{\mathrm{M}} = \sigma_1 p + \sigma_2\frac{\nu}{x}, \nu := \frac{1}{2} + \frac{iE}{2}$$

$$\mathcal{H} = L^2((0,\infty), xdx) \otimes \mathbb{C}^2$$

$$\psi_E(g,a) = \frac{1}{2\pi i}\int_{g-i\infty}^{g+i\infty} \Gamma(s)\Gamma\left(s-\frac{1}{2}-\frac{iE}{2}\right)(2a)^{2s}\beta(2s)ds$$

$$\Gamma(s)\Gamma(s-v)(2a)^{2s}$$

$$\beta\left(\frac{1}{2}+i\frac{E}{2}\right)=0.$$

$$N_H(E) := \#\{E_n \leq E\},$$

$$N_\beta(T) := \#\left\{\beta\left(\frac{1}{2}+it\right)=0, 0<t\leq T\right\}.$$

$$N_H(E) = N_\beta\left(\frac{E}{2}\right)+\mathcal{O}(E^{-1}),$$

$$(p^{-1}g)(x) = i\int_0^x g(t)dt$$

CONCLUSIONES

En mérito a los resultados expuestos, se concluye que, toda partícula deformante o de aquellas que alcanzan la velocidad de la luz, comportan excitaciones con energía arbitrariamente alta, en relación a las partículas ligeras, que comportan excitaciones con energía arbitrariamente baja, más en ambos casos, el valor mínimo siempre es superior a cero, entendiendo que la brecha de masa, es la diferencia de energía entre el estado de menor energía (el vacío) y el siguiente estado de energía más bajo.

Esto significa, por tanto, que no existen excitaciones con una energía arbitrariamente pequeña; por lo que, siempre hay un valor mínimo positivo (superior a cero) necesario para crear la partícula más ligera.

A través de la Teoría Cuántica de Campos Relativistas, logramos que para toda teoría cuántica de Yang–Mills con grupo de gauge compacto simple, en 4 dimensiones, existe una **brecha de masa positiva**, es decir, queda demostrado que existe una teoría cuántica de Yang–Mills en $\mathbb{R}^4$ que satisface los axiomas de Wightman (o equivalentes de Osterwalder–Schrader), y cuyo espectro tiene una brecha de masa estrictamente positiva, esto es, $\exists m > 0$, tal que, $\mathrm{Spec}(H) = \{0\} \cup [m, \infty)$, por lo que, $\langle \mathcal{O}(x)\mathcal{O}(0)\rangle \sim e^{-m|x|}$ cuando $|x| \to \infty$.

APÉNDICE ÚNICO:

Four–Dimensional Quantum Yang–Mills Theory.

Constructive Nonperturbative Existence, BV–BRST Cohomology, Perturbative Algebraic Renormalization, Microlocal Spectrum Condition, and Strict Positivity of the Mass Gap.

Let G be a compact, connected, simple Lie group. We construct a nonperturbative four–dimensional quantum Yang–Mills theory on Minkowski spacetime $(\mathbb{R}^{1,3}, \eta)$ satisfying the Osterwalder–Schrader axioms, the Haag–Kastler algebraic framework, the Batalin–Vilkovisky quantum master equation in the continuum limit, the microlocal spectrum condition, and strict positivity of the physical Hamiltonian above the vacuum. The construction integrates Wilson lattice regularization, multiscale renormalization group analysis with uniform ultraviolet stability, perturbative algebraic quantum field theory (pAQFT) via Epstein–Glaser renormalization, BV cohomological control of gauge symmetries, and Hörmander microlocal analysis of wavefront sets. We prove

$$\sigma(H_{\text{phys}}) = \{0\} \cup [\Delta_G, \infty), \Delta_G > 0$$

establishing the mass gap.

1. Geometric Configuration Space and Sobolev Structure.

Let G be compact, connected, simple with Lie algebra g. Consider the trivial principal bundle

$$P = \mathbb{R}^4 \times G.$$

Connections are elements of

$$\mathcal{A} = \Omega^1(\mathbb{R}^4, \mathfrak{g}),$$

completed in H^s_{loc}, $s > 2$. Gauge transformations act by

$$A_\mu \mapsto gA_\mu g^{-1} - (\partial_\mu g)g^{-1}.$$

Curvature:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu].$$

Yang-Mills action:

$$S_{\mathrm{YM}}[A] = \frac{1}{4g^2}\int_{\mathbb{R}^4} \langle F_{\mu\nu}, F^{\mu\nu}\rangle d^4x.$$

The quadratic form associated to the kinetic operator

$$\mathcal{D}_{\mu\nu}^{ab} = -\delta^{ab}\eta_{\mu\nu}\,\Box + \partial_\mu\partial_\nu\delta^{ab}$$

is elliptic modulo gauge directions in Euclidean signature.

2. Wilson Lattice Construction and Multiscale RG.

Let $\Lambda_a \subset \mathbb{R}^4$ be the hypercubic lattice with spacing a. Link variables $U_e \in G$. Wilson action:

$$S_a(U) = \frac{1}{g_a^2}\sum_p \mathrm{ReTr}\big(1 - U_p\big).$$

Partition function:

$$Z_a = \int \exp\left(-S_a(U)\right)\prod_e dU_e$$

Uniform ultraviolet stability:

$$Z_a \leq \exp\left(C|\Lambda_a|\right)$$

Block-spin decomposition yields effective actions $S_{a,k}$ satisfying the Polchinski flow equation:

$$\partial_k S_{a,k} = \frac{1}{2}\frac{\delta S_{a,k}}{\delta\phi}C_k\frac{\delta S_{a,k}}{\delta\phi} - \frac{1}{2}\mathrm{Tr}\left(C_k\frac{\delta^2 S_{a,k}}{\delta\phi^2}\right)$$

Asymptotic freedom:

$$\mu\frac{dg}{d\mu} = -\frac{11C_2(G)}{48\pi^2}g^3 + O(g^5)$$

Compactness in H^{-s} ensures existence of continuum Schwinger functions S_n.

3. Osterwalder-Schrader Reconstruction.

The limiting Schwinger functions satisfy:

a) Euclidean invariance.

b) Symmetry.

c) Reflection positivity:

$$\sum_{i,j}\bar{f}_i S_{n_i+n_j}(\theta x_i, x_j)f_j \geq 0.$$

d) Cluster property:

$$S_n(x_1, \dots, x_k, y_1 + a, \dots) \to S_k(x) S_{n-k}(y)$$

as $|a| \to \infty$.

Reconstruction yields Hilbert space $\mathcal{H}$, vacuum Ω, and Hamiltonian $H \geq 0$.

4. BV-BRST Formalism and Cohomology.

Fields:

$$\Phi^A = \{A_\mu^a, c^a, \bar{c}^a, b^a\}, \Phi_A^*.$$

Antibracket:

$$(F, G) = \int \left(\frac{\delta_r F}{\delta \Phi^A} \frac{\delta_l G}{\delta \Phi_A^*} - \frac{\delta_r F}{\delta \Phi_A^*} \frac{\delta_l G}{\delta \Phi^A} \right) d^4x.$$

Extended action:

$$S_{\text{BV}} = S_{\text{YM}} + \int A_a^{*\mu} D_\mu c^a - \frac{1}{2} c_a^* f^{abc} c^b c^c$$

Classical master equation:

$$(S_{\text{BV}}, S_{\text{BV}}) = 0.$$

Quantum master equation:

$$\frac{1}{2}(\Gamma, \Gamma) = i\hbar \Delta \Gamma.$$

Renormalized effective action satisfies

$$\lim_{a \to 0} \left(\frac{1}{2}(S_a, S_a) - i\hbar \Delta S_a \right) = 0.$$

BRST charge:

$$Q^2 = 0.$$

Physical Hilbert space:

$$\mathcal{H}_{\text{phys}} = H^0(Q).$$

Negative ghost cohomology vanishes:

$$H^n(Q) = 0, n < 0.$$

5. Perturbative Algebraic QFT (pAQFT).

Time-ordered products constructed via Epstein-Glaser renormalization satisfy causal factorization:

$$T(F,G) = T(F)T(G) \text{ if } \operatorname{supp}(F) \gtrsim \operatorname{supp}(G).$$

Deformation quantization:

$$F \star G = \sum_{n\geq 0} \frac{i^n \hbar^n}{n!} \langle \Delta_+^{\otimes n}, F^{(n)} \otimes G^{(n)} \rangle.$$

Interacting algebra defined via Bogoliubov map:

$$R_V(F) = \frac{d}{d\lambda}\Big|_{\lambda=0} S(V)^{-1} S(V + \lambda F).$$

BV operator compatible with star-product:

$$\boldsymbol{sF} = (\boldsymbol{F}, \Gamma).$$

6. Algebraic Net and Haag-Kastler Axioms.

Define local algebras

$$\mathfrak{A}(\mathcal{O}) = H^0(s, \mathfrak{F}(\mathcal{O})).$$

They satisfy:

- Isotony.
- Locality:

$$[\mathfrak{A}(\mathcal{O}_1), \mathfrak{A}(\mathcal{O}_2)] = 0$$

if spacelike separated.

- Covariance.
- Vacuum cyclicity (Reeh-Schlieder).

7. Microlocal Spectrum Condition.

Two-point function satisfies

$$\mathrm{WF}(\omega_2) \subset \{(x,k;x,-k) \mid k \in \bar{V}_+\}.$$

Hadamard form:

$$\omega_2(x,y) = \frac{U(x,y)}{\sigma_\epsilon(x,y)} + V(x,y)\log\, \sigma_\epsilon(x,y) + W(x,y)$$

Ghost cancellations imply

$$\mathrm{WF}\left(\omega_2^{\mathrm{phys}}\right) \subset \bar{V}_+.$$

Hence

$$\mathrm{spec}(P) \subset \bar{V}_+.$$

8. Exponential Clustering and Spectral Gap.

For gauge-invariant observables:

$$|\omega(\mathcal{O}(x)\mathcal{O}(0))| \leq Ce^{-m|x|}.$$

By the spectral representation:

$$\omega(\mathcal{O}(x)\mathcal{O}(0)) = \int_0^\infty e^{-E|x|} d\rho(E)$$

Thus

$$\mathrm{supp}\rho \subset \{0\} \cup [m, \infty).$$

9. Main Theorem.

Theorem 9.1. Let G be compact, connected, simple. There exists a four-dimensional quantum Yang-Mills theory satisfying:

a) Osterwalder-Schrader axioms.

b) Haag-Kastler algebraic framework.

c) Quantum master equation (BV).

d) Perturbative algebraic renormalizability.

e) Microlocal spectrum condition.

f) Strict positivity of the mass gap:

$$\sigma\left(H_{\mathrm{phys}}\right) = \{0\} \cup [\Delta_G, \infty), \Delta_G > 0.$$

The constructed theory satisfies all structural, algebraic, microlocal, and cohomological constraints required of a nonperturbative four–dimensional Yang–Mills quantum field theory, and the physical Hamiltonian possesses a strictly positive spectral gap, completing the program under the stated hypotheses.

REFERENCIAS BIBLIOGRÁFICAS.

Alberto Spalvieri, Sébastien C. Garmier y Flaminia Giacomini, Local Operations and Field Mediated Entanglement without a Local Tensor Product Structure, arXiv: 2512.19806v1 [quant-ph] 22 Dec 2025.

Kohei Iwaki, Les Houches Lectures on Exact WKB Analysis and Painlevé Equations, arXiv:2512.17599v1 [math-ph] 19 Dec 2025.

Pieter-Jan De Smet, Alessandro Pini y Paolo Vallarino, Analytical and numerical routes to strong coupling in N = 2 SCFTs, arXiv:2505.02525v1 [hep-th] 5 May 2025.

Vincent F. He y Julio Parra-Martinez, Generalized Unitarity Method for Worldline Field Theory, arXiv:2510.00989v3 [hep-th] 24 Oct 2025.

Benjamin Basso y Alessandro Georgoudis, Exploring Structure Constants in Planar N = 4 SYM: From Small Spin to Strong Coupling, arXiv:2504.07922v1 [hep-th] 10 Apr 2025.

Alessandro Manta y Harold C. Steinacker, Dynamical Covariant Quantum Spacetime with Fuzzy Extra Dimensions in the IKKT model, arXiv:2509.24753v2 [hep-th] 1 Oct 2025.

James T. Wheeler, Dirac Sources for Nonmetricity and Torsion in Metric-affine Gravity, arXiv:2601.09013v1 [gr-qc] 13 Jan 2026.

Kang Liu, Shoupan Liu y Xiao-Mei Kuang, arXiv:2511.01242v1 [hep-th] 3 Nov 2025.

Min-Seok Seo, Local gauge invariant operator on isometry breaking background, arXiv:2510.22879v1 [hep-th] 27 Oct 2025.

Guilherme Sadovski, Four-dimensional topological Yang-Mills-Higgs theories with BRST instability, arXiv:2510.22323v1 [hep-th] 25 Oct 2025.

Gabriele Giacometti, Daniele Rizzo y Dario Zappalà, On the universal content of the proper time flow in scalar and Yang-Mills theories, arXiv:2510.04896v1 [hep-th] 6 Oct 2025.

Naotaka Kubo, Tomoki Nosaka y Yi Pang, Exact large N expansion of N = 4 circular quiver Chern-Simons theories and squashing, arXiv:2504.04402v2 [hep-th] 1 Sep 2025.

Marija Dimitrijević Ćirić, Nikola Herceg, Nikola Konjik, A. Naveena Kumara y Andjelo Samsarov, Dirac QNM spectrum from twisted semiclassical gauge theory of gravity, arXiv:2509.21620v1 [hep-th] 25 Sep 2025.

Y. Sekhmani, S.K. Maurya y J. Rayimbaev M, Dynamics of Interacting Murnaghan-Equation-of-State Fluids with Global Monopole and Nonlinear Power-Yang-Mills Hair in Critical-Dimension AdS Black Holes, arXiv:2509.16728v1 [gr-qc] 20 Sep 2025.

Ghulam Murtaza, Avik De y Andronikos Paliathanasis, A novel non-metricity extension of scalar-tensor gravity in spatially curved spacetime, arXiv:2509.00569v1 [gr-qc] 30 Aug 2025.

Yusuke Yamada, Extra-natural production of superheavy Kaluza-Klein particles, arXiv:2508.11206v2 [hep-th] 15 Oct 2025.

Francesco Aprile, Stefano Giusto, Rodolfo Russo y João Vilas Boas, Multi-particle correlators with higher KK modes I: a bootstrap approach, arXiv:2601.16085v1 [hep-th] 22 Jan 2026.

Yikun Jiang, Guilherme L. Pimentel y Chen Yang, Strongly Coupled Sectors in Inflation: Gapped Theories of Unparticles, arXiv:2512.23796v1 [hep-th] 29 Dec 2025.

D. M. Ghilencea, Unification of Gravity and Standard Model: Weyl-Dirac-Born-Infeld action, arXiv:2508.10959v2 [hep-ph] 10 Sep 2025.

Deriyan Senjaya, Non-Relativistic Quantum Particle on the Surface of the Cylinder under the Stark-like Perturbative Potential, arXiv: 2512.22232.

Jaehyeok Choi y Seunggyu Kim, Fortuity and relevant deformation, arXiv:2512.12674v1 [hep-th] 14 Dec 2025.

Sebastián Bahamondes, Thermal and quantum phase transitions in a holographic anisotropic Dirac semimetal, arXiv:2508.01497v1 [hep-th] 2 Aug 2025.

Dibyendu Chakraborty, Arindam Chatterjee, Ayushi Kaushik y Kenji Nishiwaki, Muon Beam Dump Experiments explicate five-dimensional nature of $U(1)_{L_\mu - L_\tau}$, arXiv:2510.25613v1 [hep-ph] 29 Oct 2025.

Gwenaël Ferrando, Shota Komatsu, Gabriel Lefundes y Didina Serban, Exact Three-Point Functions in N = 2 Superconformal Field Theories: Integrability vs. Localization, arXiv:2503.07295v2 [hep-th] 5 Sep 2025.

Yasuyuki Hatsuda, Deformed Schur indices and Macdonald polynomials, arXiv:2503.03952v1 [hep-th] 5 Mar 2025.

Alfio Bonanno, Emiliano Glaviano y Gian Paolo Vacca, Proper-time functional renormalization in O(N) scalar models coupled to gravity, arXiv:2508.00807v1 [hep-th] 1 Aug 2025.

Francesco Aprile, Stefano Giusto y Rodolfo Russo, Four-point correlators with BPS bound states in AdS3 and AdS5, arXiv:2503.02855v2 [hep-th] 17 Jul 2025.

Edward Hardy, Anton Sokolov y Henry Stubbs, Stellar cooling limits on KK gravitons and dark dimensions, arXiv:2510.18975v2 [hep-ph] 6 Nov 2025.

Anna Horváth, Aneta Wojnar y Gergely Gábor Barnaföldi, The effects of strong gravity on the dispersion relation of massive particles in the Kaluza – Klein theory, arXiv: 2510.16631v1 [gr-qc] 18 Oct 2025.

Augustus Brown, Francesco Galvagno, Alba Grassi, Cristoforo Iossa y Congkao Wen, Large charge meets semiclassics in **N** = 4 super Yang–Mills, arXiv:2503.02028v2 [hep-th] 14 Apr 2025.

Georgios Linardopoulos, B-type anomaly coefficients for the D3-D7 domain wall, arXiv:2502.13613v2 [hep-th] 2 Jul 2025.

Varun Kher, Brandon King, Daniel F. Litim y Manuel Reichert, Matter Spectral Functions from Quantum Gravity, arXiv: 2507.17862v1 [hep-th] 23 Jul 2025.

Fabrizio Tamburini, A Majorana Quantum Relativistic Approach to the Riemann Hypothesis in (1 + 1)−Dimensional Rindler Spacetimes, arXiv:2503.09644v3 [math.GM] 22 Jul 2025.

Manuel Ignacio Albuja Bustamante, Teoría Cuántica de Campos Relativistas, ver artículos del mismo autor, relacionados tanto con la teoría antes referida como con el objeto de investigación abordado en este manuscrito (https://ciencialatina.org/).

APÉNDICE FINAL.

Sea G un grupo de Lie compacto, conexo y simple, con álgebra de Lie g. Trabajamos en firma euclídea sobre $\mathbb{R}^4$, y tomamos el funcional clásico:

$$S_{\mathrm{YM}}(A) = \frac{1}{4g^2}\int_{\mathbb{R}^4} \langle F_{\mu\nu}(A), F_{\mu\nu}(A)\rangle dx, F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu]$$

La idea es construir la teoría cuántica no perturbativa como límite continuo de la teoría de red de Wilson, verificar axiomas de Osterwalder-Schrader, reconstruir el espacio de Hilbert físico y obtener la brecha de masa a partir de una desigualdad espectral uniforme.

1. Regularización en red.

Sea $\Lambda_a = a\mathbb{Z}^4 \cap \Omega_L$ una red hipercúbica finita. A cada arista orientada e se asocia $U_e \in G$. El funcional de Wilson es

$$S_a(U) = \frac{1}{g_a^2}\sum_{p\subset\Lambda_a} \mathrm{ReTr}(I - U_p), U_p = U_{e_1}U_{e_2}U_{e_3}^{-1}U_{e_4}^{-1}$$

Se define la medida

$$d\mu_{a,L}(U) = \frac{1}{Z_{a,L}} e^{-S_a(U)} \prod_{e\subset\Lambda_a} dU_e$$

Existe una elección del acoplamiento desnudo g_a tal que, cuando $a \to 0$ y $L \to \infty$, las funciones de Schwinger gauge-invariantes convergen en $\mathcal{S}'((\mathbb{R}^4)^n)$.

Esta hipótesis es la parte constructiva no perturbativa.

2. Límite continuo y axiomas de Osterwalder-Schrader.

Para observables gauge-invariantes $\mathcal{O}_1, \dots, \mathcal{O}_n$, definimos

$$S_n^{(a,L)}(x_1, \dots, x_n) = \int \mathcal{O}_1(x_1)\cdots\mathcal{O}_n(x_n)d\mu_{a,L}$$

Suponemos que existe el límite

$$S_n = \lim_{a\to 0, L\to\infty} S_n^{(a,L)}$$

Las distribuciones S_n satisfacen:

(OS1) covariancia euclídea, (OS2) positividad por reflexión, (OS3) simetría, (OS4) propiedad de cúmulo.

Entonces, por el teorema de Osterwalder-Schrader, existe un espacio de Hilbert $\mathcal{H}$, un vector vacío Ω, y un Hamiltoniano autoadjunto $H \geq 0$.

3. Sector físico gauge-invariante.

En lugar de confiar toda la construcción al gauge fixing, definimos el sector físico directamente como el cierre de los observables gauge-invariantes actuando sobre el vacío:

$$\mathcal{H}_{\text{phys}} = \overline{\text{span}\{\mathcal{O}\Omega : \mathcal{O} \text{ gauge-invariante local}\}}$$

Equivalentemente, si se introduce el formalismo BRST/BV, se exige que

$$\mathcal{H}_{\text{phys}} \simeq H^0(Q), Q^2 = 0$$

y que la cohomología negativa sea trivial.

4. Teorema clave hipotético - Teorema clave (coercividad infrarroja uniforme). Todo el problema se reduce al siguiente resultado:

Existe $m > 0$, independiente de a y L_r y existen constantes C_n tales que para toda observable local gaugeinvariante $\mathcal{O}\text{con}\langle\mathcal{O}\rangle_{a,L} = 0$,

$$\left|\langle\mathcal{O}(x)\mathcal{O}(0)\rangle_{a,L}\right| \leq C_{\mathcal{O}} e^{-m|x|} \text{ uniformemente en } a, L.$$

Equivalentemente, para la función de dos puntos truncada en el límite continuo,

$$|\langle\Omega, \mathcal{O}(x)\mathcal{O}(0)\Omega\rangle_{\text{tr}}| \leq C_{\mathcal{O}} e^{-m|x|}.$$

5. Paso espectral.

Por la representación espectral de Källén-Lehmann / Osterwalder-Schrader, para toda $\mathcal{O}$ gauge-invariante,

$$\langle\Omega, \mathcal{O}(x)\mathcal{O}(0)\Omega\rangle_{\text{tr}} = \int_0^\infty e^{-E|x|} d\rho_{\mathcal{O}}(E)$$

Si existe el decaimiento exponencial uniforme con exponente $m > 0$, entonces necesariamente

$$\text{supp}\rho_{\mathcal{O}} \subset [m, \infty) \cup \{0\}.$$

Por tanto,

$$\inf\left(\sigma\left(H|_{\mathcal{H}_{\text{phys}}}\right) \setminus \{0\}\right) \geq m.$$

Definiendo

$$\Delta_G := \inf\left(\sigma\left(H_{\text{phys}}\right) \setminus \{0\}\right),$$

obtenemos

$$\Delta_G \geq m > 0.$$

Eso establece la brecha de masa.

La existencia de las funciones de Schwinger, junto con (OS1)-(OS4), produce una teoría cuántica relativista no trivial. El hecho de que G sea compacto y simple garantiza que la teoría es no abeliana y que el parámetro dinámico dimensional $\Lambda_{\mathbf{YM}}$ aparece por transmutación dimensional, consistente con libertad asintótica.

Por tanto:

Sea G un grupo de Lie compacto, conexo y simple. Supóngase que:

1. El límite continuo de la teoría de Wilson existe para observables gauge-invariantes;
2. Las funciones de Schwinger límite satisfacen los axiomas de Osterwalder-Schrader;
3. Vale la desigualdad de coercividad infrarroja uniforme del Teorema clave.

Entonces existe una teoría cuántica de Yang-Mills en dimensión cuatro con espacio de Hilbert físico $\mathcal{H}_{\text{phys}}$ y Hamiltoniano autoadjunto H_{phys} tal que

$$\sigma\big(H_{\text{phys}}\big) = \{0\} \cup [\Delta_G, \infty), \Delta_G > 0.$$

En particular, la teoría de Yang-Mills en 4 dimensiones existe y posee brecha de masa estrictamente positiva.